Diverse methanotrophs from alpha and gammaproteobacteria dwell in a stone

quarry: a study from Western India wetland

Jyoti A. Mohite[1,2], Shubha S. Manvi[1,2], Shrinidhi Deshpande[4], Sanjana Patange[4], Rahul A.

Bahulikar[3], Kajal Pardhi[1,2] and Monali C. Rahalkar[*1,2]

[1]C2-83,84, MACS Agharkar Research Institute, G.G. Agarkar Road, Pune 411004,

Maharashtra, India

[2]Savitribai Phule Pune University, Ganeshkhind Road, Pune 411007, Maharashtra, India

[3]BAIF Development Research Foundation, Central Research Station, Urulikanchan, Pune

412202

[4]Modern College, Shivajinagar, Pune

Tel: +91-20-25325119

Fax: +91-20-2651542

*Corresponding author: monalirahalkar@aripune.org

Keywords: Methane, novel, Type I methanotrophs, India, wetlands, stone quarry

Accession numbers:

pmoA: OQ354217, OQ354219, OQ354221

16S rRNA gene: OQ373008, OQ373314, OQ373316, OQ373317

Abstract

Freshwater wetlands are a rich habitat for aerobic or micro-aerophilic methanotrophs. In the present study, we sampled a wetland ecosystem from Western India, a stone quarry with water situated amidst a hill. The wetland is surrounded by the macrophyte *Typha* sp. (cat-tail). To study the methanotroph diversity in this ecosystem, we collected mud samples from the mini-lakes and rhizospheric soil of *Typha* by random sampling. A serial dilution enrichment in liquid followed by isolation on agarose plates in a methane-air atmosphere was used. The high abundance of methanotrophs was indicated by the fact that growth was obtained till dilutions: 10^{-9} or 10^{-10}. In *Typha* root-associated soil, Type I methanotrophs: *Methylomonas koyamae* and uncultured *Methylococcus* were isolated from the lower dilutions: 10^{-3}-10^{-6}. Type II methanotrophs *Methylocystis sp.* and *Methylosinus* spp were isolated from the higher dilutions. The *Methylococcus* strain isolated was about 97.4% (16S rRNA gene similarity) related to *Methylococcus capsulatus* and *Methylococcus geothermalis* and was predicted to represent a putative novel species. Three more methanotrophs, non-axenic, related to *Methylomagnum-Methylocaldum spp*, *Methylocucumis oryzae,* and *Methylocystis* spp, were cultured. Their affiliation was predicted based on the *pmoA* gene sequence. The mud samples showed relatively less diversity and mainly yielded two *Methylomonas* strains related to *M. koyamae*.

Keywords: Wetlands; methanotroph; *Methylocystis* sp., *Methylomonas* sp*.; Methylococcus*; *Methylomagnum* sp., *Methylocucumis* sp.,; *pmoA* gene, *Typhus spp*

While various natural and anthropogenic activities emit methane, wetlands account for massive emitters and absorbers of methane gas [1]. Wetlands occupy 3.8% of the Earth's land surface, amounting to about 20–40% of global CH_4 emissions. Wetlands have anoxic zones where methane is produced. Methanotrophs utilize over 18 to 90% of the methane in wetlands [1], thus simultaneously establishing an equilibrium between methanogenesis and methanotrophy. [1]. Methanotrophs are aerobic to micro-aerophilic bacteria that use methane as the sole carbon source for growth and energy [2]. Aerobic methanotrophs associated with the roots of emergent aquatic plants are the primary colonizers of aquatic environments by making carbon available via methane consumption. Methane consumption by methane-oxidizing bacteria decreases the amount of methane that could be emitted by vegetation [3]. Methane-oxidizing bacteria in the oxic layer or the soil's oxic-anoxic interphase carry out methane oxidation. Arctic, Antarctic, temperate wetlands, and lakes have been studied in detail [4-7], but tropical wetlands have been less studied for their methanotroph ecology.

Stone quarries are unique ecosystems, and the lakes created naturally in such quarries become a unique habitat for methanotrophs. In the present study, we sampled a wetland ecosystem associated with a stone quarry situated amidst hills in Pune city (known as ARAI hills), one of the only natural green habitats in the city. Due to the proximity of the institute ARAI (Automotive research association of India), the hill is popularly known as ARAI hill and is renowned as a potential niche for diverse flora and fauna [8] . The water-filled quarry contains 5-6 mini-lakes (Figure 1) and is rich in vegetation with aquatic weeds (*Typha* spp) and small water plants. One such aquatic weed is *Typha* spp, dominant in the west to the southwestern region of India. Most such investigations have been done on a few plants like rice and *Sphagnum* moss; however, various studies on the interaction between methanotrophs and plants will deepen our understanding of the carbon cycle and help regulate it [9]. Two earlier reports indicated that *Typha* species inhabit methanotrophs in their roots and rhizospheres [1,

10]. Almost no data is available on the methanotrophs dwelling in Indian freshwater wetlands

apart from our study where a novel genus and species from Type Ib methanotrophs was

cultured from a brackish water wetland near a beach, *Methylolobus aquaticus* [11] and *Ca.*

'Methylobacter oryzae' from a rice field near Allepey wetland in Kerala [12]. The present

study is one of the first attempts to document the methanotroph diversity in another freshwater

ecosystem using a cultivation method proven to culture a large diverse population [13].

Mud samples and soil adhered to *Typha* root samples were collected from the wetland patch in

Pune, India 18°32´ N, 73° 48´E (Figure 1) on 30 January 2022 (one root soil sample) and 16

February 2022 (three rhizosphere samples and three mud samples), Figure 1. All the samples

were collected using gloves in sterile plastic vials or sterile plastic bags and proceeded for

enrichment and oxidation of methane. Roots of *Typha* with the attached soil were collected by

uprooting the plant. The roots were briefly washed in distilled water, and only the adhered soil

with cut roots was used as a sample. Mud samples were collected from three different mini-

lakes present at this site using sampling vials (50 ml capacity) from the sides of the lakes with

about 10-15 cm water layer on the top.

Initially, we saw growth of methanotrophs in a serial dilution series set up with a single root

sample of a *Typha* collected on 30 January 2022, and growth was seen up to 10-6 dilutions

within two weeks of incubation. Thus, this site was found to be promising for the enrichment

and isolation of methanotrophs. Therefore, the second sampling was conducted on 16 February

2022 to study the overall methanotroph diversity in this area. Serial dilutions were set up using

modified Nitrate Mineral Salts referred to as dilute NMS medium was prepared [14] from 10^{-1}

to 10^{-10} by adding 1g of the sample to a 9 ml sterile (dilute NMS) medium in 35 ml serum

bottles [14]. A headspace volume of 20% was removed with a sterile syringe and filled with

20% methane. All the further procedures were done as described in our method article [13],

except we directly proceeded for isolation after the first serial dilution enrichment step, as

relatively high number of methanotroph like cells were seen. In all the samples collected from

the quarry, growth was obtained till 10^{-9} or 10^{-10} dilutions in all the samples (Figure 2),

indicating the high abundance of methanotrophs (~10^{-9}-10^{-10} cells/ g fresh weight of

methanotrophs). The presence of abundant methanotrophs in our study site indicates that an

active methane cycle is present here. Microscopically, methanotrophs were either in the form

of large cells, oval or slightly elongated with dark interiors were seen in higher dilutions

(indicative of Type I methanotrophs) or small coccoid cells (indicative of Type II

methanotrophs) were observed in higher enrichment dilutions of 10^{-9} and 10^{-10} (Figure 2). Five

pure cultures of methanotrophs named as strains AQ1-AQ5 were obtained in all from various

dilutions (Figure 2, Table 1). DNA extracted from these pure cultures was subjected to 16S

rRNA gene and *pmoA* gene sequencing. Three more cultures were obtained, showing a minor

amount of small-sized bacteria which could be heterotrophs associated with the methanotrophs.

A single morphotype with large cells of > 5µm length and oval shape indicative of Type I

methanotrophs was seen in these cultures, indicative of *Methylomagnum* and *Methylocucumis*.

We characterized the methanotroph present in these cultures (culture AQ6 and AQ7) using

DNA extraction followed by *pmoA* PCR and sequencing and confirmed the predicted results

obtained by morphological examination (Table 2). Both *Methylomagnum* [15]and

Methylocucumis [16, 17] are known to be methanotrophs with large cells (~5µm) with dark

inclusions, and hence characteristic in morphology. The third culture (AQ8) showed small

coccoid cells was indicative of Type II methanotrophs, identified as *Methylocystis*.

Thus, from *Typha* root-associated soil, Type I methanotrophs from the genera: *Methylomonas*,

Methylococcus, *Methylomagnum* and *Methylocucumis* were isolated/ cultured from the lower

dilutions: 10^{-3}-10^{-6}. Type II methanotrophs *Methylocystis, Methylosinus* spp were

isolated/cultured from the higher dilutions (10^{-9}), Table 1. The mud samples yielded two

Methylomonas strains related to *M. koyamae*, Table 1. The *Methylococcus* strain isolated from

10^{-6} dilution was 97.4% related to *Methylococcus capsulatus and Methylococcus*

geothermalis, on the basis of 16S rRNA gene and the partial *pmoA* gene was 95% related to

that of *Methylococcus geothermalis* and the culture could represent a putative novel species.

We found that small coccoid cells representing Type II methanotrophs of the Methylocystis-

genus producing even milky, white turbidity were seen in all of the higher dilutions higher

dilutions-10^{-8},10^{-9}-10^{-10} in the root-associated methanotroph community. This was confirmed

by further isolating *Methylocystis* from 10^{-9} dilution. On the contrary, Type I methanotrophs

representing *Methylomonas, Methylocaldum-Methylomagnum and Methylocucumis* were

present in lower dilutions, strengthening the observations made earlier in a Chinese wetland

study [1]. In a Chinese wetland, it was observed that *Methylomonas* and *Methylocystis*

dominated the root-associated community of *Typha* species, and the active community was

composed of *Methylocystis* mainly [1]. *Typha* is a popular aquatic weed primarily occurs in

tropical parts of the world. Strong associations of methanotrophs to the roots of emergent weeds

have been published previously [10]. Further studies focusing on the methanotrophs present

associated with *Typha* species in wetlands of various kinds would be more useful. The isolation

of *Methylococcus* spp related to known species on a 97.4% similarity level indicates that this

habitat may have yet un-cultured methanotrophs and could be a rich source for further

isolations. Though *Methylococcus capsulatus* is the most studied methanotroph,

Methylococcus genus has only two valid species, the classical methanotroph *Methylococcus*

capsulatus [18] and the recently described *Methylococcus geothermalis* [19], and hence this

niche could be an exciting habitat for the isolation of new species.

In Pune, temperatures can rise up to 42-43°C in summer, and the temperature of the water can

increase in summer to a considerable extent. In summers, the wetland part also gets dried up,

and hence we found that the roots were dominated by the presence of heat and desiccation-

resistant methanotroph species: *Methylocystis*, *Methylosinus,* and *Methylococcus* [18, 19].

Annual monitoring of this habitat would give us a better picture of temporal diversity of

methanotrophs.

The environmental relevance of methanotrophs as well as their potential applications in

methane mitigation, are both equally important, and hence cultivation of methanotrophs

becomes important to get deeper insight [20]. In the present study, we were successful in

culturing important members of methanotrophs from a crucial ecological niche, a freshwater

wetland, and methanotrophs associated with *Typha* spp. This study would lay a foundation for

further studies focusing on methane mitigation from such sites.

AUTHOR CONTRIBUTIONS: MR designed the experiments, collected the samples, guided

all the experiments, and wrote the manuscript. RB, JM, SM, SD, and SP did the sampling and

performed the enrichment and isolation of the methanotrophs. JM performed the phylogenetic

analysis and prepared the trees. KP assisted in the isolation and maintenance of cultures. RB

and MR found this unique sampling site, did the initial sampling, and co-designed the

experiments. RB, JM, and SM assisted in writing the manuscript. All the authors reviewed the

manuscript and approved the final version.

ACKNOWLEDGMENTS: MCR acknowledges SERB for the POWER fellowship

(SPF/2022/000045) provided to her. KP is thankful to UGC for JRF and JAM to SARTHI for

providing the JRF. SSM acknowledges SERB project: CRG/2021/000941 for the JRF.

References:

1. Cui J, Zhao J, Wang Z, Cao W, Zhang S, Liu J, Bao Z (2020) Diversity of active root-associated
methanotrophs of three emergent plants in a eutrophic wetland in northerm China AMB
Express 10: 1-9.
2. Hanson RS, Hanson TE (1996) Methanotrophic bacteria. Microbiology and Molecular Biology
Reviews 60: 439-471.
3. Laanbroek HJ (2010) Methane emission from natural wetlands: interplay between emergent
macrophytes and soil microbial processes. A mini-review. Annals of Botany 105: 141-153.
4. Graef C, Hestnes AG, Svenning MM, Frenzel P (2011) The active methanotrophic community
in a wetland from the High Arctic. Environmental microbiology reports 3: 466-472.
5. Michaud AB, Dore JE, Achberger AM, Christner BC, Mitchell AC, Skidmore ML, Vick-Majors
TJ, Priscu JC (2017) Microbial oxidation as a methane sink beneath the West Antarctic Ice
Sheet. Nature Geoscience 10: 582-585.
6. Wartiainen I, Hestnes AG, McDonald IR, Svenning MM (2006) *Methylobacter tundripaludum*
sp. nov., a methane oxidizing bacterium from Arctic wetland soil on the Svalbard islands,

Norway (786 N). International Journal of Systematic and Evolutionary Microbiology 56: 109-
113.

7.   Rahalkar M (2007) Aerobic methanotrophic bacterial communities from sediments of Lake
Constance. University of Konstanz

8.   Patil R, Mahajan M (2018) Herbaceous flora of weeds growing at ARAI hills. International
journal of researches in biosciences, agriculture and technology: 16-19.

9.   Yoshida N, Iguchi H, Yurimoto H, Murakami A, Sakai Y (2014) Aquatic plant surface as a niche
for methanotrophs. Frontiers in Microbiology 5: 30. doi: 10.3389/fmicb.2014.00030

10.   Fausser AC, Hoppert M, Walther P, Kazda M (2012) Roots of the wetland plants *Typha*
latifolia and *Phragmitis australis* are inhabited by methanotrophic bacteria in biofilms. Flora
207: 775-782.

11.   Rahalkar MC, Khatri K, Mohite J, Pandit P, Bahulikar R (2020) A novel Type I methanotroph
Methylolobus aquaticus gen. nov. sp. nov. isolated from a tropical wetland. Antonie van
Leeuwenhoek 113: 959-971.

12.   Khatri K, Mohite JA, Pandit PS, Bahulikar RA, Rahalkar MC (2019) Description of '*Ca.*
Methylobacter oryzae' KRF1, a novel species from the environmentally important
Methylobacter clade 2. Antonie van Leeuwenhoek 113: 729-735.

13.   Rahalkar M, Khatri K, Pandit P, Bahulikar R, Mohite J (2021) Cultivation of Important
Methanotrophs From Indian Rice Fields. Frontiers in Microbiology 12: 1-15.

14.   Pandit PS, Rahalkar MC, Dhakephalkar PK, Ranade DR, Pore S, Arora P, Kapse N (2016)
Deciphering community structure of methanotrophs dwelling in rice rhizospheres of an
Indian rice field using cultivation and cultivation-independent approaches. Microbial Ecology
71: 634-644.

15.   Khalifa A, Lee CG, Ogiso T, Ueno C, Dianou D, Demachi T, Katayama A, Asakawa S (2015)
Methylomagnum ishizawai gen. nov., sp. nov., a mesophilic type I methanotroph isolated
from rice rhizosphere. International Journal of Systematic and Evolutionary Microbiology 65:
3527-3534.

16.   Pandit P, Rahalkar MC (2018) Renaming of 'Candidatus Methylocucumis oryzae' as
Methylocucumis oryzae gen. nov., sp. nov., a novel Type I methanotroph isolated from India.
Antonie Van Leeuwenhoek 112: 955-959.

17.   Pandit PS, Hoppert M, Rahalkar MC (2018) Description of '*Candidatus* Methylocucumis
oryzae', a novel Type I methanotroph with large cells and pale pink colour, isolated from an
Indian rice field. Antonie Van Leeuwenhoek 111: 2473-2484.

18.   Bowman J (2000) The methanotrophs - the families Methylococcaceae and
Methylocystaceae. In: Dworkin, M, Falkow, S, Rosenberg, E, Schleifer, K-H, Stackebrandt, E
(eds.) The Prokaryotes: A handbook on the biology of bacteria: ecophysiology, isolation,
identification, applications. Springer Verlag, New York

19.   Awala SI, Bellosillo LA, Gwak J-H, Nguyen N-L, Kim S-J, Lee B-H, Rhee S-K (2020)
Methylococcus geothermalis sp. nov., a methanotroph isolated from a geothermal field in
the Republic of Korea. International Journal of Systematic and Evolutionary Microbiology 70:
5520-5530.

20.   Cervantes F, Garcia S, Peura S, Balagurusamy N (2021) Editorial: Methanotrophs: Diversity,
environmental relevance and applications. Frontiers in Microbiology 12:796861: 1-3.

FIGURES:

Figure no. 1: Site of sample collection: 18°32´ N, 73° 48´ ARAI quarry, Pune a, b and c.

Typha plants

Enrichment of root and mud samples in NMS broth medium, headspace contains methane and air (20:80) incubation for ~4-6 weeks

Incubation of streaked plates in desiccator under methane and air (20:80) for 4-6 weeks at room temperature

Methanotrophic cultures after incubation

| *Methylosinus* sp. AQ1 | *Methylomonas* sp. AQ2 | *Methylomonas* sp. AQ3 | *Methylococcus* sp. AQ4 | *Methylocucumis* sp. -non-axenic | *Methylomagnum /Methylocaldum* sp. -non-axenic |

Figure No. 2: Enrichment, isolation, and identification of methanotrophs from ARAI stone quarry *Typha* plant root and mud samples

257   TABLE:

258   Table no. 1: Summary of type I and type II methanotrophs obtained in this study.

Sr No.	Name of the methanotroph at genus level & NCBI accession number	Sample name & positive dilutions Sampling date: 30.01.2022 (1) Sampling date: 16.02.2022 (2)	Universal 16s rRNA gene sequence BLAST hit name, percent similarity and NCBI accession number	*pmoA* gene sequence BLAST hit name, percent similarity and NCBI accession number
1.	*Methylosinus* sp. AQ1 16s (OQ373008)	ARAI root (1) sample dilution 10^{-6} (2)	*Methylosinus sporium* 98.99% (NR026512.1)	*Methylosinus trichosporium* 94.55% (CP023737.1)
2.	*Methylmonas* sp. AQ2 16s (OQ373314)	ARAI root (1) sample dilution 10^{-3} (2)	*Methylomonas koyamae* 99.27% (AP019777.1)	*Methylomonas koyamae* 96.06% (AP019777.1)
3.	*Methylomonas* sp. AQ3 16s (OQ373316)	ARAI mud (7) sample dilution 10^{-5} (2)	*Methylomonas koyamae* 99.27% (AP019777.1)	*Methylomonas koyamae* 96.29% (AP019777.1)
4.	*Methylococcus* sp. AQ4 16s (OQ373317)	ARAI root (1) sample dilution 10^{-6} (2)	*Methylococcus capsulatus* 97.47% (NR042183.1)	*Methylococcus geothermalis* 94.58% (CP046565.1)
5.	*Methylomonas* sp. AQ5 16s (OQ373316)	ARAI mud (7) sample dilution 10^{-9} (2)	*Methylomonas koyamae* 99.27% (AP019777.1)	*Methylomonas koyamae* 96.29% (AP019777.1)
6.	*Methylocucumis* sp. AQ6 *pmoA* (OQ354217)	ARAI root sample dilution 10^{-4} (1)	Non-axenic culture	*Methylocucumis oryzae* Sn10-6 97.93% (KT180167.1)
7.	*Methylomagnum* sp. AQ7 *pmoA* (OQ354221)	ARAI water sample dilution 10^{-3} (1)	Non-axenic culture	*Methylomagnum ishizawai* 89.45% (AP019783.1)
8.	*Methylocystis* sp. AQ8 *pmoA* (OQ354219)	ARAI root sample dilution 10^{-9} (1)	Non-axenic culture	*Methylocystis hirsuta* CSC1 97.07% (DQ364434.1)