Area postrema neurons mediate interleukin-6 function in cancer-associated cachexia

Qingtao Sun[1,#], Daniëlle van de Lisdonk[1,2,#], Miriam Ferrer[1,#], Bruno Gegenhuber[1], Melody Wu[1], Jessica Tollkuhn[1], Tobias Janowitz[1], Bo Li[1*]

1. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY 11724, USA
2. Center for Neuroscience, University of Amsterdam, Amsterdam, the Netherlands

\# These authors contributed equally

*Correspondence: bli@cshl.edu (B.L.)

Interleukin-6 (IL-6) has been long considered a key player in cancer-associated cachexia[1-

15]. It is believed that sustained elevation of IL-6 production during cancer progression

causes brain dysfunctions, which ultimately result in cachexia[16-20]. However, how

peripheral IL-6 influences the brain remains poorly understood. Here we show that

neurons in the area postrema (AP), a circumventricular structure in the hindbrain,

mediate the function of IL-6 in cancer-associated cachexia in mice. We found that

circulating IL-6 can rapidly enter the AP and activate AP neurons. Peripheral tumor,

known to increase circulating IL-6[1-5,15,18,21-23], leads to elevated IL-6 and neuronal

hyperactivity in the AP, and causes potentiated excitatory synaptic transmission onto AP

neurons. Remarkably, neutralization of IL-6 in the brain of tumor-bearing mice with an

IL-6 antibody prevents cachexia, reduces the hyperactivity in an AP network, and

markedly prolongs lifespan. Furthermore, suppression of *Il6ra*, the gene encoding IL-6

receptor, specifically in AP neurons with CRISPR/dCas9 interference achieves similar

effects. Silencing of Gfral-expressing AP neurons also ameliorates the cancer-associated

cachectic phenotypes and AP network hyperactivity. Our study identifies a central

mechanism underlying the function of peripheral IL-6, which may serve as a target for

treating cancer-associated cachexia.

Cancer-associated cachexia is a devastating metabolic wasting syndrome characterized by

anorexia, fatigue, and dramatic involuntary bodyweight loss[18,19,24,25]. It affects 50-80% of cancer

patients, lowering the quality of life, reducing tolerance to anticancer therapies, and drastically

accelerating death[19,20]. The brain is known to have an important role in the pathogenesis of

cancer-associated cachexia[16-18]. In particular, recent studies implicate the hypothalamus,

parabrachial nucleus, area postrema and other hindbrain structures in the development of

cachectic phenotypes in animal models of cancer, such as anorexia, weight loss, and accelerated

catabolic processes[26-32]. However, how the brain senses and reacts to peripheral cancers, thereby

contributing to the development of cachectic phenotypes, is not well understood.

Possible mediators of cancer-associated cachexia that may act as messengers to engage the brain

during cancer progression include tumor-derived factors, metabolites from organs indirectly

affected by tumor, and immune or inflammatory factors altered by tumor[16-18,24,33,34]. One such

messenger is the pleiotropic cytokine IL-6[18-20,23,24,35,36]. Indeed, elevated levels of circulating IL-

6 are associated with cancer progression and cachexia in patients and animal models[1-5,15].

Systemic administration of antibodies against IL-6 or IL-6 receptor shows anticachectic effects

in human case reports[6,7,37,38]. Consistently, cancer-associated cachexia in mouse models can be

ameliorated by peripheral administration of antibodies against IL-6[8-13] or IL-6 receptor[14], or by

deletion of the *Il6* gene[9,10]. These findings strongly indicate that IL-6 is a key mediator of cancer-

associated cachexia.

Most studies and therapeutic explorations on IL-6 in cancer-associated cachexia has focused on

its functions in peripheral organs, including the skeletal muscle, liver, and gut[23]. Although

previous studies suggest that IL-6 may also influence brain functions – such as the regulation of

food intake[39-41], fever[42] and the hypothalamic-pituitary-adrenal (HPA) axis[43] – how peripheral

IL-6 is involved in these functions is unclear. In principle, IL-6 can activate its receptors on the

terminals of peripheral nerves, which then transmit the signals to the brain[44]. Alternatively,

circulating IL-6 may cross the blood-brain barrier (BBB) or reach circumventricular organs that

lack or have a weak BBB, thereby acting within the brain[43,45,46].

The area postrema senses circulating IL-6

To determine if circulating IL-6 can enter the brain, we administered biotinylated IL-6 to the

venous sinus of mice through retro-orbital injection (Methods). Three hours later, we prepared

brain sections from these mice in which the presence of the exogenous IL-6 was examined on the

basis of avidin-biotin interactions. In the entire brain, we detected the peripherally administered

IL-6 only in the area postrema (AP) (Fig. 1a-d; Extended Data Fig. 1), which is a

circumventricular organ located outside of the BBB and has been implicated in nausea and

vomiting response to emetic agents entering the circulation[47-50]. Furthermore,

immunohistochemistry in the AP revealed that the IL-6 administration markedly increased the

expression of Fos (Fig. 1e, f), an immediate early gene product linked to recent neuronal

activation[51,52].

Single molecule fluorescent *in situ* hybridization (smFISH) revealed that the *Il6ra*-expressing

(*Il6ra*[+]) cells in the AP partially overlapped with *glucagon-like peptide 1 receptor*-expressing

(*Glp1r*[+]) neurons (Fig. 1g-j), the major excitatory neuronal type in the AP[50]. About 17-18% of all

the detected AP cells (which likely included glia cells) expressed both *Il6ra* and *Glp1r* (Fig. 1h

& j). These *Il6ra*[+] cells also partially overlapped with *Gfral*-expressing (*Gfral*[+]) neurons (Fig. 1g,

h), a subpopulation of *Glp1r*[+] neurons in the AP that have been implicated in nausea and cancer-

associated cachexia[28,29,50]. Interestingly, intravenous IL-6 administration induced *Fos* expression

mainly in the *Il6ra*[+] cells, and these cells partially overlapped with the *Glp1r*[+] neurons (Fig. 1i,

j). These results demonstrate that increased IL-6 in circulation is readily "sensed" by the AP,

where it leads to neuronal activation within hours.

Tumor causes AP hyperactivity

To investigate whether cancer, which is known to increase circulating IL-6[1-5,15], affects AP

neurons, we used mice inoculated with the C26 adenocarcinoma (Fig. 2a, Methods). Mice in this

model show persistent increase in blood IL-6 levels, followed by robust cachectic phenotypes,

including anorexia and dramatic bodyweight loss[15,18,21-23]. We first measured IL-6 levels in the

AP at different timepoints in this model. Notably, IL-6 was increased in the AP on day 7

following tumor inoculation, as well as after the animals had developed cachexia (Fig. 2b).

We then examined Fos expression in the brain in this model before the onset of cachexia (11

days after tumor inoculation). The number of Fos[+] cells in the AP was markedly increased in the

tumor-bearing mice compared with control mice (Fig. 2c, d). Interestingly, the parabrachial

nucleus (PBN), the paraventricular nucleus of the hypothalamus (PVN), the bed nucleus of the

stria terminalis (BNST), and the central amygdala (CeA) – which are structures previously

implicated in cancer cachexia[17,27-29] – also showed tumor-induced increase in Fos[+] cells. Given

these structures are interconnected[53-56] and receive monosynaptic or di-synaptic inputs from the

AP[50,57], these results suggest that cancer progression leads to increased activities in a network of

brain areas encompassing the AP.

Next, we tested whether cachexia is associated with lasting functional changes in AP neurons.

We prepared acute brain slices from healthy control mice as well as the tumor-bearing mice

already showing cachectic phenotypes, and recorded synaptic responses from AP neurons (Fig.

2e). We found that the amplitude of miniature excitatory postsynaptic currents (EPSCs) was

markedly increased, while the frequency of the EPSCs was unchanged in cachectic mice relative

to control mice (Fig. 2f, g). In contrast, there was no difference in the inhibitory postsynaptic

currents (IPSCs) between cachectic mice and control mice (Fig. 2h, i). These results demonstrate

that cancer-associated cachexia is accompanied by a potentiation of excitatory synaptic

transmission onto AP neurons, which may lead to hyperactivity in these neurons.

Neutralizing IL-6 in the brain prevents cancer-associated cachexia and AP hyperactivity

We reasoned that increased circulating IL-6 during cancer progression readily enters the AP and

results in AP neuron activation – like the intravenously administered exogenous IL-6 (Fig. 1) –

leading to hyperactivity in these neurons, which contributes to the development of cachexia.

As a first step to test this hypothesis, we neutralized IL-6 in the brain by intracerebroventricular

(i.c.v.) infusion of an antibody against IL-6, which was achieved using an implanted miniature

pump (Methods). Continuous infusion of the IL-6 antibody, or an isotype control antibody, was

initiated at 10 or 12 days after tumor inoculation (Fig. 3a, b; Extended Data Fig. 2a), a stage

when AP neurons show elevated activity (Fig. 2c-d) but cachexia has not yet started.

Remarkably, compared with the control antibody, the IL-6 antibody prevented the cachectic

phenotypes in almost all the mice, prolonging lifespans (Fig. 3c), reducing bodyweight and tissue

loss (Fig. 3d, e; Extended Data Fig. 2b, c), increasing food and water intake (Fig. 3f), and

increasing blood glucose levels (Extended Data Fig. 2d). Moreover, IL-6 antibody infusion

reduced Fos expression in the AP, PBN, PVN, BNST, and, to a lesser extent, CeA (Fig. 3g, h).

As expected, i.c.v. infusion of IL-6 antibodies did not stop tumor from growing (Extended Data

Fig. 2e, f).

Thus, reducing IL-6 levels in the brain during cancer progression effectively prevents cachexia,

and also dampens the cancer-associated hyperactivity in the AP network.

Suppression of *Il6ra* expression in AP neurons ameliorates cancer-associated cachexia

To investigate whether AP neurons mediate the functions of IL-6 in the development of cancer-

associated cachexia, we sought to suppress *Il6ra*, the gene encoding IL-6Rα, in these neurons

using a recently developed CRISPR/dCas9 interference system. This system consists of dCas9-

KRAB-MeCP2 [a fusion protein including the nuclease-dead Cas9 (dCas9), a Krüppel-

associated box (KRAB) repressor domain, and the methyl-CpG binding protein 2 (MeCP2)] for

transcriptional repression, and a CRISPR sgRNA (single guide RNA) for targeting the promoter

region of genes of interest[58,59]. We designed and identified two sgRNAs, *Il6ra*-sgRNA-4 and -6,

that resulted in suppression of *Il6ra* transcription when co-expressed with dCas9-KRAB-MeCP2

in *in vitro* screening (Extended Data Fig. 3).

We then injected the AP with a lentivirus expressing dCas9-KRAB-MeCP2 in neurons (lenti-

SYN-FLAG-dCas9-KRAB-MeCP2), together with a lentivirus expressing *Il6ra*-sgRNA-4 or -6

(lenti-U6-*Il6ra* sgRNA-4/EF1α-mCherry or lenti-U6-*Il6ra* sgRNA-6/EF1α-mCherry,

respectively), or a sgRNA targeting the bacterial gene *lacZ* as a control (lenti-U6-*lacZ*

sgRNA/EF1α-mCherry)[58] (Fig. 4a, b). Two weeks later, these mice were inoculated with the

C26 tumor.

Compared with the *lacZ* sgRNA control group, the *Il6ra*-sgRNA-4 group had reduced *Il6ra*

expression in AP neurons, including the *Gfral*[+] neurons (Fig. 4c, d). Notably, the *Il6ra*-sgRNA-4

group had markedly increased lifespans (Fig. 4e), an effect that was inversely correlated with the

percentage of *Gfral*[+] neurons that had detectable *Il6ra* expression (Fig. 4f). The *Il6ra*-sgRNA-4

group also had reduced bodyweight loss (Fig. 4g, h), increased food and water intake (Fig. 4i, j),

increased blood glucose levels (Extended Data Fig. 4a), and had a tendency to reduce tissue loss

(Extended Data Fig. 4b, c). At the endpoint of the experiment, mice in the *Il6ra*-sgRNA-4 group

had larger tumor and spleen compared with mice in the control group (Extended Data Fig. 4d, e),

presumably because of the increase in lifespan in the former group.

In a separate experiment, the *Il6ra*-sgRNA-4 mice and the *lacZ*-sgRNA control mice were

sacrificed on the same day, as soon as the latter had developed cachexia (Extended Data Fig. 5a).

This design ensured that the two groups had tumor for the same duration and thus had

comparable tumor and spleen mass (Extended Data Fig. 5g, h). Consistent with the above

observations, the *Il6ra*-sgRNA-4 group had reduced bodyweight loss and tissue loss (Extended

Data Fig. 5b-d), even though these mice had similar food and water intake to the control mice

before the termination of this experiment (Extended Data Fig. 5e, f). Interestingly, the *Il6ra*-

sgRNA-4 mice had reduced Fos expression compared with the *lacZ* sgRNA mice in the AP,

PBN, and PVN (Extended Data Fig. 5i, j), suggesting that suppression of *Il6ra* expression in AP

neurons lowers the hyperactivity in the AP network.

Additional experiments showed that the other sgRNA, the *Il6ra* sgRNA-6, had similar effects to

those of *Il6ra* sgRNA-4 (Extended Data Fig. 6). Together, these results indicate that the IL-6Rα

on AP neurons, especially that on *Gfral+* neurons, is a critical mediator of IL-6 function in the

development of cancer-associated cachexia.

Inhibition of Gfral+ neurons ameliorates cancer-induced anorexia

Given the result that the survival of mice in the C26 model was correlated with the suppression

of *Il6ra* in *Gfral+* neurons in the AP (Fig. 4f), and previous findings that Gfral and its ligand

GDF-15 (Growth/differentiation factor 15) are involved in cancer-associated cachexia[28,29], we

reasoned that the activity of *Gfral+* AP neurons contributes to the development of this syndrome.

To test this hypothesis, we sought to selectively manipulate these neurons using the *Gfral-p2a-*

Cre mice[50] in combination with a Cre-dependent viral approach. As the C26 model necessitates

the use of Balb/c or CD2F1 mice[60], it is incompatible with the *Gfral-p2a-Cre* mice which have a

C57BL/6 genetic background. Therefore, for this experiment, we used the implantable Lewis

lung carcinoma (LLC) model, another established murine cancer model exhibiting features of

cachexia, albeit milder than those of the C26 model[18,23,27].

We first measured the levels of IL-6 and GDF-15 in the blood at different timepoints of cancer

progression in this model. Plasma IL-6 and GDF-15 were both increased at around two weeks

following tumor inoculation (Fig. 5a). To specifically inhibit the activity of *Gfral+* neurons, we

injected the AP of *Gfral-p2a-Cre* mice bilaterally with an adeno-associated virus (AAV)

expressing the tetanus toxin light chain (TeLC, which blocks neurotransmitter release[61]), or GFP

(as a control) in a Cre-dependent manner. Two weeks later, these mice were inoculated with the

LLC (Fig. 5b, c). Interestingly, compared with the GFP mice, the TeLC mice exhibited increased

food intake at the late stage of cancer progression (Fig. 5d) and increased fat and muscle mass at

the endpoint (Fig. 5e). Moreover, the TeLC mice showed reduced Fos expression than the GFP

mice in the AP, PBN, PVN, CeA and BNST (Fig. 5f, g). The two groups had similar tumor and

spleen mass (Fig. 5h). These results indicate that reducing $Gfral^+$ AP neuron activity ameliorates

the cachectic phenotypes. In addition, other areas in the AP network are likely also involved in

this process.

Discussion

Our study identifies AP neurons as a critical mediator of the function of IL-6 that leads to

cancer-associated cachexia in mice. We show that, first, circulating IL-6 rapidly enters the AP

and causes AP neuron activation within hours. Second, periphery tumor results in increased IL-6

and hyperactivity in the AP, and induces enhanced excitatory synaptic drive onto AP neurons.

Third, neutralization of IL-6 in the brain of tumor-bearing mice, via i.c.v. infusion of an IL-6

antibody, prevents cachexia, counteracts the cachexia-associated hyperactivity in the AP

network, and markedly prolongs lifespan. Fourth, specific suppression of $Il6ra$ in AP neurons

with CRISPR/dCas9 interference has similar effects. Lastly, specific suppression of the activities

of $Gfral^+$ neurons, which partially overlap with $Il6ra^+$ cells in the AP (Fig. 1h) and are involved

in the effects of $Il6ra$ suppression (Fig. 4f), also ameliorates cancer-associated cachectic

phenotypes and AP network hyperactivity.

The AP network showing cachexia-associated hyperactivity includes the PBN, the PVN, the

BNST, and the CeA besides the AP. These structures are interconnected[50,53-57] and have been

implicated in regulating feeding behavior and metabolism[50,56,62-69]. In particular, the AP and the

neighboring nucleus tractus solitarii (NTS), as well as the PBN and the PVN have previously

been implicated in cancer-associated cachexia[17,27-29]. The AP sends direct projections to the PBN

and the NTS, and the NTS also directly projects to the PBN as well as the PVN[50,57,70]. Given

previous findings that both the PBN[27,50,55,56,70-72] and the PVN[16,73,74] are involved in feeding

suppression, it is possible that the AP drives cancer-associated anorexia via the AP→PBN, the

AP→NTS→PBN, or the AP→NTS→PVN pathway.

A notable observation is that the AP also drives weight loss independent of anorexia during

cancer progression (Extended Data Fig. 5), consistent with findings that cancer-associated

cachexia involves active catabolic processes in addition to anorexia, and the tissue wasting can

only be partially reversed by nutritional support[18-20]. Interestingly, multiple nuclei in the AP

network are connected with mechanisms that promote catabolic process in peripheral

organs[62,63,66,69,75], providing an anatomical basis for this function of the AP.

Recent studies indicate that Gfral is exclusively expressed by neurons in the AP and the NTS[76-

79], and systemic administration of GDF-15 activates GFRAL$^+$ neurons in the AP and induces

vomiting and anorexia[28,50,80-83]. Furthermore, neutralization of Gfral or GDF-15 with antibodies

ameliorates cancer-associated cachectic phenotypes in animals[28,29]. Thus, GDF-15 may also

influence cancer-associated cachexia, like IL-6, through the AP network. However, as GDF-15

functions as a central alert to the organism in response to a broad range of stressors[84], including

infection, blockade of GDF-15/GFRAL is likely to have detrimental effects if used as a

therapeutic strategy. Indeed, it has recently been shown that GDF-15 is essential for surviving

bacterial and viral infections[65].

IL-6 has long been known as a key contributor to cancer-associated cachexia[18-20,23,24,35,36]. Efforts

exploring IL-6 as a potential therapeutic target thus far have been focused on peripheral IL-6 or

IL-6 receptors, and relied on systemic application of antibodies against these molecules[18,23,38].

However, such systemic approach may not be effective in reducing IL-6 signaling in the brain.

Furthermore, as IL-6 is a pleiotropic cytokine essential for immune and metabolic functions,

with receptors widely distributed in the entire organism[23,85], systemic neutralization of IL-6 or its

receptors will compromise these functions globally and likely cause severe side effects[86,87]. Our

results suggest that targeting IL-6 signaling in the brain, or more specifically in the AP, could be

an effective treatment for cancer-associated cachexia.

References

1   Knupfer, H. & Preiss, R. Serum interleukin-6 levels in colorectal cancer patients--a
summary of published results. *Int J Colorectal Dis* **25**, 135-140, doi:10.1007/s00384-009-
0818-8 (2010).
2   Salgado, R. *et al.* Circulating interleukin-6 predicts survival in patients with metastatic
breast cancer. *Int J Cancer* **103**, 642-646, doi:10.1002/ijc.10833 (2003).
3   Kumari, N., Dwarakanath, B. S., Das, A. & Bhatt, A. N. Role of interleukin-6 in cancer
progression and therapeutic resistance. *Tumour Biol* **37**, 11553-11572,
doi:10.1007/s13277-016-5098-7 (2016).
4   Scott, H. R., McMillan, D. C., Crilly, A., McArdle, C. S. & Milroy, R. The relationship
between weight loss and interleukin 6 in non-small-cell lung cancer. *Br J Cancer* **73**,
1560-1562, doi:10.1038/bjc.1996.294 (1996).
5   Mantovani, G. *et al.* Cytokine activity in cancer-related anorexia/cachexia: role of
megestrol acetate and medroxyprogesterone acetate. *Semin Oncol* **25**, 45-52 (1998).
6   Ando, K. *et al.* Possible role for tocilizumab, an anti-interleukin-6 receptor antibody, in
treating cancer cachexia. *J Clin Oncol* **31**, e69-72, doi:10.1200/JCO.2012.44.2020 (2013).
7   Prado, B. L. & Qian, Y. Anti-cytokines in the treatment of cancer cachexia. *Ann Palliat*
Med **8**, 67-79, doi:10.21037/apm.2018.07.06 (2019).

8   Strassmann, G., Fong, M., Kenney, J. S. & Jacob, C. O. Evidence for the involvement of
interleukin 6 in experimental cancer cachexia. *J Clin Invest* **89**, 1681-1684,
doi:10.1172/JCI115767 (1992).
9   Baltgalvis, K. A. *et al.* Interleukin-6 and cachexia in ApcMin/+ mice. *Am J Physiol Regul*
Integr Comp Physiol **294**, R393-401, doi:10.1152/ajpregu.00716.2007 (2008).
10 Baltgalvis, K. A. *et al.* Muscle wasting and interleukin-6-induced atrogin-I expression in
the cachectic Apc (Min/+) mouse. *Pflugers Arch* **457**, 989-1001, doi:10.1007/s00424-
008-0574-6 (2009).
11 Zaki, M. H., Nemeth, J. A. & Trikha, M. CNTO 328, a monoclonal antibody to IL-6, inhibits
human tumor-induced cachexia in nude mice. *Int J Cancer* **111**, 592-595,
doi:10.1002/ijc.20270 (2004).
12 Fujita, J. *et al.* Anti-interleukin-6 receptor antibody prevents muscle atrophy in colon-26
adenocarcinoma-bearing mice with modulation of lysosomal and ATP-ubiquitin-
dependent proteolytic pathways. *Int J Cancer* **68**, 637-643, doi:10.1002/(SICI)1097-
0215(19961127)68:5<637::AID-IJC14>3.0.CO;2-Z (1996).
13 White, J. P. *et al.* The regulation of skeletal muscle protein turnover during the
progression of cancer cachexia in the Apc(Min/+) mouse. *PLoS One* **6**, e24650,
doi:10.1371/journal.pone.0024650 (2011).
14 Ando, K. *et al.* Tocilizumab, a proposed therapy for the cachexia of Interleukin6-
expressing lung cancer. *PLoS One* **9**, e102436, doi:10.1371/journal.pone.0102436
(2014).
15 Flint, T. R. *et al.* Tumor-Induced IL-6 Reprograms Host Metabolism to Suppress Anti-
tumor Immunity. *Cell Metab* **24**, 672-684, doi:10.1016/j.cmet.2016.10.010 (2016).
16 Grossberg, A. J., Scarlett, J. M. & Marks, D. L. Hypothalamic mechanisms in cachexia.
Physiol Behav **100**, 478-489, doi:10.1016/j.physbeh.2010.03.011 (2010).
17 Olson, B., Diba, P., Korzun, T. & Marks, D. L. Neural Mechanisms of Cancer Cachexia.
Cancers (Basel) **13**, doi:10.3390/cancers13163990 (2021).
18 Biswas, A. K. & Acharyya, S. Cancer-Associated Cachexia: A Systemic Consequence of
Cancer Progression. *Annual Review of Cancer Biology* **4**, 391–411 (2020).
19 Baracos, V. E., Martin, L., Korc, M., Guttridge, D. C. & Fearon, K. C. H. Cancer-associated
cachexia. *Nat Rev Dis Primers* **4**, 17105, doi:10.1038/nrdp.2017.105 (2018).
20 Fearon, K., Arends, J. & Baracos, V. Understanding the mechanisms and treatment
options in cancer cachexia. *Nat Rev Clin Oncol* **10**, 90-99,
doi:10.1038/nrclinonc.2012.209 (2013).
21 Tanaka, Y. *et al.* Experimental cancer cachexia induced by transplantable colon 26
adenocarcinoma in mice. *Cancer Res* **50**, 2290-2295 (1990).
22 Strassmann, G. & Kambayashi, T. Inhibition of experimental cancer cachexia by anti-
cytokine and anti-cytokine-receptor therapy. *Cytokines Mol Ther* **1**, 107-113 (1995).
23 Narsale, A. A. & Carson, J. A. Role of interleukin-6 in cachexia: therapeutic implications.
Curr Opin Support Palliat Care **8**, 321-327, doi:10.1097/SPC.0000000000000091 (2014).
24 Fearon, K. C., Glass, D. J. & Guttridge, D. C. Cancer cachexia: mediators, signaling, and
metabolic pathways. *Cell Metab* **16**, 153-166, doi:10.1016/j.cmet.2012.06.011 (2012).
25 Fearon, K. C. & Carter, D. C. Cancer cachexia. *Ann Surg* **208**, 1-5, doi:10.1097/00000658-
198807000-00001 (1988).

26 Braun, T. P. *et al.* Central nervous system inflammation induces muscle atrophy via
activation of the hypothalamic-pituitary-adrenal axis. *J Exp Med* **208**, 2449-2463,
doi:10.1084/jem.20111020 (2011).
27 Campos, C. A. *et al.* Cancer-induced anorexia and malaise are mediated by CGRP
neurons in the parabrachial nucleus. *Nat Neurosci* **20**, 934-942, doi:10.1038/nn.4574
(2017).
28 Lerner, L. *et al.* MAP3K11/GDF15 axis is a critical driver of cancer cachexia. *J Cachexia*
Sarcopenia Muscle **7**, 467-482, doi:10.1002/jcsm.12077 (2016).
29 Suriben, R. *et al.* Antibody-mediated inhibition of GDF15-GFRAL activity reverses cancer
cachexia in mice. *Nat Med* **26**, 1264-1270, doi:10.1038/s41591-020-0945-x (2020).
30 Borner, T. *et al.* Anorexia-cachexia syndrome in hepatoma tumour-bearing rats requires
the area postrema but not vagal afferents and is paralleled by increased MIC-1/GDF15. *J*
Cachexia Sarcopenia Muscle **8**, 417-427, doi:10.1002/jcsm.12169 (2017).
31 Borner, T., Liberini, C. G., Lutz, T. A. & Riediger, T. Brainstem GLP-1 signalling contributes
to cancer anorexia-cachexia syndrome in the rat. *Neuropharmacology* **131**, 282-290,
doi:10.1016/j.neuropharm.2017.12.024 (2018).
32 Navarro, I. B. K., Schraner, M. & Riediger, T. Brainstem prolactin-releasing peptide
contributes to cancer anorexia-cachexia syndrome in rats. *Neuropharmacology* **180**,
108289, doi:10.1016/j.neuropharm.2020.108289 (2020).
33 Mueller, T. C., Bachmann, J., Prokopchuk, O., Friess, H. & Martignoni, M. E. Molecular
pathways leading to loss of skeletal muscle mass in cancer cachexia--can findings from
animal models be translated to humans? *BMC Cancer* **16**, 75, doi:10.1186/s12885-016-
2121-8 (2016).
34 Tisdale, M. J. Molecular pathways leading to cancer cachexia. *Physiology (Bethesda)* **20**,
340-348, doi:10.1152/physiol.00019.2005 (2005).
35 Tan, B. H. *et al.* Identification of possible genetic polymorphisms involved in cancer
cachexia: a systematic review. *J Genet* **90**, 165-177, doi:10.1007/s12041-011-0027-4
(2011).
36 Stephens, N. A., Skipworth, R. J. & Fearon, K. C. Cachexia, survival and the acute phase
response. *Curr Opin Support Palliat Care* **2**, 267-274,
doi:10.1097/SPC.0b013e3283186be2 (2008).
37 Hirata, H. *et al.* Favorable responses to tocilizumab in two patients with cancer-related
cachexia. *J Pain Symptom Manage* **46**, e9-e13, doi:10.1016/j.jpainsymman.2013.01.009
(2013).
38 Paval, D. R. *et al.* A systematic review examining the relationship between cytokines and
cachexia in incurable cancer. *J Cachexia Sarcopenia Muscle* **13**, 824-838,
doi:10.1002/jcsm.12912 (2022).
39 Benrick, A. *et al.* Interleukin-6 gene knockout influences energy balance regulating
peptides in the hypothalamic paraventricular and supraoptic nuclei. *J Neuroendocrinol*
21, 620-628, doi:10.1111/j.1365-2826.2009.01879.x (2009).
40 Plata-Salaman, C. R. Anorexia induced by activators of the signal transducer gp 130.
Neuroreport **7**, 841-844, doi:10.1097/00001756-199602290-00038 (1996).

41   Bossola, M., Luciani, G., Giungi, S. & Tazza, L. Anorexia, fatigue, and plasma interleukin-6
levels in chronic hemodialysis patients. *Ren Fail* **32**, 1049-1054,
doi:10.3109/0886022X.2010.504910 (2010).
42   Rothwell, N. J. *et al.* Interleukin-6 is a centrally acting endogenous pyrogen in the rat.
Can J Physiol Pharmacol **69**, 1465-1469, doi:10.1139/y91-219 (1991).
43   Rothwell, N. J. & Hopkins, S. J. Cytokines and the nervous system II: Actions and
mechanisms of action. *Trends Neurosci* **18**, 130-136, doi:10.1016/0166-2236(95)93890-a
(1995).
44   Pavlov, V. A., Chavan, S. S. & Tracey, K. J. Molecular and Functional Neuroscience in
Immunity. *Annu Rev Immunol* **36**, 783-812, doi:10.1146/annurev-immunol-042617-
053158 (2018).
45   Banks, W. A., Kastin, A. J. & Broadwell, R. D. Passage of cytokines across the blood-brain
barrier. *Neuroimmunomodulation* **2**, 241-248, doi:10.1159/000097202 (1995).
46   Hopkins, S. J. & Rothwell, N. J. Cytokines and the nervous system. I: Expression and
recognition. *Trends Neurosci* **18**, 83-88 (1995).
47   Borison, H. L. Area postrema: chemoreceptor circumventricular organ of the medulla
oblongata. *Prog Neurobiol* **32**, 351-390, doi:10.1016/0301-0082(89)90028-2 (1989).
48   Miller, A. D. & Leslie, R. A. The area postrema and vomiting. *Front Neuroendocrinol* **15**,
301-320, doi:10.1006/frne.1994.1012 (1994).
49   Price, C. J., Hoyda, T. D. & Ferguson, A. V. The area postrema: a brain monitor and
integrator of systemic autonomic state. *Neuroscientist* **14**, 182-194,
doi:10.1177/1073858407311100 (2008).
50   Zhang, C. *et al.* Area Postrema Cell Types that Mediate Nausea-Associated Behaviors.
Neuron **109**, 461-472 e465, doi:10.1016/j.neuron.2020.11.010 (2021).
51   Clayton, D. F. The genomic action potential. *Neurobiol Learn Mem* **74**, 185-216,
doi:10.1006/nlme.2000.3967 (2000).
52   Guzowski, J. F. *et al.* Mapping behaviorally relevant neural circuits with immediate-early
gene expression. *Curr Opin Neurobiol* **15**, 599-606, doi:10.1016/j.conb.2005.08.018
(2005).
53   Dong, H. W., Petrovich, G. D. & Swanson, L. W. Topography of projections from
amygdala to bed nuclei of the stria terminalis. *Brain Res Brain Res Rev* **38**, 192-246,
doi:10.1016/s0165-0173(01)00079-0 (2001).
54   Dong, H. W., Petrovich, G. D., Watts, A. G. & Swanson, L. W. Basic organization of
projections from the oval and fusiform nuclei of the bed nuclei of the stria terminalis in
adult rat brain. *J Comp Neurol* **436**, 430-455, doi:10.1002/cne.1079 (2001).
55   Palmiter, R. D. The Parabrachial Nucleus: CGRP Neurons Function as a General Alarm.
Trends Neurosci **41**, 280-293, doi:10.1016/j.tins.2018.03.007 (2018).
56   Roman, C. W., Derkach, V. A. & Palmiter, R. D. Genetically and functionally defined NTS
to PBN brain circuits mediating anorexia. *Nat Commun* **7**, 11905,
doi:10.1038/ncomms11905 (2016).
57   Song, C. K., Schwartz, G. J. & Bartness, T. J. Anterograde transneuronal viral tract tracing
reveals central sensory circuits from white adipose tissue. *Am J Physiol Regul Integr
Comp Physiol* **296**, R501-511, doi:10.1152/ajpregu.90786.2008 (2009).

58   Duke, C. G. *et al.* An Improved CRISPR/dCas9 Interference Tool for Neuronal Gene
408   Suppression. *Front Genome Ed* **2**, 9, doi:10.3389/fgeed.2020.00009 (2020).
409 59   Yeo, N. C. *et al.* An enhanced CRISPR repressor for targeted mammalian gene regulation.
410   *Nat Methods* **15**, 611-616, doi:10.1038/s41592-018-0048-5 (2018).
411 60   Bonetto, A., Rupert, J. E., Barreto, R. & Zimmers, T. A. The Colon-26 Carcinoma Tumor-
412   bearing Mouse as a Model for the Study of Cancer Cachexia. *J Vis Exp*,
413   doi:10.3791/54893 (2016).
414 61   Xu, W. *et al.* Distinct neuronal coding schemes in memory revealed by selective erasure
415   of fast synchronous synaptic transmission. *Neuron* **73**, 990-1001,
416   doi:10.1016/j.neuron.2011.12.036 (2012).
417 62   Cardoso, F. *et al.* Neuro-mesenchymal units control ILC2 and obesity via a brain-adipose
418   circuit. *Nature* **597**, 410-414, doi:10.1038/s41586-021-03830-7 (2021).
419 63   Deuchars, S. A. & Lall, V. K. Sympathetic preganglionic neurons: properties and inputs.
420   *Compr Physiol* **5**, 829-869, doi:10.1002/cphy.c140020 (2015).
421 64   Koren, T. *et al.* Insular cortex neurons encode and retrieve specific immune responses.
422   *Cell* **184**, 5902-5915 e5917, doi:10.1016/j.cell.2021.10.013 (2021).
423 65   Luan, H. H. *et al.* GDF15 Is an Inflammation-Induced Central Mediator of Tissue
424   Tolerance. *Cell* **178**, 1231-1244 e1211, doi:10.1016/j.cell.2019.07.033 (2019).
425 66   Nandi, J. *et al.* Central mechanisms involved with catabolism. *Curr Opin Clin Nutr Metab*
426   *Care* **5**, 407-418, doi:10.1097/00075197-200207000-00010 (2002).
427 67   Sternson, S. M. & Eiselt, A. K. Three Pillars for the Neural Control of Appetite. *Annu Rev*
428   *Physiol* **79**, 401-423, doi:10.1146/annurev-physiol-021115-104948 (2017).
429 68   Zhang, X. *et al.* Brain control of humoral immune responses amenable to behavioural
430   modulation. *Nature* **581**, 204-208, doi:10.1038/s41586-020-2235-7 (2020).
431 69   Papazoglou, I. *et al.* A distinct hypothalamus-to-beta cell circuit modulates insulin
432   secretion. *Cell Metab* **34**, 285-298 e287, doi:10.1016/j.cmet.2021.12.020 (2022).
433 70   Xie, Z. *et al.* The gut-to-brain axis for toxin-induced defensive responses. *Cell* **185**, 4298-
434   4316 e4221, doi:10.1016/j.cell.2022.10.001 (2022).
435 71   Chen, J. Y., Campos, C. A., Jarvie, B. C. & Palmiter, R. D. Parabrachial CGRP Neurons
436   Establish and Sustain Aversive Taste Memories. *Neuron* **100**, 891-899 e895,
437   doi:10.1016/j.neuron.2018.09.032 (2018).
438 72   Sabatini, P. V. *et al.* GFRAL-expressing neurons suppress food intake via aversive
439   pathways. *Proc Natl Acad Sci U S A* **118**, doi:10.1073/pnas.2021357118 (2021).
440 73   Yoo, E. S. *et al.* Galpha(i/o)-coupled Htr2c in the paraventricular nucleus of the
441   hypothalamus antagonizes the anorectic effect of serotonin agents. *Cell Rep* **37**, 109997,
442   doi:10.1016/j.celrep.2021.109997 (2021).
443 74   Pei, H., Sutton, A. K., Burnett, K. H., Fuller, P. M. & Olson, D. P. AVP neurons in the
444   paraventricular nucleus of the hypothalamus regulate feeding. *Mol Metab* **3**, 209-215,
445   doi:10.1016/j.molmet.2013.12.006 (2014).
446 75   Morrison, S. F. RVLM and raphe differentially regulate sympathetic outflows to
447   splanchnic and brown adipose tissue. *Am J Physiol* **276**, R962-973,
448   doi:10.1152/ajpregu.1999.276.4.R962 (1999).
449 76   Emmerson, P. J. *et al.* The metabolic effects of GDF15 are mediated by the orphan
450   receptor GFRAL. *Nat Med* **23**, 1215-1219, doi:10.1038/nm.4393 (2017).

77   Hsu, J. Y. *et al.* Non-homeostatic body weight regulation through a brainstem-restricted
receptor for GDF15. *Nature* **550**, 255-259, doi:10.1038/nature24042 (2017).
78   Mullican, S. E. *et al.* GFRAL is the receptor for GDF15 and the ligand promotes weight
loss in mice and nonhuman primates. *Nat Med* **23**, 1150-1157, doi:10.1038/nm.4392
(2017).
79   Yang, L. *et al.* GFRAL is the receptor for GDF15 and is required for the anti-obesity
effects of the ligand. *Nat Med* **23**, 1158-1166, doi:10.1038/nm.4394 (2017).
80   Johnen, H. *et al.* Tumor-induced anorexia and weight loss are mediated by the TGF-beta
superfamily cytokine MIC-1. *Nat Med* **13**, 1333-1340, doi:10.1038/nm1677 (2007).
81   Breen, D. M. *et al.* GDF-15 Neutralization Alleviates Platinum-Based Chemotherapy-
Induced Emesis, Anorexia, and Weight Loss in Mice and Nonhuman Primates. *Cell Metab*
32, 938-950 e936, doi:10.1016/j.cmet.2020.10.023 (2020).
82   Altena, R. *et al.* Growth differentiation factor 15 (GDF-15) plasma levels increase during
bleomycin- and cisplatin-based treatment of testicular cancer patients and relate to
endothelial damage. *PLoS One* **10**, e0115372, doi:10.1371/journal.pone.0115372 (2015).
83   Tsai, V. W. W., Husaini, Y., Sainsbury, A., Brown, D. A. & Breit, S. N. The MIC-1/GDF15-
GFRAL Pathway in Energy Homeostasis: Implications for Obesity, Cachexia, and Other
Associated Diseases. *Cell Metab* **28**, 353-368, doi:10.1016/j.cmet.2018.07.018 (2018).
84   Lockhart, S. M., Saudek, V. & O'Rahilly, S. GDF15: A Hormone Conveying Somatic
Distress to the Brain. *Endocr Rev* **41**, doi:10.1210/endrev/bnaa007 (2020).
85   Heinrich, P. C. *et al.* Principles of interleukin (IL)-6-type cytokine signalling and its
regulation. *Biochem J* **374**, 1-20, doi:10.1042/BJ20030407 (2003).
86   Berti, A., Boccalatte, F., Sabbadini, M. G. & Dagna, L. Assessment of tocilizumab in the
treatment of cancer cachexia. *J Clin Oncol* **31**, 2970, doi:10.1200/JCO.2012.48.4147
(2013).
87   Kopf, M. *et al.* Impaired immune and acute-phase responses in interleukin-6-deficient
mice. *Nature* **368**, 339-342, doi:10.1038/368339a0 (1994).
88   Labun, K. *et al.* CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome
editing. *Nucleic Acids Res* **47**, W171-W174, doi:10.1093/nar/gkz365 (2019).
89   Savell, K. E. *et al.* A Neuron-Optimized CRISPR/dCas9 Activation System for Robust and
Specific Gene Regulation. *eNeuro* **6**, doi:10.1523/ENEURO.0495-18.2019 (2019).
90   Savell, K. E., Sultan, F. A. & Day, J. J. A Novel Dual Lentiviral CRISPR-based Transcriptional
Activation System for Gene Expression Regulation in Neurons. *Bio Protoc* **9**, e3348,
doi:10.21769/BioProtoc.3348 (2019).
91   Gordon, M. G. *et al.* lentiMPRA and MPRAflow for high-throughput functional
characterization of gene regulatory elements. *Nat Protoc* **15**, 2387-2412,
doi:10.1038/s41596-020-0333-5 (2020).
488

Methods

Mice

Male mice aged 2–4 months were used in all the experiments. Mice were housed under a 12-h

light/dark cycle (7 a.m. to 7 p.m. light) in groups of 2–5 animals, with a room temperature (RT)

of 22 °C and humidity of 50%. Food and water were available *ad libitum* before experiments. All

experimental procedures were approved by the Institutional Animal Care and Use Committee of

Cold Spring Harbor Laboratory and performed in accordance with the US National Institutes of

Health guidelines.

The Balbc mice (strain number: 000651) were purchased from Jackson laboratory. The *Gfral-*

p2a-Cre mice was generated by Stephen Liberles[50].

Immunohistochemistry

Immunohistochemistry experiments were performed following standard procedures. Briefly,

mice were anesthetized with Euthasol (0.2 ml; Virbac, Fort Worth, Texas, USA) and

transcardially perfused with 30 ml of PBS, followed by 30 ml of 4% paraformaldehyde (PFA) in

PBS. Brains were extracted and further fixed in 4% PFA overnight followed by cryoprotection in

a 30% PBS-buffered sucrose solution for 36 h at 4 °C. Coronal sections (50 μm in thickness)

were cut using a freezing microtome (Leica SM 2010R). Brain sections were first washed in PBS

(3×5 min), incubated in PBST (0.3% Triton X-100 in PBS) for 30 min at RT and then washed

with PBS (3×5 min). Next, sections were blocked in 5% normal goat serum in PBST for 30 min

at RT and then incubated with the primary antibody overnight at 4 °C. Sections were washed

with PBS (5×15 min) and incubated with the fluorescent secondary antibody at RT for 2 h. After

washing with PBS (5×15 min), sections were mounted onto slides with Fluoromount-G

(eBioscience, San Diego, California, USA). Images were taken using an LSM 780 laser-scanning

confocal microscope (Carl Zeiss, Oberkochen, Germany).

The primary antibodies and dilutions used in this study were: rabbit anti-Fos (1:500, Santa Cruz,

sc-52), mouse anti-FLAG (1:1000, Sigma-Aldrich, F1804), rabbit anti-mCherry (1:1,000;

Abcam, ab167453, GR3213077-3). The fluorophore-conjugated secondary antibodies and

dilutions used were Alexa Fluor 488 goat anti-rabbit IgG (H + L; 1:500; A-11008, Invitrogen),

Alexa Fluor 647 goat anti-rabbit IgG (H + L; 1:500; A-21244, Invitrogen), Alexa Fluor 594 goat

anti-mouse IgG (H + L; 1:500; A-11005, Invitrogen).

Retro-orbital injection of exogenous IL-6 and its detection in the brain

Biotinylated human IL-6 solution (Acrobiosystems, IL-6-H8218; 2 µg/ml dissolved in saline)

was injected into Balbc mice (100 µl per mouse) via retro-orbital injection. For retro-orbital

injection, briefly, the animal was anaesthetized with 2% isoflurane. A 27-gauge needle on a 0.5

mL insulin syringe was used for the injection. The animal was placed on its side on a heat pad.

The gauge needle was inserted at approximately a 30-45° angle to the eye, lateral to the medial

canthus, through the conjunctival membrane. There is a bit of resistance that causes the eye to

retreat back into the sinus until the needle pierces through the conjunctiva. The needle was

positioned behind the globe of the eye in the retro-bulbar sinus. The biotinylated human IL-6

solution was injected slowly into the retro-bulbar sinus. After the injection, the needle was

removed gently and the animal was returned to homecage for recovery. 3 hours after the

injection, the animals were sacrificed and transcardially perfused with 30 ml of PBS, followed by

30 ml of 4% paraformaldehyde (PFA) in PBS. Brains were extracted and further fixed in 4%

PFA overnight followed by cryoprotection in a 30% PBS-buffered sucrose solution for 36 h at

4 °C. Coronal sections (50 μm in thickness) were cut using a freezing microtome (Leica SM

2010R). Brain sections were incubated in Streptavidin solution (1:1000, ThermoFisher, Alexa

Fluor™ 647 conjugate, dissolved in 0.3% PBST) in room temperature for 2 hours. After washing

with PBS (5 × 15 min), sections were mounted onto slides with Fluoromount-G (eBioscience).

Images were taken using an LSM 780 laser-scanning confocal microscope (Carl Zeiss).

Fluorescence *in situ* hybridization

Single molecule fluorescent *in situ* hybridization (smFISH) (RNAscope, ACDBio) was used to

detect the expression of *Glp1r*, *Il6ra*, *Fos*, and *Gfral* mRNAs in the area postrema of Balbc mice.

For tissue preparation, mice were first anesthetized under isoflurane and then decapitated. Their

brain tissue was first embedded in cryomolds (Sakura Finetek, Catalog number 4566) filled with

M-1 Embedding Matrix (Thermo Scientific, Catalog number 1310) then quickly fresh-frozen on

dry ice. The tissue was stored at −80 °C until it was sectioned with a cryostat. Cryostat-cut

sections (16-μm thick) containing the entire area postrema were collected through the rostro-

caudal axis in a series of four slides, and quickly stored at −80 °C until processed. Hybridization

was carried out using the RNAscope kit (ACDBio). On the day of the experiment, frozen

sections were postfixed in 4% PFA in RNA-free PBS (hereafter referred to as PBS) at RT for

15 min, then washed in PBS, dehydrated using increasing concentrations of ethanol in water

(50%, once; 70%, once; 100%, twice; 5 min each). Sections were then dried at RT and incubated

with Protease IV for 30 min at RT. Sections were washed in PBS three times (5 min each) at RT,

then hybridized. Probes against *Glp1r* (Catalog number 418851-C3, dilution 1:50), *Il6ra*

(Catalog number 438931-O1, dilution 1:50), *Fos* (Catalog number 316921-C2, dilution 1:50),

and *Gfral* (Catalog number 417021-C2, dilution 1:50) were applied to the area postrema

sections. Hybridization was carried out for 2 h at 40 °C. After that, sections were washed twice in

1× Wash Buffer (Catalog number 310091; 2 min each) at RT, then incubated with the

amplification reagents for three consecutive rounds (30 min, 15 min and 30 min, at 40 °C). After

each amplification step, sections were washed twice in 1× Wash Buffer (2 min each) at RT.

Finally, fluorescence detection was carried out for 15 min at 40 °C. Sections were then washed

twice in 1× Wash Buffer (2 min each), incubated with DAPI for 2 min, washed twice in 1× Wash

Buffer (2 min each), then mounted with a coverslip using mounting medium. Images were

acquired using an LSM780 confocal microscope equipped with 20x and 40x lenses, and

visualized and processed using ImageJ and Adobe Illustrator. Cell counting and mean

fluorescence intensity quantification of the images were performed with ImageJ.

Single guide RNA (sgRNA) design and lentiviral production for CRISPR/dCas9

interference

sgRNAs targeting the *Il6ra* transcription start site (TSS) were designed using CHOPCHOP[88].

Seven *Il6ra* sgRNAs (sgRNA-1 to sgRNA-7) as well as a sgRNA targeting the *lacZ* promoter

(*LacZ* sgRNA) were cloned into the Lenti U6-sgRNA/Ef1a-mCherry plasmid (Addgene

#114199), as described previously[89,90]. The eight sgRNA plasmids, Lenti SYN-FLAG-dCas9-

KRAB-MeCP2 plasmid (Addgene #155365), and the two helper plasmids pCMV-VSV-G

(Addgene #8454) and psPAX2 (Addgene #12260) were purified with the NucleoBond Xtra Midi

EF kit (Takara 740420). *Il6ra* knockdown efficiency was assessed by transient transfection of

sgRNA and dCas9-KRAB-MeCP2 into the mHypoA hypothalamic neural cell line (Cedarlane

Labs, clone clu-175). 3 μg of each sgRNA plasmid and 3 μg of dCas9-KRAB-MeCP2 plasmid

were co-nucleofected into 1 x 10^6 mHypoA cells resuspended in a 1:1 mixture of Ingenio

Electroporation reagent (Mirus Bio 50111) and OptiMEM (Gibco 31985062), using program A-

033 on the Nucleofector 2b (Lonza). The cells were harvested 60 hours post-nucleofection.

DAPI- and mCherry-positive cells were collected by FACS. The *Il6ra* mRNA was extracted and

the knockdown efficiency was measured by RT-qPCR. The two most effective sgRNAs, sgRNA-

4 (-23 to -41 of TSS) and sgRNA-6 (-163 to -182 of TSS), resulting in 67% and 35% knockdown

of *Il6ra* expression in mHypoA cells, respectively, were used for *in vivo* experiments. FLAG-

dCas9-KRAB-MeCP2, *Il6ra* sgRNA-4, *Il6ra* sgRNA-6, and *lacZ* sgRNA lentiviruses were

produced in HEK293T cells. Lentiviral pellets were resuspended in 30 uL DPBS, aliquoted and

flash-frozen on dry ice, and stored at -80°C. Physical and functional titers were obtained using

the Lenti-X qRT-PCR Titration Kit (Takara 631235) and qPCR of genomic DNA following

HEK293T transduction[91], respectively.

Lentiviral preparation	Physical titer (genomic copies/uL)	Functional titer (transducing units/uL)
FLAG-dCas9-KRAB-MeCP2 (prep 1)	3.0E+11	1.2E+07
FLAG-dCas9-KRAB-MeCP2 (prep 2)	8.0E+10	1.1E+07
Il6ra sgRNA-4	3.0E+12	1.4E+08
Il6ra sgRNA-6	5.0E+11	3.8E+07
lacZ sgRNA	3.0E+12	8.8E+07

Plasmids for lentiviral production:

lenti SYN-FLAG-dCas9-KRAB-MeCP2 was a gift from Jeremy Day (Addgene plasmid #

155365; http://n2t.net/addgene:155365 ; RRID:Addgene_155365)

lenti U6-sgRNA/EF1a-mCherry was a gift from Jeremy Day (Addgene plasmid # 114199;

http://n2t.net/addgene:114199 ; RRID:Addgene_114199)

pCMV-VSV-G was a gift from Bob Weinberg (Addgene plasmid # 8454;

http://n2t.net/addgene:8454 ; RRID:Addgene_8454)

psPAX2 was a gift from Didier Trono (Addgene plasmid # 12260; http://n2t.net/addgene:12260;

RRID:Addgene_12260)

Adeno-associated viruses (AAVs)

The AAV-CMV-DIO-EGFP-2A-TeLC vector was a gift from Dr. Wei Xu at UT Southwestern.

A custom virus (AAV-DJ) based on this vector was produced by WZ Biosciences Inc (Rockville,

MD 20855). pAAV-hSyn-DIO-EGFP was purchased from Addgene (Watertown, MA 02472,

USA). All viruses were aliquoted and stored at −80 °C until use.

Stereotaxic surgery

All surgery was performed under aseptic conditions and body temperature was maintained with a

heating pad. Standard surgical procedures were used for stereotaxic injection. Briefly, mice were

anesthetized with isoflurane (3% at the beginning and 1% for the rest of the surgical procedure),

and were positioned in a stereotaxic injection frame and on top of a heating pad maintained at

35 °C. A digital mouse brain atlas was linked to the injection frame to guide the identification

and targeting of different brain areas (Angle Two Stereotaxic System, http://myNeuroLab.com).

We used the following coordinates for injections in the area postrema: -7.65 mm from bregma,

0 mm lateral from the midline, and 4.7 mm vertical from the skull surface. For virus injection, we

made a small cranial window (1–2 mm^2) for each mouse, through which a glass micropipette (tip

diameter, ~5 μm) containing viral solution was lowered down to the target. For AAVs, about

0.3 μl of viral solution was injected. For lentiviruses, 0.2-0.3 μl of viral mixture (the dCas9 and

the sgRNA viruses were mixed at a volume:volume ratio of 2:1) was injected. Viral solutions

were delivered with pressure applications (5–20 psi, 5–20 ms at 1 Hz) controlled by a

Picospritzer III (General Valve) and a pulse generator (Agilent). The speed of injection was

~0.1 μl/10 min. We waited for at least 10 min following the injection before slowly removing the

injection pipette. After injection, the incision was sealed by surgical sutures and the animal was

returned to homecage for recovery.

Colon-26 (C26) adenocarcinoma cells

C26 cells were cultured in complete growth medium consisting of RPMI 1640 medium with

Glutamine (#11-875-093; Thermo Fisher) containing 10% of heat-inactivated Fetal Bovine

Serum (FBS) (#10-438-026; Thermo Fisher) and 1x Penicillin-Streptomycin solution (#15-140-

122; Thermo Fisher) under sterile conditions. 1x Trypsin-EDTA (#15400054; Thermo Fisher)

was used for cell dissociation. Cells were resuspended in FBS-free RPMI and viable cells were

counted using a Vi-Cell counter prior to subcutaneous injection of 2×10^6 viable cells diluted in

100 μL RPMI into the right flank of each BALB/c mouse.

Lewis lung carcinoma (LLC) cells

LL/2 (LLC1) cells were obtained from ATCC (American Type Culture Collection; #CRL-1642)

and cultured in Dulbecco's Modified Eagle's Medium (DMEM) (#30-2002; ATCC) complete

growth medium, with 10% of heat-inactivated FBS (#10-438-026; Thermo Fisher) and 1x

Penicillin-Streptomycin solution (#15-140-122; Thermo Fisher) under sterile conditions. 1x

Trypsin-EDTA (#15400054; Thermo Fisher) was used for cell dissociation. Cells were

resuspended in FBS-free DMEM and viable cells were counted using a Vi-Cell counter prior to

subcutaneous injection of. 2×10^6 viable cells were diluted in 100 μL FBS-free DMEM and were

subcutaneous injected into the right flank of each C57BL/6 mouse.

Intracerebroventricular (i.c.v.) infusion of anti-IL-6

On day 10 or 12 post-C26 injection, an osmotic device of 200 uL volume and a release rate of

0.5 uL/hour consisting of a cannula, connecting line, metal flow moderator and pump (#AP-

2001; Alzet) was placed in a subcutaneous pocket and stereotactically implanted into the right

lateral ventricle of the C26-tumour bearing BALB/c mice for a period of 14 days. Prior to use,

the infusion device was assembled and equilibrated in saline overnight. The pump was filled

with either the InVivoMAb rat anti-mouse IL-6 (clone MP5-20F3, #BE0046; BioXCell) or an

InVivoMAb rat IgG1 isotype control (anti-HRPN, #BE0088; BioXCell). Both antibodies were

diluted in PBS to achieve continuous infusion of a 5 mg/mL dose. Pump replacement surgery

was performed after 14 days. The coordinate for targeting the lateral ventricle was -0.5 mm from

bregma, 1.25 mm lateral from the midline, and 2.5 mm vertical from the skull surface.

Measuring bodyweight, food intake, and water intake

Food and water intake monitoring cages (BioDAQ Unplugged, Research Diets, Inc., New

Brunswick, NJ 08901 USA) were used to measure the food intake and water intake of the

animals. Mice were singly housed in these cages. Food and water were placed in an extended

hopper which can be reached by the animal. The bodyweight of the animal, weight of the food

and water in the hopper were measured daily at 4 pm. The cachectic mice which lost >20% of

bodyweight were sacrificed and the tissues were collected for further analysis. Blood glucose

concentrations were measured from whole venous blood using an automatic glucose monitor

(Bayer HealthCare Ascensia Contour).

Blood and plasma measurements

Blood glucose concentrations were measured from whole venous blood using an automatic glucose

monitor (Bayer HealthCare Ascensia Contour). Tail vein bleeding was performed using a scalpel

via tail venesection without restraint. Blood samples were collected from tail bleed using heparin-

coated hematocrit capillary tubes to avoid coagulation. Samples were then centrifuged at 14,000

rpm for 5 min at 4°C. Plasma was collected in a new tube, snap frozen in liquid nitrogen and stored

at -80°C. IL-6 and GDF-15 levels were measured in plasma using the mouse IL-6 Quantikine

ELISA Kit (#M6000B; R&D Systems) and the Mouse/Rat GDF-15 Quantikine ELISA Kit

(#MGD150; R&D) respectively.

Brain tissue lysis and IL-6 quantification

Mice were transcardially perfused with saline and the area postrema was collected, snap frozen in

liquid nitrogen and stored at -80°C until further analysis. Tissue was placed into 2-mL round-

bottom homogenizer tubes pre-loaded with Stainless Steel beads (#69989; Qiagen) and filled up

with lysis buffer (#AA-LYS-16ml; RayBiotech) supplemented with Protease Inhibitor Cocktail

(#AA-PI; Raybiotech) and Phosphatase Inhibitor Cocktail Set I (#AA-PHI-I; RayBiotech).

Samples were homogenized in Tissue Lyser II (#85300; Qiagen) for 5 minutes and then lysates

were centrifuged at 4°C for 20 minutes at maximum speed. The supernatant was harvested and

kept on ice if testing fresh or sored at -80°C. The Bicinchoninic Acid (BCA) Method was used to

determine protein concentration in lysates. IL-6 levels were quantified in the lysates using a Mouse

IL-6 ELISA specific for lysates (#ELM-IL6-CL-1; RayBiotech).

In vitro **electrophysiology**

Acute slices were obtained from two- to three-month-old mice. Mice were anaesthetized with

isoflurane (4%) before rapid decapitation. The brain was rapidly removed, and coronal slices

(300 μm) containing the AP were cut using a HM650 Vibrating-blade Microtome (Thermo

Fisher Scientific). Slices were cut in ice-cold dissection buffer (110.0 mM Choline chloride, 25.0

mM $NaHCO_3$, 1.25 mM NaH_2PO_4, 2.5 mM KCl, 25.0 mM glucose, 0.5 mM $CaCl_2$, 7.0 mM

$MgCl_2$, 11.6 mM ascorbic acid, and 3.1 mM pyruvic acid, and bubbled with 95% O_2 and 5%

CO_2) and subsequently transferred to a recovery chamber containing artificial cerebrospinal fluid

(ACSF) solution (containing 118 mM NaCl, 2.5 mM KCl, 26.2 mM $NaHCO_3$, 1 mM NaH_2PO_4,

20 mM Glucose, 2 mM $CaCl_2$ and 2 mM $MgCl_2$, pH 7.4, and saturated with 95% O_2 and 5%

CO_2) at 34°C. The slices were maintained at 34°C for at least 40 minutes and subsequently at

room temperature (20-24°C). Recordings were made in a continuously flow of ACSF and

bubbled with 95% O_2/5% CO_2.

Whole-cell patch-clamp recordings were obtained with Multiclamp 700B amplifiers and

pCLAMP 10 software (Molecular Devices; Sunnyvale, California, USA) and was guided using

an Olympus BX51 Microscope equipped with both transmitted and epifluorescence light sources

(Olympus Corporation, Shinjuku, Tokyo, Japan).

Synaptic responses were recorded at holding potentials of -70mV (for AMPA receptor-mediated

responses), and 0 mV (for GABAA receptor-mediated responses) and were low-pass filtered at 1

kHz. The internal solution for voltage-clamp experiments contained 115 mM Cesium

methanesulfonate, 20 mM CsCl, 10 mM HEPES, 2.5 mM $MgCl_2$, 4 mM Na_2-ATP, 0.4 mM Na_3-

GTP, 10 mM Na-phosphocreatine, and 0.6 mM EGTA, pH 7.2. Miniature EPSCs were recorded

in the presence of tetrodotoxin (1 μM) and picrotoxin (100 μM). Spontaneous IPSCs were

recorded in the presence of AP-5 (100 μM) and CNQX (5 μM). The EPSCs and IPSCs were

analyzed using Mini Analysis software (Synaptosoft).

Statistical Analysis

All statistics are indicated where used. Statistical analyses were conducted using GraphPad

Prism version 6.0 (GraphPad Software, Inc., La Jolla, CA). Statistical comparisons were

performed using Student's t test or ANOVA. All comparisons were two tailed. Statistic

hypothesis testing was conducted at a significance level of 0.05.

Data availability

The authors declare that the data supporting the findings of this study are available within the

paper and its supplementary information files.

Acknowledgements

We thank Dr. Stephen D. Liberles for sharing the *Gfral-p2a-Cre* mice. We thank Dr. Roberto

Malinow and Dr. Chuchu Zhang for comments on earlier versions of the manuscript. We thank

Radhashree Sharma for technical assistance and members of the Li laboratory for helpful

discussions. This work was supported by The Mark Foundation for Cancer Research (33300111,

T.J.), developmental funds from CSHL Cancer Center Support Grant (5P30CA045508, T.J.),

National Institutes of Health (NIH) (R01MH101214, R01MH108924, R01NS104944,

R01DA050374, B.L.), and the Cold Spring Harbor Laboratory and Northwell Health Affiliation

(B.L.).

Author Contributions

Q.S., D.V.D.L., M.F., B.G, M.W., J.T., T.J. and B.L. designed research; Q.S., D.V.D.L., M.F.,

B.G, and M.W. performed research; Q.S., D.V.D.L., M.F., B.G, M.W., J.T., T.J. and B.L.

analyzed data; and Q.S. and B.L. wrote the paper with inputs from all authors.

Competing interests

The authors declare no competing interests.

Correspondence and requests for materials should be addressed to B.L. (bli@cshl.edu).

Area postrema neurons mediate interleukin-6 function in cancer-associated cachexia

Qingtao Sun[1, #], Daniëlle van de Lisdonk[1,2, #], Miriam Ferrer[1,#], Bruno Gegenhuber[1], Melody Wu[1], Jessica Tollkuhn[1], Tobias Janowitz[1], Bo Li[1*]

FIGURES and SUPPLEMENTARY FIGURES

Figure 1 | Circulating IL-6 can reach the area postrema (AP) and activate AP neurons. a. A schematic of the approach. **b.** Confocal images showing the binding of the exogenous biotinylated IL-6 to cells in the AP. **c.** Quantification of the fluorescence signals from

fluorescence-conjugated avidin in the AP, which recognizes the biotinylated exogenous IL-6 (n = 3 mice for each group, t = 24.71, ****P = 0.000016, unpaired t test). **d.** A diagram showing the position of the AP in a coronal brain section. **e.** Confocal images showing Fos expression in the AP. **f.** Quantification of Fos-expressing (Fos$^+$) cells in the AP (n = 3 mice for each group, t = 9.246, ***P = 0.0008, unpaired t test). **g.** Confocal images showing the expression of different genes in AP cells, which was detected with single molecule fluorescent *in situ* hybridization (smFISH). At the bottom are higher magnification images of the boxed area in the overlay image on the top. Arrowheads indicate a neuron that expresses all three genes. **h.** A Venn diagram showing the relationships among cells expressing *Il6ra*, *Gfral*, and *Glp1r* in the AP. **i.** Characterization of the types of *Fos$^+$* cells in the AP by smFISH. At the bottom are higher magnification images of the boxed areas in images on the top. **j.** A Venn diagram showing the relationships among cells expressing *Fos*, *Il6ra*, and *Glp1r* in the AP.

Figure 2 | C26 cancer causes increased IL-6 in the AP and AP neuron hyperactivity. a. A schematic of the experimental procedure. **b.** IL-6 levels in the area postrema (AP) during cancer progression. IL-6 levels were normalized to total protein levels (n = 4-5 mice in each group, F =

2.9, P = 0.0445, *P < 0.05; One-way ANOVA followed by Tukey's post-hoc test). **c.** Confocal images showing Fos expression in different brain areas in tumor-bearing (top) or control (bottom) mice. **d.** Quantification of Fos$^+$ cells in different brain areas (n = 4 mice in each group; AP, t = 2.91, *P = 0.0268, PBN, t = 11.65, ****P = 2.41 × 10^{-5}, CeA, t = 4.37, **P = 0.0047, PVN, t = 3.65, *P = 0.011, BNST, t = 5.86, **P = 0.0011; unpaired t test). **e.** A diagram showing the AP in a coronal brain section for electrophysiological recording. **f.** Representative miniature EPSC traces from AP neurons in control (top) and cachectic (bottom) mice. **g.** Quantification of miniature EPSC frequency (left) and amplitude (right) (control, n = 30 cells / 6 mice, cachexia, n = 32 cells / 7 mice; frequency, n.s. (nonsignificant), P = 0.2513, amplitude, **P = 0.0017, Mann-Whitney test). **h.** Representative spontaneous IPSC traces from AP neurons in control (top) and cachectic (bottom) mice. **i.** Quantification of spontaneous IPSC frequency (left) and amplitude (right) (control, n = 36 cells / 7 mice, cachexia, n = 34 cells / 6 mice; frequency, n.s., P = 0.1637, amplitude, n.s., P = 0.4580, Mann-Whitney test).

Figure 3 | Intracerebroventricular (i.c.v.) infusion of IL-6 antibody prevents cachexia in the C26 cancer model. a. A schematic of the experimental procedure. **b.** A confocal image of a coronal brain section from a representative mouse, showing the location of the infusion cannula above the lateral ventricle (VL). **c.** Survival curves of the mice after tumor inoculation (IL-6

antibody group, n = 10, isotype control antibody group, n = 6; P = 0.0003, Mantel-Cox test). **d.** Bodyweight of individual mice relative to their bodyweight on the day of tumor inoculation. **e.** Average bodyweight normalized to that on day -5 (IL-6 antibody, n = 10, isotype control antibody, n = 6; $F(1,84) = 75.13$, $P = 2.77 \times 10^{-13}$; day -1, ****$P = 1.2 \times 10^{-6}$, day 0, ****$P < 1 \times 10^{-15}$; two-way repeated-measures (RM) ANOVA followed by Sidak's post hoc test). **f.** Cumulative food (left) and water (right) intake of the mice in the 5 days before sacrifice (IL-6 antibody, n = 10, isotype control antibody, n = 6 mice; food: $F(1,84) = 70.87$, $P = 8.78 \times 10^{-13}$, day -2, **$P = 0.007$, day -1, ****$P = 2 \times 10^{-6}$, day 0, ****$P = 8 \times 10^{-11}$; water: $F(1,84) = 106.6$, $P < 1 \times 10^{-15}$, day -2, **$P = 0.002$, day -1, ****$P = 3.6 \times 10^{-10}$, day 0, ****$P < 1 \times 10^{-15}$; two-way RM ANOVA with Sidak's post hoc test). **g.** Confocal images showing Fos expression in different brain areas in the mice infused with the IL-6 antibody (top) and the control antibody (bottom). **h.** Quantification of Fos$^+$ cells in different brain areas (IL-6 antibody, n = 9 mice, isotype control antibody, n = 6 mice; AP, $t = 6.44$, ****$P = 2.2 \times 10^{-5}$, PBN, $t = 3.43$, **$P = 0.0045$, CeA, $t = 1.49$, $P = 0.16$, PVN, $t = 8.89$, ****$P = 6.95 \times 10^{-7}$, BNST, $t = 4.309$, *** $P = 0.0008$; unpaired t test).

Figure 4 | Suppression of *Il6ra* expression in AP neurons ameliorates cachexia in the C26 cancer model. a. A schematic of the experimental procedure. **b.** Confocal immunohistochemical images of a coronal brain section from a representative mouse, showing the infection of AP cells with lentiviruses expressing the sgRNA (tagged with mCherry) and dCas9-KRAB-MeCP2 (tagged with FLAG). mCherry and FLAG were recognized by antibodies. **c.** Confocal images

showing the expression of *Gfral*, *Il6ra*, and *mCherry* in AP cells detected with smFISH. *mCherry* marks the cells infected with the CRISPR/dCas9 system containing the *lacZ* sgRNA (top) or *Il6ra* sgRNA-4 (bottom). **d.** Quantification of the fractions of $Gfral^+$ neurons expressing *Il6ra* in the AP of individual mice (*lacZ* sgRNA group, n = 7, *Il6ra* sgRNA-4 group, n = 10; t = 12.42, ****P = 2.7×10^{-9}, t test). **e.** Survival curves of the mice after tumor inoculation (*lacZ* sgRNA group, n = 8, *Il6ra* sgRNA-4 group, n = 7; P < 0.0001, Mantel-Cox test). **f.** Relationship between survival days and the fractions of $Gfral^+$ neurons expressing *Il6ra* (n = 17 mice, R^2 = 0.806, P < 0.0001 by a linear regression). **g.** Bodyweight of individual mice relative to their bodyweight on the day of tumor inoculation. **h.** Average bodyweight normalized to that on day -5 (*lacZ* sgRNA group, n = 8, *Il6ra* sgRNA-4 group, n = 7; F(1,78) = 23.63, P = 9.6×10^{-6}, *P = 0.012, ****P = 0.00004, two-way repeated-measures (RM) ANOVA with Sidak's post hoc test). **i.** Cumulative food (left) and water (right) intake of the mice after tumor inoculation (food: F(1,130) = 253.6, P < 1×10^{-15}; day 4, ***P = 0.0007, day 5, ****P = 9.2×10^{-6}, day 6, ****P = 5.5×10^{-7}, day 7, ****P = 3.6×10^{-8}, day 8, ****P = 4.23×10^{-9}, day 9, ****P = 9.21×10^{-13}, day 10, ****P = 2.4×10^{-14}; water: F(1,130) = 125.2, P < 1×10^{-15}; day 5, **P = 0.009, day 6, **P = 0.0019, day 7, ***P = 0.00024, day 8, ****P = 0.00003, day 9, ****P = 2.65×10^{-7}, day 10, ****P = 3.58×10^{-9}; two-way RM ANOVA with Sidak's post hoc test). **j.** Cumulative food (left) and water (right) intake of the mice in the 5 days before sacrifice (food: F(1, 65) = 24.83, P = 4.9×10^{-6}; *P = 0.033, **P = 0.0034; water: F(1, 65) = 31.32, P = 4.7×10^{-7}; **P = 0.0064, ***P = 0.0006; two-way RM ANOVA with Sidak's post hoc test).

Figure 5 | Inhibition of *Gfral*+ AP neurons ameliorates cachexia in the Lewis lung cancer (LLC) model. a. Plasma IL-6 (left) and GDF-15 (right) concentrations during cancer progression (IL-6: control, 7 mice at all timepoints except day 2 and day 8, where there are 5 mice; LLC, 8 mice at all timepoints except day 4 where there are 6 mice, day 6 and day 12 where there are 11 mice, and day 10 and day 18 where there are 7 mice; *P < 0.05; GDF-15: control, 5 mice at all timepoints except day 0 where there are 8 mice, and day 14 where there are 3 mice; LLC, 5 mice at all timepoints except day 0 where there are 7 mice; *P < 0.05, **P < 0.01;

multiple unpaired t-tests at each timepoint with false discovery rate adjusted with the two-stage step-up method). **b.** A schematic of the experimental procedure. **c.** Confocal immunohistochemical images of coronal brain sections from two representative mice, showing the infection of *Gfral*[+] AP neurons with an AAV expressing TeLC (left) or GFP only (right). **d.** Cumulative food (left) and water (right) intake of the mice in the 7 days before sacrifice (TeLC group, n = 9, GFP group, n = 8; food: $F(1,120) = 78.04$, $P = 10^{-14}$; day -4, *$P = 0.03$, day -3, ***$P = 0.0002$, day -2, ****$P = 0.00002$, day -1, ****$P = 5 \times 10^{-7}$, day 0, ****$P = 5.8 \times 10^{-8}$; water: $F(1,120) = 1.256$, $P = 0.26$; two-way repeated-measures ANOVA with Sidak's post hoc test. **e.** Inguinal fat (left) and quadriceps muscle (right) mass of the mice at 23 days after tumor inoculation (TeLC group, n = 9, GFP group, n = 8; fat: $t = 2.3$, *$P = 0.0363$; muscle: $t = 4.04$, **$P = 0.001$; t test). **f.** Confocal images showing Fos expression in different brain areas in the mice where *Gfral*[+] AP neurons were infected with the AAV expressing TeLC (top) or GFP only (bottom). **g.** Quantification of Fos[+] cells in different brain areas (TeLC group, n = 9 mice, GFP group, n = 8 mice; AP, $t = 2.33$, *$P = 0.0343$; PBN, $t = 4.04$, **$P = 0.0011$; PVN, $t = 3.09$, **$P = 0.0094$; CeA, $t = 2.62$, *$P = 0.019$; BNST, $t = 2.55$, *$P = 0.022$; unpaired t test). **h.** Tumor (left) and spleen (right) mass of the mice at 23 days after tumor inoculation (TeLC group, n = 9, GFP group, n = 8; tumor: $t = 1.34$, $P = 0.2$; spleen: $t = 0.65$, $P = 0.525$; t test).

Extended Data Figure 1 | Circulating IL-6 does not reach brain areas other than the AP. a. Confocal images showing the lack of exogenous IL-6 signals in different brain areas of mice received biotinylated IL-6 (left) compared with mice received saline (right) via retro-orbital injection. **b.** Quantification of the fluorescence signals from fluorescence-conjugated avidin in different brain areas, which recognizes the biotinylated exogenous IL-6 (n = 3 mice for each group; LS, t = 0.765, P = 0.487; BNST, t = 0.282, P = 0.792; PVN, t = 1.277, P = 0.271; CeA, t = 0.892, P = 0.423; t test).

Extended Data Figure 2 | Intracerebroventricular (i.c.v.) infusion of IL-6 antibody improves the physiological conditions of mice despite tumor growth in the C26 cancer model. a. IL-6 levels in the plasma (left) and cerebrospinal fluid (CSF; right). Plasma IL-6: anti-IL-6 group, n = 9 mice, isotype control antibody group, n = 8 mice; t = 1.664, P = 0.1168; CSF IL-6: anti-IL-6 group, n = 8 mice, isotype control antibody group, n = 7 mice; t = 2.002, P = 0.0666; t test. **b.** Inguinal fat mass (anti-IL-6 group, n = 10, isotype control antibody group, n = 6; t = 2.334, *P = 0.035, t test). **c.** Quadriceps muscle mass (anti-IL-6 group, n = 10, isotype control antibody group, n = 6; t = 2.538, *P = 0.0237, t test). **d.** Blood glucose levels at the endpoint (anti-IL-6 group, n = 9 mice, isotype control antibody group, n = 6 mice; t = 3.555, **P = 0.0035, t test). **e.** Tumor mass (anti-IL-6 group, n = 10, isotype control antibody group, n = 6; t = 2.864, *P = 0.0125, t test). **f.** Spleen mass (anti-IL-6 group, n = 10, isotype control antibody group, n = 6; t = 3.274, **P = 0.0055, t test).

Extended Data Figure 3 | Design and characterization of sgRNAs for the CRISPR/dCas9 system to suppress *Il6ra* expression. a. Transcription start site (TSS) targeting positions of the different sgRNAs. Cyan, template strand; red, non-template strand. **b.** *In vitro* characterization of the efficacy of the CRISPR/dCas9 system with different sgRNAs. Plasmids expressing dCas9-KRAB-MeCP2 and each of the sgRNAs were co-transfected into mHypoA cell line. 60 hours after the transfection, the expression of *Il6ra* in these cells was measured by qPCR (*Il6ra* sgRNA-4, n = 6 plates, *lacZ* sgRNA, n = 6 plates, P = 0.0005785; *Il6ra* sgRNA-6, n = 3 plates, *lacZ* sgRNA, n = 3 plates, P = 0.1965; t test).

Extended Data Figure 4 | Suppression of *Il6ra* expression in AP neurons improves the physiological conditions of mice despite tumor growth in the C26 cancer model. *Il6ra* sgRNA-4 group, n = 7 mice, *lacZ* sgRNA group, n = 8 mice. **a.** Blood glucose levels at different time points ($F_{(1,49)}$ = 9.834, P = 0.0029; two-way repeated-measures ANOVA with Sidak's post hoc test). **b.** Inguinal fat mass (t = 1.657, P = 0.12, t test). **c.** Quadriceps muscle mass (t = 1.92, P = 0.078, t test). **d.** Tumor mass (t = 7.32, ****P = 5.86 × 10^{-6}, t test). **e.** Spleen mass (t = 2.71, P = 0.018, t test).

Extended Data Figure 5 | Suppression of *Il6ra* expression in AP neurons ameliorates cachexia and reduces the hyperactivity in the AP network in the C26 cancer model. *lacZ* sgRNA group, n = 4 mice, *Il6ra* sgRNA-4 group, n = 4 mice. **a.** A schematic of the experimental

procedure. When one animal in the *lacZ* sgRNA (control) group became cachectic, that animal and a randomly selected animal in the *Il6ra* sgRNA-4 group were sacrificed to check Fos expression and other phenotypes. **b.** Average bodyweight normalized to that on day -5 ($F_{(1,36)}$ = 10.45, P = 0.0026, **P = 0.0072, two-way repeated-measures (RM) ANOVA with Sidak's post hoc test). **c.** Inguinal fat mass (t = 3.941, **P = 0.0079, t test). **d.** Quadriceps muscle mass (t = 5.26, **P = 0.0019, t test). **e, f.** Cumulative food (e) and water (f) intake of the mice in the 5 days before sacrifice (food: $F_{(1, 36)}$ = 3.239, P = 0.0803; water: $F_{(1, 36)}$ = 0.3045, P = 0.5845. two-way RM ANOVA with Sidak's post hoc test). **g, h.** Tumor (g) and spleen (h) mass of the mice (tumor: t = 0.444, P = 0.6725; spleen: t = 0.898, P = 0.404; t test). **i.** Confocal images showing Fos expression in different brain areas in representative mice of the two groups. **j.** Quantification of Fos$^+$ cells in different brain areas (AP, t = 4.617, **P = 0.0036; PBN, t = 2.849, *P = 0.029; PVN, t = 5.332, **P = 0.0018; CeA, t = 0.5267, P = 0.617; BNST, t = 0.27, P = 0.796; unpaired t test).

Extended Data Figure 6 | Suppression of *Il6ra* expression in AP neurons ameliorates cachexia in the C26 cancer model. *Il6ra* sgRNA-6 group, n = 5 mice, *lacZ* sgRNA group, n = 7 mice. **a.** Survival curves of the mice after tumor inoculation (P = 0.0008, Mantel-Cox test). **b.** Bodyweight of individual mice relative to their bodyweight on the day of tumor inoculation. **c.** Average of bodyweight normalized to that on day -5 ($F_{(1,60)}$ = 8.287, P = 0.0055; day 0, *P = 0.011, two-way repeated-measures (RM) ANOVA with Sidak's post hoc test). **d.** Cumulative food (left) and water (right) intake of the mice after tumor inoculation (food: $F_{(1,100)}$ = 33.61, P = 7.89×10^{-8}, *P = 0.049; water: $F_{(1,100)}$ = 1.977, P = 0.163; two-way RM ANOVA with Sidak's post hoc test). **e.** Cumulative food (left) and water (right) intake of the mice in the 4 days before sacrifice (food: $F_{(1,70)}$ = 0.5572, P = 0.458; water: $F_{(1,50)}$ = 0.1192, P = 0.73; two-way RM ANOVA with Sidak's post hoc test). **f.** Blood glucose levels at different time points ($F_{(1,38)}$ = 29.7, P = 3.24×10^{-6}, *P = 0.013, **P = 0.0056, two-way RM ANOVA with Sidak's post hoc

test). **g.** Inguinal fat mass (t = 2.78, *P = 0.0195, t test). **h.** Quadriceps muscle mass (t = 4.65, ***P = 0.0009, t test). **i.** Tumor mass (t = 7, ****P = 0.000037, t test). **j.** Spleen mass (t = 3.43, **P = 0.0064, t test).