Lytic bacteriophage vB_KmiS-Kmi2C disrupts biofilms formed by members of

the *Klebsiella oxytoca* complex, and represents a novel virus family and genus

Fiona Newberry[1], Preetha Shibu[2]#, Thomas Smith-Zaitlik[1], Mohamed Eladawy[1,3],

Anne L. McCartney[1], Lesley Hoyles[1]*, David Negus[1]*

[1] Department of Biosciences, Nottingham Trent University, United Kingdom

[2] Life Sciences, University of Westminster, United Kingdom

[3] Microbiology and Immunology Department, Faculty of Pharmacy, Mansoura

University, Egypt

* **Corresponding authors:** David Negus, david.negus@ntu.ac.uk; Lesley Hoyles,

lesley.hoyles@ntu.ac.uk

Present address: Berkshire and Surrey Pathology Services, Frimley Health NHS

Trust, Wexham Park Hospital, Slough, United Kingdom.

Abbreviations: CARD, comprehensive antibiotic resistance database; KoC,

Klebsiella oxytoca complex; PC, protein cluster; RBP, receptor binding protein; SD,

standard deviation; TEM, transmission electron microscopy; TSA, tryptone soy agar;

TSBG, tryptone soy broth supplemented with 1 % glucose; VC, viral cluster; VFDB,

virulence factor database.

Keywords: multidrug-resistant, *Klebsiella michiganensis*, siphovirus, phage

taxonomy, biofilm.

Running headline: *Klebsiella* phage vB_KmiS-Kmi2C.

Data availability: The genome sequence of vB_KmiS-Kmi2C has been deposited

with GenBank under accession <u>OP495736</u>. Updated genome data for *K.*

michiganensis strains PS_Koxy1, PS_Koxy2 and PS_Koxy4 are available from

BioProject <u>PRJNA562720</u>. The Cytoscape file used to create **Figure 5** is available

as Supplementary Material from <u>figshare</u>.

ABSTRACT

<u>AIMS</u>

This study aimed to characterise the lytic phage vB_KmiS-Kmi2C, isolated from

sewage water on a GES-positive strain of *Klebsiella michiganensis*.

<u>METHODS AND RESULTS</u>

Comparative phylogenetic and network-based analyses were used to characterise

the genome of phage vB_KmiS-Kmi2C (circular genome of 42,234 bp predicted to

encode 55 genes), demonstrating it shared little similarity with other known phages.

The phage was lytic on clinical strains of *K. oxytoca* (n=2) and *K. michiganensis*

(n=4), and was found to both prevent biofilm formation and disrupt established

biofilms produced by these strains.

<u>CONCLUSIONS</u>

We have identified a phage capable of killing clinically relevant members of the

Klebsiella oxytoca complex (KoC). The phage represents a novel virus family

(proposed name *Dilsviridae*) and genus (proposed name *Dilsvirus*).

<u>SIGNIFICANCE AND IMPACT OF THE STUDY</u>

Identification a novel lytic phage active against clinically relevant strains of the KoC

provides an alternative to antibiotics to treat these increasingly antimicrobial-resistant

opportunistic pathogens. The unusual way in which the phage can disrupt

established biofilms may allow us to identify novel phage-based approaches for

biofilm remediation in the future.

INTRODUCTION

Bacteriophages (phages) are viruses of bacteria that infect their bacterial hosts for the purpose of replication. In the case of lytic phages, this process results in lysis and destruction of the host. The potential for applying the bactericidal action of lytic phages in a therapeutic setting was realised shortly after their discovery, with d'Herelle performing the first recorded clinical studies in 1918 (d'Herelle, 1918). Although interest in phage therapy diminished in Western countries following the introduction of antibiotics, the emergence of multidrug-resistant pathogens has resulted in a resurgence of interest.

Members of the *Klebsiella oxytoca* complex (KoC) are emerging as a serious health concern. The complex currently consists of several distinct phylogroups, representing *K. oxytoca* (Ko2), *K. michiganensis* (Ko1, Ko5), *K. grimontii* (Ko6), *K. huaxiensis* (Ko8), *K. pasteurii* (Ko4), *K. spallanzanii* (Ko3), and three unnamed novel species (Merla *et al.*, 2019; Yang *et al.*, 2022). Members are phylotyped based on the sequence of their chromosomally encoded β-lactamase (*bla*$_{oxy}$) gene (Cosic *et al.*, 2021). KoC bacteria can cause a range of serious infections in both humans and animals and are of particular concern in healthcare settings where immunocompromised patients are predisposed to potential infection. Isolates have been associated with urinary tract infections, lower respiratory tract infections, septicaemia, and soft-tissue injuries (Sohn, Seo & Jung, 2012; Paasch, Wilczek & Strik, 2017; Shakya *et al.*, 2017; Lee *et al.*, 2019). Certain members of the complex (*K. oxytoca, K. pasteurii, K. grimontii, K. michiganensis*) also encode the kleboxymycin biosynthetic gene cluster associated with antibiotic-associated haemorrhagic colitis (Shibu *et al.*, 2021). Worryingly, antibiotic resistance amongst

KoC members is increasing and resistance mechanisms include extended-spectrum

beta-lactamases and carbapenemases (Lowe *et al.*, 2012; White *et al.*, 2016).

New treatment options for tackling drug-resistant infections caused by KoC

isolates would be a welcome addition to our therapeutic armoury. This study aimed

to characterise the morphology, genome and host range of a lytic phage, vB_KmiS-

Kmi2C, isolated against a GES-5-encoding strain of *K. michiganensis* (PS_Koxy2)

(Shibu *et al.*, 2021). The anti-biofilm properties of the phage were investigated to

further determine therapeutic utility with respect to preventing or resolving biofouling

by KoC bacteria such as that which occurs on medical devices, particularly

indwelling catheters and ventilators. Further extensive genomic and phylogenetic

analyses of vB_KmiS-Kmi2C reveal that it represents a novel genus of phage.

MATERIALS AND METHODS

Strain and cultivation information

Details of all strains included in this study can be found in **Table 1**. All strains

were grown on nutrient agar (Sigma Aldrich) unless stated otherwise. Nutrient broth

(Sigma Aldrich) was used for overnight cultures unless stated otherwise, incubated

aerobically at 37 °C. All media used for phage assays were supplemented with

$CaCl_2$ and $MgCl_2$ (both at final concentration of 0.5 mM). The study of anonymised

clinical isolates provided by the Nottingham University Hospitals NHS Trust (NUH)

Pathogen Bank was approved by NUH Research and Innovation (19MI001).

Generation of hybrid genome sequences

Cell pellets from 10-ml overnight cultures grown in nutrient broth were sent to

microbesNG (Birmingham, UK) for DNA extraction, library preparation and

sequencing according to the sequencing provider's strain-submission procedures

(refer to

https://microbesng.com/documents/24/MicrobesNG_Sequencing_Service_Methods_

v20210419.pdf for the protocol).

For Illumina sequencing, the Nextera XT Library Preparation Kit (Illumina, San

Diego, USA) was used and paired-end reads (HiSeq/NovaSeq; 2x250 bp were

generated for PS_Koxy1 (coverage 141×), PS_Koxy2 (coverage 52×) and

PS_Koxy4 (coverage 57×). Reads were adapter-trimmed using Trimmomatic 0.30

(Bolger, Lohse & Usadel, 2014) with a sliding window quality cut-off of Q15. For

Illumina/MinION hybrid genomes, long-read genomic DNA libraries were prepared

with Oxford Nanopore SQK-RBK004 kit and/or SQK-LSK109 kit with Native

Barcoding EXP-NBD104/114 (ONT, United Kingdom) using 400–500 ng of high-

molecular-weight DNA. Barcoded samples were pooled together into a single

sequencing library and loaded into a FLO-MIN106 (R.9.4.1) flow cell in a GridION

(ONT, United Kingdom). *De novo* genome assembly was performed using Unicycler

v0.4.0 (Wick *et al.*, 2017).

The genome assemblies returned to us by microbesNG were annotated in-

house using Bakta v1.6.1, database v4.0 (Schwengers *et al.*, 2021). RGI 6.0.0,

CARD 3.2.5 was used to predict antimicrobial resistance genes encoded within

genomes (Alcock *et al.*, 2020). COPLA and plaSquid were used to characterise

plasmids harboured by strains (Redondo-Salvo *et al.*, 2021; Giménez, Ferrés &

Iraola, 2022).

Biolog GEN III MicroPlate assays

All strains were grown aerobically overnight on blood-free BUG agar (Biolog)

at 37 °C. For each assay, one to two colonies were used to inoculate Biolog

Inoculation Fluid B to a turbidity of 95 % (Biolog turbidimeter). Biolog Gen III plates

were inoculated according to the manufacturer's instructions, and incubated

aerobically at 37 °C for 22 h. Results were read at 540 nm using a BioTek Cytation |[3]

imaging reader spectrophotometer. All Biolog reagents and kits were purchased from

Technopath. Assays were carried out in triplicate for strains PS_Koxy1, PS_Koxy2

and PS_Koxy4.

Biofilm assays and their interpretation

Biofilm assays were performed as described previously (Stepanovic *et al.*,

2000; Merritt, Kadouri & O'Toole, 2005; Eladawy *et al.*, 2021). In brief, a single

colony of each strain was used to inoculate 5 ml of tryptone soy broth supplemented

with 1 % glucose (TSBG). Cultures were incubated aerobically for 24 h at 37 °C

without shaking. The overnight cultures were diluted to 1:100 using TSBG, then

aliquots (100 µl) of the diluted cultures were introduced into wells of a 96-well plate.

The plates were incubated aerobically for 24 h at 37 °C without shaking. Then, the

spent medium was carefully removed from each well. The wells were washed three

times with 200 µl sterile phosphate-buffered saline (pH 7.4; Oxoid) to remove any

non-adherent planktonic cells. The adherent cells were fixed by heat treatment at 60

°C for 60 min to prevent widespread detachment of biofilms prior to dye staining. The

adhered biofilms were then stained by addition of 1 % (w/v) crystal violet (150 µl per

well) and the 96-well plate was left to incubate for 20 min. The excess stain was then

carefully removed from the wells and discarded. The 96-well plate was then carefully

rinsed with distilled water three times, and the plate was inverted and left at room

temperature until the wells were dry. The stained biofilms were solubilised by adding

33 % (v/v) glacial acetic acid (Sigma Aldrich) to each well (150 µl per well). After

solubilisation of the stained biofilms, the OD_{540} was measured and recorded for all

samples using a BioTek Cytation |^3 imaging reader spectrophotometer. *K.*

michiganensis Ko14 and uninoculated medium were used as negative controls in

biofilm assays. Biological (*n*=3) and technical (*n*=4) replicates were done for all

strains. The mean of each isolate's OD quadruplicate readings (OD_i) was calculated

and compared with the control cut-off OD (OD_C), which was defined as three

standard deviations (SD) above the mean of the negative control (3SD + mean). The

amount of biofilm formed was scored as non-adherent ($OD_i \leq OD_C$), weakly adherent

($OD_C < OD_i \leq 2\ OD_C$), moderately adherent ($2\ OD_C < OD_i \leq 4\ OD_C$) or strongly

adherent ($4\ OD_C < OD_i$).

Isolation of lytic phage and host range analysis

Filter-sterilised sewage samples (0.45 µm cellulose acetate filter; Millipore)

collected from mixed-liquor tanks at Mogden Sewage Treatment Works (March

2017) were screened against *K. michiganensis* strain PS_Koxy2 as described

previously (Smith-Zaitlik *et al.*, 2022). Host range analysis was carried out as

described previously (Smith-Zaitlik *et al.*, 2022). After overnight incubation, overlay

assay plates were inspected for lysis, with results recorded according to a

modification of (Haines *et al.*, 2021): ++, complete lysis; +, hazy lysis; 0, no visible

plaques.

Phage concentration and transmission electron microscopy (TEM)

The Vivaspin 20 50 kDa centrifugal concentrator (Cytiva) was used to

concentrate filter-sterilised propagated phage as described previously (Smith-Zaitlik

et al., 2022). For TEM, formvar/carbon-coated 200 mesh copper grids (Agar

Scientific) were prepared via glow discharge (10 mA, 10 s) using a Q150R ES

sputter coater (Quorum Technologies Ltd) and processed as described previously

(Smith-Zaitlik *et al.*, 2022). Samples were visualised using a JEOL JEM-2100Plus

(JEOL Ltd) TEM and an accelerating voltage of 200 kV. Images were analysed and

annotated using ImageJ (https://imagej.net/Fiji).

Phage DNA extraction and sequencing

Phage DNA was extracted from concentrated phage lysate using the Qiagen

DNeasy Blood & Tissue Kit as described previously (Smith-Zaitlik *et al.*, 2022).

Sequence data were generated on our in-house Illumina MiSeq platform, with the

Nextera XT DNA library preparation kit (Illumina) to produce fragments of

approximately 500 bp. Fragmented and indexed samples were run on the sequencer

using a Micro flow cell with the MiSeq Reagent Kit v2 (Illumina; 150-bp paired-end

reads) following Illumina's recommended denaturation and loading procedures.

Phage genome assembly and annotation

Quality of raw sequence data was assessed using FastQC v0.11.9. Reads

had a mean phred score above 30 and contained no adapter sequences, so data

were not trimmed. The genome of the novel phage was assembled using SPAdes

v3.15.4 (default settings) (Bankevich *et al.*, 2012) and visualised using Bandage

v0.8.1 (Wick *et al.*, 2015). Contamination and completeness of the genome was

determined using CheckV v0.8.1 (CheckV database v1.0) (Nayfach *et al.*, 2021).

Gene annotations were made using Prokka (v1.14.6) with the PHROGs database

(v3) (Terzian *et al.*, 2021). The genome was screened for antimicrobial resistance

genes using the Resistance Gene Identifier (v5.2.0) of the Comprehensive Antibiotic

Resistance Database (CARD) (v3.1.4) (Alcock *et al.*, 2020), and for virulence genes

using the Virulence Factor Database (VFDB) (accessed: 6/6/2022) (Liu *et al.*, 2019).

Presence of temperate-lifestyle genes was determined using PhageLeads

(Yukgehnaish *et al.*, 2022). An annotated genome map was generated using

GenoPlotR (v0.8.11) and predicted protein function from the PHROGs database

(Guy, Kultima & Andersson, 2010). To identify potential depolymerase-associated

genes encoded within the phage genome, predicted protein names were searched

for the following terms: 'depolymerase', 'pectin', 'pectate', 'sialidase', 'levanase',

'xylosidase', 'rhamnosidase', 'dextranase', 'alginate', 'hyaluronidase', 'hydrolase',

'lyase'.

Comparative genome analysis

To generate a gene-sharing network, 21,903 phage genomes from the

INPHARED database (April 2022) and the query phage were clustered using

vConTACT2 (v0.9.22; default settings) and visualised in Cytoscape (v.3.9.1) (Bolduc

et al., 2017; Cook *et al.*, 2021). The viral cluster (VC) containing the query phage

was determined and a proteomic phylogenetic tree generated from all phage within

the same VC using VipTree (v1.1.2; default settings) (Nishimura *et al.*, 2017).

Klebsiella oxytoca phage genomes vB_KmiM-2Dii (accession: MZ707156) and

vB_KmiM-4Dii (accession: MZ707157) were used as outliers. An additional

proteomic phylogenetic tree was generated from all phages within the VC and three

representative phage genomes from distantly related genera *Nonagvirus* (JenK1;

NC_029021.1), *Seuratvirus* (CaJan; NC_028776.1) and *Nipunavirus* (NP1;

NC_031058.1). The large terminase protein sequences were used to generate a

maximum likelihood phylogenetic tree with ClustalW (v.2.1) and IQTree with 1000

bootstraps (v1.6.12) (Thompson, Higgins & Gibson, 1994; Minh, Nguyen & von

Haeseler, 2013). IQModelFinder (v.1.4.2) was used with IQTree to determine the

model of best fit (Kalyaanamoorthy *et al.*, 2017). The percentage of shared proteins

within the VC was determined using all-versus-all BLASTP ($\geq$30 % identity and $\geq$50

% sequence coverage; v2.12.0) (Turner, Kropinski & Adriaenssens, 2021). The

intergenomic similarity between phage within the same VC was determined using

VIRIDIC (v1.0) (Moraru, Varsani & Kropinski, 2020). ComplexHeatmap was used to

generated all heatmaps (Gu, Eils & Schlesner, 2016).

Phage–biofilm assays

The titre of the phage stock was determined by plaque assay using the

double-layer agar technique. Briefly, phage vB_KmiS-Kmi2C was serially diluted in

phosphate-buffered saline (pH 7.4; Oxoid) and 100 µl of each phage dilution was

combined with 100 µl of an overnight culture of *K. michiganensis* PS_Koxy2 and 5 ml

of 0.6 % tryptone soy agar (TSA) supplemented with $CaCl_2$ and $MgCl_2$ both at a final

concentration of 1 mM. The mixture was gently swirled and poured onto solid TSA

plates. Plates were incubated overnight at 37 °C and pfu/ml determined by

enumeration of visible plaques.

The ability of vB_KmiS-Kmi2C to prevent and disrupt biofilms was examined

using a modification of a previously described protocol (Taha *et al.*, 2018). Hosts of

vB_KmiS-Kmi2C identified as moderately ($2\ OD_C < OD_i \leq 4\ OD_C$) adherent were

included in the assay. For prevention of biofilms, host cultures were incubated

aerobically for 24 h at 37 °C without shaking in TSBG. Overnight cultures were

diluted 1:100 with TSBG and aliquots (100 µl) of diluted culture were introduced into

wells of a 96-well plate with or without phage vB_KmiS-Kmi2C (added to a final

concentration of 4.5 x 10^8 pfu/ml to each test well). Plates were incubated without

shaking for 24 h at 37 °C. Then, the supernatants were discarded, the biofilm of each

well was washed to remove planktonic cells and biofilms stained as described

above.

To investigate the disruption of established biofilms, host cultures were grown

and prepared as described above prior to inoculating a 96-well plate. Plates were

incubated without shaking for 24 h at 37 °C to allow biofilms to form. Unattached

planktonic cells were carefully aspirated without disrupting the biomass. Phage

vB_KmiS-Kmi2C diluted in 100 µl TSBG was added to test wells (final titre of 4.5×10^8

pfu/ml) whereas control wells received only TSBG without phage. Plates were

incubated for a further 24 h at 37 °C without shaking. Supernatants were carefully

discarded; the biofilm of each well was washed to remove planktonic cells and

biofilms stained as described above. Biological (*n*=3) and technical (*n*=4) replicates

were completed for all strains.

RESULTS

Host range of phage vB_KmiS-Kmi2C

Phage vB_KmiS-Kmi2C was isolated from sewage water on *K. michiganensis* PS_Koxy2 (ST138), a multidrug-resistant, GES-5-positive isolate recovered from human urine (Shibu *et al.*, 2021). The phage was screened against 56 clinical and 28 animal isolates representing a range of *Klebsiella* spp. (*K. michiganensis* n=49, *K. oxytoca* n=25, *K. grimontii* n=9, *K. huaxiensis* n=1) (Shibu *et al.*, 2021; Smith-Zaitlik *et al.*, 2022). It did not infect the closely related *K. michiganensis* strains PS_Koxy1 and PS_Koxy4 (Shibu *et al.*, 2021). vB_KmiS-Kmi2C did not infect animal isolates of *Klebsiella* spp. (Smith-Zaitlik *et al.*, 2022), but did infect some of the clinical *K. oxytoca* (*n*=2) and *K. michiganensis* (*n*=4) strains within our extended in-house collection (Smith-Zaitlik *et al.*, 2022) (**Table 1**). From strain information provided by the Pathogen Bank (EUCAST testing) all these strains were resistant to amoxycillin, and sensitive to amikacin, ceftazidime, ceftazidime, ceftriaxone, cefuroxime, ciprofloxacin, ertapenem, gentamicin, meropenem, piptazobactam and trimethoprim. Strains Ko13, Ko22, Ko43 and Ko53 were sensitive to co-amoxiclav; Ko21 was resistant to co-amoxiclav; no data were supplied for Ko37 and co-amoxiclav. Phage vB_KmiS-Kmi2C did not exhibit depolymerase activity (i.e. it did not form haloes around plaques) on any of the strains that it infected.

Genotypic and phenotypic characterisation of the GES-positive strains

Our previous work had shown PS_Koxy1, PS_Koxy2 and PS_Koxy4 to be very similar based on the analysis of draft genome sequence data (Shibu *et al.*, 2021). In this study, we generated hybrid assemblies of their genomes (**Table 2**).

The initial description of the strains' draft genomes (Ellington *et al.*, 2019)

predicted they each harboured an identical IncQ 8,300-bp circular GES-5-positive

plasmid, confirmed in this study. PS_Koxy2 and PS_Koxy4 both carried an identical

76,870-bp circular plasmid (replication initiation protein domain Rep_3, replicon type

IncR). The 4,448-bp plasmid carried by all three strains shared high identity with

seven other plasmids, harboured by *Salmonella enterica*, *Enterobacter* spp. and *K.*

michiganensis (**Table 3**). A comparison (progressiveMauve, not shown) of the

sequence of pPSKoxy2_4 with those of the other 4,448-bp plasmids in **Table 3**

showed they shared 4,446 identical sites (99.98 % pairwise identity) across their

length. A PlasmidMLST search returned no hits for the plasmid sequence. It was

found to be PTU-E3 (score 96.97 %) by COPLA, exceeding the 90 % threshold to

validate the plasmid taxonomic unit assignment [16], and to belong to mobility group

MOBP1 with a Col440I replicon type by plaSquid (Giménez *et al.*, 2022). The

plasmid was predicted by Bakta to encode six genes (MobA, MobC, two YgdI/YgdR

family lipoproteins, two hypothetical proteins), an origin of replication, *oriT* and a non-

coding RNAI (**Figure 1a**). UniProt BLASTP analyses showed MobA shared 55 %

identity (across 517 aa) with the DNA relaxase MbeA of *Escherichia coli* (reviewed

UniProt record P13658); MobC shared 63.6 % identity (across 115 aa) with the

mobilisation protein MbeC of *E. coli* (reviewed UniProt record P13657). The 39-aa

hypothetical protein was predicted by I-TASSER (Zhang, 2008) to be an alpha helix,

while the 122-aa hypothetical protein was found to include a transmembrane

segment (HHPRED, UniRef30 (Zimmermann *et al.*, 2018)) that had limited structural

homology with only the *Escherichia coli* signal transduction protein PmrD

(PDB:4HN7) and the Vps26 dimer region of a fungal vacuolar protein sorting-

associated protein (PDB:7BLQ).

PS_Koxy4 could be differentiated from PS_Koxy1 and PS_Koxy2 by

phenotypic traits determined using the Biolog GEN III system (**Table 4**). PS_Koxy1

and PS_Koxy2 were resistant to the tetracycline antibiotic minocycline; both were

predicted to encode the tetracycline efflux pump *tet(A)* on incompletely assembled

plasmids (contig_3 PS_Koxy1, predicted to be FIIK by PlasmidMLST; contig_2

PS_Koxy2, predicted to be IncA/C ST12 by PlasmidMLST), but this gene was absent

from the genome of PS_Koxy4, with the strain sensitive to minocycline (**Figure 1b**,

Table 4).

Biofilm-forming ability of strains used in this study

We determined the biofilm-forming ability of strains used in this study as

described previously (Taha *et al.*, 2018). Our results show a wide range of biofilm-

forming abilities among the strains (**Figure 2a**; **Supplementary Table 1**), yet there

was no difference in the virulence factor profiles of PS_Koxy1, PS_Koxy2 and

PS_Koxy4 according to a VFDB (Liu *et al.*, 2019) analysis (**Supplementary Table

2**). Strains *K. michiganensis* PS_Koxy1, *K. michiganensis* Ko13, *K. michiganensis*

Ko21 and *K. michiganensis* Ko22 were identified as weakly adherent ($OD_C < OD_i \leq 2$

OD_C). Strains *K. michiganensis* PS_Koxy2, *K. michiganensis* PS_Koxy4, *K. oxytoca*

Ko37, *K. oxytoca* Ko43 and *K. oxytoca* Ko53 were identified as moderately adherent

($2 OD_C < OD_i \leq 4 OD_C$). *K. michiganensis* Ko14 was used as a non-adherent

negative control ($OD_i \leq OD_C$).

Capacity of phage vB_KmiS-Kmi2C to prevent and disrupt biofilms

Phage vB_KmiS-Kmi2C was found to be highly effective at both preventing

(**Figure 2b**) and disrupting (**Figure 2c**) biofilms formed by isolates identified as

susceptible to lysis. Presence of vB_KmiS-Kmi2C was found to reduce biofilm

formation for all strains included in the study. This reduction was found to be

significant ($P < 0.01$) for three of the isolates tested (PS_Koxy2, Ko37 and Ko53)

compared to no-phage controls. Addition of vB_KmiS-Kmi2C to pre-established

biofilms also resulted in biofilm disruption, seen as a reduction in measured biofilm

compared to non-phage-treated controls. Biofilm disruption was determined to be

significant ($P < 0.05$) for isolates PS_Koxy2 and Ko37 compared to non-phage

treated controls.

Morphology of phage vB_KmiS-Kmi2C

TEM showed vB_KmiS-Kmi2C to have a siphovirus-like morphology (**Figure**

3). It had a long, non-contractile tail. The capsid diameter was 59.6 nm (SD 2.3 nm),

the tail and baseplate were 157.8 nm in length (SD 5.1 nm), and the phage had a

total length of 218.11 nm (SD 4.8) (n=3 measurements).

Genome of phage vB_KmiS-Kmi2C

The genome of phage vB_KmiS-Kmi2C was assembled into a single contig,

with 100 % completeness according to CheckV (Nayfach *et al.*, 2021). Assembly of

the phage was confirmed using visualisation with Bandage (**Supplementary Figure**

1). It comprised 42,234 bp (816 × coverage) and was predicted to encode 55 genes

(upper panel **Figure 4**). The genome is arranged into functional modules as typically

seen in phage genomes: replication/regulation, viral structure, DNA packaging and

lysis. Interesting genomic features of the genome included a Sak4-like ssDNA

annealing protein with single-strand DNA-binding protein, a Cas4-domain

exonuclease, a holin and a Rz-like spanin.

Analysis using NCBI BLASTN showed the genome of vB_KmiS-Kmi2C

shared 85.6 % identity (80 % query coverage) with a phage metagenome-assembled

genome (accession OP072809) recovered from human faeces in Japan (Nishijima *et*

al., 2022). Preliminary analysis of the genome sequence using the online version of

ViPTree suggested the phage was novel (not shown). Further support for this came

from genomic similarity analysis (**Supplementary Figure 2**) and analyses of

protein–protein network data using vConTACT (**Figure 5**). According to vConTACT,

vB_KmiS-Kmi2C shared a VC with 38 other known phage from a range of genera:

Septimatrevius (n=24), *Lokivirus* (n=6), *Titanvirus* (n=4), *Pradovirus* (n=1),

Kilunavirus (n=1) and 2 unclassified genera. The VC did not contain any other phage

isolated using *Klebsiella* and the majority of phage had *Pseudomonas* as an

assigned host (**Figure 6**). The intergenomic similarity within the VC ranged from 100

% to 2 %, with the highest similarity noted between phage within the same genus

(**Supplementary Figure 2**). vB_KmiS-Km2iC showed the highest genomic similarity

(18.8 %) to *Septimatrevirus* vB_Pae-Kakheti25, a *Pseudomonas* phage. The genera

Titanvirus, *Lokivirus*, *Septimatrevirus* do not currently belong to any recognised

phage family, whereas *Pradovirus* belongs to the *Autographiviridae*, which are

podovirus-like in terms of their morphology. Therefore, we suggest vB_KmiS-Kmi2C

does not belong to any currently recognised phage family.

The vConTACT output was used to determine the shared protein clusters

(PCs) within the VC compared to phage vB_KmiS-Kmi2C (lower panel **Figure 4**). Of

the 53 PCs identified in vB_KmiS-Kmi2C, 14 were shared between all phage within

the VC (007793 Cas4-domain exonuclease (KJBENDCP_00001); 006166 DNA

helicase (KJBENDCP_00002); 006824 DNA polymerase processivity factor

(KJBENDCP_00003); 004149 DNA polymerase (KJBENDCP_00004); 005059 tail

length tape measure protein (KJBENDCP_00011); 005663 Neck1 protein

(KJBENDCP_00012); 005850 minor tail protein (KJBENDCP_00015); 008503 tail

terminator (KJBENDCP_00016); 005899 head-tail adaptor Ad1

(KJBENDCP_00018); 003930 head scaffolding protein (KJBENDCP_00022);

002520 portal protein (KJBENDCP_00023); 005106 terminase large subunit

(KJBENDCP_00024); 007093 Sak4-like ssDNA annealing protein

(KJBENDCP_00053); 008553 ssDNA binding protein (KJBENDCP_000154)). Seven

of the shared PCs were associated with viral structure and six were associated with

viral replication/regulation. vB_KmiS-Kmi2C shared the most PCs with *Erwinia*

phage AH03, including most of the viral structure genome module.

Protein-based phylogenetic analysis with VipTree showed phages within the

VC grouped according to the assigned genus, with vB_Kmi2S_Kmi2C not within the

defined genus groups (**Figure 6**). vB_KmiS-Kmi2C showed the closest relationship

to *Erwinia* phage AH03, as previously noted in the lower panel of **Figure 4**. A

representative species from three closely related genera were included in a further

protein-based phylogenetic analysis to determine if vB_KmiS-Kmi2C grouped with

genera outside the VC defined by vConTACT: *Nipunavirus* NP1 (NC_031058),

Nonagavirus JenK1 (NC_024146) and *Seuratvirus* Cajan (NC_028776)

(**Supplementary Figure 3**). Analysis of the shared protein percentage within the

viral cluster was consistent with the intergenomic similarity and protein phylogenetic

analysis (**Supplementary Figure 4**). The phage belonging to the same genera

shared the highest percentage of proteins. vB_KmiS-Kmi2C shared the highest

percentage of proteins with *Pseudomonas* phage *Septimatrevirus* KP1 and *Erwinia*

phage AH03. This confirmed that vB_KmiS-Kmi2C does not belong to any currently

defined genus of phage.

Given that vB_KmiS-Kmi2C represents a novel family, genus and species of

phage, we propose the following taxonomy: *Duplodnaviria* › *Heunggongvirae* ›

Uroviricota › *Caudoviricetes* › *Dilsviridae* › *Dilsvirus* › *Dilsvirus Kmi2C.*

DISCUSSION

There has been a significant research effort into the investigation of phages

for the treatment of *K. pneumoniae* infections and we have recently authored an

extensive review on phages of *Klebsiella* spp. (Herridge *et al.*, 2020). However, there

are relatively few studies documenting the isolation and characterisation of lytic

phages against members of the KoC (Kęsik-Szeloch *et al.*, 2013; Townsend *et al.*,

2021; Smith-Zaitlik *et al.*, 2022), an emerging group of clinical and veterinary

pathogens with growing antibiotic resistance. As such, we set out to identify phages

with the capacity to lyse clinically relevant members of the KoC. We have previously

reported on the characterisation of two such phages with lytic activity against

numerous species within the KoC (Smith-Zaitlik *et al.*, 2022). Here, we report on the

characterisation of phage vB_KmiS-Kmi2C isolated against a carbapenem-resistant

isolate of *K. michiganensis* (PS_Koxy2) (Shibu *et al.*, 2021), for which we generated

a hybrid genome assembly and phenotypic data to allow it to be differentiated from

closely related strains (PS_Koxy1, PS_Koxy4) from our in-house collection.

There were clear differences in the biofilm-forming abilities of the three GES-5

-positive strains, yet all strains encoded the same virulence factors (**Supplementary**

Table 2). We did not compare protein sequences of the virulence factors across the

strains. It is acknowledged that our understanding of KoC phenotypes is poor (Yang

et al., 2022), and our future work will look to address how differences in biofilm-

forming abilities of KoC members and their genetic differences contribute to survival

and potentially pathogenicity.

While the three strains all carried two small (4,448 and 8,300 bp) plasmids,

only PS_Koxy2 and PS_Koxy4 carried a 76,860-bp plasmid. This and the other

larger plasmids carried by the strains will be described elsewhere as part of ongoing

work. The 8,300-bp IncQ plasmid carried by the three strains has been described

elsewhere (Ellington *et al.*, 2019). While the 4,448-bp plasmid (mobility group

MOBP1, replicon type Col440I) has been detected in a strain of *K. michiganensis*

previously (AbuOun *et al.*, 2021), it has not been described in detail. Based on its

gene content (specifically, MobA/MobC), we predict the plasmid to be conjugative

plasmid. The MobA and MobC functional homologues MbeA and MbeC,

respectively, are required for efficient mobilisation of the ColE1 plasmid (Varsaki *et*

al., 2012).

Host range analysis and anti-biofilm properties

Understanding the lytic profile of any individual phage is important for its

potential deployment in a clinical setting, either in an isolated form or for inclusion in

a phage cocktail that may increase host range and potentially suppress phage

resistance (Yang *et al.*, 2020). Ordinarily, phage host range is narrow, sometimes

down to the strain level. In accordance with this, we found that vB_KmiS-Kmi2C

exhibits a relatively narrow host range, lysing six clinical isolates from the KoC out of

a panel of 84 isolates representing a range of clinical and animal *Klebsiella* spp.

To further characterise the therapeutic potential of vB_KmiS-Kmi2C,

experiments were performed to determine the phage's capacity to prevent and

disrupt biofilms formed by the clinical isolates susceptible to lysis in our host range

analysis. Biofilm formation can reduce the efficacy of antibiotic chemotherapy and

contributes to the successful colonisation of wounds and indwelling medical devices

(Wang *et al.*, 2020). Therefore, phages with anti-biofilm properties may make

particularly attractive therapeutics. We found that vB_KmiS-Kmi2C is highly effective

at both preventing biofilm formation and disrupting established biofilms. It is likely

that prevention of biofilm formation is largely due to the lytic action of the phage,

resulting in cell death and the prevention of biofilm formation. However, established

biofilms can be extremely recalcitrant to disruption. This is likely due to the complex

nature of mature biofilms which are typically composed of numerous

macromolecules including, polysaccharides, proteins, nucleic acids and

metabolically inactive cells (Chhibber, Nag & Bansal, 2013). Phage-mediated

disruption of *K. michiganensis* biofilms has been previously reported in the literature

(Ku *et al.*, 2021). The lytic phage KMI8 was shown to disrupt a

K. michiganensis mono-biofilm produced by a strain expressing the polysaccharide

capsule KL70 locus. It was hypothesised this antibiofilm activity was largely due to

the activity of a potential phage-associated depolymerase, identified as a putative

endosialidase encoded within the KMI8 genome. The antibiofilm effects of phages

against *K. oxytoca* have also been investigated (Townsend, Moat & Jameson, 2020).

A recent study showed that phages originally isolated against *K. pneumoniae* were

able to reduce the viability of *K. oxytoca* biofilms when added at the start of biofilm

formation. However, the same phages were unable to reduce the viability of

established *K. oxytoca* biofilms. Here, we found that vB_KmiS-Kmi2C is highly

effective at disrupting established biofilms formed by both *K. michiganensis* and *K.*

oxytoca. In the present study, we found no evidence of depolymerase activity

associated with vB_KmiS-Kmi2C. Plaques on susceptible host lawns did not show

evidence of haloes which are typically indicative of depolymerase activity.

Furthermore, our bioinformatic analysis did not identify any genes associated with

depolymerase activity. Studies are therefore ongoing to determine the mechanism of

biofilm disruption used by vB_KmiS-Kmi2C.

Phage vB_KmiS-Kmi2C represents a new family and genus

Whole-genome sequencing is important for the identification of features

relevant to phage therapy and for aiding taxonomic classification. Genomic analysis

using the resistance gene identifier tool available through the comprehensive

antibiotic resistance database did not identify the presence of known antimicrobial

resistance mechanisms within the genome of vB_KmiS-Kmi2C. The presence of

such genes would suggest the potential to transfer antibiotic resistance markers

between bacteria if vB_KmiS-Kmi2C were to be applied therapeutically in a

polymicrobial setting. PhageLeads did not detect the presence of classical lysogeny

genes, suggesting vB_KmiS-Kmi2C is exclusively lytic.

Initial genome sequence analysis of vB_KmiS-Kmi2C using BLASTN and

ViPTree suggested that our phage was novel, sharing little sequence similarity with

other deposited phage genomes. To aid with taxonomy, we employed the gene-

sharing network analysis tool vConTACT2. Phage taxonomic assignment using

whole genome gene-sharing profiles has been shown to be highly accurate; a recent

study showed that vConTACT2 produces near-identical replication of existing genus-

level viral taxonomy assignments from the International Committee on Taxonomy of

Viruses (ICTV) (Bin Jang *et al.*, 2019). Our analysis using vConTACT2 and the

INPHRARED database of phage genomes identified that vB_KmiS-Kmi2C shared a

VC with 38 other sequenced phages. The VC was polyphyletic, containing phage

from at least five different genera, suggesting that vB_KmiS-Km2iC did not group

with any existing genus. Recently published guidelines have suggested that for

genome-based phage taxonomy, assignment to taxonomic ranks should be based

on nucleotide sequence identity across entire genomes (Adriaenssens & Brister,

2017; Turner *et al.*, 2021). These guidelines suggest any two phages belong to the

same species if they are more than 95 % identical across their genome and belong

to the same genus if they are more than 70 % identical. We used VIRIDIC to

determine the intergenomic similarity between vB_KmiS-Kmi2C and the 38 other

phages identified within the VC. Our findings show that based on these criteria,

vB_KmiS-Kmi2C cannot be assigned to any current species or genus as it shares

only 18.8 % genomic similarity to *Septimatrevirus* vB_Pae-Kakheti25, a

Pseudomonas phage, the closest nucleotide sequence relative we could identify.

Most of the phages within the VC (21/38) were identified as infecting bacteria

belonging to the genus *Pseudomonas*. According to the GenBank accession details

associated with the genomes of these phages, they were all isolated-on strains of *P.*

aeruginosa. A preliminary screen of vB_KmiS-Kmi2C against a panel of 11 *P.*

aeruginosa isolates did not identify any lytic activity against members of this species

(data not shown).

In addition to identifying phages belonging to a VC, vConTACT2 can also

identify proteins shared by members within the VC. Our analysis found that 14

proteins were shared between all phages within the VC, of which seven were

associated with viral structure and six were associated with viral

replication/regulation. At least 35 of the phages within the VC are from genera with

siphovirus-like morphology. It is therefore unsurprising that phages within the cluster

share several proteins related to general structure/morphology.

Phage vB_KmiS-Kmi2C is the only member of the VC known to infect

members of the genus *Klebsiella*. Host-cell tropism is often determined by receptor-

binding proteins (RBPs) which mediate recognition and attachment to host cells

(Dams *et al.*, 2019). RBPs are typically identified as phage tail fibres or spikes

(Nobrega *et al.*, 2018). The genomic region of vB_KmiS-Kmi2C encoding

hypothetical tail fibre proteins can be seen in **Figure 4** and comprises proteins

088411, 095417, 028210, 033709, 031062 and 060189 (KJBENDCP_00005 to

KJBENDCP_00010). Unannotated proteins within this region were confirmed to

share homology with phage tail fibre proteins by HHPred analysis using an e-value

cut-off of >10^{-5}. The tail fibre proteins of vB_KmiS-Kmi2C are not widely present

within the VC and are typically shared by only one or two phages within the cluster.

This may explain the host tropism of vB_KmiS-Kmi2C compared to other members

of the VC but further experimentation will be required to determine this. Protein

005059 (KJBENDCP_00011) is a predicted tail length tape measure protein; such

proteins are involved in determining tail length and participate in DNA injection into

the host-cells and are not involved in host recognition (Mahony *et al.*, 2016).

Our additional protein-based phylogenetic analyses, including phages from

outside the VC, highlighted that vB_Kmi2S_Kmi2C does not group with any genera

included in the analyses and as such does not belong to any currently defined genus

of phage.

Phages are recognised as the most abundant biological entities on the planet

and exhibit extensive biological diversity (Hendrix, 2002). With an estimated 10^{31}

phage virions in the biosphere (Hatfull, 2015) and 14,244 complete phage genomes

sequenced as of 2021 (Cook *et al.*, 2021), it is clear we have characterised a

relatively small number of distinct phages. However, with the cost of genome

sequencing falling and the capacity to assemble phage genomes from metagenome

sequence data, it is likely that the number of novel phages identified will increase

rapidly. The classification of phages representing entirely novel genera or

families/sub-families poses a challenge for taxonomy, especially when new isolates

share little sequence similarity with previously characterised phages. Previously,

tailed phages were assigned to a family based on morphology, with the families

Podoviridae, *Myoviridae* and *Siphoviridae* belonging to the order *Caudovirales*.

However, the increase in available phage sequence data has shown that these

families are paraphyletic. It has, therefore, been suggested that they are abolished,

and that phage morphology should no longer play a role in classification (Turner *et*

al., 2021). Based on recently updated guidelines for the classification of phage

based on nucleotide sequence identity, and subsequent protein based phylogenetic

analyses, we show here that phage vB_KmiS-Kmi2C represents a new genus.

AUTHOR CONTRIBUTION STATEMENT

Conceptualisation: FN, PS, LH, DN. Data curation: all authors. Formal

analysis: FN, TSZ, PS, ME, ALM, LH, DN. Funding acquisition: PS, ALM, LH.

Investigation: FN, TSZ, PS, ME, ALM, LH, DN. Methodology: FN, ME, LH, DN.

Resources: ALM, LH, DN. Supervision: ALM, LH, DN. Visualisation: FN, LH, DN.

Writing – original draft: FN, ME, LH, DN. Writing – reviewing and editing: all authors.

ACKNOWLEDGEMENTS

PS was in receipt of an IBMS Research Grant (project title "Isolation of lytic

bacteriophages active against antibiotic-resistant *Klebsiella pneumoniae*"). Imperial

Health Charity is thanked for contributing to registration fees for the Professional

Doctorate studies of PS. ME is supported by the Egyptian Government – Ministry of

Higher Education/Egyptian Cultural and Education Bureau. This work used

computing resources funded by the Research Contingency Fund of the Department

of Biosciences, Nottingham Trent University (NTU). TSZ completed this work as part

of an MRes degree at NTU.

REFERENCES

AbuOun, M., Jones, H., Stubberfield, E., Gilson, D., Shaw, L.P., Hubbard, A.T.M., Chau, K.K., Sebra, R.,
Peto, T.E.A., Crook, D.W., Read, D.S., Gweon, H.S., Walker, A.S., Stoesser, N., Smith, R.P.,
Anjum, M.F. and On Behalf Of The Rehab Consortium, null (2021) A genomic
epidemiological study shows that prevalence of antimicrobial resistance in Enterobacterales
is associated with the livestock host, as well as antimicrobial usage. *Microb Genomics*. **7**,
000630.

Adriaenssens, E. and Brister, J.R. (2017) How to Name and Classify Your Phage: An Informal Guide.
Viruses. **9**, E70.

Alcock, B.P., Raphenya, A.R., Lau, T.T.Y., Tsang, K.K., Bouchard, M., Edalatmand, A., Huynh, W.,
Nguyen, A.-L.V., Cheng, A.A., Liu, S., Min, S.Y., Miroshnichenko, A., Tran, H.-K., Werfalli, R.E.,
Nasir, J.A., Oloni, M., Speicher, D.J., Florescu, A., Singh, B., Faltyn, M., Hernandez-
Koutoucheva, A., Sharma, A.N., Bordeleau, E., Pawlowski, A.C., Zubyk, H.L., Dooley, D.,
Griffiths, E., Maguire, F., Winsor, G.L., Beiko, R.G., Brinkman, F.S.L., Hsiao, W.W.L.,
Domselaar, G.V. and McArthur, A.G. (2020) CARD 2020: antibiotic resistome surveillance
with the comprehensive antibiotic resistance database. *Nucleic Acids Res*. **48**, D517–D525.

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin, V.M., Nikolenko,
S.I., Pham, S., Prjibelski, A.D., Pyshkin, A.V., Sirotkin, A.V., Vyahhi, N., Tesler, G., Alekseyev,
M.A. and Pevzner, P.A. (2012) SPAdes: a new genome assembly algorithm and its
applications to single-cell sequencing. *J Comput Biol J Comput Mol Cell Biol*. **19**, 455–477.

Bin Jang, H., Bolduc, B., Zablocki, O., Kuhn, J.H., Roux, S., Adriaenssens, E.M., Brister, J.R., Kropinski,
A.M., Krupovic, M., Lavigne, R., Turner, D. and Sullivan, M.B. (2019) Taxonomic assignment
of uncultivated prokaryotic virus genomes is enabled by gene-sharing networks. *Nat
Biotechnol*. **37**, 632–639.

Bolduc, B., Jang, H.B., Doulcier, G., You, Z.-Q., Roux, S. and Sullivan, M.B. (2017) vConTACT: an iVirus
tool to classify double-stranded DNA viruses that infect Archaea and Bacteria. *PeerJ*. **5**,
e3243.

Bolger, A.M., Lohse, M. and Usadel, B. (2014) Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinforma Oxf Engl*. **30**, 2114–2120.

Chhibber, S., Nag, D. and Bansal, S. (2013) Inhibiting biofilm formation by Klebsiella pneumoniae
B5055 using an iron antagonizing molecule and a bacteriophage. *BMC Microbiol*. **13**, 174.

Clokie, M.R., Millard, A.D., Letarov, A.V. and Heaphy, S. (2011) Phages in nature. *Bacteriophage*. **1**,
31–45.

Cook, R., Brown, N., Redgwell, T., Rihtman, B., Barnes, M., Clokie, M., Stekel, D.J., Hobman, J., Jones,
M.A. and Millard, A. (2021) INfrastructure for a PHAge REference Database: Identification of
Large-Scale Biases in the Current Collection of Cultured Phage Genomes. *PHAGE*. **2**, 214–223.

Cosic, A., Leitner, E., Petternel, C., Galler, H., Reinthaler, F.F., Herzog-Obereder, K.A., Tatscher, E.,
Raffl, S., Feierl, G., Högenauer, C., Zechner, E.L. and Kienesberger, S. (2021) Variation in
Accessory Genes Within the *Klebsiella oxytoca* Species Complex Delineates Monophyletic
Members and Simplifies Coherent Genotyping. *Front Microbiol*. **12**, 692453.

Dams, D., Brøndsted, L., Drulis-Kawa, Z. and Briers, Y. (2019) Engineering of receptor-binding
proteins in bacteriophages and phage tail-like bacteriocins. *Biochem Soc Trans*. **47**, 449–460.

Eladawy, M., El-Mowafy, M., El-Sokkary, M.M.A. and Barwa, R. (2021) Antimicrobial resistance and
virulence characteristics in ERIC-PCR typed biofilm forming isolates of P. aeruginosa. *Microb*
Pathog. **158**, 105042.

Ellington, M.J., Davies, F., Jauneikaite, E., Hopkins, K.L., Turton, J.F., Adams, G., Pavlu, J., Innes, A.J.,
Eades, C., Brannigan, E.T., Findlay, J., White, L., Bolt, F., Kadhani, T., Chow, Y., Patel, B.,
Mookerjee, S., Otter, J.A., Sriskandan, S., Woodford, N. and Holmes, A. (2019) A multi-
species cluster of GES-5 carbapenemase producing Enterobacterales linked by a
geographically disseminated plasmid. *Clin Infect Dis Off Publ Infect Dis Soc Am*.

Giménez, M., Ferrés, I. and Iraola, G. (2022) Improved detection and classification of plasmids from
circularized and fragmented assemblies. *bioRxiv*.

Gu, Z., Eils, R. and Schlesner, M. (2016) Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinforma Oxf Engl*. **32**, 2847–2849.

Guy, L., Kultima, J.R. and Andersson, S.G.E. (2010) genoPlotR: comparative gene and genome
visualization in R. *Bioinforma Oxf Engl*. **26**, 2334–2335.

Haines, M.E.K., Hodges, F.E., Nale, J.Y., Mahony, J., van Sinderen, D., Kaczorowska, J., Alrashid, B.,
Akter, M., Brown, N., Sauvageau, D., Sicheritz-Pontén, T., Thanki, A.M., Millard, A.D., Galyov,
E.E. and Clokie, M.R.J. (2021) Analysis of Selection Methods to Develop Novel Phage Therapy
Cocktails Against Antimicrobial Resistant Clinical Isolates of Bacteria. *Front Microbiol*. **12**,
613529.

Hatfull, G.F. (2015) Dark Matter of the Biosphere: the Amazing World of Bacteriophage Diversity. *J*
Virol. **89**, 8107–8110.

Hendrix, R.W. (2002) Bacteriophages: Evolution of the Majority. *Theor Popul Biol*. **61**, 471–480.

d'Herelle, F. (1918) Sur le rôle du microbe filtrant bactériophage dans la dysentérie bacillaire.
Comptes Rendus Académie Sci. **167**, 970–972.

Herridge, W.P., Shibu, P., O'Shea, J., Brook, T.C. and Hoyles, L. (2020) Bacteriophages of Klebsiella
spp., their diversity and potential therapeutic uses. *J Med Microbiol*. **69**, 176–194.

Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., von Haeseler, A. and Jermiin, L.S. (2017)
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods*. **14**,
587–589.

Kęsik-Szeloch, A., Drulis-Kawa, Z., Weber-Dąbrowska, B., Kassner, J., Majkowska-Skrobek, G.,
Augustyniak, D., Łusiak-Szelachowska, M., Żaczek, M., Górski, A. and Kropinski, A.M. (2013)
Characterising the biology of novel lytic bacteriophages infecting multidrug resistant
Klebsiella pneumoniae. *Virol J*. **10**, 100.

Ku, H., Kabwe, M., Chan, H.T., Stanton, C., Petrovski, S., Batinovic, S. and Tucci, J. (2021) Novel
Drexlerviridae bacteriophage KMI8 with specific lytic activity against Klebsiella michiganensis
and its biofilms. *PLOS ONE*. **16**, e0257102.

Lee, S.H., Ruan, S.-Y., Pan, S.-C., Lee, T.-F., Chien, J.-Y. and Hsueh, P.-R. (2019) Performance of a
multiplex PCR pneumonia panel for the identification of respiratory pathogens and the main
determinants of resistance from the lower respiratory tract specimens of adult patients in
intensive care units. *J Microbiol Immunol Infect*. **52**, 920–928.

Liu, B., Zheng, D., Jin, Q., Chen, L. and Yang, J. (2019) VFDB 2019: a comparative pathogenomic
platform with an interactive web interface. *Nucleic Acids Res*. **47**, D687–D692.

Lowe, C., Willey, B., O'Shaughnessy, A., Lee, W., Lum, M., Pike, K., Larocque, C., Dedier, H., Dales, L.,
Moore, C., McGeer, A., and the Mount Sinai Hospital Infection Control Team (2012)
Outbreak of Extended-Spectrum β-Lactamase–producing *Klebsiella oxytoca* Infections
Associated with Contaminated Handwashing Sinks1. *Emerg Infect Dis*. **18**, 1242–1247.

Mahony, J., Alqarni, M., Stockdale, S., Spinelli, S., Feyereisen, M., Cambillau, C. and Sinderen, D. van
(2016) Functional and structural dissection of the tape measure protein of lactococcal phage
TP901-1. *Sci Rep*. **6**, 36667.

Merla, C., Rodrigues, C., Passet, V., Corbella, M., Thorpe, H.A., Kallonen, T.V.S., Zong, Z., Marone, P.,
Bandi, C., Sassera, D., Corander, J., Feil, E.J. and Brisse, S. (2019) Description of Klebsiella
spallanzanii sp. nov. and of Klebsiella pasteurii sp. nov. *Front Microbiol*. **10**, 2360.

Merritt, J.H., Kadouri, D.E. and O'Toole, G.A. (2005) Growing and analyzing static biofilms. *Curr*
Protoc Microbiol. **Chapter 1**, Unit 1B.1.

Minh, B.Q., Nguyen, M.A.T. and von Haeseler, A. (2013) Ultrafast Approximation for Phylogenetic
Bootstrap. *Mol Biol Evol*. **30**, 1188–1195.

Moraru, C., Varsani, A. and Kropinski, A.M. (2020) VIRIDIC—A Novel Tool to Calculate the
Intergenomic Similarities of Prokaryote-Infecting Viruses. *Viruses*. **12**, 1268.

Nayfach, S., Camargo, A.P., Schulz, F., Eloe-Fadrosh, E., Roux, S. and Kyrpides, N.C. (2021) CheckV
assesses the quality and completeness of metagenome-assembled viral genomes. *Nat*
Biotechnol. **39**, 578–585.

Nishijima, S., Nagata, N., Kiguchi, Y., Kojima, Y., Miyoshi-Akiyama, T., Kimura, M., Ohsugi, M., Ueki,
K., Oka, S., Mizokami, M., Itoi, T., Kawai, T., Uemura, N. and Hattori, M. (2022) Extensive gut
virome variation and its associations with host and environmental factors in a population-
level cohort. *Nat Commun*. **13**, 5252.

Nishimura, Y., Yoshida, T., Kuronishi, M., Uehara, H., Ogata, H. and Goto, S. (2017) ViPTree: the viral
proteomic tree server. *Bioinforma Oxf Engl*. **33**, 2379–2380.

Nobrega, F.L., Vlot, M., de Jonge, P.A., Dreesens, L.L., Beaumont, H.J.E., Lavigne, R., Dutilh, B.E. and
Brouns, S.J.J. (2018) Targeting mechanisms of tailed bacteriophages. *Nat Rev Microbiol*. **16**,
760–773.

Paasch, C., Wilczek, S. and Strik, M.W. (2017) Liver abscess and sepsis caused by *Clostridium*
perfringens and *Klebsiella oxytoca*. *Int J Surg Case Rep*. **41**, 180–183.

Redondo-Salvo, S., Bartomeus-Peñalver, R., Vielva, L., Tagg, K.A., Webb, H.E., Fernández-López, R.
and de la Cruz, F. (2021) COPLA, a taxonomic classifier of plasmids. *BMC Bioinformatics*. **22**,
390.

Schwengers, O., Jelonek, L., Dieckmann, M.A., Beyvers, S., Blom, J. and Goesmann, A. (2021) Bakta:
rapid and standardized annotation of bacterial genomes via alignment-free sequence
identification. *Microb Genomics*. **7**.

Shakya, P., Shrestha, D., Maharjan, E., Sharma, V.K. and Paudyal, R. (2017) ESBL Production Among *E.*
coli and *Klebsiella* spp. Causing Urinary Tract Infection: A Hospital Based Study. *Open*
Microbiol J. **11**, 23–30.

Shibu, P., McCuaig, F., McCartney, A.L., Kujawska, M., Hall, L.J. and Hoyles, L. (2021) Improved
molecular characterization of the *Klebsiella oxytoca* complex reveals the prevalence of the
kleboxymycin biosynthetic gene cluster. *Microb Genomics*. **7**, 000592.

Smith-Zaitlik, T., Shibu, P., McCartney, A.L., Foster, G., Hoyles, L. and Negus, D. (2022) Extended
genomic analyses of the broad-host-range phages vB_KmiM-2Di and vB_KmiM-4Dii reveal
slopekviruses have highly conserved genomes. *Microbiology*. **168**, 001247.

Sohn, W.-I., Seo, B.F. and Jung, S.-N. (2012) Forehead Abscess Caused by *Klebsiella oxytoca* With
Undiagnosed Type 2 Diabetes: *J Craniofac Surg*. **23**, 247–249.

Stepanovic, S., Vukovic, D., Dakic, I., Savic, B. and Svabic-Vlahovic, M. (2000) A modified microtiter-
plate test for quantification of staphylococcal biofilm formation. *J Microbiol Methods*. **40**,
175–179.

Taha, O.A., Connerton, P.L., Connerton, I.F. and El-Shibiny, A. (2018) Bacteriophage ZCKP1: A
Potential Treatment for Klebsiella pneumoniae Isolated From Diabetic Foot Patients. *Front*
Microbiol. **9**, 2127.

Terzian, P., Olo Ndela, E., Galiez, C., Lossouarn, J., Pérez Bucio, R.E., Mom, R., Toussaint, A., Petit, M.-
A. and Enault, F. (2021) PHROG: families of prokaryotic virus proteins clustered using remote
homology. *NAR Genomics Bioinforma*. **3**, lqab067.

Thompson, J.D., Higgins, D.G. and Gibson, T.J. (1994) CLUSTAL W: improving the sensitivity of
progressive multiple sequence alignment through sequence weighting, position-specific gap
penalties and weight matrix choice. *Nucleic Acids Res*. **22**, 4673–4680.

Townsend, E.M., Kelly, L., Gannon, L., Muscatt, G., Dunstan, R., Michniewski, S., Sapkota, H.,
Kiljunen, S.J., Kolsi, A., Skurnik, M., Lithgow, T., Millard, A.D. and Jameson, E. (2021) Isolation
and Characterization of *Klebsiella* Phages for Phage Therapy. *PHAGE*. **2**, 26–42.

Townsend, E.M., Moat, J. and Jameson, E. (2020) CAUTI's next top model - Model dependent
Klebsiella biofilm inhibition by bacteriophages and antimicrobials. *Biofilm*. **2**, 100038.

Turner, D., Kropinski, A.M. and Adriaenssens, E.M. (2021) A Roadmap for Genome-Based Phage
Taxonomy. *Viruses*. **13**, 506.

Varsaki, A., Lamb, H.K., Eleftheriadou, O., Vandera, E., Thompson, P., Moncalián, G., de la Cruz, F.,
Hawkins, A.R. and Drainas, C. (2012) Interaction between relaxase MbeA and accessory
protein MbeC of the conjugally mobilizable plasmid ColE1. *FEBS Lett*. **586**, 675–679.

Wang, G., Zhao, G., Chao, X., Xie, L. and Wang, H. (2020) The Characteristic of Virulence, Biofilm and
Antibiotic Resistance of Klebsiella pneumoniae. *Int J Environ Res Public Health*. **17**, 6278.

Webb, H.E., Kim, J.Y., Tagg, K.A., de la Cruz, F., Peñil-Celis, A., Tolar, B., Ellison, Z., Schwensohn, C.,
Brandenburg, J., Nichols, M. and Folster, J.P. (n.d.) Genome Sequences of 18 Salmonella
enterica Serotype Hadar Strains Collected from Patients in the United States. *Microbiol*
Resour Announc. **11**, e00522-22.

White, L., Hopkins, K.L., Meunier, D., Perry, C.L., Pike, R., Wilkinson, P., Pickup, R.W., Cheesbrough, J.
and Woodford, N. (2016) Carbapenemase-producing *Enterobacteriaceae* in hospital
wastewater: a reservoir that may be unrelated to clinical isolates. *J Hosp Infect*. **93**, 145–151.

Wick, R.R., Judd, L.M., Gorrie, C.L. and Holt, K.E. (2017) Unicycler: Resolving bacterial genome
assemblies from short and long sequencing reads. *PLoS Comput Biol*. **13**, e1005595.

Wick, R.R., Schultz, M.B., Zobel, J. and Holt, K.E. (2015) Bandage: interactive visualization of de novo
genome assemblies. *Bioinforma Oxf Engl*. **31**, 3350–3352.

Yachison, C.A., Yoshida, C., Robertson, J., Nash, J.H.E., Kruczkiewicz, P., Taboada, E.N., Walker, M.,
Reimer, A., Christianson, S., Nichani, A., PulseNet Canada Steering Committee and Nadon, C.
(2017) The Validation and Implications of Using Whole Genome Sequencing as a
Replacement for Traditional Serotyping for a National Salmonella Reference Laboratory.
Front Microbiol. **8**, 1044.

Yang, J., Long, H., Hu, Y., Feng, Y., McNally, A. and Zong, Z. (2022) Klebsiella oxytoca Complex:
Update on Taxonomy, Antimicrobial Resistance, and Virulence. *Clin Microbiol Rev*. **35**,
e0000621.

Yang, Y., Shen, W., Zhong, Q., Chen, Q., He, X., Baker, J.L., Xiong, K., Jin, X., Wang, J., Hu, F. and Le, S.
(2020) Development of a Bacteriophage Cocktail to Constrain the Emergence of Phage-
Resistant *Pseudomonas aeruginosa*. *Front Microbiol*. **11**, 327.

Yukgehnaish, K., Rajandas, H., Parimannan, S., Manickam, R., Marimuthu, K., Petersen, B., Clokie,
M.R.J., Millard, A. and Sicheritz-Pontén, T. (2022) PhageLeads: Rapid Assessment of Phage
Therapeutic Suitability Using an Ensemble Machine Learning Approach. *Viruses*. **14**, 342.

Zhang, Y. (2008) I-TASSER server for protein 3D structure prediction. *BMC Bioinformatics*. **9**, 40.

Zimmermann, L., Stephens, A., Nam, S.-Z., Rau, D., Kübler, J., Lozajic, M., Gabler, F., Söding, J., Lupas,
A.N. and Alva, V. (2018) A Completely Reimplemented MPI Bioinformatics Toolkit with a
New HHpred Server at its Core. *J Mol Biol*. **430**, 2237–2243.

Table 1. *K. oxytoca* complex strains used in this study

Strain *	Species	Source	Reference
PS_Koxy1	*K. michiganensis*	Throat swab	(Shibu *et al.*, 2021)
PS_Koxy2 §#	*K. michiganensis*	Urine	(Shibu *et al.*, 2021)
PS_Koxy4	*K. michiganensis*	Rectal swab	(Shibu *et al.*, 2021)
Ko13 §	*K. michiganensis*	Blood culture	(Smith-Zaitlik *et al.*, 2022)
Ko14	*K. michiganensis*	Blood culture	(Smith-Zaitlik *et al.*, 2022)
Ko21 §	*K. michiganensis*	Perfusion fluid	(Smith-Zaitlik *et al.*, 2022)
Ko22 §	*K. michiganensis*	Perfusion fluid	(Smith-Zaitlik *et al.*, 2022)
Ko37 §	*K. oxytoca*	Blood culture	(Smith-Zaitlik *et al.*, 2022)
Ko43 §	*K. michiganensis*	Bile fluid	(Smith-Zaitlik *et al.*, 2022)
Ko53 §	*K. oxytoca*	Blood culture	(Smith-Zaitlik *et al.*, 2022)

* Ko prefix indicates strains came from the Pathogen Bank of Queen's Medical

Centre Nottingham, UK.

§ Infected by phage vB_KmiS-Kmi2C.

Strain on which phage vB_KmiS-Kmi2C was isolated; all other strains used in host

range analysis but not infected by the phage are described in (Smith-Zaitlik *et al.*,

2022).

Table 2. Summary statistics for hybrid genome assemblies of the GES-5-positive

strains

Strain (accession)	Contigs	Size (bp)	Status *	N50	CDS	Plasmids (RIP; MOB; REP; PTU) #
PS_Koxy1 (GCA_014050555.2)	7	6,392,093	CCC; 4 CCPs; ? incomplete plasmid	6,023,296	5,952	pPSKoxy1_1 (RepA_C; MOBH; IncAC, IncN; –) pPSKoxy1_2 (Rep_3; –; IncR; –) pPSKoxy1_3 [–; MOBP1; –; PTU-Q2 (IncQ)] pPSKoxy1_4 (–; MOBP1; Col440I; PTU-E3)
PS_Koxy2 (GCA_014050535.2)	10	6,357,486	Draft chromosome; 4 CCPs; ? incomplete plasmid	5,917,750	5,910	pPSKoxy2_1 (IncFII_repA, IncFII_repA, RepA_C, RepB–RCR_reg; MOBF, MOBH; IncFII_1_pKP91, IncAC, IncN; –) pPSKoxy2_2 (Rep_3; –; IncR; –) pPSKoxy2_3 [–; MOBP1; –; PTU-Q2 (IncQ)] pPSKoxy2_4 (–; MOBP1; Col440I; PTU-E3)
PS_Koxy4 (GCA_014050515.2)	6	6,406,207	CCC; 5 CCPs	6,075,855	5,969	pPSKoxy4_1 [IncFII_repA, IncFII_repA, RepB-RCR_reg; MOBF; IncFII_1_pKP91; PTU-FK (IncFII)] pPSKoxy4_2 [–; MOBH; IncAC; PTU-C (IncA)] pPSKoxy4_3 (Rep_3; –; IncR; –) pPSKoxy4_4 [–; MOBP1; –; PTU-Q2 (IncQ)] pPSKoxy4_5 (–; MOBP1; Col440I; PTU-E3)

*CCC, complete circular chromosome; CCP, complete circular plasmid.

RIP, replication initiation protein domain; MOB, mobility group; REP, replicon type;

PTU, plasmid taxonomy unit; –, no data available. RIP, MOB and REP determined

using plaSquid (Giménez *et al.*, 2022), PTU determined using COPLA (Redondo-

Salvo *et al.*, 2021).

Table 3. Closest relatives of the circular 4,448 bp plasmid as determined using NCBI BLASTN

Species, strain and plasmid	Identity	Size (nt)	Accession	Source	Location	Reference
S. enterica PNUSAS021403 pPNUSAS021403_3	3511/3513 (99 %)	4,448	CP093106.1	Patient	USA	(Webb *et al.*, 2022)
S. enterica subsp. *enterica* serovar Hadar PNUSAS018090 pPNUSAS018090_3	2662/2664 (99 %)	4,448	CP093099.1	Patient	USA	(Webb *et al.*, 2022)
S. enterica 2016K-0377 p2016K-0377_3	4387/4389 (99 %)	4,448	CP093079.1	Patient	USA	(Webb *et al.*, 2022)
S. enterica subsp. *enterica* serovar Hadar 12-2388 p12-2388.3	2662/2664 (99 %)	4,448	CP038598.1	Reference laboratory	Canada	(Yachison *et al.*, 2017)
E. cloacae 78 pP6	3403/3403 (100 %)	9,923	OW849463.1	Hospital	Spain	BioProject PRJEB42440
E. hormaechei RHBSTW-00070 pRHBSTW-00070_3	3403/3403 (100 %)	9,923	CP058166.1	Wastewater influent	UK	(AbuOun *et al.*, 2021)
K. michiganensis RHBSTW-00676 pRHBSTW-00676_6	3211/3211 (100 %)	4,448	CP055990.1	Wastewater influent	UK	(AbuOun *et al.*, 2021)

800   **Table 4.** Phenotypic traits that differentiate PS_Koxy1 and PS_Koxy2 from

801   PS_Koxy4

Growth in presence of	PS_Koxy1	PS_Koxy2	PS_Koxy4
D-Serine	–	–	+
L-Alanine	+	+	–
Minocycline	+	+	–
Acetic acid	–	–	+

802   +, Positive according to the Biolog GEN III system; –, negative according to the

803   Biolog GEN III system.

Figure 1. (a) Illustration of the circular conjugative 4,448-bp plasmid harboured by *K.*

michiganensis strains PS_Koxy1, PS_Koxy2 and PS_Koxy4. **(b)** Tetracycline-

antibiotic-associated genes predicted to be encoded within the genomes of strains

PS_Koxy1, PS_Koxy2 and PS_Koxy4 by RGI/CARD. All hits were considered 'strict'

by CARD.

Figure 2. Biofilm-forming abilities of the strains used in this study. **(a)** The biofilm-

forming abilities of all strains listed in **Table 1** were assessed. *K. michiganensis*

Ko14, a strain that does not form biofilms in TSBG, was used as the negative

control. Differences in biofilm formation in comparison with the control were

statistically significant for all strains (unpaired *t* test, adjusted *P* value < 0.05,

Benjamini-Hochberg; **Supplementary Table 1**). The ability of phage vB_KmiS-

Kmi2C to **(b)** prevent biofilm formation or **(c)** disrupt established biofilms was

assessed using those strains infected by the phage and forming the strongest

biofilms **(a)**. **(b, c)** Statistical significance of the differences in biofilm formation of

strains in the presence and absence of phage was assessed using unpaired *t* test.

(a–c) Data are shown as mean for four technical and three biological replicates per

strain.

Figure 3. Transmission electron micrograph of phage vB_KmiS-Kmi2C. Scale bar,

50 nm.

Figure 4. Genome map and predicted function of vB_KmiS-Kmi2C (upper image) and presence/absence of PCs within VC (lower

image). The genome map shows the predicted function of the coding regions, direction of transcription and PC ID for each gene

segment. The phage genus and host genus are shown to the left of the PC heatmap (see legend for annotation) and phage

genome size on the right-hand side of the heatmap.

Figure 5. Protein-sharing network generated from vB_KmiS-Kmi2C and INPHARED database. Each node represents a phage genome and connection between the genomes shown by a grey line. The nodes are annotated according to viral family. The VC containing vB_KmiS-Kmi2C is highlighted with a black circle.

Figure 6. Protein phylogenetic analysis generated with VipTree from all phage within the VC. The inner circle is coloured according to phage host and outer circle according to assigned phage genus. The tree was rooted at the midpoint and scale shown in the bottom right. *Klebsiella* phage vB_KmiM-4Dii (KiM 4Dii), a *Slopekvirus* phage (Smith-Zaitlik *et al.*, 2022), was used as an outlier.