Phage vB_KmiS-Kmi2C infects members of the *Klebsiella oxytoca* complex and represents a novel genus of lytic bacteriophage

Fiona Newberry[1], Preetha Shibu[2]#, Thomas Smith-Zaitlik[1], Mohamed Eladawy[1,3], Anne L. McCartney[1], Lesley Hoyles[1]*, David Negus[1]*

[1] Department of Biosciences, Nottingham Trent University, United Kingdom

[2] Life Sciences, University of Westminster, United Kingdom

[3] Microbiology and Immunology Department, Faculty of Pharmacy, Mansoura University, Egypt

* **Corresponding authors:** David Negus, david.negus@ntu.ac.uk; Lesley Hoyles, lesley.hoyles@ntu.ac.uk

Present address: Berkshire and Surrey Pathology Services, Frimley Health NHS Trust, Wexham Park Hospital, Slough, United Kingdom.

Abbreviations: CARD, comprehensive antibiotic resistance database; KoC, *Klebsiella oxytoca* complex; PC, protein cluster; RBP, receptor binding protein; SD, standard deviation; TEM, transmission electron microscopy; TSA, tryptone soy agar; TSBG, tryptone soy broth supplemented with 1 % glucose; VC, viral cluster; VFDB, virulence factor database.

Keywords: multidrug-resistant, *Klebsiella michiganensis*, siphovirus, phage taxonomy.

Data availability: The genome sequence of vB_KmiS-Kmi2C has been deposited with GenBank under accession OP495736. Updated genome data for *K.*

26   *michiganensis* strains PS_Koxy1, PS_Koxy2 and PS_Koxy4 are available from

27   BioProject PRJNA562720.

ABSTRACT

Members of the *Klebsiella oxytoca* complex (KoC) are emerging opportunistic

pathogens responsible for causing a range of serious infections, especially in

healthcare settings. Increasing levels of antimicrobial resistance within the complex

could be circumvented by the therapeutic use of lytic bacteriophage to tackle

infections caused by KoC members. Here, we report the phenotypic and genomic

characterisation of phage vB_KmiS-Kmi2C, a siphovirus-like phage with lytic

properties against clinical isolates of the KoC and isolated on a GES-5-positive,

ST138 strain of *K. michiganensis*. Application of phage vB_KmiS-Kmi2C was found

to both prevent biofilm formation and disrupt established biofilms produced by

members of the KoC. The genome of vB_KmiS-Kmi2C comprises 42,234 bp and is

predicted to encode 55 genes. Comparative genomic and phylogenetic analyses

shows that vB_KmiS-Kmi2C shares little sequence similarity with other known

phages and represents a novel genus. Further gene-sharing network analysis using

vConTACT showed vB_KmiS-Kmi2C forms a viral cluster with 38 other known

phages, none of which were isolated using *Klebsiella* as a host.

INTRODUCTION

Bacteriophages (phages) are viruses of bacteria that infect their bacterial hosts for the purpose of replication. In the case of lytic phages, this process results in lysis and destruction of the host. The potential for applying the bactericidal action of lytic phages in a therapeutic setting was realised shortly after their discovery, with d'Herelle performing the first recorded clinical studies in 1918 [1]. Although interest in phage therapy diminished in Western countries following the introduction of antibiotics, the emergence of multidrug-resistant pathogens has resulted in a resurgence of interest.

Members of the *Klebsiella oxytoca* complex (KoC) are emerging as a serious health concern. The complex currently consists of several distinct phylogroups, representing *K. oxytoca* (Ko2), *K. michiganensis* (Ko1, Ko5), *K. grimontii* (Ko6), *K. huaxiensis* (Ko8), *K. pasteurii* (Ko4), *K. spallanzanii* (Ko3), and three unnamed novel species [2, 3]. Members are phylotyped based on the sequence of their chromosomally encoded β-lactamase (*bla*$_{oxy}$) gene [4]. KoC bacteria can cause a range of serious infections in both humans and animals and are of particular concern in healthcare settings where immunocompromised patients are predisposed to potential infection. Isolates have been associated with urinary tract infections, lower respiratory tract infections, septicaemia, and soft-tissue injuries [5–8]. Certain members of the complex (*K. oxytoca, K. pasteurii, K. grimontii, K. michiganensis*) also encode the kleboxymycin biosynthetic gene cluster associated with antibiotic-associated haemorrhagic colitis [9]. Worryingly, antibiotic resistance amongst KoC members is increasing and resistance mechanisms include extended-spectrum beta-lactamases and carbapenemases [10, 11].

New treatment options for tackling drug-resistant infections caused by KoC

isolates would be a welcome addition to our therapeutic armoury. This study aimed

to characterise the morphology, genome and host range of a lytic phage, vB_KmiS-

Kmi2C, isolated against a GES-5-encoding strain of *K. michiganensis* (PS_Koxy2)

[9]. The anti-biofilm properties of the phage were investigated to further determine

therapeutic utility with respect to preventing or resolving biofouling by KoC bacteria

such as that which occurs on medical devices, particularly indwelling catheters and

ventilators. Further extensive genomic and phylogenetic analyses of vB_KmiS-

Kmi2C reveal that it represents a novel genus of phage.

METHODS

Strain and cultivation information

Details of all strains included in this study can be found in **Table 1**. All strains

were grown on nutrient agar (Sigma Aldrich) unless stated otherwise. Nutrient broth

(Sigma Aldrich) was used for overnight cultures unless stated otherwise, incubated

aerobically at 37 °C. All media used for phage assays were supplemented with

$CaCl_2$ and $MgCl_2$ (both at final concentration of 0.5 mM). The study of anonymised

clinical isolates provided by the Nottingham University Hospitals NHS Trust (NUH)

Pathogen Bank was approved by NUH Research and Innovation (19MI001).

Generation of hybrid genome sequences

Cell pellets from 10-ml overnight cultures grown in nutrient broth were sent to

microbesNG (Birmingham, UK) for DNA extraction, library preparation and

sequencing according to the sequencing provider's strain-submission procedures

(refer to

https://microbesng.com/documents/24/MicrobesNG_Sequencing_Service_Methods_

v20210419.pdf for the protocol).

For Illumina sequencing, the Nextera XT Library Preparation Kit (Illumina, San

Diego, USA) was used and paired-end reads (HiSeq/NovaSeq; 2x250 bp were

generated for PS_Koxy1 (coverage 141×), PS_Koxy2 (coverage 52×) and

PS_Koxy4 (coverage 57×). Reads were adapter-trimmed using Trimmomatic 0.30

[12] with a sliding window quality cut-off of Q15. For Illumina/MinION hybrid

genomes, long-read genomic DNA libraries were prepared with Oxford Nanopore

SQK-RBK004 kit and/or SQK-LSK109 kit with Native Barcoding EXP-NBD104/114

(ONT, United Kingdom) using 400–500 ng of high-molecular-weight DNA. Barcoded

samples were pooled together into a single sequencing library and loaded into a

FLO-MIN106 (R.9.4.1) flow cell in a GridION (ONT, United Kingdom). *De novo*

genome assembly was performed using Unicycler v0.4.0 [13].

The genome assemblies returned to us by microbesNG were annotated in-

house using Bakta v1.6.1, database v4.0 [14]. RGI 6.0.0, CARD 3.2.5 was used to

predict antimicrobial resistance genes encoded within genomes [15]. COPLA and

plaSquid were used to characterise plasmids harboured by strains [16, 17].

Biolog GEN III MicroPlate assays

All strains were grown aerobically overnight on blood-free BUG agar (Biolog)

at 37 °C. For each assay, one to two colonies were used to inoculate Biolog

Inoculation Fluid B to a turbidity of 95 % (Biolog turbidimeter). Biolog Gen III plates

were inoculated according to the manufacturer's instructions, and incubated

aerobically at 37 °C for 22 h. Results were read at 540 nm using a BioTek Cytation |[3]

imaging reader spectrophotometer. All Biolog reagents and kits were purchased from

Technopath. Assays were carried out in triplicate for strains PS_Koxy1, PS_Koxy2

and PS_Koxy4.

Biofilm assays and their interpretation

Biofilm assays were performed as described previously [18–20]. In brief, a

single colony of each strain was used to inoculate 5 ml of tryptone soy broth

supplemented with 1 % glucose (TSBG). Cultures were incubated aerobically for 24

h at 37 °C without shaking. The overnight cultures were diluted to 1:100 using TSBG,

then aliquots (100 µl) of the diluted cultures were introduced into wells of a 96-well

plate. The plates were incubated aerobically for 24 h at 37 °C without shaking. Then,

the spent medium was carefully removed from each well. The wells were washed

three times with 200 µl sterile phosphate-buffered saline (pH 7.4; Oxoid) to remove

any non-adherent planktonic cells. The adherent cells were fixed by heat treatment

at 60 °C for 60 min to prevent widespread detachment of biofilms prior to dye

staining. The adhered biofilms were then stained by addition of 1 % (w/v) crystal

violet (150 µl per well) and the 96-well plate was left to incubate for 20 min. The

excess stain was then carefully removed from the wells and discarded. The 96-well

plate was then carefully rinsed with distilled water three times, and the plate was

inverted and left at room temperature until the wells were dry. The stained biofilms

were solubilised by adding 33 % (v/v) glacial acetic acid (Sigma Aldrich) to each well

(150 µl per well). After solubilisation of the stained biofilms, the OD_{540} was measured

and recorded for all samples using a BioTek Cytation |[3] imaging reader

spectrophotometer. *K. michiganensis* Ko14 and uninoculated medium were used as

negative controls in biofilm assays. Biological (n=3) and technical (n=4) replicates

were done for all strains. The mean of each isolate's OD quadruplicate readings

(OD_i) was calculated and compared with the control cut-off OD (OD_C), which was

defined as three standard deviations (SD) above the mean of the negative control

(3SD + mean). The amount of biofilm formed was scored as non-adherent $(OD_i \leq$

$OD_C)$, weakly adherent $(OD_C < OD_i \leq 2\ OD_C)$, moderately adherent $(2\ OD_C < OD_i \leq 4$

$OD_C)$ or strongly adherent $(4\ OD_C < OD_i)$.

Isolation of lytic phage and host range analysis

Filter-sterilised sewage samples (0.45 µm cellulose acetate filter; Millipore)

collected from mixed-liquor tanks at Mogden Sewage Treatment Works (March

2017) were screened against *K. michiganensis* strain PS_Koxy2 as described

previously [21]. Host range analysis was carried out as described previously [21].

After overnight incubation, overlay assay plates were inspected for lysis, with results

recorded according to a modification of [22]: ++, complete lysis; +, hazy lysis; 0, no

visible plaques.

Phage concentration and transmission electron microscopy (TEM)

The Vivaspin 20 50 kDa centrifugal concentrator (Cytiva) was used to

concentrate filter-sterilised propagated phage as described previously [21]. For TEM,

formvar/carbon-coated 200 mesh copper grids (Agar Scientific) were prepared via

glow discharge (10 mA, 10 s) using a Q150R ES sputter coater (Quorum

Technologies Ltd) and processed as described previously [21]. Samples were

visualised using a JEOL JEM-2100Plus (JEOL Ltd) TEM and an accelerating voltage

of 200 kV. Images were analysed and annotated using ImageJ

(https://imagej.net/Fiji).

Phage DNA extraction and sequencing

Phage DNA was extracted from concentrated phage lysate using the Qiagen

DNeasy Blood & Tissue Kit as described previously [21]. Sequence data were

generated on our in-house Illumina MiSeq platform, with the Nextera XT DNA library

preparation kit (Illumina) to produce fragments of approximately 500 bp. Fragmented

and indexed samples were run on the sequencer using a Micro flow cell with the

MiSeq Reagent Kit v2 (Illumina; 150-bp paired-end reads) following Illumina's

recommended denaturation and loading procedures.

Phage genome assembly and annotation

Quality of raw sequence data was assessed using FastQC v0.11.9. Reads

had a mean phred score above 30 and contained no adapter sequences, so data

were not trimmed. Genomes were assembled using SPAdes v3.15.4 [23] and

visualised using Bandage v0.8.1 [24]. Contamination and completeness of genomes

were determined using CheckV v0.8.1 (CheckV database v1.0) [25]. Gene

annotations were made using Prokka (v1.14.6) with the PHROGs database (v3) [26].

Genomes were screened for antimicrobial resistance genes using the Resistance

Gene Identifier (v5.2.0) of the Comprehensive Antibiotic Resistance Database

(CARD) (v3.1.4) [15], and for virulence genes using the Virulence Factor Database

(VFDB) (accessed: 6/6/2022) [27]. An annotated genome map was generated using

GenoPlotR (v0.8.11) and predicted protein function from the PHROGs database [28].

Comparative genome analysis

To generate a gene-sharing network, 21,903 phage genomes from the

INPHARED database (April 2022) and the query phage were clustered using

vConTACT2 (v0.9.22; default settings) and visualised in Cytoscape (v.3.9.1) [29, 30].

The viral cluster (VC) containing the query phage was determined and a proteomic

phylogenetic tree generated from all phage within the same VC using VipTree

(v1.1.2; default settings) [31]. *Klebsiella oxytoca* phage genomes vB_KmiM-2Dii

(accession: MZ707156) and vB_KmiM-4Dii (accession: MZ707157) were used as

outliers. An additional proteomic phylogenetic tree was generated from all phages

within the VC and three representative phage genomes from distantly related genera

Nonagvirus (JenK1; NC_029021.1), *Seuratvirus* (CaJan; NC_028776.1) and

Nipunavirus (NP1; NC_031058.1). The large terminase protein sequences were

used to generate a maximum likelihood phylogenetic tree with ClustalW (v.2.1) and

IQTree with 1000 bootstraps (v1.6.12) [32, 33]. IQModelFinder (v.1.4.2) was used

with IQTree to determine the model of best fit [34]. The percentage of shared

proteins within the VC was determined using all-versus-all BLASTP (≥30 % identity

and ≥50 % sequence coverage; v2.12.0) [35]. The intergenomic similarity between

phage within the same VC was determined using VIRIDIC (v1.0) [36].

ComplexHeatmap was used to generated all heatmaps [37].

Phage–biofilm assays

The titre of the phage stock was determined by plaque assay using the

double-layer agar technique. Briefly, phage vB_KmiS-Kmi2C was serially diluted in

phosphate-buffered saline (pH 7.4; Oxoid) and 100 µl of each phage dilution was

combined with 100 µl of an overnight culture of *K. michiganensis* PS_Koxy2 and 5 ml

of 0.6 % tryptone soy agar (TSA) supplemented with $CaCl_2$ and $MgCl_2$ both at a final

concentration of 1 mM. The mixture was gently swirled and poured onto solid TSA

plates. Plates were incubated overnight at 37 °C and pfu/ml determined by

enumeration of visible plaques.

The ability of vB_KmiS-Kmi2C to prevent and disrupt biofilms was examined

using a modification of a previously described protocol [38]. Hosts of vB_KmiS-

Kmi2C identified as moderately ($2\ OD_C < OD_i \leq 4\ OD_C$) adherent were included in

the assay. For prevention of biofilms, host cultures were incubated aerobically for 24

h at 37 °C without shaking in TSBG. Overnight cultures were diluted 1:100 with

TSBG and aliquots (100 µl) of diluted culture were introduced into wells of a 96-well

plate with or without phage vB_KmiS-Kmi2C (added to a final concentration of 4.5 x

10^8 pfu/ml to each test well). Plates were incubated without shaking for 24 h at 37

°C. Then, the supernatants were discarded, the biofilm of each well was washed to

remove planktonic cells and biofilms stained as described above.

To investigate the disruption of established biofilms, host cultures were grown

and prepared as described above prior to inoculating a 96-well plate. Plates were

incubated without shaking for 24 h at 37 °C to allow biofilms to form. Unattached

planktonic cells were carefully aspirated without disrupting the biomass. Phage

vB_KmiS-Kmi2C diluted in 100 µl TSBG was added to test wells (final titre of 4.5×10^8

pfu/ml) whereas control wells received only TSBG without phage. Plates were

incubated for a further 24 h at 37 °C without shaking. Supernatants were carefully

discarded; the biofilm of each well was washed to remove planktonic cells and

biofilms stained as described above. Biological (*n*=3) and technical (*n*=4) replicates

were completed for all strains.

RESULTS

Host range of phage vB_KmiS-Kmi2C

Phage vB_KmiS-Kmi2C was isolated from sewage water on *K. michiganensis* PS_Koxy2 (ST138), a multidrug-resistant, GES-5-positive isolate recovered from human urine [9]. The phage was screened against 56 clinical and 28 animal isolates representing a range of *Klebsiella* spp. (*K. michiganensis* n=49, *K. oxytoca* n=25, *K. grimontii* n=9, *K. huaxiensis* n=1) [9, 21]. It did not infect the closely related *K. michiganensis* strains PS_Koxy1 and PS_Koxy4 [9]. vB_KmiS-Kmi2C did not infect animal isolates of *Klebsiella* spp. [21], but did infect some of the clinical *K. oxytoca* (*n*=2) and *K. michiganensis* (*n*=4) strains within our extended in-house collection [21] (**Table 1**). From strain information provided by the Pathogen Bank (EUCAST testing) all these strains were resistant to amoxycillin, and sensitive to amikacin, ceftazidime, ceftazidime, ceftriaxone, cefuroxime, ciprofloxacin, ertapenem, gentamicin, meropenem, piptazobactam and trimethoprim. Strains Ko13, Ko22, Ko43 and Ko53 were sensitive to co-amoxiclav; Ko21 was resistant to co-amoxiclav; no data were supplied for Ko37 and co-amoxiclav. Phage vB_KmiS-Kmi2C did not exhibit depolymerase activity (i.e. it did not form haloes around plaques) on any of the strains that it infected.

Genotypic and phenotypic characterisation of the GES-positive strains

Our previous work had shown PS_Koxy1, PS_Koxy2 and PS_Koxy4 to be very similar based on the analysis of draft genome sequence data [9]. In this study, we generated hybrid assemblies of their genomes (**Table 2**).

The initial description of the strains' draft genomes [39] predicted they each harboured an identical IncQ 8,300-bp circular GES-5-positive plasmid, confirmed in

this study. PS_Koxy2 and PS_Koxy4 both carried an identical 76,870-bp circular

plasmid (replication initiation protein domain Rep_3, replicon type IncR). The 4,448-

bp plasmid carried by all three strains shared high identity with seven other plasmids,

harboured by *Salmonella enterica*, *Enterobacter* spp. and *K. michiganensis* (**Table**

3). A comparison (progressiveMauve, not shown) of the sequence of pPSKoxy2_4

with those of the other 4,448-bp plasmids in **Table 3** showed they shared 4,446

identical sites (99.98 % pairwise identity) across their length. A PlasmidMLST search

returned no hits for the plasmid sequence. It was found to be PTU-E3 (score 96.97

%) by COPLA, exceeding the 90 % threshold to validate the plasmid taxonomic unit

assignment [16], and to belong to mobility group MOBP1 with a Col440I replicon

type by plaSquid [17]. The plasmid was predicted by Bakta to encode six genes

(MobA, MobC, two YgdI/YgdR family lipoproteins, two hypothetical proteins), an

origin of replication, *oriT* and a non-coding RNAI (**Figure 1a**). UniProt BLASTP

analyses showed MobA shared 55 % identity (across 517 aa) with the DNA relaxase

MbeA of *Escherichia coli* (reviewed UniProt record P13658); MobC shared 63.6 %

identity (across 115 aa) with the mobilisation protein MbeC of *E. coli* (reviewed

UniProt record P13657). The 39-aa hypothetical protein was predicted by I-TASSER

[40] to be an alpha helix, while the 122-aa hypothetical protein was found to include

a transmembrane segment (HHPRED, UniRef30 [41]) that had limited structural

homology with only the *Escherichia coli* signal transduction protein PmrD

(PDB:4HN7) and the Vps26 dimer region of a fungal vacuolar protein sorting-

associated protein (PDB:7BLQ).

PS_Koxy4 could be differentiated from PS_Koxy1 and PS_Koxy2 by

phenotypic traits determined using the Biolog GEN III system (**Table 3**). PS_Koxy1

and PS_Koxy2 were resistant to the tetracycline antibiotic minocycline; both were

predicted to encode the tetracycline efflux pump *tet(A)* on incompletely assembled

plasmids (contig_3 PS_Koxy1, predicted to be FIIK by PlasmidMLST; contig_2

PS_Koxy2, predicted to be IncA/C ST12 by PlasmidMLST), but this gene was absent

from the genome of PS_Koxy4, with the strain sensitive to minocycline (**Figure 1b**,

Table 3).

Biofilm-forming ability of strains used in this study

We determined the biofilm-forming ability of strains used in this study as

described previously [38]. Our results show a wide range of biofilm-forming abilities

among the strains (**Figure 1a**; **Supplementary Table 1**), yet there was no difference

in the virulence factor profiles of PS_Koxy1, PS_Koxy2 and PS_Koxy4 according to

a VFDB [27] analysis (**Supplementary Table 2**). Strains *K. michiganensis*

PS_Koxy1, *K. michiganensis* Ko13, *K. michiganensis* Ko21 and *K. michiganensis*

Ko22 were identified as weakly adherent ($OD_C < OD_i \leq 2\ OD_C$). Strains *K.*

michiganensis PS_Koxy2, *K. michiganensis* PS_Koxy4, *K. oxytoca* Ko37, *K. oxytoca*

Ko43 and *K. oxytoca* Ko53 were identified as moderately adherent ($2\ OD_C < OD_i \leq 4$

OD_C). *K. michiganensis* Ko14 was used as a non-adherent negative control ($OD_i \leq$

OD_C).

Capacity of phage vB_KmiS-Kmi2C to prevent and disrupt biofilms

Phage vB_KmiS-Kmi2C was found to be highly effective at both preventing

(**Figure 1b**) and disrupting (**Figure 1c**) biofilms formed by isolates identified as

susceptible to lysis. Presence of vB_KmiS-Kmi2C was found to reduce biofilm

formation for all strains included in the study. This reduction was found to be

significant ($P < 0.01$) for three of the isolates tested (PS_Koxy2, Ko37 and Ko53)

compared to no-phage controls. Addition of vB_KmiS-Kmi2C to pre-established

biofilms also resulted in biofilm disruption, seen as a reduction in measured biofilm

compared to non-phage-treated controls. Biofilm disruption was determined to be

significant ($P < 0.05$) for isolates PS_Koxy2 and Ko37 compared to non-phage

treated controls.

Morphology of phage vB_KmiS-Kmi2C

TEM showed vB_KmiS-Kmi2C to have a siphovirus-like morphology (**Figure**

2). It had a long, non-contractile tail. The capsid diameter was 59.6 nm (SD 2.3 nm),

the tail and baseplate were 157.8 nm in length (SD 5.1 nm), and the phage had a

total length of 218.11 nm (SD 4.8) (n=3 measurements).

Genome of phage vB_KmiS-Kmi2C

The genome of phage vB_KmiS-Kmi2C was assembled into a single contig,

with 100 % completeness according to CheckV [25]. Assembly of the phage was

confirmed using visualisation with Bandage (**Supplementary Figure 1**). It comprised

42,234 bp (816 $\times$ coverage) and was predicted to encode 55 genes (upper panel

Figure 3). The genome is arranged into functional modules as typically seen in

phage genomes: replication/regulation, viral structure, DNA packaging and lysis.

Interesting genomic features of the genome included a Sak4-like ssDNA annealing

protein with single-strand DNA-binding protein, a Cas4-domain exonuclease, a holin

and a Rz-like spanin.

Analysis using NCBI BLASTN showed the genome of vB_KmiS-Kmi2C

shared 85.6 % identity (80 % query coverage) with a phage metagenome-assembled

genome (accession OP072809) recovered from human faeces in Japan [42].

Preliminary analysis of the genome sequence using the online version of ViPTree suggested the phage was novel (not shown). Further support for this came from genomic similarity analysis (**Supplementary Figure 2**) and analyses of protein–protein network data using vConTACT (**Figure 4**). According to vConTACT, vB_KmiS-Kmi2C phage formed a VC with 38 other known phage from a range of genera: 24 *Septimatrevius*, 6 *Lokivirus*, 4 *Titanvirus*, 1 *Pradovirus*, 1 *Kilunavirus* and 2 unclassified. The VC did not contain any other phage isolated using *Klebsiella* and the majority of phage had *Pseudomonas* as an assigned host (**Figure 5**). The intergenomic similarity within the VC ranged from 100 % to 2 %, with the highest similarity noted between phage within the same genus (**Supplementary Figure 2**). vB_KmiS-Km2iC showed the highest genomic similarity (18.8 %) to *Septimatrevirus* vB_Pae-Kakheti25, a *Pseudomonas* phage.

The vConTACT output was used to determine the shared protein clusters (PCs) within the VC compared to phage vB_KmiS-Kmi2C (lower panel **Figure 3**). Of the 53 PCs identified in vB_KmiS-Kmi2C, 14 were shared between all phage within the VC (007793 Cas4-domain exonuclease; 006166 DNA helicase; 006824 DNA polymerase processivity factor; 004149 DNA polymerase; 005059 tail length tape measure protein; 005663 Neck1 protein; 005850 minor tail protein; 008503 tail terminator; 005899 head-tail adaptor Ad1; 003930 head scaffolding protein; 002520 portal protein; 005106 terminase large subunit; 007093 Sak4-like ssDNA annealing protein; 008553 ssDNA binding protein). Seven of the shared PCs were associated with viral structure and six were associated with viral replication/regulation. vB_KmiS-Kmi2C shared the most PCs with *Erwinia* phage AH03, including the majority of the viral structure genome module.

Protein-based phylogenetic analysis with VipTree showed phages within the

VC grouped according to the assigned genus, with vB_Kmi2S_Kmi2C not within the

defined genus groups. vB_KmiS-Kmi2C showed the closest relationship to *Erwinia*

phage AH03, as previously noted in the lower panel of **Figure 3**. A representative

species from three closely related genera were included in a further protein-based

phylogenetic analysis to determine if vB_KmiS-Kmi2C grouped with genera outside

the VC defined by vConTACT: *Nipunavirus* NP1 (NC_031058), *Nonagavirus* JenK1

(NC_024146) and *Seuratvirus* Cajan (NC_028776) (**Supplementary Figure 3**).

Analysis of the shared protein percentage within the viral cluster was consistent with

the intergenomic similarity and protein phylogenetic analysis (**Supplementary**

Figure 4). The phage belonging to the same genera shared the highest percentage

of proteins. vB_KmiS-Kmi2C shared the highest percentage of proteins with

Pseudomonas phage *Septimatrevirus* KP1 and *Erwinia* phage AH03. This confirmed

that vB_KmiS-Kmi2C does not belong to any currently defined genus of phage.

DISCUSSION

There has been a significant research effort into the investigation of phages

for the treatment of *K. pneumoniae* infections and we have recently authored an

extensive review on phages of *Klebsiella* spp. [43]. However, there are relatively few

studies documenting the isolation and characterisation of lytic phages against

members of the KoC [21, 44, 45], an emerging group of clinical and veterinary

pathogens with growing antibiotic resistance. As such, we set out to identify phages

with the capacity to lyse clinically relevant members of the KoC. We have previously

reported on the characterisation of two such phages with lytic activity against

numerous species within the KoC [21]. Here, we report on the characterisation of

phage vB_KmiS-Kmi2C isolated against a carbapenem-resistant isolate of *K.*

michiganensis (PS_Koxy2) [9], for which we generated a hybrid genome assembly

and phenotypic data to allow it to be differentiated from closely related strains

(PS_Koxy1, PS_Koxy4) from our in-house collection.

There were clear differences in the biofilm-forming abilities of the three GES-5

-positive strains, yet all strains encoded the same virulence factors (**Supplementary**

Table 2). We did not compare protein sequences of the virulence factors across the

strains. It is acknowledged that our understanding of KoC phenotypes is poor [3],

and our future work will look to address how differences in biofilm-forming abilities of

KoC members and their genetic differences contribute to survival and potentially

pathogenicity.

While the three strains all carried two small (4,448 and 8,300 bp) plasmids,

only PS_Koxy2 and PS_Koxy4 carried a 76,860-bp plasmid. This and the other

larger plasmids carried by the strains will be described elsewhere as part of ongoing

work. The 8,300-bp IncQ plasmid carried by the three strains has been described

elsewhere [39]. While the 4,448-bp plasmid (mobility group MOBP1, replicon type

Col440I) has been detected in a strain of *K. michiganensis* previously [46], it has not

been described in detail. Based on its gene content (specifically, MobA/MobC), we

predict the plasmid to be conjugative plasmid. The MobA and MobC functional

homologues MbeA and MbeC, respectively, are required for efficient mobilisation of

the ColE1 plasmid [47].

Host range analysis and anti-biofilm properties

Understanding the lytic profile of any individual phage is important for its

potential deployment in a clinical setting, either in an isolated form or for inclusion in

a phage cocktail that may increase host range and potentially suppress phage

resistance [48]. Ordinarily, phage host range is narrow, sometimes down to the strain

level. In accordance with this, we found that vB_KmiS-Kmi2C exhibits a relatively

narrow host range, lysing six clinical isolates from the KoC out of a panel of 84

isolates representing a range of clinical and animal *Klebsiella* spp.

To further characterise the therapeutic potential of vB_KmiS-Kmi2C,

experiments were performed to determine the phage's capacity to prevent and

disrupt biofilms formed by the clinical isolates susceptible to lysis in our host range

analysis. Biofilm formation can reduce the efficacy of antibiotic chemotherapy and

contributes to the successful colonisation of wounds and indwelling medical devices

[49]. Therefore, phages with anti-biofilm properties may make particularly attractive

therapeutics. We found that vB_KmiS-Kmi2C is highly effective at both preventing

biofilm formation and disrupting established biofilms. It is likely that prevention of

biofilm formation is largely due to the lytic action of the phage, resulting in cell death

and the prevention of biofilm formation. However, established biofilms can be

extremely recalcitrant to disruption. This is likely due to the complex nature of mature

biofilms which are typically composed of numerous macromolecules including,

polysaccharides, proteins, nucleic acids and metabolically inactive cells [50]. Phage-

mediated disruption of *K. michiganensis* biofilms has been previously reported in the

literature [51]. The lytic phage KMI8 was shown to disrupt a *K. michiganensis* mono-

biofilm produced by a strain expressing the polysaccharide capsule KL70 locus. It

was hypothesised this antibiofilm activity was largely due to the activity of a potential

phage-associated depolymerase, identified as a putative endosialidase encoded

within the KMI8 genome. In the present study, we found no evidence of

depolymerase activity associated with vB_KmiS-Kmi2C. Plaques on susceptible host

lawns did not show evidence of haloes which are typically indicative of depolymerase

activity. Furthermore, our bioinformatic analysis did not identify any genes

associated with depolymerase activity. Studies are therefore ongoing to determine

the mechanism of biofilm disruption used by vB_KmiS-Kmi2C.

Phage vB_KmiS-Kmi2C represents a new genus

Whole-genome sequencing is important for the identification of features

relevant to phage therapy and for aiding taxonomic classification. Genomic analysis

using the resistance gene identifier tool available through the comprehensive

antibiotic resistance database did not identify the presence of known antimicrobial

resistance mechanisms within the genome of vB_KmiS-Kmi2C. The presence of

such genes would suggest the potential to transfer antibiotic resistance markers

between bacteria if vB_KmiS-Kmi2C were to be applied therapeutically in a

polymicrobial setting.

Initial genome sequence analysis of vB_KmiS-Kmi2C using BLASTN and

ViPTree suggested that our phage was novel, sharing little sequence similarity with

other deposited phage genomes. To aid with taxonomy, we employed the gene-

sharing network analysis tool vConTACT2. Phage taxonomic assignment using

whole genome gene-sharing profiles has been shown to be highly accurate; a recent

study showed that vConTACT2 produces near-identical replication of existing genus-

level viral taxonomy assignments from the International Committee on Taxonomy of

Viruses (ICTV) [52]. Our analysis using vConTACT2 and the INPHRARED database

of phage genomes identified that vB_KmiS-Kmi2C shared a VC with 38 other

sequenced phages. The VC was polyphyletic, containing phage from at least five

different genera, suggesting that vB_KmiS-Km2iC did not group with any existing

genus. Recently published guidelines have suggested that for genome-based phage

taxonomy, assignment to taxonomic ranks should be based on nucleotide sequence

identity across entire genomes [35, 53]. These guidelines suggest any two phages

belong to the same species if they are more than 95 % identical across their genome

and belong to the same genus if they are more than 70 % identical. We used

VIRIDIC to determine the intergenomic similarity between vB_KmiS-Kmi2C and the

38 other phages identified within the VC. Our findings show that based on these

criteria, vB_KmiS-Kmi2C cannot be assigned to any current species or genus as it

shares only 18.8 % genomic similarity to *Septimatrevirus* vB_Pae-Kakheti25, a

Pseudomonas phage, the closest nucleotide sequence relative we could identify.

Most of the phages within the VC (21/38) were identified as infecting bacteria

belonging to the genus *Pseudomonas*. According to the GenBank accession details

associated with the genomes of these phages, they were all isolated-on strains of *P.*

aeruginosa. A preliminary screen of vB_KmiS-Kmi2C against a panel of 11 *P.*

aeruginosa isolates did not identify any lytic activity against members of this species

(data not shown).

In addition to identifying phages belonging to a VC, vConTACT2 can also

identify proteins shared by members within the VC. Our analysis found that 14

proteins were shared between all phages within the VC, of which seven were

associated with viral structure and six were associated with viral

replication/regulation. At least 35 of the phages within the VC are from genera with

siphovirus-like morphology. It is therefore unsurprising that phages within the cluster

share several proteins related to general structure/morphology.

Phage vB_KmiS-Kmi2C is the only member of the VC known to infect

members of the genus *Klebsiella*. Host-cell tropism is often determined by receptor-

binding proteins (RBPs) which mediate recognition and attachment to host cells [54].

RBPs are typically identified as phage tail fibres or spikes [55]. The genomic region

of vB_KmiS-Kmi2C encoding hypothetical tail fibre proteins can be seen in **Figure 3**

and comprises proteins 088411, 095417, 028210, 033709, 031062 and 060189.

Unannotated proteins within this region were confirmed to share homology with

phage tail fibre proteins by HHPred analysis using an e-value cut-off of $>10^{-5}$. The

tail fibre proteins of vB_KmiS-Kmi2C are not widely present within the VC and are

typically shared by only one or two phages within the cluster. This may explain the

host tropism of vB_KmiS-Kmi2C compared to other members of the VC but further

experimentation will be required to determine this. Protein 005059 is a predicted tail

length tape measure protein; such proteins are involved in determining tail length

and participate in DNA injection into the host-cells and are not involved in host

recognition [56].

Our additional protein-based phylogenetic analyses, including phages from

outside the VC, highlighted that vB_Kmi2S_Kmi2C does not group with any genera

included in the analyses and as such does not belong to any currently defined genus

of phage.

Phages are recognised as the most abundant biological entities on the planet

[57] and exhibit extensive biological diversity [58]. With an estimated 10^{31} phage

virions in the biosphere [59] and 14,244 complete phage genomes sequenced as of

2021 [29], it is clear we have characterised a relatively small number of distinct

phages. However, with the cost of genome sequencing falling and the capacity to

assemble phage genomes from metagenome sequence data, it is likely that the

number of novel phages identified will increase rapidly. The classification of phages

representing entirely novel genera or families/sub-families poses a challenge for

taxonomy, especially when new isolates share little sequence similarity with

previously characterised phages. Previously, tailed phages were assigned to a family

based on morphology, with the families *Podoviridae*, *Myoviridae* and *Siphoviridae*

belonging to the order *Caudovirales*. However, the increase in available phage

sequence data has shown that these families are paraphyletic. It has, therefore,

been suggested that they are abolished, and that phage morphology should no

longer play a role in classification [35]. Based on recently updated guidelines for the

classification of phage based on nucleotide sequence identity, and subsequent

protein based phylogenetic analyses, we show here that phage vB_KmiS-Kmi2C

represents a new genus.

ACKNOWLEDGEMENTS

Conceptualisation: FN, PS, LH, DN. Data curation: all authors. Formal

analysis: FN, TSZ, PS, ME, ALM, LH, DN. Funding acquisition: PS, ALM, LH.

Investigation: FN, TSZ, PS, ME, ALM, LH, DN. Methodology: FN, ME, LH, DN.

Resources: ALM, LH, DN. Supervision: ALM, LH, DN. Visualisation: FN, LH, DN.

Writing – original draft: FN, ME, LH, DN. Writing – reviewing and editing: all authors.

The authors declare that there are no conflicts of interest.

PS was in receipt of an IBMS Research Grant (project title "Isolation of lytic

bacteriophages active against antibiotic-resistant *Klebsiella pneumoniae*"). Imperial

Health Charity is thanked for contributing to registration fees for the Professional

Doctorate studies of PS. ME is supported by the Egyptian Government – Ministry of

Higher Education/Egyptian Cultural and Education Bureau. This work used

computing resources funded by the Research Contingency Fund of the Department

of Biosciences, Nottingham Trent University (NTU). TSZ completed this work as part

of an MRes degree at NTU.

REFERENCES

1.   **d'Herelle F**. Sur le rôle du microbe filtrant bactériophage dans la dysentérie bacillaire. *Comptes*
Rendus Académie Sci 1918;167:970–972.

2.   **Merla C, Rodrigues C, Passet V, Corbella M, Thorpe HA, *et al.*** Description of Klebsiella
spallanzanii sp. nov. and of Klebsiella pasteurii sp. nov. *Front Microbiol* 2019;10:2360.

3.   **Yang J, Long H, Hu Y, Feng Y, McNally A, *et al.*** Klebsiella oxytoca Complex: Update on
Taxonomy, Antimicrobial Resistance, and Virulence. *Clin Microbiol Rev* 2022;35:e0000621.

4.   **Cosic A, Leitner E, Petternel C, Galler H, Reinthaler FF, *et al.*** Variation in Accessory Genes
Within the *Klebsiella oxytoca* Species Complex Delineates Monophyletic Members and Simplifies
Coherent Genotyping. *Front Microbiol* 2021;12:692453.

5.   **Shakya P, Shrestha D, Maharjan E, Sharma VK, Paudyal R**. ESBL Production Among *E. coli* and
Klebsiella spp. Causing Urinary Tract Infection: A Hospital Based Study. *Open Microbiol J*
2017;11:23–30.

6.   **Sohn W-I, Seo BF, Jung S-N**. Forehead Abscess Caused by *Klebsiella oxytoca* With Undiagnosed
Type 2 Diabetes: *J Craniofac Surg* 2012;23:247–249.

7.   **Paasch C, Wilczek S, Strik MW**. Liver abscess and sepsis caused by *Clostridium perfringens* and
Klebsiella oxytoca. Int J Surg Case Rep 2017;41:180–183.

8.   **Lee SH, Ruan S-Y, Pan S-C, Lee T-F, Chien J-Y, *et al.*** Performance of a multiplex PCR pneumonia
panel for the identification of respiratory pathogens and the main determinants of resistance
from the lower respiratory tract specimens of adult patients in intensive care units. *J Microbiol*
Immunol Infect 2019;52:920–928.

9.   **Shibu P, McCuaig F, McCartney AL, Kujawska M, Hall LJ, *et al.*** Improved molecular
characterization of the *Klebsiella oxytoca* complex reveals the prevalence of the kleboxymycin
biosynthetic gene cluster. *Microb Genomics* 2021;7:000592.

10.   **White L, Hopkins KL, Meunier D, Perry CL, Pike R, *et al.*** Carbapenemase-producing
Enterobacteriaceae in hospital wastewater: a reservoir that may be unrelated to clinical isolates.
J Hosp Infect 2016;93:145–151.

11.   **Lowe C, Willey B, O'Shaughnessy A, Lee W, Lum M, *et al.*** Outbreak of Extended-Spectrum β-
Lactamase–producing *Klebsiella oxytoca* Infections Associated with Contaminated Handwashing
Sinks1. *Emerg Infect Dis* 2012;18:1242–1247.

12.   **Bolger AM, Lohse M, Usadel B**. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinforma Oxf Engl 2014;30:2114–2120.

13.   **Wick RR, Judd LM, Gorrie CL, Holt KE**. Unicycler: Resolving bacterial genome assemblies from
short and long sequencing reads. *PLoS Comput Biol* 2017;13:e1005595.

14. **Schwengers O, Jelonek L, Dieckmann MA, Beyvers S, Blom J, *et al.*** Bakta: rapid and
standardized annotation of bacterial genomes via alignment-free sequence identification.
Microb Genomics;7. Epub ahead of print November 2021. DOI: 10.1099/mgen.0.000685.

15. **Alcock BP, Raphenya AR, Lau TTY, Tsang KK, Bouchard M, *et al.*** CARD 2020: antibiotic
resistome surveillance with the comprehensive antibiotic resistance database. *Nucleic Acids Res*
2020;48:D517–D525.

16. **Redondo-Salvo S, Bartomeus-Peñalver R, Vielva L, Tagg KA, Webb HE, *et al.*** COPLA, a
taxonomic classifier of plasmids. *BMC Bioinformatics* 2021;22:390.

17. **Giménez M, Ferrés I, Iraola G**. Improved detection and classification of plasmids from
circularized and fragmented assemblies. *bioRxiv*. Epub ahead of print 2022. DOI:
10.1101/2022.08.04.502827.

18. **Stepanovic S, Vukovic D, Dakic I, Savic B, Svabic-Vlahovic M**. A modified microtiter-plate test
for quantification of staphylococcal biofilm formation. *J Microbiol Methods* 2000;40:175–179.

19. **Eladawy M, El-Mowafy M, El-Sokkary MMA, Barwa R**. Antimicrobial resistance and virulence
characteristics in ERIC-PCR typed biofilm forming isolates of P. aeruginosa. *Microb Pathog*
2021;158:105042.

20. **Merritt JH, Kadouri DE, O'Toole GA**. Growing and analyzing static biofilms. *Curr Protoc Microbiol*
2005;Chapter 1:Unit 1B.1.

21. **Smith-Zaitlik T, Shibu P, McCartney AL, Foster G, Hoyles L, *et al.*** Extended genomic analyses of
the broad-host-range phages vB_KmiM-2Di and vB_KmiM-4Dii reveal slopekviruses have highly
conserved genomes. *Microbiology* 2022;168:001247.

22. **Haines MEK, Hodges FE, Nale JY, Mahony J, van Sinderen D, *et al.*** Analysis of Selection
Methods to Develop Novel Phage Therapy Cocktails Against Antimicrobial Resistant Clinical
Isolates of Bacteria. *Front Microbiol* 2021;12:613529.

23. **Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, *et al.*** SPAdes: a new genome
assembly algorithm and its applications to single-cell sequencing. *J Comput Biol J Comput Mol*
Cell Biol 2012;19:455–477.

24. **Wick RR, Schultz MB, Zobel J, Holt KE**. Bandage: interactive visualization of de novo genome
assemblies. *Bioinforma Oxf Engl* 2015;31:3350–3352.

25. **Nayfach S, Camargo AP, Schulz F, Eloe-Fadrosh E, Roux S, *et al.*** CheckV assesses the quality and
completeness of metagenome-assembled viral genomes. *Nat Biotechnol* 2021;39:578–585.

26. **Terzian P, Olo Ndela E, Galiez C, Lossouarn J, Pérez Bucio RE, *et al.*** PHROG: families of
prokaryotic virus proteins clustered using remote homology. *NAR Genomics Bioinforma*
2021;3:lqab067.

27. **Liu B, Zheng D, Jin Q, Chen L, Yang J**. VFDB 2019: a comparative pathogenomic platform with an
interactive web interface. *Nucleic Acids Res* 2019;47:D687–D692.

28. **Guy L, Kultima JR, Andersson SGE**. genoPlotR: comparative gene and genome visualization in R.
Bioinforma Oxf Engl 2010;26:2334–2335.

29. **Cook R, Brown N, Redgwell T, Rihtman B, Barnes M, *et al.*** INfrastructure for a PHAge REference
Database: Identification of Large-Scale Biases in the Current Collection of Cultured Phage
Genomes. *PHAGE* 2021;2:214–223.

30. **Bolduc B, Jang HB, Doulcier G, You Z-Q, Roux S, *et al.*** vConTACT: an iVirus tool to classify
double-stranded DNA viruses that infect Archaea and Bacteria. *PeerJ* 2017;5:e3243.

31. **Nishimura Y, Yoshida T, Kuronishi M, Uehara H, Ogata H, *et al.*** ViPTree: the viral proteomic
tree server. *Bioinforma Oxf Engl* 2017;33:2379–2380.

32. **Thompson JD, Higgins DG, Gibson TJ**. CLUSTAL W: improving the sensitivity of progressive
multiple sequence alignment through sequence weighting, position-specific gap penalties and
weight matrix choice. *Nucleic Acids Res* 1994;22:4673–4680.

33. **Minh BQ, Nguyen MAT, von Haeseler A**. Ultrafast Approximation for Phylogenetic Bootstrap.
Mol Biol Evol 2013;30:1188–1195.

34. **Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS**. ModelFinder: fast model
selection for accurate phylogenetic estimates. *Nat Methods* 2017;14:587–589.

35. **Turner D, Kropinski AM, Adriaenssens EM**. A Roadmap for Genome-Based Phage Taxonomy.
Viruses 2021;13:506.

36. **Moraru C, Varsani A, Kropinski AM**. VIRIDIC—A Novel Tool to Calculate the Intergenomic
Similarities of Prokaryote-Infecting Viruses. *Viruses* 2020;12:1268.

37. **Gu Z, Eils R, Schlesner M**. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinforma Oxf Engl* 2016;32:2847–2849.

38. **Taha OA, Connerton PL, Connerton IF, El-Shibiny A**. Bacteriophage ZCKP1: A Potential
Treatment for Klebsiella pneumoniae Isolated From Diabetic Foot Patients. *Front Microbiol*
2018;9:2127.

39. **Ellington MJ, Davies F, Jauneikaite E, Hopkins KL, Turton JF, *et al.*** A multi-species cluster of
GES-5 carbapenemase producing Enterobacterales linked by a geographically disseminated
plasmid. *Clin Infect Dis Off Publ Infect Dis Soc Am*. Epub ahead of print 20 November 2019. DOI:
10.1093/cid/ciz1130.

40. **Zhang Y**. I-TASSER server for protein 3D structure prediction. *BMC Bioinformatics* 2008;9:40.

41. **Zimmermann L, Stephens A, Nam S-Z, Rau D, Kübler J, *et al.*** A Completely Reimplemented MPI
Bioinformatics Toolkit with a New HHpred Server at its Core. *J Mol Biol* 2018;430:2237–2243.

42. **Nishijima S, Nagata N, Kiguchi Y, Kojima Y, Miyoshi-Akiyama T, *et al.*** Extensive gut virome
variation and its associations with host and environmental factors in a population-level cohort.
Nat Commun 2022;13:5252.

43. **Herridge WP, Shibu P, O'Shea J, Brook TC, Hoyles L**. Bacteriophages of Klebsiella spp., their
diversity and potential therapeutic uses. *J Med Microbiol* 2020;69:176–194.

44. **Townsend EM, Kelly L, Gannon L, Muscatt G, Dunstan R, *et al.*** Isolation and Characterization of
Klebsiella Phages for Phage Therapy. *PHAGE* 2021;2:26–42.

45. **Kęsik-Szeloch A, Drulis-Kawa Z, Weber-Dąbrowska B, Kassner J, Majkowska-Skrobek G, *et al.***
Characterising the biology of novel lytic bacteriophages infecting multidrug resistant *Klebsiella*
pneumoniae. *Virol J* 2013;10:100.

46. **AbuOun M, Jones H, Stubberfield E, Gilson D, Shaw LP, *et al.*** A genomic epidemiological study
shows that prevalence of antimicrobial resistance in Enterobacterales is associated with the
livestock host, as well as antimicrobial usage. *Microb Genomics* 2021;7:000630.

47. **Varsaki A, Lamb HK, Eleftheriadou O, Vandera E, Thompson P, *et al.*** Interaction between
relaxase MbeA and accessory protein MbeC of the conjugally mobilizable plasmid ColE1. *FEBS*
Lett 2012;586:675–679.

48. **Yang Y, Shen W, Zhong Q, Chen Q, He X, *et al.*** Development of a Bacteriophage Cocktail to
Constrain the Emergence of Phage-Resistant *Pseudomonas aeruginosa*. *Front Microbiol*
2020;11:327.

49. **Wang G, Zhao G, Chao X, Xie L, Wang H**. The Characteristic of Virulence, Biofilm and Antibiotic
Resistance of Klebsiella pneumoniae. *Int J Environ Res Public Health* 2020;17:6278.

50. **Chhibber S, Nag D, Bansal S**. Inhibiting biofilm formation by Klebsiella pneumoniae B5055 using
an iron antagonizing molecule and a bacteriophage. *BMC Microbiol* 2013;13:174.

51. **Ku H, Kabwe M, Chan HT, Stanton C, Petrovski S, *et al.*** Novel Drexlerviridae bacteriophage
KMI8 with specific lytic activity against Klebsiella michiganensis and its biofilms. *PLOS ONE*
2021;16:e0257102.

52. **Bin Jang H, Bolduc B, Zablocki O, Kuhn JH, Roux S, *et al.*** Taxonomic assignment of uncultivated
prokaryotic virus genomes is enabled by gene-sharing networks. *Nat Biotechnol* 2019;37:632–
639.

53. **Adriaenssens E, Brister JR**. How to Name and Classify Your Phage: An Informal Guide. *Viruses*
2017;9:E70.

54. **Dams D, Brøndsted L, Drulis-Kawa Z, Briers Y**. Engineering of receptor-binding proteins in
bacteriophages and phage tail-like bacteriocins. *Biochem Soc Trans* 2019;47:449–460.

55. **Nobrega FL, Vlot M, de Jonge PA, Dreesens LL, Beaumont HJE, *et al.*** Targeting mechanisms of
tailed bacteriophages. *Nat Rev Microbiol* 2018;16:760–773.

56. **Mahony J, Alqarni M, Stockdale S, Spinelli S, Feyereisen M, *et al.*** Functional and structural
dissection of the tape measure protein of lactococcal phage TP901-1. *Sci Rep* 2016;6:36667.

57. **Clokie MR, Millard AD, Letarov AV, Heaphy S**. Phages in nature. *Bacteriophage* 2011;1:31–45.

58. **Hendrix RW**. Bacteriophages: Evolution of the Majority. *Theor Popul Biol* 2002;61:471–480.

59. **Hatfull GF**. Dark Matter of the Biosphere: the Amazing World of Bacteriophage Diversity. *J Virol*
2015;89:8107–8110.

60. **Webb HE, Kim JY, Tagg KA, de la Cruz F, Peñil-Celis A, *et al.*** Genome Sequences of 18
Salmonella enterica Serotype Hadar Strains Collected from Patients in the United States.
Microbiol Resour Announc;11:e00522-22.

61. **Yachison CA, Yoshida C, Robertson J, Nash JHE, Kruczkiewicz P,** *et al.* The Validation and
Implications of Using Whole Genome Sequencing as a Replacement for Traditional Serotyping
for a National Salmonella Reference Laboratory. *Front Microbiol* 2017;8:1044.

Table 1. *K. oxytoca* complex strains used in this study

Strain *	Species	Source	Reference
PS_Koxy1	*K. michiganensis*	Throat swab	[9]
PS_Koxy2 §#	*K. michiganensis*	Urine	[9]
PS_Koxy4	*K. michiganensis*	Rectal swab	[9]
Ko13 §	*K. michiganensis*	Blood culture	[21]
Ko14	*K. michiganensis*	Blood culture	[21]
Ko21 §	*K. michiganensis*	Perfusion fluid	[21]
Ko22 §	*K. michiganensis*	Perfusion fluid	[21]
Ko37 §	*K. oxytoca*	Blood culture	[21]
Ko43 §	*K. michiganensis*	Bile fluid	[21]
Ko53 §	*K. oxytoca*	Blood culture	[21]

* Ko prefix indicates strains came from the Pathogen Bank of Queen's Medical

Centre Nottingham, UK.

§ Infected by phage vB_KmiS-Kmi2C.

Strain on which phage vB_KmiS-Kmi2C was isolated; all other strains used in host

range analysis but not infected by the phage are described in [21].

Table 2. Summary statistics for hybrid genome assemblies of the GES-5-positive

strains

Strain (accession)	Contigs	Size (bp)	Status *	N50	CDS	Plasmids (RIP; MOB; REP; PTU) #
PS_Koxy1 (GCA_014050555.2)	7	6,392,093	CCC; 4 CCPs; ? incomplete plasmid	6,023,296	5,952	pPSKoxy1_1 (RepA_C; MOBH; IncAC, IncN; –) pPSKoxy1_2 (Rep_3; –; IncR; –) pPSKoxy1_3 [–; MOBP1; –; PTU-Q2 (IncQ)] pPSKoxy1_4 (–; MOBP1; Col440I; PTU-E3)
PS_Koxy2 (GCA_014050535.2)	10	6,357,486	Draft chromosome; 4 CCPs; ? incomplete plasmid	5,917,750	5,910	pPSKoxy2_1 (IncFII_repA, IncFII_repA, RepA_C, RepB–RCR_reg; MOBF, MOBH; IncFII_1_pKP91, IncAC, IncN; –) pPSKoxy2_2 (Rep_3; –; IncR; –) pPSKoxy2_3 [–; MOBP1; –; PTU-Q2 (IncQ)] pPSKoxy2_4 (–; MOBP1; Col440I; PTU-E3)
PS_Koxy4 (GCA_014050515.2)	6	6,406,207	CCC; 5 CCPs	6,075,855	5,969	pPSKoxy4_1 [IncFII_repA, IncFII_repA, RepB-RCR_reg; MOBF; IncFII_1_pKP91; PTU-FK (IncFII)] pPSKoxy4_2 [–; MOBH; IncAC; PTU-C (IncA)] pPSKoxy4_3 (Rep_3; –; IncR; –) pPSKoxy4_4 [–; MOBP1; –; PTU-Q2 (IncQ)] pPSKoxy4_5 (–; MOBP1; Col440I; PTU-E3)

702 *CCC, complete circular chromosome; CCP, complete circular plasmid.

703 # RIP, replication initiation protein domain; MOB, mobility group; REP, replicon type;

704 PTU, plasmid taxonomy unit; –, no data available. RIP, MOB and REP determined

705 using plaSquid [17], PTU determined using COPLA [16].

707 **Table 3.** Closest relatives of the circular 4,448 bp plasmid as determined using NCBI BLASTN

Species, strain and plasmid	Identity	Size (nt)	Accession	Source	Location	Reference
S. enterica PNUSAS021403 pPNUSAS021403_3	3511/3513 (99 %)	4,448	CP093106.1	Patient	USA	[60]
S. enterica subsp. *enterica* serovar Hadar PNUSAS018090 pPNUSAS018090_3	2662/2664 (99 %)	4,448	CP093099.1	Patient	USA	[60]
S. enterica 2016K-0377 p2016K-0377_3	4387/4389 (99 %)	4,448	CP093079.1	Patient	USA	[60]
S. enterica subsp. *enterica* serovar Hadar 12-2388 p12-2388.3	2662/2664 (99 %)	4,448	CP038598.1	Reference laboratory	Canada	[61]
E. cloacae 78 pP6	3403/3403 (100 %)	9,923	OW849463.1	Hospital	Spain	BioProject PRJEB42440
E. hormaechei RHBSTW-00070 pRHBSTW-00070_3	3403/3403 (100 %)	9,923	CP058166.1	Wastewater influent	UK	[46]
K. michiganensis RHBSTW-00676 pRHBSTW-00676_6	3211/3211 (100 %)	4,448	CP055990.1	Wastewater influent	UK	[46]

709 **Table 4.** Phenotypic traits that differentiate PS_Koxy1 and PS_Koxy2 from

710 PS_Koxy4

Growth in presence of	PS_Koxy1	PS_Koxy2	PS_Koxy4
D-Serine	–	–	+
L-Alanine	+	+	–
Minocycline	+	+	–
Acetic acid	–	–	+

+, Positive according to the Biolog GEN III system; –, negative according to the

Biolog GEN III system.

Figure 1. (a) Illustration of the circular conjugative 4,448-bp plasmid harboured by *K.*

michiganensis strains PS_Koxy1, PS_Koxy2 and PS_Koxy4. **(b)** Tetracycline-

antibiotic-associated genes predicted to be encoded within the genomes of strains

PS_Koxy1, PS_Koxy2 and PS_Koxy4 by RGI/CARD. All hits were considered 'strict'

by CARD.

Figure 2. Biofilm-forming abilities of the strains used in this study. **(a)** The biofilm-

forming abilities of all strains listed in **Table 1** were assessed. *K. michiganensis*

Ko14, a strain that does not form biofilms in TSBG, was used as the negative

control. Differences in biofilm formation in comparison with the control were

statistically significant for all strains (unpaired *t* test, adjusted *P* value < 0.05,

Benjamini-Hochberg; **Supplementary Table 1**). The ability of phage vB_KmiS-

Kmi2C to **(b)** prevent biofilm formation or **(c)** disrupt established biofilms was

assessed using those strains infected by the phage and forming the strongest

biofilms **(a)**. **(b, c)** Statistical significance of the differences in biofilm formation of

strains in the presence and absence of phage was assessed using unpaired *t* test.

(a–c) Data are shown as mean for four technical and three biological replicates per

strain.

734

Figure 3. Transmission electron micrograph of phage vB_KmiS-Kmi2C. Scale bar,

50 nm.

Figure 4. Genome map and predicted function of vB_KmiS-Kmi2C (upper image) and presence/absence of PCs within VC (lower

image). The genome map shows the predicted function of the coding regions, direction of transcription and PC ID for each gene

segment. The phage genus and host genus are shown to the left of the PC heatmap (see legend for annotation) and phage

genome size on the right-hand side of the heatmap.

Figure 5. Protein-sharing network generated from vB_KmiS-Kmi2C and INPHARED

database. Each node represents a phage genome and connection between the

genomes shown by a grey line. The nodes are annotated according to viral family.

The VC containing vB_KmiS-Kmi2C is highlighted with a black circle.

Figure 6. Protein phylogenetic analysis generated with VipTree from all phage within

the VC. The inner circle is coloured according to assigned viral genus and outer

circle according to phage host. The tree was rooted at the midpoint and scale shown

in the bottom right. *Klebsiella* phage vB_KmiM-4Dii (KiM 4Dii), a *Slopekvirus* phage

[21], was used as an outlier.

SUPPLEMENTARY MATERIAL ASSOCIATED WITH NEWBERRY *ET AL.*

Newberry F, Shibu P, Smith-Zaitlik T, McCartney AL, Hoyles L, Negus D. Phage vB_KmiS-Kmi2C infects members of the *Klebsiella oxytoca* complex and represents a novel genus of lytic bacteriophage.

Supplementary Table 1. Comparison of biofilm-forming abilities of strains with the negative control Ko14

P values were calculated using unpaired t tests of each biofilm-forming strain against the negative control (Ko14). P values were adjusted to account for multiple testing using the Benjamini–Hochberg (BH) method. Mean data are from four technical replicates per biological (Expt) replicate.

	Mean			SD	P value	BH P value
	Expt 1	Expt 2	Expt 3			
K. michiganensis Ko14	0.068	0.073	0.074	0.004		
K. michiganensis Ko13	0.110	0.154	0.113	0.025	3.21E-06	2.89E-05
K. michiganensis Ko43	0.169	0.198	0.206	0.020	8.77E-06	3.95E-05
K. michiganensis PS_Koxy4	0.238	0.276	0.295	0.029	3.14E-04	9.43E-04
K. michiganensis PS_Koxy2	0.156	0.158	0.160	0.002	4.86E-04	1.09E-03
K. oxytoca Ko37	0.108	0.116	0.103	0.007	8.28E-04	1.49E-03
K. michiganensis Ko21	0.107	0.136	0.104	0.017	1.13E-03	1.69E-03
K. oxytoca Ko53	0.178	0.225	0.194	0.024	1.06E-02	1.36E-02
K. michiganensis Ko22	0.296	0.323	0.315	0.014	1.31E-02	1.47E-02
K. michiganensis PS_Koxy1	0.186	0.319	0.322	0.078	2.06E-02	2.06E-02

Supplementary Table 2. Summary of VFDB data for *K. michiganensis* strains

PS_Koxy1, PS_Koxy2 and PS_Koxy4

Available as an Excel file (**Supplementary Table 2**).

Supplementary Figure 1. Bandage image generated from SPAdes assembly of the phage vB_KmiS-Kmi2C genome. The complete circular genome represents the phage, whereas the smaller linear contigs represent host DNA.

Supplementary Figure 2. Intergenomic similarity of all phage within the viral cluster shown by a red gradient (darker red = higher similarity) and percentage within each square. The phage and host genus are represented to the left of the heatmap and coloured annotates in the legend.

Supplementary Figure 3. Protein phylogenetic analysis generated with VipTree

from all phage within the viral cluster and 3 representative phage from related

genera (*Nipunavirus, Nonagavirus, Seuratvirus*). The inner circle is coloured

according to assigned viral genus and outer circle according to phage host. The tree

was rooted at the midpoint and scale shown in the bottom right.

Supplementary Figure 4. Shared protein percentage of all phage within the viral

cluster shown by a purple gradient (darker purple = higher protein percentage

shared). The phage and host genus are represented to the left of the heatmap and

coloured annotates in the legend. Dendrogram shown along the top *x* axis.