A sensation for inflation: initial swim bladder inflation in larval zebrafish is mediated

by the mechanosensory lateral line.

Alexandra Venuto[1], Stacey Thibodeau-Beganny[2], Josef G. Trapani[2], Timothy Erickson[3]

1. Department of Biology, East Carolina University, Greenville, NC, USA.

2. Department of Biology and Neuroscience Program, Amherst College, Amherst, MA, USA.

3. Department of Biology, University of New Brunswick, Fredericton, NB, Canada.

Abstract

Larval zebrafish achieve neutral buoyancy by swimming up to the surface and taking in air

through their mouths to inflate their swim bladders. We define this behavior as 'surfacing'. Little

is known about the sensory basis for this underappreciated behavior of larval fish. A strong

candidate is the mechanosensory lateral line, a hair cell-based sensory system that detects

hydrodynamic information from sources like water currents, predators, prey, and surface waves.

However, a role for the lateral line in mediating initial inflation of the swim bladder has not been

reported.

To explore the connection between the lateral line and surfacing, we utilized a genetic

mutant (*lhfpl5b*[-/-]) that renders the zebrafish lateral line insensitive to mechanical stimuli. We

observe that approximately half of these lateral line mutants over-inflate their swim bladders during

initial inflation and become positively buoyant. Thus, we hypothesize that larval zebrafish use their

lateral line to moderate interactions with the air-water interface during surfacing to regulate swim

bladder inflation. To test the hypothesis that lateral line defects are responsible for swim bladder

over-inflation, we show exogenous air is required for the hyperinflation phenotype and transgenic

rescue of hair cell function restores normal inflation. We also find that chemical ablation of anterior

lateral line hair cells in wild type larvae causes hyperinflation. Furthermore, we show that

manipulation of lateral line sensory information results in abnormal inflation. Finally, we report

spatial and temporal differences in the surfacing behavior between wild type and lateral line mutant

larvae. In summary, we propose a novel sensory basis for achieving neutral buoyancy where larval

zebrafish use their lateral line to sense the air-water interface and regulate initial swim bladder

inflation.

Keywords: behavior, buoyancy, hair cells, hydrodynamics, lateral line, lhfpl5, swim bladder,

sensory system, zebrafish

Introduction

The swim bladder of teleost fish provides for neutral buoyancy in the water column, thereby minimizing the energy expenditure associated with locomotion [1]. Larval teleosts initate swim bladder inflation by swimming up to the surface to ingest air using a set of behaviors we refer to as "surfacing" [2–4]. Although surfacing poses a significant risk of predation, larval fish are highly motivated to reach the surface to intake air. For example, pre-inflated lake trout larvae will swim hundreds of feet at a constant rate in attempts to access the surface [5]. Surfacing is a complex task that requires a combination of newly developed physical features, sensory systems, and motor skills [6–10] – (i) Larvae must discriminate up from down and (ii) be able to move directionally towards the surface; (iii) they must appropriately sense the air-water interface, and (iv) be able to intake air through the mouth upon arrival at the surface. Ingested air is then moved by peristalsis into the swim bladder through the pneumatic duct that connects the gut and swim bladder [5,10,11]. Together, surfacing represents perhaps the most complex behavior that larvae perform prior to achieving neutral buoyancy [12].

Little is known about the sensory cues that instruct the surfacing behavior. Perturbations to the vestibular system typically result in hypoinflation [13,14]. The inability to detect gravity by the utricular otolith is specifically responsible for the phenotype, presumably because larvae are unable to orient their movements upwards towards the surface [15]. Photosensory cues may also play a role in directional orientation and surface detection. However, which photosensory systems are involved has not been defined and the effects of altering lighting conditions varies between species [16–18]. Even less is known about how larval fish detect the air-water interface itself. Fish and some amphibians possess an additional hair cell-based sensory system - the *mechanosensory lateral line* - that contributes to detection of near-field hydrodynamic information originating from water currents, predators and prey, conspecifics, abiotic objects, and surface waves [19–26]. Specific to surface detection, surface-feeding fish use their lateral line to detect surface waves created by their prey [27]. Furthermore, Japanese flying fish are predicted to use their ventrally-located lateral line to sense transitions through the air-water interface [28,29]. While the lateral line is a strong candidate to aid in surfacing behaviors that lead to initial swim bladder inflation in larval fish, its specific role in this important behavior has not been explored.

A CRISPR-Cas9 knockout of *LHFPL tetraspan subfamily member 5b* (*lhfpl5b*) is the first genetic zebrafish mutant where the lateral line is non-functional from birth but hearing and balance

are normal [30]. These mutants provide an opportunity to uncover novel roles for the lateral line.

Unexpectedly, homozygous mutant larvae present with a hyperinflation phenotype characterized

by over-filled swim bladders and positive buoyancy, a very rare finding among swim bladder

defects [31,32]. Using several genetic and chemical techniques to manipulate lateral line function,

we show that lateral line sensory information is used by larval zebrafish to appropriately fill their

swim bladders, likely by detecting surface tension at the air-water interface using mechanosensory

neuromasts located on their heads. Overall, our study uncovers a novel role for the mechanosensory

lateral line mediating initial inflation of the swim bladder.

Methods

Animal husbandry and ethics statement

Adult zebrafish (*Danio rerio*) were maintained and bred using standard procedures [33].

All experiments used larvae at 2-6 days post-fertilization (dpf), which are of indeterminate sex at

this stage. Except where otherwise stated, animals were placed in clear, plastic tubs (dimensions:

15 cm x 10 cm x 4 cm, E3 volume: 460 mL) at 2 dpf and maintained at 28.5°C on a 14:10 light/dark

cycle at 500-700 lux in E3 embryo media (5 mM NaCl, 0.17 mM KCl, 0.33 mM CaCl2, 0.33 mM

MgCl2, buffered with NaHCO3). Animal research complied with guidelines stipulated by the

Institutional Animal Care and Use Committees at East Carolina University (protocol D372) and

Amherst College (assurance number 3925-1 with the Office of Laboratory Animal Welfare).

Mutant and transgenic fish lines

The zebrafish mutant allele *lhfpl5b^{vo35}* and transgenic lines *Tg(myo6b:EGFP-lhfpl5a)vo23*

and *Tg(myo6b:EGFP-PA)vo68* were used in this study [30,34]. For all experiments, *lhfpl5b^{vo35}*

homozygotes were identified from their hetero- and homozygous wild type siblings by lack of FM

1-43 (ThermoFisher) labeling of the lateral line hair cells.

To create a fish line that expresses the Channelrhodopsin-2 (ChR2) optogenetic protein

solely in the hair cells of the lateral line, approximately 2 kb of DNA upstream of the start codon

of the *lhfpl5b* gene was cloned into the p5E plasmid of the recombination-based multisite Gateway

cloning system (ThermoFisher). A Tol2 transposon backbone was then used to create the injectable

expression plasmid *lhfpl5b2028:ChR2-EYFP-PA*. This plasmid was then co-injected with

transposase mRNA into single-cell embryos [35] to create the *Tg(lhfpl5b2028:ChR2-EYFP-PA)*

zebrafish line.

 Buoyancy tests

To ascertain the buoyancy status of larval fish, larvae were anesthetized with MS-222

(Western Chemical Inc., Ferndale, WA) and placed at the vertical midway point of a 50 mL conical

tube containing E3 media. At the expiration of a 30-second timer, the end position of the fish was

recorded and the fish was removed from the tube for imaging under the microscope. If the fish

reached the top or bottom of the tube before 30 seconds, then the time it took to reach that location

was recorded. The total height of the water column was 8.9 cm, and the fish was placed half-way

at 4.45 cm. The rate of buoyancy was determined by dividing the centimeters travelled by the

seconds it took to travel that distance (cm/s).

Imaging and swim bladder measurements

Swim bladders were imaged using a SteREO Discovery.V8 microscope (Zeiss) equipped

with a Gryphax Arktur camera (Jenoptik). Lateral and dorsal images of the swim bladder were

taken for each fish. Swim bladder measurements were conducted using the measurement tool in

Adobe Photoshop where measurements were calibrated according to the image of the stage

micrometer taken when imaging experiments. The major axis and minor axis of the lateral view of

the swim bladder was measured as well as the minor axis of the dorsal view. All measurements

were taken from numbered images so that the measurer was blinded to the genotype/condition.

These meaurements were halved and used in the equation for the volume of an ellipsoid: $V =$

$4/3\pi abc$ to find the volume of the swim bladder [2]

Manipulation of photosensory cues during surfacing

At 0 dpf, zebrafish embryos were collected and immediately placed in plastic tubs filled

with E3 (detailed above) in an unlit incubator. Larvae were not brought into light until 6 dpf, at

which point lateral line mutants and wild type siblings were identified by FM 1-43 labeling. After

sorting, larvae were analyzed for buoyancy. The experiment was repeated three times using 20-21

larvae (combined mutant and wild type) per trial.

Blocking access to the air-water interface

Mutant larvae and wild type siblings were sorted at 2 dpf into glass cylinders (diameter: 9.5

cm, E3 volume: 425 mL) equipped with fitted, wire mesh filter plungers (Bodum Inc., USA).

Conditions were as follows: wild type with open access to the surface, wild type with blocked

access to the surface, mutant with open access to the surface, and mutant with blocked access to

the surface. All containers were filled half way with E3 and in the experimental condition ("blocked

access"), the filter was pressed down below the surface for both wild type and mutant larvae. The

control condition used the same amount of E3 but the filter remained above the surface so that

larvae could access the exogenous air. Conditions remained constant until 6 dpf when larvae were

imaged and analyzed for swim bladder inflation. The experiment was repeated three times using

15-25 larvae per condition.

Analysis of surfacing behavior

Wild type and *lhfpl5b^{vo35}* larvae were blocked from having access to the surface until 9 AM

on 4 dpf when individuals were immediately placed into clear rectangular containers (dimensions:

3 cm x 3 cm x 3 cm, E3 volume: 20 mL) for video recording. Recordings lasted 12 hours, from 9

AM to 9 PM on 4 dpf. At the end of filming, larvae were imaged and analyzed for buoyancy and

swim bladder volume, as described. All surfacing videos were numbered for blind analysis.

Surfacing analysis was conducted by manual observation of each individual larvae from the videos

(30 frames per second) and manually counting visits made to the surface (counted as reaching the

meniscus), the timeframe of the video at which the surfacing event occurred, and the duration of

time spent at the surface per surfacing event. Recording was repeated six times with 3 mutant and

2 wild type larvae per recording. Larvae that did not inflate their swim bladders by the end of

filming were excluded from this analysis due to lack of surfacing events.

Lateral line hair cell ablations

For all experiments, wild type and *lhfpl5b* mutant larvae were treated with ototoxin with

untreated siblings acting as controls (n = 7-15 larvae per condition per trial, 4 trials) For single

treatments, surface accessed was blocked as described above for untreated and treated larvae until

9 AM on 4 dpf when treatment occurred. Single 30 minute treatments with neomycin sulfate (EMD

Millipore Corp., USA) were done with a 100 μm solution in E3 while repeated neomycin

treatments [36] were done as previously described with a concentration of 50 μM. For full ablation

with copper sulfate ($CuSO_4$; Ward's Science, Rochester, NY), the treatment concentration was 10

uM with a 30 minute exposure. For experiments where head and trunk neuromasts were selectively

ablated, we anesthetized the larvae and pipetted them on agarose gel. Excess water was removed

and a thin layer of petroleum jelly (Vaseline, Unilever USA) was placed horizontally at the back

of the head. Next, 30 uM $CuSO_4$ or E3 was pipetted onto the appropriate region depending on the

experimental condition. Immediately follow the 13 minute treatment, larvae were placed in fresh

E3 in clear tubs detailed above for 24 hours until 9 AM on 5 dpf when larvae were imaged and analyzed for swim bladder inflation. For all experiments, larvae were imaged and analyzed for swim bladder inflation at 9 AM on 5 dpf. Each experiment was repeated three times using 10-24 larvae per condition.

Transgenic rescue of lateral line mutant

Larval zebrafish from crosses of *lhfpl5b+/-* and *Tg(myo6b:EGFP-lhfpl5a)vo23Tg; lhfpl5b+/-* were sorted at 2 dpf for transgene expression. At 6 dpf, larvae were numbered, imaged for swim bladder inflation and genotyped for the *lhfpl5bvo35* allele as previously described [30]. The experiment was repeated twice using 26-48 larvae per condition.

Optogenetic activation of lateral line hair cells with Channelrhodopsin-2 (ChR2)

After blocking access to the surface until 4 dpf, we exposed wild type and lateral line mutants, with or without the *lhfpl5b2028:ChR2-EYFP-PA* transgene, to LED blue light (470 nm) with an intensity of 730 lux over their holding tanks against a dark background. Light flashed for 25 milliseconds with a one second delay between each pulse. Control larvae were kept in the same background, without blue light exposure. Conditions were held constant until 6 dpf when larvae were imaged and analyzed for swim bladder inflation. The experiment was repeated three times using 15-25 larvae per condition.

Oil-water interface experiments

Mutant larvae and wild type siblings were sorted into clear cylindrical containers (diameter: 5 cm , volume: 60 mL) with embryo media only or embryo media with a 2 mm thick surface layer of lab grade mineral oil (Ward's Science, Rochester, NY). Oil on the surface decreases surface tension [37] and has been previously used in experiments involving the zebrafish swim bladder [38]. Conditions remained constant until 6 dpf when larvae were imaged and analyzed for swim bladder volume. Because oil-filled larvae are negatively buoyant, we did not perform buoyancy tests and classified larvae as over-filled based on swim bladder volume alone. The experiment was repeated three times using 15-25 larvae per condition.

Graphs and statistical tests

All graphs and statistical tests were done using R [39]. One-way ANOVAs were used to compare the conditions as a whole (genotype independent variable plus environment independent variable) with Tukey post-hoc test. Chi-squared tests were used to analyze proportion data for over

and under-inflation. For the over-inflation comparisons, the two categories used in the comparison

between conditions/genotypes were (i) over-inflation and (ii) all other inflation (regular and under),

meaning that totals per category/genotype accounted for 100% of the population. Similarly, for the

under-inflation comparisons, the two categories used in the comparison between

conditions/genotypes were (i) under-inflation and (ii) all other inflation (regular and over). P-values

less than 0.05 were considered significant.

Results

Lateral line mutants exhibit a swim bladder hyperinflation phenotype

The lateral line has not been previously implicated in initial inflation of the larval swim

bladder. As such, it was unexpected to observe that lateral line mutants (*lhfpl5b$^{-/-}$*) exhibit

hyperinflated swim bladders by 5-6 dpf (Figure 1A). We measured swim bladder volume and

buoyancy for each larva in the mutant and wild type populations (Figure 1B) and found that, on

average, 53.7% ($SD = 6.4\%$, $n = 41$) of lateral line mutants over-inflate their swim bladders and

7.3% ($SD = 7.1\%$, $n = 41$) under-inflate. For wild type larvae (including *lhfpl5b^{vo35}* heterozygotes),

0.0% ($SD = 0.0\%$, $n = 41$) over-inflate and 4.9% ($SD = 3.7\%$, $n = 41$) under-inflate their swim

bladders during initial inflation on average (Figure 1C). The proportion of over-inflation is

significantly different between mutant and wild type larvae (chi-square statistic: 70.45, $p <$

0.00001), while the proportion of under-inflation is not (chi-square statistic: 0.3546, $p = 0.552$).

Average swim bladder volume of mutant fish is larger ($M = 0.012$ mm^3, $SD = 0.002$ mm^3) than

wild type siblings ($M = 0.0058$ mm^3, $SD = 0.00043$ mm^3, $t(80) = 17.4$, $p < 0.0001$) (Figure 1D).

The proportions of neutrally and positively buoyant larvae do not change when raised in total

darkness (Supplemental Figure 1). This unexpected hyperinflation phenotype in *lhfpl5b^{vo35}*

homozygotes led us to investigate if the lateral line is contributing important sensory information

during the surfacing behavior.

Access to the air-water interface is required for the hyperinflation phenotype of lateral line mutants

For most teleosts, initial swim bladder inflation requires an intake of exogenous air,

typically from the water's surface. To test if the hyperinflation phenotype of lateral line mutants

requires access to surface air, we physically blocked access to the surface from 2-6 dpf (Figure 2).

On average, 0% ($SD = 0.0\%$, $n = 43$) of blocked *lhfpl5b* mutants over-inflate their swim bladders,

which is significantly different from mutants with access to the surface (chi-square statistic: 70.45,

$p < 0.00001$; Figure 2B). The average swim bladder volume of blocked mutants ($M = 0.0033$ mm^3,

$SD = 0.0015$ mm^3) is significantly smaller than mutants with open access ($M = 0.014$ mm^3, $SD =$

0.0027 mm^3) (One-way ANOVA with Tukey post-test, $p < 0.00001$). Blocked mutant larvae had

statistically similar swim bladder volumes to blocked wild type larvae ($M = 0.0037$ mm^3, $SD =$

0.001 mm^3) (One-way ANOVA with Tukey post-test, $p = 0.927$) (Figure 2C). From these data, we

conclude that the *lhfpl5b* mutant hyperinflation phenotype requires intake of exogenous air and is

not caused by abnormal gas exchange from surrounding media nor excess internal gas production.

Transgenic rescue of lateral line mutants restores normal inflation

We next examined if lateral line defects are responsible for the hyper-inflation phenotype

of *lhfpl5b* mutants by restoring lateral line function in mutant larvae. We used the

Tg(myo6b:EGFP-lhfpl5a)vo23 transgene, which uses a hair cell specific promoter (*myo6b*) to drive

a GFP-tagged *lhfpl5a* gene [34]. It has been shown that this *lhfpl5a* transgene rescues lateral line

hair cell function in *lhfpl5b*$^{-/-}$ *(or* lateral line) mutant fish [30]. Using this rescue method, we

observe no cases of hyperinflation in transgenic mutants (0%, $SD = 0.0\%$, $n = 17$), which is

significantly different from non-transgenic mutants (chi-square statistic: 82.12, $p < 0.00001$; Figure

3A). The average volume of the transgenic mutant swim bladder ($M = 0.0057$ mm^3, $SD = 0.00001$

mm^3) is significantly smaller than non-transgenic, *lhfpl5b* mutant siblings ($M = 0.0109$ mm^3, $SD =$

0.0031 mm^3) (One-way ANOVA with Tukey post-test, $p = 0.000003$), but statistically similar to

that of wild type transgenics ($M = 0.0057$ mm^3, $SD = 0.0005$ mm^3) (One-way ANOVA with Tukey

post-test, $p = 0.979$, Figure 3B). Because the restoration of *lhfpl5* function specifically in hair cells

mitigates hyperinflation in *lhfpl5b* mutants, we conclude that the phenotype is due to defects in

sensory hair cell function and not caused by an unrecognized role for *lhfpl5b* in another sensory

organ or the swim bladder itself.

Selective ablation of head neuromasts produces a hyperinflation phenotype in wild type larvae

To further test if lateral line defects are responsible for the hyper-inflation phenotype seen

in *lhfpl5b* mutants, we ablated lateral line hair cells in wild type larvae using either neomycin

sulfate (neo) or copper sulfate (CuSO4) (Supplemental Figure 2 and Figure 4). Administering either

single or repeated neomycin treatments during the initial swim bladder inflation period of 3 - 4 dpf

results in a significant increase in the proportion of positively buoyant larvae (single neo chi-square

statistic: 9.95, $p = 0.0016$, repeated neo chi-square statistic: 34.88, $p < 0.0001$; Supplemental Figure

2A), but a non-significant increase in the average swim bladder volume (One-way ANOVA with

Tukey post-test, single neo $p = 1.0$, repeated neo p $= 0.578$; Supplemental Figure 2B). By

comparison, a single CuSO4 treatment results in a hyperinflation phenotype that is statistically indistinguishable from lateral line mutants. An average of 59.6% ($SD = 7.7\%$, $n = 72$) of CuSO4-treated wild types over inflate their swim bladders, which is significantly different from wild type controls (chi-square statistic: 39.69, $p < 0.0001$; Supplemental Figure 2A). The average swim bladder volume of the CuSO4-treated wild type larvae ($M = 0.0126$ mm^3, $SD = 0.0027$ mm^3) is statistically similar to the average swim bladder volume of untreated lateral line mutants ($M = 0.0139$ mm^3, $SD = 0.0007$ mm^3) (One-way ANOVA with Tukey post-test, $p = 0.813$) (Supplemental Figure 2B).

Since the anterior lateral line is likely to be the only part of the sensory organ to interact with the air-water interface during surfacing, we predict that disruption of the head neuromasts will result in hyperinflation and that trunk neuromasts will not play a role (Figure 4, Supplemental Figure 3). Selective ablation of the anterior region with CuSO4 results in an average of 35.9% ($SD = 10.1\%$, $n = 38$) hyperinflated larvae, which is significantly different from unablated controls (chi-square statistic: 39.69, $p < 0.0001$; Figure 4B). Wild type larvae with head-specific ablations exhibit a significant increase in average swim bladder volume ($M = 0.016$ mm^3, $SD = 0.0013$ mm^3) compared to untreated wild type siblings ($M = 0.0012$ mm^3, $SD = 0.0013$ mm^3) (One-way ANOVA with Tukey post-test, $p = 0.0365$) and statistically comparable swim bladder volumes to that of full CuSO4-treated larvae ($M = 0.016$ mm^3, $SD = 0.0029$ mm^3) (One-way ANOVA with Tukey post-test, $p = 0.987$) (Figure 4C). Conversely, tail-specific CuSO4 treatments do not cause hyperinflation in wild type fish and result in statistically similar swim bladder volumes to untreated siblings ($M = 0.012$ mm^3, $SD = 0.0031$ mm^3) (One-way ANOVA with Tukey post-test, $p = 0.835$) (Figure 4C). The *lhfpl5b* mutant hyperinflation phenotype was not affected by either neomycin or CuSO4 treatments (Supplemental Figure 2). Taken together, we conclude that the anterior region of the lateral line system is critical to surface detection during initial swim bladder inflation.

Decreasing surface tension results in over-filling of the swim bladder

Surface tension is a characteristic that makes the air-water interface distinct from the rest of the water column. We predict that larval fish sense this stimulus with their anterior lateral line neuromasts and that by decreasing the interfacial tension, we would observe abnormal swim bladder inflation in wild type larvae while lateral line mutants would be unaffected. The interfacial tension between oil and water is approximately half that of air and water [37]. We layered oil onto the surface of the embryo media and allowed larvae to perform surfacing behaviors between 2-6

dpf (Figure 5). Using this experimental design, we find that the average swim bladder volume of

wild types exposed to an oil-water interface (M = 0.016 mm^3, SD = 0.0029 mm^3) is significantly

greater than the control group with access to an air-water interface (M = 0.007 mm^3, SD = 0.001

mm^3) (One-way ANOVA with Tukey post-test, p < 0.0001). Furthermore, oil-exposed wild types

have statistically similar swim bladder volumes to that of *lhfpl5b* mutants (M = 0.014 mm^3, SD =

0.0038 mm^3) (One-way ANOVA with Tukey post-test, p = 0.886) (Figure 5C). Lateral line mutants

overfill their swim bladders to statistically similar proportions when presented with either an air-

water or oil-water interface (M = 0.01 mm^3, SD = 0.0026 mm^3) (One-way ANOVA with Tukey

post-test, p = 0.133). We conclude that surface tension is one of the surface characteristics the

lateral line is capable of detecting during initial swim bladder inflation.

Optogenetic activation of lateral line hair cells prevents hyperinflation in lateral line mutant larvae

If hydrodynamic information is informing the surfacing behaviors of larval fish, then

stimulus-independent manipulations of lateral line activity should alter their ability to correctly

inflate their swim bladders. To test this, we created the *Tg(lhfpl5b2028:ChR2-EYFP-PA)*

transgenic line that expresses the blue light activated channel, Channelrhodopsin-2, specifically in

lateral line hair cells (See Methods, Supplemental Figure 4). We predicted that optogenetic

stimulation of *lhfpl5b* mutants, whose lateral lines are insensitive to physical stimuli, would reduce

the frequency of the hyperinflation phenotype. Consistent with these predictions, under blue-light

stimulation, lateral line mutants with the *lhfpl5b:ChR2* transgene do not exhibit positive buoyancy

and have a significant decrease in average swim bladder volume (M = 0.0046 mm^3, SD = 0.0034

mm^3) compared to unstimulated transgenic mutants (M = 0.012 mm^3, SD = 0.0031 mm^3) (One-

way ANOVA with Tukey post-test, p = 0.00127) (Figure 6B). Transgenic wild types under blue

light exhibit a non-significant decrease in swim bladder volume (M = 0.0052 mm^3, SD = 0.004

mm^3) compared to unstimulated transgenic siblings (M = 0.0079 mm^3, SD = 0.002 mm^3) (One-way

ANOVA with Tukey post-test, p = 0.458). To demonstrate that blue light stimulation is reducing

swim bladder volumes via activation of lateral line hair cells, we treated wild type *lhfpl5b:ChR2*

larvae with CuSO4. As expected, we observe hyperinflation following CuSO4 treatment in

stimulated wild type *lhfpl5b:ChR2* larvae, with an average swim bladder volume (WT: M = 0.012

mm^3, SD = 0.0037 mm^3) that is statistically similar to experimental non-transgenic *lhfpl5b* mutants

(M = 0.011 mm^3, SD = 0.0024 mm^3) (One-way ANOVA with Tukey post-test, p = 0.998)

(Supplemental Figure 4B). Overall, optogenetic activation of mechanically-insensitive lateral line

organs in *lhfpl5b* mutants prevents the hyperinflation phenotype. Together, these data support the

hypothesis that the lateral line provides crucial sensory information during initial swim bladder

inflation in larval zebrafish.

Lateral line mutants exhibit abnormal surfacing behaviors

Our results suggest that larval fish are using their head neuromasts to sense the interfacial

tension between air and water at the surface. Even in wild type zebrafish, the interaction with the

surface during initial swim bladder inflation is not well characterized. To study this behavior in

more detail, we quantified the number of unique visits to the surface and the amount of time per

visit spent within the meniscus region from videos of wild type (n = 7) and mutant (n = 14) larvae

at 4 dpf. *lhfpl5b* mutants that will over-inflate their swim bladders make more visits (M = 87.3, SD

= 17.4) to the surface compared to both wild type (M = 42.0, SD = 16.3) and mutant siblings that

will inflate to normal proportions (M = 31.9, SD = 19.2) (One-way ANOVA with Tukey post-test,

p = 0.0004) (Figure 7A). Mutants destined for hyperinflation also spend more time (M = 139.0 s,

SD = 31.6 s) at the surface than normally-inflated wild type (M = 51.9 s, SD = 24.7 s) and mutant

siblings (M = 42.3 s, SD = 29.5 s) (One-way ANOVA with Tukey post-test, p = 0.00006) (Figure

7B). Since excess surfacing could be due to some mutants initiating the behavior earlier, we also

documented the time of the first surfacing event and found no significant difference between

neutrally buoyant and over-inflated larvae (Figure 7C). We conclude there is a correlation between

how larvae interact with the surface during the initial swim bladder inflation period and their

resulting swim bladder volume and buoyancy status.

Discussion

The intake of exogenous air from the surface is a common feature for intitial swim bladder

inflation in many species of fish [40–44]. However, few studies have considered what sensory

information is required to achieve neutral buoyancy. In this study, we describe a new role for the

mechanosensory lateral line in mediating initial inflation of the swim bladder in zebrafish (Figure

8). Our data support a model in which larval zebrafish use their head neuromasts to detect the air-

water interface in support of their transition to neutral buoyancy.

Lateral line defects in lhfpl5b mutants are responsible for hyperinflation phenotypes

lhfpl5b mutant zebrafish are the first genetic model for the specific loss of lateral line

function in any organism [30]. These "lateral line mutants" exhibit an uncommon swim bladder

hyperinflation phenotype that prompted us to investigate this novel connection between lateral line
sensation and initial inflation of the swim bladder (Figure 1). Previous studies have demonstrated
that restricting access to surface air during the initial inflation period prevents swim bladder
inflation in larval zebrafish [40,45]. In addition, one study showed that some *solute carrier family*
12 member 2 (*slc12a2 / nkcc1)* mutant zebrafish over-fill their swim bladders and that blocking
access to the surface prevents this phenotype [32], though further studies are required to determine
the cause of over-inflation in *slc12a2* mutants. Likewise, we find that blocking access to the air-
water interface eliminates over-inflation in lateral line mutants and greatly decreases average swim
bladder volumes of both mutant and wild type larvae (Figure 2). Our method of blocking access to
the air-water interface resulted in a small proportion of larvae (both wild type and mutant) inflating
their swim bladder to normal proportions, though hyperinflation was never observed. Inflation was
likely achieved by larvae accessing air bubbles that remain in the water after submerging the filter
[40]. Overall, the hyperinflation phenotype in lateral line mutants requires an intake of exogenous
air, pointing to a sensory defect in *lhfpl5b* mutants rather than a physiological one.

In support of this sensory-deficit hypothesis, we show the hyperinflation phenotype is
prevented by either transgenic rescue of *lhfpl5* function specifically in hair cells or optogenetic
stimulation of mechanically-insensitive lateral line hair cells in *lhfpl5b* mutants (Figures 3 and 6).
Furthermore, chemical ablation of lateral line hair cells in wild type larvae prior to initial inflation
can phenocopy the *lhfpl5b* mutant. A single treatment with CuSO4 caused similar swim bladder
over-inflation proportions to lateral line mutants (Supplemental Figure 2). Repeated neomycin
treatments over 3-4 dpf caused the next highest percentage of swim bladder over-inflation and a
single neomycin treatment only resulted in a few cases of wild type over-inflation. These results
agree with previous work showing rapid hair cell regeneration following neomycin treatment and
comparatively slower recovery after CuSO4 exposure [36,46–49]. Thus, via multiple genetic and
chemical approaches, our data show that lateral line hair cell activity can modulate swim bladder
inflation in larval zebrafish.

Ototoxins have been extensively used to study lateral line-mediated behaviors in free-
swimming larvae and adult fish [19,20,50–52]. However, we are not aware of any behavioral
studies where the lateral line was disrupted prior to larvae achieving neutral buoyancy, likely
explaining why this connection between the lateral line and initial swim bladder inflation has not
been noted in current literature. Interestingly, an 1896 study predicted a relationship between the

lateral line and swim bladder inflation, since disruption of the lateral line in adult goldfish resulted

in "swelling of the swim bladder" and positive buoyancy [53]. However, it is not clear if the

phenotypic similarities between these adult goldfish and our experimental larval zebrafish share

the same underlying causes. Regarding roles for the lateral line in early life behaviors, a close

parallel to our findings is the report that unhatched red-eyed treefrogs use their lateral line organ to

mediate mechanosensory-cued hatching [54]. In this case, the frog lateral line can sense vibrational

stimuli, such as that produced by predators, and promote escape hatching behaviors even before

the onset of vestibular function. Similarly, our data suggest that the lateral line is modulating early

locomotor activities in zebrafish prior to the development of fully formed vestibular or rheotactic

responses [55–57]. As such, the involvement of lateral line in the surfacing behavior may represent

the earliest behavioral role for the lateral line yet described.

Head neuromasts are required for normal swim bladder inflation, likely by sensing surface tension

We expanded on the full-body lateral line ablation experiments with specific ablations of

either the head or truck neuromasts with CuSO4. Previous studies have shown that head neuromasts

are involved with detection of surface features and prey capture [21,58–60]. Indeed, ablation of the

head neuromasts in wild type larvae recapitulated the hyperinflation phenotype of *lhfpl5b* mutants

while ablation of the tail neuromasts had no effect on either swim bladder volume or buoyancy

(Figure 4). These findings suggest that the head neuromasts interact with the air-water interface

during surfacing and that the posterior lateral line is dispensible for achieving neutral buoyancy.

As such, our findings also provide additional support for the anterior lateral line as a central

mediator of surface-related behaviors in fish.

The interfacial tension between air and water represents a unique hydrodynamic stimulus

that may be detected by the lateral line. The tension between oil and water is half that of air and

water [37]. Thus, by adding a layer of oil to the surface, we predicted that surface detection would

become increasingly difficult and result in over-filling of the swim bladder. We find that, when

given access to an oil-water interface, wild type larvae over-fill their swim bladders with oil to a

statistically similar volume as lateral line mutants who overinflate with air (Figure 6). We interpret

this result as support for the hypothesis that the lateral line senses interfacial tension between air

and water as larvae interact with the surface during initial swim bladder inflation. This

interpretation is further supported by the observation that *lhfpl5b* mutants (who are already

insensitive to interfacial tension) exhibit similar swim bladder volumes when exposed to either oil

or air. However, a possible complementary interpretation for these data is that larvae who fill their

swim bladders with oil may be deprived of the normal feedback on buoyancy levels as they intake

from the surface. Since oil is denser than air, oil-filled larvae are negatively buoyant. In this

scenario, larvae may continue to visit the surface for air intake in an attempt to achieve neutral

buoyancy. However, future studies are required to examine if such a feedback mechanism exists in

larval zebrafish.

Swim bladder over-inflation is correlated with excess time spent at the air-water interface

Since our data suggest that the air-water interface is providing a distinct stimulus to the

lateral line, we investigated whether surface interactions differed between hyperinflated *lhfpl5b*

mutants and their wild type counterparts. As predicted, over-inflated mutants took more visits *and*

spent more time at the surface than neutrally-buoyant mutant and wild type siblings (Figure 7).

Excess breaching of the surface occurred despite over-inflated and neutrally buoyant larvae

exhibiting similar timing for their first surfacing attempts. While it is unclear whether larvae ingest

air during the entirety of the duration spent above the air-water interface, it is known that surface

breaches are observed during initial inflation [2,11,40]. Therefore, the increase in surface visits

likely leads to an increase in air deposition to the swim bladder in lateral line mutants.

Multisensory integration during the surfacing behavior

Interestingly, approximately half of all lateral line mutants inflate their swim bladders to

normal proportions and continue to develop into viable adults. To account for this fact, we

hypothesize that additional somatosensory, photosensory, or mechanosensory systems may

compensate for the loss of lateral line sensation and help mutant larvae achieve neutral buoyancy

[15,16,61]. The lateral line is involved several multi-sensory behaviors, including flow orientation

[62–64], prey capture [21,65,66], and predator avoidance [23,54,67], and we predict that surfacing

can be added to this list.

The photosensory and mechanosensory systems exhibit distinct organizational patterns for

primary input to the central nervous system — even the anterior and posterior lateral line ganglia

primary projections are somatotopically organized in zebrafish larvae [68–73]. The Mauthner

neurons integrate multisensory input at the level of the hindbrain, but the surfacing behavior is

unlikely to use this circuit. For higher order processing however, information from the lateral line,

vestibular, and photosensory systems are integrated in various brain regions. The thalamus is one

such sensory integration center in zebrafish [74]. Socially-relevant mechanosensory and

photosensory information is represented in a thalamic region marked by the expression of

parathyroid hormone 2 (*pth2*) and *somatostatin 7* (*sst7*) [24,26,75]. Vestibular information is also

represented in the thalamus [76,77], though it is not clear if the exact same region is involved.

Photosensory, lateral line, and vestibular/auditory information is also integrated in the tectum [76–

79]. In fact, neurons in the periventricular layer (PVL) of the tectum, which is similar to the superior

colliculus in mammals, respond to visual, auditory, and lateral line inputs [80–83]. However, it is

hypothesized that most of these modalities are processed in parallel at younger stages of

development [83]. It is worth noting that these sensory systems are in developing stages during

surfacing behaviors.

Photosensory cues did not alter the buoyancy status of either wild type or *lhfpl5b* mutant

larvae (Supplemental Figure 1), suggesting that, at most, photosensory systems play a subtle

modulatory role in the surfacing behavior of larval zebrafish. The vestibular system is likely

required to coordinate the vertical orientation and climbing portions of surfacing [38,56,84].

Meanwhile, our data support the idea that the lateral line is responsible for the surface detection

and breaching aspects of surfacing. However, given that both the vestibular and lateral line systems

are providing spatial information in during the surfacing behavior, there may be crosstalk between

them. For example, the vestibular system must quickly adapt to a dynamic increase in buoyancy

over the course of multiple breaching events, so feedback from the lateral line during surface

interactions may help fine tune the vestibular sense. However, the link between these systems have

not been well-investigated during deliberate behaviors like surfacing [85,86]. So, while sensory

integration exists in zebrafish and seems necessary for larvae to achieve neutral buoyancy, the

integration dynamics are unclear during the intial swim bladder inflation timeframe and require

future investigation.

Acknowledgements

The authors would like to thank Dr. Fadi Issa (ECU) for his support and technical advice. The

authors also thank the Issa, Trapani, and Erickson lab members for fish care and maintenance.

Author contributions

Conceptualization: A.V., T.E.; Methodology: A.V., J.G.T., T.E.; Validation: A.V.; Formal

analysis: A.V.; Investigation: A.V.; Resources: S.T-B., J.G.T; Writing - original draft: A.V., T.E.;

Writing - review & editing: A.V., J.G.T., T.E..; Visualization: A.V.; Supervision: T.E.; Project

administration: T.E.; Funding acquisition: J.G.T., T.E.

Funding

This work was supported by funding from East Carolina University's Division of Research,

Economic Development and Engagement (REDE) and a Natural Science and Engineering

Research Council (NSERC) Discovery Grant (RGPIN-2021-03166) to T.E., a National Institute on

Deafness and Other Communication Disorders (NIDCD) award to J.G.T. (NIH R15DC014843),

and East Carolina Department of Biology funding provided to A.V..

References

1. Alexander RM. Physical aspects of swimbladder function. Biological reviews of the Cambridge Philosophical Society. John Wiley & Sons, Ltd; 1966. pp. 141–176. doi:10.1111/j.1469-185x.1966.tb01542.x

2. Lindsey BW, Smith FM, Croll RP. From inflation to flotation: Contribution of the swimbladder to whole-body density and swimming depth during development of the zebrafish (Danio rerio). Zebrafish. 2010;7: 85–96. doi:10.1089/zeb.2009.0616

3. Evans HM, Damant GCC. Observations on the Physiology of the Swim Bladder in Cyprinoid Fishes. Journal of Experimental Biology. 1928;6: 42–55.

4. v. Ledebur JF. Beiträge zur Physiologie der Schwimmblase der Fische. Z Vgl Physiol. 1928;8: 445–460. doi:10.1007/BF00338968

5. Tait JS. The First Filling Of The Swim Bladder In Salmonoids. Can J Zool. 1960;38: 179–187. doi:10.1139/z60-021

6. Bailey HC, Doroshov SI. The duration of the interval associated with successful inflation of the swimbladder in larval striped bass (Morone saxatilis). Aquaculture. 1995;131: 135–143. doi:10.1016/0044-8486(94)00215-A

7. Kitajima C, Watanabe T, Tsukashima Y, Fujita S. Lordotic Deformation and Abnormal Development of Swim Bladders in Some Hatchery-Bred Marine Physoclistous Fish in Japan. J World Aquac Soc. 1994;25: 64–77. doi:10.1111/j.1749-7345.1994.tb00806.x

8. Chatain B. Problems related to the lack of functional swimbladder in intensive rearing of Dicentrarchus labrax and Sparus auratus. Advances in Tropical Aquaculture, Workshop at Tahiti, French Polynesia, 20 Feb - 4 Mar 1989. 1989 [cited 16 Jan 2021]. Available: https://archimer.ifremer.fr/doc/00000/1432/

9. Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. Stages of embryonic development of the zebrafish. Developmental Dynamics. 1995;203: 253–310. doi:10.1002/aja.1002030302

10. Doroshev S, Cornacchia J, Hogan K. Initial swim bladder inflation in the larvae of physoclistous fishes and its importance for larval culture. Rapports et Proces-verbaux des Réunions Cons int Explor Mer. 1981;178: 495–500.

11. Rieger PW, Summerfelt RC. Microvideography of gas bladder inflation in larval walleye. J Fish Biol. 1998;53: 93–99. doi:10.1111/J.1095-8649.1998.TB00112.X

12. Fero K, Yokogawa T, Burgess HA. The behavioral repertoire of larval zebrafish. Neuromethods. 2011;52: 249–291. doi:10.1007/978-1-60761-922-2_12

13. Schoppik D, Bianco IH, Prober DA, Douglass AD, Robson DN, Li JMB, et al. Gaze-stabilizing central vestibular neurons project asymmetrically to extraocular motoneuron pools. Journal of Neuroscience. 2017;37: 11353–11365. doi:10.1523/JNEUROSCI.1711-17.2017

14. Nicolson T, Rüsch A, Friedrich RW, Granato M, Ruppersberg JP, Nüsslein-Volhard C. Genetic analysis of vertebrate sensory hair cell mechanosensation: The zebrafish circler mutants. Neuron. 1998;20: 271–283. doi:10.1016/S0896-6273(00)80455-9

15. Riley BB, Moorman SJ. Development of utricular otoliths, but not saccular otoliths, is necessary for vestibular function and survival in zebrafish. J Neurobiol. 2000;43: 329–337. doi:10.1002/1097-4695(20000615)43:4<329::AID-NEU2>3.0.CO;2-H

16. Trotter AJ, Battaglene SC, Pankhurst PM. Effects of photoperiod and light intensity on initial swim bladder inflation, growth and post-inflation viability in cultured striped

trumpeter (Latris lineata) larvae. Aquaculture. 2003;224: 141–158. doi:10.1016/S0044-
8486(03)00212-6

17. Villamizar N, García-Alcazar A, Sánchez-Vázquez FJ. Effect of light spectrum and
photoperiod on the growth, development and survival of European sea bass (Dicentrarchus
labrax) larvae. Aquaculture. 2009;292: 80–86. doi:10.1016/j.aquaculture.2009.03.045

18. Stuart KR, Drawbridge M. The effect of photoperiod on larval culture performance of two
marine finfish species. Aquaculture. 2012;360–361: 54–57.
doi:10.1016/j.aquaculture.2012.07.017

19. Montgomery JC, Baker CF, Carton AG. The lateral line can mediate rheotaxis in fish.
Nature. 1997;389: 960–963. doi:10.1038/40135

20. Bleckmann H. The Lateral Line System of Fish. Fish Physiology. 2006. pp. 411–453.
doi:10.1016/S1546-5098(06)25010-6

21. Carrillo A, McHenry MJ. Zebrafish learn to forage in the dark. Journal of Experimental
Biology. 2016;219: 582–589. doi:10.1242/jeb.128918

22. Mogdans J. Sensory ecology of the fish lateral-line system: Morphological and
physiological adaptations for the perception of hydrodynamic stimuli. J Fish Biol.
2019;95: 53–72. doi:10.1111/jfb.13966

23. Stewart WJ, Cardenas GS, McHenry MJ. Zebrafish larvae evade predators by sensing
water flow. J Exp Biol. 2013;216: 388–98. doi:10.1242/jeb.072751

24. Anneser L, Alcantara IC, Gemmer A, Mirkes K, Ryu S, Schuman EM. The neuropeptide
Pth2 dynamically senses others via mechanosensation. Nature. 2020;588: 653–657.
doi:10.1038/s41586-020-2988-z

25. Butler JM, Maruska KP. Mechanosensory signaling as a potential mode of communication
during social interactions in fishes. J Exp Biol. 2016;219: 2781–2789.
doi:10.1242/jeb.133801

26. Venuto A, Smith CP, Cameron-Pack M, Erickson T. Alone in a crowd: effect of a
nonfunctional lateral line on expression of the social hormone parathyroid hormone 2. Biol
Open. 2022;11. doi:10.1242/bio.059432

27. Bleckmann H, Schwartz E. The functional significance of frequency modulation within a
wave train for prey localization in the surface-feeding fishAplocheilus lineatus
(Cyprinodontidae). Journal of Comparative Physiology A. 1982;145: 331–339.
doi:10.1007/BF00619337

28. Gibbs MA. Lateral Line Receptors: Where Do They Come from Developmentally and
Where Is Our Research Going? Brain Behav Evol. 2004;64: 163–181.
doi:10.1159/000079745

29. Tsukamoto Y, Yoshino S. A Study of the Lateral-line System in Fish II. On the Relation
between the Function of the Lateral-line Organ and the Habits of Fish. Japanese Journal of
Ichthyology. 1957;6: 131–135. doi:10.11369/jji1950.6.131

30. Erickson T, Pacentine IV, Venuto A, Clemens R, Nicolson T. The lhfpl5 Ohnologs lhfpl5a
and lhfpl5b Are Required for Mechanotransduction in Distinct Populations of Sensory
Hair Cells in Zebrafish. Front Mol Neurosci. 2020;12: 320. doi:10.3389/fnmol.2019.00320

31. McCune AR, Carlson RL. Twenty ways to lose your bladder: Common natural mutants in
zebrafish and widespread convergence of swim bladder loss among teleost fishes. Evol
Dev. 2004;6: 246–259. doi:10.1111/J.1525-142X.2004.04030.X

32. Abbas L, Whitfield TT. Nkcc1 (Slc12a2) is required for the regulation of endolymph
volume in the otic vesicle and swim bladder volume in the zebrafish larva. Development.
2009;136: 2837–2848. doi:10.1242/DEV.034215

33.   Westerfield M. The zebrafish book. A guide for the laboratory use of zebrafish (Danio
rerio). 4th ed. Eugene OR: University of Oregon; 2000. Available:
http://zfin.org/zf_info/zfbook/zfbk.html
34.   Erickson T, Morgan CP, Olt J, Hardy K, Busch-Nentwich E, Maeda R, et al. Integration of
Tmc1/2 into the mechanotransduction complex in zebrafish hair cells is regulated by
Transmembrane O-methyltransferase (Tomt). Elife. 2017;6. doi:10.7554/eLife.28474
35.   Kwan KM, Fujimoto E, Grabher C, Mangum BD, Hardy ME, Campbell DS, et al. The
Tol2kit: a multisite gateway-based construction kit for Tol2 transposon transgenesis
constructs. Dev Dyn. 2007;236: 3088–3099. doi:10.1002/DVDY.21343
36.   Venuto A, Erickson T. Evaluating the Death and Recovery of Lateral Line Hair Cells
Following Repeated Neomycin Treatments. Life 2021, Vol 11, Page 1180. 2021;11: 1180.
doi:10.3390/LIFE11111180
37.   Johansen EM. The Interfacial Tension between Petroleum Products and Water. Ind Eng
Chem. 1924;16: 132–135.
38.   Ehrlich DE, Schoppik D. Control of Movement Initiation Underlies the Development of
Balance. Current Biology. 2017;27: 334–344. doi:10.1016/j.cub.2016.12.003
39.   R Core Team. R: A Language and Environment for Statistical Computing. Vienna,
Austria; 2022. Available: https://www.r-project.org/
40.   Goolish EM, Okutake K. Lack of gas bladder inflation by the larvae of zebrafish in the
absence of an air-water interface. J Fish Biol. 1999;55: 1054–1063. doi:10.1111/j.1095-
8649.1999.tb00740.x
41.   v. Ledebur JF, Wunder W. Beiträge zur Physiologie der Schwimmblase der Fische.
Zeitschrift für vergleichende Physiologie 1937 25:2. 1937;25: 149–155.
doi:10.1007/BF00339307
42.   Jones FRH, Marshall NB. The structure and functions of the teleostean swimbladder.
Biological Reviews. 1953;28: 16–82. doi:10.1111/j.1469-185X.1953.tb01370.x
43.   Furukawa S, Machida Y, Takeuchi K, Hoshikawa Y, Irie K. Failure to gulp surface air
induces swim bladder adenomas in Japanese medaka (Oryzias latipes). J Toxicol Pathol.
2022;35: 237–246. doi:10.1293/TOX.2022-0030
44.   Korsøen ØJ, Fosseidengen JE, Kristiansen TS, Oppedal F, Bui S, Dempster T. Atlantic
salmon (Salmo salar L.) in a submerged sea-cage adapt rapidly to re-fill their swim
bladders in an underwater air filled dome. Aquac Eng. 2012;51: 1–6.
doi:10.1016/J.AQUAENG.2012.04.001
45.   Lindsey BW, Dumbarton TC, Moorman SJ, Smith FM, Croll RP. Effects of simulated
microgravity on the development of the swimbladder and buoyancy control in larval
zebrafish (Danio rerio). J Exp Zool A Ecol Genet Physiol. 2011;315A: 302–313.
doi:10.1002/jez.677
46.   Murakami SL, Cunningham LL, Werner LA, Bauer E, Pujol R, Raible DW, et al.
Developmental differences in susceptibility to neomycin-induced hair cell death in the
lateral line neuromasts of zebrafish (Danio rerio). Hear Res. 2003;186: 47–56.
doi:10.1016/S0378-5955(03)00259-4
47.   Santos F, MacDonald G, Rubel EW, Raible DW. Lateral line hair cell maturation is a
determinant of aminoglycoside susceptibility in zebrafish (Danio rerio). Hear Res.
2006;213: 25–33. doi:10.1016/j.heares.2005.12.009
48.   Hernández PP, Moreno V, Olivari FA, Allende ML. Sub-lethal concentrations of
waterborne copper are toxic to lateral line neuromasts in zebrafish (Danio rerio). Hear Res.
2006;213: 1–10. doi:10.1016/j.heares.2005.10.015

49. Mackenzie SM, Raible DW. Proliferative Regeneration of Zebrafish Lateral Line Hair
Cells after Different Ototoxic Insults. PLoS One. 2012;7: 47257.
doi:10.1371/JOURNAL.PONE.0047257
50. Baker CF, Montgomery JC. The sensory basis of rheotaxis in the blind Mexican cave fish,
Astyanax fasciatus. J Comp Physiol A. 1999;184: 519–527. doi:10.1007/s003590050351
51. Pavlov DS, Tyuryukov SN. The role of lateral-line organs and equilibrium in the behavior
and orientation of the dace, Leuciscus leuciscus, in a turbulent flow. J Ichthyol. 1993;33:
45–55.
52. Coffin AB, Ramcharitar J. Chemical Ototoxicity of the Fish Inner Ear and Lateral Line
BT - Fish Hearing and Bioacoustics: An Anthology in Honor of Arthur N. Popper and
Richard R. Fay. In: Sisneros JA, editor. Cham: Springer International Publishing; 2016.
pp. 419–437. doi:10.1007/978-3-319-21059-9_18
53. Richard MJ. Comptes rendus des séances de la Société de biologie et de ses filiales.
Paris,Masson [etc.]; 1896. Available: https://www.biodiversitylibrary.org/item/100656
54. Jung J, Serrano-Rojas SJ, Warkentin KM. Multimodal mechanosensing enables treefrog
embryos to escape egg-predators. J Exp Biol. 2020;223. doi:10.1242/jeb.236141
55. Mo W, Chen F, Nechiporuk A, Nicolson T. Quantification of vestibular-induced eye
movements in zebrafish larvae. BMC Neurosci. 2010;11: 110. doi:10.1186/1471-2202-11-
110
56. Bagnall MW, Schoppik D. Development of vestibular behaviors in zebrafish. Curr Opin
Neurobiol. 2018;53: 83–89. doi:10.1016/j.conb.2018.06.004
57. Olive R, Wolf S, Dubreuil A, Bormuth V, Debrégeas G, Candelier R. Rheotaxis of Larval
Zebrafish: Behavioral Study of a Multi-Sensory Process. Front Syst Neurosci. 2016;10.
doi:10.3389/fnsys.2016.00014
58. Müller U, Schwartz E. Influence of single neuromasts on prey localizing behavior of the
surface feeding fish, Aplocheilus lineatus. Journal of Comparative Physiology A.
1982;149: 399–408. doi:10.1007/BF00619155
59. Schwartz E, Hasler AD. Perception of Surface Waves by the Blackstripe Topminnow,
Fundulus notatus. Journal of the Fisheries Research Board of Canada. 1966;23: 1331–
1352. doi:10.1139/f66-124
60. Bleckmann H. Prey Identification and Prey Localization in Surface-feeding Fish and
Fishing Spiders. Sensory Biology of Aquatic Animals. New York, NY: Springer New
York; 1988. pp. 619–641. doi:10.1007/978-1-4612-3714-3_24
61. Suchocki CR, Sepulveda-Villet OJ. The role of phototaxis in the initial swim bladder
inflation of larval yellow perch (Perca flavescens). Int Aquat Res. 2019;11: 33–42.
doi:10.1007/s40071-019-0217-x
62. Liao JC. The role of the lateral line and vision on body kinematics and hydrodynamic
preference of rainbow trout in turbulent flow. J Exp Biol. 2006;209: 4077–90.
doi:10.1242/jeb.02487
63. Bak-Coleman J, Court A, Paley DA, Coombs S. The spatiotemporal dynamics of
rheotactic behavior depends on flow speed and available sensory information. J Exp Biol.
2013;216: 4011–24. doi:10.1242/jeb.090480
64. Coombs S, Bak-Coleman J, Montgomery J. Rheotaxis revisited: a multi-behavioral and
multisensory perspective on how fish orient to flow. Journal of Experimental Biology.
2020;223. doi:10.1242/JEB.223008
65. New JG. Multimodal Integration in the Feeding Behaviors of Predatory Teleost Fishes.
Brain Behav Evol. 2002;59: 177–189. doi:10.1159/000064905

66.   Nelson ME, MacIver MA, Coombs S. Modeling Electrosensory and Mechanosensory
Images during the Predatory Behavior of Weakly Electric Fish. Brain Behav Evol.
2002;59: 199–210. doi:10.1159/000064907

67.   Free BA, McHenry MJ, Paley DA. Probabilistic analytical modelling of predator–prey
interactions in fishes. J R Soc Interface. 2019;16: 20180873. doi:10.1098/rsif.2018.0873

68.   McCormick CA. Central Lateral Line Mechanosensory Pathways in Bony Fish. The
Mechanosensory Lateral Line. 1989; 341–364. doi:10.1007/978-1-4612-3560-6_17

69.   Mirjany M, Faber DS. Characteristics of the anterior lateral line nerve input to the
Mauthner cell. Journal of Experimental Biology. 2011;214: 3368–3377.
doi:10.1242/JEB.056226

70.   Hale ME, Katz HR, Peek MY, Fremont RT. Neural circuits that drive startle behavior,
with a focus on the Mauthner cells and spiral fiber neurons of fishes. https://doi-
org.proxy.hil.unb.ca/101080/0167706320161182526. 2016;30: 89–100.
doi:10.1080/01677063.2016.1182526

71.   Alexandre D, Ghysen A, Ghysen A. Somatotopy of the lateral line projection in larval
zebrafish. Proc Natl Acad Sci U S A. 1999;96: 7558–62. doi:10.1073/pnas.96.13.7558

72.   Liao JC, Haehnel M. Physiology of afferent neurons in larval zebrafish provides a
functional framework for lateral line somatotopy. J Neurophysiol. 2012;107: 2615–2623.
doi:10.1152/jn.01108.2011

73.   Fame RM, Brajon C, Ghysen A. Second-order projection from the posterior lateral line in
the early zebrafish brain. Neural Dev. 2006;1: 4. doi:10.1186/1749-8104-1-4

74.   Mueller T. What is the thalamus in zebrafish? Frontiers in Neuroscience. Frontiers; 2012.
p. 64. doi:10.3389/fnins.2012.00064

75.   Kappel JM, Förster D, Slangewal K, Shainer I, Svara F, Donovan JC, et al. Visual
recognition of social signals by a tectothalamic neural circuit. Nature 2022 608:7921.
2022;608: 146–152. doi:10.1038/s41586-022-04925-5

76.   Migault G, Plas TL van der, Trentesaux H, Panier T, Candelier R, Proville R, et al. Whole-
Brain Calcium Imaging during Physiological Vestibular Stimulation in Larval Zebrafish.
Current Biology. 2018;28: 3723-3735.e6. doi:10.1016/J.CUB.2018.10.017

77.   Favre-Bulle IA, Vanwalleghem G, Taylor MA, Rubinsztein-Dunlop H, Scott EK. Cellular-
Resolution Imaging of Vestibular Processing across the Larval Zebrafish Brain. Current
Biology. 2018;28: 3711-3722.e3. doi:10.1016/j.cub.2018.09.060

78.   Isa T, Marquez-Legorreta E, Grillner S, Scott EK. The tectum/superior colliculus as the
vertebrate solution for spatial sensory integration and action. Current Biology. 2021;31:
R741–R762. doi:10.1016/j.cub.2021.04.001

79.   Filosa A, Barker AJ, Dal Maschio M, Baier H. Feeding State Modulates Behavioral
Choice and Processing of Prey Stimuli in the Zebrafish Tectum. Neuron. 2016;90: 596–
608. doi:10.1016/J.NEURON.2016.03.014

80.   Perrault TJ, Stein BE, Rowland BA. Non-stationarity in multisensory neurons in the
superior colliculus. Front Psychol. 2011;2. doi:10.3389/FPSYG.2011.00144

81.   King AJ, Schnupp JWH, Carlile S, Smith AL, Thompson ID. The development of
topographically-aligned maps of visual and auditory space in the superior colliculus. Prog
Brain Res. 1996;112: 335–350. doi:10.1016/S0079-6123(08)63340-3

82.   Ingle D. Evolutionary perspectives on the function of the optic tectum. Brain Behav Evol.
1973;8: 211–237. doi:10.1159/000124355

83.   Thompson AW, Vanwalleghem GC, Heap LA, Scott EK. Functional profiles of visual-,
699   auditory-, and water flow-responsive neurons in the Zebrafish Tectum. Current Biology.
700   2016;26: 743–754. doi:10.1016/j.cub.2016.01.041
701   84.   Ehrlich DE, Schoppik D. A primal role for the vestibular sense in the development of
702   coordinated locomotion. Elife. 2019;8. doi:10.7554/eLife.45839
703   85.   Liu Z, Hildebrand DGC, Morgan JL, Slimmon N, Bagnall MW. The organization of the
704   gravity-sensing system in zebrafish. bioRxiv. 2021; 2021.07.09.451839.
705   doi:10.1101/2021.07.09.451839
706   86.   Chagnaud BP, Engelmann J, Fritzsch B, Glover JC, Straka H. Sensing External and Self-
707   Motion with Hair Cells: A Comparison of the Lateral Line and Vestibular Systems from a
708   Developmental and Evolutionary Perspective. Brain Behav Evol. 2017;90: 98–116.
709   doi:10.1159/000456646
710
711

Figures

Figure 1. Hyperinflation of the swim bladder in lateral line mutants (*lhfpl5b*[-/-]). A) Representative examples of over-, regular, and under-inflated swim bladders (arrows) in larval zebrafish at 6 dpf (top: *lhfpl5b*[-/-], others are wild type). B) Correlation between buoyancy and swim bladder volume. Line of best fit applied through the combination of mutant and wild type data (linear model, R-squared = 0.8574). C) Percentage of larvae that exhibit over-, regular, and under-inflation of the swim bladder, as defined by positive, neutral, and negative buoyancy. A Chi-Squared Test was used to determine significance. D) Swim bladder volume (mm^3) in wild type and *lhfpl5b* mutant larvae at 6 dpf. A Welch's t-test was used to determine significance. *** = $p < 0.001$, **** = $p < 0.0001$. Scale bar = 0.2 mm.

725
Figure 2: Blocking access to the surface eliminates the hyperinflation phenotype in lateral line
mutants (*lhfpl5b*[-/-]). A) Diagram of experimental set up with a filter blocking access to the surface.
B) Percentage of larvae that exhibit over-, regular, and under-inflation of the swim bladder at 6
dpf. A Chi-Squared Test was used to determine significance. C) Swim bladder volume (mm^3) of
larvae at 6 dpf. A One-Way ANOVA was used to determine significance. **** = p < 0.0001, ,
**** = p < 0.0001, ns = no significance. Full ANOVA table in supplemental tables.
732

733
734 **Figure 3**: Rescue of lateral line mutants (*lhfpl5b⁻/⁻*) with the *Tg(myo6b:EGFP-lhfpl5a)vo23*
735 transgene. A) Percentage of larvae that exhibit swim bladder under, regular, and over-inflation at
736 6 dpf. A Chi-Squared Test was used to determine significance. B) Swim bladder volume (mm^3) of
737 larvae at 6 dpf. A One-Way ANOVA was used to determine significance. **** = $p < 0.0001$, ns =
738 no significance. Full ANOVA table in supplemental tables.
739

740
Figure 4: Head-specific ablations of the lateral line recapitulates the *lhfpl5b* mutant hyperinflation
phenotype. A) Diagram of experiment. B) Percentage of larvae that exhibit swim bladder under,
regular, and over-inflation at 5 dpf. A Chi-Squared Test was used to determine significance. C)
Swim bladder volume (mm^3) of larvae at 5 dpf. A One-Way ANOVA was used to determine
significance. * = $p < 0.05$, , **** = $p < 0.0001$, ns = no significance. Full ANOVA table in
supplemental tables. Images of live larvae following treatment in Supplemental Figure 3.
747

748
749 **Figure 5**: Decreasing interfacial surface tension with mineral oil results in over-filling of the swim
750 bladder. A) Diagram of experiment with 2 mm layer of oil on surface. B) Images of swim bladder
751 phenotypes resulting from surface oil and surface air exposure in wild type larvae at 6 dpf. C) Swim
752 bladder volume (mm^3) at 6 dpf. A One-Way ANOVA was used to determine significance. * = p <
753 0.05, ns = no significance. Scale bar = 0.1 mm. Full ANOVA table in supplemental tables.
754

Figure 6: Optogenetic activation of lateral line hair cells with Channelrhodopsin-2 (ChR2) in wild type and *lhfpl5b* mutant larval zebrafish. A) Diagram of experiment with blue light overhead flashing for 25 ms with 1 s intervals. Treatments were started at 4 dpf and swim bladder volume was assessed at 6 dpf. B) Swim bladder volume (mm^3) of larvae at 6 dpf, n = 16-21 per condition. One-Way ANOVA was used to determine significance. ** = $p < 0.001$, ns = no significance. Full ANOVA table in supplemental tables.

763
Figure 7: Quantification of the surfacing behaviors of wild type and *lhfpl5b* mutant zebrafish larvae
during initial swim bladder inflation. A) Total number of individual visits taken by larval fish to
the air-water interface on 4 dpf for 12 hours (9 AM – 9 PM). B) Total amount of time larval fish
spend at the surface for all visits combined on 4 dpf. C) Time to first surface visit after access was
allowed. A One-Way ANOVA was used to determine significance. *** = p < 0.001, ns = no
significance.
770

771
Figure 8. Summary of the surfacing behavior of wild type and lateral line-deficient zebrafish
larvae. A) Wild type larvae use their lateral line hair cells (in green) to mediate interactions with
the air-water interface during surfacing. As a result of accurate surface detection, larvae take in an
appropriate volume of air for swim bladder (SB) inflation and achieve neutral buoyancy. B) Larvae
with a genetic loss of lateral line function (hair cells in white) misinterpret the air-water interface,
leading to increased interactions with the surface. Consequently, lateral line mutants *(lhfpl5b⁻/⁻)*
take in an excess volume of air, resulting in hyperinflation of the swim bladder for approximately
half of mutant larvae. Selective chemical ablations of the head neuromasts produce similar results,
implicating the anterior lateral line as the primary sensory organ for interactions with the air-water
interface during the surfacing behavior.
782

783   **Supplemental Figures**

784
Supplemental Figure 1. Buoyancy status of larvae raised in total darkness. Percentage of larvae
that exhibit swim bladder under, regular, and over-inflation raised in either a 14:10 light/dark
cycle (Control) or total darkness (0:24 light/dark cycle). A Chi-Squared Test was used to assess
significance. Both Wild Type Control and Wild Type Dark over-inflation proportions: $M = 0.0\%$,
$SD = 0.0\%$, $n = 32$. *lhfpl5b*$^{-/-}$ Control over-inflation proportion: $M = 57.5\%$, $SD = 6.6\%$, $n = 30$.
lhfpl5b$^{-/-}$ Dark over-inflation proportion: $M = 52.4\%$, $SD = 4.1\%$, $n = 30$.
791

Supplemental Figure 2. Chemical ablations of lateral line hair cells phenocopies the hyperinflation phenotype of *lhfpl5b* mutants. A) Percentage of larvae treated with either neomycin or CuSO4 that exhibit swim bladder under, regular, and over-inflation. A Chi-Squared Test was used to assess significance. B) Swim bladder volume (mm^3) in 5 dpf larvae treated with either neomycin or CuSO4. A One-Way ANOVA was used to determine significance. **** $= p < 0.0001$, *** $= p < 0.001$, ** $= p < 0.01$, * $= p < 0.05$, ns = no significance. A full ANOVA table can be found in the supplemental tables.

801
802 **Supplemental Figure 3.** Head and trunk-specific lateral line ablations with CuSO4. Representative
803 images of live *Tg(myo6b:EGFP-PA)vo68* transgenics at 4 dpf, two hours after CuSO4 treatment.
804 This transgene labels hair cells of the lateral line and inner ear (indicated by white brackets on the
805 Control image). CuSO4 treatment ablates lateral line hair cells only, leaving inner ear hair cells
806 intact.
807

808
Supplemental Figure 4: A) Representative images of Channelrhodopsin-2 (ChR2) expression in
neuromast, utricle, and crista hair cells of *Tg(lhfpl5b2028:ChR2-EYFP-PA)* transgenics at 2 dpf.
ChR2-YFP expression is detectable only in the lateral line neuromasts, indicating specificity of the
lhfpl5b regulatory region used to drive ChR2 expression. First panel is YFP only (false-colored
green) and the second panel is a merge of YFP and DIC. Scale bars = 10 uM. B) Swim bladder
volume (mm^3) of 6 dpf wild type and *lhfpl5b^{-/-}* larvae, with and without the *ChR2* transgene. Groups
of larvae were also treated with CuSO4 to ablate lateral line hair cells. All larvae were stimulated
with blue light. The impacts of blue light stimulation on wild type ChR2 transgenics is abolished
by chemical ablation of lateral line hair cells. As expected, *lhfpl5b* mutants are unaffected by
CuSO4 treatment. A One-Way ANOVA was used to determine significance. ** = $p < 0.00724$, ns
= no significance. A full ANOVA table can be found in the supplemental tables.
820