A sensation for inflation: initial swim bladder inflation in larval zebrafish is mediated

by the mechanosensory lateral line.

Alexandra Venuto[1], Stacey Thibodeau-Beganny[2], Josef G. Trapani[2], Timothy Erickson[3]

1. Department of Biology, East Carolina University, Greenville, NC, USA.

2. Department of Biology and Neuroscience Program, Amherst College, Amherst, MA, USA.

3. Department of Biology, University of New Brunswick, Fredericton, NB, Canada.

Abstract

Larval zebrafish achieve neutral buoyancy by swimming up to the surface and taking in air

through their mouths to inflate their swim bladders. We define this behavior as 'surfacing'. Little

is known about the sensory basis for this underappreciated behavior of larval fish. A strong

candidate is the mechanosensory lateral line, a hair cell-based sensory system that detects

hydrodynamic information from sources like water currents, predators, prey, and surface waves.

However, a role for the lateral line in mediating initial inflation of the swim bladder has not been

reported.

To explore the connection between the lateral line and surfacing, we utilized a genetic

mutant (*lhfpl5b*[-/-]) that renders the zebrafish lateral line insensitive to mechanical stimuli. We

observe that approximately half of these lateral line mutants over-inflate their swim bladders during

initial inflation and become positively buoyant. Thus, we hypothesize that larval zebrafish use their

lateral line to moderate interactions with the air-water interface during surfacing to regulate swim

bladder inflation. To test the hypothesis that lateral line defects are responsible for swim bladder

over-inflation, we show exogenous air is required for the hyperinflation phenotype and transgenic

rescue of hair cell function restores normal inflation. We also find that chemical ablation of anterior

lateral line hair cells in wild type larvae causes hyperinflation. Furthermore, we show that

manipulation of lateral line sensory information results in abnormal inflation. Finally, we report

spatial and temporal differences in the surfacing behavior between wild type and lateral line mutant

larvae. In summary, we propose a novel sensory basis for achieving neutral buoyancy where larval

zebrafish use their lateral line to sense the air-water interface and regulate initial swim bladder

inflation.

Keywords: behavior, buoyancy, hair cells, hydrodynamics, lateral line, lhfpl5, swim bladder,

sensory system, zebrafish

Introduction

The swim bladder of teleost fish provides for neutral buoyancy in the water column, thereby minimizing the energy expenditure associated with locomotion (Alexander, 1966). Larval teleosts initate swim bladder inflation by swimming up to the surface to ingest air using a set of behaviors we refer to as "surfacing" (Evans and Damant, 1928; Lindsey et al., 2010; von Ledebur, 1928). Although surfacing poses a significant risk of predation, larval fish are highly motivated to reach the surface to intake air. For example, pre-inflated lake trout larvae will swim hundreds of feet at a constant rate in attempts to access the surface (Tait, 1960). Surfacing is a complex task that requires a combination of newly developed physical features, sensory systems, and motor skills (Bailey and Doroshov, 1995; Chatain, 1989; Doroshev et al., 1981; Kimmel et al., 1995; Kitajima et al., 1994) – (i) Larvae must discriminate up from down and (ii) be able to move directionally towards the surface; (iii) they must appropriately sense the air-water interface, and (iv) be able to intake air through the mouth upon arrival at the surface. Ingested air is then moved by peristalsis into the swim bladder through the pneumatic duct that connects the gut and swim bladder (Doroshev et al., 1981; Rieger and Summerfelt, 1998; Tait, 1960). Together, surfacing represents perhaps the most complex behavior that larvae perform prior to achieving neutral buoyancy (Fero et al., 2011).

Little is known about the sensory cues that instruct the surfacing behavior. Perturbations to the vestibular system typically result in hypoinflation (Nicolson et al., 1998; Schoppik et al., 2017). The inability to detect gravity by the utricular otolith is specifically responsible for the phenotype, presumably because larvae are unable to orient their movements upwards towards the surface (Riley and Moorman, 2000). Photosensory cues may also play a role in directional orientation and surface detection. However, which photosensory systems are involved has not been defined and the effects of altering lighting conditions varies between species (Stuart and Drawbridge, 2012; Trotter et al., 2003; Villamizar et al., 2009). Even less is known about how larval fish detect the air-water interface itself. Fish and some amphibians possess an additional hair cell-based sensory system - the *mechanosensory lateral line* - that contributes to detection of near-field hydrodynamic information originating from water currents, predators and prey, conspecifics, abiotic objects, and surface waves (Anneser et al., 2020; Bleckmann, 2006; Butler and Maruska, 2016; Carrillo and McHenry, 2016; Mogdans, 2019; Montgomery et al., 1997; Stewart et al., 2013; Venuto et al., 2022). Specific to surface detection, surface-feeding fish use their lateral line to detect surface

waves created by their prey (Bleckmann and Schwartz, 1982). Furthermore, Japanese flying fish

are predicted to use their ventrally-located lateral line to sense transitions through the air-water

interface (Gibbs, 2004; Tsukamoto and Yoshino, 1957). While the lateral line is a strong candidate

to aid in surfacing behaviors that lead to initial swim bladder inflation in larval fish, its specific

role in this important behavior has not been explored.

A CRISPR-Cas9 knockout of *LHFPL tetraspan subfamily member 5b* (*lhfpl5b*) is the first

genetic zebrafish mutant where the lateral line is non-functional from birth but hearing and balance

are normal (Erickson et al., 2020). These mutants provide an opportunity to uncover novel roles

for the lateral line. Unexpectedly, homozygous mutant larvae present with a hyperinflation

phenotype characterized by over-filled swim bladders and positive buoyancy, a very rare finding

among swim bladder defects (Abbas and Whitfield, 2009; McCune and Carlson, 2004). Using

several genetic and chemical techniques to manipulate lateral line function, we show that lateral

line sensory information is used by larval zebrafish to appropriately fill their swim bladders, likely

by detecting surface tension at the air-water interface using mechanosensory neuromasts located

on their heads. Overall, our study uncovers a novel role for the mechanosensory lateral line

mediating initial inflation of the swim bladder.

Methods

Animal husbandry and ethics statement

Adult zebrafish (*Danio rerio*) were maintained and bred using standard procedures

(Westerfield, 2000). All experiments used larvae at 2-6 days post-fertilization (dpf), which are of

indeterminate sex at this stage. Except where otherwise stated, animals were placed in clear, plastic

tubs (dimensions: 15 cm x 10 cm x 4 cm, E3 volume: 460 mL) at 2 dpf and maintained at 28.5°C

on a 14:10 light/dark cycle at 500-700 lux in E3 embryo media (5 mM NaCl, 0.17 mM KCl, 0.33

mM CaCl2, 0.33 mM MgCl2, buffered with NaHCO3). Animal research complied with guidelines

stipulated by the Institutional Animal Care and Use Committees at East Carolina University

(protocol D372) and Amherst College (assurance number 3925-1 with the Office of Laboratory

Animal Welfare).

Mutant and transgenic fish lines

The zebrafish mutant allele *lhfpl5b^{vo35}* and transgenic lines *Tg(myo6b:EGFP-lhfpl5a)vo23*

and *Tg(myo6b:EGFP-PA)vo68* were used in this study (Erickson et al., 2017; Erickson et al.,

2020). For all experiments, *lhfpl5b^{vo35}* homozygotes were identified from their hetero- and

homozygous wild type siblings by lack of FM 1-43 (ThermoFisher) labeling of the lateral line hair

cells.

To create a fish line that expresses the Channelrhodopsin-2 (ChR2) optogenetic protein

solely in the hair cells of the lateral line, approximately 2 kb of DNA upstream of the start codon

of the *lhfpl5b* gene was cloned into the p5E plasmid of the recombination-based multisite Gateway

cloning system (ThermoFisher). A Tol2 transposon backbone was then used to create the injectable

expression plasmid *lhfpl5b2028:ChR2-EYFP-PA*. This plasmid was then co-injected with

transposase mRNA into single-cell embryos (Kwan et al., 2007) to create the

Tg(lhfpl5b2028:ChR2-EYFP-PA) zebrafish line.

Buoyancy tests

To ascertain the buoyancy status of larval fish, larvae were anesthetized with MS-222

(Western Chemical Inc., Ferndale, WA) and placed at the vertical midway point of a 50 mL conical

tube containing E3 media. At the expiration of a 30-second timer, the end position of the fish was

recorded and the fish was removed from the tube for imaging under the microscope. If the fish

reached the top or bottom of the tube before 30 seconds, then the time it took to reach that location

was recorded. The total height of the water column was 8.9 cm, and the fish was placed half-way

at 4.45 cm. The rate of buoyancy was determined by dividing the centimeters travelled by the

seconds it took to travel that distance (cm/s).

Imaging and swim bladder measurements

Swim bladders were imaged using a SteREO Discovery.V8 microscope (Zeiss) equipped

with a Gryphax Arktur camera (Jenoptik). Lateral and dorsal images of the swim bladder were

taken for each fish. Swim bladder measurements were conducted using the measurement tool in

Adobe Photoshop where measurements were calibrated according to the image of the stage

micrometer taken when imaging experiments. The major axis and minor axis of the lateral view of

the swim bladder was measured as well as the minor axis of the dorsal view. All measurements

were taken from numbered images so that the measurer was blinded to the genotype/condition.

These meaurements were halved and used in the equation for the volume of an ellipsoid: $V =$

$4/3\pi abc$ to find the volume of the swim bladder (Lindsey et al., 2010).

Manipulation of photosensory cues during surfacing

At 0 dpf, zebrafish embryos were collected and immediately placed in plastic tubs filled

with E3 (detailed above) in an unlit incubator. Larvae were not brought into light until 6 dpf, at

which point lateral line mutants and wild type siblings were identified by FM 1-43 labeling. After

sorting, larvae were analyzed for buoyancy. The experiment was repeated three times using 20-21

larvae (combined mutant and wild type) per trial.

Blocking access to the air-water interface

Mutant larvae and wild type siblings were sorted at 2 dpf into glass cylinders (diameter: 9.5

cm, E3 volume: 425 mL) equipped with fitted, wire mesh filter plungers (Bodum Inc., USA).

Conditions were as follows: wild type with open access to the surface, wild type with blocked

access to the surface, mutant with open access to the surface, and mutant with blocked access to

the surface. All containers were filled half way with E3 and in the experimental condition ("blocked

access"), the filter was pressed down below the surface for both wild type and mutant larvae. The

control condition used the same amount of E3 but the filter remained above the surface so that

larvae could access the exogenous air. Conditions remained constant until 6 dpf when larvae were

imaged and analyzed for swim bladder inflation. The experiment was repeated three times using

15-25 larvae per condition.

Analysis of surfacing behavior

Wild type and *lhfpl5b^{vo35}* larvae were blocked from having access to the surface until 9 AM

on 4 dpf when individuals were immediately placed into clear rectangular containers (dimensions:

3 cm x 3 cm x 3 cm, E3 volume: 20 mL) for video recording. Recordings lasted 12 hours, from 9

AM to 9 PM on 4 dpf. At the end of filming, larvae were imaged and analyzed for buoyancy and

swim bladder volume, as described. All surfacing videos were numbered for blind analysis.

Surfacing analysis was conducted by manual observation of each individual larvae from the videos

(30 frames per second) and manually counting visits made to the surface (counted as reaching the

meniscus), the timeframe of the video at which the surfacing event occurred, and the duration of

time spent at the surface per surfacing event. Recording was repeated six times with 3 mutant and

2 wild type larvae per recording. Larvae that did not inflate their swim bladders by the end of

filming were excluded from this analysis due to lack of surfacing events.

Lateral line hair cell ablations

For all experiments, wild type and *lhfpl5b* mutant larvae were treated with ototoxin with

untreated siblings acting as controls (n = 7-15 larvae per condition per trial, 4 trials) For single

treatments, surface accessed was blocked as described above for untreated and treated larvae until

9 AM on 4 dpf when treatment occurred. Single 30 minute treatments with neomycin sulfate (EMD

Millipore Corp., USA) were done with a 100 µm solution in E3 while repeated neomycin

treatments (Venuto and Erickson, 2021) were done as previously described with a concentration of

50 µM. For full ablation with copper sulfate (CuSO4; Ward's Science, Rochester, NY), the

treatment concentration was 10 uM with a 30 minute exposure. For experiments where head and

trunk neuromasts were selectively ablated, we anesthetized the larvae and pipetted them on agarose

gel. Excess water was removed and a thin layer of petroleum jelly (Vaseline, Unilever USA) was

placed horizontally at the back of the head. Next, 30 uM CuSO4 or E3 was pipetted onto the

appropriate region depending on the experimental condition. Immediately follow the 13 minute

treatment, larvae were placed in fresh E3 in clear tubs detailed above for 24 hours until 9 AM on 5

dpf when larvae were imaged and analyzed for swim bladder inflation. For all experiments, larvae

were imaged and analyzed for swim bladder inflation at 9 AM on 5 dpf. Each experiment was

repeated three times using 10-24 larvae per condition.

Transgenic rescue of lateral line mutant

Larval zebrafish from crosses of *lhfpl5b^{+/-}* and *Tg(myo6b:EGFP-lhfpl5a)vo23Tg; lhfpl5b^{+/-}*

were sorted at 2 dpf for transgene expression. At 6 dpf, larvae were numbered, imaged for swim

bladder inflation and genotyped for the *lhfpl5b^{vo35}* allele as previously described (Erickson et al.,

2020). The experiment was repeated twice using 26-48 larvae per condition.

Optogenetic activation of lateral line hair cells with Channelrhodopsin-2 (ChR2)

After blocking access to the surface until 4 dpf, we exposed wild type and lateral line

mutants, with or without the *lhfpl5b2028:ChR2-EYFP-PA* transgene, to LED blue light (470 nm)

with an intensity of 730 lux over their holding tanks against a dark background. Light flashed for

25 milliseconds at one second intervals. Control larvae were kept in the same background, without

blue light exposure. Conditions were held constant until 6 dpf when larvae were imaged and

analyzed for swim bladder inflation. The experiment was repeated three times using 15-25 larvae

per condition.

Oil-water interface experiments

Mutant larvae and wild type siblings were sorted into clear cylindrical containers (diameter:
5 cm , volume: 60 mL) with embryo media only or embryo media with a 2 mm thick surface layer
of lab grade mineral oil (Ward's Science, Rochester, NY). Oil on the surface decreases surface
tension (Johansen, 1924) and has been previously used in experiments involving the zebrafish swim
bladder (Ehrlich and Schoppik, 2017). Conditions remained constant until 6 dpf when larvae were
imaged and analyzed for swim bladder volume. Because oil-filled larvae are negatively buoyant,
we did not perform buoyancy tests and classified larvae as over-filled based on swim bladder
volume alone. The experiment was repeated three times using 15-25 larvae per condition.

Graphs and statistical tests

All graphs and statistical tests were done using R (R Core Team, 2022). One-way ANOVAs
were used to compare the conditions as a whole (genotype independent variable plus environment
independent variable) with Tukey post-hoc test. Chi-squared tests were used to analyze proportion
data for over and under-inflation. For the over-inflation comparisons, the two categories used in
the comparison between conditions/genotypes were (i) over-inflation and (ii) all other inflation
(regular and under), meaning that totals per category/genotype accounted for 100% of the
population. Similarly, for the under-inflation comparisons, the two categories used in the
comparison between conditions/genotypes were (i) under-inflation and (ii) all other inflation
(regular and over). P-values less than 0.05 were considered significant.

Results

Lateral line mutants exhibit a swim bladder hyperinflation phenotype

The lateral line has not been previously implicated in initial inflation of the larval swim
bladder. As such, it was unexpected to observe that lateral line mutants ($lhfpl5b^{-/-}$) exhibit
hyperinflated swim bladders by 5-6 dpf (Figure 1A). We measured swim bladder volume and
buoyancy for each larva in the mutant and wild type populations (Figure 1B) and found that, on
average, 53.7% ($SD = 6.4\%$, $n = 41$) of lateral line mutants over-inflate their swim bladders and
7.3% ($SD = 7.1\%$, $n = 41$) under-inflate. For wild type larvae (including $lhfpl5b^{vo35}$ heterozygotes),
0.0% ($SD = 0.0\%$, $n = 41$) over-inflate and 4.9% ($SD = 3.7\%$, $n = 41$) under-inflate their swim
bladders during initial inflation on average (Figure 1C). The proportion of over-inflation is
significantly different between mutant and wild type larvae (chi-square statistic: 70.45, $p <$
0.00001), while the proportion of under-inflation is not (chi-square statistic: 0.3546, $p = 0.552$).

Average swim bladder volume of mutant fish is larger ($M = 0.012$ mm^3, $SD = 0.002$ mm^3) than

wild type siblings ($M = 0.0058$ mm^3, $SD = 0.00043$ mm^3, $t(80) = 17.4$, $p < 0.0001$) (Figure 1D).

The proportions of neutrally and positively buoyant larvae do not change when raised in total

darkness (Supplemental Figure 1). This unexpected hyperinflation phenotype in *lhfpl5b^{vo35}*

homozygotes led us to investigate if the lateral line is contributing important sensory information

during the surfacing behavior.

Access to the air-water interface is required for the hyperinflation phenotype of lateral line mutants

For most teleosts, initial swim bladder inflation requires an intake of exogenous air,

typically from the water's surface. To test if the hyperinflation phenotype of lateral line mutants

requires access to surface air, we physically blocked access to the surface from 2-6 dpf (Figure 2).

On average, 0% ($SD = 0.0\%$, $n = 43$) of blocked *lhfpl5b* mutants over-inflate their swim bladders,

which is significantly different from mutants with access to the surface (chi-square statistic: 70.45,

$p < 0.00001$; Figure 2B). The average swim bladder volume of blocked mutants ($M = 0.0033$ mm^3,

$SD = 0.0015$ mm^3) is significantly smaller than mutants with open access ($M = 0.014$ mm^3, $SD = $

0.0027 mm^3) (One-way ANOVA with Tukey post-test, $p < 0.00001$). Blocked mutant larvae had

statistically similar swim bladder volumes to blocked wild type larvae ($M = 0.0037$ mm^3, $SD = $

0.001 mm^3) (One-way ANOVA with Tukey post-test, $p = 0.927$) (Figure 2C). From these data, we

conclude that the *lhfpl5b* mutant hyperinflation phenotype requires intake of exogenous air and is

not caused by abnormal gas exchange from surrounding media nor excess internal gas production.

Transgenic rescue of lateral line mutants restores normal inflation

We next examined if lateral line defects are responsible for the hyper-inflation phenotype

of *lhfpl5b* mutants by restoring lateral line function in mutant larvae. We used the

Tg(myo6b:EGFP-lhfpl5a)vo23 transgene, which uses a hair cell specific promoter (*myo6b*) to drive

a GFP-tagged *lhfpl5a* gene (Erickson et al., 2017). It has been shown that this *lhfpl5a* transgene

rescues lateral line hair cell function in *lhfpl5b$^{-/-}$ (or* lateral line) mutant fish (Erickson et al., 2020).

Using this rescue method, we observe no cases of hyperinflation in transgenic mutants (0%, $SD = $

0.0%, $n = 17$), which is significantly different from non-transgenic mutants (chi-square statistic:

82.12, $p < 0.00001$; Figure 3A). The average volume of the transgenic mutant swim bladder ($M = $

0.0057 mm^3, $SD = 0.00001$ mm^3) is significantly smaller than non-transgenic, *lhfpl5b* mutant

siblings ($M = 0.0109$ mm^3, $SD = 0.0031$ mm^3) (One-way ANOVA with Tukey post-test, $p = $

0.000003), but statistically similar to that of wild type transgenics ($M = 0.0057$ mm^3, $SD = 0.0005$

mm^3) (One-way ANOVA with Tukey post-test, p = 0.979, Figure 3B). Because the restoration of

lhfpl5 function specifically in hair cells mitigates hyperinflation in *lhfpl5b* mutants, we conclude

that the phenotype is due to defects in sensory hair cell function and not caused by an unrecognized

role for *lhfpl5b* in another sensory organ or the swim bladder itself.

Selective ablation of head neuromasts produces a hyperinflation phenotype in wild type larvae

To further test if lateral line defects are responsible for the hyper-inflation phenotype seen

in *lhfpl5b* mutants, we ablated lateral line hair cells in wild type larvae using either neomycin

sulfate (neo) or copper sulfate (CuSO4) (Supplemental Figure 2 and Figure 4). Administering either

single or repeated neomycin treatments during the initial swim bladder inflation period of 3 - 4 dpf

results in a significant increase in the proportion of positively buoyant larvae (single neo chi-square

statistic: 9.95, p = 0.0016, repeated neo chi-square statistic: 34.88, p < 0.0001; Supplemental Figure

2A), but a non-significant increase in the average swim bladder volume (One-way ANOVA with

Tukey post-test, single neo p = 1.0, repeated neo p = 0.578; Supplemental Figure 2B). By

comparison, a single CuSO4 treatment results in a hyperinflation phenotype that is statistically

indistinguishable from lateral line mutants. An average of 59.6% (SD = 7.7%, n = 72) of CuSO4-

treated wild types over inflate their swim bladders, which is significantly different from wild type

controls (chi-square statistic: 39.69, p < 0.0001; Supplemental Figure 2A). The average swim

bladder volume of the CuSO4-treated wild type larvae (M = 0.0126 mm^3, SD = 0.0027 mm^3) is

statistically similar to the average swim bladder volume of untreated lateral line mutants (M =

0.0139 mm^3, SD = 0.0007 mm^3) (One-way ANOVA with Tukey post-test, p = 0.813)

(Supplemental Figure 2B).

Since the anterior lateral line is likely to be the only part of the sensory organ to interact

with the air-water interface during surfacing, we predict that disruption of the head neuromasts will

result in hyperinflation and that trunk neuromasts will not play a role (Figure 4, Supplemental

Figure 3). Selective ablation of the anterior region with CuSO4 results in an average of 35.9% (SD

= 10.1%, n = 38) hyperinflated larvae, which is significantly different from unablated controls (chi-

square statistic: 39.69, p < 0.0001; Figure 4B). Wild type larvae with head-specific ablations exhibit

a significant increase in average swim bladder volume (M = 0.016 mm^3, SD = 0.0013 mm^3)

compared to untreated wild type siblings (M = 0.0012 mm^3, SD = 0.0013 mm^3) (One-way ANOVA

with Tukey post-test, p = 0.0365) and statistically comparable swim bladder volumes to that of full

CuSO4-treated larvae (M = 0.016 mm^3, SD = 0.0029 mm^3) (One-way ANOVA with Tukey post-

test, *p* = 0.987) (Figure 4C). Conversely, tail-specific CuSO4 treatments do not cause

hyperinflation in wild type fish and result in statistically similar swim bladder volumes to untreated

siblings (M = 0.012 mm^3, SD = 0.0031 mm^3) (One-way ANOVA with Tukey post-test, *p* = 0.835)

(Figure 4C). The *lhfpl5b* mutant hyperinflation phenotype was not affected by either neomycin or

CuSO4 treatments (Supplemental Figure 2). Taken together, we conclude that the anterior region

of the lateral line system is critical to surface detection during initial swim bladder inflation.

Decreasing surface tension results in over-filling of the swim bladder

Surface tension is a characteristic that makes the air-water interface distinct from the rest

of the water column. We predict that larval fish sense this stimulus with their anterior lateral line

neuromasts and that by decreasing the interfacial tension, we would observe abnormal swim

bladder inflation in wild type larvae while lateral line mutants would be unaffected. The interfacial

tension between oil and water is approximately half that of air and water (Johansen, 1924). We

layered oil onto the surface of the embryo media and allowed larvae to perform surfacing behaviors

between 2-6 dpf (Figure 5). Using this experimental design, we find that the average swim bladder

volume of wild types exposed to an oil-water interface (M = 0.016 mm^3, SD = 0.0029 mm^3) is

significantly greater than the control group with access to an air-water interface (M = 0.007 mm^3,

SD = 0.001 mm^3) (One-way ANOVA with Tukey post-test, *p* < 0.0001). Furthermore, oil-exposed

wild types have statistically similar swim bladder volumes to that of *lhfpl5b* mutants (M = 0.014

mm^3, SD = 0.0038 mm^3) (One-way ANOVA with Tukey post-test, *p* = 0.886) (Figure 5C). Lateral

line mutants overfill their swim bladders to statistically similar proportions when presented with

either an air-water or oil-water interface (M = 0.01 mm^3, SD = 0.0026 mm^3) (One-way ANOVA

with Tukey post-test, *p* = 0.133). We conclude that surface tension is one of the surface

characteristics the lateral line is capable of detecting during initial swim bladder inflation.

Optogenetic activation of lateral line hair cells prevents hyperinflation in lateral line mutant larvae

If hydrodynamic information is informing the surfacing behaviors of larval fish, then

stimulus-independent manipulations of lateral line activity should alter their ability to correctly

inflate their swim bladders. To test this, we created the *Tg(lhfpl5b2028:ChR2-EYFP-PA)*

transgenic line that expresses the blue light activated channel, Channelrhodopsin-2, specifically in

lateral line hair cells (See Methods, Supplemental Figure 4). We predicted that optogenetic

stimulation of *lhfpl5b* mutants, whose lateral lines are insensitive to physical stimuli, would reduce

the frequency of the hyperinflation phenotype. Consistent with these predictions, under blue-light

stimulation, lateral line mutants with the *lhfpl5b:ChR2* transgene do not exhibit positive buoyancy and have a significant decrease in average swim bladder volume (M = 0.0046 mm^3, SD = 0.0034 mm^3) compared to unstimulated transgenic mutants (M = 0.012 mm^3, SD = 0.0031 mm^3) (One-way ANOVA with Tukey post-test, p = 0.00127) (Figure 6B). Transgenic wild types under blue light exhibit a non-significant decrease in swim bladder volume (M = 0.0052 mm^3, SD = 0.004 mm^3) compared to unstimulated transgenic siblings (M = 0.0079 mm^3, SD = 0.002 mm^3) (One-way ANOVA with Tukey post-test, p = 0.458). To demonstrate that blue light stimulation is reducing swim bladder volumes via activation of lateral line hair cells, we treated wild type *lhfpl5b:ChR2* larvae with CuSO4. As expected, we observe hyperinflation following CuSO4 treatment in stimulated wild type *lhfpl5b:ChR2* larvae, with an average swim bladder volume (WT: M = 0.012 mm^3, SD = 0.0037 mm^3) that is statistically similar to experimental non-transgenic *lhfpl5b* mutants (M = 0.011 mm^3, SD = 0.0024 mm^3) (One-way ANOVA with Tukey post-test, p = 0.998) (Supplemental Figure 4B). Overall, optogenetic activation of mechanically-insensitive lateral line organs in *lhfpl5b* mutants prevents the hyperinflation phenotype. Together, these data support the hypothesis that the lateral line provides crucial sensory information during initial swim bladder inflation in larval zebrafish.

Lateral line mutants exhibit abnormal surfacing behaviors

Our results suggest that larval fish are using their head neuromasts to sense the interfacial tension between air and water at the surface. Even in wild type zebrafish, the interaction with the surface during initial swim bladder inflation is not well characterized. To study this behavior in more detail, we quantified the number of unique visits to the surface and the amount of time per visit spent within the meniscus region from videos of wild type (n = 7) and mutant (n = 14) larvae at 4 dpf. *lhfpl5b* mutants that will over-inflate their swim bladders make more visits (M = 87.3, SD = 17.4) to the surface compared to both wild type (M = 42.0, SD = 16.3) and mutant siblings that will inflate to normal proportions (M = 31.9, SD = 19.2) (One-way ANOVA with Tukey post-test, p = 0.0004) (Figure 7A). Mutants destined for hyperinflation also spend more time (M = 139.0 s, SD = 31.6 s) at the surface than normally-inflated wild type (M = 51.9 s, SD = 24.7 s) and mutant siblings (M = 42.3 s, SD = 29.5 s) (One-way ANOVA with Tukey post-test, p = 0.00006) (Figure 7B). Since excess surfacing could be due to some mutants initiating the behavior earlier, we also documented the time of the first surfacing event and found no significant difference between neutrally buoyant and over-inflated larvae (Figure 7C). We conclude there is a correlation between

how larvae interact with the surface during the initial swim bladder inflation period and their
resulting swim bladder volume and buoyancy status.

Discussion

The intake of exogenous air from the surface is a common feature for intitial swim bladder
inflation in many species of fish (Furukawa et al., 2022; Goolish and Okutake, 1999; Jones and
Marshall, 1953; Korsøen et al., 2012; v. Ledebur and Wunder, 1937). However, few studies have
considered what sensory information is required to achieve neutral buoyancy. In this study, we
describe a new role for the mechanosensory lateral line in mediating initial inflation of the swim
bladder in zebrafish (Figure 8). Our data support a model in which larval zebrafish use their head
neuromasts to detect the air-water interface in support of their transition to neutral buoyancy.

Lateral line defects in lhfpl5b mutants are responsible for hyperinflation phenotypes

lhfpl5b mutant zebrafish are the first genetic model for the specific loss of lateral line
function in any organism (Erickson et al., 2020). These "lateral line mutants" exhibit an uncommon
swim bladder hyperinflation phenotype that prompted us to investigate this novel connection
between lateral line sensation and initial inflation of the swim bladder (Figure 1). Previous studies
have demonstrated that restricting access to surface air during the initial inflation period prevents
swim bladder inflation in larval zebrafish (Goolish and Okutake, 1999; Lindsey et al., 2011). In
addition, one study showed that some *solute carrier family 12 member 2 (slc12a2 / nkcc1)* mutant
zebrafish over-fill their swim bladders and that blocking access to the surface prevents this
phenotype (Abbas and Whitfield, 2009), though further studies are required to determine the cause
of over-inflation in *slc12a2* mutants. Likewise, we find that blocking access to the air-water
interface eliminates over-inflation in lateral line mutants and greatly decreases average swim
bladder volumes of both mutant and wild type larvae (Figure 2). Our method of blocking access to
the air-water interface resulted in a small proportion of larvae (both wild type and mutant) inflating
their swim bladder to normal proportions, though hyperinflation was never observed. Inflation was
likely achieved by larvae accessing air bubbles that remain in the water after submerging the filter
(Goolish and Okutake, 1999). Overall, the hyperinflation phenotype in lateral line mutants requires
an intake of exogenous air, pointing to a sensory defect in *lhfpl5b* mutants rather than a
physiological one.

In support of this sensory-deficit hypothesis, we show the hyperinflation phenotype is
prevented by either transgenic rescue of *lhfpl5* function specifically in hair cells or optogenetic

stimulation of mechanically-insensitive lateral line hair cells in *lhfpl5b* mutants (Figures 3 and 6). Furthermore, chemical ablation of lateral line hair cells in wild type larvae prior to initial inflation can phenocopy the *lhfpl5b* mutant. A single treatment with CuSO4 caused similar swim bladder over-inflation proportions to lateral line mutants (Supplemental Figure 2). Repeated neomycin treatments over 3-4 dpf caused the next highest percentage of swim bladder over-inflation and a single neomycin treatment only resulted in a few cases of wild type over-inflation. These results agree with previous work showing rapid hair cell regeneration following neomycin treatment and comparatively slower recovery after CuSO4 exposure (Hernández et al., 2006; Mackenzie and Raible, 2012; Murakami et al., 2003; Santos et al., 2006; Venuto and Erickson, 2021). Thus, via multiple genetic and chemical approaches, our data show that lateral line hair cell activity can modulate swim bladder inflation in larval zebrafish.

Ototoxins have been extensively used to study lateral line-mediated behaviors in free-swimming larvae and adult fish (Baker and Montgomery, 1999; Bleckmann, 2006; Coffin and Ramcharitar, 2016; Montgomery et al., 1997; Pavlov and Tyuryukov, 1993). However, we are not aware of any behavioral studies where the lateral line was disrupted prior to larvae achieving neutral buoyancy, likely explaining why this connection between the lateral line and initial swim bladder inflation has not been noted in current literature. Interestingly, an 1896 study predicted a relationship between the lateral line and swim bladder inflation, since disruption of the lateral line in adult goldfish resulted in "swelling of the swim bladder" and positive buoyancy (Richard, 1896). However, it is not clear if the phenotypic similarities between these adult goldfish and our experimental larval zebrafish share the same underlying causes. Regarding roles for the lateral line in early life behaviors, a close parallel to our findings is the report that unhatched red-eyed treefrogs use their lateral line organ to mediate mechanosensory-cued hatching (Jung et al., 2020). In this case, the frog lateral line can sense vibrational stimuli, such as that produced by predators, and promote escape hatching behaviors even before the onset of vestibular function. Similarly, our data suggest that the lateral line is modulating early locomotor activities in zebrafish prior to the development of fully formed vestibular or rheotactic responses (Bagnall and Schoppik, 2018; Mo et al., 2010; Olive et al., 2016). As such, the involvement of lateral line in the surfacing behavior may represent the earliest behavioral role for the lateral line yet described.

Head neuromasts are required for normal swim bladder inflation, likely by sensing surface tension

We expanded on the full-body lateral line ablation experiments with specific ablations of

either the head or truck neuromasts with CuSO4. Previous studies have shown that head neuromasts

are involved with detection of surface features and prey capture (Bleckmann, 1988; Carrillo and

McHenry, 2016; Müller and Schwartz, 1982; Schwartz and Hasler, 1966). Indeed, ablation of the

head neuromasts in wild type larvae recapitulated the hyperinflation phenotype of *lhfpl5b* mutants

while ablation of the tail neuromasts had no effect on either swim bladder volume or buoyancy

(Figure 4). These findings suggest that the head neuromasts interact with the air-water interface

during surfacing and that the posterior lateral line is dispensible for achieving neutral buoyancy.

As such, our findings also provide additional support for the anterior lateral line as a central

mediator of surface-related behaviors in fish.

The interfacial tension between air and water represents a unique hydrodynamic stimulus

that may be detected by the lateral line. The tension between oil and water is half that of air and

water (Johansen, 1924). Thus, by adding a layer of oil to the surface, we predicted that surface

detection would become increasingly difficult and result in over-filling of the swim bladder. We

find that, when given access to an oil-water interface, wild type larvae over-fill their swim bladders

with oil to a statistically similar volume as lateral line mutants who overinflate with air (Figure 6).

We interpret this result as support for the hypothesis that the lateral line senses interfacial tension

between air and water as larvae interact with the surface during initial swim bladder inflation. This

interpretation is further supported by the observation that *lhfpl5b* mutants (who are already

insensitive to interfacial tension) exhibit similar swim bladder volumes when exposed to either oil

or air. However, a possible complementary interpretation for these data is that larvae who fill their

swim bladders with oil may be deprived of the normal feedback on buoyancy levels as they intake

from the surface. Since oil is denser than air, oil-filled larvae are negatively buoyant. In this

scenario, larvae may continue to visit the surface for air intake in an attempt to achieve neutral

buoyancy. However, future studies are required to examine if such a feedback mechanism exists in

larval zebrafish.

Swim bladder over-inflation is correlated with excess time spent at the air-water interface

Since our data suggest that the air-water interface is providing a distinct stimulus to the

lateral line, we investigated whether surface interactions differed between hyperinflated *lhfpl5b*

mutants and their wild type counterparts. As predicted, over-inflated mutants took more visits *and*

spent more time at the surface than neutrally-buoyant mutant and wild type siblings (Figure 7).

Excess breaching of the surface occurred despite over-inflated and neutrally buoyant larvae

exhibiting similar timing for their first surfacing attempts. While it is unclear whether larvae ingest
air during the entirety of the duration spent above the air-water interface, it is known that surface
breaches are observed during initial inflation (Goolish and Okutake, 1999; Lindsey et al., 2010;
Rieger and Summerfelt, 1998). Therefore, the increase in surface visits likely leads to an increase
in air deposition to the swim bladder in lateral line mutants.

Multisensory integration during the surfacing behavior

Interestingly, approximately half of all lateral line mutants inflate their swim bladders to
normal proportions and continue to develop into viable adults. To account for this fact, we
hypothesize that additional somatosensory, photosensory, or mechanosensory systems may
compensate for the loss of lateral line sensation and help mutant larvae achieve neutral buoyancy
(Riley and Moorman, 2000; Suchocki and Sepulveda-Villet, 2019; Trotter et al., 2003). The lateral
line is involved several multi-sensory behaviors, including flow orientation (Bak-Coleman et al.,
2013; Coombs et al., 2020; Liao, 2006), prey capture (Carrillo and McHenry, 2016; Nelson et al.,
2002; New, 2002), and predator avoidance (Free et al., 2019; Jung et al., 2020; Stewart et al., 2013),
and we predict that surfacing can be added to this list.

The photosensory and mechanosensory systems exhibit distinct organizational patterns for
primary input to the central nervous system — even the anterior and posterior lateral line ganglia
primary projections are somatotopically organized in zebrafish larvae (Alexandre et al., 1999;
Fame et al., 2006; Hale et al., 2016; Liao and Haehnel, 2012; McCormick, 1989; Mirjany and
Faber, 2011). The Mauthner neurons integrate multisensory input at the level of the hindbrain, but
the surfacing behavior is unlikely to use this circuit. For higher order processing however,
information from the lateral line, vestibular, and photosensory systems are integrated in various
brain regions. The thalamus is one such sensory integration center in zebrafish (Mueller, 2012).
Socially-relevant mechanosensory and photosensory information is represented in a thalamic
region marked by the expression of *parathyroid hormone 2* (*pth2*) and *somatostatin 7* (*sst7*)
(Anneser et al., 2020; Kappel et al., 2022; Venuto et al., 2022). Vestibular information is also
represented in the thalamus (Favre-Bulle et al., 2018; Migault et al., 2018), though it is not clear if
the exact same region is involved. Photosensory, lateral line, and vestibular/auditory information
is also integrated in the tectum (Favre-Bulle et al., 2018; Filosa et al., 2016; Isa et al., 2021; Migault
et al., 2018). In fact, neurons in the periventricular layer (PVL) of the tectum, which is similar to
the superior colliculus in mammals, respond to visual, auditory, and lateral line inputs (Ingle, 1973;

King et al., 1996; Perrault et al., 2011; Thompson et al., 2016). However, it is hypothesized that most of these modalities are processed in parallel at younger stages of development (Thompson et al., 2016). It is worth noting that these sensory systems are in developing stages during surfacing behaviors.

Photosensory cues did not alter the buoyancy status of either wild type or *lhfpl5b* mutant larvae (Supplemental Figure 1), suggesting that, at most, photosensory systems play a subtle modulatory role in the surfacing behavior of larval zebrafish. The vestibular system is likely required to coordinate the vertical orientation and climbing portions of surfacing (Bagnall and Schoppik, 2018; Ehrlich and Schoppik, 2017; Ehrlich and Schoppik, 2019). Meanwhile, our data support the idea that the lateral line is responsible for the surface detection and breaching aspects of surfacing. However, given that both the vestibular and lateral line systems are providing spatial information in during the surfacing behavior, there may be crosstalk between them. For example, the vestibular system must quickly adapt to a dynamic increase in buoyancy over the course of multiple breaching events, so feedback from the lateral line during surface interactions may help fine tune the vestibular sense. However, the link between these systems have not been well-investigated during deliberate behaviors like surfacing (Chagnaud et al., 2017; Liu et al., 2021). So, while sensory integration exists in zebrafish and seems necessary for larvae to achieve neutral buoyancy, the integration dynamics are unclear during the intial swim bladder inflation timeframe and require future investigation.

Acknowledgements

The authors would like to thank Dr. Fadi Issa (ECU) for his support and technical advice. The authors also thank the Issa, Trapani, and Erickson lab members for fish care and maintenance.

Author contributions

Conceptualization: A.V., T.E.; Methodology: A.V., J.G.T., T.E.; Validation: A.V.; Formal analysis: A.V.; Investigation: A.V.; Resources: S.T-B., J.G.T; Writing - original draft: A.V., T.E.; Writing - review & editing: A.V., J.G.T., T.E..; Visualization: A.V.; Supervision: T.E.; Project administration: T.E.; Funding acquisition: J.G.T., T.E.

Funding

This work was supported by funding from East Carolina University's Division of Research, Economic Development and Engagement (REDE) and a Natural Science and Engineering

491 Research Council (NSERC) Discovery Grant (RGPIN-2021-03166) to T.E., a National Institute on

492 Deafness and Other Communication Disorders (NIDCD) award to J.G.T. (NIH R15DC014843),

493 and East Carolina Department of Biology funding provided to A.V..

References

Abbas, L. and Whitfield, T. T. (2009). Nkcc1 (Slc12a2) is required for the regulation of
endolymph volume in the otic vesicle and swim bladder volume in the zebrafish larva.
Development **136**, 2837–2848.
Alexander, R. M. (1966). Physical aspects of swimbladder function. *Biol Rev Camb Philos Soc*
41, 141–176.
Alexandre, D., Ghysen, A. and Ghysen, A. (1999). Somatotopy of the lateral line projection in
larval zebrafish. *Proc Natl Acad Sci U S A* **96**, 7558–62.
Anneser, L., Alcantara, I. C., Gemmer, A., Mirkes, K., Ryu, S. and Schuman, E. M. (2020).
The neuropeptide Pth2 dynamically senses others via mechanosensation. *Nature* **588**, 653–
657.
Bagnall, M. W. and Schoppik, D. (2018). Development of vestibular behaviors in zebrafish.
Curr Opin Neurobiol **53**, 83–89.
Bailey, H. C. and Doroshov, S. I. (1995). The duration of the interval associated with successful
inflation of the swimbladder in larval striped bass (Morone saxatilis). *Aquaculture* **131**, 135–
143.
Bak-Coleman, J., Court, A., Paley, D. A. and Coombs, S. (2013). The spatiotemporal
dynamics of rheotactic behavior depends on flow speed and available sensory information. *J*
Exp Biol **216**, 4011–24.
Baker, C. F. and Montgomery, J. C. (1999). The sensory basis of rheotaxis in the blind
Mexican cave fish, Astyanax fasciatus. *J Comp Physiol A* **184**, 519–527.
Bleckmann, H. (1988). Prey Identification and Prey Localization in Surface-feeding Fish and
Fishing Spiders. In *Sensory Biology of Aquatic Animals*, pp. 619–641. New York, NY:
Springer New York.
Bleckmann, H. (2006). The Lateral Line System of Fish. *Fish Physiology* **25**, 411–453.
Bleckmann, H. and Schwartz, E. (1982). The functional significance of frequency modulation
within a wave train for prey localization in the surface-feeding fishAplocheilus lineatus
(Cyprinodontidae). *Journal of Comparative Physiology A* **145**, 331–339.
Butler, J. M. and Maruska, K. P. (2016). Mechanosensory signaling as a potential mode of
communication during social interactions in fishes. *J Exp Biol* **219**, 2781–2789.
Carrillo, A. and McHenry, M. J. (2016). Zebrafish learn to forage in the dark. *Journal of*
Experimental Biology **219**, 582–589.
Chagnaud, B. P., Engelmann, J., Fritzsch, B., Glover, J. C. and Straka, H. (2017). Sensing
External and Self-Motion with Hair Cells: A Comparison of the Lateral Line and Vestibular
Systems from a Developmental and Evolutionary Perspective. *Brain Behav Evol* **90**, 98–
116.
Chatain, B. (1989). Problems related to the lack of functional swimbladder in intensive rearing
of Dicentrarchus labrax and Sparus auratus. *Advances in Tropical Aquaculture, Workshop at*
Tahiti, French Polynesia, 20 Feb - 4 Mar 1989.
Coffin, A. B. and Ramcharitar, J. (2016). Chemical Ototoxicity of the Fish Inner Ear and
Lateral Line BT - Fish Hearing and Bioacoustics: An Anthology in Honor of Arthur N.

Popper and Richard R. Fay. In (ed. Sisneros, J. A.), pp. 419–437. Cham: Springer
International Publishing.
Coombs, S., Bak-Coleman, J. and Montgomery, J. (2020). Rheotaxis revisited: a multi-
behavioral and multisensory perspective on how fish orient to flow. *Journal of Experimental*
Biology **223**,.
Doroshev, S., Cornacchia, J. and Hogan, K. (1981). Initial swim bladder inflation in the larvae
of physoclistous fishes and its importance for larval culture. *Rapports et Proces-verbaux des*
Réunions Cons. int. Explor. Mer **178**, 495–500.
Ehrlich, D. E. and Schoppik, D. (2017). Control of Movement Initiation Underlies the
Development of Balance. *Current Biology* **27**, 334–344.
Ehrlich, D. E. and Schoppik, D. (2019). A primal role for the vestibular sense in the
development of coordinated locomotion. *Elife* **8**,.
Erickson, T., Morgan, C. P., Olt, J., Hardy, K., Busch-Nentwich, E., Maeda, R., Clemens,
R., Krey, J. F., Nechiporuk, A., Barr-Gillespie, P. G., et al. (2017). Integration of Tmc1/2
into the mechanotransduction complex in zebrafish hair cells is regulated by
Transmembrane O-methyltransferase (Tomt). *Elife* **6**,.
Erickson, T., Pacentine, I. V., Venuto, A., Clemens, R. and Nicolson, T. (2020). The lhfpl5
Ohnologs lhfpl5a and lhfpl5b Are Required for Mechanotransduction in Distinct
Populations of Sensory Hair Cells in Zebrafish. *Front Mol Neurosci* **12**, 320.
Evans, H. M. and Damant, G. C. C. (1928). Observations on the Physiology of the Swim
Bladder in Cyprinoid Fishes. *Journal of Experimental Biology* **6**, 42–55.
Fame, R. M., Brajon, C. and Ghysen, A. (2006). Second-order projection from the posterior
lateral line in the early zebrafish brain. *Neural Dev* **1**, 4.
Favre-Bulle, I. A., Vanwalleghem, G., Taylor, M. A., Rubinsztein-Dunlop, H. and Scott, E.
K. (2018). Cellular-Resolution Imaging of Vestibular Processing across the Larval Zebrafish
Brain. *Current Biology* **28**, 3711-3722.e3.
Fero, K., Yokogawa, T. and Burgess, H. A. (2011). The behavioral repertoire of larval
zebrafish. *Neuromethods* **52**, 249–291.
Filosa, A., Barker, A. J., Dal Maschio, M. and Baier, H. (2016). Feeding State Modulates
Behavioral Choice and Processing of Prey Stimuli in the Zebrafish Tectum. *Neuron* **90**,
596–608.
Free, B. A., McHenry, M. J. and Paley, D. A. (2019). Probabilistic analytical modelling of
predator–prey interactions in fishes. *J R Soc Interface* **16**, 20180873.
Furukawa, S., Machida, Y., Takeuchi, K., Hoshikawa, Y. and Irie, K. (2022). Failure to gulp
surface air induces swim bladder adenomas in Japanese medaka (Oryzias latipes). *J Toxicol*
Pathol **35**, 237–246.
Gibbs, M. A. (2004). Lateral Line Receptors: Where Do They Come from Developmentally and
Where Is Our Research Going? *Brain Behav Evol* **64**, 163–181.
Goolish, E. M. and Okutake, K. (1999). Lack of gas bladder inflation by the larvae of zebrafish
in the absence of an air-water interface. *J Fish Biol* **55**, 1054–1063.
Hale, M. E., Katz, H. R., Peek, M. Y. and Fremont, R. T. (2016). Neural circuits that drive
startle behavior, with a focus on the Mauthner cells and spiral fiber neurons of fishes.
https://doi-org.proxy.hil.unb.ca/10.1080/01677063.2016.1182526 **30**, 89–100.
Hernández, P. P., Moreno, V., Olivari, F. A. and Allende, M. L. (2006). Sub-lethal
concentrations of waterborne copper are toxic to lateral line neuromasts in zebrafish (Danio
rerio). *Hear Res* **213**, 1–10.

Ingle, D. (1973). Evolutionary perspectives on the function of the optic tectum. *Brain Behav Evol* **8**, 211–237.

Isa, T., Marquez-Legorreta, E., Grillner, S. and Scott, E. K. (2021). The tectum/superior colliculus as the vertebrate solution for spatial sensory integration and action. *Current Biology* **31**, R741–R762.

Johansen, E. M. (1924). The Interfacial Tension between Petroleum Products and Water. *Ind Eng Chem* **16**, 132–135.

Jones, F. R. H. and Marshall, N. B. (1953). The structure and functions of the teleostean swimbladder. *Biological Reviews* **28**, 16–82.

Jung, J., Serrano-Rojas, S. J. and Warkentin, K. M. (2020). Multimodal mechanosensing enables treefrog embryos to escape egg-predators. *J Exp Biol* **223**,.

Kappel, J. M., Förster, D., Slangewal, K., Shainer, I., Svara, F., Donovan, J. C., Sherman, S., Januszewski, M., Baier, H. and Larsch, J. (2022). Visual recognition of social signals by a tectothalamic neural circuit. *Nature 2022 608:7921* **608**, 146–152.

Kimmel, C. B., Ballard, W. W., Kimmel, S. R., Ullmann, B. and Schilling, T. F. (1995). Stages of embryonic development of the zebrafish. *Developmental Dynamics* **203**, 253–310.

King, A. J., Schnupp, J. W. H., Carlile, S., Smith, A. L. and Thompson, I. D. (1996). The development of topographically-aligned maps of visual and auditory space in the superior colliculus. *Prog Brain Res* **112**, 335–350.

Kitajima, C., Watanabe, T., Tsukashima, Y. and Fujita, S. (1994). Lordotic Deformation and Abnormal Development of Swim Bladders in Some Hatchery-Bred Marine Physoclistous Fish in Japan. *J World Aquac Soc* **25**, 64–77.

Korsøen, Ø. J., Fosseidengen, J. E., Kristiansen, T. S., Oppedal, F., Bui, S. and Dempster, T. (2012). Atlantic salmon (Salmo salar L.) in a submerged sea-cage adapt rapidly to re-fill their swim bladders in an underwater air filled dome. *Aquac Eng* **51**, 1–6.

Kwan, K. M., Fujimoto, E., Grabher, C., Mangum, B. D., Hardy, M. E., Campbell, D. S., Parant, J. M., Yost, H. J., Kanki, J. P. and Chien, C. bin (2007). The Tol2kit: a multisite gateway-based construction kit for Tol2 transposon transgenesis constructs. *Dev Dyn* **236**, 3088–3099.

Liao, J. C. (2006). The role of the lateral line and vision on body kinematics and hydrodynamic preference of rainbow trout in turbulent flow. *J Exp Biol* **209**, 4077–90.

Liao, J. C. and Haehnel, M. (2012). Physiology of afferent neurons in larval zebrafish provides a functional framework for lateral line somatotopy. *J Neurophysiol* **107**, 2615–2623.

Lindsey, B. W., Smith, F. M. and Croll, R. P. (2010). From inflation to flotation: Contribution of the swimbladder to whole-body density and swimming depth during development of the zebrafish (Danio rerio). *Zebrafish* **7**, 85–96.

Lindsey, B. W., Dumbarton, T. C., Moorman, S. J., Smith, F. M. and Croll, R. P. (2011). Effects of simulated microgravity on the development of the swimbladder and buoyancy control in larval zebrafish (Danio rerio). *J Exp Zool A Ecol Genet Physiol* **315A**, 302–313.

Liu, Z., Hildebrand, D. G. C., Morgan, J. L., Slimmon, N. and Bagnall, M. W. (2021). The organization of the gravity-sensing system in zebrafish. *bioRxiv* 2021.07.09.451839.

Mackenzie, S. M. and Raible, D. W. (2012). Proliferative Regeneration of Zebrafish Lateral Line Hair Cells after Different Ototoxic Insults. *PLoS One* **7**, 47257.

McCormick, C. A. (1989). Central Lateral Line Mechanosensory Pathways in Bony Fish. *The Mechanosensory Lateral Line* 341–364.

McCune, A. R. and Carlson, R. L. (2004). Twenty ways to lose your bladder: Common natural
mutants in zebrafish and widespread convergence of swim bladder loss among teleost fishes.
Evol Dev **6**, 246–259.
Migault, G., Plas, T. L. van der, Trentesaux, H., Panier, T., Candelier, R., Proville, R.,
Englitz, B., Debrégeas, G. and Bormuth, V. (2018). Whole-Brain Calcium Imaging during
Physiological Vestibular Stimulation in Larval Zebrafish. *Current Biology* **28**, 3723-
3735.e6.
Mirjany, M. and Faber, D. S. (2011). Characteristics of the anterior lateral line nerve input to
the Mauthner cell. *Journal of Experimental Biology* **214**, 3368–3377.
Mo, W., Chen, F., Nechiporuk, A. and Nicolson, T. (2010). Quantification of vestibular-
induced eye movements in zebrafish larvae. *BMC Neurosci* **11**, 110.
Mogdans, J. (2019). Sensory ecology of the fish lateral-line system: Morphological and
physiological adaptations for the perception of hydrodynamic stimuli. *J Fish Biol* **95**, 53–72.
Montgomery, J. C., Baker, C. F. and Carton, A. G. (1997). The lateral line can mediate
rheotaxis in fish. *Nature* **389**, 960–963.
Mueller, T. (2012). What is the thalamus in zebrafish? *Front Neurosci* **6**, 64.
Müller, U. and Schwartz, E. (1982). Influence of single neuromasts on prey localizing behavior
of the surface feeding fish, Aplocheilus lineatus. *Journal of Comparative Physiology A* **149**,
399–408.
Murakami, S. L., Cunningham, L. L., Werner, L. A., Bauer, E., Pujol, R., Raible, D. W. and
Rubel, E. W. (2003). Developmental differences in susceptibility to neomycin-induced hair
cell death in the lateral line neuromasts of zebrafish (Danio rerio). *Hear Res* **186**, 47–56.
Nelson, M. E., MacIver, M. A. and Coombs, S. (2002). Modeling Electrosensory and
Mechanosensory Images during the Predatory Behavior of Weakly Electric Fish. *Brain*
Behav Evol **59**, 199–210.
New, J. G. (2002). Multimodal Integration in the Feeding Behaviors of Predatory Teleost Fishes.
Brain Behav Evol **59**, 177–189.
Nicolson, T., Rüsch, A., Friedrich, R. W., Granato, M., Ruppersberg, J. P. and Nüsslein-
Volhard, C. (1998). Genetic analysis of vertebrate sensory hair cell mechanosensation: The
zebrafish circler mutants. *Neuron* **20**, 271–283.
Olive, R., Wolf, S., Dubreuil, A., Bormuth, V., Debrégeas, G. and Candelier, R. (2016).
Rheotaxis of Larval Zebrafish: Behavioral Study of a Multi-Sensory Process. *Front Syst*
Neurosci **10**,.
Pavlov, D. S. and Tyuryukov, S. N. (1993). The role of lateral-line organs and equilibrium in
the behavior and orientation of the dace, Leuciscus leuciscus, in a turbulent flow. *J Ichthyol*
33, 45–55.
Perrault, T. J., Stein, B. E. and Rowland, B. A. (2011). Non-stationarity in multisensory
neurons in the superior colliculus. *Front Psychol* **2**,.
R Core Team (2022). R: A Language and Environment for Statistical Computing.
Richard, M. J. (1896). *Comptes rendus des séances de la Société de biologie et de ses filiales.*
Paris,Masson [etc.].
Rieger, P. W. and Summerfelt, R. C. (1998). Microvideography of gas bladder inflation in
larval walleye. *J Fish Biol* **53**, 93–99.
Riley, B. B. and Moorman, S. J. (2000). Development of utricular otoliths, but not saccular
otoliths, is necessary for vestibular function and survival in zebrafish. *J Neurobiol* **43**, 329–
337.

Santos, F., MacDonald, G., Rubel, E. W. and Raible, D. W. (2006). Lateral line hair cell
maturation is a determinant of aminoglycoside susceptibility in zebrafish (Danio rerio).
Hear Res **213**, 25–33.
Schoppik, D., Bianco, I. H., Prober, D. A., Douglass, A. D., Robson, D. N., Li, J. M. B.,
Greenwood, J. S. F., Soucy, E., Engert, F. and Schier, A. F. (2017). Gaze-stabilizing
central vestibular neurons project asymmetrically to extraocular motoneuron pools. *Journal*
of Neuroscience **37**, 11353–11365.
Schwartz, E. and Hasler, A. D. (1966). Perception of Surface Waves by the Blackstripe
Topminnow, *Fundulus notatus*. *Journal of the Fisheries Research Board of Canada* **23**,
1331–1352.
Stewart, W. J., Cardenas, G. S. and McHenry, M. J. (2013). Zebrafish larvae evade predators
by sensing water flow. *J Exp Biol* **216**, 388–98.
Stuart, K. R. and Drawbridge, M. (2012). The effect of photoperiod on larval culture
performance of two marine finfish species. *Aquaculture* **360–361**, 54–57.
Suchocki, C. R. and Sepulveda-Villet, O. J. (2019). The role of phototaxis in the initial swim
bladder inflation of larval yellow perch (Perca flavescens). *Int Aquat Res* **11**, 33–42.
Tait, J. S. (1960). The First Filling Of The Swim Bladder In Salmonoids. *Can J Zool* **38**, 179–
187.
Thompson, A. W., Vanwalleghem, G. C., Heap, L. A. and Scott, E. K. (2016). Functional
profiles of visual-, auditory-, and water flow-responsive neurons in the Zebrafish Tectum.
Current Biology **26**, 743–754.
Trotter, A. J., Battaglene, S. C. and Pankhurst, P. M. (2003). Effects of photoperiod and light
intensity on initial swim bladder inflation, growth and post-inflation viability in cultured
striped trumpeter (Latris lineata) larvae. *Aquaculture* **224**, 141–158.
Tsukamoto, Y. and Yoshino, S. (1957). A Study of the Lateral-line System in Fish II. On the
Relation between the Function of the Lateral-line Organ and the Habits of Fish. *Japanese*
Journal of Ichthyology **6**, 131–135.
v. Ledebur, J. F. and Wunder, W. (1937). Beiträge zur Physiologie der Schwimmblase der
Fische. *Zeitschrift für vergleichende Physiologie 1937 25:2* **25**, 149–155.
Venuto, A. and Erickson, T. (2021). Evaluating the Death and Recovery of Lateral Line Hair
Cells Following Repeated Neomycin Treatments. *Life 2021, Vol. 11, Page 1180* **11**, 1180.
Venuto, A., Smith, C. P., Cameron-Pack, M. and Erickson, T. (2022). Alone in a crowd:
effect of a nonfunctional lateral line on expression of the social hormone parathyroid
hormone 2. *Biol Open* **11**,.
Villamizar, N., García-Alcazar, A. and Sánchez-Vázquez, F. J. (2009). Effect of light
spectrum and photoperiod on the growth, development and survival of European sea bass
(Dicentrarchus labrax) larvae. *Aquaculture* **292**, 80–86.
von Ledebur, J. Frhr. (1928). Beiträge zur Physiologie der Schwimmblase der Fische. *Z Vgl*
Physiol **8**, 445–460.
Westerfield, M. (2000). *The zebrafish book. A guide for the laboratory use of zebrafish (Danio*
rerio). 4th ed. Eugene OR: University of Oregon.
713
714

Figures

Figure 1. Hyperinflation of the swim bladder in lateral line mutants (*lhfpl5b*[-/-]). A) Representative
examples of over-, regular, and under-inflated swim bladders (arrows) in larval zebrafish at 6 dpf
(top: *lhfpl5b*[-/-], others are wild type). B) Correlation between buoyancy and swim bladder volume.
Line of best fit applied through the combination of mutant and wild type data (linear model, R-
squared = 0.8574). C) Percentage of larvae that exhibit over-, regular, and under-inflation of the
swim bladder, as defined by positive, neutral, and negative buoyancy. A Chi-Squared Test was
used to determine significance. D) Swim bladder volume (mm^3) in wild type and *lhfpl5b* mutant
larvae at 6 dpf. A Welch's t-test was used to determine significance. *** = p < 0.001, **** = p <
0.0001. Scale bar = 0.2 mm.
727

728
Figure 2: Blocking access to the surface eliminates the hyperinflation phenotype in lateral line
mutants (*lhfpl5b*$^{-/-}$). A) Diagram of experimental set up with a filter blocking access to the surface.
B) Percentage of larvae that exhibit over-, regular, and under-inflation of the swim bladder at 6
dpf. A Chi-Squared Test was used to determine significance. C) Swim bladder volume (mm^3) of
larvae at 6 dpf. A One-Way ANOVA was used to determine significance. **** = p < 0.0001, ,
**** = p < 0.0001, ns = not significant. Full ANOVA table in supplemental tables.
735

Figure 3: Rescue of lateral line mutants (*lhfpl5b*^{-/-}) with the *Tg(myo6b:EGFP-lhfpl5a)vo23* transgene. A) Percentage of larvae that exhibit swim bladder under, regular, and over-inflation at 6 dpf. A Chi-Squared Test was used to determine significance. B) Swim bladder volume (mm^3) of larvae at 6 dpf. A One-Way ANOVA was used to determine significance. **** = $p < 0.0001$, ns = not significant. Full ANOVA table in supplemental tables.

743
Figure 4: Head-specific ablations of the lateral line recapitulates the *lhfpl5b* mutant hyperinflation
phenotype. A) Diagram of experiment. B) Percentage of larvae that exhibit swim bladder under,
regular, and over-inflation at 5 dpf. A Chi-Squared Test was used to determine significance. C)
Swim bladder volume (mm^3) of larvae at 5 dpf. A One-Way ANOVA was used to determine
significance. * = $p < 0.05$, , **** = $p < 0.0001$, ns = not significant. Full ANOVA table in
supplemental tables. Images of live larvae following treatment in Supplemental Figure 3.
750

751
752 **Figure 5**: Decreasing interfacial surface tension with mineral oil results in over-filling of the swim
753 bladder. A) Diagram of experiment with 2 mm layer of oil on surface. B) Images of swim bladder
754 phenotypes resulting from surface oil and surface air exposure in wild type larvae at 6 dpf. C) Swim
755 bladder volume (mm^3) at 6 dpf. A One-Way ANOVA was used to determine significance. * = p <
756 0.05, ns = not significant. Scale bar = 0.1 mm. Full ANOVA table in supplemental tables.
757

758
Figure 6: Optogenetic activation of lateral line hair cells with Channelrhodopsin-2 (ChR2) in wild
type and *lhfpl5b* mutant larval zebrafish. A) Diagram of experiment with blue light stimuli
delivered for 25 milliseconds at 1 second intervals. Treatments were started at 4 dpf and swim
bladder volume was assessed at 6 dpf. B) Swim bladder volume (mm^3) of larvae at 6 dpf, n = 16-
21 per condition. One-Way ANOVA was used to determine significance. ** = $p < 0.001$, ns = not
significant. Full ANOVA table in supplemental tables.
765

766
Figure 7: Quantification of the surfacing behaviors of wild type and *lhfpl5b* mutant zebrafish larvae
during initial swim bladder inflation. A) Total number of individual visits taken by larval fish to
the air-water interface on 4 dpf for 12 hours (9 AM – 9 PM). B) Total amount of time larval fish
spend at the surface for all visits combined on 4 dpf. C) Time to first surface visit after access was
allowed. A One-Way ANOVA was used to determine significance. *** = p < 0.001, ns = not
significant.
773

774
Figure 8. Summary of the surfacing behavior of wild type and lateral line-deficient zebrafish
larvae. A) Wild type larvae use their lateral line hair cells (in green) to mediate interactions with
the air-water interface during surfacing. As a result of accurate surface detection, larvae take in an
appropriate volume of air for swim bladder (SB) inflation and achieve neutral buoyancy. B) Larvae
with a genetic loss of lateral line function (hair cells in white) misinterpret the air-water interface,
leading to increased interactions with the surface. Consequently, lateral line mutants *(lhfpl5b$^{-/-}$)*
take in an excess volume of air, resulting in hyperinflation of the swim bladder for approximately
half of mutant larvae. Selective chemical ablations of the head neuromasts produce similar results,
implicating the anterior lateral line as the primary sensory organ for interactions with the air-water
interface during the surfacing behavior.
785

786 **Supplemental Figures**

787
Supplemental Figure 1. Buoyancy status of larvae raised in total darkness. Percentage of larvae
that exhibit swim bladder under, regular, and over-inflation raised in either a 14:10 light/dark
cycle (Control) or total darkness (0:24 light/dark cycle). A Chi-Squared Test was used to assess
significance. Both Wild Type Control and Wild Type Dark over-inflation proportions: $M = 0.0\%$,
$SD = 0.0\%$, $n = 32$. *lhfpl5b*$^{-/-}$ Control over-inflation proportion: $M = 57.5\%$, $SD = 6.6\%$, $n = 30$.
lhfpl5b$^{-/-}$ Dark over-inflation proportion: $M = 52.4\%$, $SD = 4.1\%$, $n = 30$.
794

Supplemental Figure 2. Chemical ablations of lateral line hair cells phenocopies the hyperinflation phenotype of *lhfpl5b* mutants. A) Percentage of larvae treated with either neomycin or CuSO4 that exhibit swim bladder under, regular, and over-inflation. A Chi-Squared Test was used to assess significance. B) Swim bladder volume (mm^3) in 5 dpf larvae treated with either neomycin or CuSO4. A One-Way ANOVA was used to determine significance. **** = $p < 0.0001$, *** = $p < 0.001$, ** = $p < 0.01$, * = $p < 0.05$, ns = not significant. A full ANOVA table can be found in the supplemental tables.

805 **Supplemental Figure 3.** Head and trunk-specific lateral line ablations with CuSO4. Representative
806 images of live *Tg(myo6b:EGFP-PA)vo68* transgenics at 4 dpf, two hours after CuSO4 treatment.
807 This transgene labels hair cells of the lateral line and inner ear (indicated by white brackets on the
808 Control image). CuSO4 treatment ablates lateral line hair cells only, leaving inner ear hair cells
809 intact.
810

811
Supplemental Figure 4: A) Representative images of Channelrhodopsin-2 (ChR2) expression in
neuromast, utricle, and crista hair cells of *Tg(lhfpl5b2028:ChR2-EYFP-PA)* transgenics at 2 dpf.
ChR2-YFP expression is detectable only in the lateral line neuromasts, indicating specificity of the
lhfpl5b regulatory region used to drive ChR2 expression. First panel is YFP only (false-colored
green) and the second panel is a merge of YFP and DIC. Scale bars = 10 uM. B) Swim bladder
volume (mm^3) of 6 dpf wild type and *lhfpl5b$^{-/-}$* larvae, with and without the *ChR2* transgene. Groups
of larvae were also treated with CuSO4 to ablate lateral line hair cells. All larvae were stimulated
with blue light. The impacts of blue light stimulation on wild type ChR2 transgenics is abolished
by chemical ablation of lateral line hair cells. As expected, *lhfpl5b* mutants are unaffected by
CuSO4 treatment. A One-Way ANOVA was used to determine significance. ** = $p < 0.00724$, ns
= not significant. A full ANOVA table can be found in the supplemental tables.
823