Real-time feedback reduces participant motion during task-based fMRI

Chad S. Rogers[1], Michael S. Jones[2], Sarah McConkey[2],
Drew J. McLaughlin[3], Jonathan E. Peelle[4, 5, 6]

[1] Department of Psychology, Union College

[2] Department of Otolaryngology, Washington University in St. Louis

[3] Basque Center on Cognition, Brain and Language

[4] Center for Cognitive and Brain Health, Northeastern University

[5] Department of Communication Sciences and Disorders, Northeastern University

[6] Department of Psychology, Northeastern University

Running title: Real-time feedback reduces fMRI motion

Data availability statement: The data that support the findings of this study are openly available in OpenNeuro at https://doi.org/10.18112/openneuro.ds004285.v1.0.0, reference number ds004285.

IRB statement: The study reported here was conducted according to a protocol approved by the Washington University in Saint Louis Institutional Review Board.

Please address post-publication correspondence to:

Dr. Chad Rogers   Dr. Jonathan Peelle
Department of Psychology   Center for Cognitive and Brain Health
Union College   Northeastern University
rogersc@union.edu   j.peelle@northeastern.edu

Abstract

The potential negative impact of head movement during fMRI has long been appreciated.
Although a variety of prospective and retrospective approaches have been developed to help
mitigate these effects, reducing head movement in the first place remains the most appealing
strategy for optimizing data quality. Real-time interventions, in which participants are provided
feedback regarding their scan-to-scan motion, have recently shown promise in reducing motion
during resting state fMRI. However, whether feedback might similarly reduce motion during
task-based fMRI is an open question. In particular, it is unclear whether participants can
effectively monitor motion feedback while attending to task-related demands. Here we assessed
whether a combination of real-time and between-run feedback could reduce head motion during
task-based fMRI. During an auditory word repetition task, 78 adult participants (aged 18–81)
were pseudorandomly assigned to receive feedback or not. We quantified movement using
framewise displacement (FD). We found that motion feedback resulted in a statistically
significant reduction in participant head motion, with a small-to-moderate effect size (reducing
average FD from 0.347 to 0.282). There was some evidence for a change of effect over the
course of six runs, but no clear evidence for practice effects. We conclude that under some
circumstances real-time feedback may reduce head motion during task-based fMRI, although its
effectiveness may depend on the specific participant population and task demands of a given
study.
68

Introduction

Head movement during fMRI can introduce a number of different challenges to data analysis.
Although long appreciated (Friston et al., 1996), the detrimental influence of head motion on
resting state functional connectivity has only recently been widely recognized (Power et al.,
2015). Rigid body realignment—a mainstay of fMRI analysis for decades—goes some way
towards improving correspondence across images (Ashburner and Friston, 2004), but does not
remove extraneous signal components introduced by movement (Friston et al., 1996). A
common approach for mitigating motion-related artifacts is to include the 6 realignment
parameters (translation and rotation around the X, Y, and Z axes) as nuisance regressors in
first-level models. Additional approaches include wavelet despiking (Patel et al., 2014), ICA
(Pruim et al., 2015), robust-weighted least squares (Diedrichsen and Shadmehr, 2005),
Bayesian approaches (Eklund et al., 2017), and frame censoring (Lemieux et al., 2007).
However, recent work investigating motion correction strategies in multiple data sets suggests
that the optimal strategy may depend on the specific data set and output metric (Jones et al.,
2022), precluding a simple one-size-fits-all solution.
Given the lack of certainty regarding how to best handle head motion during analysis,
reducing head motion in the first place is a particularly appealing option. Common approaches
to reducing head motion during MRI scanning include foam padding and instructing participants
to not move during the scan. Other existing strategies include using custom head molds (Power
et al., 2019) and playing participants a relaxing abstract movie (Vanderwal et al., 2015).
A complementary approach for reducing head motion is to provide participants with
feedback regarding their head movement so they can learn to minimize motion. FIRMM
software[1] (Dosenbach et al., 2017) is one implementation of motion feedback. FIRMM uses
rapid image reconstruction and rigid-body alignment to estimate frame-by-frame movement,
providing visual cues to participants based on estimated movement. Feedback can either be
provided in real time, or between scanning runs. FIRMM has been shown to reduce movement
during resting state scans in both adults (Dosenbach et al., 2017) and in young children
(Greene et al., 2018; Badke D'Andrea et al., 2022).
Despite its success in resting state fMRI, it is unclear whether motion feedback would
lead to similar improvements during a typical task-based fMRI paradigm. On the one hand, if
participants are able to modulate their motion in one experimental paradigm (namely, resting
state studies) it stands to reason they could do so in a different experimental context (a task-
based study). On the other hand, task-based fMRI studies require participants to attend to the
experimental task at hand, which will reduce the cognitive resources available for noting and
processing the feedback, as well as the ability to attend to proprioceptive cues related to self-
motion. In addition, if experimental stimuli and motion feedback are in the same modality,
subjects may also have difficulty accurately processing motion-related feedback.
To evaluate the degree to which feedback might affect head motion during task-based
fMRI, we assigned participants in a task-based fMRI study of spoken word recognition to
receive feedback (the feedback group) or not (the no-feedback group). All other instructions,
scanning parameters, and task requirements were identical. We included both young and older
adults, and a task (word repetition) that we anticipated would typically require head movement.
Finally, we were able to capitalize on the fact that each subject has many observations
(hundreds of frames) by using linear mixed effects analysis.

---

[1] Now a commercial product offered by Turing Medical (St. Louis, MO).

Method

MRI data are available from OpenNeuro (Markiewicz et al., 2021) as ds004285 (Rogers et al.,
2023), and motion data and analysis scripts are available from https://github.com/jpeelle/motion-
feedback. (Although we collected these data in the context of a task-based fMRI study,
described below, here we focus only on effects of feedback on head motion and not the task
results.)

Participants
We tested 78 participants aged 19–81. All reported themselves to have normal hearing, to be
right handed, and not have a history of neurological disease. There were no exclusions based
on sex, gender, race, or ethnicity. Participants were limited to adults because of our scientific
focus on adult language processing. Because the task involved spoken language, all were
native speakers of American English.
Participants were randomly assigned to control (no motion feedback) or intervention
(real-time motion feedback) groups. Demographic characteristics for these groups are shown in
Table 1. Because data collection was ended early due to COVID-19 safety requirements the
groups did not have equal numbers of participants. All provided informed consent under a
process approved by the Washington University in Saint Louis Institutional Review Board.
130
131

Table 1. Participant characteristics

		No feedback	Feedback
Young adults	N	17	31
	Age (mean ± SD)	23.3 (3.06)	22.4 (2.64)
	Sex	13 F, 4 M	28 F, 3 M
Older adults	N	5	25
	Age (mean ± SD)	70.4 (5.22)	71.3 (3.81)
	Sex	5 F, 0 M	17 F, 8 M

133 **Procedure**
134 The procedure is modeled after that in Rogers et al. (2020) and illustrated in **Figure 1**.
135 Participants performed a word repetition task in which on every trial they heard a spoken word,
136 presented in stationary background noise (3 dB SNR), and repeated it back aloud.
137 The no-feedback group was given the following instructions:
138
139 During this task, it is important that you hold your body and head very still.
140 Please stay relaxed, stay alert, and keep your eyes open and on the fixation
141 cross.
142

Figure 1. Summary of task design and processing pipeline. **a)** Schematic of the word repetition task. Participants heard a word presented in the midst of background noise during the gap between volume acquisitions; during the following gap, they repeated the word aloud. **b)** Illustration of the two groups to which a participant could be pseudorandomly assigned. The group without feedback received standard instructions to keep still at the beginning of the study; the feedback group received instructions regarding real-time feedback, a colored crosshair indicating their scan-to-scan motion (framewise displacement; FD) during each run; and a visual report on feedback at the end of each run. **c)** Illustration of time variables available for modeling.

144
145   FIRMM was run in the background for experimenters to monitor motion during scanning
146   sequences and participants were given verbal feedback after each sequence.
147   The group receiving motion feedback was given the following instructions:
148
149   It is very important to remain still during your MRI so that we can obtain clear
150   images. Even very small movements that you are not aware of can affect the
151   image quality. While performing the task, you will be receiving feedback
152   corresponding to your ability to remain still. This is to help you be aware of any
153   movements you may be making. You will see a white fixation cross on the
154   screen. The cross will change to yellow and then red depending on how much
155   you are moving. It will go back to white if you become still again. Sometimes
156   even if you are doing your best you will still see a red cross. This may mean that

the computer is too strict and not that you are necessarily doing anything wrong.

Just keep trying your best to keep the cross in the white.

During the task, participants viewed a colored cross in the middle of the screen. FD

thresholds were set to show a white cross at < 0.2 mm, a yellow cross at 0.2 mm to < 0.3 mm,

and a red cross at ≥ 0.3 mm. At the conclusion of each run, participants viewed a Head Motion

Report (**Figure 2**) which displayed their performance on a gauge of 0–100 (a percentage score

of 0% to 100%), and a graph of their motion level over time to help visualize total movement

during the session. Participants were encouraged to bring their score closer to 100% on

subsequent sequences.

Figure 2. Head motion report provided to participants following each run of the experiment.

MRI acquisition and analysis

MRI data were acquired using a Siemens Prisma scanner (Siemens Medical Systems) at 3 T

equipped with a 32-channel head coil. Scan sequences began with a T1-weighted structural

volume using an MPRAGE sequence [repetition time (TR) = 2.4 s, echo time (TE) = 2.2 ms, flip

angle = 8°, 300 × 320 matrix, voxel size = 0.8 mm isotropic]. Blood oxygenation level-dependent

(BOLD) functional MRI images were acquired using a multiband echo planar imaging sequence
(Feinberg et al., 2010) [TR = 3.07 s, TA = 0.770 s, TE = 37 ms, flip angle = 90º, voxel size = 2
mm isotropic, multiband factor = 8]. We used a sparse imaging design in which there was a 2.3
second delay between scanning acquisitions and the TR was longer than the acquisition time to
allow for minimal scanning noise during stimulus presentation and audio recording of participant
responses (Edmister et al., 1999; Hall et al., 1999). Due to slight variations in scanning
sessions, participants had 763–846 frames of data (median = 794).
Analysis of the MRI data was performed using Automatic Analysis version 5.8.1 (Cusack
et al., 2015) (RRID:SCR_003560) which scripted analyses run in SPM12 (Wellcome Trust
Centre for Human Neuroimaging) version 7487 (RRID:SCR_007037). Functional images
underwent rigid body realignment (i.e, "motion correction"). We used framewise displacement
(FD) to parsimoniously summarize frame-by-frame motion, calculated as the sum of the six
head motion estimates obtained from realignment, with a dimensional conversion of the three
rotations assuming the head is a 50 mm sphere (Power et al., 2012). We supplemented FD
values with differential variance (DVARS), a measure of variations in image intensity. DVARS
was calculated as the root-mean-squared of the time difference in the BOLD signal calculated
across the entire brain, before realignment (Smyser et al., 2011). Although FD and DVARS are
highly correlated, they are not identical (Jones et al., 2022), and we anticipated that these two
metrics might differ in sensitivity to motion feedback.

194   **Results**

195   Movement parameters for young and older adults as a function of feedback is shown in **Figure
196   3**. **Figure 3a** shows FD values for two subjects (our first young adult without feedback, and our
197   first young adult with feedback), as well as the mean FD for all subjects in each feedback group.
198   Although we used all time points in the statistical analysis, for illustrative purposes we also
199   plotted summary values in **Figure 3b** (each point denoting mean FD per subject).

223
224

Figure 3. Comparison of average framewise displacement (FD) values for feedback and no-feedback groups. **a)** *Top:* FD values for two example subjects, one with feedback and one without. *Bottom:* Mean FD values for all participants in the feedback or no-feedback group. **b)** Summary of mean FD values as a function of age and feedback. Individual subjects are dots, mean ± SE shown in bars.

225
226
227   We analyzed these data using R version 4.2 (R Core Team, 2020) (RRID:SCR_001905).
228   We conducted linear mixed effects analysis using the *nlme* package (Pinheiro and Bates, 2000).
229   We accounted for temporal autocorrelation (that is, motion at one frame is related to motion
230   during the following frame) using a first-order autoregressive model. The model specification
231   was:

232
```
233    m2 <- lme(FD ~ 1 + frame + age_group * feedback,
234                    random = ~ frame | subject_number,
235                    data = df,
236                    method = "ML",
237                    correlation = corAR1(value =
238    autocorrelation_estimate, form = ~ frame | subject_number, fixed
239    = FALSE))
```
240
241
242

243
244 where `df` is the data frame, `FD` is framewise displacement, and `feedback` is a dichotomous
245 variable denoting whether feedback was provided or not. We calculated the
246 `autocorrelation_estimate` from the data (value of 0.52). Model results are shown in
247 **Table 2**.
248
249

Table 2. Fixed effects results for motion model by frame

	Est	SE	df	t	p	
Intercept	0.4504154	0.3766628	61778	11.958053	< .0001	***
frame	0.0000442	0.00001716	61778	2.572848	0.0101	*
age group	-0.1659770	0.04270876	74	-3.886252	0.0002	***
feedback	-0.1193043	0.04112688	74	-2.900883	0.0049	**
age group x feedback	0.0507396	0.04830427	74	1.050417	0.2969	

*** p < .001; ** p < .01; * p < .05

250
251
252
253   Although we first included time (frame) in the model, in fMRI research it is often
254 convenient to think about continuous scanning runs (sometimes also referred to as "sessions").
255 In the current study, each participant had 6 runs of data. We thus ran an additional model to see
256 whether the effect of feedback was comparable across run. Because time was expressed as 6
257 runs, we did not account for temporal autocorrelation in this model:
258
259 ```
m3 <- lme(FD ~ 1 + run + age_group * feedback,
260   random = ~ run | subject_number,
261   data = df,
262   method = "ML",
263   correlation = NULL)
```
264
265
266        Results of this analysis by run are shown in **Table 3**, and FD as a function of run shown
267 in **Figure 4**. We again found a significant effect of feedback, but the effect of feedback did not
268 significantly differ by run (as one might expect if there were learning effects).

**Figure 4.** Mean FD values for all participants in the feedback or no-feedback group as a function of run. Individual subjects are dots, mean ± SE shown in bars.

276
277

**Table 3.** Fixed effects results for motion model by run

|  | Est | SE | df | t | p | |
|---|---|---|---|---|---|---|
| Intercept | 0.4524204 | 0.03491125 | 61774 | 12.959158 | < .0001 | *** |
| run2 | 0.0198209 | 0.01118544 | 61774 | 1.772030 | 0.0764 | |
| run3 | 0.0294646 | 0.00999432 | 61774 | 2.948129 | 0.0032 | ** |
| run4 | 0.0358940 | 0.01053085 | 61774 | 3.408465 | 0.0007 | *** |
| run5 | 0.0352148 | 0.01335915 | 61774 | 2.636004 | 0.0084 | ** |
| run6 | 0.0278481 | 0.01217601 | 61774 | 2.287125 | 0.0222 | * |
| age group | -0.1769315 | 0.03945612 | 74 | -4.484260 | < .0001 | *** |
| feedback | -0.1344319 | 0.03799784 | 74 | -3.537883 | 0.0007 | *** |
| age_group x feedback | 0.0724903 | 0.04462674 | 74 | 1.624370 | 0.1085 | |

*** p < .001; ** p < .01; * p < .05

279        Although our analysis was focused on changes in mean FD, we wanted to look in more
280    detail at how feedback affected the distribution of movement. We thus plotted the density of FD
281    values for all scans as a function of feedback (**Figure 5**) for FD values up to 2.0. We used
282    density rather than the count to control for the different numbers of frames in the two groups
283    (resulting from different numbers of subjects). Although qualitative, this analysis highlights that
284    participants receiving feedback showed fewer high-FD scans than those receiving lower
285    feedback, with this effect becoming more apparent at higher FD values—for FD values above
286    1.5, the feedback group had less than a third of these scans than the group without feedback.
287
288

**Figure 5. a)** Density plot of all FD values as a function of feedback. **b)** Density of FD values in the feedback condition relative to the no feedback condition. A value of 1 (dashed line) indicates equal density of FD values in the two conditions. The plot illustrates a decrease in high-FD values in the feedback condition, and an increase in low FD values, relative to the no feedback condition.

289
290        We next repeated our primary FD analyses, but using DVARS (rather than FD) as a
291    dependent measure, under the logic that FD and DVARS may be differently sensitive to effects
292    of motion, and one or the other may be of more interest in any specific study. Summary results
293    are shown in **Figure 6** and **Table 4**. Although DVARS was numerically lower for the feedback
294    group than the no-feedback group, the difference was not statistically significant. Results as a
295    function of run is shown in **Figure 7** and **Table 5**. Again, though we saw numerical reductions in
296    DVARS for the feedback group, the difference was not significant.
297
298
299

**Figure 6.** Comparison of average DVARS values for feedback and no-feedback groups. **a)** *Top:* DVARS values for two example subjects, one with feedback and one without. *Bottom:* Mean DVARS values for all participants in the feedback or no-feedback group. **b)** Summary of mean DVARS values as a function of age and feedback. Individual subjects are dots, mean ± SE shown in bars.

300
301
302

**Table 4.** Fixed effects results for DVARS model by frame

|  | Est | SE | df | t | p |  |
| --- | --- | --- | --- | --- | --- | --- |
| Intercept | 244.41952 | 13.012674 | 61778 | 18.783188 | <.0001 | *** |
| frame | 0.00611 | 0.004908 | 61778 | 1.244030 | 0.2135 | |
| age_group | -31.76489 | 14.664553 | 74 | -2.166100 | 0.0335 | * |
| feedback | -24.40635 | 14.121338 | 74 | -1.728331 | 0.0881 | |
| age_group x feedback | 2.21691 | 16.585800 | 74 | 0.133663 | 0.8940 | |

*** p < .001; ** p < .01; * p < .05

**Figure 7.** Mean DVARS values for all participants in the feedback or no-feedback group as a function of run. Individual subjects are dots, mean ± SE shown in bars.

304
305

**Table 5.** Fixed effects results for DVARS model by run

|  | Est | SE | df | t | p |  |
|---|---|---|---|---|---|---|
| Intercept | 243.49427 | 12.192275 | 61774 | 19.971192 | < .0001 | *** |
| run 2 | 3.08907 | 2.623021 | 61774 | 1.177675 | 0.2389 |  |
| run 3 | 5.56690 | 2.696872 | 61774 | 2.064205 | 0.0390 | * |
| run 4 | 7.33254 | 2.863635 | 61774 | 2.560571 | 0.0105 | * |
| run 5 | 5.65629 | 3.768866 | 61774 | 1.500794 | 0.1334 |  |
| run 6 | 2.94432 | 3.157047 | 61774 | 0.932620 | 0.3510 |  |
| age group | -32.41583 | 13.721953 | 74 | -2.362334 | 0.0208 | * |
| feedback | -27.06506 | 13.213980 | 74 | -2.048214 | 0.0441 | * |
| age group x feedback | 6.10355 | 15.519814 | 74 | 0.393274 | 0.6952 |  |

*** p < .001; ** p < .01; * p < .05

306
307

308         In addition to framewise measurements of FD and DVARS, we investigated whether
309   motion feedback affected the temporal signal-to-noise ratio (tSNR) of the data. We first created
310   a brain mask using each subject's tissue class segmentation, binarizing at a gray matter
311   probability of 0.8. We then calculated tSNR (the mean of the signal divided by its standard
312   deviation, over frames) in every voxel within the brain mask. We then performed two
313   complementary analyses. First, for each run, we took the mean tSNR over voxels, and used the
314   same statistical model as for FD and DVARS, shown in **Figure 8** and shown in **Table 6**.
315   Although the tSNR values were numerically higher in the feedback group, there was no
316   significant effect of feedback ($p = 0.0814$).
317
318
319

**Figure 8.** Comparison of average temporal SNR (tSNR) values for feedback and no-feedback groups. Each participant is represented by a single point per run, averaging the tSNR over all voxels in the gray matter mask.

**Table 6.** Fixed effects results for tSNR model

|  | Est | SE | df | t | p |  |
|---|---|---|---|---|---|---|
| Intercept | 35.62189 | 2.3494343 | 389 | 15.161899 | < .0001 | *** |
| run | -0.01283 | 0.1116692 | 389 | -0.114888 | 0.9086 |  |
| age group | 5.61719 | 2.6542796 | 74 | 2.116276 | 0.0377 | * |
| feedback | 4.51521 | 2.5559386 | 74 | 1.766556 | 0.0814 |  |
| age group x feedback | -0.11883 | 3.0020112 | 74 | -0.039584 | 0.9685 |  |

*** p < .001; ** p < .01; * p < .05

327
328
329       Finally, we assessed the degree to which feedback may have affected performance on
330  the behavioral task. Accuracy on the word repetition task is shown in **Figure 9** as a function of
331  age group and feedback. We used a linear model to test for the impact of feedback on
332  behavioral accuracy (averaging over experimental runs). We found no significant effect of age
333  group, no significant effect of feedback, and no significant age x feedback interaction, although
334  in older adults there was a small numerical decrease in accuracy during the feedback condition.
335

**Figure 9.** Behavioral accuracy for young and older adults as a function of feedback.

337        **Discussion**

338    Given the pernicious and sometimes unpredictable effects of head movement in fMRI studies,
339    reducing motion in the first place is on its face an appealing strategy. In the current study we
340    evaluated the degree to which a combination of real-time and summary motion feedback could
341    reduce head movement in adult participants by pseudorandomly assigning participants to
342    feedback or no feedback groups. We found that motion feedback was associated with a
343    statistically significant reduction in average FD of approximately 0.06, with consistent (though
344    not significant) changes in DVARS and tSNR. We discuss these findings and their implications
345    below.
346        First, our primary finding that providing motion feedback can significantly reduce head
347    motion in participants undergoing task-based MRI is notable. These findings add motion
348    feedback to the limited set of tools MRI researchers have for reducing head motion during
349    acquisition. Although numerous strategies exist for accounting for head motion in preprocessing
350    and analysis, none are perfect, and it is difficult to ascertain the most appropriate approach
351    (Ardekani et al., 2001; Oakes et al., 2005; Johnstone et al., 2006; Jones et al., 2022). Reducing
352    motion during acquisition may therefore be a preferable alternative in many circumstances.
353        It is important to highlight the magnitude of the effect we observed. Compared to the
354    participants with no motion feedback, participants with feedback showed a reduction in FD of
355    approximately .06. Although this may be helpful in reducing motion-related artifacts in data
356    processing, it is not a large effect (though in theory it might be increased if different types of
357    feedback were used, or in the context of a different task). It is also important to remember that
358    we tested healthy adults aged 19–81. It could be that the effect of motion feedback would be
359    larger (or smaller) in different populations.
360        That being said, the principal effect of feedback may not be the *average* reduction in FD
361    observed, but rather a reduction in the number of large motion events. Rapid head motion tends
362    to have a larger effect on the BOLD signal than slower motion. Thus, reducing the number of
363    large, quick movements may improve signal quality to a greater extent than a simple reduction
364    in mean FD. Indeed, we saw the biggest proportional difference between feedback groups at
365    higher FD values. Interestingly, the reduction in high motion events (**Figure 5b**) roughly
366    corresponds to an FD of 0.3, the value at which participants received "red" feedback.
367        An important consideration when implementing real-time feedback during fMRI is
368    whether doing so significantly changes task demands. Although we found accuracy was
369    statistically comparable across feedback groups, we cannot rule out other changes in task
370    demands. Indeed, we observed a numeric (though not significant) decrease in task accuracy in
371    older adults when they were presented with real-time feedback relative to the no-feedback
372    group. Broadly speaking, we might expect monitoring the display for feedback, and adjusting
373    ones movement in response, to engage systems related to cognitive control (sometimes also
374    referred to as attentional control or executive function). In one framing of cognitive control,
375    cognitive control is proposed to operate in at least two complementary modes: a *proactive* mode
376    that is concerned with maintaining goal-directed behavior, and a *reactive* mode that is engaged
377    when goal-directed performance falters (Braver, 2012). There are two implications of such a
378    framework. First, participants who are less able to engage cognitive control may benefit less
379    from motion feedback compared to those who are better able to engage control systems.
380    Second, researcher concerns about how task demands may affect imaging results of interest
381    may depend on the nature of the main fMRI task. For example, tasks tapping cognitive control
382    or executive function may be more impacted by motion feedback than those focused on primary
383    sensory or motor systems. In our case, we used a speech-in-noise task, one that is
384    hypothesized to engage cognitive control networks (Peelle, 2018; McLaughlin et al., 2021).
385        In our implementation, participants received brief instructions before entering the
386    scanner, and once in the scanner the display was explained to them. One interesting future

387  direction would be to provide participants with motion training over a longer period to see
388  whether they might learn to better limit their motion in the absence of feedback (for example, by
389  training in a mock scanner). Such an approach would take more time, but would also circumvent
390  some of the challenges associated with real-time feedback (such as introducing a "dual-task"
391  situation).
392      Although we found a significant effect of motion feedback on FD, we did not observe a
393  significant effect on motion feedback on DVARS (although DVARS values were numerically
394  lower in the feedback group). FD and DVARS are typically strongly correlated (Jones et al.,
395  2022), but not identical, and thus the divergence in findings is not entirely unexpected. The
396  numerical (albeit not significant) reductions in DVARS in participants receiving motion feedback
397  is consistent with the effect we saw for FD.
398      In addition, we found that feedback led to increased whole-brain tSNR, which we used
399  as a proxy of overall image quality. Despite the tSNR being numerically larger in the group
400  receiving motion feedback, the difference was not statistically significant. However, it is still
401  possible that small improvements in tSNR would result in appreciable improvements in the
402  accuracy or reliability of statistical models.
403      In conclusion, we found that motion feedback as implemented in FIRMM slightly but
404  significantly reduced the amount of head motion we observed during task-based fMRI. Real-
405  time feedback may thus be well-suited to complement other approaches to motion reduction
406  depending on the specific needs of a given study.
407

408          **Acknowledgments**

409     Work reported here was supported by grant R01 DC014281 from the US National Institutes of
410     Health. The multiband echo planar imaging sequence was provided by the University of
411     Minnesota Center for Magnetic Resonance Research. We are grateful to Linda Hood for
412     assistance with data collection, and to Nico Dosenbach and Jackie Hampton for assistance
413     implementing FIRMM.

414          **References**

415     Ardekani BA, Bachman AH, Helpern JA (2001) A quantitative comparison of motion detection
416          algorithms in fMRI. Magn Reson Imaging 19:959–963.

417     Ashburner J, Friston KJ (2004) Rigid Body Registration. In: Human Brain Function, Second.
418          (Frackowiak RSJ, Friston KJ, Frith CD, Dolan RJ, Price CJ, Zeki S, Ashburner J, Penny W,
419          eds), pp 635–653. New York: Elsevier.

420     Badke D'Andrea C, Kenley JK, Montez DF, Mirro AE, Miller RL, Earl EA, Koller JM, Sung S,
421          Yacoub E, Elison JT, Fair DA, Dosenbach NUF, Rogers CE, Smyser CD, Greene DJ
422          (2022) Real-time motion monitoring improves functional MRI data quality in infants. Dev
423          Cogn Neurosci 55:101116.

424     Braver TS (2012) The variable nature of cognitive control: a dual mechanisms framework.
425          Trends Cogn Sci 16:106–113.

426     Cusack R, Vicente-Grabovetsky A, Mitchell DJ, Wild CJ, Auer T, Linke AC, Peelle JE (2015)
427          Automatic analysis (aa): Efficient neuroimaging workflows and parallel processing using
428          Matlab and XML. Front Neuroinform 8:90.

429     Diedrichsen J, Shadmehr R (2005) Detecting and adjusting for artifacts in fMRI time series data.
430          Neuroimage 27:624–634.

431     Dosenbach NUF, Koller JM, Earl EA, Miranda-Dominguez O, Klein RL, Van AN, Snyder AZ,
432          Nagel BJ, Nigg JT, Nguyen AL, Wesevich V, Greene DJ, Fair DA (2017) Real-time motion
433          analytics during brain MRI improve data quality and reduce costs. Neuroimage 161:80–93.

434     Edmister WB, Talavage TM, Ledden PJ, Weisskoff RM (1999) Improved auditory cortex imaging
435          using clustered volume acquisitions. Hum Brain Mapp 7:89–97.

436     Eklund A, Lindquist MA, Villani M (2017) A Bayesian heteroscedastic GLM with application to
437          fMRI data with motion spikes. Neuroimage 155:354–369.

438     Feinberg DA, Moeller S, Smith SM, Auerbach E, Ramanna S, Glasser MF, Miller KL, Ugurbil K,
439          Yacoub E (2010) Multiplexed echo planar imaging for sub-second whole brain FMRI and
440          fast diffusion imaging. PLoS One 5:e15710.

441     Friston KJ, Williams S, Howard R, Frackowiak RS, Turner R (1996) Movement-related effects in
442          fMRI time-series. Magn Reson Med 35:346–355.

443     Greene DJ, Koller JM, Hampton JM, Wesevich V, Van AN, Nguyen AL, Hoyt CR, McIntyre L,
444          Earl EA, Klein RL, Shimony JS, Petersen SE, Schlaggar BL, Fair DA, Dosenbach NUF
445          (2018) Behavioral interventions for reducing head motion during MRI scans in children.

446     Neuroimage 171:234–245.

447   Hall DA, Haggard MP, Akeroyd MA, Palmer AR, Summerfield AQ, Elliott MR, Gurney EM,
448     Bowtell RW (1999) "Sparse" temporal sampling in auditory fMRI. Hum Brain Mapp 7:213–
449     223.

450   Johnstone T, Ores Walsh KS, Greischar LL, Alexander AL, Fox AS, Davidson RJ, Oakes TR
451     (2006) Motion correction and the use of motion covariates in multiple-subject fMRI analysis.
452     Hum Brain Mapp 27:779–788.

453   Jones M, Zhou Z, Bajracharya A, Luor A, Peelle JE (2022) A multi-dataset evaluation of frame
454     censoring for motion correction in task-based fMRI. Aperture Neuro 2:1–25.

455   Lemieux L, Salek-Haddadi A, Lund TE, Laufs H, Carmichael D (2007) Modelling large motion
456     events in fMRI studies of patients with epilepsy. Magn Reson Imaging 25:894–901.

457   Markiewicz CJ, Gorgolewski KJ, Feingold F, Blair R, Halchenko YO, Miller E, Hardcastle N,
458     Wexler J, Esteban O, Goncavles M, Jwa A, Poldrack R (2021) The OpenNeuro resource for
459     sharing of neuroscience data. Elife 10:e71774.

460   McLaughlin DJ, Braver TS, Peelle JE (2021) Measuring the Subjective Cost of Listening Effort
461     Using a Discounting Task. J Speech Lang Hear Res 64:337–347.

462   Oakes TR, Johnstone T, Ores Walsh KS, Greischar LL, Alexander AL, Fox AS, Davidson RJ
463     (2005) Comparison of fMRI motion correction software tools. Neuroimage 28:529–543.

464   Patel AX, Kundu P, Rubinov M, Jones PS, Vertes PE, Ersche KD, Suckling J, Bullmore ET
465     (2014) A wavelet method for modeling and despiking motion artifacts from resting-state
466     fMRI time series. Neuroimage 95:287–304.

467   Peelle JE (2018) Listening effort: How the cognitive consequences of acoustic challenge are
468     reflected in brain and behavior. Ear Hear 39:204–214.

469   Pinheiro JC, Bates DM (2000) Mixed-Effects Models in S and S-PLUS. Springer Science &
470     Business Media.

471   Power JD, Barnes KA, Snyder AZ, Schlaggar BL, Petersen SE (2012) Spurious but systematic
472     correlations in functional connectivity MRI networks arise from subject motion. Neuroimage
473     59:2142–2154.

474   Power JD, Schlaggar BL, Petersen SE (2015) Recent progress and outstanding issues in
475     motion correction in resting state fMRI. Neuroimage 105:536–551.

476   Power JD, Silver BM, Silverman MR, Ajodan EL, Bos DJ, Jones RM (2019) Customized head
477     molds reduce motion during resting state fMRI scans. Neuroimage 189:141–149.

478   Pruim RHR, Mennes M, van Rooij D, Llera A, Buitelaar JK, Beckmann CF (2015) ICA-AROMA:
479     A robust ICA-based strategy for removing motion artifacts from fMRI data. Neuroimage
480     112:267–277.

481   R Core Team (2020) R: A language and environment for statistical computing. Available at:
482     http://r-project.org.

483    Rogers CS, Jones MS, McConkey S, Peelle JE (2023) Listening task, OpenNeuro. Available at:
484         http://dx.doi.org/10.18112/openneuro.ds004285.v1.0.0.

485    Rogers CS, Jones MS, McConkey S, Spehar B, Van Engen KJ, Sommers MS, Peelle JE (2020)
486         Age-Related Differences in Auditory Cortex Activity During Spoken Word Recognition.
487         Neurobiol Lang (Camb) 1:452–473.

488    Smyser CD, Snyder AZ, Neil JJ (2011) Functional connectivity MRI in infants: exploration of the
489         functional organization of the developing brain. Neuroimage 56:1437–1452.

490    Vanderwal T, Kelly C, Eilbott J, Mayes LC, Castellanos FX (2015) Inscapes: A movie paradigm
491         to improve compliance in functional magnetic resonance imaging. Neuroimage 122:222–
492         232.