Real-time feedback reduces participant motion during task-based fMRI

Chad S. Rogers[1], Michael S. Jones[2], Sarah McConkey[2],
Drew J. McLaughlin[3], Jonathan E. Peelle[4, 5, 6]

[1] Department of Psychology, Union College

[2] Department of Otolaryngology, Washington University in St. Louis

[3] Basque Center on Cognition, Brain and Language

[4] Center for Cognitive and Brain Health, Northeastern University

[5] Department of Communication Sciences and Disorders, Northeastern University

[6] Department of Psychology, Northeastern University

Running title: Real-time feedback reduces fMRI motion

Data availability statement: The data that support the findings of this study are openly available in OpenNeuro at https://doi.org/10.18112/openneuro.ds004285.v1.0.0, reference number ds004285.

IRB statement: The study reported here was conducted according to a protocol approved by the Washington University in Saint Louis Institutional Review Board.

Please address post-publication correspondence to:

Dr. Chad Rogers   Dr. Jonathan Peelle
Department of Psychology   Center for Cognitive and Brain Health
Union College   Northeastern University
rogersc@union.edu   j.peelle@northeastern.edu

Abstract

The potential negative impact of head movement during fMRI has long been appreciated.
Although a variety of prospective and retrospective approaches have been developed to help
mitigate these effects, reducing head movement in the first place remains the most appealing
strategy for optimizing data quality. Real-time interventions, in which participants are provided
feedback regarding their scan-to-scan motion, have recently shown promise in reducing motion
during resting state fMRI. However, whether feedback might similarly reduce motion during
task-based fMRI is an open question. In particular, it is unclear whether participants can
effectively monitor motion feedback while attending to task-related demands. Here we assessed
whether a combination of real-time and between-run feedback could reduce head motion during
task-based fMRI. During an auditory word repetition task, 78 adult participants (aged 18–81)
were pseudorandomly assigned to receive feedback, implemented in FIRMM software, or not.
We quantified movement using framewise displacement (FD). We found that motion feedback
resulted in a statistically significant reduction in participant head motion, with a small-to-
moderate effect size (reducing average FD from 0.347 to 0.282). There was some evidence for
a change of effect over the course of six runs, but no clear evidence for practice effects based
on motion feedback. We conclude that under some circumstances real-time feedback may
reduce head motion during task-based fMRI, although its effectiveness may depend on the
specific participant population and task demands of a given study.
66

Introduction

Head movement during fMRI can introduce a number of different challenges to data analysis.
Although long appreciated (Friston et al., 1996), the detrimental influence of head motion on
resting state functional connectivity has only recently been widely recognized (Power et al.,
2015). Rigid body realignment—a mainstay of fMRI analysis for decades—goes some way
towards improving correspondence across images (Ashburner and Friston, 2004), but does not
remove extraneous signal components introduced by movement (Friston et al., 1996). A
common approach for mitigating motion-related artifacts is to include the 6 realignment
parameters (translation and rotation around the X, Y, and Z axes) as nuisance regressors in
first-level models. Additional approaches include wavelet despiking (Patel et al., 2014), ICA
(Pruim et al., 2015), robust-weighted least squares (Diedrichsen and Shadmehr, 2005),
Bayesian approaches (Eklund et al., 2017), and frame censoring (Lemieux et al., 2007).
However, recent work investigating motion correction strategies in multiple data sets suggests
that the optimal strategy may depend on the specific data set and output metric (Jones et al.,
2022), precluding a simple one-size-fits-all solution.
Given the lack of certainty regarding how to best handle head motion during analysis,
reducing head motion in the first place is a particularly appealing option. Common approaches
to reducing head motion during MRI scanning include foam padding and instructing participants
to not move during the scan. Other existing strategies include using custom head molds (Power
et al., 2019) and playing participants a relaxing abstract movie (Vanderwal et al., 2015).
A complementary approach for reducing head motion is to provide participants with
feedback regarding their head movement so they can learn to minimize motion. FIRMM
software[1] (Dosenbach et al., 2017) is one implementation of motion feedback. FIRMM uses
rapid image reconstruction and rigid-body alignment to estimate frame-by-frame movement,
providing visual cues to participants based on estimated movement. Feedback can either be
provided in real time, or between scanning runs. FIRMM has been shown to reduce movement
during resting state scans in both adults (Dosenbach et al., 2017) and in young children
(Greene et al., 2018; Badke D'Andrea et al., 2022).
Despite its success in resting state fMRI, it is unclear whether motion feedback would
lead to similar improvements during a typical task-based fMRI paradigm. Task-based paradigms
have been shown to produce greater head motion both in terms of displacement and rotation
relative to resting state scans (Huang, 2019), and could represent a natural target for motion
feedback. On the one hand, if participants are able to modulate their motion in one experimental
paradigm (namely, resting state studies) it stands to reason they could do so in a different
experimental context (a task-based study). On the other hand, task-based fMRI studies require
participants to attend to the experimental task at hand, which will reduce the cognitive resources
available for noting and processing the feedback, as well as the ability to attend to
proprioceptive cues related to self-motion. In addition, if experimental stimuli and motion
feedback are in the same modality, subjects may also have difficulty accurately processing
motion-related feedback.
To evaluate the degree to which feedback might affect head motion during task-based
fMRI, we assigned participants in a task-based fMRI study of spoken word recognition to
receive feedback (the feedback group) or not (the no-feedback group). All other instructions,
scanning parameters, and task requirements were identical. We included both young and older
adults, and a task (word repetition) that we anticipated would typically require head movement.
Finally, we were able to capitalize on the fact that each subject has many observations
(hundreds of frames) by using linear mixed effects analysis.

---

[1] Now a commercial product offered by Turing Medical (St. Louis, MO).

Method

MRI data are available from OpenNeuro(Markiewicz et al., 2021) at
https://doi.org/10.18112/openneuro.ds004285.v1.0.0, reference number ds004285, and motion
data and analysis scripts are available from https://github.com/jpeelle/motion-feedback.
(Although we collected these data in the context of a task-based fMRI study, described below,
here we focus only on effects of feedback on head motion and not the task results.)

Participants
We tested 78 participants aged 19–81. All reported themselves to have normal hearing, to be
right handed, and not have a history of neurological disease. There were no exclusions based
on sex, gender, race, or ethnicity. Participants were limited to adults because of our scientific
focus on adult language processing. Because the task involved spoken language, all were
native speakers of American English.
Participants were randomly assigned to control (no motion feedback) or intervention
(real-time motion feedback) groups. Demographic characteristics for these groups are shown in
Table 1. Because data collection was ended early due to COVID-19 safety requirements the
groups did not have equal numbers of participants. All provided informed consent under a
process approved by the Washington University in Saint Louis Institutional Review Board.
131
132

Table 1. Participant characteristics

		No feedback	Feedback
Young adults	N	17	31
	Age (mean ± SD)	23.3 (3.06)	22.4 (2.64)
	Sex	13 F, 4 M	28 F, 3 M
Older adults	N	5	25
	Age (mean ± SD)	70.4 (5.22)	71.3 (3.81)
	Sex	5 F, 0 M	17 F, 8 M

134 **Procedure**
135 The procedure is modeled after that in Rogers et al. (2020) and illustrated in **Figure 1**.
136 Participants performed a word repetition task in which on every trial they heard a spoken word,
137 presented in stationary background noise (3 dB SNR), and repeated it back aloud.
138 The no-feedback group was given the following instructions:
139
140 During this task, it is important that you hold your body and head very still.
141 Please stay relaxed, stay alert, and keep your eyes open and on the fixation
142 cross.
143

Figure 1. Summary of task design and processing pipeline. **a)** Schematic of the word repetition task. Participants heard a word presented in the midst of background noise during the gap between volume acquisitions; during the following gap, they repeated the word aloud. **b)** Illustration of the two groups to which a participant could be pseudorandomly assigned. The group without feedback received standard instructions to keep still at the beginning of the study; the feedback group received instructions regarding real-time feedback, a colored crosshair indicating their scan-to-scan motion (framewise displacement; FD) during each run; and a visual report on feedback at the end of each run. **c)** Illustration of time variables available for modeling.

145
146   FIRMM was run in the background for experimenters to monitor motion during scanning
147 sequences and participants were given verbal feedback after each sequence.
148   The group receiving motion feedback was given the following instructions:
149
It is very important to remain still during your MRI so that we can obtain clear
images. Even very small movements that you are not aware of can affect the
image quality. While performing the task, you will be receiving feedback
corresponding to your ability to remain still. This is to help you be aware of any
movements you may be making. You will see a white fixation cross on the
screen. The cross will change to yellow and then red depending on how much
you are moving. It will go back to white if you become still again. Sometimes
even if you are doing your best you will still see a red cross. This may mean that

the computer is too strict and not that you are necessarily doing anything wrong.
Just keep trying your best to keep the cross in the white.
160

During the task, participants viewed a colored cross in the middle of the screen. FD

thresholds were set to show a white cross at < 0.2 mm, a yellow cross at 0.2 mm to < 0.3 mm,

and a red cross at ≥ 0.3 mm. At the conclusion of each run, participants viewed a Head Motion

Report (**Figure 2**) which displayed their performance on a gauge of 0–100 (a percentage score

of 0% to 100%), and a graph of their motion level over time to help visualize total movement

during the session. Participants were encouraged to bring their score closer to 100% on

subsequent sequences.

Figure 2. Head motion report provided to participants following each run of the experiment. The upper right quadrant indicates how many minutes of data with acceptable motion has been collected (under the logic that in a typical resting state paradigm, additional data might be collected to meet a threshold for amount of good data) (Dosenbach et al., 2017).

169
170

MRI acquisition and analysis
MRI data were acquired using a Siemens Prisma scanner (Siemens Medical Systems) at 3 T
equipped with a 32-channel head coil. Scan sequences began with a T1-weighted structural
volume using an MPRAGE sequence [repetition time (TR) = 2.4 s, echo time (TE) = 2.2 ms, flip
angle = 8°, 300 × 320 matrix, voxel size = 0.8 mm isotropic]. Blood oxygenation level-dependent
(BOLD) functional MRI images were acquired using a multiband echo planar imaging sequence
(Feinberg et al., 2010) [TR = 3.07 s, TA = 0.770 s, TE = 37 ms, flip angle = 90º, voxel size = 2
mm isotropic, multiband factor = 8]. We used a sparse imaging design in which there was a 2.3
second delay between scanning acquisitions and the TR was longer than the acquisition time to
allow for minimal scanning noise during stimulus presentation and audio recording of participant
responses (Edmister et al., 1999; Hall et al., 1999). Due to slight variations in scanning
sessions, participants had 763–846 frames of data (median = 794).
Analysis of the MRI data was performed using Automatic Analysis version 5.8.1 (Cusack
et al., 2015) (RRID:SCR_003560) which scripted analyses run in SPM12 (Wellcome Trust
Centre for Human Neuroimaging) version 7487 (RRID:SCR_007037). Functional images
underwent rigid body realignment (i.e, "motion correction"). We used framewise displacement
(FD) to parsimoniously summarize frame-by-frame motion, calculated as the sum of the six
head motion estimates obtained from realignment, with a dimensional conversion of the three
rotations assuming the head is a 50 mm sphere (Power et al., 2012). We supplemented FD
values with differential variance (DVARS), a measure of variations in image intensity. DVARS
was calculated as the root-mean-squared of the time difference in the BOLD signal calculated
across the entire brain, before realignment (Smyser et al., 2011). Although FD and DVARS are
highly correlated, they are not identical (Jones et al., 2022), and we anticipated that these two
metrics might differ in sensitivity to motion feedback.

195   **Results**

196 Movement parameters for young and older adults as a function of feedback are shown in **Figure
197 3**. **Figure 3a** shows FD values for two subjects (our first young adult without feedback, and our
198 first young adult with feedback), as well as the mean FD for all subjects in each feedback group.
199 Although we used all time points in the statistical analysis, for illustrative purposes we also
200 plotted summary values in **Figure 3b** (each point denoting mean FD per subject).

Figure 3. Comparison of average framewise displacement (FD) values for feedback and no-feedback groups. **a)** *Top:* FD values for two example subjects, one with feedback and one without. *Bottom:* Mean FD values for all participants in the feedback or no-feedback group. **b)** Summary of mean FD values as a function of age and feedback. Individual subjects are dots, mean ± SE shown in bars.

We analyzed these data using R version 4.2 (R Core Team, 2020) (RRID:SCR_001905). We conducted linear mixed effects analysis using the *nlme* package (Pinheiro and Bates, 2000). We accounted for temporal autocorrelation (that is, motion at one frame is related to motion during the following frame) using a first-order autoregressive model. The model specification was:

```
m2 <- lme(FD ~ 1 + frame + age_group * feedback,
                random = ~ frame | subject_number,
                data = df,
                method = "ML",
                correlation = corAR1(value =
autocorrelation_estimate, form = ~ frame | subject_number, fixed
= FALSE))
```

238 where `df` is the data frame, `FD` is framewise displacement, and `feedback` is a dichotomous
239 variable denoting whether feedback was provided or not (reference: no feedback). We
240 calculated the `autocorrelation_estimate` from the data (value of 0.52). Model results are
241 shown in **Table 2**.
242
243

Table 2. Fixed effects results for motion (FD) model by frame

	Est	SE	df	t	p	
Intercept	0.4504154	0.3766628	61778	11.958053	< .0001	***
frame	0.0000442	0.00001716	61778	2.572848	0.0101	*
age group	-0.1659770	0.04270876	74	-3.886252	0.0002	***
feedback	-0.1193043	0.04112688	74	-2.900883	0.0049	**
age group x feedback	0.0507396	0.04830427	74	1.050417	0.2969	

*** p < .001; ** p < .01; * p < .05

244
245
246
247   Although we first included time (frame) in the model, in fMRI research it is often
248 convenient to think about continuous scanning runs (sometimes also referred to as "sessions").
249 In the current study, each participant had 6 runs of data. We thus ran an additional model to see
250 whether the effect of feedback was comparable across run. Because time was expressed as 6
251 runs, we did not account for temporal autocorrelation in this model:
252
```
253 m3 <- lme(FD ~ 1 + run + age_group * feedback,
254          random = ~ run | subject_number,
255          data = df,
256          method = "ML",
257          correlation = NULL)
```
258
259
260   Results of this analysis by run are shown in **Table 3**, and FD as a function of run are
261 shown in **Figure 4**. We again found a significant effect of feedback, but the effect of feedback
262 did not significantly differ by run (as one might expect if there were learning effects of the motion
263 feedback).

Figure 4. Mean FD values for all participants in the feedback or no-feedback group as a function of run. Individual subjects are dots, mean ± SE shown in bars.

271
272

Table 3. Fixed effects results for motion (FD) model by run

	Est	SE	df	t	p	
Intercept	0.4524204	0.03491125	61774	12.959158	< .0001	***
run2	0.0198209	0.01118544	61774	1.772030	0.0764	
run3	0.0294646	0.00999432	61774	2.948129	0.0032	**
run4	0.0358940	0.01053085	61774	3.408465	0.0007	***
run5	0.0352148	0.01335915	61774	2.636004	0.0084	**
run6	0.0278481	0.01217601	61774	2.287125	0.0222	*
age group	-0.1769315	0.03945612	74	-4.484260	< .0001	***
feedback	-0.1344319	0.03799784	74	-3.537883	0.0007	***
age group x feedback	0.0724903	0.04462674	74	1.624370	0.1085	

*** p < .001; ** p < .01; * p < .05

273
274
275
Although our analysis was focused on changes in mean FD, we wanted to look in more
detail at how feedback affected the distribution of movement. We thus plotted the density of FD
values for all scans as a function of feedback (**Figure 5**) for FD values up to 2.0. We used
density rather than the count to control for the different numbers of frames in the two groups
(resulting from different numbers of subjects). Although qualitative, this analysis highlights that
participants receiving feedback showed fewer high-FD scans than those receiving lower
feedback, with this effect becoming more apparent at higher FD values—for FD values above
1.5, the feedback group had less than a third of these scans than the group without feedback.
284
285

Figure 5. a) Density plot of all FD values as a function of feedback. **b)** Density of FD values in the feedback condition relative to the no feedback condition. A value of 1 (dashed line) indicates equal density of FD values in the two conditions. The plot illustrates a decrease in high-FD values in the feedback condition, and an increase in low FD values, relative to the no feedback condition.

286
287
We next repeated our primary FD analyses, but using DVARS (rather than FD) as a
dependent measure, under the logic that FD and DVARS may be differently sensitive to effects
of motion, and one or the other may be of more interest in any specific study. Summary results
are shown in **Figure 6** and **Table 4**. Although DVARS was numerically lower for the feedback
group than the no-feedback group, the difference was not statistically significant. Results as a

function of run are shown in **Figure 7** and **Table 5**. Here, the feedback-based reduction in
DVARS was statistically significant (p < .05).
295

Figure 6. Comparison of average DVARS values for feedback and no-feedback groups. **a)** *Top:*
DVARS values for two example subjects, one with feedback and one without. *Bottom:* Mean
DVARS values for all participants in the feedback or no-feedback group. **b)** Summary of mean
DVARS values as a function of age and feedback. Individual subjects are dots, mean ± SE
shown in bars.

Table 4. Fixed effects results for DVARS model by frame

	Est	SE	df	t	p	
Intercept	244.41952	13.012674	61778	18.783188	<.0001	***
frame	0.00611	0.004908	61778	1.244030	0.2135	
age group	-31.76489	14.664553	74	-2.166100	0.0335	*
feedback	-24.40635	14.121338	74	-1.728331	0.0881	
age group x feedback	2.21691	16.585800	74	0.133663	0.8940	

*** p < .001; ** p < .01; * p < .05

Figure 7. Mean DVARS values for all participants in the feedback or no-feedback group as a function of run. Individual subjects are dots, mean ± SE shown in bars.

298
299

Table 5. Fixed effects results for DVARS model by run

	Est	SE	df	t	p	
Intercept	243.49427	12.192275	61774	19.971192	< .0001	***
run 2	3.08907	2.623021	61774	1.177675	0.2389	
run 3	5.56690	2.696872	61774	2.064205	0.0390	*
run 4	7.33254	2.863635	61774	2.560571	0.0105	*
run 5	5.65629	3.768866	61774	1.500794	0.1334	
run 6	2.94432	3.157047	61774	0.932620	0.3510	
age group	-32.41583	13.721953	74	-2.362334	0.0208	*
feedback	-27.06506	13.213980	74	-2.048214	0.0441	*
age group x feedback	6.10355	15.519814	74	0.393274	0.6952	

*** p < .001; ** p < .01; * p < .05

301
In addition to framewise measurements of FD and DVARS, we investigated whether
motion feedback affected the temporal signal-to-noise ratio (tSNR) of the data. We first created
a brain mask using each subject's tissue class segmentation, binarizing at a gray matter
probability of 0.8. We then calculated tSNR (the mean of the signal divided by its standard
deviation, over frames) in every voxel within the brain mask. We then performed two
complementary analyses. First, for each run, we took the mean tSNR over voxels, and used the
same statistical model as for FD and DVARS, shown in **Figure 8** and shown in **Table 6**.
Although the tSNR values were numerically higher in the feedback group, there was no
significant effect of feedback (p = 0.0814).

311
312
313

Figure 8. Comparison of average temporal SNR (tSNR) values for feedback and no-feedback groups. Each participant is represented by a single point per run, averaging the tSNR over all voxels in the gray matter mask.

Table 6. Fixed effects results for tSNR model

	Est	SE	df	t	p	
Intercept	35.62189	2.3494343	389	15.161899	< .0001	***
run	-0.01283	0.1116692	389	-0.114888	0.9086	
age group	5.61719	2.6542796	74	2.116276	0.0377	*
feedback	4.51521	2.5559386	74	1.766556	0.0814	
age group x feedback	-0.11883	3.0020112	74	-0.039584	0.9685	

*** p < .001; ** p < .01; * p < .05

320
321
Finally, we assessed the degree to which feedback may have affected performance on
the behavioral task. Accuracy on the word repetition task is shown in **Figure 9** as a function of
age group and feedback. We used a linear model to test for the impact of feedback on
behavioral accuracy (averaging over experimental runs). We found no significant effect of age
group, no significant effect of feedback, and no significant age x feedback interaction, although
in older adults there was a small numerical decrease in accuracy during the feedback condition.
328

Figure 9. Behavioral accuracy for young and older adults as a function of feedback.

Discussion

Given the pernicious and sometimes unpredictable effects of head movement in fMRI studies,
reducing motion in the first place is on its face an appealing strategy. In the current study we
evaluated the degree to which a combination of real-time and summary motion feedback could
reduce head movement in adult participants by pseudorandomly assigning participants to
feedback or no feedback groups. We found that motion feedback was associated with a
statistically significant reduction in average FD of approximately 0.06, with consistent (though
not significant) changes in DVARS and tSNR. We discuss these findings and their implications
below.
First, our primary finding that providing motion feedback can significantly reduce head
motion in participants undergoing task-based MRI is notable. These findings add motion
feedback to the limited set of tools MRI researchers have for reducing head motion during
acquisition. Although numerous strategies exist for accounting for head motion in preprocessing
and analysis, none are perfect, and it is difficult to ascertain the most appropriate approach
(Ardekani et al., 2001; Oakes et al., 2005; Johnstone et al., 2006; Jones et al., 2022). Reducing
motion during acquisition may therefore be a preferable alternative in many circumstances.
It is important to highlight the magnitude of the effect we observed. Compared to the
participants with no motion feedback, participants with feedback showed a reduction in FD of
approximately .06. Although this may be helpful in reducing motion-related artifacts in data
processing, it is not a large effect (though in theory it might be increased if different types of
feedback were used, or in the context of a different task). It is also important to remember that
we tested healthy adults aged 19–81. It could be that the effect of motion feedback would be
larger (or smaller) in different populations.
That being said, the principal effect of feedback may not be the *average* reduction in FD
observed, but rather a reduction in the number of large motion events. Rapid head motion tends
to have a larger effect on the BOLD signal than slower motion. Thus, reducing the number of
large, quick movements may improve signal quality to a greater extent than a simple reduction
in mean FD. Indeed, we saw the biggest proportional difference between feedback groups at
higher FD values. Interestingly, the reduction in high motion events (**Figure 5b**) roughly
corresponds to an FD of 0.3, the value at which participants received "red" feedback.
An important consideration when implementing real-time feedback during fMRI is
whether doing so significantly changes task demands. Although we found accuracy was
statistically comparable across feedback groups, we cannot rule out other changes in task
demands. Indeed, we observed a numeric (though not significant) decrease in task accuracy in
older adults when they were presented with real-time feedback relative to the no-feedback
group. Broadly speaking, we might expect monitoring the display for feedback, and adjusting
one's movement in response, to engage systems related to cognitive control (sometimes also
referred to as attentional control or executive function) (see also McCabe et al., 2010). In one
framing of cognitive control, cognitive control is proposed to operate in at least two
complementary modes: a *proactive* mode that is concerned with maintaining goal-directed
behavior, and a *reactive* mode that is engaged when goal-directed performance falters (Braver,
2012). There are two implications of such a framework. First, participants who are less able to
engage cognitive control may benefit less from motion feedback compared to those who are
better able to engage control systems. Second, researcher concerns about how task demands
may affect imaging results of interest may depend on the nature of the main fMRI task. For
example, tasks tapping cognitive control or executive function may be more impacted by motion
feedback than those focused on primary sensory or motor systems. In our case, we used a
speech-in-noise task, one that is hypothesized to engage cognitive control networks (Peelle,
2018; McLaughlin et al., 2021).

In our implementation, participants received brief instructions before entering the
scanner, and once in the scanner the display was explained to them. One interesting future
direction would be to provide participants with motion training over a longer period to see
whether they might learn to better limit their motion in the absence of feedback (for example, by
training in a mock scanner). Such an approach would take more time, but would also circumvent
some of the challenges associated with real-time feedback (such as introducing a "dual-task"
situation).
Although we found a significant effect of motion feedback on FD, we did not observe a
significant effect on motion feedback on DVARS (although DVARS values were numerically
lower in the feedback group). FD and DVARS are typically strongly correlated (Jones et al.,
2022), but not identical, and thus the divergence in findings is not entirely unexpected. The
numerical (albeit not significant) reductions in DVARS in participants receiving motion feedback
is consistent with the effect we saw for FD.
In addition, we found that feedback led to increased whole-brain tSNR, which we used
as a proxy of overall image quality. Despite the tSNR being numerically larger in the group
receiving motion feedback, the difference was not statistically significant. However, it is still
possible that small improvements in tSNR would result in appreciable improvements in the
accuracy or reliability of statistical models.
In conclusion, we found that motion feedback as implemented in FIRMM significantly
reduced the amount of head motion we observed during task-based fMRI. Real-time feedback
may thus be well-suited to complement other approaches to motion reduction depending on the
specific needs of a given study.
401
402

Acknowledgments

Work reported here was supported by grant R01 DC014281 from the US National Institutes of
Health. Support was also provided by the Basque Government through the BERC 2022-2025
program and by the Spanish State Research Agency through BCBL Severo Ochoa excellence
accreditation CEX2020-001010-S. The multiband echo planar imaging sequence was provided
by the University of Minnesota Center for Magnetic Resonance Research. We are grateful to
Linda Hood for assistance with data collection, and to Nico Dosenbach and Jackie Hampton for
assistance implementing FIRMM.

References

Ardekani BA, Bachman AH, Helpern JA (2001) A quantitative comparison of motion detection
algorithms in fMRI. Magn Reson Imaging 19:959–963.

Ashburner J, Friston KJ (2004) Rigid Body Registration. In: Human Brain Function, Second.
(Frackowiak RSJ, Friston KJ, Frith CD, Dolan RJ, Price CJ, Zeki S, Ashburner J, Penny W,
eds), pp 635–653. New York: Elsevier.

Badke D'Andrea C, Kenley JK, Montez DF, Mirro AE, Miller RL, Earl EA, Koller JM, Sung S,
Yacoub E, Elison JT, Fair DA, Dosenbach NUF, Rogers CE, Smyser CD, Greene DJ
(2022) Real-time motion monitoring improves functional MRI data quality in infants. Dev
Cogn Neurosci 55:101116.

Braver TS (2012) The variable nature of cognitive control: a dual mechanisms framework.
Trends Cogn Sci 16:106–113.

Cusack R, Vicente-Grabovetsky A, Mitchell DJ, Wild CJ, Auer T, Linke AC, Peelle JE (2015)
Automatic analysis (aa): Efficient neuroimaging workflows and parallel processing using
Matlab and XML. Front Neuroinform 8:90.

Diedrichsen J, Shadmehr R (2005) Detecting and adjusting for artifacts in fMRI time series data.
Neuroimage 27:624–634.

Dosenbach NUF, Koller JM, Earl EA, Miranda-Dominguez O, Klein RL, Van AN, Snyder AZ,
Nagel BJ, Nigg JT, Nguyen AL, Wesevich V, Greene DJ, Fair DA (2017) Real-time motion
analytics during brain MRI improve data quality and reduce costs. Neuroimage 161:80–93.

Edmister WB, Talavage TM, Ledden PJ, Weisskoff RM (1999) Improved auditory cortex imaging
using clustered volume acquisitions. Hum Brain Mapp 7:89–97.

Eklund A, Lindquist MA, Villani M (2017) A Bayesian heteroscedastic GLM with application to
fMRI data with motion spikes. Neuroimage 155:354–369.

Feinberg DA, Moeller S, Smith SM, Auerbach E, Ramanna S, Glasser MF, Miller KL, Ugurbil K,
Yacoub E (2010) Multiplexed echo planar imaging for sub-second whole brain FMRI and
fast diffusion imaging. PLoS One 5:e15710.

Friston KJ, Williams S, Howard R, Frackowiak RS, Turner R (1996) Movement-related effects in
fMRI time-series. Magn Reson Med 35:346–355.

Greene DJ, Koller JM, Hampton JM, Wesevich V, Van AN, Nguyen AL, Hoyt CR, McIntyre L,

Earl EA, Klein RL, Shimony JS, Petersen SE, Schlaggar BL, Fair DA, Dosenbach NUF
(2018) Behavioral interventions for reducing head motion during MRI scans in children.
Neuroimage 171:234–245.

Hall DA, Haggard MP, Akeroyd MA, Palmer AR, Summerfield AQ, Elliott MR, Gurney EM,
Bowtell RW (1999) "Sparse" temporal sampling in auditory fMRI. Hum Brain Mapp 7:213–
223.

Huang J (2019) Greater brain activity during the resting state and the control of activation during
the performance of tasks. Sci Rep 9:5027.

Johnstone T, Ores Walsh KS, Greischar LL, Alexander AL, Fox AS, Davidson RJ, Oakes TR
(2006) Motion correction and the use of motion covariates in multiple-subject fMRI analysis.
Hum Brain Mapp 27:779–788.

Jones M, Zhou Z, Bajracharya A, Luor A, Peelle JE (2022) A multi-dataset evaluation of frame
censoring for motion correction in task-based fMRI. Aperture Neuro 2:1–25.

Lemieux L, Salek-Haddadi A, Lund TE, Laufs H, Carmichael D (2007) Modelling large motion
events in fMRI studies of patients with epilepsy. Magn Reson Imaging 25:894–901.

Markiewicz CJ, Gorgolewski KJ, Feingold F, Blair R, Halchenko YO, Miller E, Hardcastle N,
Wexler J, Esteban O, Goncavles M, Jwa A, Poldrack R (2021) The OpenNeuro resource for
sharing of neuroscience data. Elife 10:e71774.

McCabe DP, Roediger HLI, McDaniel MA, Balota DA, Hambrick DZ (2010) The relationship
between working memory capacity and executive functioning: Evidence for a common
executive attention construct. Neuropsychology 24:222–243.

McLaughlin DJ, Braver TS, Peelle JE (2021) Measuring the Subjective Cost of Listening Effort
Using a Discounting Task. J Speech Lang Hear Res 64:337–347.

Oakes TR, Johnstone T, Ores Walsh KS, Greischar LL, Alexander AL, Fox AS, Davidson RJ
(2005) Comparison of fMRI motion correction software tools. Neuroimage 28:529–543.

Patel AX, Kundu P, Rubinov M, Jones PS, Vertes PE, Ersche KD, Suckling J, Bullmore ET
(2014) A wavelet method for modeling and despiking motion artifacts from resting-state
fMRI time series. Neuroimage 95:287–304.

Peelle JE (2018) Listening effort: How the cognitive consequences of acoustic challenge are
reflected in brain and behavior. Ear Hear 39:204–214.

Pinheiro JC, Bates DM (2000) Mixed-Effects Models in S and S-PLUS. Springer Science &
Business Media.

Power JD, Barnes KA, Snyder AZ, Schlaggar BL, Petersen SE (2012) Spurious but systematic
correlations in functional connectivity MRI networks arise from subject motion. Neuroimage
59:2142–2154.

Power JD, Schlaggar BL, Petersen SE (2015) Recent progress and outstanding issues in
motion correction in resting state fMRI. Neuroimage 105:536–551.

Power JD, Silver BM, Silverman MR, Ajodan EL, Bos DJ, Jones RM (2019) Customized head

molds reduce motion during resting state fMRI scans. Neuroimage 189:141–149.

Pruim RHR, Mennes M, van Rooij D, Llera A, Buitelaar JK, Beckmann CF (2015) ICA-AROMA:
A robust ICA-based strategy for removing motion artifacts from fMRI data. Neuroimage
112:267–277.

R Core Team (2020) R: A language and environment for statistical computing. Available at:
http://r-project.org.

Rogers CS, Jones MS, McConkey S, Spehar B, Van Engen KJ, Sommers MS, Peelle JE (2020)
Age-Related Differences in Auditory Cortex Activity During Spoken Word Recognition.
Neurobiol Lang (Camb) 1:452–473.

Smyser CD, Snyder AZ, Neil JJ (2011) Functional connectivity MRI in infants: exploration of the
functional organization of the developing brain. Neuroimage 56:1437–1452.

Vanderwal T, Kelly C, Eilbott J, Mayes LC, Castellanos FX (2015) Inscapes: A movie paradigm
to improve compliance in functional magnetic resonance imaging. Neuroimage 122:222–
232.

493