Sustainability of seed harvesting in wild plant populations: an insight from a global database of
matrix population models

Anna Bucharova[1],*, Oliver Bossdorf[2], J. F. Scheepens[3] and Roberto Salguero-Gómez[4]

[1] Department of Biology, Philipps-University Marburg, Karl-von-Frisch-Straße 8

35032 Marburg, Germany

[2] Institute of Evolution and Ecology, University of Tübingen, Auf der Morgenstelle 5, 72076 Tübingen,

Germany

[3] Faculty of Biological Sciences, Goethe University Frankfurt, Max-von-Laue-Str. 13, 60438 Frankfurt

am Main, Germany

[4] Department of Biology, University of Oxford, 11a Mansfield Rd, OX1 3SZ, Oxford, UK

*Contact author: anna.lampei-bucharova@biologie.uni-marburg.de, +49 6421 28-26586

Abstract

Seed harvesting from wild plant populations is key for ecological restoration, but may threaten the persistence of source populations. Consequently, several countries have set guidelines limiting the proportions of harvestable seeds. However, these guidelines are so far inconsistent, and they lack a solid empirical basis. Here, we use high-resolution data from 298 plant species to model the demographic consequences of seed harvesting. We find that the current guidelines do not protect populations of annuals and short-lived perennials, while they are overly restrictive for long-lived plants. We show that the maximum possible fraction of seed production – what can be harvested without compromising the long-term persistence of populations – is strongly related to the generation time of the target species. When harvesting every year, this safe seed fraction ranges from 80% in long-lived species to 2% in most annuals. Less frequent seed harvesting substantially increases the safe seed fraction: In the most vulnerable annual species, it is safe to harvest 5%, 10% or 30% of population seed production when harvesting every two, five or ten years, respectively. Our results provide a quantitative basis for seed harvesting legislations worldwide, based on species' generation time and harvesting regime.

Significance:

The UN Decade on Ecosystem Restoration, 2021-2030, foresees upscaling restoration, and the demand for native seed is skyrocketing. Seeds for restoring native vegetation are often harvested in wild, but too intensive harvest can threaten the donor populations. Existing guidelines that set limits to wild seed harvest are mostly based on expert opinions, yet they commonly lack empirical basis and vary among regions in one order of magnitude. We show that the current guidelines urgently need to be reformulated, because they are overly restrictive in long-lived species, while they do not protect annual plants from extinction. Using matrix population models of nearly 300 plant species, we provide a quantitative basis for a new seed harvesting legislation world-wide.

Introduction

The restoration of degraded ecosystems is a major goal of global nature conservation (1). We are in

the middle of the 'UN Decade on Ecosystem Restoration' (2), with a key goal to reverse the

destruction and degradation of billions of hectares of ecosystems. However, ecological restoration at

such scales requires high volumes of plant seeds for the re-establishment of native vegetation (3).

Although there is a growing industry for the production of wild plant seeds in specialised seed

orchards (4, 5), large-scale harvesting of seeds from wild populations is still common in ecological

restoration, and is projected to continue growing (6). Seed harvesting is particularly common for

plant species that are long-lived or difficult to cultivate (7–10).

With increasing demands for wild plant seeds, there is a growing risk of driving source populations to

local extinction (11, 12). Moreover, donor populations are often remnants of habitats with high

conservation value (11, 13). Some regions, in particular the US (14), Australia (15), and Europe (16,

17), have therefore begun to set limits for the maximum fraction of seeds that can be harvested

annually from wild plant populations, to prevent significant negative effects on their long-term

viability ('*safe seed fraction*', hereafter). Notably, though, the safe seed fraction guidelines are

inconsistent across countries, with *e.g.* 20% harvest allowed in the US (14) and 10% in Australia (15),

but only 2-10% in Germany, depending on plant growth type (16). When the harvest does not take

place annually, some guidelines permit higher safe seed fractions (16). In general, however, these

guidelines are mostly based on expert opinion and lack a solid quantitative basis.

Only a few studies have experimentally tested the effects of seed harvesting on wild populations.

However, these studies are either focused on individual species or specific ecosystems (11, 18, 19).

More effective rules would require collection of data across multiple species and ecosystems, but

this course of action is labour- and cost-demanding. As an alternative to collecting new data, Menges

and colleagues (12) used published plant matrix population models to link seed harvesting to the

probabilities of population extinction for 22 perennial species. While this study is widely used to back

up seed collection guidelines for rare species for ex-situ conservation (e.g., (15, 20), the species set is

limited to mostly herbaceous perennials of temperate and subtropical North America. To obtain a

quantitative basis for predicting the effects seed harvesting on wild populations globally, data from

many more species across life histories and ecosystems are essential.

Here, we employed a modelling approach and simulated seed harvesting for 298 plant species

ranging from annuals to long-lived trees from many habitats around the globe using matrix

population models stored in the the COMPADRE Plant Matrix Database (21, 22), Table S1.

Specifically, we (1) tested the efficacy of current guidelines at safeguarding long-term population

persistence, (2) identified traits that are associated with species vulnerability to seed harvesting, and

(3) used the trait that best determines species vulnerability to seed harvesting, generation time, to

predict safe seed fraction, and formulated quantitative basis for seed harvesting in wild plant

populations world-wide.

Results

To test how well the current safe seed fraction guidelines protect source populations from

overharvesting, we modelled the maximal possible harvest fractions permitted in the US, Australia,

and Germany. To allow comparison across species, we expressed effects of seed harvesting as

relative population sizes, where 1 indicates no effect, 0 indicates extinction, and *e.g.* 0.8 represents a

20% reduction of population size in comparison to the population size that would be reached without

seed harvesting. Seed harvesting according the existing safe seed fraction guidelines results in rather

variable relative population sizes among species (Figure 1). For instance, the current US guidelines

(20% seed harvesting allowed) protect long-lived palms, with relative population sizes of 0.6 to 1

after 30 years, but would drive all 10 annual plants in our data to extinction (Fig. 1). With the more

restrictive German guidelines (2% seed harvesting allowed), annual plants are projected to persist,

with relative population sizes of 0.54 to 0.63 after 30 years. Within all other plant growth types, the

effects of seed harvesting on the relative population sizes are much more variable. For example, with

the 20% seed harvesting currently allowed in the US, the predicted relative population sizes of

herbaceous perennials would range from 0 (local extinction) to 1 (no effect) after 30 years, while that

of shrubs would range from 0.12 to 0.99, of succulents from 0.27 to 0.99, and of trees from 0.18 to

0.99 (Fig. 1).

We next examined whether and which life history traits are better predictors of seed harvesting

impacts (Figure 2). We found out that generation time, the mean age of reproductive individuals in

the population, is the strongest predictor of population vulnerability to seed harvesting. This life

history trait alone explains 52.3% of the variation in harvesting vulnerability, and vulnerability to seed

harvesting decreases with increasing generation time (Fig. 2B). Four other life history traits are also

significantly related to seed harvesting vulnerability (Fig. 2B) – species that reproduce more

frequently and/or postpone their first reproductive event are more vulnerable to seed harvesting,

while species with clonal reproduction and/or persistent seed banks are less vulnerable – but the

predictive power of these traits is low (Fig. 2A, Table S3). Population vulnerability also differs

significantly among plant growth types, but with minor effects (Fig. 2C, Table S3). All five life history

traits together explain 62.3% variability in vulnerability to seed harvesting among species.

To improve the efficacy of seed harvesting regulation, we then used the best predictor of species

vulnerability to seed harvesting, generation time, to estimate safe seed fraction across species. For

annual harvesting, the safe seed fraction ranges from close to 0% to 100%, with an average of 2.3%

(95% CI: 0.5-4.1%) for annual and biennial plants, 10.1% (6.8-14.2%) for species with a 5-year

generation time, and 40.1% (36.4-43.7%) for species with generation times of 20 years (Fig. 3A). With

simulated harvesting only every two years, the safe seed fraction for annuals and biennials increases

from 2.3% to 5.3 % (2.7-7.9%), and with a 5-year or 10-year harvesting interval to 11.3% (6.5-16.0%)

and 30.3% (23.8-36.8%), respectively (Fig 3B-D). For plant species with generation times above two

years, a 5-year harvesting cycle resulted in an average safe seed fraction of >30% (Fig. 3C). While safe

seed fraction critically depends on generation time, there is substantial residual variation among

species.

The estimated safe seed fraction for each species was not substantially affected by environmental

stochasticity. The median of safe seed fractions based on models that included environmental

stochasticity (see methods) was on average 1.8% larger than the safe seed fraction based on the

mean models for each species, yet they were closely correlated (Figure S3).

Discussion

Seed harvesting in wild population is indispensable for ex-situ conservation and ecosystem

restoration, but overharvesting can threaten source populations (13). Consequently, some countries

have introduced limits that restrict wild seed harvesting (14–16). Here, using data from wild

populations of 298 plant species from five continents, we show that the current seed harvesting

guidelines are often ineffective: existing guidelines do not protect populations of annuals and short-

lived perennials, while they are overly restrictive for long-lived plants. Based on generation time, the

trait that best predicts seed harvesting vulnerability, we estimate that safe seed fraction varies from

2% in annual and biennial plants to 80-100% in long-lived plants, when seeds are harvested annually.

Lower frequency of harvesting allows for higher seed fractions in a viable way. The safe seed

fractions presented here can serve as a solid quantitative basis for seed harvesting regulations

globally.

When wild seed harvesting follows the existing safe seed fraction guidelines, the effects on

population sizes can vary from no effect to extinction, depending on the species. For example, annual

seed harvesting of 20% of the annual seed production, as currently recommended in the US (14),

would have small effect on palms, trees or some herbaceous perennials, but it would drive all

annuals plants to extinction within three decades. In reality, extinction will be less common because

we modelled an extreme scenario when seeds are harvested every growing season for 30

consecutive years from the same population, which is possible but uncommon (13). Nevertheless,

the high variability in model outcomes highlights that effective safe seed fraction guidelines must be

more nuanced than one-size-fits-all – one safe seed fraction for all species – as currently

implemented in many regions (14, 15, 17).

The current German safe seed fractions guidelines are plant growth-type specific (16). For annual

plant species, the safe seed fraction is 2% when harvesting annually, which in our modelling does not

cause unacceptable population declines (Figure 1). For herbaceous perennials, the safe seed fraction

in Germany is set to 10% for annual harvest, yet this threshold leads to a wide range of relative

population sizes, from substantial population declines to no effects. The variability within the

herbaceous perennials is even stronger when following the US guidelines (20% of annual seed

production). Plant growth type alone is thus a poor predictor of species vulnerability to seed

harvesting.

Over 60% of the vulnerability to seed harvesting is predicted by life history traits. The highest

predictive value in our analyses offers generation time, which alone predicts the seed harvesting

vulnerability by more than 50%. Population growth rates in long-lived species are generally

insensitive to changes in fecundity (23, 24). Indeed, (12) showed that long-lived plants are relatively

insensitive to seed harvesting. Other life history traits in the present study have much smaller

predictive power for seed harvesting impacts. For instance, species with higher iteroparity (*i.e.*

reproducing more than once during their life cycle), and species that are later sexually mature, are

more vulnerable to seed harvesting, while clonal species and species with permanent soil seed banks

are less vulnerable. The buffering effect of seed bank against the effects of seed harvesting are well

supported by the literature (25). However, the relatively small effect of clonality on the impacts of

seed harvesting is surprising, since clonality provides an alternative reproduction independent of

seed production, and has been experimentally identified as a major predictor of vulnerability to seed

harvesting in grassland plants (18). This discrepancy is likely because many matrix population models

calculate generation times of individuals originated from seeds, *i.e.* genets. Clonal reproduction thus

leads to longer generation times of the genets (24, 26), and explains little additional variability in

vulnerability to seed harvesting above what is already explained by generation time as the more universal predictor.

To provide a universal quantitative basis for seed harvesting guidelines, we estimated safe seed fraction as a function of generation time, the best predictor of vulnerability to seed harvesting (Figure 3). The lowest safe seed fractions are in annuals and biennial, 2.3% for annual harvest, which is close to the current German guidelines of 2%, (16). The safe seed fraction continuously increases with generation time, but remains below 10% for plants with generation times of five years and less. Adhering to such low seed safe fractions is possible only when collecting seed manually, yet this is very labor intensive. In grasslands, seeds are commonly harvested using combine harvesters, which typically removes 30% of the ripe seeds (27). Such a high proportion is safe for annual harvesting only in species with generation time above 15 years. Grasslands are dominated by annuals and herbaceous perennials, of which 60% in our dataset have generation times below 15 years. Annual seed removal with combine harvesters thus threatens a substantial proportion of grasslands species, especially non-clonal forbs and annuals (18).

Less frequent harvesting allows higher safe seed fractions. Harvesting seeds less often is already suggested as a precautional principle in some guidelines (*e.g.* (13, 17), although mostly without a clear specification of safe seed fractions and harvesting frequencies. Less frequent harvesting is relevant especially for species with short generation times, where the safe seed fraction is the lowest. In annual and biennials, the safe seed fraction increases from 2.3% for annual harvesting to 5% when harvesting every second year, 11% every five years and 30% every 10 years. Importantly, harvesting at 10-year intervals allows to collect 30 % of the seed production even in the most vulnerable species. Seed harvesting with combine harvesters, which collects on average 30% of the seed (27), should be sustainable even in drylands with high proportion of annual plants, if done at sufficiently long intervals.

Seed harvesting is less problematic in species with long generation times. In species with generation

times above 20 years (most trees and palms, many shrubs and some herbaceous perennials (28)),

safe seed fractions are above 40% when harvesting every year, and above 80% when harvesting less

frequently. Previous empirical and modelling studies also reported that long-lived species are rather

insensitive to seed harvesting (12, 18, 29), although too frequent and too intense harvesting can

deplete populations of seedlings (19). Even in long-lived species, it might thus be beneficial to omit

seed harvesting in some years to give populations opportunities for juvenile recruitment.

Our results demonstrate the demographic impact of seed harvesting, and how it depends on plant

life histories. Yet, we could have overestimated harvesting impacts for three reasons. First, our

analyses are based on matrix population models of species averaged across years and sites, but

temporal or spatial variation in demographic rates could buffer some impacts of seed harvesting (30).

Indeed, incorporating environmental and demographic stochasticity into our models in a subset of

species resulted in safe seed fractions on average 1.8% larger, confirming that matrix averaging may

cause overestimation, but the effect was small. Second, our approach assumes plant populations to

be seed-limited. However, longer-lived plants are often limited by safe sites rather than seeds,

whereas seed limitation is more common in short-lived species (31). It is thus likely that in longer-

lived species the effects of seed harvesting are even less severe than our findings suggest, but for

annuals and short-lived forbs – the most vulnerable to seed harvesting – our results are more likely

to be accurate. A specific case of safe-site limited habitats are European seminatural meadows that

are annually mown with the biomass, including a large proportion of seed, used as fodder for

domestic animals. Species growing in this ecosystem are likely adapted to regular seed removal and

thus less vulnerable to seed harvesting than predicted by our models. Third, our models do not

incorporate maximal carrying capacities, because this information is rarely available for matrix

population models. In populations with high population growth rates and close to carrying capacity

of the environment, matrix models still predict population growth even though the population

already reached maximal space occupancy. In such cases, seed harvesting might have much smaller

effect than predicted.

Seed harvesting in wild populations should be generally accompanied by monitoring of the harvested

sites. Our results provide the currently best quantitative basis for sustainable seed harvesting in wild

populations. Yet, they are model results, and all models are simplifications of the reality as it is

impossible to capture the full complexity of the real world (32). As a precaution, and to be able to

adjust harvesting practice if necessary, it is therefore important to monitor the harvested sites. The

safe seed fractions presented here cause only very slow population declines, maximum 2% per year,

and monitoring every few years should be sufficient to detect unexpected negative effects on

population sizes before the population would be irreversibly damaged.

In summary, we show that seed harvesting in wild populations is possible and allows long-term

population persistence, but the harvesting must be guided by the critical factors of plant generation

time and harvesting frequency. For longer-lived species, harvesting large fractions of seeds is unlikely

to harm wild populations, particularly if seeds are not harvested every year. For short-lived species,

though, more caution is necessary. A profitable harvesting of 30% of the seeds of annual species may

only be possible if the harvesting takes place only every 10 or more years. However, ultimately, even

with improved guidelines, seed harvesting from wild populations is unlikely to cover the growing

worldwide needs of ecological restoration (33). The ambitious targets of the UN Decade on

Ecosystem Restoration (2) may only be reached with professional, large-scale seed production in

seed orchards (4, 34, 35).

Methods

We used data stored in The COMADRE Plant Matrix Database (version 5.0.0. last accessed 25.8.2019

(22), and selected matrix population models for 298 species (SM, section 1). As the ultimate goal of

this study was to simulate seed harvesting, we selected field-based models for angiosperms with

clearly defined sexual reproduction (SM 1.3 for details). For the majority of studies in COMPADRE,

matrix population models are available for several annual transitions and populations. For all

calculations, except the stochastic simulations (see below), we used a single MPM per species

averaged across all years and populations available for that species. Below we briefly outline our

methods; a more detailed description is available in online supplementary information.

To test how well the current guidelines safeguard long-term populations persistence, we used matrix

population models to calculate 30-year projections of population sizes. We simulate seed harvesting

as a reduction of the sexually produced new recruits. We generally modelled the most extreme

scenario: the highest permitted seed fraction harvested every year. To allow comparison across

species, we expressed effects of seed harvesting as relative population sizes, where *e.g.* 0.8

represents a 20% reduction of population size and 0.3 a 70% reduction over 30 years, in comparison

to the population sizes that would be reached without seed harvesting (SM, section 4). As the effects

of seed harvesting were independent of the biogeographic origins of the examined species (Table

S2), we generally used all species in our dataset to test the guidelines of specific countries. We

present the results separately for different growth types, as in the German the guidelines the

recommended safe seed fractions are growth-type specific (16).

To find a better predictor of safe seed fraction than the growth types, we examined whether and

which life history traits were better predictors of seed harvesting impacts (Figure 2). To enable

practitioners to apply our findings, we restricted our analyses to five key life history traits readily

available from public databases (21, 22, 36) or easy to estimate in the field: generation time, mean

age at sexual maturity, the degree of iteroparity (frequency of reproduction) and clonality, and seed

bank persistence (Figure 2, SM section 5). We then related these traits to the vulnerability of our 298

species to seed harvesting, defined as the slope of the relative decrease in population size with

increasing seed harvesting (SM sections 3 and 6, Table S3).

To provide a quantitative basis for improving seed harvesting guidelines, we used generation time,

the best predictor of species vulnerability to seed harvesting, to estimate safe seed fractions across

species (SM section 7). The safe seed fractions were defined as the proportions of seed production

where annual removal caused a <50% decrease of population sizes during 30 years of continuous

seed harvesting, compared to the same populations without seed harvesting. A 50% decrease over

30 years corresponds to an annual decrease of about 2%. Importantly, this threshold ensures a >95%

probability of population viability under environmental stochasticity in all analysed species but one

(Figure S4).

To understand how environmental stochasticity affected our prediction for seed harvesting based on

mean matrix population models, we simulated the effects of environmental stochasticity on

population dynamics (SM section 8). This was possible in 108 species for which we had at least three

spatial or temporal replicate matrix population models (so called individual models). We simulated

environmental stochasticity as projecting population vector by randomly drawn individual matrix

population models in each step, replicated 1000 times to obtain probability distributions of seed

harvesting impacts. To understand how robust our estimates were to environmental stochasticity,

we compared the safe seed fractions based on the mean matrix models to the respective medians of

the safe seed fractions based on stochastic simulations (SM section 8.1). We also used stochastic

simulations to test whether the thresholds of 50% population declines (see above) effectively

prevented populations from extinction (SM section 8.2).

1.   IPBES, *The IPBES assessment report on land degradation and restoration.*, L. Montanarella, R.

Scholes, A. Brainich, Eds. (Secretariat of the Intergovernmental Science-Policy Platform on

Biodiversity and Ecosystem Services, 2018).

2.   UN Decade on Ecosystem Restoration 2021-2030. Available at:

https://www.decadeonrestoration.org/.

3.   N. Hölzel, E. Buisson, T. Dutoit, Species introduction - a major topic in vegetation restoration.

Applied Vegetation Science **15**, 161–165 (2012).

4.   M. Conrady, *et al.*, Plants cultivated for ecosystem restoration can evolve toward a

domestication syndrome. *Proceedings of the National Academy of Sciences* **120**,

e2219664120 (2023).

5.   P. G. Nevill, *et al.*, Seed production areas for the global restoration challenge. *Ecology and*

Evolution (2016).

6.   T. A. Jones, Native Seeds in the Marketplace: Meeting Restoration Needs in the

Intermountain West, United States in *Rangeland Ecology and Management*, (Elsevier Inc.,

2019), pp. 1017–1029.

7.   *An Assessment of Native Seed Needs and the Capacity for Their Supply: Final Report* (National

Academies Press, 2023).

8.   L. Broadhurst, *et al.*, Seeding the future - the issues of supply and demand in restoration in

Australia. *Ecological Management & Restoration* **16**, 29–32 (2015).

9.   R. M. Stevenson, Commercial Wildland Harvested Seed and the Utah Connection.

Rangelands **38**, 273–277 (2016).

10.   A. White, J. B. Fant, K. Havens, M. Skinner, A. T. Kramer, Restoring species diversity: assessing

capacity in the U.S. native plant industry. *Restoration Ecology* **26**, 605–611 (2018).

11.   J. C. Meissen, S. M. Galatowitsch, M. W. Cornett, Risks of overharvesting seed from native

tallgrass prairies. *Restoration Ecology* **23**, 882–891 (2015).

12. E. S. Menges, E. O. Guerrant, S. Hamze, "Effects of seed collection on the extinction risk of

perennial plants" in *Ex Situ Plant Conservation: Supporting Species Survival in the Wild*, E. O.

Guerrant, K. Havens, M. Maunde, Eds. (Island Press, 2004), pp. 305–324.

13. S. Pedrini, *et al.*, Collection and production of native seeds for ecological restoration.

Restoration Ecology **28**, S228–S238 (2020).

14. Plant Conservation Alliance, "National seed strategy for rehabilitation and restoration."

(2015).

15. M. A. J. Yenson, *et al.*, Plant Germplasm Conservation in Australia: strategies and guidelines

for developing, managing and utilising ex situ collections. Third edition. (2021).

16. R. Prasse, D. Kunzmann, R. Schröder, "Development and practical implementation of minimal

requirements for the verification of origin of native seeds of herbaceous plants (in German).

In cooperation with Verband Deutscher Wildsamen- und Wildpflanzenproduzenten. DBU,

reference no. 23931, 166p." (2010).

17. ENSCONET, "Seed collecting manual for wild species" (2009).

18. J. C. Meissen, S. M. Galatowitsch, M. W. Cornett, Assessing long-term risks of prairie seed

harvest: what is the role of life-history? *Botany* **95**, 1081–1092 (2017).

19. C. A. Peres, *et al.*, Demographic threats to the sustainability of Brazil nut exploitation. *Science*

(New York, N.Y.) **302**, 2112–4 (2003).

20. Center for Plant Conservation., CPC Best Plant Conservation Practices to Support Species

Survival in the Wild. Center for Plant Conservation, Escondido, CA. (2019).

21. R. Salguero-Gómez, *et al.*, The compadre Plant Matrix Database: an open online repository

for plant demography. *Journal of Ecology* **103**, 202–218 (2015).

22. COMPADRE Plant Matrix Database (2019). Available from: https://www.compadre-db.org

[25.8.2019, Version 5.0.0]. (2019). [Accessed 25 August 2019].

23. M. Franco, J. Silvertown, A comparative demography of plants based upon elasticities of vital

rates. *Ecology* **85**, 531–538 (2004).

24. Z. Janovský, T. Herben, Reaching similar goals by different means – Differences in life-history

strategies of clonal and non-clonal plants. *Perspectives in Plant Ecology, Evolution and*

Systematics **44**, 125534 (2020).

25. V. M. Adams, D. M. Marsh, J. S. Knox, Importance of the seed bank for population viability

and population monitoring in a threatened wetland herb. *Biological Conservation* **124**, 425–

436 (2005).

26. L. C. de Witte, J. Stöcklin, Longevity of clonal plants: why it matters and how to measure it.

Annals of Botany **106**, 859–870 (2010).

27. M. Scotton, M. Ševčíková, Efficiency of mechanical seed harvesting for grassland restoration.

Agriculture, Ecosystems and Environment **247**, 195–204 (2017).

28. R. Salguero-Gómez, *et al.*, Fast-slow continuum and reproductive strategies structure plant

life-history variation worldwide. *Proceedings of the National Academy of Sciences of the*

United States of America **113**, 230–5 (2016).

29. J. C. Meissen, S. M. Galatowitsch, M. W. Cornett, Risks of overharvesting seed from native

tallgrass prairies. *Restoration Ecology* **23**, 882–891 (2015).

30. J. Villellas, D. F. Doak, M. B. García, W. F. Morris, Demographic compensation among

populations: what is it, how does it arise and what are its implications? *Ecology Letters* **18**,

1139–1152 (2015).

31. L. A. Turnbull, M. J. Crawley, M. Rees, Are plant populations seed-limited? A review of seed

sowing experiments. *Oikos* **88**, 225–238 (2000).

32. G. E. P. Box, Science and Statistics. *Journal of the American Statistical Association* **71**, 791–

799 (1976).

33. S. Díaz, *et al.*, Pervasive human-driven decline of life on Earth points to the need for

transformative change. *Science (New York, N.Y.)* **366** (2019).

34.   M. Conrady, C. Lampei, O. Bossdorf, W. Durka, A. Bucharova, Evolution during seed

production for ecological restoration? A molecular analysis of 19 species finds only minor

genomic changes. *Journal of Applied Ecology* **59**, 1383–1393 (2022).

35.   E. K. Espeland, *et al.*, Evolution of plant materials for ecological restoration: Insights from the

applied and basic literature. *Journal of Applied Ecology* **54**, 102–115 (2017).

36.   J. Kattge, *et al.*, TRY - a global database of plant traits. *Global Change Biology* **17**, 2905–2935

(2011).

Figure 1. Predicted effects of 30 years of continuous seed harvesting on the relative population sizes

of 298 plant species worldwide, using the current guidelines of countries where legislation exists:

USA, Germany, and Australia. Points represent individual species. The data result from simulation of

seed harvesting using matrix population models parameterised with data from natural populations.

Herb. = Herbaceous; n = numbers of species included.

Figure 2. Associations of plant life histories and growth forms with variation in seed harvesting

vulnerability across 298 plant species, as calculated from matrix population models parameterised

with data from natural populations. (A) Proportion of variability explained by different life history

traits, and (B) their effect estimates. (C) The fitted values of vulnerability for different growth types.

Estimates in (B) and (C) are presented with their 95% credible intervals. As both vulnerability to seed

harvesting and all explanatory variables were standardised prior the analysis, the slope estimates are

in arbitrary units. (D) Definitions of the five examined life history traits (for calculation see Table S2).

Herb. = herbaceous. Significance levels: * P<0.05, ** P<0.01, *** P<0.001, See Table S3 for detailed

model results.

(D)
Generation time: the mean age of reproductive individuals in a population.
Iteroparity: the frequency of reproduction throughout the lifespan of an individual, with high/low Demetrius' entropy (S) values for highly iteroparous/semelparous populations.
Age at sexual maturity: the average number of years after which individuals in a population become sexually reproductive.
Seed bank: the mean life expectancy of seeds in the soil seed bank.
Clonality: the per-capita clonal contributions, weighted by the relative frequency of individuals in each stage along the life cycle of the species.

386 **Figure 3.** Relationships between the generation times of 298 plant species and their safe fractions for

387 seed harvesting, estimated at different harvesting frequencies. The safe seed fraction is the

388 maximum proportion of annual seed production of a population that can be harvested without

389 reducing the relative population size to below 50% in 30 years.

Supplementary Materials for

Sustainable seed harvesting in wild plant populations

Anna Bucharova*, Oliver Bossdorf, J. F. Scheepens and Roberto Salguero-Gómez

*Corresponding author: anna.lampei-bucharova@uni-marburg.de

Materials and Methods, Supplementary results

To quantify the effect of seed harvesting on wild plant populations, we used matrix population models (*35*). We first tested the impacts of seed harvesting by simulating the regulatory recommendations on seed harvesting in the wild of three regions where such regulations are in place (Australia, Germany and USA). Second, we calculated the population vulnerability to seed harvesting for each of the 280 plant species examined. Third, we related those effects to plant key life history traits (*i.e.* defining characteristics of their life cycles; e.g. generation time, age at maturity). In the fourth step, we used the life history traits that explained most of the vulnerability of natural populations to seed harvesting to formulate biologically-sound management recommendations. The ultimate goal of these recommendations is to introduce a threshold to seed harvesting so that (i) the population size does not decline more than by 50% over 30 years of consecutive (*i.e.* annual) seed harvest and (ii) the population may still have a 95% probability of persistence. All calculations and statistics were performed in R (*36*), and the reproducible, commented scripts are found as *Auxiliary material and will be available at Zonedo upon acceptance.*

1   Matrix population models

1.1   General introduction

Matrix population models (MPMs, hereafter) are a widely used tool for investigating population dynamics (*35*). Briefly, an MPM describes the life cycle of an organism in terms of age, size and/or developmental stages along its life cycle and the transitions between stages, usually from one year to the next, as well as the sexual and clonal per-capita contributions to the population by individuals in each of those stages (Figure S1). One of the many applications of MPMs is to project the dynamics of a population through time (*35*), whereby a long-term population growth rate can be estimated (Figure S1). Importantly here, MPMs can also be used to calculate a wide range of population characteristics such as life history traits (*37*), extinction probability (*38*), and the effects of different hypothetical events (such as seed harvesting) on the long-term viability of a population (*39*, *40*).

In this study, we used MPMs to simulate seed harvesting as reduction of the per-capita contribution(s) describing seed production (Figure S1). We did so by simulating the harvesting of newly produced seeds while keeping all other demographic processes unaltered. The resulting MPM thus describes the population dynamics in a year where seed harvesting took place.

1.2 COMPADRE database

We used data stored in THE COMPADRE Plant Matrix Database (version 5.0.0.), last accessed 25.8.2019 (20). In this version, COMPADRE contains 9121 MPMs from 647 published works describing life cycles of 760 plant species, ranging from algae to trees worldwide. MPMs in the database are accompanied by extensive metadata including the continent where the study was carried out, whether it was carried out in captivity or in the wild, and standardized information about each life cycle stage in three categories: propagules, individuals photosynthetically active, and individuals in vegetative dormancy. In the vast majority of MPMs in COMPADRE, the full MPM **A** is divided into three submatrices (37): **U** includes demographic processes that depend on survival of individuals alive at the beginning of the census (i.e., progressive growth, stasis, retrogressive growth, seed bank persistence, and vegetative dormancy), **F** includes sexual reproduction (e.g. production of seeds and juveniles), and **C** includes clonal reproduction (*i.e.* vegetative reproduction through ramets), such that

$$\boldsymbol{A} = \boldsymbol{U} + \boldsymbol{F} + \boldsymbol{C} \qquad\qquad eq.\ 1$$

1.3 Selection of the MPMs

We selected species and MPMs from COMPADRE based on the following criteria to allow for inter-specific comparisons to answer our questions:

- Only angiosperms and gymnosperms, since the ultimate goal of this study is to simulate the effect of seed harvesting on seed-producing plants.
- MPMs parameterised from field data from wild populations and under unmanipulated conditions, because the aim of this study is to understand the effect of seed harvest on natural, wild populations.
- MPMs for which the sexual reproduction component had been quantified explicitly, and separated from other processes in order to allow us to accurately perturb sexual reproduction (seed production; see below).
- MPMs that are irreducible, ergodic, and primitive, so the dominant eigenvalue (population growth rate) and other key properties could be calculated (35).
- When multiple studies per species were available (n = 235 species), we selected the single study per species that:
  - documented a seed bank, because inclusion of this transition in MPMs is vital to correct estimation of life history traits (41)

465   ▪   contained the highest number of individual MPMs (*i.e.*, from more populations or
466   more years, see SM section 1.4) to use the most representative demographic
467   information for the target species.

These selection criteria resulted in 467 MPMs from 467 plant species. Next, we checked the reliability
of incorporating a seed bank in them or not. While survival of seeds in the seed bank is well
documented in many demographic studies (*42*), between 42.9% and 47.3% of studies using MPMs in
plant species unjustifiably exclude seed banks (8), thus assigning seedlings in year *t* to reproductive
plants in *t*-1 (e.g. (*43*)). However, this assumption is only correct in species with a transient seed bank,
i.e. seeds survive in the soil less than one year and thus, do not form a permanent soil seed bank (*41*).
For those studies in our list where seed banks were not explicitly considered in their MPMs, we verified
whether the species indeed have only a transient seed bank or not. We did so by carefully examining
the original source of the MPM(s). If the source did not mention a seed bank, we further searched in
the TRY database (*21*) for its potential existence. Consequently, we excluded 169 species where seed
banks were unjustifiably excluded from their MPMs.

In twelve species, the simulated seed harvesting (SM section 2) did not cause any changes of
population sizes, which suggests that generative reproduction was not correctly incorporated in these
MPMs. We excluded these species from the further analysis.

This final selection criterion resulted in a dataset of 280 species (each with a representative MPM)
from 83 plant families. This is the final set of species and data that were used for the simulations
described below (**Error! Reference source not found.**).

1.4   Mean MPMs vs individual MPMs

For the majority of studies in COMPADRE, MPMs are available for several annual transitions and
populations. This was also the case in our final dataset. For all calculations, except in the case of
stochastic simulations (Section 8), we used a single *mean* MPM per species across all years and
populations of demographic data available for that species. This mean MPM was calculated as the
element-by-element arithmetic mean of the aforementioned MPMs, or pooled directly (e.g. weighted
mean by sample size) from the individual-level data when provided by the author in the publication or
through personal communications with the COMPADRE team.

For the stochastic simulations we used *individual* MPMs, which represented the population dynamics
during a given annual transition and at a given population. We only used species that were represented

in the database by at least three individual MPMs (Section 8), resulting in 1578 individual MPMs from

across 108 plant species in our dataset.

2   Simulating seed harvesting

We used the selected MPMs to simulate the impact of seed harvesting on populations. We first used

the mean MPM (Section 1.4) for each species, and simulated seed harvesting as a reduction in the

values describing reproduction via seed in the sexual reproduction matrix **F** (see equation 1).

Specifically, we created a modified MPM **A'** with reduced per-capita contributions of seed production

in **F**. To carry out our projections, we initiated the population vector n_0 as the stable stage distribution

of the original MPM **A**. This vector n_0 was obtained as the right-eigenvector of **A** following methods

described by Caswell (2001). We then projected n_0 over 30 years using the modified MPM **A'** and the

chain rule (*35*). We chose this period of time for our projections because it is long enough to observe

even minor changes in the overall population size *N* that are not typically possible to quantify by short-

term monitoring (*44*), while it is of sufficient length to fit within the active career of a land manager or

conservation practitioner. We benchmarked the resulting population size $N_{30\ harvest}$ relative to the

population size $N_{30\ no\ harvest}$ that would have been achieved in the absence of seed harvesting as in

equation 2:

$$N_{30\ relative} = \frac{N_{30\ harvest}}{N_{30\ no\ harvest}} \qquad\qquad eq.\,2$$

The relative population size $N_{30\ relative}$ thus ranges between 1 (when seed harvesting has no effect on

population size; $N_{30\ harvest} = N_{30\ no\text{-}harvest}$) to 0 (when the effect is so drastic it drives *N* to 0 within 30

years). For example, a value of $N_{30\ relative}$ = 0.1 means that the population size achieved with seed

harvesting is 10% of the population size that would have been achieved without seed harvesting. The

use of this metric as measure of seed harvesting impact allowed us to implement intra- and inter-

specific comparisons, regardless of the variable population growth rates of each species' population.

When calculating the population sizes with and without harvest ($N_{30\ harvest}$ and $N_{30\ no\ harvest}$), we included

only the active but not dormant (seed bank, dormant vegetative) life stages of the population vectors

N_{30} because practitioners and scientists commonly evaluate population size based on counting active,

standing individuals.

3   Vulnerability to seed harvesting

We used mean MPMs to calculate species vulnerabilities to seed harvesting. For each species, we

created 101 MPMs that describe the population dynamics when harvesting 0-100% of seed

production, in 1% steps (Figure S1). As in section 2, we used the virtual MPMs to project population

sizes over 30 years. We then fitted an exponential-decay model to quantify the effects of the varying

proportion of harvested seed (p) on the relative population size in 30 years ($N_{30\ relative}$) as follows:

$$N_{30\ relative} = e^{p(-b)}$$ eq. 3

where b determines how steeply the relative population size ($N_{30\ relative}$) decreases with increasing

proportion of harvest pressure, such that the larger b, the steeper this decrease is. We refer to this

coefficient as *vulnerability to seed harvesting* (**Error! Reference source not found.**).

4   Testing current recommendations

Next, we used MPMs to simulate the impact of seed harvesting according to the current rules on the

relative population size $N_{30\ relative}$. As far as we are aware of, explicit recommendations for the maximal

proportion of seeds that can be harvested from natural populations so far exist only in three countries.

In USA and Australia, this value is 20% and 10%, respectively, for common plant species when

harvesting seeds for restoration projects (*13*, *14*). German rules are available for herbaceous plants:

2% for annual and 10% for perennial species when harvested every year (*15*).

As the current recommendations are partly growth-form specific (*15*), we examined the reduction in

relative population size as a function of plant growth form: annuals, herbaceous perennials,

epiphytes, lianas, palms, succulents, shrubs, and trees, as indicated in the COMPADRE metadata. We

excluded growth forms represented by less than 5 species: epiphytes (n=4) and lianas (n=1), as well

as plant species whose generation time disagreed with the metadata of the species, in particular

annual species with generation times larger than two years (n=4). As the vulnerability to seed

harvesting of individual species (Section 3) depended neither on a continent nor on the interaction

between a continent and plant growth form (Table S2), we grouped species only by growth form and

used the same set of species to test the recommendations from Australia, USA and Germany (Figure

1 in the main text).

5   Life history traits

We used life history traits to explain species vulnerability to seed harvesting. A life history trait is a

key feature that describes the life cycle of the organism (e.g. generation time, survival of seeds in the

seed bank, clonal propagation). As our ultimate motivation was to facilitate the translation of our

findings to land managers and practitioners, out of the wide range of life history traits that can be

derived from MPMs (e.g (*35*, *37*)), we selected the traits that are readily available in trait databases

or easy to estimate in the field (Table S3). All life history traits were calculated based on the matrix A
of the mean MPM of each of our 280 species.

6   The effect of life history traits on vulnerability to seed harvesting

We used linear models to determine which life history traits (generation time, degree of iteroparity,
age at sexual maturity, seed bank residence, clonality) best explained species' vulnerability to seed
harvesting (Section 3). We also added plant growth form as an explanatory variable (as defined in the
COMPADRE database (*20*)) to the model to test whether it explains any additional variability.
Restricting the model to key life history traits allowed us to keep the full model and avoid model
selection, which is known to produce exaggerated effect sizes and spurious effects (*45*). Species
vulnerability to seed harvesting was log-transformed prior analysis to achieve normality. Other
explanatory variables except plant growth type (factor) were log-transformed and standardised to
adhere to the model assumptions of normally distributed errors.

To illustrate the importance of the life history traits for predicting the species vulnerability to seed
harvesting (Figure 2 in the main text), we expressed the relative importance of each predictor in the
model as the proportion of explained variability assigned to each predictor. As the explained variability
can depend on the sequential order of the predictors in the model, we averaged the explained
variability for each predictor across all possible ordering of the predictors using the R package *relaimp*
(*46*). To visualize effect sizes of the effects of life history traits on species vulnerability to seed
harvesting, as well as uncertainty of these effects, we used 95% credible intervals, a Bayesian analogue
of confidence intervals. These were calculated from 10,000 simulations of the mean and variance of
each estimate, using the *sim* function in the R package *arm* with non-informative prior (*47*).

We also ran a model including the phylogenetic relationships among species to test the extent to which
the explanatory power of life history traits on species' vulnerability to seed harvesting is in fact driven
by the phylogenetic inertia in plant life history traits (*48*). We used a phylogenetic generalized least
square model to include the phylogeny of our species. We obtained the phylogeny from COMPADRE,
following methods detailed elsewhere (*37*). With this model, we estimated Pagel's λ (not to be
confused with the population growth rate, also referred to as λ in the demographic literature (*35*)), a
measure of phylogenetic signal in the trait structure. Briefly, Pagel's λ=0 indicates no effect of the
phylogenetic structure in the dataset in explaining variation in a given trait, while Pagel's λ=1 indicates
that the phylogenetic structure perfectly predicts, i.e. is responsible for, the life history trait structure.

Negative values suggest that closely related species have more different traits than would be expected

by chance ((*48*). We found that the phylogenetic signal was overall weak and negative (Pagel's λ=-0.1).

Based on this result, we opted to present in this paper results from the linear model without

phylogenetic correction.

7   Assessing limits of seed collection

We used the mean MPM per species to estimate what fraction of seed production one can collect from

a natural population while only moderately affecting its dynamics. As a moderate effect we defined a

reduction in population size *N* to not below 50% of the size that would have been achieved without

seed harvesting during 30 years of a constant annual harvest intensity. While a reduction of population

size by up to 50% over 30 years may seem relatively high, it corresponds to an annual decline of <2%.

This threshold also allows for the persistence of the natural population under environmental

stochasticity in >99% of species (see section 8.2).

For each species' MPM, we simulated the effect of seed harvesting as a reduction of seed production

transition by 0-100%, in 1% intervals. We used such reduced, virtual MPMs to simulate population

dynamics across 30 years, and we recorded the final population size and expressed it as relative to

population size that would be achieved without seed harvesting (see Section 2, note this calculation is the same

as the first step of the calculation of vulnerability to seed harvesting, Section 3). Besides annual harvests, we also modelled

the effect of harvesting seeds every 2, 5 or 10 years because reducing harvesting frequency up to once

in 10 years is sometimes recommended to limit negative effects of seed harvesting on population

dynamics (*11*). In this case, we modelled population dynamics with the original mean MPM while the

reduced MPM was used every 2^{nd}, 5^{th} or 10^{th} run. As the safe fraction for seed harvesting, we

considered the largest proportion of seed that was possible to harvest without exceeding the 50 %

reduction of the relative population size.

We related the safe fraction for seed harvesting to the generation time of plants – the most important

predictor of species vulnerability to seed harvesting, which alone explained 52.3% of total variability

in species vulnerability to seed harvesting. We used non-linear regression in *R* (*nsl*) to describe the

sigmoid relationship between the safe fractions of seed harvesting and the generation time, and used

function *PredFit* in package *investr* (*49*) to generate confidence intervals for the relationship (Figure 3

in the main text).

8 Effect of environmental stochasticity

In a subset of our studied species, we simulated the effects of environmental stochasticity on population dynamics to understand how environmental stochasticity affects our prediction for seed harvesting based on mean MPMs. We used all species in our dataset represented by at least three individual MPMs (Section 1.4), resulting in 1,578 MPMs across 108 plant species. We simulated environmental stochasticity as projecting vector of stable stage distribution of the mean MPM by randomly drawn individual MPM in each step. To obtain a probability distribution of results under environmental stochasticity, we repeated this process 1,000 times. We expressed the results as $N_{30\ relative}$ (equation eq. 2). The effects of seed harvesting were simulated as above (Section 4), with the difference that in each of the 30 annual time-steps in each of the 1,000 simulation runs, we randomly drew an individual MPM from the set of individual MPMs available for a given species.

8.1 The effect of seed harvesting on population size based on environmental stochasticity versus mean MPM

To understand how environmental stochasticity affected our results, we estimated the robustness of our results in stochastic environments. As an example, we used the effect of harvest of 20% of seed production, expressed as $N_{30\ relative}$, and simulated seed harvesting either using mean MPMs or stochastic simulation. We then compared the safe fraction for seed harvesting ($N_{30\ relative} > 0.5$) based on the mean MPMs to the median of safe seed fraction based on the stochastic simulations.

The median of relative population sizes $N_{30\ relative}$ based on 1,000 permutations of stochastic simulations (y axis in Figure S3) closely correlated with the $N_{30\ relative}$ based on the mean MPMs. Interestingly, the relative population size $N_{30\ relative}$ based on stochastic simulation (orange points in Figure S3) was slightly higher than the $N_{30\ relative}$ based on mean MPMs (black line in Figure S3), especially in species that are more vulnerable to seed harvesting. Consequently, the safe fraction for seed harvesting based on the median of stochastic simulations was on average 0.017 higher that safe fraction based on the mean MPMs (Figure S4). This suggests that environmental stochasticity partly buffers the predicted decrease of population size caused by seed harvesting.

8.2 Threshold for seed harvesting based on mean MPM versus extinction probability

In the models above, we set a threshold for seed harvesting so that the relative population size $N_{30\ relative}$ decreased not below 50% of the population size that would be achieved without seed harvesting. In this section, we tested whether this threshold also prevented populations from extinctions. For each species, we computed what fraction of seeds could be sustainably harvested without causing

extinction in at least 95% of stochastic simulations. We considered a population to go locally extinct

when $N_{30\ relative}$ < 0.01 (see Section 2 for definition of $N_{30\ relative}$). For each species, we compared the

threshold based on the 95% probability of population survival with the threshold based on mean MPM

and $N_{30\ relative}$ > 0.5.

In the vast majority (>99%) of examined species, the threshold based on $N_{30\ relative}$ > 0.5 (as calculated

using mean MPMs, black line in the Figure S5) allowed for the collection of a lower proportion of seeds

than the threshold based on 95% probability population survival when using stochastic simulations

(individual points in Figure 5). This suggests that the rules based on $N_{30\ relative}$ > 0.5 derived from the

mean MPMs prevent populations from going locally extinct.

Fig. S1.

Figure S1: Life cycle of a hypothetical plant species with three stages (seedbank, juvenile, and adult) and its corresponding matrix population model (MPM), with λ indicating its long-term population growth rate, which is a function of population size (N) between two time-points t and t+1. Seed harvesting in this study was simulated by manipulating the transitions that describe generative reproduction.

Fig. S2.

Figure S2. Vulnerability of population dynamics to seed harvesting (*b* in equation S3) in two

of our 280 examined plant species. Note how the larger the value of *b*, the more vulnerable

the given species is to seed harvesting. Black dots: simulated values; red line: fitted

exponential-decay model as per equation 3.

Fig. S3.

Figure S3: Comparison of relative population sizes ($N_{30\ relative}$) when 20% seeds were harvested

based on mean MPMs (x axis and the 1:1 black line) versus from calculations with

environmental stochasticity (y axis).

Fig. S4.

Figure S4: The safe fractions for seed harvesting based on $N_{30\,relative}$ > 0.5 as calculated from

the mean MPM (x-axis and 1:1 black line) versus the same safe fraction based on stochastic

simulations (with 95% CI).

Fig. S5.

Figure S5: Comparison of the threshold for maximal seed harvest based on $N_{30\,relative} > 0.5$ as calculated

from the mean MPM (x-axis and the 1:1 black line), with the maximal seed harvest that allows 95%

probability of population survival of each considered species, as based on stochastic simulation.

Table S1.

696 Table S1: The final set of species used in this study and the original study that was the source of the

697 MPMs.

Species	Publication
Abies concolor	Van Mantgem, P. J., & Stephenson, N. L. (2005). The accuracy of matrix population model projections for coniferous trees in the Sierra Nevada, California. Journal of Ecology, 93(4), 737–747. Portico. https://doi.org/10.1111/j.1365-2745.2005.01007.x
Abies magnifica	Van Mantgem, P. J., & Stephenson, N. L. (2005). The accuracy of matrix population model projections for coniferous trees in the Sierra Nevada, California. Journal of Ecology, 93(4), 737–747. Portico. https://doi.org/10.1111/j.1365-2745.2005.01007.x
Abies sachalinensis	Hiura, T., & Fujiwara, K. (1999). Density-dependence and co-existence of conifer and broad-leaved trees in a Japanese northern mixed forest. Journal of Vegetation Science, 10(6), 843–850. Portico. https://doi.org/10.2307/3237309
Acacia bilimekii	Jiménez-Lobato, V., & Valverde, T. (2006). Population dynamics of the shrub *Acacia bilimekii* in a semi-desert region in central Mexico. Journal of Arid Environments, 65(1), 29–45. https://doi.org/10.1016/j.jaridenv.2005.07.002
Acacia suaveolens	Warton, D. I., & Wardle, G. M. (2003). Site-to-site variation in the demography of a fire-affected perennial, *Acacia suaveolens*, at Ku-ring-gai Chase National Park, New South Wales, Australia. Austral Ecology, 28(1), 38–47. https://doi.org/10.1046/j.1442-9993.2003.01246.x
Acer amoenum	Tanaka, H., Shibata, M., Masaki, T., Iida, S., Niiyama, K., Abe, S., Kominami, Y., & Nakashizuka, T. (2008). Comparative demography of three coexisting *Acer* species in gaps and under closed canopy. Journal of Vegetation Science, 19(1), 127–138. Portico. https://doi.org/10.3170/2007-8-18342

Species	Publication
Acer mono	Tanaka, H., Shibata, M., Masaki, T., Iida, S., Niiyama, K., Abe, S., Kominami, Y., & Nakashizuka, T. (2008). Comparative demography of three coexisting *Acer* species in gaps and under closed canopy. Journal of Vegetation Science, 19(1), 127–138. Portico. https://doi.org/10.3170/2007-8-18342
Acer rufinerve	Tanaka, H., Shibata, M., Masaki, T., Iida, S., Niiyama, K., Abe, S., Kominami, Y., & Nakashizuka, T. (2008). Comparative demography of three coexisting *Acer* species in gaps and under closed canopy. Journal of Vegetation Science, 19(1), 127–138. Portico. https://doi.org/10.3170/2007-8-18342
Acer saccharum	Lin, Y., & Augspurger, C. K. (2008). Impact of spatial heterogeneity of neighborhoods on long-term population dynamics of sugar maple (*Acer saccharum*). Forest Ecology and Management, 255(10), 3589–3596. https://doi.org/10.1016/j.foreco.2008.02.040
Actaea spicata	Fröborg, H., & Eriksson, O. (2003). Predispersal seed predation and population dynamics in the perennial understorey herb *Actaea spicata*. Canadian Journal of Botany, 81(11), 1058–1069. https://doi.org/10.1139/b03-099
Adesmia volckmannii	Cipriotti, P. A., & Aguiar, M. R. (2011). Direct and indirect effects of grazing constrain shrub encroachment in semi-arid Patagonian steppes. Applied Vegetation Science, 15(1), 35–47. https://doi.org/10.1111/j.1654-109x.2011.01138.x
Aeschynomene virginica	Griffith, A. B., & Forseth, I. N. (2005). Population matrix models of *Aeschynomene virginica*, a rare annual plant: implications for conservation. Ecological Applications, 15(1), 222–233. https://doi.org/10.1890/02-5219
Aesculus turbinata	Kaneko, Y., Takada, T., & Kawano, S. (1999). Population biology of *Aesculus turbinata* Blume: A demographic analysis using transition matrices on a natural population along a riparian environmental gradient. Plant Species Biology, 14(1), 47–68. https://doi.org/10.1046/j.1442-1984.1999.00007.x
Agrimonia eupatoria	Mondragón Chaparro, D., & Ticktin, T. (2011). Demographic effects of harvesting epiphytic bromeliads and an alternative approach to collection.

Species	Publication
	Conservation Biology, 25(4), 797–807. https://doi.org/10.1111/j.1523-1739.2011.01691.x
Ailanthus altissima	Bullock, J. M., White, S. M., Prudhomme, C., Tansey, C., Perea, R., & Hooftman, D. A. P. (2011). Modelling spread of British wind-dispersed plants under future wind speeds in a changing climate. Journal of Ecology, 100(1), 104–115. https://doi.org/10.1111/j.1365-2745.2011.01910.x
Alliaria petiolata	Evans, J. A., Davis, A. S., Raghu, S., Ragavendran, A., Landis, D. A., & Schemske, D. W. (2012). The importance of space, time, and stochasticity to the demography and management of *Alliaria petiolata*. Ecological Applications, 22(5), 1497–1511. https://doi.org/10.1890/11-1291.1
Allium tricoccum	Nault, A., & Gagnon, D. (1993). Ramet demography of *Allium tricoccum*, a spring ephemeral, perennial forest herb. The Journal of Ecology, 81(1), 101. https://doi.org/10.2307/2261228
Alyxia stellata	Wong, T. M., & Ticktin, T. (2014). Using population dynamics modelling to evaluate potential success of restoration: a case study of a Hawaiian vine in a changing climate. Environmental Conservation, 42(1), 20–30. https://doi.org/10.1017/s0376892914000204
Andropogon brevifolius	Canales, J., Trevisan, M.C., Silva, J.F. & Caswell,H. (1994): A demographic-study of an annual grass (*Andropogon brevifolius* Schwarz) in burnt and unburnt savanna. Acta Oecologica 15(3): 261-273
Androsace elongata	Dostál, P. (2007). Population dynamics of annuals in perennial grassland controlled by ants and environmental stochasticity. Journal of Vegetation Science, 18(1), 91–102. Portico. https://doi.org/10.1111/j.1654-1103.2007.tb02519.x
Anemone patens	Williams, J. L., & Crone, E. E. (2006). The impact of invasive grasses on the population growth of *Anemone patens*, a long-lived native forb. Ecology, 87(12), 3200–3208. https://doi.org/10.1890/0012-9658(2006)87[3200:tioigo]2.0.co;2

Species	Publication
Anthericum ramosum	Černá, L., & Münzbergová, Z. (2013). Comparative population dynamics of two closely related species differing in ploidy level. PLoS ONE, 8(10), e75563. https://doi.org/10.1371/journal.pone.0075563
Anthyllis vulneraria	Davison, R., Jacquemyn, H., Adriaens, D., Honnay, O., de Kroon, H., & Tuljapurkar, S. (2010). Demographic effects of extreme weather events on a short-lived calcareous grassland species: stochastic life table response experiments. Journal of Ecology, 98(2), 255–267. https://doi.org/10.1111/j.1365-2745.2009.01611.x
Aquilaria crassna	Zhang, L., Brockelman, W. Y., & Allen, M. A. (2008). Matrix analysis to evaluate sustainability: The tropical tree *Aquilaria crassna*, a heavily poached source of agarwood. Biological Conservation, 141(6), 1676–1686. https://doi.org/10.1016/j.biocon.2008.04.015
Aquilaria malaccensis	Soehartono, T., & C. Newton, A. (2001). Conservation and sustainable use of tropical trees in the genus *Aquilaria* II. The impact of gaharu harvesting in Indonesia. Biological Conservation, 97(1), 29–41. https://doi.org/10.1016/s0006-3207(00)00089-6
Aquilaria microcarpa	Soehartono, T., & C. Newton, A. (2001). Conservation and sustainable use of tropical trees in the genus *Aquilaria* II. The impact of gaharu harvesting in Indonesia. Biological Conservation, 97(1), 29–41. https://doi.org/10.1016/s0006-3207(00)00089-6
Aquilegia chrysantha	Stubben, C.J. (2007). Projecting the response of yellow columbine populations to climate change. PhD thesis, New Mexico State University, Las Cruces, New Mexico.
Aquilegia sp*.*	Stubben, C. & Milligan, B. (2007): Estimating and analyzing demographic models using the popbio package in R. Journal of Statistics Software 22(11): 1-23
Araucaria araucana	Bekessy, S., Newton, A., Fox, J., Lara, A., Premoli, A., Cortes, M., Gonzalez, M., & Burgman, M. (2004). Monkey puzzle tree (*Araucaria araucana*) in southern Chile: Effects of timber and seed harvest, volcanic activity, and

Species	Publication
	fire. Species Conservation and Management: Case Studies (pp. 48-63). RMIT University.
Araucaria cunninghamii	Enright, N., & Ogden, J. (1979). Applications of transition matrix models in forest dynamics: *Araucaria* in Papua New Guinea and Nothofagus in New Zealand. Austral Ecology, 4(1), 3–23. https://doi.org/10.1111/j.1442-9993.1979.tb01195.x
Araucaria hunsteinii	Enright, N. J. (1982). Does *Araucaria hunsteinii* compete with its neighbours? Austral Ecology, 7(1), 97–99. https://doi.org/10.1111/j.1442-9993.1982.tb01304.x
Araucaria muelleri	Enright, N. J., Miller, B. P., Perry, G. L. W., Goldblum, D., & Jaffré, T. (2013). Stress-tolerator leaf traits determine population dynamics in the endangered New Caledonian conifer *Araucaria muelleri*. Austral Ecology, 39(1), 60–71. https://doi.org/10.1111/aec.12045
Arenaria bolosii	Iriondo, J.M., Albert M. J., Gimenez-Benavides, L., Dominguez-Lozano, F. & Escudero, A. [Eds.] (2009): Poblaciones en peligro: viabilidad demográfica de la flora vascular amenazada de España. Dirección General de Medio Natural y Política Forestal (Ministerio de Medio Ambiente, y Medio Rural y Marino), Madrid, 242 pp.
Arenaria serpyllifolia	Dostál, P. (2007). Population dynamics of annuals in perennial grassland controlled by ants and environmental stochasticity. Journal of Vegetation Science, 18(1), 91–102. Portico. https://doi.org/10.1111/j.1654-1103.2007.tb02519.x
Argyroxiphium sandwicense subsp. *macrocephalum*	Forsyth, S. A. (2003). Density-dependent seed set in the *Haleakala silversword*: evidence for an Allee effect. Oecologia, 136(4), 551–557. https://doi.org/10.1007/s00442-003-1295-3
Armeria maritima	Lefebvre, C., & Chandler-Mortimer, A. (1984). Demographic characteristics of the perennial herb *Armeria maritima* on zinc lead mine wastes. The Journal of Applied Ecology, 21(1), 255. https://doi.org/10.2307/2403051

Species	Publication
Armeria merinoi	Iriondo, J.M., Albert M. J., Gimenez-Benavides, L., Dominguez-Lozano, F. & Escudero, A. [Eds.] (2009): Poblaciones en peligro: viabilidad demográfica de la flora vascular amenazada de España. Dirección General de Medio Natural y Política Forestal (Ministerio de Medio Ambiente, y Medio Rural y Marino), Madrid, 242 pp.
Artemisia genipi	Marcante, S., Winkler, E., & Erschbamer, B. (2009). Population dynamics along a primary succession gradient: do alpine species fit into demographic succession theory? Annals of Botany, 103(7), 1129–1143. https://doi.org/10.1093/aob/mcp047
Asarum canadense	Damman, H., & Cain, M. L. (1998). Population growth and viability analyses of the clonal woodland herb, *Asarum canadense*. Journal of Ecology, 86(1), 13–26. https://doi.org/10.1046/j.1365-2745.1998.00242.x
Astragalus alopecurus	Nicole F. (2005): Biologie de la conservation appliquée aux plantes menacées des Alpes. PhD thesis, University of Grenoble, Grenoble, France.
Astragalus australis var. *olympicus*	Kaye, T. (1990). Autecology, reproductive ecology, and demography of *Astragalus australis* var. *olympicus* (Fabaceae). MSc. Thesis, Oregon State University.
Astragalus peckii	Martin, E. F., & Meinke, R. J. (2012). Variation in the demographics of a rare central Oregon endemic, *Astragalus peckii* Piper (Fabaceae), with fluctuating levels of herbivory. Population Ecology, 54(3), 381–390. Portico. https://doi.org/10.1007/s10144-012-0318-5
Astragalus scaphoides	Lesica, P (1995): Demography of *Astragalus scaphoides* and effects of herbivory on population-growth. Great Basin Naturalis 55(2): 142-150
Astragalus tyghensis	Kaye, T. N., & Pyke, D. A. (2003). The effect of stochastic technique on estimates of population viability from transition matrix models. Ecology, 84(6), 1464–1476. https://doi.org/10.1890/0012-9658(2003)084[1464:teosto]2.0.co;2

Species	Publication
Astrophytum asterias	Martínez-Ávalos J.G. (2007): Estudio demográfico del "star cactus" *Astrophytum asterias* (Lem.) Zucc. (Cactaceae) una especie en riesgo de extinción. PhD Thesis, Universidad Autónoma de Nuevo León, México.
Astrophytum capricorne	Mandujano, M. C., Bravo, Y., Verhulst, J., Carrillo-Angeles, I., & Golubov, J. (2015). The population dynamics of an endemic collectible cactus. Acta Oecologica, 63, 1–7. https://doi.org/10.1016/j.actao.2014.12.004
Astrophytum ornatum	Zepeda-Martínez, V., Mandujano, M. C., Mandujano, F. J., & Golubov, J. K. (2013). What can the demography of *Astrophytum ornatum* tell us of its endangered status? Journal of Arid Environments, 88, 244–249. https://doi.org/10.1016/j.jaridenv.2012.08.006
Avicennia germinans	López-Hoffman, L., Ackerly, D. D., Anten, N. P. R., Denoyer, J. L., & Martinez-Ramos, M. (2007). Gap-dependence in mangrove life-history strategies: a consideration of the entire life cycle and patch dynamics. Journal of Ecology, 95(6), 1222–1233. https://doi.org/10.1111/j.1365-2745.2007.01298.x
Banksia ericifolia	Bradstock, R. A., & O'Connell, M. A. (1988). Demography of woody plants in relation to fire: *Banksia ericifolia* L.f. and *Petrophile pulchella* (Schrad) R.Br. Austral Ecology, 13(4), 505–518. https://doi.org/10.1111/j.1442-9993.1988.tb00999.x
Borassus aethiopum	Barot, S., Gignoux, J., Vuattoux, R., & Legendre, S. (2000). Demography of a savanna palm tree in Ivory Coast (Lamto): population persistence and life-history. Journal of Tropical Ecology, 16(5), 637–655. https://doi.org/10.1017/s0266467400001620
Borderea chouardii	García, M. B., Guzmán, D., & Goñi, D. (2002). An evaluation of the status of five threatened plant species in the Pyrenees. Biological Conservation, 103(2), 151–161. https://doi.org/10.1016/s0006-3207(01)00113-6
Boswellia papyrifera	Groenendijk, P., Eshete, A., Sterck, F. J., Zuidema, P. A., & Bongers, F. (2011). Limitations to sustainable frankincense production: blocked regeneration, high adult mortality and declining populations. Journal of

Species	Publication
	Applied Ecology, 49(1), 164–173. Portico. https://doi.org/10.1111/j.1365-2664.2011.02078.x
Brassica insularis	Noël, F., Maurice, S., Mignot, A., Glémin, S., Carbonell, D., Justy, F., Guyot, I., Olivieri, I., & Petit, C. (2010). Interaction of climate, demography and genetics: a ten-year study of *Brassica insularis*, a narrow endemic Mediterranean species. Conservation Genetics, 11(2), 509–526. https://doi.org/10.1007/s10592-010-0056-1
Brassica napus	Garnier, A., & Lecomte, J. (2006). Using a spatial and stage-structured invasion model to assess the spread of feral populations of transgenic oilseed rape. Ecological Modelling, 194(1–3), 141–149. https://doi.org/10.1016/j.ecolmodel.2005.10.009
Braya fernaldii	Squires, S. (2010): Insect pests and pathogens compromise the persistence of two endemic and rare *Braya* (Brassicaceae). PhD Dissertation, memorial University of Newfoundland, Newfoundland and labrador, 184 pp.
Braya longii	Squires, S. (2010): Insect pests and pathogens compromise the persistence of two endemic and rare *Braya* (Brassicaceae). PhD Dissertation, memorial University of Newfoundland, Newfoundland and Labrador, 184 pp.
Bursera glabrifolia	Hernández-Apolinar, M., Valverde, T., & Purata, S. (2006). Demography of *Bursera glabrifolia*, a tropical tree used for folk woodcrafting in Southern Mexico: An evaluation of its management plan. Forest Ecology and Management, 223(1–3), 139–151. https://doi.org/10.1016/j.foreco.2005.10.072
Calathea ovandensis	Horvitz, C. C., & Schemske, D. W. (1995). Spatiotemporal variation in demographic transitions of a tropical understory herb: projection matrix analysis. Ecological Monographs, 65(2), 155–192. Portico. https://doi.org/10.2307/2937136
Callitris intratropica	Price, O., & Bowman, D. M. J. S. (1994). Fire-stick forestry: a matrix model in support of skillful fire management of *Callitris intratropica* R. T. Baker by

Species	Publication
	north Australian Aborigenes. Journal of Biogeography, 21(6), 573. https://doi.org/10.2307/2846032
Calluna vulgaris	Scandrett, E., & Gimingham, C. H. (1989). A model of *Calluna* population dynamics; the effects of varying seed and vegetative regeneration. Vegetatio, 84(2), 143–152. https://doi.org/10.1007/bf00036515
Calocedrus decurrens	Van Mantgem, P. J., & Stephenson, N. L. (2005). The accuracy of matrix population model projections for coniferous trees in the Sierra Nevada, California. Journal of Ecology, 93(4), 737–747. Portico. https://doi.org/10.1111/j.1365-2745.2005.01007.x
Calochortus albus	Fiedler, P. L. (1987). Life history and population dynamics of rare and common mariposa lilies (*Calochortus pursh*: Liliaceae). The Journal of Ecology, 75(4), 977. https://doi.org/10.2307/2260308
Calochortus obispoensis	Fiedler, P. L. (1987). Life history and population dynamics of rare and common mariposa lilies (*Calochortus pursh*: Liliaceae). The Journal of Ecology, 75(4), 977. https://doi.org/10.2307/2260308
Calochortus pulchellus	Fiedler, P. L. (1987). Life history and population dynamics of rare and common mariposa lilies (*Calochortus pursh*: Liliaceae). The Journal of Ecology, 75(4), 977. https://doi.org/10.2307/2260308
Calochortus tiburonensis	Fiedler, P. L. (1987). Life history and population dynamics of rare and common mariposa lilies (*Calochortus pursh*: Liliaceae). The Journal of Ecology, 75(4), 977. https://doi.org/10.2307/2260308
Carduus nutans	Shea, K., Kelly, D., Sheppard, A. W., & Woodburn, T. L. (2005). Context-dependent biological control of an invasive thistle. Ecology, 86(12), 3174–3181. https://doi.org/10.1890/05-0195
Carlina vulgaris	Jongejans, E., Jorritsma-Wienk, L. D., Becker, U., Dostál, P., Mildén, M., & de Kroon, H. (2010). Region versus site variation in the population dynamics of three short-lived perennials. Journal of Ecology, 98(2), 279–289. https://doi.org/10.1111/j.1365-2745.2009.01612.x

Species	Publication
Carnegiea gigantea	Steenbergh, W. F., & Lowe, C. H. (1969). Critical factors during the first years of life of the saguaro (*Cereus giganteus*) at Saguaro National Monument, Arizona. Ecology, 50(5), 825–834. Portico. https://doi.org/10.2307/1933696
Cassia nemophila	Silander, J. A. (1983). Demographic variation in the Australian desert cassia under grazing pressure. Oecologia, 60(2), 227–233. https://doi.org/10.1007/bf00379524
Castanea dentata	Davelos, A. L., & Jarosz, A. M. (2004). Demography of American chestnut populations: effects of a pathogen and a hyperparasite. Journal of Ecology, 92(4), 675–685. Portico. https://doi.org/10.1111/j.0022-0477.2004.00907.x
Cecropia obtusifolia	Alvarez-Buylla, E. R. (1994). Density dependence and patch dynamics in tropical rain forests: matrix models and applications to a tree species. The American Naturalist, 143(1), 155–191. https://doi.org/10.1086/285599
Centaurea maculosa	Emery, S. M., & Gross, K. L. (2005). Effects of timing of prescribed fire on the demography of an invasive plant, spotted knapweed *Centaurea maculosa*. Journal of Applied Ecology, 42(1), 60–69. https://doi.org/10.1111/j.1365-2664.2004.00990.x
Cephalanthera longifolia	Shefferson, R. P., Kull, T., Tali, K., & Kellett, K. M. (2012). Linking vegetative dormancy to fitness in two long-lived herbaceous perennials. Ecosphere, 3(2), art13. https://doi.org/10.1890/es11-00328.1
Chaerophyllum aureum	Magda, D., Duru, M., & Theau, J.-P. (2004). Defining management rules for grasslands using weed demographic characteristics. Weed Science, 52(3), 339–345. https://doi.org/10.1614/p2202-067
Chamaecrista keyensis	Liu, H., Menges, E. S., & Quintana-Ascencio, P. F. (2005). Population viability analyses of *Chamaecrista keyensis*: Effects of fire season and frequency. Ecological Applications, 15(1), 210–221. https://doi.org/10.1890/03-5382
Chamaedorea elegans	Valverde, T., Hernandez-Apolinar, M., & Mendoza-Amarom, S. (2006). Effect of leaf harvesting on the demography of the tropical palm

Species	Publication
	Chamaedorea elegans in South-Eastern Mexico. Journal of Sustainable Forestry, 23(1), 85–105. https://doi.org/10.1300/j091v23n01_05
Chamaedorea radicalis	Endress, B. A., Gorchov, D. L., & Noble, R. B. (2004).Non-timber forest product extraction: effects of harvest and browsing on an understory palm. Ecological Applications, 14(4), 1139–1153. https://doi.org/10.1890/02-5365
Choerospondias axillaris	Brodie, J. F., Helmy, O. E., Brockelman, W. Y., & Maron, J. L. (2009). Functional differences within a guild of tropical mammalian frugivores. Ecology, 90(3), 688–698. Portico. https://doi.org/10.1890/08-0111.1
Cimicifuga elata	Kaye, T. N., & Pyke, D. A. (2003). The effect of stochastic technique on estimates of population viability from transition matrix models. Ecology, 84(6), 1464–1476. https://doi.org/10.1890/0012-9658(2003)084[1464:teosto]2.0.co;2
Cirsium acaule	Bullock, J. M., White, S. M., Prudhomme, C., Tansey, C., Perea, R., & Hooftman, D. A. P. (2011). Modelling spread of British wind-dispersed plants under future wind speeds in a changing climate. Journal of Ecology, 100(1), 104–115. https://doi.org/10.1111/j.1365-2745.2011.01910.x
Cirsium dissectum	Jongejans, E., de Vere, N., & de Kroon, H. (2008). Demographic vulnerability of the clonal and endangered meadow thistle. Plant Ecology, 198(2), 225–240. https://doi.org/10.1007/s11258-008-9397-y
Cirsium palustre	Ramula, S. (2008). Population dynamics of a monocarpic thistle: simulated effects of reproductive timing and grazing of flowering plants. Acta Oecologica, 33(2), 231–239. https://doi.org/10.1016/j.actao.2007.11.005
Cirsium perplexans	Dodge, G.J. 2005. Ecological effects of the biocontrol insects, *Larinus planus* and *Rhinocyllus conicus*, on native thistles. PhD Thesis, University of Maryland, Maryland
Cirsium scariosum	Dodge, G.J. 2005. Ecological effects of the biocontrol insects, *Larinus planus* and *Rhinocyllus conicus*, on native thistles. PhD Thesis, University of Maryland, Maryland

Species	Publication
Cirsium undulatum var. *tracyi*	Dodge, G.J. 2005. Ecological effects of the biocontrol insects, *Larinus planus* and *Rhinocyllus conicus*, on native thistles. PhD Thesis, University of Maryland, Maryland
Cirsium vulgare	Bullock, J. M., Hill, B. C., & Silvertown, J. (1994). Demography of *Cirsium vulgare* in a grazing experiment. The Journal of Ecology, 82(1), 101. https://doi.org/10.2307/2261390
Cochlearia bavarica	Abs, C. (1999). Differences in the life histories of two *Cochlearia species*. Folia Geobotanica, 34(1), 33–45. https://doi.org/10.1007/bf02803075
Cochlearia pyrenaica	Abs, C. (1999). Differences in the life histories of two *Cochlearia species*. Folia Geobotanica, 34(1), 33–45. https://doi.org/10.1007/bf02803075
Colchicum autumnale	Winter, S., Jung, L. S., Eckstein, R. L., Otte, A., Donath, T. W., & Kriechbaum, M. (2014). Control of the toxic plant *Colchicum autumnale* in semi-natural grasslands: effects of cutting treatments on demography and diversity. Journal of Applied Ecology, 51(2), 524–533. https://doi.org/10.1111/1365-2664.12217
Collinsia verna	Kalisz, S., & McPeek, M. A. (1992). Demography of an age-structured annual: resampled projection matrices, elasticity analyses, and seed bank effects. Ecology, 73(3), 1082–1093. Portico. https://doi.org/10.2307/1940182
Conyza canadensis	Bullock, J. M., White, S. M., Prudhomme, C., Tansey, C., Perea, R., & Hooftman, D. A. P. (2011). Modelling spread of British wind-dispersed plants under future wind speeds in a changing climate. Journal of Ecology, 100(1), 104–115. https://doi.org/10.1111/j.1365-2745.2011.01910.x
Coryphantha robbinsorum	Schmalzel, R.J., Reichenbacher F.W. & Rutman S. (1995). Demographic study of the rare *Coryphantha robbinsorum* (Cactaceae) in southeastern Arizona. Madroño 42: 332-348
Cynoglossum officinale	Boorman, L. A., & Fuller, R. M. (1984). The comparative ecology of two sand dune biennials: *Lactuca virosa* L. and *Cynoglossum officinale* L. New

Species	Publication
	Phytologist, 96(4), 609–629. https://doi.org/10.1111/j.1469-8137.1984.tb03596.x
Cypripedium calceolus	Nicolè, F., Brzosko, E., & Till-Bottraud, I. (2005). Population viability analysis of *Cypripedium calceolus* in a protected area: longevity, stability and persistence. Journal of Ecology, 93(4), 716–726. Portico. https://doi.org/10.1111/j.1365-2745.2005.01010.x
Cypripedium parviflorum var. *parviflorum*	Shefferson, R. P., Warren, R. J., & Pulliam, H. R. (2014). Life-history costs make perfect sprouting maladaptive in two herbaceous perennials. Journal of Ecology, 102(5), 1318–1328. https://doi.org/10.1111/1365-2745.12281
Cytisus scoparius	da Silveira Pontes, L., Magda, D., Jarry, M., Gleizes, B., & Agreil, C. (2012). Shrub encroachment control by browsing: Targeting the right demographic process. Acta Oecologica, 45, 25–30. https://doi.org/10.1016/j.actao.2012.08.006
Dactylorhiza lapponica	Sletvold, N., Øien, D.-I., & Moen, A. (2010). Long-term influence of mowing on population dynamics in the rare orchid *Dactylorhiza lapponica*: The importance of recruitment and seed production. Biological Conservation, 143(3), 747–755. https://doi.org/10.1016/j.biocon.2009.12.017
Daphne rodriguezii	Rodríguez-Ortega C. (2008). Consecuencias demográficas y evolutivas del secuestro de semillas en tres especies del género *Mammillaria* (Cactaceae). PhD Dissertation, Universidad Autónoma Metropolitana, Mexico.
Daucus carota	Verkaar, H. J., & Schenkeveld, A. J. (1984). On the ecology of short-lived forbs in chalk grasslands: life-history characteristics. New Phytologist, 98(4), 659–672. https://doi.org/10.1111/j.1469-8137.1984.tb04155.x
Dicerandra frutescens	Menges, E. S., Quintana Ascencio, P. F., Weekley, C. W., & Gaoue, O. G. (2006). Population viability analysis and fire return intervals for an endemic Florida scrub mint. Biological Conservation, 127(1), 115–127. https://doi.org/10.1016/j.biocon.2005.08.002

Species	Publication
Digitalis purpurea	van Baalen, J., & Prins, E. G. M. (1983). Growth and reproduction of *Digitalis purpurea* in different stages of succession. Oecologia, 58(1), 84–91. https://doi.org/10.1007/bf00384546
Dioon merolae	Lázaro-Zermeño, J. M., González-Espinosa, M., Mendoza, A., Martínez-Ramos, M., & Quintana-Ascencio, P. F. (2011). Individual growth, reproduction and population dynamics of *Dioon merolae* (Zamiaceae) under different leaf harvest histories in Central Chiapas, Mexico. Forest Ecology and Management, 261(3), 427–439. https://doi.org/10.1016/j.foreco.2010.10.028
Dipsacus sylvestris	Werner, P. A., & Caswell, H. (1977). Population rowth rates and age versus stage-distribution models for teasel (*Dipsacus sylvestris* Huds.). Ecology, 58(5), 1103–1111. Portico. https://doi.org/10.2307/1936930
Draba asterophora	Putnam, E. R. S. (2013): Ecology, phylogenetics, and conservation of *Draba asterophora* complex: a rare, alpine, endemic from Lake Tahoe, USA. PhD Thesis, Birgham Young University, Utah.
Dracocephalum austriacum	Dostálek, T., & Münzbergová, Z. (2012).Comparative population biology of critically endangered *Dracocephalum austriacum* (Lamiaceae) in two distant regions. Folia Geobotanica, 48(1), 75–93. https://doi.org/10.1007/s12224-012-9132-2
Echeveria longissima	Martorell, C., Garcillán, P. P., & Casillas, F. (2012). Ruderality in extreme-desert cacti? Population effects of chronic anthropogenic disturbance on *Echinocereus lindsayi*. Population Ecology, 54(2), 335–346. Portico. https://doi.org/10.1007/s10144-012-0307-8
Echinocactus platyacanthus	Jiménez-Sierra, C., Mandujano, M. C., & Eguiarte, L. E. (2007). Are populations of the candy barrel cactus (*Echinocactus platyacanthus*) in the desert of Tehuacán, Mexico at risk? Population projection matrix and life table response analysis. Biological Conservation, 135(2), 278–292. https://doi.org/10.1016/j.biocon.2006.10.038

Species	Publication
Echinospartum algibicum	Iriondo, J.M., Albert M. J., Gimenez-Benavides, L., Dominguez-Lozano, F. & Escudero, A. [Eds.] (2009): Poblaciones en peligro: viabilidad demográfica de la flora vascular amenazada de España. Dirección General de Medio Natural y Política Forestal (Ministerio de Medio Ambiente, y Medio Rural y Marino), Madrid, 242 pp.
Echium vulgare	Klemow, K. M., & Raynal, D. J. (1985). Demography of two facultative biennial plant species in an unproductive habitat. The Journal of Ecology, 73(1), 147. https://doi.org/10.2307/2259775
Encephalartos cycadifolius	Raimondo, D. C., & Donaldson, J. S. (2003). Responses of cycads with different life histories to the impact of plant collecting: simulation models to determine important life history stages and population recovery times. Biological Conservation, 111(3), 345–358. https://doi.org/10.1016/s0006-3207(02)00303-8
Epipactis atrorubens	Hens, H., Pakanen, V.-M., Jäkäläniemi, A., Tuomi, J., & Kvist, L. (2017). Low population viability in small endangered orchid populations: Genetic variation, seedling recruitment and stochasticity. Biological Conservation, 210, 174–183. https://doi.org/10.1016/j.biocon.2017.04.019
Eriogonum longifolium var. *gnaphalifolium*	Satterthwaite, W. H., Menges, E. S., & Quintana-Ascencio, P. F. (2002). Assessing scrub buckwheat population viability in relation to fire using multiple modeling techniques. Ecological Applications, 12(6), 1672–1687. https://doi.org/10.1890/1051-0761(2002)012[1672:asbpvi]2.0.co;2
Eryngium alpinum	Andrello, M., Bizoux, J.-P., Barbet-Massin, M., Gaudeul, M., Nicolè, F., & Till-Bottraud, I. (2012). Effects of management regimes and extreme climatic events on plant population viability in *Eryngium alpinum*. Biological Conservation, 147(1), 99–106. https://doi.org/10.1016/j.biocon.2011.12.012
Eryngium cuneifolium	Menges, E. S., & Quintana-Ascencio, P. F. (2004). Population viability with fire in *Eryngium cuneifolium*: Deciphering a decade of demographic data. Ecological Monographs, 74(1), 79–99. https://doi.org/10.1890/03-4029

Species	Publication
Erythronium japonicum	Kawano S., Takada T., Nakayama S., Hiratsuka A. (1987) Demographic differentiation and life-history evolution in temperate woodland plants. In: Urbanska K. M. (ed.). Differentiation Patterns in Higher Plants. Academic Press, Harcourt Brace Jovanovich Publishers, London, pp. 153– 181.
Escontria chiotilla	Ortega, B. (2001). Demografía de la cactácea columnar *Escontria chiotilla*. MSc. Thesis, Universidad Autónoma Metropolitana, Mexico.
Espeletia spicata	Silva, J. F., Trevisan, M. C., Estrada, C. A., & Monasterio, M. (2000). Comparative demography of two giant caulescent rosettes (*Espeletia timotensis* and *E. spicata*) from the high tropical Andes. Global Ecology and Biogeography, 9(5), 403–413. Portico. https://doi.org/10.1046/j.1365-2699.2000.00187.x
Espeletia timotensis	Silva, J. F., Trevisan, M. C., Estrada, C. A., & Monasterio, M. (2000). Comparative demography of two giant caulescent rosettes (*Espeletia timotensis* and *E. spicata*) from the high tropical Andes. Global Ecology and Biogeography, 9(5), 403–413. Portico. https://doi.org/10.1046/j.1365-2699.2000.00187.x
Eupatorium perfoliatum	Byers, D. L., & Meagher, T. R. (1997).A comparison of demographic characteristics in a rare and a common species of Eupatorium. Ecological Applications, 7(2), 519–530. https://doi.org/10.1890/1051-0761(1997)007[0519:acodci]2.0.co;2
Eupatorium resinosum	Byers, D. L., & Meagher, T. R. (1997).A comparison of demographic characteristics in a rare and a common species of *Eupatorium*. Ecological Applications, 7(2), 519–530. https://doi.org/10.1890/1051-0761(1997)007[0519:acodci]2.0.co;2
Fagus grandifolia	Batista, W. B., Platt, W. J., & Macchiavelli, R. E. (1998). Demography of a shade-tolerant tree (*Fagus grandifolia*) in a hurricane-disturbed forest. Ecology, 79(1), 38. https://doi.org/10.2307/176863

Species	Publication
Fritillaria meleagris	Zhang, L., & Hytteborn, H. (1985). Effect of ground water regime on development and distribution of *Fritillaria meleagri*s. Ecography, 8(4), 237–244. https://doi.org/10.1111/j.1600-0587.1985.tb01174.x
Gardenia actinocarpa	Osunkoya, O. O. (2003). Two-sex population projection of the endemic and dioecious rainforest shrub, *Gardenia actinocarpa* (Rubiaceae). Biological Conservation, 114(1), 39–51. https://doi.org/10.1016/s0006-3207(02)00417-2
Gentianella campestris	Lennartsson, T., & Oostermeijer, J. G. B. (2001). Demographic variation and population viability in *Gentianella campestris*: effects of grassland management and environmental stochasticity. Journal of Ecology, 89(3), 451–463. Portico. https://doi.org/10.1046/j.1365-2745.2001.00566.x
Geonoma brevispatha	Souza, A. F., & Martins, F. R. (2006). Demography of the clonal palm *Geonoma brevispatha* in a Neotropical swamp forest. Austral Ecology, 31(7), 869–881. https://doi.org/10.1111/j.1442-9993.2006.01650.x
Geonoma orbignyana	Rodríguez-Buriticá, S., Orjuela, M. A., & Galeano, G. (2005). Demography and life history of *Geonoma orbignyana*: An understory palm used as foliage in Colombia. Forest Ecology and Management, 211(3), 329–340. https://doi.org/10.1016/j.foreco.2005.02.052
Geonoma schottiana	Sampaio, M. B., & Scariot, A. (2010). Effects of stochastic herbivory events on population maintenance of an understorey palm species (*Geonoma schottiana*) in riparian tropical forest. Journal of Tropical Ecology, 26(2), 151–161. https://doi.org/10.1017/s0266467409990599
Geranium sylvaticum	Ramula, S., Toivonen, E., & Mutikainen, P. (2007). Demographic consequences of pollen limitation and inbreeding depression in a gynodioecious herb. International Journal of Plant Sciences, 168(4), 443–453. https://doi.org/10.1086/512040
Geum rivale	Kiviniemi, K. (2002). Population dynamics of *Agrimonia eupatoria* and *Geum rivale*, two perennial grassland species. Plant Ecology, 159(2), 153–169. https://doi.org/10.1023/a:1015506019670

Species	Publication
Gilia tenuiflora subsp. hoffmannii	Levine, J. M., McEachern, A. K., & Cowan, C. (2008). Rainfall effects on rare annual plants. Journal of Ecology, 96(4), 795–806. https://doi.org/10.1111/j.1365-2745.2008.01375.x
Grias peruviana	Peters, C.M. 1990b. Population ecology and management of forest fruit trees in Peruvian Amazonian. In A.B. Anderson (ed.), Alternatives to Deforestation: Steps Toward Sustainable Use of the Amazon Rain Forest, pp. 86-98. Columbia University Press, New York
Guaiacum sanctum	Lopez-Toledo, L., Burslem, D. F. R. P., Martinez-Ramos, M. & Garcia-Naranno, A. (2008): Non-detriment findings report on *Guaiacum sanctum* in Mexico. NDF Workshop Case Studies, WG1 - Trees, Case Study 7. CITES Plant Commitee, Mexico
Haplopappus radiatus	Kaye, T. N., & Pyke, D. A. (2003). The effect of stochastic technique on estimates of population viability from transition matrix models. Ecology, 84(6), 1464–1476. https://doi.org/10.1890/0012-9658(2003)084[1464:teosto]2.0.co;2
Helenium virginicum	Adams, V. M., Marsh, D. M., & Knox, J. S. (2005). Importance of the seed bank for population viability and population monitoring in a threatened wetland herb. Biological Conservation, 124(3), 425–436. https://doi.org/10.1016/j.biocon.2005.02.001
Helianthemum juliae	Marrero-Gómez, M. V., Oostermeijer, J. G. B., Carqué-Álamo, E., & Bañares-Baudet, Á. (2007). Population viability of the narrow endemic *Helianthemum juliae* (Cistaceae) in relation to climate variability. Biological Conservation, 136(4), 552–562. https://doi.org/10.1016/j.biocon.2007.01.010
Helianthemum polygonoides	Iriondo, J.M., Albert M. J., Gimenez-Benavides, L., Dominguez-Lozano, F. & Escudero, A. [Eds.] (2009): Poblaciones en peligro: viabilidad demográfica de la flora vascular amenazada de España. Dirección General de Medio Natural y Política Forestal (Ministerio de Medio Ambiente, y Medio Rural y Marino), Madrid, 242 pp.

Species	Publication
Heliconia acuminata	Bruna, E. M. (2003). Are plant populations in fragmented habitats recruitment limited? Tests with an Amazonian herb. Ecology, 84(4), 932–947. https://doi.org/10.1890/0012-9658(2003)084[0932:appifh]2.0.co;2
Heracleum mantegazzianum	Nehrbass, N., Winkler, E., Pergl, J., Perglova, I., & Pysek, P. (2006). Empirical and virtual investigation of the population dynamics of an alien plant under the constraints of local carrying capacity: *Heracleum mantegazzianum* in the Czech Republic. Perspectives in Plant Ecology, Evolution and Systematics, 7(4), 253–262. https://doi.org/10.1016/j.ppees.2005.11.001
Hieracium floribundum	Thomas, A. G., & Dale, H. M. (1975). The role of seed reproduction in the dynamics of established populations of *Hieracium floribundum* and a comparison with that of vegetative reproduction. Canadian Journal of Botany, 53(24), 3022–3031. https://doi.org/10.1139/b75-331
Himantoglossum hircinum	Pfeifer, M., Wiegand, K., Heinrich, W., & Jetschke, G. (2006). Long-term demographic fluctuations in an orchid species driven by weather: implications for conservation planning. Journal of Applied Ecology, 43(2), 313–324. https://doi.org/10.1111/j.1365-2664.2006.01148.x
Hymenoxys herbacea	Campbell, L. G., & Husband, B. C. (2005). Impact of clonal growth on effective population size in *Hymenoxys herbacea* (Asteraceae). Heredity, 94(5), 526–532. https://doi.org/10.1038/sj.hdy.6800653
Hypericum cumulicola	Quintana-Ascencio, P. F., Menges, E. S., & Weekley, C. W. (2003). A fire-explicit population viability analysis of *Hypericum cumulicola* in Florida rosemary scrub. Conservation Biology, 17(2), 433–449. https://doi.org/10.1046/j.1523-1739.2003.01431.x
Ipomopsis tenuituba	Campbell, D. R., & Waser, N. M. (2007). Evolutionary dynamics of an *Ipomopsis* hybrid zone: Confronting models with lifetime fitness data. The American Naturalist, 169(3), 298–310. https://doi.org/10.1086/510758
Iris germanica	Burns, J. H., Pardini, E. A., Schutzenhofer, M. R., Chung, Y. A., Seidler, K. J., & Knight, T. M. (2013). Greater sexual reproduction contributes to differences

Species	Publication
	in demography of invasive plants and their noninvasive relatives. Ecology, 94(5), 995–1004. https://doi.org/10.1890/12-1310.1
Jurinea fontqueri	Iriondo, J.M., Albert M. J., Gimenez-Benavides, L., Dominguez-Lozano, F. & Escudero, A. [Eds.] (2009): Poblaciones en peligro: viabilidad demográfica de la flora vascular amenazada de España. Dirección General de Medio Natural y Política Forestal (Ministerio de Medio Ambiente, y Medio Rural y Marino), Madrid, 242 pp.
Khaya senegalensis	Gaoue, O. G., & Ticktin, T. (2010). Effects of harvest of nontimber forest products and ecological differences between sites on the demography of african mahogany. Conservation Biology, 24(2), 605–614. https://doi.org/10.1111/j.1523-1739.2009.01345.x
Knautia arvensis	Johansen, L., Wehn, S., & Hovstad, K. A. (2016). Clonal growth buffers the effect of grazing management on the population growth rate of a perennial grassland herb. Flora, 223, 11–18. https://doi.org/10.1016/j.flora.2016.04.007
Kosteletzkya pentacarpos	Pino, J., Picó, F. X., & De Roa, E.(2007). Population dynamics of the rare plant *Kosteletzkya pentacarpos* (Malvaceae): a nine-year study. Botanical Journal of the Linnean Society, 153(4), 455–462. https://doi.org/10.1111/j.1095-8339.2007.00628.x
Kummerowia striata	Levin, S. C., Crandall, R. M., & Knight, T. M. (2019). Population projection models for 14 alien plant species in the presence and absence of aboveground competition. Ecology, e02681. Portico. https://doi.org/10.1002/ecy.2681
Lactuca serriola	Bullock, J. M., White, S. M., Prudhomme, C., Tansey, C., Perea, R., & Hooftman, D. A. P. (2011). Modelling spread of British wind-dispersed plants under future wind speeds in a changing climate. Journal of Ecology, 100(1), 104–115. https://doi.org/10.1111/j.1365-2745.2011.01910.x
Lantana camara	Raghu, S., Osunkoya, O. O., Perrett, C., & Pichancourt, J.-B. (2014). Historical demography of *Lantana camara* L. reveals clues about the influence of land use and weather in the management of this widespread

Species	Publication
	invasive species. Basic and Applied Ecology, 15(7), 565–572. https://doi.org/10.1016/j.baae.2014.08.006
Lathyrus vernus	de Vries, C., & Caswell, H. (2017). Demography when history matters: construction and analysis of second-order matrix population models. Theoretical Ecology, 11(2), 129–140. https://doi.org/10.1007/s12080-017-0353-0
Lathyrus vernus	Ehrlen, J. (1995). Demography of the perennial herb *Lathyrus vernus*. II. Herbivory and population dynamics. The Journal of Ecology, 83(2), 297. https://doi.org/10.2307/2261568
Lechea cernua	Maliakal Witt, S. (2004): Microhabitat distribution and demography of two Florida scrub endemic plants with comparisons to their habitat-generalist congeners. PhD Thesis, Louisiana State Eniversity, Louisiana.
Lechea deckertii	Maliakal Witt, S. (2004): Microhabitat distribution and demography of two Florida scrub endemic plants with comparisons to their habitat-generalist congeners. PhD Thesis, Louisiana State Eniversity, Louisiana.
Leucopogon setiger	Swab R. M. (2014): Increasing understanding of species responses to global changes through modeling plant metapopulation dynamics. PhD Thesis, University of California, Riverside.
Limonium carolinianum	Baltzer, J. L., Reekie, E. G., Hewlin, H. L., Taylor, P. D., & Boates, J. S. (2002). Impact of flower harvesting on the salt marsh plant *Limonium carolinianum*. Canadian Journal of Botany, 80(8), 841–851. https://doi.org/10.1139/b02-070
Limonium delicatulum	Hegazy, A. K. (1992). Age-specific survival, mortality and reproduction, and prospects for conservation of *Limonium delicatulum*. The Journal of Applied Ecology, 29(3), 549. https://doi.org/10.2307/2404462
Limonium erectum	Iriondo, J.M., Albert M. J., Gimenez-Benavides, L., Dominguez-Lozano, F. & Escudero, A. [Eds.] (2009): Poblaciones en peligro: viabilidad demográfica de la flora vascular amenazada de España. Dirección General de Medio

Species	Publication
	Natural y Política Forestal (Ministerio de Medio Ambiente, y Medio Rural y Marino), Madrid, 242 pp.
Limonium geronense	Iriondo, J.M., Albert M. J., Gimenez-Benavides, L., Dominguez-Lozano, F. & Escudero, A. [Eds.] (2009): Poblaciones en peligro: viabilidad demográfica de la flora vascular amenazada de España. Dirección General de Medio Natural y Política Forestal (Ministerio de Medio Ambiente, y Medio Rural y Marino), Madrid, 242 pp.
Limonium malacitanum	Iriondo, J.M., Albert M. J., Gimenez-Benavides, L., Dominguez-Lozano, F. & Escudero, A. [Eds.] (2009): Poblaciones en peligro: viabilidad demográfica de la flora vascular amenazada de España. Dirección General de Medio Natural y Política Forestal (Ministerio de Medio Ambiente, y Medio Rural y Marino), Madrid, 242 pp.
Lindera umbellate subsp. *membrancea*	Hara, M., Kanno, H., Hirabuki, Y., & Takehara, A. (2004). Population dynamics of four understorey shrub species in beech forest. Journal of Vegetation Science, 15(4), 475–484. Portico. https://doi.org/10.1111/j.1654-1103.2004.tb02286.x
Linum catharticum	Verkaar, H. J., & Schenkeveld, A. J. (1984). On the ecology of short-lived forbs in chalk grasslands: life-history characteristics. New Phytologist, 98(4), 659–672. https://doi.org/10.1111/j.1469-8137.1984.tb04155.x
Linum flavum	Münzbergová, Z. (2013). Comparative demography of two co-occurring *Linum* species with different distribution patterns. Plant Biology, 15(6), 963–970. https://doi.org/10.1111/plb.12007
Linum tenuifolium	Münzbergová, Z. (2013). Comparative demography of two co-occurring *Linum* species with different distribution patterns. Plant Biology, 15(6), 963–970. https://doi.org/10.1111/plb.12007
Lithospermum ruderale	Bricker, M., & Maron, J. (2012). Postdispersal seed predation limits the abundance of a long-lived perennial forb (*Lithospermum ruderale*). Ecology, 93(3), 532–543. https://doi.org/10.1890/11-0948.1

Species	Publication
Lomatium bradshawii	Kaye, T. N., Pendergrass, K. L., Finley, K., & Kauffman, J. B. (2001). The effect of fire on the population viability of an endangered prairie plant. Ecological Applications, 11(5), 1366–1380. https://doi.org/10.1890/1051-0761(2001)011[1366:teofot]2.0.co;2
Lomatium cookii	Kaye, T. N., & Pyke, D. A. (2003). The effect of stochastic technique on estimates of population viability from transition matrix models. Ecology, 84(6), 1464–1476. https://doi.org/10.1890/0012-9658(2003)084[1464:teosto]2.0.co;2
Lophophora diffusa	Diaz Segura O. (2013): Dinámica poblacional de *Lophophora diffusa* "peyote" (Cactaceae) en una localidad del Estado de Querétaro. MSc. Thesis, Universidad Autónoma Metropolitana, Mexico.
Lupinus arboreus	Kauffman, M. J., & Maron, J. L. (2006). Consumers limit the abundance and dynamics of a perennial shrub with a seed bank. The American Naturalist, 168(4), 454–470. https://doi.org/10.1086/507877
Lupinus lepidus var. *lobii*	Bishop, J. G. (1996): Demographic and population genetic variation during colonization by the herb *Lupinus lepidus* on Mount St. Helens. PhD Thesis, University of Washington.
Lupinus tidestromii	Dangremond, E. M., Pardini, E. A., & Knight, T. M. (2010). Apparent competition with an invasive plant hastens the extinction of an endangered lupine. Ecology, 91(8), 2261–2271. https://doi.org/10.1890/09-0418.1
Magnolia dealbata	Sánchez-Velásquez, L. R., & Pineda-López, M. del R. (2009). Comparative demographic analysis in contrasting environments of Magnolia dealbata: an endangered species from Mexico. Population Ecology, 52(1), 203–210. Portico. https://doi.org/10.1007/s10144-009-0161-5
Malacothrix indecora	Levine, J. M., McEachern, A. K., & Cowan, C. (2008). Rainfall effects on rare annual plants. Journal of Ecology, 96(4), 795–806. https://doi.org/10.1111/j.1365-2745.2008.01375.x

Species	Publication
Mammillaria hernandezii	Rodríguez-Ortega C. (2008). Consecuencias demográficas y evolutivas del secuestro de semillas en tres especies del género *Mammillaria* (Cactaceae). PhD Dissertation, Universidad Autónoma Metropolitana, Mexico.
Mammillaria huitzilopochtli	Martínez, A. F., Medina, G. I. M., Golubov, J., Montaña, C., & Mandujano, M. C. (2010). Demography of an endangered endemic rupicolous cactus. Plant Ecology, 210(1), 53–66. https://doi.org/10.1007/s11258-010-9737-6
Mammillaria magnimamma	Valverde, T., Quijas, S., López-Villavicencio, M., & Castillo, S. (2004). Population dynamics of *Mammillaria magnimamma* Haworth. (Cactaceae) in a lava-field in central Mexico. Plant Ecology (Formerly Vegetatio), 170(2), 167–184. https://doi.org/10.1023/b:vege.0000021662.78634.de
Mammillaria solisioides	Rodríguez-Ortega C. (2008). Consecuencias demográficas y evolutivas del secuestro de semillas en tres especies del género *Mammillaria* (Cactaceae). PhD Dissertation, Universidad Autónoma Metropolitana, Mexico.
Manilkara zapota	Cruz-Rodríguez, J. A., López-Mata, L., & Valverde, T. (2009). A comparison of traditional elasticity and variance-standardized perturbation analyses: a case study with the tropical tree species *Manilkara zapota* (Sapotaceae). Journal of Tropical Ecology, 25(2), 135–146. https://doi.org/10.1017/s0266467408005713
Miconia albicans	Hoffmann, W. A. (1999).Fire and population dynamics of woody plants in a neotropical savanna: matrix model projections. Ecology, 80(4), 1354–1369. https://doi.org/10.1890/0012-9658(1999)080[1354:fapdow]2.0.co;2
Mimulus cardinalis	Angert, A. L. (2006). Demography of central and marginal populations of monkeyflowers (*Mimulus cardinalisandm*. Lewisii). Ecology, 87(8), 2014–2025. https://doi.org/10.1890/0012-9658(2006)87[2014:docamp]2.0.co;2
Mimulus lewisii	Angert, A. L. (2006). Demography of central and marginal populations of monkeyflowers (*Mimulus cardinalisandm*. Lewisii). Ecology, 87(8), 2014–2025. https://doi.org/10.1890/0012-9658(2006)87[2014:docamp]2.0.co;2

Species	Publication
Miscanthus giganteus	Matlaga, D. P., & Davis, A. S. (2013). Minimizing invasive potential of *Miscanthus × giganteus* grown for bioenergy: identifying demographic thresholds for population growth and spread. Journal of Applied Ecology, 50(2), 479–487. Portico. https://doi.org/10.1111/1365-2664.12057
Mitrocereus fulviceps	Vite González, F. & J. Zavala Hurtado, J. A. (1998). Estatus ecológicos de *Mammillaria pectinifera* Weber y *Pachycereus fulviceps* Weber en el Valle de Zapotitlán, Puebla. Universidad Autónoma Metropolitana-Iztapalapa. División de Ciencias Biológicas y de la Salud. Informe final SNIB- CONABIO proyecto No. G022. México D. F.
Molinia caerulea	Jacquemyn, H., Brys, R., & Neubert, M. G. (2005). Fire increases invasive spread of *Molinia caerulea* mainly through changes in demographic parameters. Ecological Applications, 15(6), 2097–2108. https://doi.org/10.1890/04-1762
Mulinum spinosum	Cipriotti, P. A., & Aguiar, M. R. (2011). Direct and indirect effects of grazing constrain shrub encroachment in semi-arid Patagonian steppes. Applied Vegetation Science, 15(1), 35–47. https://doi.org/10.1111/j.1654-109x.2011.01138.x
Myosotis ramosissima	Dostál, P. (2007). Population dynamics of annuals in perennial grassland controlled by ants and environmental stochasticity. Journal of Vegetation Science, 18(1), 91–102. Portico. https://doi.org/10.1111/j.1654-1103.2007.tb02519.x
Narcissus pseudonarcissus	Barkham, J. P. (1980). Population dynamics of the wild daffodil (Narcissus pseudonarcissus): I. Clonal growth, seed reproduction, mortality and the effects of density. The Journal of Ecology, 68(2), 607. https://doi.org/10.2307/2259425
Neobuxbaumia macrocephala	Esparza-Olguín, L., Valverde, T., & Mandujano, M. C. (2005). Comparative demographic analysis of three *Neobuxbaumia species* (Cactaceae) with differing degree of rarity. Population Ecology, 47(3), 229–245. Portico. https://doi.org/10.1007/s10144-005-0230-3

Species	Publication
Neobuxbaumia mezcalaensis	Esparza-Olguín, L., Valverde, T., & Mandujano, M. C. (2005). Comparative demographic analysis of three *Neobuxbaumia species* (Cactaceae) with differing degree of rarity. Population Ecology, 47(3), 229–245. Portico. https://doi.org/10.1007/s10144-005-0230-3
Neobuxbaumia polylopha	Arroyo-Cosultchi, G., Golubov, J., & Mandujano, M. C. (2016). Pulse seedling recruitment on the population dynamics of a columnar cactus: Effect of an extreme rainfall event. Acta Oecologica, 71, 52–60. https://doi.org/10.1016/j.actao.2016.01.006
Neodypsis decaryi	Ratsirarson, J., Silander, J. A., & Richard, A. F. (1996). Conservation and management of a threatened Madagascar palm species, Neodypsis decaryi, Jumelle. Conservation Biology, 10(1), 40–52. https://doi.org/10.1046/j.1523-1739.1996.10010040.x
Oenothera deltoides subsp. howellii	Thompson, D. M. (2005). Matrix models as a tool for understanding invasive plant and native plant interactions. Conservation Biology, 19(3), 917–928. https://doi.org/10.1111/j.1523-1739.2005.004108.x
Orchis purpurea	Jacquemyn, H., Brys, R., & Jongejans, E. (2010). Seed limitation restricts population growth in shaded populations of a perennial woodland orchid. Ecology, 91(1), 119–129. https://doi.org/10.1890/08-2321.1
Pachycereus pecten-aboriginum	Morales-Romero, D., Godínez-Álvarez, H., Campo-Alves, J., & Molina-Freaner, F. (2012). Effects of land conversion on the regeneration of *Pachycereus pecten-aboriginum* and its consequences on the population dynamics in northwestern Mexico. Journal of Arid Environments, 77, 123–129. https://doi.org/10.1016/j.jaridenv.2011.09.005
Paeonia officinalis	Andrieu, E., Fréville, H., Besnard, A., Vaudey, V., Gauthier, P., Thompson, J. D., & Debussche, M. (2012). Forest-cutting rapidly improves the demographic status of *Paeonia officinalis*, a species threatened by forest closure. Population Ecology, 55(1), 147–158. Portico. https://doi.org/10.1007/s10144-012-0346-1

Species	Publication
Panax quinquefolius	Farrington, S. J., Muzika, R.-M., Drees, D., & Knight, T. M.(2009).Interactive effects of harvest and deer herbivory on the population dynamics of American ginseng. Conservation Biology, 23(3), 719–728. https://doi.org/10.1111/j.1523-1739.2008.01136.x
Parkinsonia aculeata	Raghu, S., Wilson, J. R., & Dhileepan, K. (2006). Refining the process of agent selection through understanding plant demography and plant response to herbivory. Australian Journal of Entomology, 45(4), 308–316. https://doi.org/10.1111/j.1440-6055.2006.00556.x
Paronychia pulvinata	Forbis, T. A., & Doak, D. F. (2004). Seedling establishment and life history trade-offs in alpine plants. American Journal of Botany, 91(7), 1147–1153. Portico. https://doi.org/10.3732/ajb.91.7.1147
Pedicularis furbishiae	Menges, E. S. (1990). Population viability analysis for an endangered plant. Conservation Biology, 4(1), 52–62. https://doi.org/10.1111/j.1523-1739.1990.tb00267.x
Periandra mediterranea	Hoffmann, W. A., & Solbrig, O. T. (2003). The role of topkill in the differential response of savanna woody species to fire. Forest Ecology and Management, 180(1–3), 273–286. https://doi.org/10.1016/s0378-1127(02)00566-2
Persoonia bargoensis	McKenna D. J. (2007). Demographic and ecological indicators of rarity in a suite of obligate-seeding *Persoonia* (Proteaceae) shrubs. PhD Thesis, University of Wollongong.
Persoonia glaucescens	McKenna D. J. (2007). Demographic and ecological indicators of rarity in a suite of obligate-seeding *Persoonia* (Proteaceae) shrubs. PhD Thesis, University of Wollongong.
Petrocoptis pseudoviscosa	García, M. B., Guzmán, D., & Goñi, D. (2002). An evaluation of the status of five threatened plant species in the Pyrenees. Biological Conservation, 103(2), 151–161. https://doi.org/10.1016/s0006-3207(01)00113-6

Species	Publication
Petrophile pulchella	Bradstock, R. A., & O'Connell, M. A. (1988). Demography of woody plants in relation to fire: *Banksia ericifolia* L.f. and *Petrophile pulchella* (Schrad) R.Br. Austral Ecology, 13(4), 505–518. https://doi.org/10.1111/j.1442-9993.1988.tb00999.x
Phacelia insularis var. *insularis*	Levine, J. M., McEachern, A. K., & Cowan, C. (2008). Rainfall effects on rare annual plants. Journal of Ecology, 96(4), 795–806. https://doi.org/10.1111/j.1365-2745.2008.01375.x
Phaseolus lunatus	Degreef, J., Baudoin, J.-P., & Rocha, O. J. (1997). Case studies on breeding systems and its consequences for germplasm conservation. Genetic Resources and Crop Evolution, 44(5), 429–438. https://doi.org/10.1023/a:1008623521755
Phyllanthus emblica	Ellis, M. M., Williams, J. L., Lesica, P., Bell, T. J., Bierzychudek, P., Bowles, M., Crone, E. E., Doak, D. F., Ehrlén, J., Ellis-Adam, A., McEachern, K., Ganesan, R., Latham, P., Luijten, S., Kaye, T. N., Knight, T. M., Menges, E. S., Morris, W. F., Nijs, H. den, … Weekley, C. W. (2012). Matrix population models from 20 studies of perennial plant populations. Ecology, 93(4), 951–951. Portico. https://doi.org/10.1890/11-1052.1
Phyllanthus indofischeri	Ticktin, T., Ganesan, R., Paramesha, M., & Setty, S. (2012). Disentangling the effects of multiple anthropogenic drivers on the decline of two tropical dry forest trees. Journal of Applied Ecology, 49(4), 774–784. https://doi.org/10.1111/j.1365-2664.2012.02156.x
Phytelephas seemannii	Bernal, R. (1998). Demography of the vegetable ivory palm *Phytelephas seemannii* in Colombia, and the impact of seed harvesting. Journal of Applied Ecology, 35(1), 64–74. Portico. https://doi.org/10.1046/j.1365-2664.1998.00280.x
Picris hieracioides	Klemow, K. M., & Raynal, D. J. (1985). Demography of two facultative biennial plant species in an unproductive habitat. The Journal of Ecology, 73(1), 147. https://doi.org/10.2307/2259775

Species	Publication
Pimpinella saxifraga	Auestad, I., Rydgren, K., Jongejans, E., & Kroon, H. de. (2010). *Pimpinella saxifraga* is maintained in road verges by mosaic management. Biological Conservation, 143(4), 899–907. https://doi.org/10.1016/j.biocon.2009.12.037
Pinguicula alpina	Svensson, B. M., Carlsson, B. A., Karlsson, P. S., & Nordell, K. O. (1993). Comparative long-term demography of three species of *Pinguicula*. The Journal of Ecology, 81(4), 635. https://doi.org/10.2307/2261662
Pinguicula villosa	Svensson, B. M., Carlsson, B. A., Karlsson, P. S., & Nordell, K. O. (1993). Comparative long-term demography of three species of *Pinguicula*. The Journal of Ecology, 81(4), 635. https://doi.org/10.2307/2261662
Pinus albicaulis	Ettl, G., & N. Cottone (2004). Whitebark pine (*Pinus albicaulis*) in Mt. Rainier National Park: response to blister rust infection. Pages 36–47 in H. Akc akaya, M. Burgman, O.Kindvall, C. Wood, P. Sjogren-Gulve, J. Hatfield, and M.McCarthy (Eds.). Species conservation and management. Oxford University Press, New York, New York, USA
Pinus kwangtungensis	Chien, P. D., Zuidema, P. A., & Nghia, N. H. (2008). Conservation prospects for threatened Vietnamese tree species: results from a demographic study. Population Ecology, 50(2), 227–237. Portico. https://doi.org/10.1007/s10144-008-0079-3
Pinus lambertiana	Van Mantgem, P. J., & Stephenson, N. L. (2005). The accuracy of matrix population model projections for coniferous trees in the Sierra Nevada, California. Journal of Ecology, 93(4), 737–747. Portico. https://doi.org/10.1111/j.1365-2745.2005.01007.x
Pinus maximartinezii	López-Mata, L. (2013). The impact of seed extraction on the population dynamics of *Pinus maximartinezii*. Acta Oecologica, 49, 39–44. https://doi.org/10.1016/j.actao.2013.02.010
Pinus nigra subsp. *lauricio*	Buckley, Y. M., Brockerhoff, E., Langer, L., Ledgard, N., North, H., & Rees, M.(2005). Slowing down a pine invasion despite uncertainty in demography

Species	Publication
	and dispersal. Journal of Applied Ecology, 42(6), 1020–1030. https://doi.org/10.1111/j.1365-2664.2005.01100.x
Pinus strobus	Münzbergová, Z., Hadincová, V., Wild, J., & Kindlmannová, J. (2013). Variability in the contribution of different life stages to population growth as a key factor in the invasion success of *Pinus strobus*. PLoS ONE, 8(2), e56953. https://doi.org/10.1371/journal.pone.0056953
Pinus sylvestris	Usher, M. B. (1966). A matrix approach to the management of renewable resources, with special reference to selection forests. The Journal of Applied Ecology, 3(2), 355. https://doi.org/10.2307/2401258
Plantago coronopus	Villellas, J., Ehrlén, J., Olesen, J. M., Braza, R., & García, M. B. (2012). Plant performance in central and northern peripheral populations of the widespread *Plantago coronopus*. Ecography, 36(2), 136–145. https://doi.org/10.1111/j.1600-0587.2012.07425.x
Plantago media	Eriksson, Å., & Eriksson, O. (2000). Population dynamics of the perennial *Plantago media* in semi-natural grasslands. Journal of Vegetation Science, 11(2), 245–252. Portico. https://doi.org/10.2307/3236803
Polemonium van-bruntiae	Hill Bermingham, L. (2010). Deer herbivory and habitat type influence long-term population dynamics of a rare wetland plant. Plant Ecology, 210(2), 359–378. https://doi.org/10.1007/s11258-010-9762-5
Polygonella basiramia	Maliakal Witt, S. (2004): Microhabitat distribution and demography of two Florida scrub endemic plants with comparisons to their habitat-generalist congeners. PhD Thesis, Louisiana State Eniversity, Louisiana.
Potentilla anserina	Eriksson, O. (1988). Ramet behaviour and population growth in the clonal herb *Potentilla anserina*. The Journal of Ecology, 76(2), 522. https://doi.org/10.2307/2260610
Primula elatior	Jacquemyn, H., & Brys, R. (2008). Effects of stand age on the demography of a temperate forest herb in post-agricultural forests. Ecology, 89(12), 3480–3489. https://doi.org/10.1890/07-1908.1

Species	Publication
Primula veris	Ehrlén, J., Syrjänen, K., Leimu, R., Begoña Garcia, M., & Lehtilä, K. (2005). Land use and population growth of *Primula veris*: an experimental demographic approach. Journal of Applied Ecology, 42(2), 317–326. https://doi.org/10.1111/j.1365-2664.2005.01015.x
Primula vulgaris	Valverde, T., & Silvertown, J. (1998). Variation in the demography of a woodland understorey herb (*Primula vulgaris*) along the forest regeneration cycle: projection matrix analysis. Journal of Ecology, 86(4), 545–562. Portico. https://doi.org/10.1046/j.1365-2745.1998.00280.x
Prosopis glandulosa	Golubov, J., Mandujano, M. D. C., Franco, M., Montana, C., Eguiarte, L. E., & Lopez-Portillo, J. (1999). Demography of the invasive woody perennial *Prosopis glandulosa* (honey mesquite). Journal of Ecology, 87(6), 955–962. https://doi.org/10.1046/j.1365-2745.1999.00420.x
Prosopis laevigata	Bernal R. (2010). Comportamiento demográfico y la dinámica espacio-temporal de la planta epífita *Tillandsia recurvata* L. (Bromeliaceae). PhD thesis. Universidad Nacional Autónoma de Mexico.
Prunus africana	Stewart, K. M. 2001. The commercial bark harvest of the African cherry (*Prunus africana*) on Mount Oku, Cameroon: effects on traditional uses and population dynamics. Ph.D. dissertation, Florida International University, Miami, FL.
Prunus serotina	Sebert-Cuvillier, E., Paccaut, F., Chabrerie, O., Endels, P., Goubet, O., & Decocq, G. (2007). Local population dynamics of an invasive tree species with a complex life-history cycle: A stochastic matrix model. Ecological Modelling, 201(2), 127–143. https://doi.org/10.1016/j.ecolmodel.2006.09.005
Pseudophoenix sargentii	Durán, R. & R. Franco. 1992. Estudio demográfico de *Pseudophoenix sargentii*. Bulletin de l'Institut Français d'Études Andines 21: 609-621.
Purshia subintegra	Maschinski, J., Baggs, J. E., Quintana-Ascencio, P. F., & Menges, E. S. (2006). Using population viability analysis to predict the effects of climate change on the extinction risk of an endangered limestone endemic shrub, arizona

Species	Publication
	cliffrose. Conservation Biology, 20(1), 218–228. https://doi.org/10.1111/j.1523-1739.2006.00272.x
Quercus crispula	Hiura, T., & Fujiwara, K. (1999). Density-dependence and co-existence of conifer and broad-leaved trees in a Japanese northern mixed forest. Journal of Vegetation Science, 10(6), 843–850. Portico. https://doi.org/10.2307/3237309
Quercus rugosa	Bonfi, C. (2006): Regeneration and population dynamics of *Quercus rugosa* at the Ajusco Volcano, Mexico. In: Kappelle, M. (eds) Ecology and Conservation of Neotropical Montane Oak Forests. Ecological Studies, vol 185. Springer, Berlin, Heidelberg. https://doi.org/10.1007/3-540-28909-7_12
Ranunculus acris	Sarukhan, J., & Harper, J. L. (1973). Studies on plant demography: *Ranunculus repens* L., *R. bulbosus* L. and *R. acris* L.: I. Population flux and survivorship. The Journal of Ecology, 61(3), 675. https://doi.org/10.2307/2258643
Rhizophora mangle	López-Hoffman, L., Ackerly, D. D., Anten, N. P. R., Denoyer, J. L., & Martinez-Ramos, M. (2007). Gap-dependence in mangrove life-history strategies: a consideration of the entire life cycle and patch dynamics. Journal of Ecology, 95(6), 1222–1233. https://doi.org/10.1111/j.1365-2745.2007.01298.x
Rhododendron ponticum	Salguero-Gomez R. 2004. Markov Chains applied to *Rhododendron ponticum* L.: ecological terminator in Great Britain ecologically terminated in Spain? MSc thesis. Kingston University, London.
Rubus discolor	Lambrecht-McDowell, S. C., & Radosevich, S. R. (2005). Population demographics and trade-offs to reproduction of an invasive and noninvasive species of *Rubus*. Biological Invasions, 7(2), 281–295. https://doi.org/10.1007/s10530-004-0870-9
Rubus ursinus	Lambrecht-McDowell, S. C., & Radosevich, S. R. (2005). Population demographics and trade-offs to reproduction of an invasive and

Species	Publication
	noninvasive species of *Rubus*. Biological Invasions, 7(2), 281–295. https://doi.org/10.1007/s10530-004-0870-9
Sabal minor	Ramp, P.F. (1989) Natural history of *Sabal minor*: Demography, population genetics, and reproductive biology. Ph.D. dissertation, Tulane University, New Orleans, Louisiana, 211 pp.
Salsola australis	Borger, C. P. D., Scott, J. K., Renton, M., Walsh, M., & Powles, S. B. (2009). Assessment of management options for *Salsola australis* in south-west Australia by transition matrix modelling. Weed Research, 49(4), 400–408. https://doi.org/10.1111/j.1365-3180.2009.00703.x
Sanicula europaea	Gustafsson, C., & Ehrlén, J. (2003). Effects of intraspecific and interspecific density on the demography of a perennial herb, *Sanicula europaea*. Oikos, 100(2), 317–324. Portico. https://doi.org/10.1034/j.1600-0706.2003.11493.x
Sarcocapnos baetica	Salinas, M. J., Suárez, V., & Blanca, G. (2002). Demographic structure of three species of *Sarcocapnos* (Fumariaceae) as a basis for their conservation. Canadian Journal of Botany, 80(4), 360–369. https://doi.org/10.1139/b02-013
Sarcocapnos enneaphylla	Salinas, M. J., Suárez, V., & Blanca, G. (2002). Demographic structure of three species of *Sarcocapnos* (Fumariaceae) as a basis for their conservation. Canadian Journal of Botany, 80(4), 360–369. https://doi.org/10.1139/b02-013
Sarcocapnos pulcherrima	Salinas, M. J., Suárez, V., & Blanca, G. (2002). Demographic structure of three species of *Sarcocapnos* (Fumariaceae) as a basis for their conservation. Canadian Journal of Botany, 80(4), 360–369. https://doi.org/10.1139/b02-013
Sarracenia alata	Brewer, J. S. (2001). A demographic analysis of fire-stimulated seedling establishment of *Sarracenia alata* (Sarraceniaceae). American Journal of Botany, 88(7), 1250–1257. Portico. https://doi.org/10.2307/3558336

Species	Publication
Saussurea medusa	Law, W., Salick, J., & Knight, T. M. (2010). The effects of pollen limitation on population dynamics of snow lotus (*Saussurea medusa* and *S. laniceps*, Asteraceae): Threatened Tibetan medicinal plants of the eastern Himalayas. Plant Ecology, 210(2), 343–357. https://doi.org/10.1007/s11258-010-9761-6
Saxifraga tridactylites	Dostál, P. (2007). Population dynamics of annuals in perennial grassland controlled by ants and environmental stochasticity. Journal of Vegetation Science, 18(1), 91–102. Portico. https://doi.org/10.1111/j.1654-1103.2007.tb02519.x
Scabiosa columbaria	Verkaar, H. J., & Schenkeveld, A. J. (1984). On the ecology of short-lived forbs in chalk grasslands: life-history characteristics. New Phytologist, 98(4), 659–672. https://doi.org/10.1111/j.1469-8137.1984.tb04155.x
Scorzonera hispanica	Münzbergová, Z. (2006). Effect of population size on the prospect of species survival. Folia Geobotanica, 41(2), 137–150. https://doi.org/10.1007/bf02806475
Senecio filaginoides	Cipriotti, P. A., & Aguiar, M. R. (2011). Direct and indirect effects of grazing constrain shrub encroachment in semi-arid Patagonian steppes. Applied Vegetation Science, 15(1), 35–47. https://doi.org/10.1111/j.1654-109x.2011.01138.x
Shorea acuminata	Yamada, T., Yamada, Y., Okuda, T., & Fletcher, C. (2012). Soil-related variations in the population dynamics of six dipterocarp tree species with strong habitat preferences. Oecologia, 172(3), 713–724. https://doi.org/10.1007/s00442-012-2529-z
Shorea bracteolata	Yamada, T., Yamada, Y., Okuda, T., & Fletcher, C. (2012). Soil-related variations in the population dynamics of six dipterocarp tree species with strong habitat preferences. Oecologia, 172(3), 713–724. https://doi.org/10.1007/s00442-012-2529-z
Silene acaulis	Morris, W. F., & Doak, D. F. (1998). Life history of the long-lived gynodioecious cushion plant *Silene acaulis* (Caryophyllaceae), inferred from

Species	Publication
	size-based population projection matrices. American Journal of Botany, 85(6), 784–793. Portico. https://doi.org/10.2307/2446413
Silene spaldingii	Lesica, P., & Crone, E. E. (2007). Causes and consequences of prolonged dormancy for an iteroparous geophyte, *Silene spaldingii*. Journal of Ecology, 95(6), 1360–1369. https://doi.org/10.1111/j.1365-2745.2007.01291.x
Sonchus pustulatus	Silva, J. L., Mejías, J. A., & García, M. B. (2015). Demographic vulnerability in cliff-dwelling *Sonchus* species endemic to the western Mediterranean. Basic and Applied Ecology, 16(4), 316–324. https://doi.org/10.1016/j.baae.2015.02.009
Stenocereus eruca	Clark-Tapia, R., Mandujano, M. C., Valverde, T., Mendoza, A., & Molina-Freaner, F. (2005). How important is clonal recruitment for population maintenance in rare plant species?: the case of the narrow endemic cactus, *Stenocereus eruca*, in Baja California, México. Biological Conservation, 124(1), 123–132. https://doi.org/10.1016/J.BIOCON.2005.01.019
Stryphnodendron excelsum	Hartshorn, G. S. 1972. The ecological life history and population dynamics of Pen-taclethra macroloba, a tropical wet forest dominant and *Stryphnodendron excel-sum*, an occasional associate. Ph.D. Thesis, University Washington, Seattle.
Succisa pratensis	Milden, M. (2005): Local and regional dynamics of Succisa pratensis. PhD Thesis, University of Stockholm.
Syzygium jambos	Brown, K. A., Spector, S., & Wu, W. (2008). Multi-scale analysis of species introductions: combining landscape and demographic models to improve management decisions about non-native species. Journal of Applied Ecology, 45(6), 1639–1648. https://doi.org/10.1111/j.1365-2664.2008.01550.x
Taxus floridana	Kwit, C., Horvitz, C. C., & Platt, W. J. (2004). Conserving slow-growing, long-lived tree species: input from the demography of a rare understory conifer, *Taxus floridana*. Conservation Biology, 18(2), 432–443. https://doi.org/10.1111/j.1523-1739.2004.00567.x

Species	Publication
Thymus webbianus	Iriondo, J.M., Albert M. J., Gimenez-Benavides, L., Dominguez-Lozano, F. & Escudero, A. [Eds.] (2009): Poblaciones en peligro: viabilidad demográfica de la flora vascular amenazada de España. Dirección General de Medio Natural y Política Forestal (Ministerio de Medio Ambiente, y Medio Rural y Marino), Madrid, 242 pp.
Tillandsia macdougallii	Mondragón Chaparro, D., & Ticktin, T. (2011). Demographic effects of harvesting epiphytic bromeliads and an alternative approach to collection. Conservation Biology, 25(4), 797–807. https://doi.org/10.1111/j.1523-1739.2011.01691.x
Tillandsia multicaulis	Winkler, M., Hülber, K., & Hietz, P. (2007). Population dynamics of epiphytic bromeliads: Life strategies and the role of host branches. Basic and Applied Ecology, 8(2), 183–196. https://doi.org/10.1016/j.baae.2006.05.003
Tillandsia punctulata	Toledo-Aceves, T., Hernández-Apolinar, M., & Valverde, T. (2014). Potential impact of harvesting on the population dynamics of two epiphytic bromeliads. Acta Oecologica, 59, 52–61. https://doi.org/10.1016/j.actao.2014.05.009
Tillandsia violacea	Mondragón Chaparro, D., & Ticktin, T. (2011). Demographic effects of harvesting epiphytic bromeliads and an alternative approach to collection. Conservation Biology, 25(4), 797–807. https://doi.org/10.1111/j.1523-1739.2011.01691.x
Trillium grandiflorum	Knight, T. M. (2003). Effects of herbivory and its timing across populations of *Trillium grandiflorum* (Liliaceae). American Journal of Botany, 90(8), 1207–1214. Portico. https://doi.org/10.3732/ajb.90.8.1207
Trollius europaeus	Lemke, T., & Salguero-Gómez, R. (2015). Land use heterogeneity causes variation in demographic viability of a bioindicator of species-richness in protected fen grasslands. Population Ecology, 58(1), 165–178. https://doi.org/10.1007/s10144-015-0519-9

Species	Publication
Trollius laxus	Scanga, S. E., & Leopold, D. J. (2012). Managing wetland plant populations: Lessons learned in Europe may apply to North American fens. Biological Conservation, 148(1), 69–78. https://doi.org/10.1016/j.biocon.2012.01.061
Tsuga canadensis	Lamar, W. R., & McGraw, J. B. (2005). Evaluating the use of remotely sensed data in matrix population modeling for eastern hemlock (*Tsuga canadensis* L.). Forest Ecology and Management, 212(1–3), 50–64. https://doi.org/10.1016/j.foreco.2005.02.056
Veronica arvensis	Dostál, P. (2007). Population dynamics of annuals in perennial grassland controlled by ants and environmental stochasticity. Journal of Vegetation Science, 18(1), 91–102. Portico. https://doi.org/10.1111/j.1654-1103.2007.tb02519.x
Verticordia fimbrilepis subsp. *fimbrilepis*	Yates, C. J., & Ladd, P. G. (2010). Using population viability analysis to predict the effect of fire on the extinction risk of an endangered shrub *Verticordia fimbrilepis* subsp. *fimbrilepis* in a fragmented landscape. Plant Ecology, 211(2), 305–319. https://doi.org/10.1007/s11258-010-9791-0
Viola fimbriatula	Solbrig, O. T., Sarandon, R., & Bossert, W. (1988). A density-dependent growth model of a perennial herb, *Viola fimbriatula*. The American Naturalist, 131(3), 385–400. https://doi.org/10.1086/284796
Vochysia ferruginea	Boucher, D. H., & Mallona, M. A. (1997). Recovery of the rain forest tree *Vochysia ferruginea* over 5 years following Hurricane Joan in Nicaragua: a preliminary population projection matrix. Forest Ecology and Management, 91(2–3), 195–204. https://doi.org/10.1016/s0378-1127(96)03890-x
Zamia inermis	Octavio-Aguilar, P., Rivera-Fernández, A., Iglesias-Andreu, L. G., Vovides, P. A., & de Cáceres-González, F. F. N. (2017). Extinction risk of *Zamia inermis*: a demographic study in its single natural population. Biodiversity and Conservation, 26(4), 787–800. https://doi.org/10.1007/s10531-016-1270-z
Zea diploperennis	Sanchez-Velasquez, L. R., Ezcurra, E., Martinez-Ramos, M., Alvarez-Buylla, E., & Lorente, R. (2002). Population dynamics of *Zea diploperennis*, an endangered perennial herb: effect of slash and burn practice. Journal of

Species	Publication
	Ecology, 90(4), 684–692. https://doi.org/10.1046/j.1365-2745.2002.00702.x

Table S2.

Table S2: The effects of continent and plant growth form on vulnerability to seed harvesting. Results

of a linear model with vulnerability to seed harvesting (log-transformed) as a response variable and

plant growth form, continent, and their interaction as explanatory variables. We report results of a

simple linear model because a generalized least square model with phylogenetic correction failed

due to singular fit. The terms were fitted sequentially. Significant values are in bold. Adjusted R^2=0.15

	df	resid. df	F	p
Plant growth form	5	243	10.37	**<0.001**
Continent	5	243	1.12	0.350
Plant growth form × Continent	13	243	0.99	0.466

 Table S3.

 Table S3: Formulation of the life history traits used to explain species vulnerability to seed harvesting

 in 280 vascular plant species. λ is the population growth rate, which corresponds to the dominant

 eigenvalue of the matrix A; l_x and m_x are stage-specific survival and fertility schedules, C is the

 submatrix describing clonal reproduction, m is the dimension of the matrix C, w is the stable stage

 distribution of the matrix A, with j column entries of the matrix population model.

Life history trait	Biological meaning	Formula
Generation time T	Number of years necessary for the individuals of a population to be fully replaced by new ones	$T = \dfrac{log(\int_{1}^{\infty} l_x m_x \, dx)}{log(\lambda)}$
Degree of iteroparity S	Spread of reproduction throughout the lifespan of the individual as quantified by Demetrius' entropy (S). High/low S values correspond to iteroparous/semelparous populations	$S = -e^{-log\lambda} l_x m_x log(e^{-log\lambda} l_x m_x)$
Age at sexual maturity L_α	Number of years that it takes an average individual in the population to become sexually reproductive	L_α as described in Caswell 2001's equation 5.41 (*35*)
Seed bank residence	Mean amount of time individuals are expected to stay in the seedbank stage	As described in Caswell 2001's equation 5.36 (*35*) according to the fundamental matrix approach for the life cycle stage(s) that correspond to the seed bank stage(s)
Clonality K	Per-capita clonal contributions weighted by the stable stage distribution of the MPM	$K = \displaystyle\sum_{1}^{m} \bar{C}_j \bar{w}_j$

 Table S4.

 Table S4: The effects of life history traits on vulnerability to seed harvesting, with significant

 values ($P<0.05$) in bold. Adjusted $R^2 = 0.61$.

	df	resid. df	F	p
Generation time	1	264	215.08	**<0.001**
Degree of iteropartity	1	264	6.34	**0.012**
Age at sexual maturity	1	264	12.30	**<0.001**
Seed bank residence	1	264	5.14	**0.024**
Clonality	1	264	10.17	**0.002**
Plant growth type	5	264	3.57	**0.003**