Low concentrations of selenium nanoparticles enhance the performance of a

generalist parasitoid and its host, with no net effect on host suppression

RUNNING TITLE: SELENIUM NANOPARTICLES AND A PARASITOID

James Rudolph Miksanek[1]*, Charles Adarkwah[1,2,3], Midori Tuda[1]*

[1]Laboratory of Insect Natural Enemies, Institute of Biological Control, Faculty of Agriculture, Kyushu

University, Fukuoka 819-0395, Japan

[2]Department of Horticulture and Crop Production, School of Agriculture and Technology, Dormaa-

Ahenkro Campus, University of Energy and Natural Resources, Sunyani, Ghana

[3]Humboldt-University of Berlin, Division Urban Plant Ecophysiology, Faculty Life Sciences, 14195,

Berlin, Germany

*Correspondence:

Laboratory of Insect Natural Enemies

Faculty of Agriculture

Kyushu University

Fukuoka 819-0395 (Japan)

email: miks0007@umn.edu

ABSTRACT

BACKGROUND: The environmental and economic costs of conventional insecticides have stirred an interest in alternative management tactics, including the use of nanotechnologies. Selenium nanoparticles (SeNPs) have many applications in agriculture but may not be compatible with biological control; however, low concentrations of SeNPs may benefit natural enemies via hormesis. This study investigates the concentration-dependent effects of SeNPs (0–1000 mg L^{-1}) on *Anisopteromalus calandrae* (Howard) (Hymenoptera: Pteromalidae), a generalist parasitoid of stored product pests.

RESULTS: The LC_{50} of SeNPs was 1540 mg L^{-1} for female parasitoids and 1164 mg L^{-1} for males. SeNPs had a significant hormetic effect; average lifespan increased by 10% at a concentration of 4.03 mg L^{-1} for females and by 35% at 13.83 mg L^{-1} for males. In a bioassay including hosts (the azuki bean beetle, *Callosobruchus chinensis* (L.) (Coleoptera: Chrysomelidae: Bruchinae)), a low concentration of SeNPs (25 mg L^{-1}) enhanced the performance of female parasitoids; lifespan increased by 23% and the number of offspring increased by 88%. However, the number of emerging hosts did not significantly decrease; in the absence of parasitism, SeNPs actually improved host emergence by 17%.

CONCLUSION: Because higher concentrations of SeNPs reduced parasitoid lifespan, whereas low concentrations enhanced not only parasitoid performance but also host emergence, practitioners should exercise caution when considering SeNPs for use in integrated pest management.

Keywords: biological control, dose-response, hormesis, integrated pest management, nanotechnology, stored product pests

1. INTRODUCTION

The complementarity of chemical, cultural, biological, and other control strategies is a key aspect of integrated pest management (IPM)[1,2]. Because there tend to be compatibility issues between conventional insecticides and natural enemies[3–5], and due to the environmental and economic costs of these pesticides[6,7], there are ongoing efforts to counter an overreliance on chemical control and incorporate safer biorational alternatives to improve management outcomes and meet sustainable development goals (SDGs)[8–10]. Emerging research is also investigating the possible roles of nanotechnology in pest management, including the use of entomotoxic nanoparticles (NPs) or nano-encapsulated pesticides or botanicals[11,12–15]. Despite this effort, the ecological risks of NPs remain relatively unknown[11,16–18]. Due to the nonspecific mode of action of elemental (or elemental oxide) NPs[19], these products might pose a risk to nontarget organisms, including natural enemies.

Even though the sublethal effects of insecticides on natural enemies are typically negative[3,20], there are also instances in which they can be positive[21,22]. The term "hormesis" (or "hormoligosis" in insects[23]) refers to the positive or stimulatory effect of low doses of a chemical stressor, which can be characterized by a biphasic dose-response curve[24]. Since pesticide residues are often present throughout the agroecological landscape, it is important to understand how these (and other stressors) influence the biological control services provided by natural enemies[24,25]. Although there are many variables that affect the persistence of anthropogenic NPs in the environment (and their relationship to naturally occurring NPs)[18,26], relatively small or residual amounts of NPs might offer a direct benefit to biological control agents by hormesis.

Anisopteromalus calandrae (Howard) (Hymenoptera: Pteromalidae) is a cosmopolitan parasitoid that attacks a broad range of hosts, including the azuki bean beetle, *Callosobruchus chinensis* (L.), and the cowpea seed beetle, *C. maculatus* (F.) (Coleoptera: Chrysomelidae: Bruchinae)[27]. As solitary idiobiont ectoparasitoids, females typically lay one egg after paralyzing a host[28], and the developing parasitoid larva feeds from outside of its host; additional parasitoid-induced mortality can arise via host feeding[29], a process through which adult female *A. calandrae* attack and consume the hemolymph of hosts to acquire nutrients for egg maturation[29–31]. Due to the importance of *A. calandrae* in the control of bruchine beetles and other coleopteran pests[32–36], it is necessary to test their compatibility with the entomotoxic NPs that have garnered recent attention—selenium nanoparticles (SeNPs), in particular[37,38]—as a possible new control strategy for these insects. Furthermore, because the risks of nano-formulated insecticides are not as well studied as those of conventional insecticides[16–18] (and more controversial[39]), these tests on *A. calandrae* may serve as an indicator of their ecotoxicity, providing insight into both the specificity of SeNPs as well as their impact on the biological control subset of ecosystem services.

SeNPs have several potential applications in agriculture: promoting plant growth, bolstering plant

defenses, and acting as an antimicrobial, nematocidal, or insecticidal agent, depending on the

concentration or formulation[40–42]. The entomotoxic effects of SeNPs arise through the slow release of

elemental selenium, which, depending on the route of entry, may amass in the Malpighian tubules and

midgut, negatively impacting the growth, development, and survival of the target insect[43–45]. While SeNPs

have shown some promise as an alternative control method for bruchine beetles, including *C.*

chinensis[37,38], they may also harm the parasitoid *A. calandrae*[38]. However, there is evidence of SeNPs

eliciting a hormetic response in the male hosts of *A. calandrae*[37], so, considering the nonspecific mode of

action of SeNPs, we hypothesize that a similar result will be observed for the parasitoid. The purpose of

the present study is to investigate the concentration-dependent effects of SeNPs on the lifespan of the

generalist parasitoid *A. calandrae* and to test the hormetic effects of a low concentration of SeNPs on

parasitoid performance (lifespan, fecundity, offspring emergence time, age-dependent parasitism rate, and

host suppression), all of which will provide better guidance on the utilization of SeNPs and *A. calandrae*

in IPM.

2. METHODS

Colonies of *C. chinensis* (strain jC, originating from Japan) and its parasitoid *A. calandrae* (also from

Japan) have been maintained in controlled environmental chambers under standard laboratory conditions

(30°C, 60% R.H., 16:8 L:D) for more than 20 years on dried azuki beans, *Vigna angularis* var. *angularis*

(Willdenow) Ohwi & Ohashi (Fabaceae) (Daiwa Grain Co., Obihiro, Japan). The *C. chinensis* colonies

are maintained in large, ventilated Petri dishes (9.5 cm in diameter, 4 cm in height), whereas *A. calandrae*

is maintained in standard-sized Petri dishes (10 cm in diameter, 1.5 cm in height) supplied with *C.*

chinensis-infested azuki beans (16-day old pupae and late-instar larvae).

106   The chemical synthesis of SeNPs[46,47] (amorphous and approximately 5–10 and 60–100 nm in

diameter) was previously described by Miksanek & Tuda[37]; in brief, a 20-mL solution of SeNPs (1000 mg

L^{-1}) was prepared by reducing sodium selenite (43.8 mg) with ascorbic acid (2 mL, 0.633 M) in ultrapure

water (17.9 mL; Milli-Q, 18.2 MΩ·cm) and stabilizing with polysorbate 20 (100 μL). Additional SeNP

concentrations of 1, 10, 25, 50, 100, 200, and 500 mg L^{-1} were made by diluting the original solution with

ultrapure water; a 0 mg L^{-1} control solution (containing only ultrapure water) was also prepared. All

chemical reagents (sodium selenate, ascorbic acid, and polysorbate 20) were obtained from Nacalai

Tesque, Inc. (Kyoto, Japan).

2.1 Concentration-dependent toxicity

Newly emerged (< 24 h) adult parasitoids (males and females, separately) were placed individually in

mini-sized Petri dishes (35 mm in diameter, 10 mm in height). Each Petri dish has an air gap around the

lid that allows for ventilation. Next, a 20-µL droplet of SeNPs (0, 1, 10, 50, 100, 200, 500, or 1000 mg L^{-1}

(0–20 µg SeNPs)) was added to the center of the dish; by holding the volume constant and manipulating

the concentration of SeNPs, this changes the overall dose. Each Petri dish was gently agitated to evenly

cover all surfaces as well as the parasitoid with SeNPs. After this initial treatment, the parasitoids were

placed in a climate-controlled chamber (30°C, 60% R.H., 16:8 L:D) and monitored daily until death.

Parasitoids were not provided supplemental nutrients before, during, or after the bioassay. All eight SeNP

concentrations were replicated 25 times for each sex.

Note that we refer to a "replicate" as a repetition of the complete set of treatment combinations

(eight SeNP concentrations × two sexes), and, to reiterate, the parasitoids (400 in total) were subjected to

treatment combinations individually rather than as part of a group. Although toxicity testing on groups of

organisms is both common and convenient (with replication at the level of each set of groups), that

approach is problematic, especially for *A. calandrae* because female *A. calandrae* exhibit high levels of

density-dependent intraspecific interference competition[35,48,49]; if these parasitoids were exposed to SeNPs

in a group, then the resulting differential rates of death across groups treated with different concentrations

would introduce an increasing asymmetry in the behavioral interactions among the individuals in each

group over time, which would be a confounding variable.

2.2 Low-concentration SeNPs and parasitoid performance

The effects of a low concentration of SeNPs on the performance of *A. calandrae*, as well as offspring

emergence timeline, were tested by exposing parasitoids to infested azuki beans treated with either SeNPs

or water. The purpose of this was to evaluate the complementarity of SeNPs and biological control. The

experimental design consisted of a full factorial for parasitism (*A. calandrae* present or absent) and SeNP

application (0 or 25 mg L^{-1} SeNPs). For this bioassay, a concentration of 25 mg L^{-1} SeNPs was chosen

because a preliminary analysis of concentration-dependent survivorship suggested that this approximate

concentration may best enhance parasitoid performance.

Adult parasitoids were collected from a laboratory colony within 24 h of emergence. Next, each

male–female pair was treated with a 20-µL droplet of SeNPs (0.5 µg SeNPs) or ultrapure water (as

previously described) and provided eight azuki beans (with a total mass of 1.1 g, on average) treated

concurrently and in the same manner (by gently agitating each dish to evenly cover all surfaces as well as

the beans with SeNPs) with a 30-µL droplet[37] (0.75 µg SeNPs) (to account for the increased surface area).

Each bean was infested by eight mixed-stage *C. chinensis* larvae (indirectly counted as the number of

eggs on the surface of the bean; beans were visually inspected prior to use in the experiment, and those

with fewer or greater than eight hatched eggs were not used); by the time of parasitoid exposure, most

larvae were 15–16 days old, which is the favored host stage for parasitism; the smaller number of younger

larvae are the preferred size for host-feeding. After treatment, beans and parasitoids were immediately

placed in a small, untreated Petri dish (55 mm in diameter, 15 mm in height); the treated beans were not

allowed to dry beforehand. After 24 h, the male parasitoid was discarded to prevent excessive sexual

harassment. Parasitoid mortality was monitored at 24-h intervals. After four days, females were provided

with a fresh set of infested beans (of the same treatment conditions), and the original sets of beans were

checked daily to record the number and sex of emerging adults (hosts and parasitoids). The second set of

beans were not monitored for daily emergence, but the total number of emerged parasitoids was recorded

three weeks after the death of the parental parasitoid in order to calculate lifetime reproduction.

Parasitoids were not provided supplemental nutrients before or during the bioassay. Each combination of

treatments was replicated 30 times; daily emergence was only recorded from a subset of replicates (24

from each of the treatments in the absence of parasitism and 25 from each of those with a parasitoid

present).

2.3 Modeling and Analyses

The concentration-dependent effects of SeNPs on parasitoid lifespan (age-at-death, x) were described

using the Cedergreen–Ritz–Streibig modification of a log-logistic model, which allows for an effect of

hormesis[50,51]. This model takes the form:

$$f(x) = \frac{d + f\exp(-1/x^{\alpha})}{1 + \exp(b(\log(x) - \log(e)))} \tag{1}$$

in which d is the upper horizontal asymptote and f is the magnitude of the effect of hormesis (with α

affecting the shape of the peak); the parameters b and e have no direct biological interpretation (in the

related three-parameter log-logistic model, b is a scaling factor and e is an inflection point that

corresponds with the EC_{50} or LC_{50})[50,51]. If the parameter f is equal to 0, then there is no hormetic effect

and the model simplifies to a three-parameter log-logistic curve[50,51]. Model fitting and analysis was

conducted in R version 4.2.2. The term α was set to 0.5 when fitting the model[50,51]. Parasitoid sex was

included as a grouping factor when estimating the parameters b and d (the model was simplified because

preliminary analyses showed no significant effect of sex on estimates of e and f). The package *drc* was

used in the analysis, which also allowed for estimating the maximum effect of hormesis from the mean

response predicted by the model (using the "MAX" function[51], as described by Cedergreen et al.[50]), lethal

concentrations (e.g. LC_{50}), and calculating t-statistics for comparisons between sexes[50,51].

The effects of SeNPs on the lifespan and fecundity of adult female parasitoids were tested with a
Cox proportional hazards model and linear regression model, respectively. A binomial GLM was used to
test the effects of SeNP treatment on the sex ratio of the resulting parasitoid offspring. The effects of
SeNPs on the development time of offspring (days to adult emergence) were assessed with a Cox
proportional hazards model, with sex and the interaction between sex and SeNP treatment as additional
factors; the identity of the maternal parasitoid was included as a random effect by clustering.

To further investigate the reproduction of female parasitoids, a linear regression was used to
assess the relationship between the log-transformed early rate of reproduction (the age-corrected number
of offspring per day, calculated as the number of offspring produced from days 0–3 (inclusive) divided by
the lifespan of the parasitoid) and SeNP treatment on the reproductive rate later in life (from day 4–death,
calculated in the same fashion); only parasitoids with a lifespan greater than four days and a nonzero early
reproductive rate were included in this analysis. Next, to investigate the influence of SeNPs on the
lifetime reproductive success (total number of offspring) of female parasitoids as a function of age, an
asymptotic Michaelis–Menten-type model was fit to the data. This model predicts the total number of
offspring of a female parasitoid living to age t as:

$$f(t) = \frac{m(t - t_{\mathrm{pre}})}{\beta \exp(-\gamma[\mathrm{Se}]) + t - t_{\mathrm{pre}}} \tag{2}$$

Analogous to the standard Michaelis-Menten model, the parameter m denotes the upper asymptote of the
saturating curve and β is associated with the midpoint of the saturating curve; biologically, m represents
the maximum number of parasitoid offspring (limited by host depletion or as a more general effect of
parasitoid senescence). The modifications on the Michaelis-Menten model include an x-intercept (t_{pre}) and
the effect of SeNPs ($\exp(-\gamma[\mathrm{Se}])$). The term t_{pre} applies a horizontal transformation to the curve;
biologically, t_{pre} represents the length of the pre-oviposition period of the adult female (a female might
use this time for mating, sperm storage, host feeding, or egg maturation). The function $\exp(-\gamma[\mathrm{Se}])$
applies an effect γ of SeNPs ([Se] = 0 or 25 mg L^{-1}) to β; thus, at a concentration of 0 mg L^{-1} (or if there
is no effect γ), then β is unaffected. For simplicity, the Michaelis–Menten-type model only considers the
effect of SeNPs on the parameter β; however, this formulation was also supported by an informal analysis
of the shortest- and longest-lived parasitoids (0–3 and 9–12 days, respectively), in which were there no
differences in the reproduction of control and SeNP-treated parasitoids in either age class, suggesting that
m and t_{pre} are likely unaffected by SeNP treatment.

Finally, the effects of SeNPs and parasitism on the number and sex ratio of emerging hosts were
tested with a linear mixed-effects model (LMM) and binomial generalized mixed-effects model (GLMM),

respectively. In both models, the identity of the source colony was included as a random effect to account

for potential variation in the stage structure or exact number of hosts; because source colony did not

improve the statistical models of parasitoid life history (described above), this random effect was only

included in these analyses of the host population.

All modeling and statistical analyses were conducted in R version 4.2.2. The Michaelis–Menten-

type model was fit using nonlinear least squares; confidence intervals for model predictions were

calculated by Monte Carlo simulation ($n = 100000$). The packages *car*, *lme4*, *merTools*, *survival*, *coxed*,

and *propagate* were also used in the analyses.

3. RESULTS

3.1 Concentration-dependent toxicity

The lifespan of adult parasitoids was measured in response to a range of SeNP concentrations (0–1000 mg

L^{-1}). In fitting the Cedergreen–Ritz–Streibig-modified log-logistic model to the concentration-dependent

effects of SeNPs on the lifespan of male ($n = 200$) and female ($n = 200$) parasitoids (with parasitoid sex as

a grouping factor when estimating b and d), the estimates for all parameters were significant (Figure 1,

Table 1). The upper asymptote d—which represents the maximum lifespan in the absence of SeNPs—was

4.55 ± 0.15 d (estimate $\pm$ SE) for females and 3.01 ± 0.16 d for males, and the difference between sexes

was significant ($t = 6.27$, $p < 0.001$) (Figure 1; Table 1); the estimates of b (a scaling factor) were

marginally different between sexes ($t = -1.91$, $p = 0.057$) (Table 1).

The LC$_{50}$ was 1540 ± 391 mg L^{-1} for females (estimate $\pm$ SE) and 1164 ± 224 mg L^{-1} for males,

but there was no significant difference in the estimates of LC$_{50}$ between sexes ($t = 1.19$, $p = 0.237$). For

hormetic effects, the estimated age at death of SeNP-treated females was 5.02 d at a concentration of 4.03

mg L^{-1}; and the estimated age at death of SeNP-treated males was 4.06 d at a concentration of 13.83 mg

L^{-1} (Figure 1; Table 1).

3.2 Low-concentration SeNPs and parasitoid performance

Parasitoid performance was assessed in response to a low concentration (25 mg L^{-1}) of SeNPs to

investigate the effects of hormesis on a generalist biological control agent. Adult female parasitoids ($n =$

60) treated with 25 mg L^{-1} SeNPs lived significantly longer than control parasitoids; adult lifespan was

7.53 ± 0.52 and 6.10 ± 0.48 d, respectively (mean $\pm$ SE) (Figure 2a; Table 2). SeNP-treated females also

produced a greater number of offspring; the total number of offspring was 13.0 ± 1.1 vs. 6.9 ± 1.1 (Figure

2b; Table 2). The sex ratio ($n = 56$) and development time ($n = 285$) of the offspring of SeNP-treated

parasitoids did not differ from those of the control (Table 2). Regardless of SeNP treatment, female

offspring (n = 233) emerged later than males (n = 52); the time to emergence was 13.66 ± 0.09 and 12.73

± 0.18 d, respectively (Table 2).

For ovipositing female parasitoids with an age of death > 4 d (n = 47), the early reproductive rate

was inversely correlated with the rate of reproduction later in life (offspring per day, days 4–death) ($F_{3,43}$

= 14.94, multiple R^2 = 0.510, p < 0.001) (Figure 3a). As the early rate of reproduction increased, there

was a decrease in the rate of reproduction later in life; treatment with 25 mg L^{-1} SeNPs increased the rate

of reproduction overall (Figure 3a; Table 3a). To investigate the lifetime reproductive success of all

female parasitoids (n = 60), a Michaelis–Menten-type model was implemented, in which all parameter

estimates were significant (Figure 3b; Table 3b). The maximum number of offspring m was 44.7 ± 17.8

(estimate ± SE) and the preoviposition period was 1.90 ± 0.30 d (Table 3b). There was also a significant

effect of SeNP treatment γ, with 25 mg L^{-1} SeNPs increasing the lifetime reproductive success of female

parasitoids (Figures 3a and 3b; Tables 3a and 3b).

Finally, the effects of SeNPs and parasitism on the host population were evaluated as the number

and sex ratio of the emerging hosts in each Petri dish (n = 120). The number of emerging hosts was lower

in the presence of the parasitoid than in the absence of the parasitoid but, when treated with SeNPs, was

higher in the absence of parasitism (Figure 4a, Table 4). When treated with 0 mg L^{-1} SeNPs, there were

an estimated 21.5 (95% prediction interval: 15.0–28.8) emerging hosts per Petri dish in the absence of

parasitism and 19.1 (12.3–26.1) emerging hosts per dish in the presence of a female parasitoid; when

treated with 25 mg L^{-1} SeNPs, there were 25.1 (18.6–31.4) emerging hosts per dish in the absence of

parasitism and 17.1 (10.9–23.4) emerging hosts with a parasitoid present (Figure 4a, Table 4). Finally, the

sex ratio of the emerging hosts shifted from a male bias (0.54 [0.51–0.57]) at 0 mg L^{-1} SeNPs to a slight

female bias (0.47 [0.45–0.51]) when treated with 25 mg L^{-1} SeNPs, independent of parasitoid presence

(Figure 4b, Table 4).

4. DISCUSSION

Because of the environmental and economic costs of conventional chemical insecticides, there is ongoing

interest in exploring alternative management tactics, including the use of nanotechnologies. Selenium

nanoparticles (SeNPs) have many applications in agriculture, but their nonspecific entomotoxic mode of

action may limit their compatibility with biological control agents; however, low concentrations may

benefit natural enemies via hormesis. The present study found that, for the generalist parasitoid *A.*

calandrae, the LC$_{50}$ of SeNPs was 1540 mg L^{-1} for female parasitoids and 1164 mg L^{-1} for males, and an

analysis of the concentration-response curves identified a significant hormetic effect; for females, the

estimated age at death increased by 10% at a concentration of 4.03 mg L^{-1} and, for males, increased by

35% at 13.83 mg L^{-1}. In a follow-up bioassay that provided female parasitoids with hosts, the low

concentration of SeNPs was found to enhance parasitoid performance; female lifespan increased by 23% and the number of offspring increased by 88%. Regardless, the number of emerging hosts (*C. chinensis*) did not significantly decrease after treatment with low-concentration SeNPs but rather, in the absence of parasitism, increased by 17%. This indicates that, while a low concentration of SeNPs did enhance parasitoid performance, there was no net effect on host suppression because *C. chinensis* responded positively in parallel.

There exists a delicate balance in the essential role of selenium as a micronutrient and its dose-dependent toxicity in both vertebrate and invertebrate animals[40–45,52–56]. In insects, diets with supplemental selenium can benefit growth, survival, and reproduction, but higher concentrations can have the opposite effect[40,43–45,53]. While insects play a role in the biogeochemical cycling of selenium[54,56], little is known of the impact of selenium on predator–prey or host–parasitoid ecology; the biotransfer or biomagnification[57] of selenium can negatively affect the development and fitness of predators[58], and higher concentrations of selenium can potentially disrupt host–parasitoid population dynamics, with *Cotesia marginiventris* (Cresson) (Hymenoptera: Braconidae) and the beet armyworm, *Spodoptera exigua* (Hubner) (Lepidoptera: Noctuidae), serving as an example[56,59]. Even though SeNPs have a lower toxicity and higher bioavailability than inorganic or organic forms of selenium[42], applications of SeNPs ranging in concentration from 100–500 mg L^{-1} can still have significant entomotoxic effects[42,60], including on *C. chinensis*[37,61]. However, in *C. chinensis*, endosymbiotic bacteria can mediate resistance to SeNPs[37]; *Wolbachia* provides females with resistance to low concentrations of SeNPs, whereas higher concentrations had an entomotoxic (but not insecticidal) effect regardless of infection status, decreasing life expectancy and reducing fecundity[37].

In the present study of the *C. chinensis–A. calandrae* system, the parasitoid *A. calandrae* also began showing adverse effects within the 100–500 mg L^{-1} range. Thus, the concurrent use of SeNPs and biological control would not be compatible in this case. Similarly, since the pest species also appears to exhibit a hormetic response to a low concentration of SeNPs, this limits the complementary of SeNPs and biological control, at least in the short term; additional studies that extend beyond the single-parasitoid, single-generation paradigm (and that also address the timing of SeNP applications and residual effects), would be required to identify any long-term consequences that might be relevant for pest suppression. But first, there is the fundamental question of whether the performance of *A. calandrae* is enhanced as a result of a response to SeNPs as a chemical stressor (i.e. hormoligosis/hormesis), via the potential benefits of selenium as a micronutrient (parasitoids could be observed drinking from dispersed droplets of SeNPs [or water] before the application completely dried, although this would not offer the same benefits as supplemental honey or sugar[30,62]), or through some combination of the two.

The positive effects of a low concentration of SeNPs on lifespan (and fecundity) were more

prominent when hosts were available for host feeding and oviposition, which suggests that SeNPs may at

least alter the nutritional physiology of female parasitoids (and, if linked to host feeding, this may

introduce an additional source of density dependence in the system[63,64]). However, because selenoproteins

have largely been lost in the related parasitoid *Nasonia vitripennis* (Walker) (Hymenoptera: Pteromalidae)

and the honey bee *Apis mellifera* L. (Hymenoptera: Apidae)[65,66], the positive effects of low-concentration

SeNPs are more likely to be a hormetic response to small amounts of oxidative stress or via a reduced

stress threshold rather than as a positive response to an essential micronutrient[67–69]. Regardless, because

most Coleoptera also lack selenoproteins[65,66,69], a hymenopteran parasitoid and its coleopteran host might

be expected to exhibit similar physiological responses to SeNPs, which could complicate control efforts.

5. CONCLUSION

Because higher concentrations of SeNPs reduced parasitoid lifespan, the concurrent use of SeNPs and

biological control do not appear to be compatible tactics in the IPM of *C. chinensis*. This is an important

finding because it offers insight into the potential ecotoxicity of SeNPs and it also raises concerns of the

nonspecific mode of action of SeNPs, which would be antithetical to their purpose as an alternative to

conventional chemical controls. Additionally, because a low-concentration treatment of SeNPs benefitted

not only the parasitoid but the host as well, SeNPs and biological control may not be complementary

tactics, either. Future studies on the residual effects of SeNP applications, on the trade-offs between

parasitoid lifespan and reproduction, or on population-level responses (extending beyond single-

parasitoid, single-generation experiments) might offer additional insight. It remains possible that, within

some broader context, SeNPs might still have a role in IPM or biological control.

6. ACKNOWLEDGEMENTS

6.1 Author contributions

All authors contributed to the study conception and design. Material preparation and data collection were

performed by JRM and CA. JRM and MT participated in data analysis. The first draft of the manuscript

was written by JRM, and all authors commented on subsequent versions of the manuscript. All authors

read and approved the final manuscript.

6.2 Data availability

The datasets and R script associated with the current study are available from the primary corresponding

author on request.

6.3 Funding

JRM is an International Research Fellow of Japan Society for the Promotion of Science (JSPS). CA was

supported by an Invitational Fellowship from JSPS. MT received additional support by KAKENHI

(19K06840) from JSPS.

7. CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest and affirm that the present study was carried out in adherence

to all applicable ethical standards.

8. REFERENCES

1. Kogan M, Integrated pest management: Historical perspectives and contemporary developments. *Annu Rev Entomol* 43:243–270 (1998).

2. Farrar JJ, Ellsworth PC, Sisco R, Baur ME, Crump A, Fournier AJ, et al., Assessing the compatibility of a pesticide in an IPM program. *J Integr Pest Manag* 9:3 (2018).

3. Desneux N, Decourtye A, Delpuech JM, The sublethal effects of pesticides on beneficial arthropods. *Annu Rev Entomol* 52:81–106 (2007).

4. Beers EH, Mills NJ, Shearer PW, Horton DR, Milickzy ER, Amarasekare KG, et al., Nontarget effects of orchard pesticides on natural enemies: Lessons from the field and laboratory. *Biol Control* 102:44–52 (2016).

5. Schmidt-Jeffris RA, Beers EH, Sater C, Meta-analysis and review of pesticide non-target effects on phytoseiids, key biological control agents. *Pest Manag Sci* 77:4848–4862 (2021).

6. Pimentel D, Environmental and economic costs of the application of pesticides primarily in the United States. *Environ Dev Sustain* 7:229–252 (2005).

7. Damalas CA, Eleftherohorinos IG, Pesticide exposure, safety issues, and risk assessment indicators. *Int J Environ Res Public Health* 8:1402–1419 (2011).

8. Rosell G, Quero C, Coll J, Guerrero A, Biorational insecticides in pest management. *J Pestic Sci* 33:103–121 (2008).

9. Dangles O, Casas J, Ecosystem services provided by insects for achieving sustainable development goals. *Ecosyst Serv* 35:109–115 (2019).

10. Struelens Q, Silvie P, Orienting insecticide research in the tropics to meet the sustainable development goals. *Curr Opin Insect Sci* 40:24–30 (2020).

11. Kah M, Hofmann T, Nanopesticide research: Current trends and future priorities. *Environ Int* 63:224–235 (2014).

12. Athanassiou CG, Kavallieratos NG, Benelli G, Losic D, Usha Rani P, Desneux N, Nanoparticles for pest control: Current status and future perspectives. *J Pest Sci* 91:1–15 (2018).

13. Sciortino M, Scurria A, Lino C, Pagliaro M, D'Agostino F, Tortorici S, et al., Silica-miroencapsulated orange oil for sustainable pest control. *Adv Sustainable Sys* 5:2000280 (2021).

14. Ricupero M, Biondi A, Cincotta F, Condurso C, Palmeri V, Verzera A, et al., Bioactivity and physico-chemistry of essential oil nanoemulsion in tomato. *Entomol Gen* 42:921–930 (2022).

15. Tortorici S, Cimino C, Ricupero M, Musumeci T, Biondi A, Siscaro G, et al., Nanostructured lipid carriers of essential oils as potential tools for the sustainable control of insect pests. *Ind Crops Prod* 181:114766 (2022).

16. Crane M, Handy RD, Garrod J, Owen R, Ecotoxicity test methods and environmental hazard
assessment for engineered nanoparticles. *Ecotoxicology* 17:421–437 (2008).

17. Hansen SF, Michelson ES, Kamper A, Borling P, Stuer-Lauridsen F, Baun A, Categorization
framework to aid exposure assessment of nanomaterials in consumer products. *Ecotoxicology*
17:438–447 (2008).

18. Wagner S, Gondikas A, Neubauer E, Hofmann T, von der Kammer F, Spot the difference:
Engineered and natural nanoparticles in the environment—release, behavior, and fate. *Angew Chem*
Int Ed 53:12398–12419 (2014).

19. Benelli G, Mode of action of nanoparticles against insects. *Environ Sci Pollut Res* 25:12329–12341
(2018).

20. Stark JD, Vargas R, Banks JE, Incorporating ecologically relevant measures of pesticide effect for
estimating the compatibility of pesticides and biocontrol agents. *J Econ Entomol* 100:1027–1032
(2007).

21. Cutler GC, Amichot M, Benelli G, Guedes RNC, Qu Y, Rix RR, et al., Hormesis and insects: Effects
and interactions in agroecosystems. *Sci Total Environ* 825:153899 (2022).

22. Wang X, Tian L, Ricupero M, Harwood JD, Liang Y, Zang LS, Wang S, Hormesis effects of
chlorantraniliprole on a key egg parasitoid used for management of rice lepidopterans. *Entomol Gen*
42:941–948 (2022).

23. Luckey TD, Insecticide hormoligosis. *J Econ Entomol* 61:7–12 (1968).

24. Cutler GC, Insects, insecticides, and hormesis: Evidence and considerations for study. *Dose–*
Response 11:154–177 (2013).

25. Guedes RNC, Cutler GC, Insecticide-induced hormesis and arthropod pest management. *Pest Manag*
Sci 70:690–697 (2013).

26. Malakar A, Kanel SR, Ray C, Snow DD, Nadagouda MN, Nanomaterials in the environment, human
exposure pathway, and health effects: A review. *Sci Total Environ* 759:143470 (2021).

27. Zhang Y, Yu F, Wu L, Dai R, Yang H, Zhang X, Hu D, Life history traits of the parasitoid
Anisopteromalus calandrae (Hymenoptera: Pteromalidae) on three beetle hosts. *J Stored Prod Res*
97:101973 (2022).

28. Perkin LC, Friesen KS, Flinn PW, Oppert B, Venom gland components of the ectoparasitoid wasp,
Anisopteromalus calandrae. *J Venom Res* 6:19–37 (2015).

29. Choi WI, Yoon TJ, Ryoo MI, Host-size-dependent feeding behavior and progeny sex ratio of
Anisopteromalus calandrae (Hym., Pteromalidae). *J Appl Entomol* 125:71–77 (2001).

30. Schmale I, Wäckers FL, Cardona C, Dorn S, Control potential of three hymenopteran parasitoid
species against the bean weevil in stored beans: The effect of adult parasitoid nutrition on longevity
and progeny production. *Biol Control* 21:134–139 (2001).

31. Menon A, Flinn PW, Dover BA, Influence of temperature on the functional response of
Anisopteromalus calandrae (Hymenoptera: Pteromalidae), a parasitoid of *Rhyzopertha dominica*
(Coleoptera: Bostrichidae). *J Stored Prod Res* 38:463–469 (2002).

32. Clausen CP, Introduced Parasites and Predators of Arthropod Pests and Weeds: A World Review.
United States Department of Agriculture Handbook (1978).

33. Van Huis A, Biological methods of bruchid control in the tropics: A review. *Insect Sci Appl* 12:87–
102 (1991).

34. Tuda M, Chou L-Y, Niyomdham C, Buranapanichpan S, Tateishi Y, Ecological factors associated
with pest status in *Callosobruchus* (Coleoptera: Bruchidae): High host specificity of non-pests to
Cajaninae (Fabaceae). *J Stored Prod Res* 41:31–45 (2005).

35. Tuda M, Shimada M, Complexity, evolution, and persistence in host–parasitoid experimental systems
with *Callosobruchus* beetles as the host. *Adv Ecol Res* 37:37–75 (2005).

36. Tuda M, Rönn J, Buranapanichpan S, Wasano N, Arnqvist G, Evolutionary diversification of the bean
beetle genus *Callosobruchus* (Coleoptera: Bruchidae): Traits associated with stored-product pest
status. *Mol Ecol* 15:3541–3551 (2006).

37. Miksanek JR, Tuda M, Endosymbiont-mediated resistance to entomotoxic nanoparticles and sex-
specific responses in a seed beetle. *J Pest Sci* 96:1257–1270 (2023).

38. San PP, Zhang YZ, Helmy EAM, Adarkwah C, Tuda M, Sex-dependent effects of biosynthesized
nanoparticles on stored bean pest beetles and their non-target parasitoid. *Crop Prot* 173:106352
(2023).

39. Ale A, Andrade VS, Gutierrez MF, Bacchetta C, Rossi AS, Orihuela PS, et al., Nanotechnology-
based pesticides: Environmental fate and ecotoxicity. *Toxicol Appl Pharmacol* 471:116560 (2023).

40. El-Ramady HR, Domokos-Szabolcsy É, Abdalla NA, Alshaal TA, Shalaby TA, Sztrik A, et al.,
Selenium and nano-selenium in agroecosystems. *Environ Chem Lett* 12:495–510 (2014).

41. Mechora Š, Selenium as a protective agent against pests: A review. *Plants* 8:262 (2019).

42. Garza-García JJO, Hernández-Díaz JA, Zamudio-Ojeda A, León-Morales JM, Guerrero-Guzmán A,
Sánchez-Chiprés DR, et al., The role of selenium nanoparticles in agriculture and food technology.
Biol Trace Elem Res 200:2528–2548 (2022).

43. Simmons TW, Jamall IS, Lockshin RA, Accumulation, distribution and toxicity of selenium in the
adult house fly, *Musca domestica*. *Comp Biochem Physiol* 91: 559–563 (1988).

44. Hogan GR, Razniak HG, Selenium-induced mortality and tissue distribution studies in *Tenebrio*
molitor (Coleoptera: Tenebrionidae). *Environ Entomol* 20:790–794 (1991).

45. Lalitha K, Rani P, Narayanaswami V, Metabolic relevance of selenium in the insect *Corcyra*
cephalonica: Uptake of ^{75}Se and subcellular distribution. Biol Trace Elem Res 41:217–233 (1994).

46. Bartůněk V, Junková J, Šuman J, Kolářová K, Rimpelová S, Ulbrich P, Sofer Z, Preparation of
amorphous antimicrobial selenium nanoparticles stabilized by odor suppressing surfactant
polysorbate 20. *Mater Lett* 152:207–209 (2015).

47. Vahdati M, Tohidi Moghadam T, Synthesis and characterization of selenium nanoparticles-lysozyme
nanohybrid system with synergistic antibacterial properties. *Sci Rep* 10:510 (2020).

48. Nakamichi Y, Tuda M, Wajnberg E, Intraspecific interference between native parasitoids modified by
a non-native parasitoid and its consequence on population dynamics. *Ecol Entomol* 45:1263–1271
(2020).

49. Zhang YY, Jin Z, Miksanek JR, Tuda M, Impact of nonnative parasitoid species on intraspecific
interference and offspring sex ratio. *Sci Rep* 11:23215 (2021).

50. Cedergreen N, Ritz C, Streibig JC, Improved empirical models describing hormesis. *Environ Toxicol*
Chem 24:3166–3172 (2005).

51. Ritz C, Baty F, Streibig JC, Gerhard D, Dose-response analysis using R. *PLoS One* 10:e0146021
(2015).

52. Golombieski RM, Graichen DÂS, da Rocha JBT, da Silva Valente VL, da Silva Loreto ÉL, Over-
activation of the *Drosophila melanogaster hsp83* gene by selenium intoxication. *Genet Mol Biol*
31:128–135 (2008).

53. Martin-Romero FJ, Kryukov GV, Lobanov AV, Carlson BA, Lee BJ, Gladyshev VN, et al., Selenium
metabolism in *Drosophila*. *J Biol Chem* 276:29798–29804 (2001).

54. Golubkina N, Sheshnitsan S, Kapitalchuk M, Ecological importance of insects in selenium biogenic
cycling. *Int J Ecol* 2014:835636 (2014).

55. Reich HJ, Hondal RJ, Why nature chose selenium. *ACS Chem Biol* 11:821–841 (2016).

56. Vickerman DB, Trumble JT, George GN, Pickering IJ, Nichol H, Selenium biotransformations in an
insect ecosystem: Effects of insects on phytoremediation. *Environ Sci Technol* 38:3581–3586 (2004).

57. Bañuelos GS, Vickerman DB, Trumble JT, Shannon MC, Davis CD, Finley JW, et al., Biotransfer
possibilities of selenium from plants used in phytoremediation. *Int J Phytoremediation* 4:315–329
(2002).

58. Vickerman DB, Trumble JT, Biotransfer of selenium: Effects on an insect predator, *Podisus*
maculiventris. *Ecotoxicology* 12:497–504 (2003).

59. Butler CD, Trumble JT, Predicting population dynamics of the parasitoid *Cotesia marginiventris*
(Hymenoptera: Braconidae) resulting from novel interactions of temperature and selenium.
Biocontrol Sci Technol 20:391–406 (2010).

60. Salem SS, Fouda MMG, Fouda A, Awad MA, Al-Olayan EM, Allam AA, et al., Antibacterial,
cytotoxicity and larvicidal activity of green synthesized selenium nanoparticles using *Penicillium*
corylophilum. *J Clust Sci* 32:351–361 (2021).

61. Helmy EAM, San PP, Zhang YZ, Adarkwah C, Tuda M, Entomotoxic efficacy of fungus-synthesized
nanoparticles against immature stages of stored bean pests. *Sci Rep* 13:8508 (2023).

62. Wäckers FL, The effect of food supplements on parasitoid–host dynamics. *Proceedings of the First*
International Symposium on Biological Control of Arthropods, Honolulu, Hawaii, pp. 226-231
(2003).

63. Miksanek JR, Heimpel GE, A matrix model describing host–parasitoid population dynamics: The
case of *Aphelinus certus* and soybean aphid. *PLoS One* 14:e0218217 (2019).

64. Miksanek JR, Heimpel GE, Density-dependent lifespan and estimation of life expectancy for a
parasitoid with implications for population dynamics. *Oecologia* 194:311–320 (2020).

65. Chapple CE, Guigó R, Relaxation of selective constraints causes independent selenoprotein extinction
in insect genomes. *PLoS One* 3:e2968 (2008).

66. Lobanov AV, Hatfield DL, Gladyshev V, Selenoproteinless animals: Selenophosphate synthetase
SPS1 functions in a pathway unrelated to selenocysteine biosynthesis. *Protein Sci* 17:176–182
(2008).

67. Alburaki M, Smith KD, Adamczyk J, Karim S, Interplay between selenium, selenoprotein genes, and
oxidative stress in the honey bee *Apis mellifera* L. *J Insect Physiol* 117:103891 (2019).

68. Shchedrina VA, Kabil H, Vorbruggen G, Lee BC, Turanov AA, Hirosawa-Takamori M, et al.,
Analyses of fruit flies that do not express selenoproteins or express the mouse selenoprotein,
methionine sulfoxide reductase B1, reveal a role of selenoproteins in stress resistance. *J Biol Chem*
286: 29449 – 29461 (2011).

69. Santesmasses D, Mariotti M, Gladyshev VN, Bioinformatics of selenoproteins. *Antioxid Redox Signal*
33:525–536 (2020).

Table 1. Parameter estimates for the Cedergreen–Ritz–Streibig-modified log-logistic model for the

concentration-dependent effect of selenium nanoparticles (0–1000 mg L^{-1}) on the lifespan of the

parasitoid *Anisopteromalus calandrae* in the absence of hosts.

Parameter	Grouping (sex)	Estimate	SE	*t*-value	*p*-value
b	Female	0.47	0.09	5.29	< 0.001
	Male	0.84	0.16	5.24	< 0.001
d (upper asymptote)	Female	4.55	0.15	29.60	< 0.001
	Male	3.01	0.16	19.21	< 0.001
e	–	494.80	118.95	4.16	< 0.001
f (effect of hormesis)	–	1.63	0.44	3.73	< 0.001

518   **Table 2.** Statistical tests of the effects of selenium nanoparticles (25 mg L^{-1}) on the life history

519   parameters of adult female parasitoids (*Anisopteromalus calandrae*) and offspring.

Response (Model)	Predictor	df	χ^2	*p*-value
Age at death (Cox prop. hazards)	SeNP treatment	1	4.27	0.039
Reproduction (linear regression)	SeNP treatment	1	16.24	< 0.001
Sex ratio (binomial GLM)	SeNP treatment	1	0.89	0.346
Development time (Cox prop. hazards)	Sex	1	18.38	< 0.001
	SeNP treatment	1	0.03	0.856
	Sex × SeNP treatment	1	0.02	0.884

521   **Table 3a.** Correlation between early and late reproduction (number of offspring per day) by female

522   *Anisopteromalus calandrae* and the effects of selenium nanoparticles (25 mg L^{-1}).

Predictor	df	χ^2	*p*-value
Early reproductive rate [log(age-adjusted number of offspring per day)]	1	42.97	< 0.001
SeNP treatment	1	6.32	0.016
Early reproductive rate × SeNP treatment	1	0.00	0.946

524   **Table 3b.** Parameter estimates for the Michaelis–Menten-type model of the lifetime reproductive success

525   of selenium nanoparticle-treated (25 mg L^{-1}) female parasitoids (*Anisopteromalus calandrae*).

Parameter	Estimate	SE	*t*-value	*p*-value
Maximum number of parasitized hosts, m	44.72	17.85	2.51	< 0.001
β	21.94	11.91	1.84	< 0.001
Preoviposition period, t_{pre}	1.90	0.30	6.35	< 0.001
Effect of SeNP treatment, γ	0.02	0.01	3.62	< 0.001

527 **Table 4.** Statistical tests of the effects of the parasitoid *Anisopteromalus calandrae* and selenium

528 nanoparticles (25 mg L^{-1}) on the host/pest (*Callosobruchus chinensis*) population.

Response (Model)	Predictor	df	χ^2	*p*-value
Host emergence (LMM)	Parasitoid presence	1	26.91	< 0.001
	SeNP treatment	1	0.61	0.436
	Parasitoid presence × SeNP treatment	1	8.14	0.004
Host sex ratio (binomial GLMM)	Parasitoid presence	1	0.06	0.808
	SeNP treatment	1	9.22	0.002
	Parasitoid presence × SeNP treatment	1	0.13	0.715

Figure 1. Concentration-dependent lifespan of the parasitoid *Anisopteromalus calandrae* (adult lifespan, i.e. age at death) treated individually with different concentrations of selenium nanoparticles (SeNPs) in the absence of hosts; the solid line plots the Cedergreen–Ritz–Streibig-modified log-logistic model (with 95% CI); points plot observed values (jittered for clarity).

Figure 2. Lifespan and fecundity of adult female *Anisopteromalus calandrae* as a function of selenium
nanoparticle (SeNP) treatment (0 or 25 mg L^{-1}). **(a)** Lifespan of adult female parasitoids; the solid line
indicates the average age at death. **(b)** Fecundity (total number of offspring); the solid line plots the linear
regression model (with 95% CI). In both panels, points plot observed values (jittered for clarity).

Figure 3. Reproduction of the generalist parasitoid *Anisopteromalus calandrae* as a function of age and selenium nanoparticle (SeNP) treatment (0 or 25 mg L^{-1}). **(a)** Correlation between the early and late reproductive rates of female parasitoids (number of offspring per day [age-adjusted], days 0–3 vs. day 4– death); the solid line plots the linear regression model (with 95% CI). **(b)** Lifetime reproduction, or the total number of offspring, per female; solid lines plot the Michaelis-Menten-type models (with 95% CI). In both panels, points plot observed values (jittered for clarity in panel b).

Figure 4. Influence of parasitism and selenium nanoparticle (SeNP) treatment (0 or 25 mg L^{-1}) on the host/pest population. **(a)** Escaped hosts, or the number of emerging adult *Callosobruchus chinensis*; solid lines plot the population-level predictions from the LMM (with 95% prediction interval). **(b)** Sex ratio of emerging beetles; the solid line plots the population-level predictions from the binomial GLMM (with 95% prediction interval). In both panels, points plot observed values (jittered for clarity).