Non-senescent species are not immortal: stress and decline in two planaria species

Jacques A. Deere[1,2]*, Penelope Holland[1,3], Aziz Aboobaker[1] & Roberto Salguero-Gómez[1,4,5]*

[1] Department of Biology, 11a Mansfield Rd, University of Oxford, Oxford, OX1 3SZ, United Kingdom.

[2] Evolutionary and Population Biology Department, Institute for Biodiversity and Ecosystem Dynamics (IBED), University

of Amsterdam,

P.O. Box 94240, 1090 GE Amsterdam, The Netherlands

[3] Division of Biology and Medicine, Brown University, Providence, RI 02912, USA.

[4] Centre for Biodiversity and Conservation Science, University of Queensland, St Lucia 4071QLD, Australia.

[5] Evolutionary Demography Laboratory, Max Plank Institute for Demographic Research, Rostock 18057, Germany.

*Corresponding authors:

Institute for Biodiversity and Ecosystem Dynamics (IBED), University of Amsterdam, P.O. Box 94249, 1090 GE

Amsterdam, The Netherlands

Department of Biology, University of Oxford, South Parks Road, OX1 3RB, Oxford, UK

E-mail addresses:

Jacques A. Deere (corresponding author):   j.a.deere@uva.nl

Penelope Holland: penelope_celeste_holland@brown.edu

Aziz Aboobaker: aziz.aboobaker@biology.ox.ac.uk

Roberto Salguero-Gómez (corresponding author): rob.salguero@biology.ox.ac.uk

Acknowledgements

This project was funded by an Oxford John Fell Funds grant to RSG. RSG was supported by a NERC

Independent Research Fellowship (NE/M018458/1).

Conflict of Interest Statement

The authors have no conflict of interest to declare.

Author Contributions

RSG conceived the idea; JAD, RSG, and AA designed the methodology; JAD and PH collected and

digitized data; JAD analysed data; JAD led the writing of the manuscript with assistance from RSG.

All authors contributed critically to the drafts and gave final approval for publication.

Data Availability

The data in this paper are published as supporting information in the open access version deposited

in bioRxiv.org (DOI: 10.1101/2023.01.12.523873; Deere et al. 2022) (data are in multiple formats;

e.g., .pdf, .csv, .cpproj).

ABSTRACT

1. Potential immortality is observed in several species (*e.g.*, prickly pear cactus, hydra, flatworms) and is indicative of their negligible or even negative senescence rates. Unlike in senescent species, which experience reduced individual performance with age due to physiological degradation, species with negligible or negative senescence display mortality rates that remain constant or decline with age, respectively. These rates vary across taxa and are correlated with life history traits. Yet, the extent to which variable resource availability, a key driver of variation in life history traits, impacts species that show negligible or negative senescence is currently unknown.

2. Here, we examine whether and how variation in the quantity, quality, and feeding interval of resources impact population structure, population performance, and life history trait trade-offs in two long-lived planaria that do not senesce: *Schmidtea mediterranea* and *Dugesia tahitiensis*. In a full factorial design, different combinations of resource quantity (reduced intake, standard intake and high intake) and quality (high and low quality) were provided in two different feeding intervals (7-day and 14-day intervals) for 19 weeks.

3. We show that variability in resource availability, via decreases in quantity, quality, and frequency of resources, does not diminish population viability in either species but does result in suboptimal conditions of stress in *S. mediterranea*.

4. The high population viability we report can be attributed to two different mechanisms: increased reproduction or increased investment into maintenance at the expense of reproduction. Moreover, which mechanism was responsible for said high population viability was context-dependent and modulated by the specific life history strategy of the two planaria species.

5. We show that suboptimal conditions can cause stress responses that have significant

impacts on non-senescent species. The context-dependent response we observe suggests

that species that do not senesce but are subject to suboptimal conditions of stress may

ultimately exhibit declines in performance and ultimately die. A clearer understanding of the

impact of suboptimal conditions of resource availability on non-senescent species is needed

to determine the extent of stress experienced and ultimately whether a species can truly be

immortal.

Keywords: immortality, diet, life history trait, planaria, population structure, senescence, trade-off,

vitality.

INTRODUCTION

A hydra (a potentially immortal metazoan (Martínez, 1998)) walks out of a bar... and is run over by a bus. Nothing lasts forever, including potential immortality. How age impacts individual performance and fitness has received much attention (Baudisch, 2011; Baudisch & Stott, 2019; Caswell & Salguero-Gómez, 2013; Kirkwood, 1977; Medvedev, 1990; J. W. Vaupel, 2010). The reduction of individual performance with age due to physiological degradation, senescence (Finch, 1994; Jones et al., 2008), was initially thought to be universal, although recent evidence shows that not all species reduce individual performance with age (review by Flatt & Partridge, 2018; (Caswell & Salguero-Gómez, 2013; Jones et al., 2008, 2014; Roper et al., 2021; but see Nelson & Masel, 2017). Indeed, some species like the ocean quahog (*Arctica islandica*) (Abele et al., 2009), sugar cone pine (*Pinus lamertiana*) or guppy (*Poecilia reticulata*) (Roper et al., 2021) have mortality rates that remain constant with age, while other species display mortality rates that decline with age (Roper et al., 2021; Vaupel et al., 2004). The former phenomenon is known as negligible senescence (Finch, 1994, 2009; Vaupel et al., 2004) and the latter as negative senescence (Jones et al., 2014; Vaupel et al., 2004). Negative senescence has been reported in hydra (Martínez, 1998), prickly pear cactus (*Opuntia rastrera*) (Roper et al., 2021), or flatworms (planaria, Phylum Platyhelminthes) (Sahu et al., 2017). Crucially, regardless whether a species is senescent, negligibly senescent, or negatively senescent, its rates of senescence typically correlate with other life history traits (*e.g.*, generation time; Healy et al., 2019; Salguero-Gómez et al., 2016), with the magnitude of actuarial (= survival) senescence largely coupled by the speed of the life history (Jones 2008).

The underlying correlation between life history traits and rates of senescence then raises the question of how drivers of variation in life history traits ultimately impact senescence. Resource availability is a key driver of individual development and population performance (Ozgul et al., 2009; Smallegange, 2011; Stearns, 1992). Yet, studies on how resource availability impacts the life

cycle of an organism have largely focused on specific life transitions (*e.g.,* age at maturity (Auer,

2010; Smallegange, 2011; Stearns, 1992) or features of the life cycle (*e.g.,* reproduction (Rutstein et

al., 2005; Williams, 1996); survival (Ozgul et al., 2009)). Survival, development, and reproduction

are key demographic processes in the life cycle of any species. Indeed, these demographic

processes define life history traits (*e.g.,* age at maturity, longevity, *etc.*) that, in turn, produce life

history strategies that shape individual fitness and dynamics of a population (Roff 1992; Stearns

1992; Capdevila & Salguero-Gómez 2019). Life history theory predicts a trade-off in resource

allocation between maintenance (survival and development) and reproduction of individuals (Roff,

1992; Stearns, 1992). However, the impact of resource availability on these demographic processes,

and therefore on life history traits, is not straightforward. The impacts of variability in resources on

demographic process can be non-linear (*e.g.,* indirect winter climate on summer foraging

conditions; Mysterud et al., 2001) but can also interact with other factors (*e.g.,* anthropogenic land

use change) resulting in demographic impacts being exacerbated or buffered (Maron et al., 2015).

Given the nuanced impact that resources have on life history traits, a key question in ecology

and evolution is to what extent negligibly and negatively senescent species are impacted by

variable resource availability. Here, we contribute to filling this gap of knowledge by examining the

impact of variability of resources in two long-lived invertebrate species, the planarians *Schmidtea*

mediterranea (Benazzi, Baguna, Ballister, Puccinelli & Del Papa, 1975) and *Dugesia tahitiensis*

(Gourbault, 1977). Obligate asexual planaria do not show physiological senescence (Sahu et al.,

2017). In these animals, stem cells also function as germ line, and so many of the well-known

somatic mechanisms of senescence (Finkel & Holbrook, 2000; Lemaître et al., 2024; Payne &

Chinnery, 2015) do not apply to obligate asexual planarian stem cells (Sahu et al., 2017). However,

individuals of planarians can still suffer suboptimal conditions of stress, shrink, and even die (*e.g.,*

Ofoegbu et al., 2016). These features provide the perfect opportunity to assess the role of variable

resource availability in non-senescent species. Specifically, we focus on the impacts of variable

resource availability on planaria population size and population structure, with the rationale for

said focus detailed below.

Variation in vital rates (*e.g.*, age-specific survival) dictate population level processes such as

population growth rate and extinction risk (Hernández-Yáñez et al., 2022; Lande & Orzack, 1988).

Where survival does not decline with age, an indication of negligible or negative senescence,

population level processes may differ from populations where survival does decline with age

(actuarial senescence). For example, in organisms where survival declines with age, the growth rate

of their populations is typically reduced compared to populations of organisms with negligible

senescence (Edelfeldt et al., 2019). Crucially, variation in survival trajectories that are caused by age

effects may also be driven by environmental factors (Edelfeldt et al., 2019; Roach et al., 2009).

Therefore, changes at the population level (*e.g.,* size) due to variable resources can be an indicator

of changes in survival trajectories of individuals and potentially "senescence" status. Population-

level assessments of the impact of variable resources on survival trajectories are a well-established

approach in organisms such as yeast (*Saccharomyces cerevisiae*, Jiang et al., 2002). In yeast, lifespan

is assessed by the number of times cells within a multicellular colony produces daughter cells

through budding (Jiang et al., 2002). Such an approach is then also amenable to assess the impacts

of variable resources on mean life expectancy of a population of other species whereby clonal

reproduction is a key life history trait (*e.g.*, planaria, Malinowski et al., 2017). Furthermore, focusing

on the population level provides the opportunity to assess how variable resources shape population

structure. Indeed, population structure can play a key role in the dynamics of a population through

the changes of the ratio of large/old individuals to smaller/younger individuals (Benton & Grant,

1999; Caswell, 2010).

Here, we develop a full factorial design where we experimentally manipulate three different

dimensions of resource availability (quantity, quality, and feeding interval) to examine how

resource variability impacts the population structure, population performance, and life history trait

trade-offs of two potentially immortal species, the planarians *S. mediterranea* and *D. tahitiensis*.

We test three hypotheses regarding the impacts of these three dimensions of resource availability

on population size and structure, and underlying life history trade-offs, as detailed in Table 1.

METHODS

Study system

To examine how resource variability impacts the population structure, population viability, and

life history trait trade-offs, we chose planaria as a study system due to their key life history traits.

Notably, the majority of planarians are capable of restorative and physiological regeneration due to

injury or damaged/sick cells (Elliott & Sánchez Alvarado, 2013). Their ability to replace damaged or

sick cells contributes to the typically long-lived life history of planarians (Elliott & Sánchez Alvarado,

2013). Indeed, under favourable, constant conditions, planaria are theoretically immortal (Elliott &

Sánchez Alvarado, 2013). Planaria are carnivorous and feed on live arthropods, annelids, and

molluscs. However, planaria also scavenge on individuals that have recently died (Abel et al., 2022;

Vila-Farré & C. Rink, 2018). Prey preference has also been shown in several planaria species (Vila-

Farré & C. Rink, 2018). Nonetheless, this preference can vary among localities and when the

preferred prey is scarce (Vila-Farré & C. Rink, 2018). Importantly, individual planarians are also able

to shrink when faced with unfavourable conditions (González-Estévez, 2009).

Here, we study responses to variability in resource availability by *Schmidtea mediterranea* and

Dugesia tahitiensis, two closely related, freshwater planarian species from the order Tricladida.

Both species are free-living and naturally occur in slow running and still water, but are easily reared

in the lab (Sousa & Adell, 2018). The two species also differ in size, with *D. tahitiensis* being larger

(≈10mm in length) than *S. mediterranea* (≈5mm). *S. mediterranea* can reproduce sexually and

asexually, whereas *D. tahitiensis* is an obligate asexual. We used asexual lines of *S. mediterranea* for

our experiments to directly compare the two species. Asexual reproduction in planaria occurs via

binary fission, which results in two genetically identical clones of smaller size from the fissioning

individual (Malinowski et al., 2017). Details of stock culture maintenance are described in the

Supporting Information.

Experiment

To test our hypotheses regarding how variability in resource availability impacts populations of

S. mediterranea and *D. tahitiensis* (Table 1), we randomly assigned populations to a full-factorial

design for each of the two species. This design included three resource factors: quality, quantity,

and feeding interval (Fig. 1, Box 1). Quality of resource had two levels, differing in energy content

and relative protein-to-carbohydrate ratio: high quality (high carbohydrate level, *HQ* hereafter) *vs.*

low quality (low carbohydrate level, *LQ* hereafter). The high quality diet was organic calf's liver and

the low quality diet bloodworm. The protein-to-carbohydrate ratio indicates resource quality, with

low carbohydrate-to-protein content corresponding to poorer quality (Lee et al. 2008). The two

diets differed in their energy content (109 kcal/100g in the liver *vs.* 14 kcal/100g in the bloodworm)

and in their relative protein-to-carbohydrate ratios (1:0.35 in liver *vs.* 1:0.17 in bloodworm).

Quantity of resources had three levels, which differed in the amount of calories available in each

feeding: standard calorie intake (*SI*: 0.001g/individual, representative of intake to maintain

laboratory populations (Jochen, 2018)), restricted calorie intake (*RI*: 0.0005g/ind.), hyper-calorie

intake (*HI*: *ad libitum* which was at least 10 fold, approx. 0.01g/individual, that of the SI). Different

combinations of resource quantity and quality were provided at two different intervals: once every

seven days (*7D*), or every 14 days (*14D*). The total amount of resources provided over a 14-day period was equivalent for both feeding interval treatments (*i.e.*, one 14-day interval equalled two 7-day intervals) (see Fig. 1). As such, across the most extreme diet treatments, there is over a 300-fold difference in calories provided (per individual per unit time). Within our experimental setup, the equivalent to a control would be either the high quality (*HQ*), standard intake (*SI*), seven days interval (*7D*) treatment (*i.e.*, *HQ-SI-7D*) or the low quality (*LQ*), standard intake (*SI*), 14 days interval (*14D*) treatment (*i.e.*, *LQ-SI-14D*). Here, we determine resources provided at an interval of every seven days (*7D*) of a high quality non-restricted (standard intake for the organism) diet as our control (i.e., *HQ-SI-7D*).

To evaluate interactive effects of resource availability on population structure, population viability, and life history trait trade-offs, we replicated all treatments in the full factorial (n = 12 levels: 2 qualities: *HQ vs. LQ*; 3 quantities: *RI*, *SI* & *HI*; 2 intervals: *7D vs. 14D*) five times for each species. A replicate consisted of a petri dish (25mm deep × 150mm Ø) containing 10 clonal individuals each, henceforth referred to as a population (total n = 600 ind./spp.). For each population, individuals were randomly isolated from one of the three stock cultures and placed in the population 7-8 days before the start of the experiment. Importantly, as individuals of both species are clonal, and stock cultures were initiated with individuals from the same cohort, individuals within each population were of the same chronological age. During this time, individuals were not fed to ensure individuals across all treatments were in a similar starved state at the start of the experiment. Treatment populations were randomised into three blocks containing 20 populations per species per block. For each species, all experimental blocks lasted 19 weeks (133 days). Blocks were staggered in three different start dates due to the logistical constraint of sampling all 60 populations on the same day, as such staggered blocks allowed a more logistically feasible sampling of 20 populations per day. Block 1 of the experiment started on the 4[th] February 2020, and block 2 on the 30[th] April 2020. In block 3, the start date for *S. mediterranea* and *D.*

tahitiensis differed due to low numbers available for *S. mediterranea* from the stock populations. As

such, block 3 for *D. tahitiensis* started on the 1st May 2020 and for *S. mediterranea* on the 19th June

2020. Part way through the experiment, due to the COVID-19 pandemic, the location of the

experiment was changed within 2 hours, while retaining the same environmental conditions, see

Fig S1 in the Supporting Information.

Within each population, every week, population extinction was determined, and asexual

reproduction (clonal reproduction) estimated by counting all living and dead individuals. The weekly

determination of population extinction and estimation of reproduction was done to determine

changes in population size. The expectation is that, in H1 and H3 (Table 1), risk of population

extinction will decrease due to increased reproduction, while in H2 (Table 1) extinction risk will

decrease due to investment in maintenance. Additionally, size (area in mm^2) of each individual

flatworm within each population was measured weekly to determine growth/shrinkage. Size

measurements were calculated from photographs taken of each population (details of image

digitization can be found in the Supporting Information). Our expectations here are that for H1,

individual size will be normally distributed as the average resource intake among individuals will not

be impacted due to the high quantity of resources available. For H2, environments will be

dominated by larger individuals due to competitive effects; and for H3, reduced feeding interval of

resources will result in greater rate of size oscillation as more frequent resources does not restrict

fissioning events to larger individuals (Nentwig & Schauble, 1974).

Size distribution analysis

To test impact of variation in resource availability on individual body size, we used the size

spectrum of populations as a method of quantifying the distribution of body size (Edwards et al.,

2017). Where populations are increasing in size, with increases in small individuals, this implies

increased reproductive output. To do so, we calculated the size spectra slope values (b) for each

population, of both species, across treatments following Carvalho et al. (2021). Briefly, b describes

the relative abundance of a population in relation to body sizes, thus providing a good proxy to

quantify the distribution of body size within a population. A steeper (more negative) b indicates

fewer large-bodied and/or more small-bodied individuals, which is an indication of a right-skewed

size population structure. Values of b can vary substantially with estimates ranging from -6.75 to -

0.40 been reported (Arranz et al., 2021; Carvalho et al., 2021). To estimate b, for each species

separately, we used the abundance density function (Carvalho et al., 2021):

250
$$N(x) = nx^b \left(\frac{b+1}{x_{max}^{b+1} - x_{min}^{b+1}} \right) \tag{1}$$

where n is the number of individuals, x the area of each individual (mm^2), x_{max} and x_{min} are the

maximum and minimum area of individuals that were digitized. To calculate the maximum and

minimum values, all populations for each treatment were pooled and maximum and minimum

values calculated separately for each treatment for each week.

For each week of the experiment, b values were calculated using the area measurements of

each individual within a given population. The b value for each of the five populations of each

treatment was calculated. As such, here b relates to how the populations responds to the varying

quality, quantity, and feeding interval of resources provided. Next, we applied a Bayesian

framework to determine changes in the mean b values using the "bmrs" package (Bürkner, 2017) in

R version 4.0.3 (R Core Team, 2020). Given that the estimated b values are based on size values,

there is the potential that the b values may show a right-skewed distribution due to the ability of

planaria to shrink. Thus, we initially compared models with a Gamma distribution to that of a

Gaussian distribution to identify the model with the best fit. We identified that in the case of both

S. mediterranea and *D. tahitiensis* a model with a gaussian distribution had the best fit (Supporting

Information, Table S2). The final model used to model the response variable *b*, had a Gaussian

distribution and adaptive priors with treatment (categorical with 12 levels), week (19 weeks) and

block (3 blocks) as fixed variables and population (60 populations) as a random effect. Presented

are the median estimate and 50% and 95% compatibility intervals of the posterior probability.

Compatibility intervals can be considered equivalent to confidence intervals in a frequentist

approach (Gelman & Greenland, 2019). See Supporting Information for model details and Table S3

& S4 for final priors of each model.

Population size analysis

To determine the impact of variation in resource availability on population size, we calculated

changes in weekly population size for each species. We applied a Bayesian framework to determine

changes in population size using the same "bmrs" package as before. Population size was modelled

with a Poisson and Gaussian distribution to determine best fit. In the case of *S. mediterranea* the

model with the best fit was identified as one with a Gaussian distribution (Supporting Information,

Table S5) and adaptive priors, with treatment (categorical with 12 levels), week (19 weeks) and

block (3 blocks) as fixed variables, population as a random effect (60 populations) and population

size the response variable. In the case of *D. tahitiensis*, the model with the best fit had a Poisson

distribution with a log link function (Supporting Information, Table S5) using adaptive priors, with

treatment (categorical with 12 levels), week (19 weeks) and block (3 blocks) as fixed variables,

population as a random effect (60 populations) and population size the response variable. As

before, median estimated values are presented with 50% and 95% compatibility intervals. See

Supporting Information for model details and Table S6 & S7 for final priors of each model.

Crucially by analysing changes in population size and population structure (as identified by the

size distribution analysis), we can provide insights in potential reproductive output given our

knowledge of the biology and life cycle of both species (Carter et al., 2015; González-Estévez, 2009;

Malinowski et al., 2017; Nentwig & Schauble, 1974). For example, a population that is increasing in

its size with increases in the number of small individuals implies increased reproductive output. In a

population where individuals are shrinking, fissioning (reproduction) is not occurring and

populations are not increasing. As such there will not be overlap with the size distributions of small

individuals due to reproductive output and small individuals as a result of shrinking.

RESULTS

Body size distributions

Over time, size spectra slope values (b) for *S. mediterranea* became more negative for most

treatments (Fig. 2), implying a more right-skewed population. The consistent decline in b values

indicates a change in the size structure of the populations with an increase in smaller individuals

and fewer larger individuals. Overall, 14-day interval treatments had more negative b values than 7-

day interval treatments (Fig. S3). In the high-quality treatments, when compared to *HQ-SI-7D* (our

control), only the 14-day feeding interval high intake (*HQ-HI-14D*) and 14-day feeding interval

reduced intake (*HQ-RI-14D*) treatment had smaller b values and as such more right-skewed size

structure, with the *HQ-RI-14D* treatment fulfilling the size prediction of H2 (Table 1). This outcome

suggests that in high quality treatments populations changed in size structure with an increase in

the number of small individuals and a decrease in the number of larger individuals over time (Fig.

2). When changes in structure were found, this change in structure was more prominent in

treatments where feeding interval was 14 days. However, in the case of the low-quality treatments,

only the 7-day feeding interval reduced intake (*LQ-RI-7D*) treatment did not show a change in size

structure that differed from that of *HQ-SI-7D* (Fig. S3). All other low quality treatments showed a

greater change in size structure (*i.e.*, more negative *b* values) with more right-skewed populations

over time (Fig. 2). In the case of *LQ-HI-7D*, the size prediction of H1 was not fulfilled as size was

predicted to be normally distributed.

For *D. tahitiensis*, size spectra slope values (*b*) varied less in the 7-day feeding interval

treatments than the 14-day feeding interval treatments (Fig. 3). When compared to *S.*

mediterranea, *D. tahitiensis* populations were less right-skewed. When assessing the outcomes in

terms of the size predictions of H1 and H2 for *D. tahitiensis*, treatment *LQ-HI-7D* fulfilled the

predictions of H1 as body size was normally distributed (Table 1). However, for treatment *HQ-RI-*

14D the size predictions of H2, a right-skewed population, was not fulfilled (Table 1). In both species

experimental block 1 differed from block 2 and block 3 in several treatments (Fig. S3 & S4).

However, this pattern was not consistent between treatments and species (see Fig. 2 and 3, light

blue points). Still, even with the block differences, the trends of all blocks were the same

(increasing, decreasing, or stable). Overall, there was no consistent trend in how *b* values across

treatments differed to that of *HQ-SI-7D*. As such, treatment effect does not predict variation in size

structure over time.

Population trends

Overall, *D. tahitiensis* populations had higher numbers than populations of *S. mediterranea* at

the end of the experiment (Fig. 4 & 5). Population size of *S. mediterranea* varied between

treatments that differed in resource quantity (*HI*: high intake, *SI*: standard intake, *RI*: reduced

intake) and interval of feeding (*7D*: every seven days, *14D*: every 14 days) but was consistent across

resource quality (*HQ*: high quality, *LQ*: low quality) (Fig. 4A & B). The exception being the high

intake resource quantity treatments with a 7-day feeding interval (*HI-7D*). In this case, the *HQ-HI-7D*

treatment, where the resource was of a high quality, the population declined compared to the *LQ-*

HI-7D treatment, where the population increased over time (Fig. 4A & B, Fig. S5). All populations

where resource was provided every 14 days showed a decrease in population size over time

compared to populations when feeding interval was every seven days, where counts either stayed

constant (*HQ-RI-7D*) or increased over time (*HQ-SI-7D, LQ-HI-7D , LQ-SI-7D*) (Fig. 4A & B Fig. S5).

The exception here was the *HQ-HI-7D* treatment, where population size declined over time. The

decline in *HQ-HI-7D* does not fulfil the prediction of H3 where the expectation was an increase in

reproduction and a larger population size. However, the decrease in population size of treatment

HQ-RI-14D supports the population size prediction of H2. As such, the outcome of *HQ-RI-14D* fulfils

the full prediction of H2 that population size will decrease and body size will be right-skewed.

In *D. tahitiensis*, populations either increased or remained stable across all treatments (Fig. 5A

& B). When comparing *HQ* and *LQ* treatments, *HQ* treatments with feeding intervals of seven days,

regardless of resource quantity, had a higher population size than *LQ* treatments with 7-day feeding

intervals, except for *HQ-HI-7D* and *LQ-HI-7D*, which had similar counts. When resource was

provided every 14 days, there was no clear trend; the *HQ-HI-14D* treatment had lower counts than

the *LQ-HI-14D* treatment, *HQ-SI-14D* had higher counts than *LQ-SI-14D* and *HQ-RI-14D* had similar

counts to *LQ-RI-14D* (Fig. 5A & B, Fig. S6). In the case of *HQ-RI-14D* population size increased which

did not fulfil the population predictions of H2. For treatments *LQ-HI*-7D and *HQ-HI-7D* population

size increased, thus fulfilling the population predictions of H1 and H3 respectively. For both species,

experimental block 1 (light blue lines, Fig. 4 & 5) had population growth rates lower than block 2

and block 3. However, for all treatments, the population response (increasing, decreasing, or stable

counts) of block 1 did not differ from that of block 2 or block 3 (Fig. 4 & Fig. 5, Fig. S5 & S6).

DISCUSSION

Compared to senescent species (Lemanski et al., 2020; Pardo et al., 2013), a comprehensive

understanding of the impact of variable resource availability on species that show negligible or

negative senescence is lacking (but see Schaible et al., 2011). Here, we contribute to this gap of

knowledge by incorporating variation in resource availability in two non-senescent species: the

flatworms *S. mediterranea* and *D. tahitiensis* (Sahu et al., 2017). Our findings show how, when

multiple factors of resource availability interact (*i.e.*, quality, quantity and frequency), these non-

senescent species show population-level responses expected from suboptimal conditions of stress.

Specifically, in *S. mediterranea*, we observe suboptimal conditions of stress through population

decline and individual shrinkage when resources are made available are every 14 days, regardless of

the resource quality or quantity. However, conditions of stress in *Dugesia tahitiensis* are less

frequent and occur only when resources are reduced, regardless of resource intervals, with only

individual-level shrinkage being noted as evidence of stress, since we observe no population

decline.

The way these factors of resource availability impacts individuals within populations is context

dependent in our experiment. The high population viability (only two of the 20 populations went

extinct in our experiment, both were *S. mediterranea*) and variable response in reproduction (as

proxied by population size and structure) across treatments is also likely driven by the specific life

history strategy of our species. Generally, planarians can restore their own damaged cells (Elliott &

Sánchez Alvarado, 2013). However, restoring damaged cells is costly and requires resources (Akashi

& Gojobori, 2002). Naturally, if most resources are used for maintenance, fewer resources will be

available for reproduction. The high population viability we report here in response to the variable

resource availability is indicative of the fundamental trade-off between survival and reproduction

under variable environmental conditions (Stearns, 1992). Here, theory predicts that, in long lived

species such as our planarians (Elliott & Sánchez Alvarado, 2013), resources will be primarily

invested in maintenance of individuals and not reproduction (Erikstad et al., 1998; Sæther et al.,

2013). As highlighted earlier, reproduction in planaria occurs via binary fission, and so individuals in

our experimental setup are clonal and thus of the same age. As such, any changes in the structure

and size of a population can be thought of as a reflection of the survival-reproduction trade-off. A

less right-skewed population, where there are fewer smaller individuals, is indicative of reduced

reproduction as new recruits are smaller. In these populations of reduced reproduction, individuals

invest in maintenance that results in reduced/no growth with little change in body size. In general,

for *S. mediterranea*, such a survival-reproduction trade-off can be observed across the 7-day

feeding treatment, but not when the feeding interval was every 14 days, where population size

decreased, indicating reduced reproduction and reduced survival of individuals. For *D. tahitiensis*,

only populations of low quantity and with a feeding interval every 14 days show a survival-

reproduction trade-off of high survival and reduced reproduction. Such context-dependant trade-

offs have also been shown in *Hydra magnipapillata*, another non-senescent species (Schaible et al.,

2011). There, the authors found a clear maintenance-reproduction trade-off when resources and

temperature were constant, whereby investment into reproduction resulted in reduced survival.

However, under fluctuating environmental conditions, the authors reported no clear maintenance-

reproduction trade-off. This context-dependent survival-reproduction trade-off then begs the

question, where there is an apparent lack of trade-off between survival and reproduction (see

Metcalf 2016), do we see a trade-off with body size?

The size structure of our populations varied across treatments, but not in a way that was

consistent with our hypotheses (Table 1). For *S. mediterranea*, populations become more right-

skewed across treatments, with two exceptions in the 7-day feeding interval treatment. The right-

skewed populations describe an increase in the number of smaller individuals and decline of larger

individuals in the populations (Carvalho et al., 2021). Such changes in population structure can

occur when there is increased reproduction, as more recruits of a smaller size enter the population.

S. mediterranea reproduce asexually through fission, where larger individuals fission into two, with

the posterior product (new recruits) generally smaller than the anterior fission product (now a

smaller individual too) (Peter et al., 2001). Thus, fissioning results in fewer larger individuals and an

increase in the number of smaller individuals. However, the right-skewed size structure due to

reproductive events will only be reflected in populations that are increasing in number as in these

populations no mortality of individuals was observed. Where *S. mediterranea* populations show a

decline or remain stable in size, the change in structure is not driven by decreased reproduction.

Instead, the life history of planaria seems to be driving the response. A key trait in the life history of

planaria is the ability to shrink when resources become scarce (Mangel et al., 2016; Rink, 2013).

The ability to shrink allows individuals of some species to survive resource-scarce periods of

time until conditions improve (Ebert, 1967; Marinovic & Mangel, 1999; Wikelski & Thom, 2000). In

planaria, when individual shrinkage occurs, cell numbers reduce through autophagy during which

energy resources are supplied to stem/regenerative (neoblasts) cells (González-Estévez, 2009;

Romero & Baguñà, 1991). This process ensures that maintenance levels can be met and as

individuals attempt to maintain homeostasis, fissioning (asexual reproduction) does not occur.

Furthermore, in *S. mediterranea* and in sister species of *D. tahitiensis* (*i.e.*, *D. japonica* and *D.*

dorotocephala) the probability of fissioning declines as individuals get smaller, with a minimum size

where fissioning can occur (Carter et al., 2015). In the case of sexually reproducing planaria,

reproductive organs are often consumed during shrinkage (Saló, 2006), thus preventing further

reproduction. Shrinkage is a strategy found in several taxa and been suggested to mitigate the

impact of fluctuating or variable environments on individual survival (Ebert, 1967; Marinovic &

Mangel, 1999; Salguero-Gómez & Casper, 2010, 2011). Indeed, shrinkage maintains relatively high

levels of survival during periods of resource scarcity compared to strategies where shrinkage does

not or cannot physically occur. As such, the high levels of survival ensure population viability. For

instance, in the Galápagos marine iguanas (*Amblyrhynchus cristatus*), individuals shrink up to 20%

of their body length when nutritional resources are low following El Niño (Wikelski & Thom, 2000);

those individuals that shrink more have a higher rate of survival, suggesting the adaptive value of

shrinkage. When nutritional resources improve during La Niña events, body length then increases

again, as is the case with planarians when they receive increased nutritional resources after periods

of low nutrient availability (González-Estévez, 2009; Thommen et al., 2019).

The impact of variable resources on body size due to shrinkage observed in *S. mediterranea*

populations is unlikely occurring in populations of *D. tahitiensis*. Fewer *D. tahitiensis* treatments

showcased populations becoming more right-skewed (*i.e.,* more smaller individuals) over the

course of the experiment. There, where populations become more right-skewed, populations do

not show a decline in population size and, unlike the case with *S. mediterranea*, changes in

population structure are likely due to fissioning. Populations of *D. tahitiensis* that have a less right-

skewed size structure increased in size during the experiment. Within these population, individuals

are likely experiencing increased competition for resources. Increased resource competition among

individuals results in contest competition, where an individual interferes with other individuals to

obtain resources (Nicholson, 1954). In competitive environments, smaller individuals are typically

outcompeted by larger individuals within the population as larger individuals tend to be more

dominant (Petersson & Järvi, 2000; Stewart & Tabak, 2011). In our study, contest competition

results in differences in average resource intake among individuals, with larger planarian individuals

able to consume more resources per feeding event. As such, competition between two individuals

of planarian naturally results in the larger of the two growing more. The competitive exclusion of

smaller individuals thereby increases the number of larger individuals in the population, resulting in less right-skewed populations. Here, competition for resources resulted in mortality of smaller individuals and faster growth in the surviving, larger individuals. Evidence of smaller individuals being outcompeted by larger individuals has been observed in several species, including other planaria (Reynoldson, 1964; Reynoldson & Sefton, 1972). For example, in a 20 month study of a lake population of *Planaria torva*, Reynoldson and Sefton (1972) found changes in the size distribution of individuals in the population that were attributed to intraspecific competition for resources. Similarly, in the brown trout (*Salmo trutta*), Petersson & Järvi (2000) observed that sea-ranch trout, which is 13% larger than its wild-type counterpart, had higher dominance than the latter.

The high population viability reported here highlights the ability of planaria to mediate impacts of variation in resource availability. However, as mentioned before, the mechanisms that allow shrinkage and regeneration of individuals do not preclude individuals from suffering suboptimal conditions of stress and ultimately dying. Said resource-availability driven stress is observed in the few *S. mediterranea* populations that went extinct and those populations declining in numbers. Why such extinction events and declining numbers only occurred in populations of *S. mediterranea* and not *D. tahitiensis* can be attributed to differences in their ability to regenerate. Regeneration is not uncommon within the Metazoa, with almost every phylum having a species with this ability (Alvarado, 2000). Furthermore, regeneration can be classified into two categories highlighting the variability of the phenomenon: 1) morphallaxis, the regeneration in the absence of active cell proliferation (*e.g.*, *Hydra*) and 2) epimorphosis, the regeneration with cell proliferation (*e.g.*, limb and tail regeneration in vertebrates) (Alvarado, 2000; Morgan, 1901). In the case of planaria, both forms of regeneration are utilised but are context dependent; morphallaxis to restore scale and proportion (*e.g.*, when shrinking occurs) and epimorphosis for the formation of a new organ (*e.g.*, due to a fission event) (Pellettieri, 2019).

The lack of local extinction and declining numbers in the examined populations of *D.*

tahitiensis, compared to *S. mediterranea*, is likely due to the former species being able to

implement epimorphosis and repair damaged cells more efficiently than the latter (Peter et al.,

2001). Indeed, *Dugesia tahitiensis* has one of the highest regenerative and fissioninig capacities and

the most neoblasts (adult stem cells found in Platyhelminthes and Acoela) within the Tricladida

taxon (Baguñá & Romero, 1981; Peter et al., 2001). High neoblast density in the cellular tissue

between the body wall and organs is a prerequisite for efficient regeneration after fission and,

when compared to *S. mediterranea*, *D. tahitiensis* have higher cell counts (Peter et al., 2001).

Moreover, neoblast self-renewal activity can be impacted by resource availability with reduced

resource levels resulting in lowered self-renewal activity (Mangel et al., 2016). Reduced neoblast

self-renewal activity will then enhance the negative impacts experienced by *S. mediterranea*. The

variation in self-renewal activity brought about by variation in resource availability will ultimately

impact survival trajectories of individuals. As such, the fittest individuals are more likely to survive

to extreme ages that increases the population's average survival, a pattern that has also been

described in other animals (Vaupel & Yashin, 1985). This age-by-environment interaction has also

been reported in some long-lived plants, *e.g.* in *Plantago lanceolata* (Roach et al., 2009) and

Fumana procumbens (Edelfeldt et al., 2019), where variation in survival trajectories are modulated

by resource availability.

No species is truly immortal. The changing environment can play a significant part in the levels

of stress and individual faces, regardless of its senescence rates. Here, we contribute to the under-

represented empirical demonstration of the impact of variation in resource availability in two

species that do not senesce. Our findings support previous work on another non-senescent species

(*Hydra magnipapillata*; Schaible et al., 2011), showing that the impacts of fluctuating resources on

the trade-off between reproduction and maintenance is context dependent. Importantly, we

highlight that two non-senescent species respond differently to variation in resource availability;

one species can suffer population extinction and the other population growth, highlighting

differential suboptimal conditions of stress under changing environmental conditions. A clearer

understanding of how fluctuating environments, particularly in resources availability, impacts

species that show the unique trait of negligible or negative senescence will be key to fully

understand the role of this life history trait and ultimately whether anything is indeed immortal.

References

Abel, C., Powers, K., Gurung, G., & Pellettieri, J. (2022). Defined diets for freshwater planarians. *Developmental Dynamics*, *251*(2), 390–402. https://doi.org/10.1002/dvdy.400

Abele, D., Brey, T., & Philipp, E. (2009). Bivalve models of aging and the determination of molluscan lifespans. *Experimental Gerontology*, *44*, 307–315. https://doi.org/10.1016/j.exger.2009.02.012

Akashi, H., & Gojobori, T. (2002). Metabolic efficiency and amino acid composition in the proteomes of Escherichia coli and Bacillus subtilis. *Proceedings of the National Academy of Sciences*, *99*(6), 3695–3700. https://doi.org/10.1073/pnas.062526999

Alvarado, A. S. (2000). Regeneration in the metazoans: Why does it happen? *BioEssays*, *22*(6), 578–590. https://doi.org/10.1002/(SICI)1521-1878(200006)22:6<578::AID-BIES11>3.0.CO;2-#

Arranz, I., Brucet, S., Bartrons, M., García-Comas, C., & Benejam, L. (2021). Fish size spectra are affected by nutrient concentration and relative abundance of non-native species across streams of the NE Iberian Peninsula. *Science of The Total Environment*, *795*, 148792. https://doi.org/10.1016/j.scitotenv.2021.148792

Auer, S. K. (2010). Phenotypic Plasticity in Adult Life-History Strategies Compensates for a Poor Start in Life in Trinidadian Guppies (Poecilia reticulata). *The American Naturalist*, *176*(6), 818–829. https://doi.org/10.1086/657061

Baguñá, J., & Romero, R. (1981). Quantitative analysis of cell types during growth, degrowth and

regeneration in the planarians Dugesia mediterranea and Dugesia tigrina. *Hydrobiologia*, *84*(1), 181–

194. https://doi.org/10.1007/BF00026179

Baudisch, A. (2011). The pace and shape of ageing. *Methods in Ecology and Evolution*, *2*(4), 375–382.

https://doi.org/10.1111/j.2041-210X.2010.00087.x

Baudisch, A., & Stott, I. (2019). A pace and shape perspective on fertility. *Methods in Ecology and Evolution*,

10(11), 1941–1951. https://doi.org/10.1111/2041-210X.13289

Bayliss, P., & Choquenot, D. (2002). The numerical response: Rate of increase and food limitation in

herbivores and predators. *Philosophical Transactions of the Royal Society of London. Series B:*

Biological Sciences, *357*(1425), 1233–1248. https://doi.org/10.1098/rstb.2002.1124

Benton, T. G., & Grant, A. (1999). Elasticity analysis as an important tool in evolutionary and population

ecology. *Trends in Ecology & Evolution*, *14*, 467–471. https://doi.org/10.1016/s0169-5347(99)01724-

3

Bürkner, P.-C. (2017). brms: An R Package for Bayesian Multilevel Models Using Stan. *Journal of Statistical*

Software, *80*(1), 1–28. https://doi.org/10.18637/jss.v080.i01

Capdevila, P., & Salguero-Gomez, R. (2019). Prototypical r-/K-Selected (Fast/Slow) Species. In T. K.

Shackelford & V. A. Weekes-Shackelford (Eds.), *Encyclopedia of Evolutionary Psychological Science*

(pp. 1–4). Springer International Publishing. https://doi.org/10.1007/978-3-319-16999-6_1934-1

Carter, J. A., Lind, C. H., Truong, M. P., & Collins, E.-M. S. (2015). To Each His Own. *Journal of Statistical*

Physics, *161*(1), 250–272. https://doi.org/10.1007/s10955-015-1310-1

Carvalho, P. G., Setiawan, F., Fahlevy, K., Subhan, B., Madduppa, H., Zhu, G., & Humphries, A. T. (2021).

Fishing and habitat condition differentially affect size spectra slopes of coral reef fishes. *Ecological*

Applications, *n/a*(n/a), e02345. https://doi.org/10.1002/eap.2345

Caswell, H. (2010). Reproductive value, the stable stage distribution, and the sensitivity of the population

growth rate to changes in vital rates. *Demographic Research*, *S8*(19), 531–548.

Caswell, H., & Salguero-Gómez, R. (2013). Age, stage and senescence in plants. *Journal of Ecology*, *101*(3),

585–595. https://doi.org/10.1111/1365-2745.12088

Ebert, T. A. (1967). Negative Growth and Longevity in the Purple Sea Urchin Strongylocentrotus purpuratus

(Stimpson). *Science, 157*(3788), 557–558. https://doi.org/10.1126/science.157.3788.557

Edelfeldt, S., Bengtsson, K., & Dahlgren, J. P. (2019). Demographic senescence and effects on population

dynamics of a perennial plant. *Ecology, 100*(8), e02742. https://doi.org/10.1002/ecy.2742

Edwards, A. M., Robinson, J. P. W., Plank, M. J., Baum, J. K., & Blanchard, J. L. (2017). Testing and

recommending methods for fitting size spectra to data. *Methods in Ecology and Evolution, 8*(1), 57–

67. https://doi.org/10.1111/2041-210X.12641

Elliott, S. A., & Sánchez Alvarado, A. (2013). The history and enduring contributions of planarians to the study

of animal regeneration. *Wiley Interdisciplinary Reviews: Developmental Biology, 2*(3), 301–326.

https://doi.org/10.1002/wdev.82

Erikstad, K. E., Fauchald, P., Tveraa, T., & Steen, H. (1998). ON THE COST OF REPRODUCTION IN LONG-LIVED

BIRDS: THE INFLUENCE OF ENVIRONMENTAL VARIABILITY. *Ecology, 79*(5), 1781–1788.

https://doi.org/10.1890/0012-9658(1998)079[1781:OTCORI]2.0.CO;2

Finch, C. E. (1994). *Longevity, Senescence, and the Genome*. University Chicago Press.

Finch, C. E. (2009). Update on Slow Aging and Negligible Senescence – A Mini-Review. *Gerontology, 55*(3),

307–313. https://doi.org/10.1159/000215589

Finkel, T., & Holbrook, N. J. (2000). Oxidants, oxidative stress and the biology of ageing. *Nature, 408*(6809),

239–247. https://doi.org/10.1038/35041687

Flatt, T., & Partridge, L. (2018). Horizons in the evolution of aging. *BMC Biology, 16*(1), 93.

https://doi.org/10.1186/s12915-018-0562-z

Gelman, A., & Greenland, S. (2019). Are confidence intervals better termed "uncertainty intervals"? *BMJ,*

366, l5381. https://doi.org/10.1136/bmj.l5381

González-Estévez, C. (2009). Autophagy meets planarians. *Autophagy, 5*(3), 290–297.

https://doi.org/10.4161/auto.5.3.7665

Healy, K., Ezard, T. H. G., Jones, O. R., Salguero-Gómez, R., & Buckley, Y. M. (2019). Animal life history is

shaped by the pace of life and the distribution of age-specific mortality and reproduction. *Nature*

Ecology & Evolution, 3(8), 1217–1224. https://doi.org/10.1038/s41559-019-0938-7

Hernández-Yáñez, H., Kim, S. Y., & Che-Castaldo, J. P. (2022). Demographic and life history traits explain

patterns in species vulnerability to extinction. *PLOS ONE*, *17*(2), e0263504.

https://doi.org/10.1371/journal.pone.0263504

Jiang, J. C., Wawryn, J., Shantha Kumara, H. M. C., & Jazwinski, S. M. (2002). Distinct roles of processes

modulated by histone deacetylases Rpd3p, Hda1p, and Sir2p in life extension by caloric restriction in

yeast. *Experimental Gerontology*, *37*(8–9), 1023–1030. https://doi.org/10.1016/S0531-

5565(02)00064-5

Jochen, R. (2018). *Planarian Regeneration: Methods and Protocols* (1st ed.). Humana Press.

Jokela, J., & Mutikainen, P. (1995). Phenotypic plasticity and priority rules for energy allocation in a

freshwater clam: A field experiment. *Oecologia*, *104*(1), 122–132.

https://doi.org/10.1007/BF00365570

Jones, O. R., Gaillard, J. M., Tuljapurkar, S., Alho, J. S., Armitage, K. B., Becker, P. H., Bize, P., Brommer, J.,

Charmantier, A., Charpentier, M., Clutton-Brock, T., Dobson, F. S., Festa-Bianchet, M., Gustafsson, L.,

Jensen, H., Jones, C. G., Lillandt, B. G., McCleery, R., Merila, J., … Coulson, T. (2008). Senescence rates

are determined by ranking on the fast-slow life-history continuum. *Ecology Letters*, *11*, 664–673.

https://doi.org/10.1111/j.1461-0248.2008.01187.x

Jones, O. R., Scheuerlein, A., Salguero-Gómez, R., Camarda, C. G., Schaible, R., Casper, B. B., Dahlgren, J. P.,

Ehrlén, J., García, M. B., Menges, E. S., Quintana-Ascencio, P. F., Caswell, H., Baudisch, A., & Vaupel,

J. W. (2014). Diversity of ageing across the tree of life. *Nature*, *505*, 169.

Kirkwood, T. B. L. (1977). Evolution of ageing. *Nature*, *270*(5635), 301–304.

https://doi.org/10.1038/270301a0

Lande, R., Engen, S., & Saether, B.-E. (2003). *Stochastic Population Dynamics in Ecology and Conservation*.

Oxford University Press.

Lande, R., & Orzack, S. H. (1988). Extinction dynamics of age-structured populations in a fluctuating

environment. *Proceedings of the National Academy of Sciences*, *85*(19), 7418–7421.

https://doi.org/10.1073/pnas.85.19.7418

606 Lee, K. P. (2015). Dietary protein:carbohydrate balance is a critical modulator of lifespan and reproduction in

607   Drosophila melanogaster: A test using a chemically defined diet. *Journal of Insect Physiology*, *75*, 12–

608   19. https://doi.org/10.1016/j.jinsphys.2015.02.007

609 Lemaître, J.-F., Moorad, J., Gaillard, J.-M., Maklakov, A. A., & Nussey, D. H. (2024). A unified framework for

610   evolutionary genetic and physiological theories of aging. *PLOS Biology*, *22*(2), e3002513.

611   https://doi.org/10.1371/journal.pbio.3002513

612 Lemanski, N. J., Bansal, S., & Fefferman, N. H. (2020). The sensitivity of a honeybee colony to worker

613   mortality depends on season and resource availability. *BMC Evolutionary Biology*, *20*(1), 139.

614   https://doi.org/10.1186/s12862-020-01706-4

615 Malinowski, P. T., Cochet-Escartin, O., Kaj, K. J., Ronan, E., Groisman, A., Diamond, P. H., & Collins, E.-M. S.

616   (2017). Mechanics dictate where and how freshwater planarians fission. *Proceedings of the National*

617   *Academy of Sciences*, *114*(41), 10888. https://doi.org/10.1073/pnas.1700762114

618 Mangel, M., Bonsall, M. B., & Aboobaker, A. (2016). Feedback control in planarian stem cell systems. *BMC*

619   *Systems Biology*, *10*, 17–17. PubMed. https://doi.org/10.1186/s12918-016-0261-8

620 Marinovic, B., & Mangel, M. (1999). Krill can shrink as an ecological adaptation to temporarily unfavourable

621   environments. *Ecology Letters*, *2*, 338–343.

622 Maron, M., McAlpine, C. A., Watson, J. E. M., Maxwell, S., & Barnard, P. (2015). Climate-induced resource

623   bottlenecks exacerbate species vulnerability: A review. *Diversity and Distributions*, *21*(7), 731–743.

624   https://doi.org/10.1111/ddi.12339

625 Martínez, D. E. (1998). Mortality Patterns Suggest Lack of Senescence in Hydra. *Experimental Gerontology*,

626   *33*(3), 217–225. https://doi.org/10.1016/S0531-5565(97)00113-7

627 Medvedev, Z. A. (1990). An attempt at a rational classification of theories of ageing. *Biological Reviews*,

628   *65*(3), 375–398. https://doi.org/10.1111/j.1469-185X.1990.tb01428.x

629 Metcalf, C. J. E. (2016). Invisible Trade-offs: Van Noordwijk and de Jong and Life-History Evolution. *The*

630   *American Naturalist*, *187*(4), iii–v. https://doi.org/10.1086/685487

631 Morgan, T. H. (1901). *Regeneration*. The Macmillan Company.

Mysterud, A., Stenseth, N. Chr., Yoccoz, N. G., Langvatn, R., & Steinheim, G. (2001). Nonlinear effects of

large-scale climatic variability on wild and domestic herbivores. *Nature*, *410*(6832), 1096–1099.

https://doi.org/10.1038/35074099

Nelson, P., & Masel, J. (2017). Intercellular competition and the inevitability of multicellular aging.

Proceedings of the National Academy of Sciences, *114*(49), 12982–12987.

https://doi.org/10.1073/pnas.1618854114

Nentwig, M. R., & Schauble, M. K. (1974). Influence of the nutritional state on repeated head regeneration,

growth, and fission in the planarian, Dugesia dorotocephala. *Journal of Experimental Zoology*, *187*(2),

295–302. https://doi.org/10.1002/jez.1401870213

Nicholson, A. (1954). An outline of the dynamics of animal populations. *Australian Journal of Zoology*, *2*(1),

9–65.

Ofoegbu, P. U., Simão, F. C. P., Cruz, A., Mendo, S., Soares, A. M. V. M., & Pestana, J. L. T. (2016). Toxicity of

tributyltin (TBT) to the freshwater planarian Schmidtea mediterranea. *Chemosphere*, *148*, 61–67.

https://doi.org/10.1016/j.chemosphere.2015.12.131

Ozgul, A., Tuljapurkar, S., Benton, T. G., Pemberton, J. M., Clutton-Brock, T. H., & Coulson, T. (2009). The

Dynamics of Phenotypic Change and the Shrinking Sheep of St. Kilda. *Science*, *325*, 464–467.

https://doi.org/10.1126/science.1173668

Pardo, D., Barbraud, C., Authier, M., & Weimerskirch, H. (2013). Evidence for an age-dependent influence of

environmental variations on a long-lived seabird's life-history traits. *Ecology*, *94*(1), 208–220.

https://doi.org/10.1890/12-0215.1

Payne, B. A. I., & Chinnery, P. F. (2015). Mitochondrial dysfunction in aging: Much progress but many

unresolved questions. *SI: Mitochondrial Dysfunction in Aging*, *1847*(11), 1347–1353.

https://doi.org/10.1016/j.bbabio.2015.05.022

Pellettieri, J. (2019). Regenerative tissue remodeling in planarians – The mysteries of morphallaxis. *Planarian*

Regeneration, *87*, 13–21. https://doi.org/10.1016/j.semcdb.2018.04.004

Peter, R., Ladurner, P., & Rieger, R. M. (2001). The Role of Stem Cell Strategies in Coping with Environmental

Stress and Choosing Between Alternative Reproductive Modes: Turbellaria Rely on a Single Cell Type

to Maintain -Individual Life and Propagate Species. *Marine Ecology*, *22*(1-2), 35–51.

https://doi.org/10.1046/j.1439-0485.2001.00743.x

Petersson, E., & Järvi, T. (2000). Both Contest and Scramble Competition Affect the Growth Performance of

Brown Trout, Salmo Trutta, Parr of Wild and of Sea-ranched Origins. *Environmental Biology of Fishes*,

59(2), 211–218. https://doi.org/10.1023/A:1007645411586

R Core Team. (2020). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical

Computing. http://www.R-project.org

Reynoldson, T. B. (1964). Evidence for Intra-Specific Competition in Field Populations of Triclads. *Journal of*

Animal Ecology, *33*, 187–201. JSTOR. https://doi.org/10.2307/2439

Reynoldson, T. B., & Sefton, A. D. (1972). The population biology ofPlanaria torva (Müller) (Turbellaria,

Tricladida). *Oecologia*, *10*(1), 1–16. https://doi.org/10.1007/BF00822758

Rink, J. C. (2013). Stem cell systems and regeneration in planaria. *Development Genes and Evolution*, *223*(1),

67–84. https://doi.org/10.1007/s00427-012-0426-4

Roach, D. A., Ridley, C. E., & Dudycha, J. L. (2009). Longitudinal analysis of Plantago: Age-by-environment

interactions reveal aging. *Ecology*, *90*(6), 1427–1433. https://doi.org/10.1890/08-0981.1

Roff, D. A. (1992). *Life History Evolution*. Sinauer Associates Incorporated.

Romero, R., & Baguñà, J. (1991). Quantitative cellular analysis of growth and reproduction in freshwater

planarians (Turbellaria; Tricladida). I. A cellular description of the intact organism. *Invertebrate*

Reproduction & Development, *19*(2), 157–165. https://doi.org/10.1080/07924259.1991.9672170

Roper, M., Capdevila, P., & Salguero-Gómez, R. (2021). Senescence: Why and where selection gradients

might not decline with age. *Proceedings of the Royal Society B: Biological Sciences*, *288*(1955),

20210851. https://doi.org/10.1098/rspb.2021.0851

Rutstein, A. N., Gilbert, L., Slater, P. J. B., & Graves, J. A. (2005). Sex-specific patterns of yolk androgen

allocation depend on maternal diet in the zebra finch. *Behavioral Ecology*, *16*(1), 62–69.

https://doi.org/10.1093/beheco/arh123

Sæther, B.-E., Coulson, T., Grøtan, V., Engen, S., Altwegg, R., Armitage, K. B., Barbraud, C., Becker, P. H.,

Blumstein, D. T., Dobson, F. S., Festa-Bianchet, M., Gaillard, J.-M., Jenkins, A., Jones, C., Nicoll, M. A.

C., Norris, K., Oli, M. K., Ozgul, A., & Weimerskirch, H. (2013). How Life History Influences Population

Dynamics in Fluctuating Environments. *The American Naturalist*, *182*(6), 743–759.

https://doi.org/10.1086/673497

Sahu, S., Dattani, A., & Aboobaker, A. A. (2017). Secrets from immortal worms: What can we learn about

biological ageing from the planarian model system? *Seminars in Cell & Developmental Biology*, *70*,

108–121. https://doi.org/10.1016/j.semcdb.2017.08.028

Salguero-Gómez, R., & Casper, B. B. (2010). Keeping plant shrinkage in the demographic loop. *Journal of*

Ecology, *98*(2), 312–323. https://doi.org/10.1111/j.1365-2745.2009.01616.x

Salguero-Gómez, R., & Casper, B. B. (2011). A hydraulic explanation for size-specific plant shrinkage:

Developmental hydraulic sectoriality. *New Phytologist*, *189*(1), 229–240.

https://doi.org/10.1111/j.1469-8137.2010.03447.x

Salguero-Gómez, R., Jones, O. R., Jongejans, E., Blomberg, S. P., Hodgson, D. J., Mbeau-Ache, C., Zuidema, P.

A., de Kroon, H., & Buckley, Y. M. (2016). Fast–slow continuum and reproductive strategies structure

plant life-history variation worldwide. *Proceedings of the National Academy of Sciences*, *113*(1), 230–

235. https://doi.org/10.1073/pnas.1506215112

Saló, E. (2006). The power of regeneration and the stem-cell kingdom: Freshwater planarians

(Platyhelminthes). *BioEssays*, *28*(5), 546–559. https://doi.org/10.1002/bies.20416

Schaible, R., Ringelhan, F., Kramer, B. H., & Miethe, T. (2011). Environmental challenges improve resource

utilization for asexual reproduction and maintenance in hydra. *Experimental Gerontology*, *46*(10),

794–802. https://doi.org/10.1016/j.exger.2011.06.004

Smallegange, I. M. (2011). Effects of paternal phenotype and environmental variability on age and size at

maturity in a male dimorphic mite. *Naturwissenschaften*, *98*, 339–346.

https://doi.org/10.1007/s00114-011-0773-4

Sousa, N. de, & Adell, T. (2018). Maintenance of Schmidtea mediterranea in the Laboratory. *Bio-Protocol*,

8(19), e3040. https://doi.org/10.21769/BioProtoc.3040

Stearns, S. C. (1992). *The Evolution of Life Histories*. Oxford University Press.

Stewart, C., & Tabak, J. (2011). Size-based dominance hierarchies in the New Zealand freshwater crayfish

(koura) Paranephrops zealandicus. *New Zealand Journal of Marine and Freshwater Research*, *45*(2),

281–285. https://doi.org/10.1080/00288330.2011.559661

Szabo, A. R. (2002). Experimental tests of intercohort competition for food and cover in the tidepool sculpin

(Oligocottus maculosus Girard). *Canadian Journal of Zoology*, *80*(1), 137–144.

https://doi.org/10.1139/z01-218

Thommen, A., Werner, S., Frank, O., Philipp, J., Knittelfelder, O., Quek, Y., Fahmy, K., Shevchenko, A.,

Friedrich, B. M., Jülicher, F., & Rink, J. C. (2019). Body size-dependent energy storage causes Kleiber's

law scaling of the metabolic rate in planarians. *eLife*, *8*, e38187. https://doi.org/10.7554/eLife.38187

Vaupel, J., Baudisch, A., Dölling, M., A. Roach, D., & Gampe, J. (2004). The case for negative senescence.

Demography in the 21st Century, *65*(4), 339–351. https://doi.org/10.1016/j.tpb.2003.12.003

Vaupel, J. W. (2010). Biodemography of human ageing. *Nature*, *464*, 536–542.

https://doi.org/10.1038/nature08984

Vaupel, J. W., & Yashin, A. I. (1985). Heterogeneity's Ruses: Some Surprising Effects of Selection on

Population Dynamics. *The American Statistician*, *39*(3), 176–185. https://doi.org/10.2307/2683925

Vila-Farré, M., & C. Rink, J. (2018). The Ecology of Freshwater Planarians. In J. C. Rink (Ed.), *Planarian*

Regeneration: Methods and Protocols (pp. 173–205). Springer New York.

https://doi.org/10.1007/978-1-4939-7802-1_3

Ward, A. J. W., Webster, M. M., & Hart, P. J. B. (2006). Intraspecific food competition in fishes. *Fish and*

Fisheries, *7*(4), 231–261. https://doi.org/10.1111/j.1467-2979.2006.00224.x

Wikelski, M., & Thom, C. (2000). Marine iguanas shrink to survive El Niño. *Nature*, *403*(6765), 37–38.

https://doi.org/10.1038/47396

Williams, T. D. (1996). Variation in Reproductive Effort in Female Zebra Finches (Taeniopygia guttata) in

Relation to Nutrient-Specific Dietary Supplements during Egg Laying. *Physiological Zoology*, *69*(5),

1255–1275. JSTOR.

Table 1. Hypotheses of the impact of the variability in three different dimensions of resource availability (quantity, quality, and feeding interval) on
population structure, population performance, and life history trait trade-offs in two species of Planaria; *Schmidtea mediterranea* and *Dugesia*
tahitiensis. Here we show results for each species, supporting (✔) or rejecting (✖) each hypothesis. High quality (🟠); low quality (🟣); high intake
(🔘🔘); reduced intake (⚫) ; feeding intervals every 14 days (14-Day); feeding intervals every seven days (7-Day).

Resource availability	Prediction	Outcome	
		D. tahitiensis	*S. mediterranea*
H1: High quantity, low quality environments 🟣 🔘🔘 **7-Day**	Population size will increase, and individual size will be normally distributed. High quantity, low quality environments are richer in protein; these environments typically promote reproduction (Lee, 2015) and, as such, maintain population viability (Bayliss & Choquenot, 2002).	✔	✖
H2: Low quantity, high quality environments 🟠 ⚫ **14-Day**	Population size will decrease and be dominated by larger individuals skewing the population size structure to the right. Environments with fewer, but higher quality resources, will likely be dominated by larger individuals that outcompete smaller individuals (Szabo, 2002; Ward et al., 2006). In environments with fewer high quality resources, available resources will be preferentially allocated to maintenance because, in organisms with indeterminate growth like planarians, energy allocation is typically prioritised for maintenance over reproduction and development (Jokela & Mutikainen, 1995), thus maintaining population viability.	✖	✔

| H3: Environments with more resources of higher quality

 7-Day | There will be an increase in clonal reproduction and subsequently larger population sizes. Where resource supply is high and constant, populations support higher numbers of individuals, typically decreasing population extinction risk (Bayliss & Choquenot, 2002; Lande et al., 2003)

 | ✔ | ✘ |

BOX 1 Glossary

HQ-HI-7D: Populations provided with high quality (HQ), high intake (HI) resources at intervals of
every seven days.

HQ-HI-14D: Populations provided with high quality (HQ), high intake (HI) resources at intervals of
every 14 days.

HQ-SI-7D: Populations provided with high quality (HQ), standard intake (SI) resources at intervals of
every seven days.

HQ-SI-14D: Populations provided with high quality (HQ), standard intake (SI) resources at intervals
of every 14 days.

HQ-RI-7D: Populations provided with high quality (HQ), reduced intake (RI) resources at intervals of
every seven days.

HQ-RI-14D: Populations provided with high quality (HQ), reduced intake (RI) resources at intervals
of every 14 days.

LQ-HI-7D: Populations provided with low quality (LQ), high intake (HI) resources at intervals of
every seven days.

LQ-HI-14D: Populations provided with low quality (LQ), high intake (HI) resources at intervals of
every 14 days.

LQ-SI-7D: Populations provided with low quality (LQ), standard intake (SI) resources at intervals of
every seven days.

LQ-SI-14D: Populations provided with low quality (LQ), standard intake (SI) resources at intervals of
every 14 days.

LQ-RI-7D: Populations provided with low quality (LQ), reduced intake (RI) resources at intervals of
every seven days.

LQ-RI-14D: Populations provided with low quality (LQ), reduced intake (RI) resources at intervals of
every 14 days.

Figure legends

Figure 1. Full factorial experimental design to examine how three key dimensions of resource

(quality, quantity, and interval of resource availability) shape population size, population structure

and life history trait trade-offs of two flatworm species, *Schmidtea mediterranea* and *Dugesia*

tahitiensis. Resource quality has two levels (top right panel): high quality (*HQ*; orange dots) and low

quality (*LQ*; purple dots). Resource quantity has three levels (middle right panel): restricted intake

(*RI*; single grey dot), standard intake (*SI*; three grey dots) and hyper caloric intake (*HI*; six grey dots).

Resource interval has two levels (bottom right panel): every seven days (*7D*; black line), every 14

days (*14D*; grey line). Each population was initiated with 10 individuals and replicated five times

resulting in a total of 600 individuals and 60 populations per species. Populations were counted and

photographed weekly.

Figure 2. Population structure in *Schmidtea mediterranea* populations changes across time in most

treatments. Weekly size spectra slope values (*b*) for *S. mediterranea*. A) Calculated size spectra

slope (*b*) values (black points) for all populations for high quality (*HQ*,) treatments over 19 weeks.

B) Calculated size spectra slope (*b*) values (black points) for all populations for low quality (*LQ*, ●)

treatments over 19 weeks. Blue points indicate estimated *b* values with compatibility intervals

(0.95) for each block (block 1 – light blue; block 2 – blue; block 3 – dark blue). Left panels in each

graph indicate feeding intervals every 14 days and right panels feeding intervals every seven days.

Top panel represent high intake (*HI*,), middles panel standard intake (*SI*,) and bottom

panel reduced intake (*RI*,) treatments.

Figure 3. Population structure in *Dugesia tahitiensis* populations has little change in structure

across calorie restriction treatments. Weekly size spectra slope values (*b*) for *D. tahitiensis*. A)

Calculated size spectra slope (*b*) values (black points) for all populations for high quality (*HQ*,   )

treatments over 19 weeks. B) Calculated size spectra slope (*b*) values (black points) for all

populations for low quality (*LQ*,   ) treatments over 19 weeks. Blue points indicate estimated *b*

values with compatibility intervals (0.95) for each block (block 1 – light blue; block 2 – blue; block 3

– dark blue). Left panels in each graph indicate feeding intervals every 14 days and right feeding

intervals every seven days. Top panel represent high intake (*HI*,   ), middles panel standard

intake (*SI*,   ) and bottom panel reduced intake (*RI*,   ) treatments.

Figure 4. Population size for *Schmidtea mediterranea* varies but with no clear patterns across

calorie restriction treatments. Weekly population size of *S. mediterranea* over the 19 weeks for A)

high quality (*HQ*,   ) treatments and B) low quality (*LQ*,   ) treatments. Black points indicate raw

data of weekly population size; blue lines indicate estimated population size for each block (block 1

– light blue; block 2 – blue; block 3 – dark blue). Shaded grey regions indicate compatibility

intervals: light grey 0.95, grey 0.8, dark grey 0.5. Left panels indicate feeding intervals every 14 days

and right panels feeding intervals every seven days. Top panel represent high intake (*HI*,   ),

middles panel standard intake (*SI*,   ) and bottom panel reduced intake (*RI*,   ) treatments.

Figure 5. Population size for *Dugesia tahitiensis* varies but with no clear patterns across calorie

restriction treatments. Weekly population size of *D. tahitiensis* over the 19 weeks for A) high quality

(*HQ*,   ) treatments and B) low quality (*LQ*,   ) treatments. Black points indicate raw data of

weekly population size; blue lines indicate estimated population size for each block (block 1 – light

blue; block 2 – blue; block 3 – dark blue). Shaded grey regions indicate compatibility intervals: light

grey 0.95, grey 0.8, dark grey 0.5. Left panels indicate feeding intervals 14 days and right

panels feeding intervals every seven days. Top panel represent high intake (*HI*,), middles

panel standard intake (*SI*,) and bottom panel reduced intake (*RI*,) treatments.

    Figure 1

Figure 2

834   **Figure 3**

Figure 4

Figure 5