Title page:

Learning functional conservation between pig and human to decipher evolutionary

mechanisms underlying gene expression and complex trait

Jinghui Li[1], Tianjing Zhao[1], Dailu Guan[1], Zhangyuan Pan[1], Zhonghao Bai[2], Jinyan Teng[3], Zhe

Zhang[3], Zhili Zheng[4,5], Jian Zeng[4], Huaijun Zhou[1], Lingzhao Fang[2]*, Hao Cheng[1]*

[1]Department of Animal Science, University of California, Davis, Davis, CA, USA

[2]Center for Quantitative Genetics and Genomics (QGG), Aarhus University, Aarhus, Denmark

[3]Guangdong Laboratory of Lingnan Modern Agriculture, National Engineering Research Center

for Breeding Swine Industry, Guangdong Provincial Key Lab of Agro-Animal Genomics and

Molecular Breeding, College of Animal Science, South China Agricultural University,

Guangzhou, China

[4]Institute for Molecular Bioscience, The University of Queensland, Brisbane, Queensland,

Australia

[5]Program in Medical and Population Genetics, Broad Institute of Harvard and MIT, Cambridge,

Massachusetts, USA

*Corresponding authors:

HC: Department of Animal Science, University of California, Davis, Davis, CA, USA

Email: qtlcheng@ucdavis.edu

LF: Center for Quantitative Genetics and Genomics (QGG), Aarhus University, Aarhus,

Denmark

Email: lingzhao.fang@qgg.au.dk

Abstract

The assessment of genomic conservation between human and pig at the functional level can help understand and improve the potential of pig as a human biomedical model. To address this, we developed a **Deep** learning-based approach to learn the **G**enomic **C**onservation at the **F**unctional level (DeepGCF) between species by integrating 386 and 374 epigenome and transcriptome profiles from human and pig, respectively. DeepGCF demonstrated a better prediction performance compared to the previous functional conservation prediction method. In addition, we showed that the resulting DeepGCF score captures the functional conservation by examining DeepGCF on chromatin states, sequence ontologies, and regulatory variants. Regions with higher DeepGCF score play a more important role in regulatory activities and show heritability enrichment in human complex traits and diseases. Our DeepGCF approach shows a promising application on the comparison of cross-species functional conservation, and the model framework can be easily adapted to other species. By expanding the model to integrate the functional profiles of multiple species, including human, mouse, pig, cattle, and other livestock animals in the future, the functional conservation information will provide additional insight into the genetic and evolutionary mechanisms behind complex traits and diseases.

Main text

Introduction

Comparative genome not only reveals evolutionary changes at the DNA sequence level[1], but

also helps with the translation of genetic and biological findings across species[2]. Compared to

lab organisms like mice, pig is more similar to human in anatomy, physiology, and genome[3],

thus is widely used as a biomedical model for human medicine and genetic diseases, such as

drug tests[4], xenotransplantation[5], Alzheimer's disease[6], breast cancer[7], and diabetes[8]. To fully

recognize the substantial potential of pig as a human biomedical model, it is essential to conduct

an extensive comparison of pig and human physiology at the molecular level for assessing to

what degree that the genetic and biological findings in pig can be extrapolated to human. Several

methods have been proposed to infer the conservation at the DNA sequence level, such as

Genomic Evolutionary Rate Profiling (GERP)[9] and Phylogenetic P-values (PhyloP)[10]. However,

the conservation at DNA sequence level is not equivalent to the conservation at functional

level[11–13].

The ongoing global efforts on functional annotation of genomes in both humans and

livestock, such as the Encyclopedia of DNA Elements[14], Roadmap Epigenomics projects[15], the

Functional Annotation of Animal Genomes (FAANG)[16], and Farm animal Genotype-Tissue

Expression (FarmGTEx) projects[17], provide an unprecedented opportunity to quantify the

genome conservation across species at the functional level. Previous studies often rely on a

single functional profile in one tissue/cell type, such as gene expression[18] or epigenome[19,20], to

infer the functional conservation of orthologous regions between human and pig. However,

integrative analysis of multi-omics is essential for unravelling how biological information

encoded in the genome is conserved or diverged across species, as the functional consequence of

genomic variants is often modulated at multiple levels of gene regulation across tissues/cells.

Artificial neural networks have been applied in the prediction and integration of multi-omics

data, such as histone marks, transcription factors, and gene expression, to investigate

transcriptional and biochemical impact of DNA sequences and their conservation across

species[21,22]. For instance, Kwon and Ernst[22] developed a neural network model, LECIF, to study

human-mouse functional conservation based on multi-omics data from Roadmap and ENCODE

databases.

In this study, to systematically evaluate the functional conservation between human and pig,

we developed a **Deep** learning-based approach to learn the **G**enomic **C**onservation at the

Functional level (DeepGCF) between species. Unlike LECIF using functional genomics data as

input, DeepGCF uses both DNA sequences and functional genomics data as input. It thus enables

us to predict the impact of sequence mutations on the functional conservation between species.

By integrating 386 and 374 epigenome and transcriptome profiles, representing 28 and 21 tissues

from human and pig, respectively, DeepGCF captures the conservation of epigenetic features and

genes across tissues between human and pig. By further examining expression/splicing

quantitative trait loci (e/sQTL) from 54 and 35 tissues in human GTEx[23] and PigGTEx[24],

respectively, and genome-wide association studies (GWAS) of 80 complex traits/diseases in

human, DeepGCF provides novel insights into the evolutionary mechanisms underlying both

molecular phenotype and complex trait variation. The DeepGCF model can be easily expanded

to multiple species for extensively understanding the genome evolution at functional genomics

level when large-scale functional annotation data is available for many other species in the near

feature.

Results

Overview of the DeepGCF model

In general, the training of DeepGCF model consists of two steps (**Fig. 1**). The first step is to

transform the binary functional features to continuous values by training a deep convolutional

network implemented in DeepSEA[25]. Binary functional feature is a common data type in the

functional genomics filed, which represents whether a genomic base overlapped with functional

annotations such as peaks or chromatin states derived from ATAC-Seq and ChIP-Seq. By taking

both DNA sequences and binary functional features as inputs, DeepSEA predicts the

probabilities of each functional feature at a single-nucleotide resolution. In this study, we

collected 309 and 294 genome-wide binary functional annotations from human and pig,

respectively (**Supplementary data 1–4**). These represented the chromatin accessibility measured

by Assay for Transposase-Accessible Chromatin (ATAC-seq), histone modifications measured

by Chromatin Immunoprecipitation sequencing (ChIP-seq) and chromatin states from 26 and 21

tissues in human and pig, respectively. We trained the DeepSEA models and predicted the

functional effect of each nucleotide in human and pig separately, which were subsequently used

as inputs in the DeepGCF for predicting the functional conservation score between these two

species. The performance of DeepSEA was evaluated using an independent validation set and

showed a strong predictive power in both species (**Supplementary Fig. 1**).

The second step of DeepGCF is to predict the functional conservation score of orthologous

regions between human and pig using a supervised deep learning approach, similar to LECIF[22].

We divided the whole-genome alignment between human and pig into non-overlapping 50-bp

regions within each alignment block, resulting in 38,961,848 paired alignments (i.e., orthologous

regions). We then selected the first base to represent the functional annotation of the 50-bp

region, because bases within such a narrow region are likely to have similar functions and the

computational burden is greatly lightened by doing so[22]. Apart from the predicted functional

effects from DeapSEA, we also included the gene expression values from 77 and 80 RNA-seq

datasets as functional annotations, representing 11 and 19 tissues in human and pig, respectively

(**Supplementary data 5** and **6**). To train the DeepSEA model, we randomly shifted the human-

pig orthologous regions to obtain the same number of non-orthologous pairs. Functional

conservation is lack of ground truth, thus as an approximation, we presume that the orthologous

regions (coded as 1) are more likely to be functionally conserved than non-orthologous regions

(coded as 0). We then trained a pseudo-Siamese neural network model[26] using both functional

effects predicted from DeepSEA and gene expression as inputs (**Fig. 1a**). We weighted non-

orthologous regions 50 times more than orthologous ones when training to highlight regions with

strong evidence of functional conservation[22]. The output, DeepGCF score, is a value between 0

and 1 quantifying the functional conservation of the paired human-pig region. Furthermore, since

the DeepGCF predicts the functional conservation based on the DNA sequence, it allows us to

conduct an *in silico* mutagenesis analysis to assess the impact of orthologous variants on the

functional conservation between species through investigating the changes of DeepGCF score

caused by a mutation (**Fig. 1b**).

The evaluation of DeepGCF model

The performance of DeepGCF was evaluated by predicting whether the paired human-pig

regions of an independent testing set are orthologous or not. Compared to LECIF, which had the

areas under receiver operating characteristic curve (AUROC) and precision-recall curve

(AUPRC) of 0.80 and 0.79, respectively, DeepGCF showed a better predictive ability with

AUROC and AUPRC of 0.89 and 0.87, respectively (**Figs. 2a, b**). Of note, we normalized the

gene expression values with a natural logarithm transformation, which showed a better predictive

ability than that without a transformation (**Supplementary Fig. 2**). Among all the 38,961,848

orthologous regions between human and pig, only a small percentage (1.2%) exhibited a

DeepGCF score greater than 0.8, while more than half with a score less than 0.1 (**Fig. 2c**),

consistent with previous findings between human and mice[22]. This result suggests that most of

orthologous regions were not functionally conserved between species.

To provide suggestions for researchers who are interested in running the DeepGCF model in

other species with limited functional annotation data available, we explored different features

that may influence the performance of DeepGCF, including sample size and diversity of

functional annotations regarding array and tissue/cell type. When training the model, we

downsampled both human and pig functional profiles. We found that using ~50% (Human: 192;

Pig: 187) and ~10% (Human: 52; Pig: 47) of the functional profiles resulted in similar AUROC

(50%: 0.88; 10%: 0.85) and AUPRC (50%: 0.87; 10%: 0.83) values compared to using all the

profiles, but using only ~1% (Human: 4; Pig: 4) of the profiles showed substantially lower

AUROC (0.69) and AUPRC (0.68) values (**Fig. 2d**). When leaving one type of functional

profiles out, the predictive ability of DeepGCF did not change too much (**Fig. 2e**).

Relationship between DNA sequence conservation and functional conservation

To fully explore whether DNA sequence conservation indicates functional conservation, we

first examined PhyloP scores, which are commonly used to measure the DNA sequence

conservation across species[10]. We observed a U-shaped relationship between PhyloP and

DeepGCF scores (**Fig. 3a**), demonstrating that both fast-evolving and slow-evolving sequences

exhibited a higher functional conservation between species, compared to evolutionary neutral or

near-neutral sequences. This agrees with previous findings on comparing individual epigenetic

marks and DNA sequence conservation[19,27]. Furthermore, we defined three types of orthologous

regions according to their PhyloP and DeepGCF scores to represent the two tails and the bottom

of the U curve: 1) regions with both high DeepGCF (> 95th percentile) and PhyloP (> 95th

percentile): high D & high P (n = 260,281), 2) those with high DeepGCF (> 95th percentile) but

low PhyloP (< 5th percentile): high D & low P (n = 152,557), and 3) those with low DeepGCF (<

5th) and medium PhyloP (between 47.5th and 52.5th): low D & med P (n = 95,231). By examining

sequence classes, which are predicted regulatory activities of DNA sequences in human genome

by a deep learning model, Sei, trained on a compendium of 21,907 epigenome profiles[28], and

Gene Ontology (GO) terms, we found that, compared to the whole genome, high D & high P

regions were more enriched in promoter, CTCF, and transcription but depleted in enhancer

(Binomial test P < 0.0001; **Fig. 3b**). Compared to other regions, high D & high P regions showed

a higher enrichment in transcription (Binomial test P < 0.0001; **Fig. 3b**), and were significantly

associated with several RNA-related regulation processes (**Supplementary Data 7**). This

indicates the similarities in transcriptional networks between pig and human[18,29]. High D & low

P regions were significantly enriched in Polycomb (Binomial test P < 0.0001; **Fig. 3b**), in

consistency with the fact that some core subunits of Polycomb protein complexes with similar

biological functions have shown a weak evolutionary conservation on DNA sequence across

species[30]. The low D & med P regions had similar sequence class compositions as the whole

genome background except promoter, which was enriched but to a less extent than high D &

high P and high D & low P (Binomial test P < 0.0001; **Fig. 3b**), and were enriched in fewer GO

terms than regions with high DeepGCF (**Supplementary Data 7–9**). In addition, we examined

six different sequence ontologies and found that 5' UTR is the most functionally conserved

element, followed by start codon, 3' UTR, stop codon, exon, and finally intron. This is consistent

between both human and pig (**Fig. 3c**).

To investigate the impact of orthologous variants on the functional conservation between

species, we examined 35,575,835 human SNPs that are located in orthologous regions between

human and pig, which were obtained from the 1,000 Genome Project[31]. We used the DeepGCF

model trained based on only predicted probabilities of binary features from DeepSEA (i.e.,

leaving RNA-seq out), as the DeepSEA model does not predict for continuous functional

features. The new score predicted from DeepGCF without RNA-seq data had a relatively well

agreement with the original DeepGCF score with a Pearson's correlation coefficient (PCC) of

0.74 (**Supplementary Fig. 3**). To measure the effect of each human SNP on functional

conservation, we recomputed the probabilities of binary features for the corresponding

orthologous human region due to the SNP mutation and kept the pig probabilities the same, and

used the new probabilities to calculate the updated DeepGCF score. The effect on functional

conservation is measured by ΔDeepGCF = DeepGCF after SNP mutation – original DeepGCF. By

classifying all the orthologous variants into eight categories[28], we found that most of the variants

had a limited effect on the functional conservation (**Fig. 3d**). We further grouped them into 40

sequence classes[28], and in general, we found that variants in functional features with larger

DeepGCF scores showed the stronger effects on the functional conservation between species

(**Fig. 3e**). Promoter and CTCF were more sensitive to variants than other sequence classes. Of

note, the average DeepGCF score of CTCF is lower than that of promoter, but it is much more

sensitive to genetic mutations regarding the functional conservation, indicating that the genetic

disruption of CTCF binding sites (chromatin conformation) may cause strong impacts on

functional genome evolution between species by altering the genome topology and consequently

the gene expression[32,33].

DeepGCF captures the evolutionary characteristics of regulatory elements

To investigate the functional conservation of distinct regulatory elements between pig and

human, we first examined the DeepGCF score of 15 chromatin states predicted from 14 pig

tissues and 12 human tissues using ChromHMM[19]. We found that strongly active promoters

showed the highest DeepGCF scores (i.e., the strongest functional conservation), followed by

poised transcription start site (TSS), chromatin states proximal to TSS, enhancers, and finally

repressed Polycomb (**Fig. 4a**). This was consistent between human and pig, which agrees with

the conservation properties of regulatory elements reported in the previous studies[19,34]. As

chromatin states that play important roles in determining the cellular functions may vary among

different tissues, we identified strongly active promoters and enhancers that were specific in each

of 12 human tissues and 14 pig tissues. Compared to promoters and enhancers shared across all

the tissues, tissue-specific ones showed significantly lower DeepGCF scores in both species

(Mann–Whitney U test $P < 2.2e\text{-}16$), indicative of their faster evolutionary rate (**Fig. 4b**). Among

eight common tissues between human and pig, we found that adipose had the strongest

functionally conserved promoters in both human and pig, followed by spleen, lung, cortex, liver,

and finally stomach (**Supplementary Fig. 4a**). This result suggests pigs could be a good model

animal for studying human obesity and metabolic traits[19]. However, the tissue-conservation

patterns of enhancers were different from those of promoters and were not consistent between

species (**Supplementary Fig. 4b**).

We further investigated the DeepGCF score on human promoters and enhancers annotated by

Sei[28]. We linked a promoter to its potential target gene and then ranked genes with the DeepGCF

scores of their promoters (from largest to smallest). We found that top 5% of genes were

significantly enriched in basic biological processes, such as anatomical structure development

and organ morphogenesis, whereas bottom 5% of genes were significantly enriched in

biosynthetic and metabolic process (**Supplementary data 10 and 11**). In addition, we ranked

enhancers according to their own DeepGCF scores and investigated the function of top 5% and

bottom 5% enhancers. Unlike promoters, top 5% of enhancers exhibited the most significant

enrichment in metabolic processes, while bottom 5% of enhancers were significantly enriched in

organ growth and development (**Supplementary data 12 and 13**). In general, we found that

promoters and enhancers with a higher DeepGCF score were enriched in much more biological

processes compared to those with a lower DeepGCF score (**Fig. 4c, d**), which indicates that

functionally conserved regions between species tend to be the hotspot of regulatory activities.

DeepGCF provide insight into the functional conservation of regulatory variants

To explore the functional conservation of regulatory variants, we systematically examined

expression QTLs (eQTLs) and splicing QTLs (sQTLs) falling in the orthologous regions in 54

human tissues and 35 pig tissues, respectively. In general, DeepGCF scores of eQTLs and sQTLs

were significantly (Mann–Whitney U test $P < 2.2e\text{-}16$) higher than the genome background

across all the tissues in both human and pig (**Fig. 5a; Supplementary Figs. 5 and 6**), which

suggests that regulatory variants are functionally conserved between species[35,36]. Of note, sQTLs

showed a higher DeepGCF score than eQTLs in both species (Mann–Whitney U test $P < 1e\text{-}8$),

probably due to their larger impacts on the transcriptome function (underlying a stronger

purifying selection). This is consistent with previous findings that sQTLs were more likely to be

enriched in 5'UTR than eQTLs (GTEx, 2020), and 5' UTR is the most functionally conserved

genomic features (**Fig. 2c**). We further observed that eGenes associated with eQTLs having a

larger absolute effect on the gene expression had a lower DeepGCF score in both species (**Fig.**

5b), which suggests that orthologous regions with smaller regulatory effects are more likely to be

functionally conserved between species, probably due to the stronger purifying selection

underlying them[37]. Moreover, regulatory variants influencing more tissues showed higher

DeepGCF scores (i.e., more functionally conserved), consistent in human and pig (**Fig. 5c, d**). In

addition, the tissue-sharing pattern of orthologous eGenes (PCC = 0.38, *P* value < 2.2e-16) and

sGenes (PCC = 0.45, *P* value < 2.2e-16) were positively correlated between human and pig.

Altogether, these results indicate that regulatory variants controlling transcriptome function in

more tissues tend to be more functionally conserved between species.

We then investigated the DeepGCF scores of 105,461 pathological and likely pathological

SNPs obtained from the ClinVar database[38]. A total 98.6% of these SNPs were in the human-pig

orthologous regions, consistent with a previous finding that reported more than 98% of

pathological variants of Mendelian diseases located in human-mouse orthologous regions[39].

Compared to random orthologous regions, these pathological SNPs were significantly more

functionally conserved (Mann–Whitney U test *P* < 2.2e-16; **Fig. 6a**). Like orthologous SNP, we

classified the ClinVar SNP into eight sequence class categories[28] and conducted an *in silico*

mutagenesis analysis to predict their impact on the functional conservation. Overall, the average

magnitude of variant effect (measured by |ΔDeepGCF|) for pathological and likely pathological

mutations is 1.5 times larger than that for random orthologous SNPs (0.0088 versus 0.0058,

Mann–Whitney U test *P* < 2.2e-16). In most of cases, the DeepGCF score did not change much

after genetic mutations, but the variance of ΔDeepGCF showed a bell-shaped curve regarding the

original DeepGCF score, indicating that SNPs with a medium-high DeepGCF (50^{th} to 80^{th}

percentile) were more sensitive to pathological mutations than those with lower or higher

DeepGCF (**Fig. 6b**). This suggests that the most functionally conserved regions (> 90^{th}

percentile) are more tolerable of mutations than less conserved ones (50^{th} to 80^{th} percentile).

Most of the ClinVar SNPs were classified as transcription (51.2%), followed by enhancer

(16.4%), Polycomb (14.8%), promoter (8.8%), transcription factor (3.3%), and CTCF (2.2%;

Fig. 6c). Among the ClinVar SNPs with top 5% of |ΔDeepGCF| (> 0.03), there were more SNPs

relevant to a decreased DeepGCF (54.4%) than an increased one (45.6%). Moreover, 9 out of 10

ClinVar SNPs with the largest effect on DeepGCF were relevant to a decreased DeepGCF (**Fig.**

6c). In summary, pathological and likely pathological SNPs are located in functionally more

conserved regions, and their impact on functional conservation tends to be related to a decreased

functional conservation between human and pig.

Application of DeepGCF on gene mapping and prediction for human complex traits

To investigate whether DeepGCF scores could advance our understanding of the evolutionary

basis of complex traits/diseases in human, we conducted a heritability partitioning analysis used

the functionally conserved genomic regions (top 5% DeepGCF scores) as a functional

annotation, along with 97 existing annotations from the baseline model of LDSC[40,41], to analyze

the GWAS summary statistics from 80 human complex traits/diseases (**Supplementary Data**

14). We found that regions with higher DeepGCF scores explained more heritability of complex

traits/diseases (**Fig. 7a**). The heritability of eight complex traits was significantly enriched in

functionally conserved regions, with the most enrichment found for coxarthrosis (enrichment =

3.5, FDR = 0.032), followed by varicose veins, height, hypertension, primary hypertension,

waist-hip ratio, weight, and BMI (**Supplementary Data 15; Fig. 7b**). Furthermore, we took

these eight traits as examples to explore whether DeepCGF can help us with fine-mapping of

causal variants. By using functionally conserved regions (top 5% of DeepCGF) as a biological

prior in the PolyFun + SuSiE model[42], we detected 33, 22, and 17 additional putative causal

variants (PIP > 0.95 and P < 5e-8) compared to the SuSiE model only without any priors in

height, BMI and weight, respectively (**Fig. 7c,** Supplementary Data 16). We further incorporated

DeepCGF in SBayesRC[43] model to conduct polygenic score prediction for 20 human complex

traits (**Supplementary Data 17**). On average, the relative prediction accuracy increased by

0.56% (**Fig. 7d; Supplementary Data 18**), and the largest increase was observed on waist-hip

ratio (3.5%), followed by body weight (1.7%). Altogether, our results showed that DeepGCF

provide additional insights into the genetic and evolutionary basis of complex phenotypes.

Discussion

In this study, we developed a two-step neural network approach, DeepGCF, to evaluate the

genomic conservation at the functional level between human and pig. DeepGCF shares a similar

model structure as LECIF[22] in the evaluation of functional conservation by comparing the

epigenome and gene expression profiles of orthologous regions between two species. But instead

of using binary epigenome profiles as the direct inputs, DeepGCF first predicts their functional

effects (i.e., the continuous probability score of each epigenome binary feature) using

DeepSEA[25], and then use them as the input to predict the functional conservation between

species. Compared to the LECIF approach, DeepSEA showed a better performance in the

ortholog prediction, probably due to a higher resolution of the model input. Similar to LECIF, we

found that the performance of DeepGCF was not sensitive to the number of functional features,

indicating that DeepGCF could be applied on other species where functional features are not

abundant.

We demonstrated that functional conservation is different from sequence conservation. The

relationship between DeepGCF and PhyloP scores confirms the U shape relationship between

functional and sequence conservation. By examining DeepGCF on chromatin states, sequence

ontologies, and regulatory variants, we verified that DeepGCF captures the functional

conservation of genome, and regions with higher DeepGCF play a more important role in

regulatory activities. We thereby expected DeepGCF to be useful in explaining complex traits

and diseases. The heritability enrichment and polygenic prediction accuracy brought by

functionally conserved regions were limited, this may because we only considered functional

conservation between human and pig compared to sequence conservation which were obtained

based on over 100 species[44]. With the increasing amount of epigenome and gene expression data

in other species in the near future, we could identify the core functionally conserved regions by

expanding the DeepGCF model structure to integrate functional profiles from multiple species.

Another limitation is that the functional conservation of the same sequence segment in different

tissues and cell types should be conceptually different, which could not be distinguished by the

current DeepGCF score. One ideal way to obtain tissue- and cell-type- specific DeepGCF scores

is to train a different model on each tissue and cell type using the respective data. However, the

current volume of functional profiles, particularly in pig, does not support the development of

tissue- and cell-type- specific DeepGCF models.

Despite the limitations, the DeepGCF approach shows a promising application on the

comparison of cross-species functional conservation. The model framework can be easily

adapted to other species. Our future work will focus on expanding the model to the comparison

of multiple species, including human, mouse, pig, cattle, and other livestock animals. The

functional conservation information among different species will provide additional insight into

the genetic and evolutionary mechanisms behind complex traits and diseases, analogous to the

sequence conservation among vertebrate animals provided by such as PhyloP score.

Methods

Genome alignment. We used the chained and netted alignments of human (GRCh38) and pig

(susScr11) genome assemblies from the UCSC genome browser[45]. The assemblies were aligned

by the lastz alignment program[46] using human as the reference.

Model inputs. We divided the whole-genome alignment between human and pig into non-

overlapping 50-bp regions within each alignment block, resulting in 38,961,848 orthologous

pairs. If an alignment block ended shorter than a 50-bp window, the window was truncated to the

end of the block, which resulted in some regions smaller than 50 bp. For each orthologous pair,

we collected the corresponding functional features, including chromatin accessibility measured

by Assay for Transposase-Accessible Chromatin (ATAC-seq), histone modifications measured

by Chromatin Immunoprecipitation sequencing (ChIP-seq), chromatin state annotations

(ChromHMM), and gene expression measured by RNA-seq for human and pig from public

resources, including ENCODE[14] and public literatures[19,20]. We only collected the functional data

at the tissue level for human, and merged those of the same data type from the same tissue, so

that the total number of human features were close to pig. For human, there were 604 ChIP-seq

and ATAC-seq files merged into 129 features, 12 ChromHMM files of 15 chromatin states (12 ×

15 = 180 features), and 77 RNA-seq features, which resulted in 386 functional annotations. For

pig, there were 287 ChIP-seq and ATAC-seq files merged into 84 features, 14 ChromHMM files

of 15 chromatin states (14 × 15 = 210 features), and 80 RNA-seq features, which resulted in 374

functional annotations. Details of features from each data type are reported in Supplementary

Data 1–6.

Prediction of binary functional features based on DeepSEA. We trained two DeepSEA

models to predict the binary functional features, including ATAC-seq, ChIP-seq and chromatin

state annotations, of human and pig using the PyTorch-based package, Selene[47]. We used the

peak calls of ATAC-seq and ChIP-seq, and one-hot encoded chromatin state annotations as the

training input. We then trained the model based on a sequence region of 1,000 bp, and the feature

must take up 50% of the center bin (200 bp) for it to be considered a feature annotated to that

sequence. All the hyperparameters were set as default (Supplementary Data 19). We created a

validation set using the data from chromosomes 6 and 7 for early stopping during training, a test

set using the data from chromosomes 8 and 9 for the generation of the receiver operating

characteristic (ROC) and precision-recall (PR) curves, and a training set using the rest data. We

then predicted the probability of each binary feature using the trained model for the first base of

all the paired regions that were at most 50 bp.

Data subsets for training and evaluation. We divided the entire data into the training,

validation, and prediction sets based on the chromosome number. To predict the DeepGCF score

of human regions from even and X chromosomes (prediction set), and the corresponding paired

pig regions, we trained a DeepGCF model based on paired regions from a subset of odd

chromosomes of human and pig. We created a validation set also from another subset of odd

chromosomes (not overlapping with the training set) for the hyper-parameter tuning and early

stopping during training. We used a subset of the test set to generate the ROC and PR curves. To

predict the DeepGCF score of human regions from odd chromosomes and the corresponding

paired pig regions, we created training and validation set similarly as above, except from even

chromosomes. We excluded Y and mitochondrial chromosomes in this study. Detailed division

of each set is shown in Supplementary Data 20.

DeepGCF training. Before training the DeepGCF model, we first randomly paired up the

human-pig orthologous regions to get the same number of non-orthologous pairs in the training

set. We then trained the DeepGCF model with a pseudo-Siamese architecture as the LECIF

model[22]. In our pseudo-Siamese neural network, for each orthologous/non-orthologous pair, two

input vectors containing the human and pig binary features (probabilities between 0 and 1)

predicted from DeepSEA and normalized RNAseq data (also between 0 and 1) were connected

to the human and pig subnetworks, respectively (Fig. 1). We performed a natural logarithm

transformation on RNAseq data given the large range before normalizing. The two subnetworks

were then fully connected to a final subnetwork, which generated the output prediction. We

weighted non-orthologous pairs 50 times more than orthologous ones during the training process.

We conducted a random grid search for hyper-parameters, including number of layers in each

subnetwork and the final subnetwork, number of neurons in each layer, learning rate, batch size,

and dropout rate. We generated 100 combinations of hyper-parameters randomly selected from

the candidate parameter pool (Supplementary Data 21), using each combination to train a

DeepGCF model based on the same random subset of 1 million aligned and 1 million unaligned

human-pig pairs from the training set. We then selected the combination of hyper-parameters

that maximized the AUROC on the validation set to train the final model based on the whole

training set. We stopped training if there was no improvement in AUROC over three epochs on

the validation set for both hyper-parameter search and training, otherwise the training stopped

when reaching the maximum number of epochs, which was set to be 100.

Human-pig orthologous SNPs. In total 73,257,633 human biallelic SNPs (GRCh38) were

obtained from 1,000 Genome Project[31]. Their positions were lifted to corresponding orthologous

positions in the pig genomes (SusScr11) using the UCSC liftover utility with chain files available

from the UCSC website[45], which resulted in 35,575,835 orthologous SNPs.

Function enrichment. To explore the Gene Ontology terms of genomic regions (e.g.,

enhancers), we used the GREAT tool[48] with default parameters and a cut-off of FDR < 0.05 for

both the binomial and the hypergeometric distribution-based tests.

Tissue specific chromatin state. For each chromatin state, we first used the merge function of

BEDtools[49] to merge any regulatory regions between two tissues overlapping by at least 1 bp

across all tissues. Then for strongly active enhancer and promoter in each tissue, if a region is

active in only one tissue and does not overlap with any active regions in other tissues, we define

the region as tissue specific regulatory element. If a region is active in all tissues (i.e., overlaps

across all tissues), we define the region as "all common" regulatory element.

Tissue-sharing of e/sGene. To explore how e/sGenes are shared across all tissues, we performed

the meta-analysis of e/sGenes using MashR (v0.2.57)[50]. We used the slope and the standard error

of slope of top e/sQTL of genes (missing slopes were set to be 0 with standard error of 1) across

49 tissues from GTEx (v8)[23] for human and 34 tissues from PigGTEx databases[24] for pig as the

input. We then obtained the estimate of effect size and the corresponding significance (local false

sign rate, LFSR) from the mash function. An e/sGene was considered active in a tissue if LFSR

< 0.05.

DeepGCF score for genes. We obtained the gene boundaries of human and pig genes from

Ensembl release 107 (GRCh38 for human and Sscrofa11 for pig), and extended them by 35 kb

upstream and 10 kb downstream to include probable cis-regulatory regions[51]. We then compute

the DeepGCF score for genes based on the average score of all orthologous regions overlapping

with the gene and the extended regions. For human genes linked to promoter sequence class, we

identified a promoter's potential target gene if the distance between the promoter and the TSS of

a gene is less than 2 kb, yielding a total of 12,044 promoter-gene pairs.

Heritability partitioning analysis. We collected the GWAS summary statistics of 80 human

complex traits from the UK Biobank and public literatures (Supplementary Data 14). We ran the

LD-score regression software ldsc (v1.0.1)[41] to partition the heritability based on two sets of

annotations: 1) one binary annotation of functionally conserved regions (top 5% of DeepGCF)

and 2) five binary annotations dividing the top 5% DeepGCF into 5 equal-width bins based on

percentiles. Both sets of annotations were analyzed with a baseline including 97 annotations[40].

Heritability enrichment was calculated as the proportion of trait heritability contributed by SNPs

in the annotation over the proportion of SNPs in that annotation.

Fine-mapping analysis. We first used PolyFun[42] to compute SNP prior causal probabilities

based on the annotation of functional conservation (top 5% DeepGCF). These prior causal

probabilities were then used as priors in SuSiE[52] for the fine-mapping analysis. To compare fine-

mapping using functional conservation as prior with not using it, we also performed a fine-

mapping analysis using SuSiE alone, which only took LD information into account. A SNP is

identified to be putative causal if the posterior causal probability (PIP) is greater than 0.95 and

the *P*-value in GWAS is smaller than 5e-8.

Polygenic prediction. We incorporated functional conservation as a prior in polygenic

prediction using the software SBayesRC[43]. The GWAS summary statistics of 20 complex traits

from UK Biobank (Supplementary Data 17) were analyzed using ~7 million common SNPs with

and without one annotation of functional conservation (top 5% DeepGCF). To compare the

prediction accuracy, we partitioned the total sample into ten equal-sized disjoint subsamples. For

each fold, we retained one subsample as the validation set and other remaining nine subsamples

as the training set. We calculated the polygenic score (PGS) using genotypes from an

independent validation set in each fold and obtained the prediction R^2 from linear regression of

true phenotype on the PGS. We then calculated the relative prediction accuracy by $(R_0^2 - R_D^2)$ /

R_0^2, where R_0^2 is the prediction R^2 without any priors, and R_D^2 is the prediction R^2 using

functional conservation as a prior.

Data availability

The DeepGCF score for human-pig orthologous regions are publicly available for download

without restrictions from https://github.com/liangend/DeepGCF. All epigenomic and gene

expression data used for model training can be found in Supplementary data 1–6. Orthologous

SNPs between human and pig are from the 1,000 Genome Project

(http://ftp.1000genomes.ebi.ac.uk/vol1/ftp/data_collections/1000_genomes_project/release/2018

1203_biallelic_SNV). GWAS summary statistics used for LDSC analysis are from UK Biobank

(http://www.ukbiobank.ac.uk), with details showing in Supplementary data 14. Summary

statistics and genotype used for polygenic score prediction from UK Biobank

(http://www.ukbiobank.ac.uk) are available through formal application.

Code availability

The code of DeepGCF is available at https://github.com/liangend/DeepGCF.

References

1. Alföldi, J. & Lindblad-Toh, K. Comparative genomics as a tool to understand evolution and disease. *Genome Res.* **23**, 1063–1068 (2013).

2. Raymond, B. *et al.* Using prior information from humans to prioritize genes and gene-associated variants for complex traits in livestock. *PLOS Genetics* **16**, e1008780 (2020).

3. Lunney, J. K. *et al.* Importance of the pig as a human biomedical model. *Science Translational Medicine* **13**, eabd5758 (2021).

4. Schelstraete, W., Devreese, M. & Croubels, S. Comparative toxicokinetics of Fusarium mycotoxins in pigs and humans. *Food and Chemical Toxicology* **137**, 111140 (2020).

5. Montgomery, R. A. *et al.* Results of Two Cases of Pig-to-Human Kidney Xenotransplantation. *New England Journal of Medicine* **386**, 1889–1898 (2022).

6. Kragh, P. M. *et al.* Hemizygous minipigs produced by random gene insertion and handmade cloning express the Alzheimer's disease-causing dominant mutation APPsw. *Transgenic Res* **18**, 545–558 (2009).

7. Luo, Y. *et al.* High efficiency of BRCA1 knockout using rAAV-mediated gene targeting: developing a pig model for breast cancer. *Transgenic Res* **20**, 975–988 (2011).

8. Renner, S. *et al.* Permanent Neonatal Diabetes in INSC94Y Transgenic Pigs. *Diabetes* **62**, 1505–1511 (2013).

9. Cooper, G. M. *et al.* Distribution and intensity of constraint in mammalian genomic sequence. *Genome Res.* **15**, 901–913 (2005).

495   10. Pollard, K. S., Hubisz, M. J., Rosenbloom, K. R. & Siepel, A. Detection of nonneutral

496   substitution rates on mammalian phylogenies. *Genome Res.* **20**, 110–121 (2010).

497   11. Bordeira-Carriço, R. *et al.* Multidimensional chromatin profiling of zebrafish pancreas to

498   uncover and investigate disease-relevant enhancers. *Nat Commun* **13**, 1945 (2022).

499   12. Kunarso, G. *et al.* Transposable elements have rewired the core regulatory network of human

500   embryonic stem cells. *Nat Genet* **42**, 631–634 (2010).

501   13. Pennacchio, L. A. & Visel, A. Limits of sequence and functional conservation. *Nat Genet* **42**,

502   557–558 (2010).

503   14. ENCODE Project Consortium. The ENCODE (ENCyclopedia Of DNA Elements) Project.

504   *Science* **306**, 636–640 (2004).

505   15. Kundaje, A. *et al.* Integrative analysis of 111 reference human epigenomes. *Nature* **518**,

506   317–330 (2015).

507   16. Andersson, L. *et al.* Coordinated international action to accelerate genome-to-phenome with

508   FAANG, the Functional Annotation of Animal Genomes project. *Genome Biology* **16**, 57

509   (2015).

510   17. Liu, S. *et al.* A multi-tissue atlas of regulatory variants in cattle. *Nat Genet* 1–10 (2022)

511   doi:10.1038/s41588-022-01153-5.

512   18. Sjöstedt, E. *et al.* An atlas of the protein-coding genes in the human, pig, and mouse brain.

513   *Science* **367**, eaay5947 (2020).

514   19. Pan, Z. *et al.* Pig genome functional annotation enhances the biological interpretation of

515   complex traits and human disease. *Nat Commun* **12**, 5848 (2021).

516   20. Zhao, Y. *et al.* A compendium and comparative epigenomics analysis of cis-regulatory

517   elements in the pig genome. *Nat Commun* **12**, 2217 (2021).

518 21. Wong, A. K., Sealfon, R. S. G., Theesfeld, C. L. & Troyanskaya, O. G. Decoding disease:

519   from genomes to networks to phenotypes. *Nat Rev Genet* **22**, 774–790 (2021).

520 22. Kwon, S. B. & Ernst, J. Learning a genome-wide score of human–mouse conservation at the

521   functional genomics level. *Nat Commun* **12**, 2495 (2021).

522 23. THE GTEX CONSORTIUM *et al.* The Genotype-Tissue Expression (GTEx) pilot analysis:

523   Multitissue gene regulation in humans. *Science* **348**, 648–660 (2015).

524 24. Consortium, T. F.-P. *et al.* A compendium of genetic regulatory effects across pig tissues.

525   2022.11.11.516073 Preprint at https://doi.org/10.1101/2022.11.11.516073 (2022).

526 25. Zhou, J. & Troyanskaya, O. G. Predicting effects of noncoding variants with deep learning–

527   based sequence model. *Nat Methods* **12**, 931–934 (2015).

528 26. Hughes, L. H., Schmitt, M., Mou, L., Wang, Y. & Zhu, X. X. Identifying Corresponding

529   Patches in SAR and Optical Images With a Pseudo-Siamese CNN. *IEEE Geoscience and*

530   *Remote Sensing Letters* **15**, 784–788 (2018).

531 27. Xiao, S. *et al.* Comparative Epigenomic Annotation of Regulatory DNA. *Cell* **149**, 1381–

532   1392 (2012).

533 28. Chen, K. M., Wong, A. K., Troyanskaya, O. G. & Zhou, J. A sequence-based global map of

534   regulatory activity for deciphering human genetics. *Nat Genet* **54**, 940–949 (2022).

535 29. Liu, Y. *et al.* Comparative Gene Expression Signature of Pig, Human and Mouse Induced

536   Pluripotent Stem Cell Lines Reveals Insight into Pig Pluripotency Gene Networks. *Stem Cell*

537   *Rev and Rep* **10**, 162–176 (2014).

538 30. Beh, L. Y., Colwell, L. J. & Francis, N. J. A core subunit of Polycomb repressive complex 1

539   is broadly conserved in function but not primary sequence. *Proceedings of the National*

540   *Academy of Sciences* **109**, E1063–E1071 (2012).

31. Lowy-Gallego, E. *et al.* Variant calling on the GRCh38 assembly with the data from phase

three of the 1000 Genomes Project. *Wellcome Open Res* **4**, 50 (2019).

32. Flavahan, W. A. *et al.* Insulator dysfunction and oncogene activation in IDH mutant gliomas.

Nature **529**, 110–114 (2016).

33. Guo, Y. *et al.* CRISPR Inversion of CTCF Sites Alters Genome Topology and

Enhancer/Promoter Function. *Cell* **162**, 900–910 (2015).

34. Villar, D. *et al.* Enhancer Evolution across 20 Mammalian Species. *Cell* **160**, 554–566

(2015).

35. Yao, Y. *et al.* Comparative transcriptome in large-scale human and cattle populations.

Genome Biology **23**, 176 (2022).

36. Zhao, R. *et al.* The conservation of human functional variants and their effects across

livestock species. *Commun Biol* **5**, 1–13 (2022).

37. Mohammadi, P., Castel, S. E., Brown, A. A. & Lappalainen, T. Quantifying the regulatory

effect size of cis-acting genetic variation using allelic fold change. *Genome Res.* **27**, 1872–

1884 (2017).

38. Landrum, M. J. *et al.* ClinVar: public archive of relationships among sequence variation and

human phenotype. *Nucleic Acids Research* **42**, D980–D985 (2014).

39. Powell, G. *et al.* Modelling the genetic aetiology of complex disease: human–mouse

conservation of noncoding features and disease-associated loci. *Biology Letters* **18**,

20210630.

40. Hujoel, M. L. A., Gazal, S., Hormozdiari, F., van de Geijn, B. & Price, A. L. Disease

Heritability Enrichment of Regulatory Elements Is Concentrated in Elements with Ancient

Sequence Age and Conserved Function across Species. *The American Journal of Human*

Genetics **104**, 611–624 (2019).

41. Finucane, H. K. *et al.* Partitioning heritability by functional annotation using genome-wide

association summary statistics. *Nat Genet* **47**, 1228–1235 (2015).

42. Weissbrod, O. *et al.* Functionally informed fine-mapping and polygenic localization of

complex trait heritability. *Nat Genet* **52**, 1355–1363 (2020).

43. Zheng, Z. *et al.* Leveraging functional genomic annotations and genome coverage to improve

polygenic prediction of complex traits within and between ancestries. 2022.10.12.510418

Preprint at https://doi.org/10.1101/2022.10.12.510418 (2022).

44. Genereux, D. P. *et al.* A comparative genomics multitool for scientific discovery and

conservation. *Nature* **587**, 240–245 (2020).

45. Lee, B. T. *et al.* The UCSC Genome Browser database: 2022 update. *Nucleic Acids Research*

50, D1115–D1122 (2022).

46. Schwartz, S. *et al.* Human–Mouse Alignments with BLASTZ. *Genome Res.* **13**, 103–107

(2003).

47. Chen, K. M., Cofer, E. M., Zhou, J. & Troyanskaya, O. G. Selene: a PyTorch-based deep

learning library for sequence data. *Nat Methods* **16**, 315–318 (2019).

48. McLean, C. Y. *et al.* GREAT improves functional interpretation of cis-regulatory regions.

Nat Biotechnol **28**, 495–501 (2010).

49. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics* **26**, 841–842 (2010).

50. Urbut, S. M., Wang, G., Carbonetto, P. & Stephens, M. Flexible statistical methods for
estimating and testing effects in genomic studies with multiple conditions. *Nat Genet* **51**,
187–195 (2019).

51. Trubetskoy, V. *et al.* Mapping genomic loci implicates genes and synaptic biology in
schizophrenia. *Nature* **604**, 502–508 (2022).

52. Wang, G., Sarkar, A., Carbonetto, P. & Stephens, M. A simple new approach to variable
selection in regression, with application to genetic fine mapping. *Journal of the Royal*
Statistical Society: Series B (Statistical Methodology) **82**, 1273–1300 (2020).

 Figures and legends

a

b

Fig. 1 Overview of the DeepGCF model. a The learning procedure of DeepGCF model consists

of two steps. The first step is to train DeepSEA models in human and pig separately to transform

the binary functional features (e.g., peaks called from ATAC-seq and ChIP-seq, and chromatin

states predicted from ChromHMM) to continuous values by predicting the functional effects of

single nucleotides through centering the target nucleotide at a genomic region of 1,000 bp. The

second step is to train a pseudo-Siamese network for predicting whether the paired human-pig

regions are orthologous or not using two corresponding vectors of functional effects predicted

from DeepSEA and normalized gene expression as inputs. The output, DeepGCF score, is a

value between 0 and 1 quantifying the functional conservation of the paired human-pig region. **b**

The DeepGCF model can be applied to predict the effect of genome variants on the functional

conservation, quantified by changes in DeepGCF scores.

Fig. 2 The performance of DeepGCF under different scenarios a Receiver operating

characteristic (ROC) curves comparing the performance of DeepGCF (this study) and LECIF[22]

methods. The ROC curve of each method is generated by predicting whether 200,000 pairs

randomly selected from the testing set, which included equal number of orthologous and non-

orthologous pairs (e.g., randomly mismatched genomics regions), were orthologous or not. **b**

Precision-recall (PR) curves generated by similar procedures as the ROC curves. **c** DeepGCF

score distribution of all 38,961,848 human-pig orthologues pairs. **d** The areas under receiver

operating characteristic curve (AUROC) and precision-recall curve (AUPRC) of DeepGCF using

all (Human: 386; Pig: 374), ~50% (Human: 192; Pig: 187), ~10% (Human: 52; Pig: 47), and

~1% (Human: 4; Pig: 4) of human and pig functional features. The subsets of the human and pig

features were randomly selected ~50%, ~10%, ~1% from each of ChIP-/ ATAC-seq,

ChromHMM, and RNAseq profiles. **e** The AUROC and AUPRC of DeepGCF using all

620   functional features (Human: 386; Pig: 374), features without ChIP-/ATAC-seq (Human: 129;

621   Pig: 84), without ChromHMM (Human: 180; Pig: 210) and without RNA-seq (Human: 77; Pig:

622   80).

Fig. 3 Comparison of functional and sequence conservations. a Relationship between

DeepGCF score and PhyloP score of 20,000 randomly selected human regions. PhyloP score is

based on multiple alignments of 99 vertebrate genomes to the human genome[10]. The blue line is

the fitted loess regression and red crosses represents 50 equally-divided percentiles of PhyloP

score corresponding to the average of DeepGCF score. **b** Enrichment fold of 8 sequence class

categories[28] for regions with high DeepGCF (> 95th percentile) and high PhyloP (> 95th

percentile; high D & high P; n = 260,281), regions with high DeepGCF but low PhyloP (< 5th

percentile; high D & low P; n = 152,557), and regions with low DeepGCF (< 5th percentile) and

medium PhyloP (between 47.5th and 52.5th percentile; low D & med P; n = 77,848). Enrichment

is equal to the proportion of a sequence class category for a type of orthologous regions divided

by that for the whole genome. The dashed line (set at 1) represents no enrichment. **c** DeepGCF

score distribution of the different sequence ontologies. The red and green dashed lines represent

the mean and the median DeepGCF score of the whole genome. The dots inside each box

represent the mean DeepGCF score. **d** ΔDeepGCF (DeepGCF after mutation – original DeepGCF)

caused by 1,000,000 randomly selected orthologous variants, which are classified into 8 sequence class

categories[28]. The red dashed line represents the fitted regression line. **e** The effect of orthologous

variants (n = 35,575,835) on DeepGCF score of regions in 40 sequence classes[28], which are

classified into 8 categories. The effect was measured by ΔDeepGCF for variants in each

sequence class. The SD of ΔDeepGCF for each sequence class quantifies the overall sensitivity

of the sequence class to variant effect.

Fig. 4 DeepGCF score of genomic regions overlapping with different regulatory elements. a
Distribution of average DeepGCF scores across human tissues ($n = 12$) and pig tissues ($n = 14$)
for each chromatin state. The red and green dashed lines represent the mean and the median
DeepGCF score of the whole genome. **b** DeepGCF scores of genomic regions overlapping with
tissue-specific strongly active promoter and enhancer for human and pig[19]. "All common"
represents promoters/enhancers shared across all tissues. **** denotes Mann–Whitney U test $P <$
2.2e-16. **c** Number of significantly enriched gene ontology terms for human of genes related to
promoters annotated by sequence class[28]. The genes were binned by DeepGCF into ten equal-
width bins, and the functional enrichment analysis was conducted on each bin. **d** Similar to **c**,
except showing the results of enhancers annotated by sequence class[28].

Fig. 5 Relationship of DeepGCF score to genetic variants. a The distribution of DeepGCF

score of eQTLs and sQTLs. The red and green dashed lines represent the mean and the median

DeepGCF score of the whole genome. The dots inside each box represent the mean DeepGCF

score. **** denotes P value < 1e-8 based on a two-sided Mann–Whitney U test. **b** Relationship

between the absolute value of eQTL effect size ($|\log_2(aFC)|$) and DeepGCF score for eGenes.

The genes were binned by DeepGCF into ten equal-width bins for human and pig, respectively.

**** denotes the group is different from all other groups with P value < 1e-8 based on a Tukey

multiple comparison. **c** DeepGCF scores of tissue-sharing e/sGenes from human at local false

sign rate (LFSR) < 5% obtained by MashR[50]. **d** Similar to **c**, except showing the results of pig.

Fig. 6 Relationship of conservation score to pathogenic variants. a The distribution of

DeepGCF scores in pathogenic and likely pathogenic SNPs (n = 104,033) obtained from

ClinVar[38], compared to the DeepGCF distribution across the whole genome. **** denotes Mann–

Whitney U test P < 2.2e-16. **b** SD of ΔDeepGCF (DeepGCF after mutation – original DeepGCF)

caused by ClinVar SNPs. The SNPs were binned by their original DeepGCF into ten equal-width

bins. **c** ClinVar SNPs classified by sequence class[28]. A polar coordinate system was used, where

the radial coordinate indicates the SNP effect on DeepGCF. The red solid circle represents zero

DeepGCF change, and two dashed circles represent ± 0.03 of DeepGCF encompassing 95% of

SNPs. Each dot represents a SNP and SNPs inside the red circle were predicted to have positive

effects (increased DeepGCF), while SNPs outside the red circle were predicted to have negative

effects (decreased DeepGCF). Dot size indicates the original DeepGCF. Within each sequence

class, SNPs were ordered by chromosomal coordinates. Top 10 SNPs with large impact on

DeepGCF associated disease and gene names were annotated.

682
Fig. 7 Application of DeepGCF on complex traits/diseases in human. a Heritability

enrichment calculated by LDSC for 80 human traits using functionally conserved regions (top

5% DeepGCF). The regions were divided into 5 equal equal-width bins and the heritability

enrichment of all traits was calculated for each bin. The dashed red line is the fitted regression

line between heritability enrichment and DeepGCF percentile, and the grey area is the 95%

confidence interval. **b** Significant heritability enrichment explained by functionally conserved

regions in 8 human traits. **c** The number of putatitive SNPs (PIP > 0.95 and P < 5e-8) identified

by PolyFun + SuSiE using functionally conserved regions as a prior and SuSiE without priors for

7 human traits. **d** The relative prediction accuracy of PRS for 20 human complex traits using

functionally conserved regions as a prior in SBayesRC[43]. Relative prediction accuracy is equal to

(prediction accuracy using the prior – prediction accuracy without priors) / prediction accuracy

without priors. A relative prediction accuracy > 0 (dashed line) indicates an accuracy higher than

without priors.