The apicoplast is important for the viability and persistence of
Toxoplasma gondii bradyzoites

Syrian G. Sanchez[1], Emilie Bassot[2], Aude Cerutti[1], Hoa Mai Nguyen[1], Amel Aïda[2], Nicolas
Blanchard[2], Sébastien Besteiro[1]*

[1] LPHI, CNRS, University of Montpellier, France

[2] Toulouse Institute for Infectious and Inflammatory Diseases, Infinity, University of Toulouse,
CNRS, Inserm, UPS, Toulouse, France

*Corresponding author: Sébastien Besteiro

Email: sebastien.besteiro@inserm.fr

Author Contributions: N. Blanchard and S. Besteiro designed research; S. Sanchez, E. Bassot,
A. Cerutti, H.M. Nguyen, A. Aïda performed research; S. Sanchez, E. Bassot, N. Blanchard and
S. Besteiro analyzed the data; S. Sanchez, N. Blanchard and S. Besteiro wrote the paper; N.
Blanchard and S. Besteiro funding acquisition.

Keywords: Toxoplasma gondii, chronic toxoplasmosis, plastid, stage differentiation

Abstract

Toxoplasma gondii is responsible for toxoplasmosis, a disease that can be serious when contracted during pregnancy, but can also be a threat for immunocompromised individuals. Acute infection is associated with the tachyzoite form that spreads rapidly within the host. However, under stress conditions, some parasites can differentiate into cyst-forming bradyzoites, residing mainly in the central nervous system, retina and muscle. Because this latent form of the parasite is resistant to all currently available treatments, and is central to persistence and transmission of the parasite, new specific therapeutic strategies targeting this developmental stage need to be discovered.

T. gondii contains a plastid of endosymbiotic origin called the apicoplast, which is an appealing drug target because it is essential for tachyzoite viability and contains several key metabolic pathways that are largely absent from the mammalian host. Its function in bradyzoites, however, is unknown. Our objective was thus to study the contribution of the apicoplast to the viability and persistence of bradyzoites during chronic toxoplasmosis.

We have used complementary strategies based on stage-specific promoters to generate conditional bradyzoite mutants of essential apicoplast genes. Our results show that specifically targeting the apicoplast in both *in vitro* or *in vivo*-differentiated bradyzoites, leads to a decrease in cyst burden or a loss of bradyzoite viability, highlighting the importance of this organelle for this developmental stage. This validates the apicoplast as a potential area to look for new therapeutic targets in bradyzoites, with the aim to interfere with this currently incurable parasite stage.

Significance Statement

In its intermediate hosts, the parasite *Toxoplasma gondii* can persist as a cyst-contained developmental form that might reactivate and cause severe pathologies. Importantly, this form is resistant to current anti-parasitic drugs. *T. gondii* harbors a plastid of endosymbiotic origin called the apicoplast, containing important and potentially druggable metabolic pathways, but whose contribution to the fitness and viability of persistent parasites has never been assessed. We have generated conditional mutants specifically affected for the homeostasis of the apicoplast in cyst-contained parasites and showed that this organelle is crucial for persistence of these particular

44 developmental forms. Our work thus validates the apicoplast as a relevant drug target in the context
45 of chronic *T. gondii* infection.
46
47
48
49 **Introduction**
50
Toxoplasma gondii is an intracellular parasitic protist responsible for toxoplasmosis, a ubiquitous
disease affecting humans, which can lead to severe forms during pregnancy or in
immunosuppressed individuals (1). Acute toxoplasmosis is associated with the tachyzoite form of
these parasites, which spreads rapidly within the host and has the ability to cross physiological
barriers, but is generally controlled by the immune system in immunocompetent hosts (2). The
genus *Toxoplasma* contains only one species, but there are many different strains which are
grouped according to their virulence and pathogenicity in mouse models (3). The pathogenic
outcome is closely linked to the ability of the parasite to convert from fast-replicating tachyzoites to
persisting encysted stages that can be found predominantly in neurons and muscle tissues and are
called bradyzoites (4). These developmental forms of the parasite, which are the hallmark of chronic
infection, are supposedly less active metabolically and slower to replicate than tachyzoites (5).
However, they play a crucial role in *T. gondii* transmission, as well as in immune escape and
survival of the parasites under stressful conditions (6).
The process of stage conversion is a continuum lasting for several days that leads to marked
differences between tachyzoites and bradyzoites. For instance the membrane of the
parasitophorous vacuole (PV, the compartment in which the parasite replicates) transforms into a
heavily-glycosylated cyst wall (7, 8). Also, several aspects of parasite metabolism change
drastically, as illustrated by the accumulation of amylopectin granules in bradyzoites (9) and the
expression of different stage-specific isoforms of metabolic enzymes (10–14). Thus unsurprisingly,
stage conversion is accompanied by a considerable change in protein expression driven by stage-
specific promoters. Transcriptomic analyses of *in vivo*-generated cysts have for instance revealed
that the expression levels of hundreds of genes are potentially changing (15, 16). The specific
metabolic requirements of bradyzoites are so far poorly characterized, largely due to the technical
challenges associated with studying this developmental stage: although *in vitro* differentiation can
be achieved by applying stresses to a cystogenic *T. gondii* strain or by infecting specific host cell
types, *in vitro*-derived bradyzoites are difficult to keep in long-term culture and they usually do not
reach the stage of complete maturity that is observed in mouse brain-derived bradyzoites (17–20).
As intracellular developmental stages, both PV-contained tachyzoites and cyst-enclosed
bradyzoites are tightly interconnected metabolically with their host cells to ensure optimal

proliferation or persistence (21). While *T. gondii* relies on specific host cell resources for survival, it
is also able to synthesize *de novo* a number of important metabolites. One key parasite organelle
for metabolite production is the apicoplast. This plastid of secondary endosymbiotic origin has lost
its photosynthetic capacity (22), but it nevertheless harbors four main metabolic pathways which
have all been shown to be essential for the *in vitro* fitness of *T. gondii* tachyzoites (23, 24). These
are important for the synthesis of fatty acids (FAs, via the FASII pathway), heme (together with the
mitochondrion), isoprenoid precursors and iron-sulfur clusters. Because of the prokaryotic origin
and the metabolic importance of most of the apicoplast-hosted pathways, this organelle is
particularly attractive as a potential drug target and compounds inhibiting translation in the
apicoplast, like clindamycin, are already used against acute toxoplasmosis (25).

However, to this day there is no drug available to clear the cyst form that persists during the chronic
phase of infection, and this has major implications for pathogenesis. Clearly, curing patients from
the infection is of paramount importance to prevent parasite reactivation. Several compounds,
including atovaquone and quinolone derivatives, or even combinatorial strategies involving
apicoplast targeting drugs like clindamycin and spiramycin, are able to limit cyst burden in animal
models of chronic toxoplasmosis (26–32). However, most fail to completely eradicate bradyzoites,
and some are even known to be potent inducers of stage conversion in vitro (33, 34). Yet, it is still
unclear whether the resistance to these drugs is due to poor accessibility of brain-localized cysts
or to the metabolically quiescent nature of the bradyzoites.

The metabolic contribution of the apicoplast to the viability and persistence of these developmental
stages is still a major conundrum. To get insights into the importance of the organelle in the context
of chronic *T. gondii* infection, we have generated conditional mutants specifically affected in
apicoplast homeostasis at the bradyzoite stage. As the apicoplast is essential to the viability of
tachyzoites, which are a necessary step prior to obtaining bradyzoites, we needed to adopt a
conditional strategy for targeting the organelle. Most conditional expression systems available for
T. gondii involve the use of small molecules (35), however the bioavailability of these compounds
may be difficult to regulate in the *in vivo* mouse model and most of them may not efficiently cross
the blood-brain barrier to reach the tissue cysts. We thus genetically-engineered parasite strains to
impact apicoplast functions specifically in bradyzoites and without the use of external compounds,
by using stage-specific promoters. We generated mutants either affected for apicoplast-related
metabolism or in the actual structure and division process of the organelle. Through *in vitro* and *in*
vivo phenotypic analyses, we showed that in both cases the viability or development of bradyzoites
was impaired. Our study thus validates the apicoplast as a potential drug target in the context of
chronic toxoplasmosis.

Results

Specific down-regulation of the APT1 apicoplast phosphate translocator at the bradyzoite stage. To generate a parasite line impacted for apicoplast homeostasis, we first targeted the Apicoplast triosephosphate translocator 1 (APT1). APT1 is involved in the import of phosphoenol pyruvate and triose phosphates, which are important precursors for the FASII system and for isoprenoid synthesis in the organelle (Fig. 1A), and as such it is essential to the survival of *T. gondii* tachyzoites (36). Although the metabolic requirements of the bradyzoites are not completely elucidated yet, we reasoned that depleting a transporter which is needed for two of the most important apicoplast-hosted metabolic pathways would be likely to have an impact. To generate a conditional mutant specifically affected in bradyzoites, we used CRISPR/Cas9 to edit the endogenous *APT1* locus in the cystogenic Prugniaud (Pru) strain: we inserted a cassette coding for a GFP-fused version of the APT1 protein driven by a tachyzoite stage-specific *SAG1* promoter (Fig. 1B). Proper integration in the resulting cell line, named pSAG1-APT1, was verified by PCR (Fig. S1A). Using co-staining with TgAtrx1, a marker of the periphery of the apicoplast (37), we also checked that the promoter change and the addition of the GFP tag were not affecting the localization of APT1 at the organelle (Fig. S1B). However, it should be noted that the *SAG1* promoter is potentially a much stronger promoter than the native *APT1* promoter according to comparative analysis of RNA expression level (6). Accordingly, we performed semi-quantitative RT-PCR analyses on tachyzoites that confirmed at least a 3-fold increase in expression for the *APT1* mRNA upon promoter change (Fig. S1C, D).

Conversion of tachyzoites into bradyzoites can be initiated *in vitro* by applying a variety of stresses like alkaline pH, heat shock, nutrient starvation or the use of specific drugs (4). For instance, alkaline pH stress-induced differentiation, which is performed in absence of CO_2, is commonly used to induce conversion to bradyzoites with minimal toxicity to host cells. We thus used this stress on the pSAG1-APT1 cell line and performed immunoblot analysis that showed a decrease in expression of the *SAG1* promoter-driven APT1 protein in *in vitro*-generated bradyzoites (Fig. 1C). We performed fluorescence microscopy on differentiating parasites using a lectin from the plant *Dolichos biflorus* (DBL) that recognizes the SRS44/CST1 cyst wall glycoprotein that accumulates during differentiation (38) and, strikingly, pSAG1-APT1 was found to be largely absent from more

146   mature cysts (Fig. 1D). This loss was quantified and the proportion of cysts displaying an intact
147   pSAG1-APT1 signal decreased over time during pH stress-induced differentiation (Fig. 1E).
148   Overall, our data show that the *SAG1* promoter replacement strategy leads to an efficient down-
149   regulation of APT1 upon *in vitro* differentiation of bradyzoites.
150
Down-regulation of APT1 leads to perturbation of apicoplast homeostasis and impacts the
development and viability of *in vitro*-generated bradyzoites. We next evaluated the impact of
APT1 down-regulation on the overall homeostasis of the apicoplast. Depleting APT1 in tachyzoites
has been shown not to have an immediate impact on the morphology of the organelle (36).
However, this translocator is important for isoprenoid production, whose perturbation leads to loss
of the organelle in the related malaria-causing *Plasmodium* parasites (39, 40), so we anticipated
that long term depletion could also have an effect in bradyzoites. Co-staining of APT1 with TgCpn60
(a marker of the apicoplast lumen (41)) during the differentiation process showed a usually
concomitant disappearance of both proteins in more mature cysts (Fig. 2A). Quantification of
apicoplast loss showed that only about 35% of the cysts still contained an intact apicoplast signal
after two weeks of pH stress-induced differentiation (Fig. 2B).
To assess the impact of APT1 down-regulation on *in vitro* cyst development, we imaged cysts after
two weeks of differentiation and then measured cyst area (Fig. 2C). We found the mean area to be
significantly lower in the pSAG1-APT1 cell line compared with the parental Pru cell line. This
suggests that APT1 depletion upon *in vitro* differentiation has a negative impact on cyst
development. We next sought to assess the impact of APT1 depletion on bradyzoite viability using
plaque assay as a proxy, as described previously (42). Cysts were generated *in vitro*, bradyzoites
were released by pepsin treatment and APT1-negative parasites were isolated with a cell sorter
based on the absence of a GFP signal, and seeded on host cell monolayers (Fig. S2). APT1-
negative bradyzoites were found to generate much less and much smaller plaques than the Pru
parental cell line (Fig. 2D, E, F). Of note, by *in vitro* plaque assay tachyzoites of the pSAG1-APT1
cell line did not show any particular fitness defect compared with the Pru parental cell line (Fig. 2D,
S3A, B).
In conclusion, our results indicate that specific depletion of APT1 in *in vitro*-generated bradyzoites
leads to visible consequences on the homeostasis of the apicoplast and on bradyzoite viability.
176
pSAG1-APT1 parasites induce a severe acute phase in mice, yet *ex vivo*-generated
bradyzoites are impaired in fitness. While *in vitro*-derived cysts are certainly a useful model,
some important differences have been observed with *in vivo*-derived cysts (20), thus we sought to
confirm our findings with additional assays in an animal model. To this end we infected CBA mice
with the pSAG1-APT1 and Pru control cell lines. In three independent experiments we observed

an excessive mortality and a more pronounced body mass loss in the surviving animals for mice
infected with pSAG1-APT1 tachyzoites (Fig. 3A, B), hinting that infection with this mutant leads to
a severe acute phase. As dendritic cells act early both as immune response mediators and as
parasite carriers that facilitate the dissemination of the infection and may play a key role in the
acute phase (43), we wanted to evaluate the ability of pSAG1-APT1 tachyzoites to infect these
cells *in vitro* using a cytometry based assay (Fig. S3C, D) (44). Yet, our results showed that pSAG1-
APT1 tachyzoites do not infect dendritic cells more efficiently than the Pru cell line (Fig. S3E).
Tissue cysts were recovered from the brains of infected mice at 40 days post-infection and
quantified by microscopic observation after DBL staining and parasite load was also quantified by
quantitative PCR (qPCR, Fig. 3C, D). Mice infected with the pSAG1-APT1 cell line had a much
higher cyst burden than the Pru parental cell line, which could either be related to the more severe
acute phase observed with these parasites, or illustrate more frequent parasite reactivation. Tissue
cysts were imaged to detect the presence of an apicoplast, and a majority of pSAG1-APT1 cysts
harbored bradyzoites missing the organelle (Fig. 3E, F). We also found that the size of pSAG1-
APT1 cysts was smaller than the size of cysts from the Pru cell line (Fig. 3G). Finally, to assess the
viability of these bradyzoites, we performed plaque assays with parasites recovered from these *ex*
vivo cysts and observed much less and much smaller plaques with pSAG1-APT1 parasites than
with the Pru parental cell line (Fig. 3H, I, J).
In conclusion, while in an *in vivo* setting, pSAG1-APT1 tachyzoites induce a more severe acute
phase and lead to a higher parasitemia in the brain, the phenotype of brain-derived pSAG1-APT1
parasites regarding cyst size, loss of the apicoplast or overall viability was very similar to the results
we obtained with *in vitro*-generated parasites. Our data thus indicate that the specific loss of APT1
function in *in vitro*- or *in vivo*-derived bradyzoites affects apicoplast homeostasis, bradyzoite
development and viability. This suggests that the apicoplast plays an important role for the fitness
of bradyzoites.
207
Generating a structural mutant of the apicoplast through a dominant-negative approach.
Although the exact nature of the metabolic requirement of bradyzoites is unknown, APT1 likely has
an important role in the metabolic contribution of the apicoplast for the development of these
parasite stages. Yet, as the inhibition of metabolic versus housekeeping functions of the organelle
can lead to different phenotypic outcomes (45), we sought to generate a second apicoplast mutant
that would be affected in its ability to divide rather than to perform a particular metabolic function.
To this end we targeted DrpA, a dynamin-related protein that was previously shown to be essential
for apicoplast fission and inheritance in dividing tachyzoites (Fig. 4A) (46). A single mutation in the

sequence of the DrpA protein (changing a lysine to an alanine in the GTP-binding site) has been
shown to disrupt dynamin function in a dominant-negative fashion in *T. gondii* tachyzoites (46).
We thus designed a strategy to specifically express the mutant version of DrpA (DrpA^{K42A}) in
bradyzoites, by using CRISPR/Cas9 to insert the sequence coding for DrpA^{K42A} at the *uracil*
phosphoribosyl transferase (*UPRT*) locus (Fig. 4B), which is dispensable for bradyzoites in normal
culture conditions and for cystogenesis *in vivo* (47), but allows selection of transgenic parasites
with fluorodeoxyribose (48). To drive the expression of the dominant-negative version of DrpA
selectively in bradyzoites, we chose the bradyzoite-specific promoter of P21, a late marker of
bradyzoite differentiation (49) that is suitable for this type of strategy (50). Our construct also
allowed the addition of a TY epitope tag (51) for subsequent immunodetection of the mutated DrpA
version. Transgenic parasites, named pP21-DrpA^{K42A}, were verified by PCR (Fig. S4) and used for
in vitro bradyzoite conversion. The UPRT is involved in pyrimidine salvage, which can be
compensated by *de novo* synthesis in the parasite (52), but this synthesis relies on CO_2, which has
to be omitted from the classical alkaline pH stress differentiation protocol. We thus used treatment
with the histone deacetylase inhibitor apicidin (53, 54) as an alternative for initiating *in vitro*
bradyzoite conversion of the pP21-DrpA^{K42A} cell line. Immunoblot analysis showed that the TY-
tagged DrpA^{K42A} protein is specifically expressed in *in vitro*-differentiated bradyzoites (Fig. 4C). This
was confirmed by immunofluorescence microscopy (Fig. 5A), showing that no DrpA^{K42A} protein was
detected in tachyzoites, while it was expressed in differentiated parasites. There, it localized to the
cytoplasm and to discrete sub-compartments of the apicoplast as expected (Fig. 5A) (46).

Dominant-negative pP21-DrpA^{K42A} parasites have their apicoplast specifically affected at the
bradyzoite stage but show no fitness defect as tachyzoites. Upon differentiation using apicidin,
detailed analysis of DrpA^{K42A}-expressing bradyzoites revealed that they display a reticulated
apicoplast labelling or absence of the organelle (Fig. 5A, B, S5), which is similar to what was
previously described upon DrpA^{K42A} expression in tachyzoites (46). Apicidin triggered stage
conversion very efficiently, but long-term incubation may have unspecific effects, so we also used
alternative methods of differentiation to drive DrpA^{K42A} expression and confirmed apicoplast
segregation and reticulation problems within cysts upon short term alkaline stress (as mentioned
above long term treatment is detrimental in absence of the *UPRT* locus) and heat stress-induced
differentiation (55) (Fig. S5A). Strikingly, when using an anti-P21 antibody on differentiating pP21-
DrpA^{K42A} parasites, we could see that cysts/vacuoles in which the morphological impact on the
apicoplast was the most visible were those with a higher expression of P21, confirming that the
dominant-negative effect is efficiently regulated by the bradyzoite-specific promoter (Fig. S5B).
However, when cyst size was measured after apicidin-induced *in vitro* differentiation, in contrast to
the results obtained with pSAG1-APT1 parasites, no significant difference was found in comparison

with the control (Fig. 5C). This highlights potential differences on the timing of phenotype initiation
or on the consequences at the cellular level when affecting the metabolism versus the ultrastructure
of the apicoplast. This also called for long term assessment of viability the pP21-DrpA^{K42A} parasites
in the mouse model. Prior to implementing these *in vivo* experiments, we assessed the *in vitro*
fitness of the tachyzoite stage of the pP21-DrpA^{K42A} cell line and found no particular problem in
their ability to form plaques on fibroblast monolayers (Fig. 5D, E). Like described earlier we also
checked by flow cytometry analysis that the pP21-DrpA^{K42A} tachyzoites were capable of infecting
dendritic cells (44).Our results showed that pP21-DrpA^{K42A} tachyzoites infect dendritic cells as
efficiently as the Pru cell line (Fig. 5F).
pP21-DrpA^{K42A} tachyzoites, when injected into mice, induced mass loss in the animals in the
timeframe corresponding to the acute phase of infection (i.e. up to 17 days, Fig. 5G), hinting that
this part of the infection process was occurring normally. However, in contrast to the pSAG1-APT1
cell line, infection with pP21-DrpA^{K42A} tachyzoites did not induce any specific mortality in mice
during the acute phase (Fig. S6A). Parasite quantification by qPCR showed the presence of
parasites in the brain as early as 12 days post-infection, confirming that pP21-DrpA^{K42A} mutant
parasites are able to disseminate and invade the brain. Yet, very little pP21-DrpA^{K42A} parasites
were detected 40 days post-infection (Fig. 5H). Importantly, there was no difference in the outcome
at 40 days when we increased the inoculum from 100 to 2000 tachyzoites (Fig. 5H). In fact, 40 days
post-infection, brain samples from mice infected with the pP21-DrpA^{K42A} cell line contained either
very few to no detectable cysts, as quantified by qPCR or estimated by microscopic counts (Fig.
5H, Fig. S6B). The fitness or invasive capacity of pP21-DrpA^{K42A} tachyzoites is not apparently
affected *in vitro* (Fig. 5D, E, F) and our *in vivo* data shows that they likely generate a normal acute
phase and are able to reach the brain (Fig. 5G, H), however long-term persistence is affected as
illustrated by the almost complete absence of detectable parasites 40 days post infection (Fig. 6B,
C, D). This suggests that long term perturbation of apicoplast division impacts bradyzoite viability
and persistence *in vivo*.
278

Discussion

The long-term persistence of encysted *T. gondii* parasites in a large proportion of the human
population, particularly in ocular tissue and in the central nervous system, has potentially important
consequences on the development of a future pathology. As neither the immune system nor current
drug-based approaches seem able to eliminate this developmental stage (31, 56), it is essential to
look for novel therapeutic approaches. Because of its evolutionary history and its metabolic
importance, the apicoplast has long been considered a promising drug target in the tachyzoite
stage, in which both pharmacological and genetic studies have shown that loss of the organelle,
loss of its genome, or loss of its metabolic function result in the death of the parasites. Here, for the

first time, we used two independent and complementary genetic approaches to investigate the
importance of the apicoplast in the encysted bradyzoite stage.

While there has been recent progress in the *in vitro* methods used to initiate stage conversion into
bradyzoites (18–20), the complexity and diversity of bradyzoites may not always be fully
recapitulated *in vitro*, so the most canonical way to experimentally generate authentic mature *T.*
gondii tissue cysts is currently through murine infections. Regulating gene expression in a cyst-
enclosed stage that is mostly confined to neuronal tissues (whose access is restricted by the blood-
brain barrier), is a technical challenge as it may not be reached efficiently by compounds commonly
used for driving conditional silencing. For this reason, we chose a stage-specific promoter-driven
strategy to conditionally control gene expression. Using the *SAG1* promoter to achieve selective
loss of expression of a gene of interest in bradyzoites is a strategy that we have already used
successfully in the past (42, 57), and it has been also recently implemented by others (58). We also
took advantage of stage-specific expression in a complementary approach, this time with the
promoter of the late bradyzoite marker P21 (50), to specifically switch on the expression of the
dominant-negative mutant version of the DrpA protein. Our data show that the regulation of
expression was efficient, both for *in vitro* and *in vivo* bradyzoite conversion models. However,
quantifications of the effect on the apicoplast morphology or the loss of the organelle, while
highlighting a strong impact on the organelle, also showed some phenotypic heterogeneity (for
instance, apicoplasts were still present in some cyst-enclosed pSAG1-APT1 bradyzoites recovered
40 days post infection -Fig. 3E). This is not surprising because stage conversion is a continuum
that may happen asynchronously. As a result, there can be some degree of heterogeneity in gene
expression, as single-cell RNAseq analyses of *in vitro*-differentiated parasites suggest (59).
Analyses on brain-derived cysts also highlighted a fairly large diversity between cysts and among
bradyzoites within the same cyst (5, 60). Our work nevertheless shows that stage-specific
promoters can be used efficiently to investigate gene function in bradyzoites through a simple
promoter replacement.

We purposely used different approaches and different types of mutants affected in apicoplast
homeostasis, with one primarily impacted for metabolism and the other one in organelle fission and
inheritance: we wanted to see if there was converging evidence on the essentiality of the organelle
in bradyzoites. Interestingly, the two cell lines we generated through these different approaches
were found to display phenotypes that are not completely overlapping (Fig. S7). One particularly
striking difference is the strong acute phase induced by the pSAG1-APT1 cell line (Fig. 3A, B, C
and D). One possible explanation is that promoter replacement likely led to an overexpression of
APT1 (Fig. S1C and D), and this might in turn have conferred a metabolic and thus a fitness
advantage to tachyzoites in the host. On the other hand, our plaque assays show that there does
not seem to be an increased fitness for tachyzoites *in vitro* (Fig. 2D, S3). Discrepancies between

in vitro and *in vivo* settings have been highlighted in several studies describing the metabolic
flexibility of *T. gondii* tachyzoites, which are able to scavenge a number of nutrients in rich culture
medium but may encounter more restrictive conditions in the animal host (61). A number of host
and parasite factors account for the heterogeneity in the ranges of mice brain cyst burden described
in the literature, and while it is difficult to establish an exact correlation between the cyst burden
and the initial inoculum, it is quite likely that a more successful acute phase can lead to increased
parasite load in the brain (62, 63). It is thus possible that a fitness advantage during the acute
phase would be the reason of the high cyst numbers generated by the pSAG1-APT1 cell line. These
large cyst numbers may also potentially reflect multiple reactivation and reinvasion events, which
might also explain the fact that these cysts were generally found to be smaller than those of the
parental cell line. However, the majority of pSAG1-APT1 cysts that were recovered showed
morphological alterations, especially regarding the apicoplast (Fig. 3E and D). Moreover,
bradyzoites that were released from these cysts were found to be largely non-viable (Fig. 3H, I and
J). So, altogether our data rather suggest that the small cysts reflect a problem in parasite
development instead of recent parasite establishment.
In contrast, the pP21-DrpA^{K42A} cell line largely failed to generate cysts (Fig. 5H). We could establish
that it was not due to an alteration in the acute phase, as the parasites were found to reach the
brain at the end of this phase of infection (Fig. 5G and H). Instead, this probably rather highlights
an increased elimination by the immune system, or/and a strong defect in cystogenesis. While both
our strategies were aimed at interfering with the function of the apicoplast, the different outcomes
in the animal model concerning cyst generation may be due to the different promoters that were
used, which respectively turn genes on or off at different times during the stage conversion process,
thus affecting the timeframe of the manifestation of the detrimental phenotypes.
Another main difference in the mutants we generated is that while for both there was a visible
morphological impact *in vitro*, the pSAG1-APT1 mutant cell line was strongly affected in cyst growth
early during *in vitro* differentiation (Fig. 3C), while for pP21-DrpA^{K42A} bradyzoites the detrimental
effect on cystogenesis only manifested during long term infection in the animal (Fig. 5C and H).
This could be related to the fact that the former targets metabolic pathways hosted by the organelle,
while the latter targets its division and inheritance in parasite progeny. It has been known for some
time that there are differences in the kinetics of parasite demise whether a metabolic pathway of
the apicoplast or a housekeeping function of the organelle are targeted (45). Targeting apicoplast-
hosted metabolism through genetic or pharmaceutical-based approaches usually lead to a rapid
death of the parasites, while affecting apicoplast division usually leads to a so-called delayed death,
where parasites are only affected after another round of reinvasion. While *in vitro*-differentiated
pP21-DrpA^{K42A} bradyzoites showed marked signs of morphological alteration of their apicoplasts
(Fig. 5A, Fig. S5), the remaining organelles potentially retained some metabolic capacity. A strong

impact on the apicoplast, which is linked to organelle inheritance defects in this mutant, likely
depend on the parasites undergoing several successive rounds of division that would only happen
over a longer period of time. It was shown that *T. gondii* tachyzoites can potentially share resources
(metabolites and proteins) through a connection at their basal pole, ensuring proper cellular division
in spite of an absence of apicoplast of several parasites within the PV, and thus explaining the
delayed death effect (64). However, bradyzoite division is largely asynchronous and there is no
connection between the parasites in mature cysts (9, 64, 65), which suggests that organelle loss
may be more detrimental for this developmental stage. Moreover, this might be particularly
important in an *in vivo* setting, where nutrient sources are potentially scarce, which could explain
why cystogenesis of pP21-DrpA^{K42A} parasites ended up being strongly affected in these conditions.
Importantly, in the mouse model, both pSAG1-APT1 parasites (that generate numerous but small
cysts, containing less fit bradyzoites) and pP21-DrpA^{K42A} parasites (that display a strong defect in
cystogenesis) showed converging evidence that the apicoplast is important for *T. gondii* survival
and persistence as bradyzoites. This now raises the question of which essential function(s) the
organelle is performing for this developmental stage. An earlier *in vivo* morphological study showed
that duplication or elongation of the apicoplast could still be observed in bradyzoites many weeks
or even months post-infection, suggesting an active role for the organelle (66). Due to the lack of
functional studies, the metabolic activities and requirements of bradyzoites are, however, still
largely uncharacterized. RNA expression data from *in vivo*-generated cysts suggest that the four
main metabolic pathways hosted by the apicoplast could be active in bradyzoites, with enzymes of
the iron-sulfur cluster and isoprenoid synthesis pathways even potentially upregulated (16). It would
now be particularly interesting to apply our conditional strategies to specifically interfere with
individual enzymes of these pathways in bradyzoites.
In summary, our work provides corroborating lines of evidence showing that the apicoplast is
important for bradyzoite fitness and persistence. Importantly, this validates the organelle as a
potential drug target in a developmental stage that is currently incurable. Of course, it should be
kept in mind that designing drugs able to interfere with the apicoplast in bradyzoites will constitute
a major challenge. For instance, any compound targeting a specific pathway hosted by the
organelle would have to be able to go through a series of physical hurdles that include the blood-
brain barrier, the host cell plasma membrane, the cyst wall, the parasite plasma membrane and,
finally, the four membranes of the organelle. Also, as bradyzoites are slow growing, they are likely
to be less sensitive to metabolic targeting, by analogy with bacterial or tumoral 'persister' cells that
are notoriously prone to resist to drugs (67). However, although cyst-enclosed bradyzoites display
some heterogeneity in their cellular and metabolic activities, they remain dynamic growing entities
(60), and our results suggest that they are likely to be vulnerable to long term interference with

apicoplast-related functions. So, the possibility to exploit this as a therapeutic strategy should not

be overlooked.

Materials and Methods

Animal care and ethics statement

Animal care and use protocols were carried out under the control of the National Veterinary

Services and in accordance with the European Union guidelines for the handling of laboratory

animals EEC Council Directive, 2010/63/EU, September 2010). The protocol inducing pain

(APAFIS#25130-2020040721346790 v3) was approved by the local Ethical Committee for Animal

Experimentation registered by the 'Comité National de Réflexion Ethique sur l'Expérimentation

Animale' under no. CEEA122. CBA/J mice mice were purchased from Janvier Labs (France).

Parasites and cells culture

Tachyzoites of the Prugniaud *T. gondii* strain (68), as well as derived transgenic parasites, were

maintained by serial passage in monolayers of human foreskin fibroblast (HFF, American Type

Culture Collection, CRL 1634) grown in Dulbecco's modified Eagle medium (DMEM, Gibco),

supplemented with 5% of decomplemented fetal bovine serum, 2 mM glutamine, and a cocktail of

penicillin-streptomycin at 100 µg/ml.

Generation of the TgAPT1 conditional knockdown pSAG1-APT1 cell line

To generate a conditional knock-down TgAPT1 line specifically depleted at the bradyzoite stage,

the endogenous promoter of the *TgAPT1* gene was replaced by a *SAG1* promoter (a tachyzoite-

specific promoter) using a CRISPR/Cas9-based strategy. The whole genomic sequence of TgAPT1

(www.toxodb.org accession number #TGME49_261070) was amplified by PCR with the Q5 High

Fidelity polymerase (New England Biolabs) from Pru genomic DNA, with primers ML3012/ML3013

(all the primers used in this study are listed in Table S1). The resulting 1059 bp fragment was then

inserted into the pLIC-CAT-GFP plasmid (69), in frame with the sequence coding for the GFP,

resulting in the pLIC-TgAPT1-GFP plasmid. The DHFR-SAG1 promoter cassette was amplified

from the DHFR-pSAG1-GFP-TgATG8 plasmid (42) using primers ML3015/ML3016, both

containing PacI restriction site. The DHFR-SAG1 promoter cassette was then digested and inserted

into PacI-digested pLIC-TgAPT1-GFP plasmid to create DHFR-pSAG1-TgAPT1-GFP plasmid.

Specific single-guide RNAs (sgRNAs) were generated to introduce a double-stranded break at the

5' and 3′ boundaries of the *TgAPT1* locus. Primers used to generate the guides were

ML3026/ML3027 and ML3030/ML3031, and the protospacer sequences were introduced in the

Cas9-expressing pU6-Universal plasmid (Addgene, ref #52694) (70). The cassette corresponding

the DHFR-pSAG1-TgAPT1-GFP cassette was amplified by PCR from the plasmid of the same

name using primers ML3088/ML3089 with the KOD DNA polymerase (Sigma-Aldrich). This donor

sequence was then used to transfected to Pru background, together with the two guide-expressing

plasmids. The Electro Cell Manipulator 630 (BTX) was used with the following settings: 2.02 kV,

50 Ω, 25µF. Transfected parasites were selected with 1µM pyrimethamine and cloned by limit

dilution. Positive parasites were first screened for GFP-positive apicoplasts by microscopy, then

positive clones were verified by PCR using primers ML3023/ML3116 (5' end of the wild-type locus),

ML3115/ML4541 (3' end of the wild-type locus), ML2873/ML3116 (5' end after integration) and

ML4525/ML4541 (3' end after integration integration). The resulting parasite line was designated

pSAG1-APT1. Several clones were isolated and tested for regulation of protein expression and

impact on the apicoplast during *in vitro* differentiation and showed no particular difference, so one

was subsequently chosen for complete phenotypic analysis.

Generation of the conditional dominant-negative pP21-DrpA^{K42A} cell line

The pP21-DrpA^{K42A} mutant cell line was generated based on the *pUPRT*-pTUB-*G13-TY* plasmid

(71) using a CRISPR-based strategy. We performed PCR with the Q5 High Fidelity polymerase

(New England Biolabs) to amplify a 1,500 bp region upstream of the start codon of the *TgP21* gene

(www.toxodb.org accession number #TGME49_238440) containing a promoter sequence, using

primers ML3655/ML3656 and cloned using NotI/BclI sites in the *pUPRT*-pTUB-*G13-TY* plasmid to

create the *pUPRT*-pP21-*G13-TY plasmid. To amplify the* DrpA *coding sequence (*www.toxodb.org

accession number #TGME49_267800)*, total cDNA of the Pru strain was generated following

purification of parasite total RNA using the NucleoSpin RNA kit (Macherey Nagel), followed by a

reverse transcription using the Super Script III first strand synthesis supermix kit (Invitrogen). This

was then used to amplify the *DrpA* cDNA by PCR with the Q5 polymerase (New England BioLabs)

and primers ML 4417/4418. *DrpA* cDNA was then subcloned in the PCR-Blunt II-TOPO vector

(Invitrogen). Site-directed mutagenesis was performed with the QuikChange II Site-directed

Mutagenesis kit (Agilent) and primers ML 4421/4422 to generate the TOPO-DrpA^{K42A} vector. The

fragment was then cloned into the BglII/AvrII digested *pUPRT*-pP21-*G13-TY* plasmid in order to

create the pUPRT-pP21-DrpA^{K42A}-Ty plasmid. This plasmid was then linearized with the AvrII

enzyme and dephosphorylated with Antartic Phosphatase (New England Biolabs) in order to insert

an extra double TY tag. The sequence coding for the tag was created by annealing primers

ML4419/4420 and cloned into the linearized and dephosphorylated pUPRT-pP21-DrpA^{K42A}-TY

vector, to yield the pUPRT-pP21-DrpA^{K42A}-3xTY plasmid.

This plasmid was then linearized by KpnI/NsiI prior to transfection. The recombination efficiency

was increased by co-transfecting with Cas9-expressing pU6-UPRT plasmids, generated by

integrating *UPRT*-specific protospacer sequences (with primers ML3445/ML3446 for the 5' and

ML2087/ML2088 for the 3') which were designed to allow a double-strand break at the *UPRT* locus.

Transgenic parasites were selected with 10 µM 5-fluorodeoxyuridine (Sigma-Aldrich) and then

cloned by serial limiting dilution in a p96 wells plate. The resulting cell line was designated pP21-

DrpA^{K42A}. Several clones were isolated and tested for regulation of protein expression and impact

on the apicoplast during *in vitro* differentiation and showed no particular difference, so one was

subsequently chosen for complete phenotypic analysis.

***In vitro* differentiation**

Several methods were used to generate bradyzoites *in vitro*. For alkaline pH-induced differentiation

(55) mostly used for conversion of the pSAG1-APT1 cell line, HFF monolayers of cells grown were

infected with freshly egressed tachyzoites for 24h in DMEM culture medium, either on T25 flasks

(with 500,000 parasites) or on coverslips in 24-well plates (with 50,000 parasites). The medium

was then replaced by Minimum Essential Medium (MEM; 10X) without NaHCO3 (Gibco),

supplemented with 50 mM HEPES, 1% penicillin-streptomycin, 1% Glutamine and 3% FBS and

adjusted to pH 8.2. Parasites were then cultured at 37°C without CO_2 the medium was changed

every two days during the whole duration of the differentiation experiment, which lasted up to two

weeks.

For the pP21-DrpA^{K42A} cell line that cannot sustain long-term CO_2 deprivation, differentiation was

induced by alkaline pH for only 3 days, or alternatively by treatment with the histone deacetylase

inhibitor apicidin (53) or by heat-stress (55). Briefly, for apicidin-induced differentiation tachyzoites

were used to infect host HFFs and after 24h they were treated with 40 nM apicidin (Santa Cruz) for

up to 6 days. It should be noted that longer incubations with apicidin led to some parasite mortality,

even in the control cell line. Alternatively, stage conversion in the pP21-DrpA^{K42A} cell line was

initiated by a temperature shift. Briefly, host cells were put at 43°C for two hours prior to infection

to acquire thermotolerance and returned at 37°C for an hour. Then tachyzoites were allowed to

invade for 2h before placing the culture at 43°C for two days and then returned at 37°C for up to

six days.

Cyst purification and bradyzoite isolation

Cysts were isolated by isopycnic centrifugation (72). Briefly, mouse brains were collected 40 days

post infection and homogenized in a Dounce tissue grinder in 3 ml of PBS. To separate parasite

cysts from brain cell debris, the homogenate was centrifuged and the pellet was resuspended in

2.5 ml of isotonic Percoll (90% Percoll, 10% NaCl 1.5 M), and PBS was added up to 7.5 ml. After

centrifugation at 500 g for 15 minutes at 4°C, cysts were recovered in the pellet for further analysis.

Bradyzoites were recovered from tissue cysts by pepsin treatment. Briefly, tissue cysts were
washed in Hanks' Balanced Salt Solution (HBSS) and incubated with a freshly prepared 2X pepsin
solution (340 mM NaCl,120 mM HCl, 0.52 mg/ml pepsin from porcine stomach mucosa – Sigma-
Aldrich), and then incubated at 37°C form 30 minutes before stopping the reaction by adding
complete DMEM.
In vitro-derived bradyzoites of the pSAG1-APT1 or Pru cell lines were removed with a cell scrapper
and subjected to a similar treatment, then passed through a 25G needle and a 40 µm filter prior to
cell sorting with a FACSCanto flow cytometer (Becton Dickinson) from the MRI cytometry facility.
510

Mouse infection, quantification of parasite burden by cyst enumeration and qPCR

Seven-week-old male CBA/J mice (Janvier Labs) were housed under specific pathogen-free
conditions and infected with 100 or 2000 tachyzoites of the appropriate strains intraperitoneally.
Infection was confirmed by assessing seropositivity of 21 days post infection using dot blot analysis
with parasite extracts. Mouse health status and body mass were monitored daily throughout acute
stage, and for up to 40 days.
At different timepoints post-infection, the brains were collected and homogenized with a glass
Potter in PBS. For parasite genomic DNA assessment, gDNA was extracted using the DNEasy
Blood & Tissue Kit (Qiagen ref 69506). As described earlier (73), a 529-bp repeat element in the
T. gondii genome was amplified using the TOX9 and TOX11 primers. The number of parasite
equivalent per µg of brain DNA was estimated by comparison with a standard curve, established
with a known number of Pru tachyzoites. For cyst enumeration, 5% of the homogenate was stained
with fluorescein-conjugated *Dolichos biflorus* agglutinin (Vector Laboratories). Cysts were counted
using an inverted fluorescence microscope.
Cyst area from both *in vivo* and *in vitro*-derived cysts was measured based on this cyst wall staining
using the ZEN software v2.5 (Zeiss).
527

Dendritic cell infection assay

MutuDC, a C57BL/6-derived dendritic cell line obtained from H. Acha-Orbea (44), were plated in a
6-well plate at 5.10^5 cells per well. Three hours later, MutuDC were infected with either parental
Pru, the pSAG1-APT1 line, or the pP21-DrpA^{K42A} line. After 24h, cells were incubated with FcR
block and eFluor 450 Fixable Viability Dye (1/1000, eBioscience) in PBS, fixed with 4% PFA,
permeabilized in PBS BSA 0,2% Saponine 0,05%, and labeled in that same buffer with AF647-
coupled rabbit antibodies that recognize the GRA6 C-terminal decapeptide HPGSVNEFDF
(custom-made, Biotem, Grenoble) for 1h at 4°C, allowing to detect vacuoles in which GRA6 was
actively secreted by live parasites. Samples were post-fixed using 1% PFA before acquisition on a
BD Fortessa and analyzed using FlowJo software.

Plaque assays

For plaque assays, confluent monolayers of HFFs seeded in 24 well plates were infected with freshly egresses tachyzoites or freshly isolated bradyzoites. Parasites were seeded in the first lane of the plate and diluted by 1/3 in each subsequent row. The number of parasites used in the first wells was as follows: 7000 (tachyzoites, manually counted), 3000 (FACS-sorted bradyzoites), or up to 4000 (qPCR-estimated tissue cyst-derived bradyzoites). Parasites which were left to grow for 10 days for tachyzoites and up to 20 days for bradyzoites. They were then fixed with 4% v/v PFA and plaques were revealed by staining with a 0.1% crystal violet solution (V5265, Sigma-Aldrich). Pictures of the plaques were acquired with an Olympus MVX10 microscope and plaques area were measured using the ZEN software v2.5 (Zeiss). Plaque area and plaque number of the mutant cell lines were expressed relatively to the mean value of those generated by the Pru control strain that was set to 100%

Immunofluorescence microscopy

Immunofluorescence assays (IFAs) of intracellular parasites were performed on coverslips containing infected HFF monolayers, fixed with 4% paraformaldehyde in PBS for 20 minutes at room temperature. Cells were washed three times with PBS, permeabilized with 0.3% Triton X-100/PBS for 15 minutes and then saturated with a 1% w/v bovine serum albumin (BSA)/PBS blocking solution for 30 minutes. Proteins were stained with primary antibodies for 1h, followed by three washes with PBS before incubation with secondary antibodies in a 1% BSA/PBS solution for 1h. Primary antibodies used in this study and their respective dilutions were mouse monoclonal anti-GFP at 1/1000 (clones 7.1 and 13.1, Roche), mouse monoclonal anti-TY at 1/250 (51), mouse anti-TgAtrx1 at 1/200 (37) and rabbit polyclonal anti-TgCpn60 at 1/2000 (41), mouse monoclonal anti-P21 at diluted 1/200 (T8 4G10) (74). Cysts were stained with a 1/300 dilution of a biotin-labelled *Dolichos biflorus* lectin (Sigma; L-6533) for 1h and revealed using a 1/300 dilution of FITC-conjugated streptavidin (Invitrogen; SNN1008). Staining of DNA was performed on fixed cells by incubating them for 5 min in a 1 µg/ml 4,6-diamidino-2-phenylindole (DAPI) solution. All images were acquired at the MRI facility from a Zeiss AXIO Imager Z2 epifluorescence microscope or a Zeiss lsm880 Fast Airyscan confocal microscope. Images were analyzed with by the ZEN software v2.5 (Zeiss) and ImageJ v1.53t (US National Institutes of Health). Z-stack acquisition and maximal intensity projection was systematically used to visualize the signal over the whole width of the cysts. Adjustments for brightness and contrast were applied uniformly on the entire image. Vacuoles or cysts were chosen randomly and identified based on the absence or presence of cyst wall labeling. The presence and the integrity of the apicoplast was evaluated through TgCpn60 staining and by comparing with the number of DAPI-stained nuclei in each cyst.

Immunoblot analysis

Protein extracts from freshly egressed tachyzoites or recently isolated bradyzoites were prepared in Laemmli sample buffer, separated by SDS-PAGE and transferred onto nitrocellulose membrane using the Mini-Transblot system (BioRad) according to the manufacturer's instructions. Primary antibodies used in this study and their respective dilutions were: mouse monoclonal anti-GFP at 1/1000 (clones 7.1 and 13.1, Roche), mouse monoclonal anti-TY at 1/200 (51), mouse monoclonal anti-P21 at diluted 1/50 (T8 4G10) (74), and mouse anti-actin at 1/100 (75).

Serological confirmation of infection by dot blot

Peripheral blood was collected intracardially without anticoagulant and centrifuged for 10 min at 14 000rpm. The serum was stored at -20°C. Tachyzoites were released from infected HFF, lysed in Laemlli buffer 1x (Biorad ref #1610737) + 5 % βmercapto-ethanol (BioRad ref #161-0710), and heated for 5 min at 70°C. Solubilized proteins were applied on a nitrocellulose membrane, which was incubated overnight at 4°C with each serum diluted at 1/1000. Immunologic detection was performed using horseradish peroxidase-conjuged anti-mouse IgG antibodies. Peroxidase activity was visualized by chemiluminescence and quantified using ChemiDoc camera (Biorad).

Semiquantitative RT-PCR

Total RNA was extracted from *T. gondii* tachyzoites using the Nucleospin RNA II kit (Macherey-Nagel). Reverse transcription was performed with the Superscript III first-strand synthesis kit (Invitrogen). 1 μg of RNA was used as a template per RT reaction. Specific primers of *T. gondii glyceraldehyde 3-phosphate dehydrogenase 2* (*GAPDH2*, ML5049/ML5680) or *TgAPT1* (ML4097/ML4098) were used for subsequent PCR amplification with GoTaq polymerase (Promega) for twenty-five cycles as follows: 15s of denaturation at 95 °C, 30s of annealing at 54 °C and 40s of elongation at 68°c. Similar volumes of samples for which no reverse transcriptase was included in the initial cDNA preparation were used as a control for genomic DNA contamination.

Statistical analyses

Statistical analyses (unpaired Student's t-test for comparisons between two groups, one-way ANOVA with post-hoc Tukey test for comparisons between three or more groups, Mann-Whitney non-parametric test for cyst size comparison, and Mantel–Cox logrank test for survival assays) were performed with Prism v8.3 (Graphpad). Unless specified, values are expressed as means + standard deviation (SD).

Acknowledgments

We are grateful to B. Striepen, D. Soldati-Favre and M. Parsons for providing plasmids and antibodies and H. Acha-Orbea for the DC cell line. We also thank M. Gissot and J.-F. Dubremetz for advice with the cyst purification protocol. Thanks to the MRI imaging and cytometry facilities for granting access to their equipment and thanks to Y. Bordat for help with formatting of the flow cytometry results. We thank R. Balouzat, R. Ecalard and the zootechnicians at Inserm UMS006-CREFRE mouse facility, and F. L'Faqihi-Olive, V. Duplan-Eche, A.-L. Iscache, and H. Garnier for assistance at the Infinity flow cytometry facility. We also thank the developers and the managers of the VeupathDB.org/ToxoDB.org databases, as well as scientists who contributed datasets. This work was supported by a grant from the Agence Nationale de la Recherche (ANR-19-CE15-0023) to N.B. and S.B. and the PIA PARAFRAP Consortium (ANR-11-LABX0024) to N.B.

References

1. S. G. Sanchez, S. Besteiro, The pathogenicity and virulence of Toxoplasma gondii. *Virulence* **12**, 3095–3114 (2021).

2. M. Sasai, M. Yamamoto, Innate, adaptive, and cell-autonomous immunity against Toxoplasma gondii infection. *Exp Mol Med* **51**, 1–10 (2019).

3. L. D. Sibley, J. C. Boothroyd, Virulent strains of Toxoplasma gondii comprise a single clonal lineage. *Nature* **359**, 82–85 (1992).

4. A. Cerutti, N. Blanchard, S. Besteiro, The bradyzoite: a key developmental stage for the persistence and pathogenesis of toxoplasmosis. *Pathogens* **9**, 234 (2020).

5. E. Watts, *et al.*, Novel Approaches Reveal that Toxoplasma gondii Bradyzoites within Tissue Cysts Are Dynamic and Replicating Entities In Vivo. *MBio* **6**, e01155-01115 (2015).

6. K. J. Pittman, L. J. Knoll, Long-term relationships: the complicated interplay between the host and the developmental stages of Toxoplasma gondii during acute and chronic infections. *Microbiol Mol Biol Rev* **79**, 387–401 (2015).

7. D. J. Ferguson, W. M. Hutchison, An ultrastructural study of the early development and tissue cyst formation of Toxoplasma gondii in the brains of mice. *Parasitol. Res.* **73**, 483–491 (1987).

8. L. Lemgruber, P. Lupetti, E. S. Martins-Duarte, W. De Souza, R. C. Vommaro, The organization of the wall filaments and characterization of the matrix structures of Toxoplasma gondii cyst form: Structure of Toxoplasma gondii cyst. *Cellular Microbiology* **13**, 1920–1932 (2011).

646   9.   J. P. Dubey, D. S. Lindsay, C. A. Speer, Structures of Toxoplasma gondii tachyzoites,
647   bradyzoites, and sporozoites and biology and development of tissue cysts. *Clin. Microbiol.*
648   *Rev.* **11**, 267–299 (1998).

649   10.   S. Yang, S. F. Parmley, Toxoplasma gondii expresses two distinct lactate dehydrogenase
650   homologous genes during its life cycle in intermediate hosts. *Gene* **184**, 1–12 (1997).

651   11.   F. Dzierszinski, *et al.*, The protozoan parasite Toxoplasma gondii expresses two functional
652   plant-like glycolytic enzymes. Implications for evolutionary origin of apicomplexans. *J. Biol.*
653   *Chem.* **274**, 24888–24895 (1999).

654   12.   F. Dzierszinski, M. Mortuaire, N. Dendouga, O. Popescu, S. Tomavo, Differential expression
655   of two plant-like enolases with distinct enzymatic and antigenic properties during stage
656   conversion of the protozoan parasite Toxoplasma gondii. *J. Mol. Biol.* **309**, 1017–1027
657   (2001).

658   13.   S. Tomavo, The differential expression of multiple isoenzyme forms during stage conversion
659   of Toxoplasma gondii: an adaptive developmental strategy. *International Journal for*
660   *Parasitology* **31**, 1023–1031 (2001).

661   14.   A. E. Abdelbaset, *et al.*, Lactate dehydrogenase in Toxoplasma gondii controls virulence,
662   bradyzoite differentiation, and chronic infection. *PLoS ONE* **12**, e0173745 (2017).

663   15.   K. R. Buchholz, *et al.*, Identification of tissue cyst wall components by transcriptome analysis
664   of in vivo and in vitro Toxoplasma gondii bradyzoites. *Eukaryotic Cell* **10**, 1637–1647 (2011).

665   16.   K. J. Pittman, M. T. Aliota, L. J. Knoll, Dual transcriptional profiling of mice and Toxoplasma
666   gondii during acute and chronic infection. *BMC Genomics* **15**, 806 (2014).

667   17.   S. Skariah, M. K. McIntyre, D. G. Mordue, Toxoplasma gondii: determinants of tachyzoite to
668   bradyzoite conversion. *Parasitol Res* **107**, 253–260 (2010).

669   18.   T. Mouveaux, *et al.*, Primary brain cell infection by Toxoplasma gondii reveals the extent
670   and dynamics of parasite differentiation and its impact on neuron biology. *Open Biol* **11**,
671   210053 (2021).

672   19.   C. Christiansen, *et al.*, In vitro maturation of Toxoplasma gondii bradyzoites in human
673   myotubes and their metabolomic characterization. *Nat Commun* **13**, 1168 (2022).

674   20.   J. Mayoral, M. Di Cristina, V. B. Carruthers, L. M. Weiss, Toxoplasma gondii: bradyzoite
675   differentiation in vitro and in vivo. *Methods Mol. Biol.* **2071**, 269–282 (2020).

676   21.   M. Blume, F. Seeber, Metabolic interactions between Toxoplasma gondii and its host.
677   *F1000Res* **7** (2018).

678   22.   G. G. van Dooren, B. Striepen, The algal past and parasite present of the apicoplast. *Annu.*
679   *Rev. Microbiol.* **67**, 271–289 (2013).

23.   G. G. van Dooren, S. V. Hapuarachchi, "The dark side of the chloroplast: biogenesis,
metabolism and membrane biology of the apicoplast" in *Advances in Botanical Research*,
(Elsevier, 2017), pp. 145–185.

24.   F. Seeber, D. Soldati-Favre, "Metabolic pathways in the apicoplast of apicomplexa" in
International Review of Cell and Molecular Biology, (2010), pp. 161–228.

25.   I. R. Dunay, K. Gajurel, R. Dhakal, O. Liesenfeld, J. G. Montoya, Treatment of toxoplasmosis:
historical perspective, animal models, and current clinical practice. *Clin Microbiol Reviews*
31, e00057-17, /cmr/31/4/e00057-17.atom (2018).

26.   O. Djurkovic-Djakovic, Efficacy of atovaquone combined with clindamycin against murine
infection with a cystogenic (Me49) strain of Toxoplasma gondii. *Journal of Antimicrobial*
Chemotherapy **50**, 981–987 (2002).

27.   I. Benmerzouga, *et al.*, Guanabenz repurposed as an antiparasitic with activity against acute
and latent toxoplasmosis. *Antimicrob. Agents Chemother.* **59**, 6939–6945 (2015).

28.   D. J. Ferguson, J. Huskinson-Mark, F. G. Araujo, J. S. Remington, An ultrastructural study of
the effect of treatment with atovaquone in brains of mice chronically infected with the
ME49 strain of Toxoplasma gondii. *Int J Exp Pathol* **75**, 111–116 (1994).

29.   R. S. R. Vidadala, *et al.*, Development of an orally available and central nervous system (CNS)
penetrant Toxoplasma gondii Calcium-Dependent Protein Kinase 1 (TgCDPK1) Inhibitor with
minimal human ether-a-go-go-related gene (hERG) activity for the treatment of
Toxoplasmosis. *J. Med. Chem.* **59**, 6531–6546 (2016).

30.   J. S. Doggett, *et al.*, Endochin-like quinolones are highly efficacious against acute and latent
experimental toxoplasmosis. *Proc Natl Acad Sci U S A* **109**, 15936–15941 (2012).

31.   N. Konstantinovic, H. Guegan, T. Stäjner, S. Belaz, F. Robert-Gangneux, Treatment of
toxoplasmosis: current options and future perspectives. *Food Waterborne Parasitol* **15**,
e00036 (2019).

32.   W. K. Chew, I. Segarra, S. Ambu, J. W. Mak, Significant reduction of brain cysts caused by
Toxoplasma gondii after treatment with spiramycin coadministered with metronidazole in a
mouse model of chronic toxoplasmosis. *Antimicrob Agents Chemother* **56**, 1762–1768
(2012).

33.   S. Tomavo, J. C. Boothroyd, Interconnection between organellar functions, development
and drug resistance in the protozoan parasite, Toxoplasma gondii. *Int J Parasitol* **25**, 1293–
1299 (1995).

34.   Y. Murata, T. Sugi, L. M. Weiss, K. Kato, Identification of compounds that suppress
Toxoplasma gondii tachyzoites and bradyzoites. *PLoS One* **12**, e0178203 (2017).

35.   S. Briquet, M. Gissot, O. Silvie, A toolbox for conditional control of gene expression in
apicomplexan parasites. *Mol Microbiol* **117**, 618–631 (2022).

36. C. F. Brooks, *et al.*, The toxoplasma apicoplast phosphate translocator links cytosolic and
apicoplast metabolism and is essential for parasite survival. *Cell Host Microbe* **7**, 62–73
(2010).

37. A. E. DeRocher, *et al.*, A thioredoxin family protein of the apicoplast periphery identifies
abundant candidate transport vesicles in Toxoplasma gondii. *Eukaryotic Cell* **7**, 1518–1529
(2008).

38. T. Tomita, *et al.*, The Toxoplasma gondii cyst wall protein CST1 is critical for cyst wall
integrity and promotes bradyzoite persistence. *PLoS Pathog.* **9**, e1003823 (2013).

39. S. C. Nair, *et al.*, Apicoplast isoprenoid precursor synthesis and the molecular basis of
fosmidomycin resistance in Toxoplasma gondii. *J. Exp. Med.* **208**, 1547–1559 (2011).

40. M. Okada, *et al.*, Critical role for isoprenoids in apicoplast biogenesis by malaria parasites.
eLife **11**, e73208 (2022).

41. S. Agrawal, G. G. van Dooren, W. L. Beatty, B. Striepen, Genetic evidence that an
endosymbiont-derived endoplasmic reticulum-associated protein degradation (ERAD)
system functions in import of apicoplast proteins. *J. Biol. Chem.* **284**, 33683–33691 (2009).

42. D. Smith, *et al.*, Toxoplasma TgATG9 is critical for autophagy and long-term persistence in
tissue cysts. *eLife* **10**, e59384 (2021).

43. A. Sanecka, E.-M. Frickel, Use and abuse of dendritic cells by Toxoplasma gondii. *Virulence* **3**,
678–689 (2012).

44. S. A. Fuertes Marraco, *et al.*, Novel murine dendritic cell lines: a powerful auxiliary tool for
dendritic cell research. *Front Immunol* **3**, 331 (2012).

45. K. Kennedy, E. M. Crisafulli, S. A. Ralph, Delayed death by plastid inhibition in apicomplexan
parasites. *Trends in Parasitology* **35**, 747–759 (2019).

46. G. G. van Dooren, *et al.*, A novel dynamin-related protein has been recruited for apicoplast
fission in Toxoplasma gondii. *Curr. Biol.* **19**, 267–276 (2009).

47. B. A. Fox, *et al.*, Type II Toxoplasma gondii KU80 knockout strains enable functional analysis
of genes required for cyst development and latent infection. *Eukaryotic Cell* **10**, 1193–1206
(2011).

48. W. Bohne, D. S. Roos, Stage-specific expression of a selectable marker in Toxoplasma gondii
permits selective inhibition of either tachyzoites or bradyzoites. *Mol. Biochem. Parasitol.* **88**,
115–126 (1997).

49. M. Soete, B. Fortier, D. Camus, J. F. Dubremetz, Toxoplasma gondii: kinetics of bradyzoite-
tachyzoite interconversion in vitro. *Exp. Parasitol.* **76**, 259–264 (1993).

50. S. K. Dogga, *et al.*, Importance of aspartyl protease 5 in the establishment of the
intracellular niche during acute and chronic infection of Toxoplasma gondii. *Mol Microbiol*
(2022) https:/doi.org/10.1111/mmi.14987.

51. P. Bastin, A. Bagherzadeh, K. R. Matthews, K. Gull, A novel epitope tag system to study
protein targeting and organelle biogenesis in Trypanosoma brucei. *Molecular and*
Biochemical Parasitology **77**, 235–239 (1996).

52. B. A. Fox, D. J. Bzik, De novo pyrimidine biosynthesis is required for virulence of Toxoplasma
gondii. *Nature* **415**, 926–929 (2002).

53. A. Bougdour, *et al.*, Drug inhibition of HDAC3 and epigenetic control of differentiation in
Apicomplexa parasites. *J. Exp. Med.* **206**, 953–966 (2009).

54. J. P. Boyle, J. P. J. Saeij, M. D. Cleary, J. C. Boothroyd, Analysis of gene expression during
development: lessons from the Apicomplexa. *Microbes and Infection* **8**, 1623–1630 (2006).

55. M. Soete, D. Camus, J. F. Dubremetz, Experimental induction of bradyzoite-specific antigen
expression and cyst formation by the RH strain of Toxoplasma gondii in vitro. *Experimental*
Parasitology **78**, 361–370 (1994).

56. N. Blanchard, I. R. Dunay, D. Schlüter, Persistence of Toxoplasma gondii in the central
nervous system: a fine-tuned balance between the parasite, the brain and the immune
system. *Parasite Immunol* **37**, 150–158 (2015).

57. A. Salvioni, *et al.*, Robust control of a brain-persisting parasite through MHC I presentation
by infected neurons. *Cell Rep* **27**, 3254-3268.e8 (2019).

58. S. Ye, M. Lunghi, D. Soldati-Favre, A signaling factor linked to Toxoplasma gondii guanylate
cyclase complex controls invasion and egress during acute and chronic infection. *mBio* **13**,
e0196522 (2022).

59. Y. Xue, *et al.*, A single-parasite transcriptional atlas of Toxoplasma gondii reveals novel
control of antigen expression. *Elife* **9** (2020).

60. A. P. Sinai, *et al.*, Reexamining chronic Toxoplasma gondii infection: surprising activity for a
"dormant" parasite. *Curr Clin Microbiol Rep* **3**, 175–185 (2016).

61. D. Walsh, N. J. Katris, L. Sheiner, C. Y. Botté, Toxoplasma metabolic flexibility in different
growth conditions. *Trends in Parasitology* **38**, 775–790 (2022).

62. E. K. Szabo, C. A. M. Finney, Toxoplasma gondii : one organism, multiple models. *Trends in*
Parasitology **33**, 113–127 (2017).

63. G. F. Watson, P. H. Davis, Systematic review and meta-analysis of variation in Toxoplasma
gondii cyst burden in the murine model. *Exp Parasitol* **196**, 55–62 (2019).

64. K. Frénal, *et al.*, Myosin-dependent cell-cell communication controls synchronicity of
division in acute and chronic stages of Toxoplasma gondii. *Nat Commun* **8**, 15710 (2017).

65. F. Dzierszinski, M. Nishi, L. Ouko, D. S. Roos, Dynamics of Toxoplasma gondii differentiation.
Eukaryotic Cell **3**, 992–1003 (2004).

66. D. J. P. Ferguson, *et al.*, Maternal inheritance and stage-specific variation of the apicoplast in
Toxoplasma gondii during development in the intermediate and definitive host. *Eukaryotic*
Cell **4**, 814–826 (2005).

67. Y. Oren, Standing on the shoulders of microbes: how cancer biologists are expanding their
view of hard-to-kill persister cells. *Molecular Systems Biology* **18** (2022).

68. P. Martrou, M. Pestre, R. Loubet, Nicolas, G. Malinvaud, La toxoplasmose congénitale (note
concernant un cas mortel). *Limousin Médical* **53**, 3–7.

69. M.-H. Huynh, V. B. Carruthers, Tagging of endogenous genes in a Toxoplasma gondii strain
lacking Ku80. *Eukaryot Cell* **8**, 530–539 (2009).

70. S. M. Sidik, C. G. Hackett, F. Tran, N. J. Westwood, S. Lourido, Efficient genome engineering
of Toxoplasma gondii using CRISPR/Cas9. *PLoS ONE* **9**, e100450 (2014).

71. L. Sheiner, *et al.*, A Systematic Screen to Discover and Analyze Apicoplast Proteins Identifies
a Conserved and Essential Protein Import Factor. *PLoS Pathog* **7** (2011).

72. A. W. C. A. Cornelissen, J. P. Overdulve, J. M. Hoenderboom, Separation of Isospora
(Toxoplasma) gondii cysts and cystozoites from mouse brain tissue by continuous density -
gradient centrifugation. *Parasitology* **83**, 103–108 (1981).

73. V. Feliu, *et al.*, Location of the CD8 T cell epitope within the antigenic precursor determines
immunogenicity and protection against the Toxoplasma gondii parasite. *PLoS Pathog.* **9**,
e1003449 (2013).

74. S. Tomavo, *et al.*, Characterization of bradyzoite-specific antigens of Toxoplasma gondii.
Infect Immun **59**, 3750–3753 (1991).

75. A. Herm-Gotz, Toxoplasma gondii myosin A and its light chain: a fast, single-headed, plus-
end-directed motor. *The EMBO Journal* **21**, 2149–2158 (2002).

809
810

  Figures and Tables

Figure 1. Conditional depletion of APT1 at the bradyzoite stage. A) Schematic representation
of APT1 function at the apicoplast and its implication upstream of two important metabolic
pathways. B) Scheme describing the strategy to edit the *APT1* locus by homologous
recombination, to express a copy of the *APT1* gene under the dependence of a tachyzoite-
specific *SAG1* promoter and allowing fusion of APT1 with a C-terminal GFP. C) Immunoblot
analysis of protein extracts from tachyzoites (Tz) and 12 days *in vitro*-differentiated (alkaline
stress) bradyzoites (Bz) with an anti-GFP antibody to detect the expression of APT1-GFP
(arrowhead) under the dependence of the *SAG1* promoter. Actin was used as a loading control.
D) Microscopic imaging of pSAG1-APT1 parasites submitted to alkaline pH stress for 7 days.
Cyst walls were labelled with *D. biflorus* lectin (DBL) or outlined with dashed lines. A more intense
DBL labelling outlines a more mature cyst (arrowhead). DNA was labelled with DAPI. Scale
bar=10 µm. E) The GFP-APT1 signal in vacuoles containing tachyzoites, or cysts generated by
alkaline pH-induced differentiation, was quantified and shows progressive signal loss upon stage
conversion. Data are mean from *n*=3 independent experiments +SD. *** $p \leq$ 0.001, **** $p \leq$
0.0001, by one-way ANOVA followed by Tukey post-hoc test.

Figure 2. Depletion of APT1 during in vitro differentiation impacts apicoplast maintenance and parasite development and viability. A) Alkaline pH-induced *in vitro* differentiation of pSAG1-APT1 parasites for 7 or 14 days leads to concomitant disappearance of APT1 and of the Cpn60 apicoplast marker (magenta) in mature cysts that are more intensely labelled with DBL (arrowhead). A 14 day-differentiated cyst of the Pru cell line is shown as a control. Cyst wall is outlined with dashed lines. DNA was stained with DAPI. Scale bar=10 μm. B) The integrity of the Cpn60-labelled apicoplast signal was quantified after 14 days of alkaline pH-induced differentiation and showed a significant loss in the pSAG1-APT1 cell line. Data are mean from *n*=6 independent experiments +SD. **** $p \leq 0.0001$, unpaired Student's *t*-test. C) Quantification of

the area of *in vitro*-differentiated cysts kept for 14 days under alkaline pH stress shows that the
pSAG1-APT1 cell line generates smaller cysts. Data are mean from *n*=3 independent
experiments. Symbols are matched between identical experimental groups. **** *p* ≤ 0.0001, non-
parametric Mann-Whitney test. D) Plaque assays were carried out by infecting HFF monolayers
with tachyzoites or bradyzoites obtained after 14 days of alkaline pH stress and FACS-based
isolation. GFP-negative bradyzoites of the pSAG1-APT1 cell line generates less and smaller
plaques than the Pru control cell lines. Measurements of lysis plaque area E) and plaque number
F) highlight a significant defect in the lytic cycle of pSAG1-APT1 bradyzoites. Values are means
of nine values (representing *n*=3 independent experiments, containing three technical triplicates)
+SD. Symbols are matched between identical experimental groups. Mean value for the Pru cell
line was set to 100% as a reference. **** *p* ≤ 0.0001, unpaired Student's t test.

852
Figure 3. Infection of mice with pSAG1-APT1 parasites leads to a severe acute phase and
high cyst burden but generates essentially non-viable bradyzoites. A) Survival curves of
CBA mice infected with 100 parasites of the Pru or pSAG1-APT1 cell lines shows that the latter
induces an excessive mortality during acute phase. Data from *n*=3 independent experiments are
plotted. * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, log-rank Mantel-Cox test. B) Monitoring of body
mass of mice infected with Pru or pSAG1-APT1 tachyzoites. Infection with pSAG1-APT1 induces
a more important mass loss than the control cell line, reflecting a more severe acute phase. Data

represent mean from n=3 independent experiments +SD. **** p ≤ 0.0001, unpaired Student's t
test. C) Quantification of cyst burden in infected mice forty days post-infection. Values are from
n=3 independent experiments and are expressed as means +SD. Symbols are matched between
identical experimental groups. *** p ≤ 0.001, unpaired Student's t test. D) Parasite burden
measured by qPCR on genomic DNA from infected mice. Values are from n=3 independent
experiments and are expressed as means +SD. Symbols are matched between identical
experimental groups. **** p ≤ 0.0001, unpaired Student's t test. E) Labelling of mouse brain-
derived cysts with anti-Cpn60 shows partial loss of the apicoplast in the pSAG1-APT1 cell line. F)
Quantification of Cpn60 signal in brain-derived pSAG1-APT1 cysts shows that they display partial
or total apicoplast loss 40 days post-infection. Shown are the mean +SD from n=5 independent
experiments, at least 12 cysts for each condition were assessed. *** p ≤ 0.01, unpaired Student's
t-test. G) Quantification of the area of brain-derived cysts obtained 40 days post-infection shows
that the pSAG1-APT1 cell line generates smaller cysts. Data are mean from n=3 independent
experiments. Symbols are matched between identical experimental groups. **** p ≤ 0.0001, non-
parametric Mann-Whitney test. H) A representative plaque assay carried out by infecting HFF
monolayers with brain-derived bradyzoites obtained after 40 days of infection shows that the
pSAG1-APT1 cell line generates very few plaques. Measurements of lysis plaque area I) and
plaque number J) confirms that there is a significant defect in the lytic cycle of brain-derived
pSAG1-APT1 bradyzoites. Values are means from nine replicates from n=2 independent
experiments +SD. Mean value for the Pru cell line was set to 100% as a reference. Symbols are
matched between identical experimental groups. *** p ≤ 0.001, unpaired Student's t test.
881

882
883
Figure 4. Generating a parasite line expressing a bradyzoite-specific dominant-negative
DrpA mutant. A) Schematic representation of DrpA function in apicoplast fission and inheritance
during parasite division. B) Scheme describing the strategy to edit the *UPRT* locus by
homologous recombination, in order to introduce a mutated version of the *DrpA* gene and
express the DrpA^{K42A} protein under the dependence of a bradyzoite-specific *P21* promoter. C)
Immunoblot analysis of protein extracts from tachyzoites (Tz) and 6 days *in vitro*-differentiated
(apicidin treatment) bradyzoites (Bz) with an anti-TY antibody to detect the expression of the
DrpA^{K42A} protein (arrowhead) under the dependence of the *P21* promoter. Actin was used as a
loading control and P21 as a differentiation control.

Figure 5. DrpA^{K42A}-expressing bradyzoites are impacted for apicoplast division and fail to
establish chronic phase in mice. A) Upon apicidin-induced *in vitro* differentiation for 6 days, the
DrpA^{K42A} protein is expressed and bradyzoites show extensive reticulation of their apicoplast
(Cpn60-labelled, magenta, magnified in insets) in DBL-labelled cysts. The Pru parental cell line
and tachyzoites of the two cell lines are shown as a control. Cyst wall is outlined with dashed
lines. DNA was stained with DAPI. Scale bar=10 µm (5 µm for insets). B) The proportion of cysts
with abnormal apicoplast signal (reticulated or lost) after 6 days of apicidin treatment was
quantified. Values are means from *n*=3 independent experiments +SD. Symbols are matched
between identical experimental groups. *** *p* ≤ 0.001, 'ns' not significant, unpaired Student's t
test. C) Quantification of the size of *in vitro*-differentiated cysts kept for 6 days in the presence of
apicidin shows no size difference between cysts of the pP21-DrpA^{K42A} cell line and the Pru
control. Data are mean from *n*=3 independent experiments. Symbols are matched between
identical experimental groups. 'ns' not significant, non-parametric Mann-Whitney test. D) A
representative plaque assay carried out by infecting HFF monolayers with tachyzoites shows that
the pP21-DrpA^{K42A} parasites are not affected in their lytic cycle *in vitro*. E) Measurement of lysis
plaque areas confirms that there is no major defect in the lytic cycle of pP21-DrpA^{K42A}
tachyzoites. Values are means from *n*=3 independent experiments (each one with three technical
replicates) +SD. Mean value for the Pru cell line was set to 100% as a reference. Symbols are
matched between identical experimental groups. **** *p* ≤ 0.0001, unpaired Student's t test. F) *In*
vitro dendritic cell infection assay was performed and analyzed by flow cytometry and shows no
impairment of the invasive capacity of pP21-DrpA^{K42A} tachyzoites. Data are mean from *n*=3
independent experiments; 'ns' not significant, unpaired Student's t test. G) Monitoring of body
mass of mice infected with Pru or of pP21-DrpA^{K42A} tachyzoites shows that the acute phase
occurs normally with of pP21-DrpA^{K42A} parasites. Data represent mean from 20 individual mice
from *n*=2 independent experiments +SD. 'ns' not significant, unpaired Student's t test. H) Parasite
burden at 12 days or 40 days post-infection measured by qPCR on genomic DNA from infected
mice. Values are from *n*=2 (12 days timepoint) or *n*=3 (40 days timepoint) independent
experiments, performed with an inoculum of 100 parasites or 2000 parasites, and are expressed
as means +SD. ** *p* ≤ 0.01, 'ns' not significant, unpaired Student's t test.