PRDM16 restricts stem cell proliferation by integrating BMP and Wnt signalling in the

developing choroid plexus

Li He[1], Jiayu Wen[2, *] and Qi Dai[1, *]

1 Department of Molecular Bioscience, the Wenner-Gren Institute, Stockholm University

2 Division of Genome Sciences and Cancer, The John Curtin School of Medical Research, The
Australian National University

* Co-corresponding authors

Abstract

Cell proliferation, quiescence and differentiation are well balanced during organismal growth and tissue homeostasis. Understanding how such balance is achieved is crucial for tackling disease mechanisms such as those in cancer and neurological disorders. Here we have identified PRDM16 as a key regulator for BMP-induced neural stem cell (NSC) quiescence. PRDM16 acts as a co-repressor in the BMP pathway to repress cell proliferation genes. Such function of PRDM16 is required for the proper specification of choroid plexus (ChP) epithelial cells. Using a single-cell resolution fluorescent *in situ* approach, we show that NSC proliferation and Wnt pathway genes are abnormally elevated in the *Prdm16* mutant ChP. These findings uncover an essential function of PRDM16 in stem cell regulation, Wnt and BMP signalling. PRDM16 is a cell fate determinant in multiple tissue types. Our finding opens a possibility that PRDM16 uses similar strategies in relevant contexts and disease conditions.

Introduction

Uncontrollable cell proliferation can lead to tumor growth, while premature differentiation can result in tissue degeneration. How stem cell proliferation and quiescence are regulated is a fundamental question in developmental and stem cell biology. BMP (Bone morphogenetic proteins) signaling is a key cell-signaling pathway that plays a role in inducing and maintaining stem cell quiescence. Additionally, BMP signaling is involved in numerous cell specification processes (Zhang and Li, 2005).

The ability of a single pathway to play a diverse range of roles relies on context-specific transcriptional outputs. SMAD4 is the common transcription factor that mediates BMP and TGF-β/Activin signaling. Upon BMP induction, SMAD4 associates with phosphorylated SMAD1/5/8, while in response to TGF-β/or Activin, it complexes with phosphorylated SMAD2/3. It is known that the SMAD protein complexes can both activate and repress target gene expression. However, the precise mechanism that determines the regulatory activity of SMADs and target gene selection remains unclear (Hill, 2016).

In the mammalian brain, BMP and Wnt signaling pattern the brain midline where an essential structure, the ChP, emerges. Neural epithelial cells at the site of presumptive ChP lose neural potential, and the majority of them exit cell cycle. Only a few cells at the boarder of the ChP primordium and cortical hem (CH) remain as slowly dividing ChP progenitors which expand the monolayered ChP epithelial cells (Liddelow et al., 2010). BMP signaling is essential for ChP epithelium specification. Conditional depletion of the BMP receptor *BMPr1a* diminishes ChP development (Hebert et al., 2002), while ectopic BMP was sufficient to transform neural progenitors into ChP progenitors (Watanabe et al., 2012). Wnt signaling peaking at CH is also required for ChPe cell specification (al, 2020). Conditional depletion of ß-Catenin resulted in defective ChP, while overexpression of ß-Catenin expands CH at the expense of the ChP epithelium. These observations suggest that the level of Wnt and BMP signaling needs to be tightly controlled to ensure proper cell specification.

The PR domain-containing (PRDM) family protein PRDM16 determines cell fate specification in various cell types (Aguilo et al., 2011; Baizabal et al., 2018; Chui et al., 2020; Chuikov et al., 2010; Inoue et al., 2017; Kajimura et al., 2008; Shimada et al., 2017). Consistent with its essential roles in normal development, mutation and dysregulation of *Prdm16* are linked with several human diseases including those identified in the patients with 1p36 chromosomal

aberrations. It has been shown that deletion of *PRDM16* has a causal role in cardiomyopathy

(Arndt et al., 2013), while co-amplification of *PRDM16* and *SKI* was shown to promote tumour

growth via suppressing TGF-β anti-proliferation activity in gastric cancer cells (Takahata et al.,

2009). Moreover, as a master regulator of the thermogenic gene expression program in brown

adipocytes, PRDM16 has been studied as a promising target to treat obesity and other metabolic

diseases (Horvath and Scheele, 2022).

PRDM16 has versatile functions at the molecular level. It regulates chromatin accessibility and

epigenetic states of its bound enhancers (Baizabal et al., 2018; He et al., 2021), and depending

on the associated cofactors, PRDM16 can repress or activate gene expression. This poses a

challenge when targeting PRDM16 is a therapeutic strategy in disease treatment because

undesired outcome may occur. Thus, it is important to get a full understanding of the regulatory

roles of this protein in each individual process.

The ChP epithelium is directly specified from highly proliferative neural epithelium, making it

a tractable model to study mechanisms that regulate the transition between stem cell

proliferation and differentiation. Here we find that *Prdm16* mutant ChP cells fail to exit cell

cycle and were still proliferating NSCs. Further we show that PRDM16 collaborates with the

BMP pathway effectors, SMAD4 and pSMAD1/5/8, to suppress cell proliferating genes

including cell cycle regulators and Wnt signaling genes in quiescent NSCs. Finally, we

demonstrate that Wnt signaling and cell proliferation are abnormally elevated in the *Prdm16*

mutant ChP. Together, our work defines the mechanism by which PRDM16 functions in the

BMP pathway and suggests a regulatory circuit composed of BMP and Wnt signaling and

PRDM16 in the control of stem cell quiescence.

Results

Prdm16 is required for the specification of the ChP epithelium

BMP signaling starts to specify the presumptive ChP at embryonic day E10.5 in mice (Hebert et al., 2002) when *Prdm16* mRNAs become detectable in this region (**Fig. 1A**). Previously we reported that animals of the *Prdm16* knockout (KO) null allele, *Prdm16^{cGT}* (*cGT*), show severely reduced ChP structure (**Fig 1B**) (Bjork et al., 2010; He et al., 2021; Strassman et al., 2017). In the *Prdm16* KO mutant, expression of the ChP marker gene, *Ttr*, is not only reduced in the lateral ventricle (tChPs) but also in the fourth ventricle (hChP) (**Fig. 1C-D**), suggesting that the function of PRDM16 is not area-restricted but generally required for ChP development.

To assess the overall patterning of the CH and ChP region, we checked expression of *Wnt2b* and *BMP4* by conventional RNA *in situ* hybridization. The *Wnt2b* signal appeared normal in mutant brains compared with the controls at E11.5 and E12.5 (**Supplementary Fig. 1A**), suggesting that the area of CH was not affected by *Prdm16* deletion. Although the morphology of *Prdm16* mutant ChP epithelium looked defective, as it remained as thick as the neural epithelium, the level and range of *BMP4* expression in wild-type and mutant brains appeared comparable (**Supplementary Fig. 1A**). This result indicates that loss of *Prdm16* does not affect overall patterning of the medial-lateral region nor expression of *Wnt2b* and *BMP4*.

Neural epithelial cells in the ChP primordium rely on PRDM16 to exit cell cycle and loss neural potential

Next, we attempted to examine cellular changes underlying defective ChP in the *Prdm16* mutant. As the mutant ChP epithelium appeared thicker than the wild-type (**Fig. 1C-D**), we examined cell proliferation of the ChP cells by a 2-hour EdU labeling at E12.5. In control embryos from the same litter, the ChP cells were mostly EdU negative and transitioning into a monolayer, suggesting that these cells have exited cell cycle. By contrast, *Prdm16* mutant ChP cells remained highly proliferative (**Fig. 1E**).

To further check the property of *Prdm16* mutant ChP cells, we stained the control and mutant brain slices with the neuronal marker Doublecortin (Dcx). The control brain lacked Dcx positive cells in the ChP, suggesting that neural epithelial cells were specified into ChP epithelial cells and thus lost neural potential (**Fig. 1F**). However, there were a number of Dcx

positive cells lining along the basal side of the epithelium in the *Prdm16* mutant ChP, indicative of continued neurogenesis.

To further examine the property of ChP cells, we stained the brain sections with the stem cell marker SOX2. SOX2 is highly expressed in NSCs adjacent to the ChP epithelium, but its expression became significantly reduced in the presumptive ChP area (**Supplementary Fig. 1C**). However, such reduction was abolished in the *Prdm16* mutant.

Together, these results demonstrate that *Prdm16* depletion disrupted the transition from neural progenitors into ChP epithelia, leading to ectopic proliferation and neurogenesis. This phenotype is a reminiscence of the BMP pathway mutant, Foxg1-cre::loxp*Bmpr1a,* where ectopic cell proliferation was found in the developing ChP (Hebert et al., 2002). We then wondered whether PRDM16 interacts with the BMP pathway to restrict cell proliferation.

PRDM16 and BMP signaling collaborate to induce NSC quiescence

It is known that BMP signaling maintains adult NSC quiescence and that BMPs are sufficient to induce cultured proliferative NSCs into quiescence (Mira et al., 2010). To determine how PRDM16 may interact with BMP signaling and regulates cell proliferation, we switched to using cultured primary NSCs (**Fig. 2A**). The PRDM16 protein is highly expressed in the nucleus of NSCs in embryonic brain tissues (**Fig. 1B**, and (He et al., 2021)). Unexpectedly, the protein becomes nearly undetectable when primary NSCs are cultured *in vitro* (**Supplementary Fig. 2A**) although *Prdm16* mRNA levels are still high (**Supplementary Fig. 2B**). This suggests that an unknown mechanism controls the stability and nuclear localization of PRDM16 *in vivo*. It was reported that post-translational modification of PRDM16 is an important regulatory switch for PRDM16 function in brown adipocytes (Wang et al., 2022). We speculate a similar mechanism is used in NSCs in the brain. To overcome this issue and examine potential molecular function of PRDM16 in NSC proliferation and gene regulation, we infected primary NSCs with a lenti-viral construct that constitutively expresses *3xNSL_Flag_Prdm16*, resulting in detectable PRDM16 protein in the nucleus and cytoplasm (**Supplementary Fig. 2A**). We referred this cell line as to *Prdm16_expressing (Prdm16_E)*. For comparison, we cultured NSCs from *Prdm16_*KO brains from the same developing stage and brain area.

Similar to previously reported (Mira et al., 2010), BMP4 was able to induce cell quiescence of proliferative *Prdm16_E*xpressing NSCs, as indicated by the reduction in the number of mKi67-

stained or EdU-labeled cells (**Fig. 2B-C**). By contrast, many of the *Prdm16*_KO cells failed to

exit cell cycle after BMP4 addition, evidenced by less reduction of the mKI67 or EdU positive

cell population (**Fig. 2B-C**) and reduced *mKi67* mRNA levels measured by reverse-

transcription followed by quantitative PCR (RT-qPCR) assay (**Fig. 2D**). Thus, similar to their

function in ChP specification, both BMP signaling and PRDM16 are required for inducing

NSCs to exit cell cycle. We then went forward to examine molecular function of PRDM16 and

BMP signaling in this context.

PRDM16 assists genomic binding of the BMP pathway transcription factors

As PRDM16 is a transcriptional regulator, we hypothesized that the interaction of PRDM16

and BMP signaling, if there is any, should occur at the genome. To this end, we monitored

changes of transcription factor binding in the genome during the transition of proliferation to

quiescence state. Using *Prdm16_E* and *Prdm16_KO* NSCs treated with BMP4 and those non-

treated, we performed CUT&TAG experiments with the PRDM16 antibody. As none of the

SMAD antibodies worked in CUT&TAG assays in our tests, we performed ChIP-seq with the

antibodies targeting all three classes of SMAD transcription factors including SMAD4,

SMAD3 and phosphorylated SMAD1/5/8. Thus, our experiment settings allow profiling of

PRDM16 binding sites together with SMAD binding sites induced by endogenous BMPs and

TGF-β/Activin and ectopic BMP4.

Using SEACR peak caller (Meers et al., 2019), we identified 3337 and 7639 PRDM16

CUT&TAG common peaks from four replicates of the BMP4-treated and non-treated

Prdm16_E samples (FDR < 10%, see Methods), respectively, representing PRDM16-bound

genomic sites at these two conditions (**Fig. 3A**). The samples from *Prdm16_KO* cells

completely lack these sites, validating the specificity of the PRDM16 CUT&TAG signal (**Fig.**

3B). There are 5936 lost and 1634 gained PRDM16 CUT&TAG peaks upon BMP4 treatment,

suggesting that PRDM16 regulates a distinct subset of genes in proliferating versus quiescent

NSCs.

In *Prdm16_E* cells without BMP4, we identified a few hundred to thousand ChIP-seq peaks for

all three classes of SMAD proteins after normalizing to the input reads (FDR < 10%),

suggesting a modest level of endogenous BMP and TGF-β signaling in these cells (**Fig. 3C,**

Supplementary Fig. 3). Following BMP4 addition, pSMAD1/5/8 gained around 750-fold

more peaks, while the number of SMAD3 peaks increased less than three folds (**Fig. 3D,**

Supplementary Fig. 3A). Moreover, in BMP4-treated cells pSMAD1/5/8 peaks largely overlap with SMAD4 peaks, but not with the SMAD3 ones (**Supplementary Fig. 3B**). For instance, pSMAD1/5/8 binding to two enhancers in the intronic region of *Prdm16* was increased in response to BMP4 (**Supplementary Fig. 3D**). This means that *Prdm16* itself is a transcriptional target of BMP signaling. We also noticed that *Prdm16* mRNA levels can be elevated by BMP4 in wild-type but not *Prdm16_KO* cells (**Supplementary Fig. 2B**). SMAD3 showed little ChIP-seq signal here in the absence and presence of BMP4. These results are consistent with the notion that pSMAD1/5/8 is the main receptor SMAD of BMP signaling while SMAD3 is generally irresponsive to BMP4. The gained Smad3 sites may derive from indirect effect of BMP signaling.

We then focused on assessing how PRDM16 influences genomic localization of the SMAD proteins. In overlapping analyses (see Methods), we found that more than 50% of SMAD4 and pSMAD1/5/8 binding peaks are consistent in *Prdm16_E* and *Prdm16_KO* cells (**Supplementary Fig. 4A-B**), suggesting that deletion of *Prdm16* does not affect general genomic binding ability of the SMAD4/pSMAD1/5/8 proteins. In *Prdm16_E* cells with BMP4, most of PRDM16 CUT&TAG peaks overlap with the SMAD4 and pSMAD1/5/8 peaks, but not much with SMAD3 (**Supplementary Fig. 4C**), suggesting that PRDM16 mainly collaborates with the SMAD4/pSMAD1/5/8 complex at the condition with BMP4.

Interestingly, we found that the enrichment of SMAD4 and pSMAD1/5/8 at the PRDM16 co-bound sites became significantly lower in *Prdm16_KO* mutant cells than that in *Prdm16_E* cells (**Fig. 3D**) (P=2.5E-6, and P=4.7E-3, respectively, two-tailed *t*-test), suggesting that SMAD binding to these sites depends on PRDM16. As a control, the sites without PRDM16 co-binding did not show such change (P>0.05). Thus, PRDM16 localizes SMAD proteins at specific genome regions.

We wondered whether there is any intrinsic difference between SMAD-bound regions with and without PRDM16 binding. It has been shown that SMAD4 can bind to two major types of DNA motifs (Hill, 2016): together with SMAD3, SMAD4 may bind to a palindromic motif, GTCTAGAC or direct repeats of GTCT like GTCTGTCTGTCT (Dennler et al., 1998; Wong et al., 1999; Zawel et al., 1998); together with pSMAD1/5/8, SMAD4 associates with GC-rich SBEs (SMAD-binding elements) including GGCGCC-AN4-GNCV and GRCGNCNNNNNGTCT (Gao et al., 2005; Pyrowolakis et al., 2004; Weiss et al., 2010). To determine how PRDM16 may influence enhancer selection of SMADs, we calculated

occurrence frequency of each of these motifs in five bins of SMAD4 ChIP-seq peaks from Prdm16_E cells treated with BMP4 (bin 1 to 5 with low to high peak scores, **Fig. 3E**). Interestingly, the palindromic motif is the most enriched one, and its occurrence frequency increases with peak scores. Notably, the frequency is lower in SMAD4 peaks without PRDM16 co-binding. The other three motifs show a different trend: they are not as highly enriched, their frequencies do not increase with higher SMAD4 peak scores, and the occurrence frequency in PRDM16 co-bound regions is either comparable to or even lower than the regions without PRDM16 binding. This result suggests that PRDM16 mainly tethers the SMAD4/pSMAD1/5/8 complex to the regions enriched of the palindromic motif.

Two candidate target genes co-regulated by PRDM16 and SMADs are *Wnt7b* and *Id1* (**Fig. 3F, Supplementary Fig. 4D**). We confirmed that these proteins simultaneously bind to the same DNA molecules by using sequential ChIP-qPCR with the PRDM16 antibody in the first round of IP and the SMAD4 antibody in the second IP at the *Wnt7b* and *Id1* gene loci (**Supplementary Fig. 4E**). SMAD4 displayed higher enrichment in the chromatin pulled by the PRDM16 antibody compared to that by the IgG control. We also validated reduced SMAD binding at one SMAD/PRDM16 co-bound site from each of the *Wnt7b* and *Id1* loci by ChIP-qPCR with the SMAD4 antibody in *Prdm16_E* and *Prdm16_KO* cells. SMAD4 binding became consistently weaker in the mutant in two biological replicates (**Fig. 3G, Supplementary Fig. 4F**).

Our careful inspection on the *Wnt7b* and *Id1* loci revealed that there appeared multiple ectopic SMAD4 and pSMAD1/5/8 peaks in the *Prdm16* mutant (arrow-indicated peaks in **Fig. 3F** and **Supplementary Fig. 4D**). We then globally assessed the extent of ectopic SMAD binding surrounding the SMAD/PRDM16 co-bound sites by quantifying differential SMAD binding intensity between *Prdm16_KO* and *Prdm16_expressing* cells (**Fig. 3H, Supplementary Fig. 4G**) (FDR < 10%). In agreement with the metaplots (**Fig. 3D**), all of the SMAD/PRDM16 co-bound sites showed reduced coverage intensity in *Prdm16* mutant cells. By contrast, the flanking regions within the 200kb range of a co-bound site displayed a trend of increase of SMAD binding in *Prdm16* mutant cells. This result indicates that PRDM16 helps to localize the SMAD complex at specific *cis* regulatory elements and that without *Prdm16*, the SMAD factors are partly redirected to neighboring genomic regions.

AP1 is the potential co-factor that interacts with relocated SMAD proteins

We then wondered how SMAD4 and pSMAD1/5/8 relocate to ectopic sites in the absence of
Prdm16. By running *de novo* motif discovery (see Methods), we identified a handful
significantly enriched DNA motifs from pSMAD1/5/8 and SMAD4 ChIP-seq peaks in
Prdm16_E and *Prdm16_KO* cells (**Fig. 4A-B**). Interestingly, in addition to the known SMAD
motifs, the only other motif identified in both pSMAD1/5/8 and SMAD4 peaks from
Prdm16_KO cells but not *Prdm16_expressing* cells is the AP1 motif. Furthermore, the AP1
motif is found more frequently in the pSMAD1/5/8 and SMAD4 regions lacking PRDM16
binding than those with PRDM16 binding (**Fig. 4C**). The interaction between the AP-1 complex
with the SMAD proteins was reported previously (Zhang et al., 1998). Our result implies that
AP-1 is a potential cofactor for the SMAD proteins in the absence of *Prdm16*. To validate this
finding, we performed ChIP-qPCR with an antibody against c-FOS, one of the subunits of AP1,
on *Prdm16_E* and *KO* cells. Supporting the global analyses, c-FOS is enriched in the Smad4-
bound peaks at the *Wnt7b* and *Id3* loci in the *Prdm16_KO* condition (**Fig. 4 D-G**). There is no
or only weak c-FOS binding to these regions in cells expressing *Prdm16*, suggesting that there
is cooperativity of SMAD and AP-1 facilitating each other's binding when PRDM16 is absent.
Alternatively, PRDM16 may suppress genomic accessibility of these regions for SMAD and
AP-1 proteins, a function PRDM16 has in cortical NSCs (He et al., 2021).

Genomic relocation of SMAD4/pSMAD de-represses cell proliferation genes

Next, we assessed gene regulatory activities of PRDM16 and SMAD4/pSMAD co-binding. As
a start, we examined expression change of *Wnt7b* using RT-qPCR, and found a drastic reduction
of *Wnt7b* expression in *Prdm16_E* cells in response to BMP4. This suggests that a high level
of BMP signaling represses *Wnt7b*. PRDM16 also directly represses *Wnt7b*, as the *Prdm16_KO*
sample displayed higher *Wnt7b* expression than the *Prdm16_E* sample even without BMP4
addition. Importantly, expression of *Wnt7b* is no longer suppressed by BMP4 addition in
Prdm16_KO cells, suggesting that BMP signaling synergizes with PRDM16 to repress *Wnt7b*
(**Fig. 5A**).

To extend this finding to all PRDM16 and BMP co-regulated genes in NSCs, we determined
the state of transcription activation using the H3K4me3 CUT&TAG signal in BMP4-treated
and non-treated *Prdm16_E* and *Prdm16_KO* cells. It is known that cell identity genes are
marked by extended breadth of H3K4me3 (Benayoun et al., 2014). We reasoned that compared
to RNA-seq profiling that measure gene expression change for all genes, changes in the
H3K4me3 signal at the promoter may bias toward cell identity genes between proliferation and

quiescence states. First, we noticed that the change of H3K4me3 intensity at the *Wnt7b* promoter faithfully recapitulated the change of *Wnt7b* mRNA levels shown by RT-qPCR (**Fig. 5B**). As a negative control, the TGF-β pathway target gene *Cdkn1a* encoding the cell cycle inhibitor *p21* showed unchanged H3K4me3 coverage between BMP4 treated and non-treated cells and between *Prdm16_E* and *KO* cells, suggesting that neither PRDM16 nor BMP signaling affects expression of *Cdkn1a* in NSCs. Next, we determined all BMP4-repressed and activated genes, by measuring changes of the H3K4me3 CUT&TAG read coverage at annotated transcription starting sites (TSS) in BMP4-treated *Prdm16_E* cells versus those non-treated, and identified 145 up-regulated and 184 down-regulated SMAD target genes (**Fig. 5C**) (P<0.01). To identify PRDM16-repressed and activated genes in the condition of BMP4, we called changes of H3K4me3 coverage at TSSs in *Prdm16_KO* cells treated with BMP4 versus *Prdm16_E* cells treated with BMP4. There were about twice as many up-regulated genes as down-regulated genes (**Fig. 5D**), suggesting that the cooperative activity of PRDM16 and BMP signaling is mainly to repress target gene expression.

To identify the function of genes regulated by PRDM16 and BMP signaling, we performed gene ontology (GO) analyses for genes that changed expression in response to BMP4 in *Prdm16_E* cells (see Methods). Remarkably, only downregulated genes (BMP-repressed genes) show significantly enriched functional categories, and nearly all GO terms are related to cell proliferation/cell cycle (**Fig. 5E**). A recent study reported that BMP2-induced genes are mostly enriched for neuronal and astrocyte differentiation (Katada et al., 2021), while our analysis did not identify these categories. The discrepancy may derive from difference in NSC culture, as ours was from E13 instead of E11 and E14, or different profiling methods, H3K4me3 versus RNA-seq. We overlapped these 184 genes with the 240 genes de-repressed in *Prdm16* mutant cells and identified 31 common ones. These 31 genes all display low H3K4me3 coverage in *Prdm16_E* cells with BMP4 but higher H3K4me3 signal in both *Prdm16_KO* cells treated with BMP4 and *Prdm16_E* cells without BMP4 (**Fig. 5F).**

We then intended to determine direct targets among these 31 genes that are regulated via PRDM16-bound enhancers. We previously showed that PRDM16 preferentially associates with distal intergenic regions (He et al., 2021), posing a challenge of pairing the PRDM16-bound enhancers to the targeting promoters. To overcome this issue, we took advantage of a public data browser, Embryonic Mouse Brain Epigenome Atlas (Rhodes et al., 2022), where promoter-enhancer pairs are predicted by the Monocle3 extension Cicero (Pliner et al., 2018). We

manually inspected all 31 genes for the presence of at least one PRDM16 CUT&TAG peak that is associated with the gene promoter, and found that 23 of them have proximal or distal PRDM16 binding peak(s) (**Fig. 5F,** text in back**).** Thus, these genes represent direct transcriptional targets synergistically repressed by PRDM16 and BMP signaling.

Together, our genomic data and validation suggest an enhancer switch model (**Fig. 5G)**: PRDM16 assists SMAD protein binding to repressive cis-regulatory elements that are enriched for GTCTAGAC palindromic motifs; loss of PRDM16 results in genomic relocation of SMAD proteins, presumably via the association with coactivators such as AP1; the consequence of SMAD relocation is de-repression of cell proliferation regulators.

SMADs and PRDM16 co-regulated genes in NSCs are expressed in the developing ChP

Next, we intended to examine whether the genes co-regulated by PRDM16 and BMP signaling in NSCs are involved in ChP development, an *in vivo* setting where BMP signaling (Hebert et al., 2002) and PRDM16 (**Fig. 1**) restrict NSC proliferation. To address this, we first examined expression of these genes in the developing mouse brain by taking advantage of the published single cell RNA-seq (scRNA-seq) data (La Manno et al., 2021). We selected a number of E12.5 cell clusters based on marker gene expression, including the ChP epithelial cell clusters (*Ttr* positive), CH clusters (BMP4 positive but *Ttr* negative) and a few forebrain-specific radial glia (RG) clusters (Sox2/Hes5 positive) (**Fig. 6A**). Among the 31 genes, six show no expression, and five have little expression in the examined clusters (**Supplementary Fig. 5A**). The remaining twenty genes display a moderate to high level of expression in at least one cluster. Interestingly, they are generally more lowly expressed in the ChP clusters than the CH and RG ones (**Fig. 6A**), suggesting expression of these genes may be repressed in the ChP. To identify which of these genes are directly regulated by PRDM16, we performed CUT&TAG assays using the PRDM16 antibody on dissected dorsal midline tissue that mainly contains CH and the ChP (**Fig 6B** and **Supplementary Fig. 5B**) and identified 3238 common peaks from three biological replicates. We then manually inspected the 31 gene loci for PRDM16 CUT&TAG peaks pairing with the target gene TSS using the browser Embryonic Mouse Brain Epigenome Atlas (Rhodes et al., 2022). This conformed that most of these genes have at least one PRDM16 binding peak in their promoter-linked enhancers (Text in black, **Fig. 6A** and **Supplementary Fig. 5A**).

Among these genes, *Wnt7b* is of particular interest, because the level of Wnt signaling was shown to be critical for ChP specification (Parichha et al., 2022), and Wnt signaling promotes proliferation of NSCs (Kalani et al., 2008). Although the sequencing coverage looks sparser compared to that from NSC culture, there are two PRDM16 binding peaks identified within the *Wnt7b* region (**Fig 6C**). The intronic peak is unique to the dorsal midline tissue, suggesting that this enhancer may be specifically regulated in the developing ChP. Thus, *Wnt7b* is a potential target of PRDM16 in the developing ChP.

PRDM16 represses Wnt genes in the developing ChP

We sought to determine whether PRDM16 regulates Wnt gene expression in the developing ChP. In addition to *Wnt7b*, there are several other Wnt ligands expressed in the CH and ChP region (Harrison-Uy and Pleasure, 2012). For example, *Wnt3a* is expressed in the dorsal-midline region but not in the forebrain, which explains why it was not identified from the forebrain-derived NSC datasets. Although there is no PRDM16 binding peak within the *Wnt3a* locus in the CUT&TAG data from NSCs, there is a peak in the intronic region of *Wnt3a* in the data from the dorsal midline tissue (**Supplementary Fig.5C**). This suggests that it may also be regulated by PRDM16. We thus examined the scRNA-seq data (La Manno et al., 2021) and confirmed that multiple components in the Wnt and BMP pathways are present in the CH and ChP clusters (**Supplementary Fig.5D**). To systematically measure expression changes of Wnt and BMP signaling components in the *Prdm16* mutant brain, we applied a multiplexed fluorescent *in situ* approach, Single-Cell Resolution in Situ Hybridization On Tissues (SCRINSHOT) (Sountoulidis et al., 2020). In this method each mRNA molecule is hybridized with three gene-specific padlock probes, amplified by rolling cycle amplification and then visualized by florescent dye-conjugated detection probes. A dot of fluorescent signal corresponds to one mRNA molecule. Thus, the number of dots in each cell correlates with the level of mRNA expression for each gene, allowing quantitative measurement of gene expression in single cell resolution.

We designed probes for ten Wnt and five BMP pathway components (**Fig. 7**). Additionally, we included probes for the ChP marker gene *Ttr* and *Foxj1*, the neural progenitor markers *Sox2*, *Hes5* and *Zbtb20,* as well as the neuronal marker Ngn2. To minimize technical variation, we performed SCRINSHOT in pairs of control and *Prdm16* mutant brains, with each pair being processed in the same reaction chamber throughout the hybridization procedure.

In the control brain, *Ttr* and Foxj1 are highly expressed in the ChP epithelium at E11.5 and

E12.5, and several Wnt (*Wnt2b*, *Wnt7b*, *Wnt5a*, *Wnt3a*, *Fzd10* and *Axin2*) and BMP

components (*BMP4* and *BMP7*) display certain tissue specificity: the Wnt genes are more

enriched at CH while the BMP genes are more at the ChP. A few representative images from

E11.5 samples are presented in **Fig. 7 A-B**. Except *Wnt2b*, *Wnt* genes also show relatively low

expression in the ChP cells, consistent with the notion that Wnt signaling is present and required

at the ChP (Parichha et al., 2022).

Next, we intended to quantify expression changes of these genes in *Prdm16* mutant CH and

ChP cells. As the expression of *Wnt2b* was relatively unchanged in the mutant (**Supplementary**

Fig. 1 and **Fig. 7A**), we used *Wnt2b*-expressing cells to define the CH cells and those expressing

Ttr to define ChP cells in the wild-type brain slice. There were about 110 cells in each region

of one brain slice. Because the mutant slice showed little *Ttr* signal, we only used the *Wnt2b*-

expressing cells at the most anterior side as the boarder and included 110 cells under the boarder

as the CH cells, and the further 110 cells as the mutant ChP cells, as shown in the images with

DAPI and cell outlines (**Fig. 7A and 7D**). We then measured the dot count of each gene in these

220 cells in each sample and summarized the changes calculated from three pairs of animals in

the violin plots (**Fig. 7 C** and **Supplementary Fig. 6A**). Expression of *Ttr* and *Foxj1* is severely

reduced, while *Sox2*, *Hes5* and *Ngn2* all become upregulated in the mutant ChP, indicating a

fate transformation of ChP epithelial to neural cells. This fully agrees with our immunostaining

and conventional *in situ* results (**Fig. 1** and **Supplementary Fig. 1**). Similar to BMP4, other

BMP components we tested including *BMP7*, *Smad6*, *Smad7 and Nog* are unchanged. As

Smad6 and *Smad7* are not only inhibitory factors of BMP signaling but also downstream targets,

this result suggests that BMP signaling is generally normal in the *Prdm16* mutant.

Remarkably, *Wnt3a* and *Wnt7b* are significantly upregulated in the mutant ChP (**Fig. 7B-E** and

Supplementary Fig. 6A-C). The Wnt target gene *Axin2* is also upregulated, indicating that the

activity of Wnt signaling is aberrantly elevated in the *Prdm16* mutant ChP. Thus, the normal

function of *Prdm16* represses WNT signaling in the developing ChP.

Levels of Wnt activity correlate with cell proliferation in the developing ChP and CH

As *Prdm16* mutant ChP cells showed elevated Wnt signaling (**Fig. 7 A-C**) and ectopic cell

proliferation (**Fig. 2**), we assessed the relationship between Wnt signaling and cell proliferation

in the ChP and CH at E12.5. We defined cell proliferation state using expression levels of

mKi67 by quantifying the number of dots of mKi67 SCRINSHOT signal in 110 CH and 110 ChP cells in each wild-type and mutant sample. There is a significant increase of *mKi67* expression in *Prdm16* mutant ChP cells at E12.5 (**Fig. 7F**), accompanied by significantly increased *Wnt3a*, *Wnt7b* and *Axin2* levels (**Fig. 7F** and **Supplementary Fig 6B-C**). In contrast, there is little or no increase of *mKi67* signal in the mutant CH, suggesting that the effect of PRDM16 is mainly cell-autonomous. We then performed linear regression and Pearson correlation analyses to see correlations between ChP markers, BMP and Wnt genes. Interestingly, the level of *Axin2* best correlates the level of m*Ki67*, even better than that with other Wnt genes in both wild-type ($r = 0.46$; $p < 2.2$ E-35) and mutant ($r = 0.56$; $p < 1.6$ E-55) CH and ChP cells **(Fig. 7G** and **Supplementary Fig. 6D)**. This result suggests that Wnt activity is responsible for cell proliferation in the CH and ChP region. In line with this finding, it was shown that ectopic Wnt signaling converts ChP cells into proliferative CH neural cells (Parichha et al., 2022).

Similarly, in the 4[th] ventricle, *Prdm16* deletion caused severe ChP defects (**Fig. 1D** and **Supplementary Fig. 7**). In this region from wild-type brains, *Wnt7b* showed little expression, while *Wnt3a* and *Wnt1* appeared abundant in the two sides that flanking the hChP. Interestingly, these two genes and *mKi67* all became ectopically expressed in *Prdm16* mutant ChP cells. Thus, PRDM16 also suppresses Wnt signaling in the hindbrain ChP. Taken together, our results suggest that specification of the ChP epithelium requires a process transitioning proliferating NSCs into quiescence, and that this process is mediated by a suppressive role of PRDM16 on Wnt signaling.

Discussion

Combinatory activities of morphogens are exploited throughout animal development and tissue

homeostasis. It remains to be determined how the right combination and levels of these

molecules are controlled in each specific setting. In this study we reveal that PRDM16

collaborates with BMP signaling to regulate cell quiescence, a prerequisite for ChP epithelial

cell specification. We show that BMP signaling up-regulates expression of PRDM16 whose

activity facilitates the regulatory activity of SMAD transcription factors to suppress expression

of the Wnt genes. Using SCRINSHOT that allows us to fine dissect gene expression change,

we find that expression of the genes encoding morphogens is under complex regulation, as

distinct Wnt ligand genes are differentially regulated even in the same tissue. In the case of ChP

specification, *Wnt7b* and *Wnt3a* were sensitive to *Prdm16* deletion in the forebrain, while *Wnt1*

and *Wnt3a* showed dramatic change in the hindbrain.

Wnt and BMP signaling pathways are integrated in many biological processes (Itasaki and

Hoppler, 2010). The crosstalk between BMP and WNT signaling is complex as it can be

synergistic or antagonistic. Surprisingly, both types of effects exist in the specification of ChP

epithelium. Here BMP signaling induces NSC quiescence, as evidenced by the presence of

ectopic proliferating cells at the ChP in *Bmpr1a* mutant mice (Hebert et al., 2002), while Wnt

signaling promotes proliferation since the gain-of-function condition of beta-Catenin

phenocopies *Bmpr1a* mutant animals (Parichha et al., 2022). On the other hand, adding a low

dose of Wnt activator to cell culture medium enhances programing efficiency of ChP epithelial

cells induced by BMP4 (Pellegrini et al., 2020), and loss-of-function of beta-Catenin resulted

in under-developed ChP structure (Parichha et al., 2022), suggesting that BMP and WNT

signaling collaborate to promote ChP epithelial cell differentiation. Our work demonstrated that

PRDM16 ensures the right balance of BMP and Wnt activity by maintaining a low level of Wnt

gene expression in the developing ChP.

Moreover, we showed that at the molecular level, PRDM16 functions as a co-repressor for the

SMAD4/pSMAD1/5/8 complex. PRDM16 was shown to physically interact with SMAD3 in

in vitro assays (Warner et al., 2007) and antagonize TGF-β induced cell cycle arrest (Takahata

et al., 2009). However, we tried hard to detect a stable PRDM16/SMAD4/pSMAD1/5/8

complex in co-immunoprecipitation experiments but never succeeded (data not shown). Instead,

we found that PRDM16 and SMAD4 bind to the same DNA molecule in the sequential ChIP

assay, suggesting that the cooperative activity of SMAD4 and PRDM16 occurs at their co-

bound *cis* regulatory elements. Remarkably, we identified the intrinsic difference of the DNA motifs bound by SMADs when associated with PRDM16 compared to those without PRDM16, with SMADs preferentially binding to the palindromic motif in the presence of PRDM16 and to the classic GC-rich SBE motifs in the absence of PRDM16. This provides a mechanistic insight into how SMADs switch its regulatory activity from repressing to activating gene expression. In contrast to the previous report indicating that PRDM16 antagonized the anti-proliferation activity of TGF-β and SMAD3, we show here that PRDM16 collaborates SMAD4 and pSMAD1/5/8 to suppress proliferation genes, which is consistent with the quiescence-inducing function of BMP signaling. Moreover, while it was reported that PRDM16 promotes Wnt signaling by stabilizing nuclear beta-Catenin in the craniofacial tissue (Shull et al., 2022), we detected the opposite outcome in the ChP and cultured NSCs, with ectopic WNT gene expression in the absence of *Prdm16*, suggesting that PRDM16 represses Wnt genes in these contexts. Notably, PRDM16, Wnt and BMP co-exist in various developmental settings, such as craniofacial development (Bjork et al., 2010; Nie et al., 2006; Shull et al., 2022), heart formation (Arndt et al., 2013; Gessert and Kuhl, 2010; van Wijk et al., 2007), limb patterning (Bjork et al., 2010; Geetha-Loganathan et al., 2008; Robert, 2007), adult intestinal stem cells (He et al., 2004; Korinek et al., 1998; Stine et al., 2019) *etc*. We speculate a similar regulatory circuit is used in some, if not all, of these settings.

ChPe epithelial cells derive from proliferating NSCs, without intermediate fate-commitment steps. This property endows ChP cells with higher plasticity and makes them more vulnerable to abnormal genetic and cellular change, to which high occurrence frequency of ChP tumors in fetuses and young children may be attributed (Wolff et al., 2002). As an essential brain structure, the ChP releases cerebrospinal fluid (CSF) and acts as a brain-blood barrier (Johansson, 2014; Lehtinen et al., 2013; Lun et al., 2015). Its dysfunction has been linked with several types of human diseases including hydrocephalus, Alzheimer's disease, multiple sclerosis (Lun et al., 2015). Deepening our knowledge on the development of ChP will provide new insights into potential therapeutics to prevent or treat ChP tumors and other ChP-related diseases.

Materials and methods

Animals

All animal procedures were approved by Swedish agriculture board (Jordbruks Verket) with document number Dnr 11553-2017 and 11766-2022. The *Prdm16*^{cGT} mice (Strassman et al., 2017) were maintained by outcrossing with the FVB/NJ line.

In situ hybridization:

To make probes for conventional RNA in situ hybridization, genomic regions covering one exon or full-length cDNA were PCR amplified to generate fragments with restriction enzyme overhangs. The sequences of all oligos were included in Supplementary table 1. The fragments were inserted to the pBluescript SK(II) vector. In vitro transcription was performed as previously described (He et al., 2021). The mouse brains at defined ages were dissected and fixed for 12 hours in 4% PFA, dehydrated in 25% sucrose, cryoprotected and embedded in O.C.T. The brain samples were then sectioned at 18 μm thickness on Leica cryostats CM3050s. RNA *in situ* hybridization was performed using digoxigenin-labeled riboprobes as described previously. Detailed protocols are available upon request. Images were taken using a Leica DMLB microscope.

Immunostaining

Immunostaining was done according to standard protocols as previously used (Dai et al., 2013a). For EdU labeling, EdU (5-ethynyl-2′-deoxyuridine) (5-20 μg/g of body weight) was injected into the peritoneal cavity of pregnant mice. EdU incorporation was detected with the Click-iT assay (Invitrogen) according to the manufacturer's instructions. Imaging was done on Zeiss confocal microscope. ZEN (ZeissLSM800), ImageJ (NIH) and Photoshop (Adobe) were used for analysis and quantification.

NSC culture

Control and mutant embryonic cortices from E13.5 animals were dissected and dissociated into single cell suspension and digested with Acutase (Sigma). Cells were maintained in proliferation media (STEMCELL Technologies). Lentivirus expressing Flag-Prdm16 and a puromycin resistant gene in the pCDH vector was produced in 293-ft cells and used to infect a

531   wild-type NSC line derived from E13.5 forebrain, and the selected NSCs were maintained in

532   puromycin-containing medium for three passages before puromycin withdrawal.

RT-qPCR

Prdm16_Expressing, wild-type and mutant NSCs were cultured and treated with or without

BMP4 (Sigma). After 48-hour treatment, total RNAs were extracted using TRIzol reagent

(Invitrogen). Total RNA was further cleaned with Turbo DNase (Ambion) and used in reverse-

transcription with RT master mix (ThermoFisher). To ensure the absence of genomic DNA,

control qPCR was performed on the mock-reverse-transcribed RNA samples. The list of qPCR

primers is included in supplementary table 1.

SCRINSHOT

Brain sections were prepared in the same way as for regular *in situ* hybridization (described

above). The SCRINSHOT experiments were carried out according to the published method

(Sountoulidis et al., 2020). In brief, 3 padlock probes and 3 corresponding detection probes

were designed for each gene of interest. The slides with cryo-sectioned brain slices were

pretreated at 45˚C to reduce moisture and fixed in 4% PFA in 1X PBS, followed by washing in

PBS Tween-20 0.05% twice. Permeabilization of tissues were done by washing slides in 0.1M

HCl for 2mins 15s, followed by washing in PBS Tween-20 0.05% twice. Then a stepwise

dehydration was performed for the slides in 70%, 85% & 100% Ethanol and air. The SecureSeal

hybridization chamber (GRACE BIO-LABS, 621501) was then mounted to cover each pair of

control and *Prdm16* mutant brain slices. Samples were then blocked in a probe-free

hybridization reaction mixture of 1XAmplifase buffer (Lucigen, A1905B), 0.05M KCl, 20%

Formamide deionized (Millipore S4117), 0.1uM Oligo-dT, 0.1ug/ul BSA (New England

Biolabs, B9000S), 1U/ul RiboLock (Thermo, EO0384), and 0.2ug/ul tRNAs (Ambion,

AM7119). Then hybridization of padlock probes was done by incubating samples with padlock

probes (with the concentration of each one 0.01uM) mixed in blocking reagents used before

(no oligo dT used in this step). The slide was then put into a PCR machine to denature at 55˚C

for 15 minutes and hybridize at 45˚C for 120 minutes. Padlock probes were then ligated by

using SplintR ligase (NEB M0375) at 25˚C for 16 hours followed by RCA (rolling cycle

amplification) at 30˚C for 16 hours by using phi29 polymerase (Lucigen, 30221–2) and RCA

primer1. Then a fixation step was applied to stabilize RCA product in 4% PFA for 15 minutes

followed by washing in PBS Tween-20 0.05%. Then the hybridization of the first 3 genes was

done by mixing all 3 3' fluorophore-conjugated detection probes of each gene in reaction

reagent (2XSSC, 20% Formamide deionized, 0.1ug/ul BSA, and 0.5 ng/µl DAPI (Biolegend,

422801) followed by hybridization at 30˚C for 1 hour. The slides were washed in 20%

formamide in 2X SSC and then in 6X SSC, followed by dehydration in 70%, 85%, 100%

Ethanol until the chamber was removed. Then the samples were preserved in SlowFade Gold

Antifade mountnant (Thermo, S36936) and kept in dark before imaging. After image

acquisition, the first 3 detection probes were removed by using Uracil-DNA Glycosylase

(Thermo, EN0362), and the slides were ready for the next round of detection probe

hybridization. The procedure was repeated until all genes were hybridized and imaged.

Images were acquired with Zeiss Axio Observer 7 fluorescent microscope with an automated

stage setting to fix imaging region for different hybridization rounds. Image analysis was done

according to the published method (Sountoulidis et al., 2020). In brief, the DAPI channel from

each round was extracted to measure the shift of imaging. The images were then aligned for all

gene channels in Zen by Creating Image Subset with the shifting value. After alignment, images

were exported into TIFF files and the threshold analysis was carried out for individual channel

one by one in CellProfiler with scripts provided by the published method (Alexandros et al.

2020). Then nuclear segmentation was done manually in Fiji ROI manager to obtain nuclear

ROIs, followed by an expansion of 2 um in CellProfiler to obtain cell ROIs. Then signal dots

were count in these cell ROIs for each gene in CellProfiler and Fiji. Summary of the dot counts

for each gene was exported to Excel files.

CUT&TAG

CUT&TAG was performed according to the published method (Kaya-Okur et al., 2019). In

brief, *Prdm16*_E and *Prdm16* mutant NSCs were cultured in NeuroCult™ Proliferation Kit

(Stem cell technologies, 05702) for 3 days with or without BMP4 at concentration of 25ng/ml

medium (Sigma, H4916). Cells on day 3 were then shortly rinsed with 1X PBS and resuspended

in 1ml cold NE1 buffer on ice. Nuclei were lightly fixed with 0.1% formaldehyde to 1ml PBS

at RT for 2 minutes and neutralized with 75mM Glycine. Nuclei were then washed for 3 times,

resuspended in 1mL wash buffer and mixed with 90uL Concanavalin A-coated magnetic beads

(Polyscience 86057). The nuclei/beads mix was then blocked with 800 uL cold Antibody buffer

for 5 minutes and resuspended in 1.2 mL Antibody buffer and aliquoted into 8 tubes with 150

uL each. 2 uL Anti-PRDM16 (Generous gift from Bryan Bjork lab) or IgG (SIGMA-ALDRICH,

I5006) was added into each tube for an overnight incubation at 4°C. The beads/nuclei/antibody

mix was then washed with Dig-wash buffer and incubated with secondary antibody (1:100

dilution) for 1 hour. After further washes with the Dig-wash buffer. a pA-Tn5 adaptor complex

in Dig-300 buffer was added to the beads for 1 hour reaction. The beads were then washed with

Dig-300 buffer before the incubation with 300ul Tagmentation buffer at 37°C for 1 hour. Then

10ul 0.5M EDTA, 3uL 10% SDS and 2.5uL 20 mg/mL proteinase K (Invitrogen) were used to

stop the reaction. Then the resultant DNA was purified with DNA clean & Concentrator kit

(ZYMO Research D4013) and eluted in 25uL Elution buffer. To generate libraries, 21uL

fragment DNA was mixed with 2uL 10uM Universal i5 primer, 2uL 10uM uniquely barcoded

i7 primer and 25uL PCR master mix (NEB Phusion® High-Fidelity PCR Master Mix with HF

Buffer, M0531). The PCR condition was as follows: 72°C 5min, 98°C 30s, repeat 12 times

(98°C 10s, 63°C 10s), 72°C 1 min and hold 4°C. The libraries were cleaned with standard

Ampure XP beads as previously described. Libraries from four biological replicates were

produced for each condition.

ChIP

ChIP was performed as previously described (Dai et al., 2013b). For each ChIP reaction, 10

million Prdm16_E, Prdm16_KO NSCs with or without 3-day BMP4 treatment were fixed,

lysed and made into chromatin extract. 2ug SMAD4 (Proteintech, 10231-1-AP), 2ug

pSMAD1/5/8 (Millipore, AB3848-1), 2ug SMAD3 (abcam, ab227223) and 2 ug c-FOS

antibodies (Invitrogen, MA5-15055) were used in each ChIP reaction. The same chromatin

lysate was used for sequential ChIP but with more antibodies in each reaction: 5 ug of IgG or

the PRDM16 antibody in the first round of immunoprecipitation. The elutes were then

precleared again using gamma-bind G beads, divided into two equal halves and

immunoprecipitated with 2 ug of IgG or the SMAD4 antibody. The precipitated DNA was

reverse cross-linked, and then purified using the Qiagen PCR purification kit.

Computation analyses

ChIP-seq libraries and analyses. 0.2% input and ChIPed DNA were made into libraries using

the NEBNext Ultra™ II DNA Library Prep Kit and sequenced on the Illumina Nextseq500

platform. Three replicates of ChIP-seq samples, after the adaptor trimming by Trimmomatic,

were mapped to the UCSC *Mus musculus* (mm10) genome assembly using Bowtie2 with the

default parameters. The uniquely mapped reads (with mapping quality >= 20) were used for

further analyses. The peaks were called by HOMER (v4.10) (Heinz et al., 2010). The

reproducibility between replicates was estimated by Irreproducibility Discovery Rate (IDR), using the HOMER IDR pipeline (https://github.com/karmel/homer-idr). As suggested by the Encode IDR guideline, we used a relatively relaxed parameter "-F 2 -fdr 0.3 -P .1 -L 3 -LP .1" for the true/pseudo/pooled replicates by the HOMER peak calling. The final confident peaks were determined by an IDR < 5%. The peaks that were overlapped with mm10 blacklist were also removed.

CUT&TAG analyses. The CUT&TAG samples of Prdm16_E and Prdm16 mutant NSCs in four replicates, and of Prdm16 from CH and ChP area at stage 12.5 embryos in three replicates, were mapped to mm10 genome assembly using Bowtie2 (bowtie2 --end-to-end --very-sensitive --no-mixed --no-discordant --phred33 -I 10 -X 700). The CUT&TAG coverage for these samples were generated by bedtools genomeCoverageBed and normalized to the library depths to give a read per million per base (RPM). By using the peak caller SEACR (Meers et al., 2019), which was designed for calling peaks from sparse chromatin profiling data such as CUT&TAG [22], we performed peak calling with FDR < 10% for each replicated sample (SEACR_1.3.sh normalized_coverage 0.1 non stringent output_peaks). To identify confident peaks, we selected the common peaks, which were called from all replicates for each condition. Peak overlapping analysis was performed by Homer mergePeaks function with default parameters. *De novo* motif discovery from SMAD4 and pSMAD1/5/8 peaks in Prdm16_E NSCs and Prdm16 mutant NSCs were performed by MEME-ChIP software (Machanick and Bailey, 2011) (parameters: -ccut 100 -meme-p 5 -dna -meme-mod anr -minw 5 -maxw 15 -filter-thresh 0.05).

The CUT&TAG samples of H3K4me3 BMP4-treated and non-treated Prdm16_E and Prdm16_KO cells in 2 or 3 replicates were mapped to mm10 genome assembly using Bowtie2 (bowtie2 --end-to-end --very-sensitive --no-mixed --no-discordant --phred33 -I 10 -X 700). Differential analysis of TSS up-stream and down-stream 500 bp between Prdm16_E BMP-treated vs non-BMP-treated and Prdm16_KO vs Prdm16_E BMP-treated were performed by Limma R package (Ritchie et al., 2015). Gene Ontology (GO) enrichment analysis of up-regulated and down-regulated regions was performed by PANTHER (Thomas et al., 2022).

scRNA-seq analyses. The scRNA-seq data of the developing mouse brain were obtained from (La Manno et al., 2021). The counts per cell were normalized by logNormCounts function in Bioconductor package "scater" and the normalized expression data per cell were used to generate gene expression violin plots. To test whether Prdm16 target gene sets of Prdm16 with and without BMP4 and ChP E12.5 are significantly enriched among the scRNA-seq gene mean

expression in each identified cell type cluster, the gene set enrichment analysis (GESA) "fgesa" Bioconductor package was used [ref: Korotkevich G. et al. Fast gene set enrichment analysis. bioRxiv, 2021. http://bioRxiv.org/content/early/2016/06/20/060012].

Acknowledgements

We thank the animal experimental core facility and the imaging facility of Stockholm University and Bioinformatics and Expression analysis core facility at Karolinska Institute, Sweden, for their service and support. We thank Christos Samakovlis and his lab members for technical help on SCRINSHOT experiments. We also appreciate technical help from Adrian Martinez Martin, financial support by the Australian Research Council (ARC) Future Fellowship (FT160100143) and ANU Future Scheme to J. W. The project was supported by the research project grant from Swedish Research Council (Vetenskapsrådet, 2020-03543) and the research grants from Swedish Cancerfonden (CAN 2017/529 and 20 1046 PjF 01 H) to Q.D.

Author contributions

Q.D. conceived and designed the project. L.H. performed all of the experiments. J.W performed all computational analysis. Q.D., J.W. and L.H. analysed and interpreted the data. Q.D. wrote the manuscript with input from L.H. and J.W.

Reference:

Aguilo, F., Avagyan, S., Labar, A., Sevilla, A., Lee, D.F., Kumar, P., Lemischka, I.R., Zhou, B.Y., and Snoeck, H.W. (2011). Prdm16 is a physiologic regulator of hematopoietic stem cells. Blood *117*, 5057-5066.

al, A.P.e. (2020). Canonical Wnt signaling exerts bidirectional control on choroid plexus epithelial development. BIORXIV.

Arndt, A.K., Schafer, S., Drenckhahn, J.D., Sabeh, M.K., Plovie, E.R., Caliebe, A., Klopocki, E., Musso, G., Werdich, A.A., Kalwa, H.*, et al.* (2013). Fine mapping of the 1p36 deletion syndrome identifies mutation of PRDM16 as a cause of cardiomyopathy. Am J Hum Genet *93*, 67-77.

Baizabal, J.M., Mistry, M., Garcia, M.T., Gomez, N., Olukoya, O., Tran, D., Johnson, M.B., Walsh, C.A., and Harwell, C.C. (2018). The Epigenetic State of PRDM16-Regulated Enhancers in Radial Glia Controls Cortical Neuron Position. Neuron *98*, 945-962 e948.

Benayoun, B.A., Pollina, E.A., Ucar, D., Mahmoudi, S., Karra, K., Wong, E.D., Devarajan, K., Daugherty, A.C., Kundaje, A.B., Mancini, E.*, et al.* (2014). H3K4me3 breadth is linked to cell identity and transcriptional consistency. Cell *158*, 673-688.

Bjork, B.C., Turbe-Doan, A., Prysak, M., Herron, B.J., and Beier, D.R. (2010). Prdm16 is required for normal palatogenesis in mice. Hum Mol Genet *19*, 774-789.

Chui, A., Zhang, Q., Dai, Q., and Shi, S.H. (2020). Oxidative stress regulates progenitor behavior and cortical neurogenesis. Development *147*.

Chuikov, S., Levi, B.P., Smith, M.L., and Morrison, S.J. (2010). Prdm16 promotes stem cell maintenance in multiple tissues, partly by regulating oxidative stress. Nat Cell Biol *12*, 999-1006.

Dai, Q., Andreu-Agullo, C., Insolera, R., Wong, L.C., Shi, S.H., and Lai, E.C. (2013a). BEND6 is a nuclear antagonist of Notch signaling during self-renewal of neural stem cells. Development *140*, 1892-1902.

Dai, Q., Ren, A., Westholm, J.O., Serganov, A.A., Patel, D.J., and Lai, E.C. (2013b). The BEN domain is a novel sequence-specific DNA-binding domain conserved in neural transcriptional repressors. Genes Dev *27*, 602-614.

Dennler, S., Itoh, S., Vivien, D., ten Dijke, P., Huet, S., and Gauthier, J.M. (1998). Direct binding of Smad3 and Smad4 to critical TGF beta-inducible elements in the promoter of human plasminogen activator inhibitor-type 1 gene. The EMBO journal *17*, 3091-3100.

Gao, S., Steffen, J., and Laughon, A. (2005). Dpp-responsive silencers are bound by a trimeric Mad-Medea complex. J Biol Chem *280*, 36158-36164.

Geetha-Loganathan, P., Nimmagadda, S., and Scaal, M. (2008). Wnt signaling in limb organogenesis. Organogenesis *4*, 109-115.

Gessert, S., and Kuhl, M. (2010). The multiple phases and faces of wnt signaling during cardiac differentiation and development. Circ Res *107*, 186-199.

Harrison-Uy, S.J., and Pleasure, S.J. (2012). Wnt signaling and forebrain development. Cold Spring Harb Perspect Biol *4*, a008094.

He, L., Jones, J., He, W., Bjork, B.C., Wen, J., and Dai, Q. (2021). PRDM16 regulates a temporal transcriptional program to promote progression of cortical neural progenitors. Development *148*.

He, X.C., Zhang, J., Tong, W.G., Tawfik, O., Ross, J., Scoville, D.H., Tian, Q., Zeng, X., He, X., Wiedemann, L.M.*, et al.* (2004). BMP signaling inhibits intestinal stem cell self-renewal through suppression of Wnt-beta-catenin signaling. Nat Genet *36*, 1117-1121.

Hebert, J.M., Mishina, Y., and McConnell, S.K. (2002). BMP signaling is required locally to
pattern the dorsal telencephalic midline. Neuron *35*, 1029-1041.
Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C., Singh,
H., and Glass, C.K. (2010). Simple combinations of lineage-determining transcription factors
prime cis-regulatory elements required for macrophage and B cell identities. Mol Cell *38*,
576-589.
Hill, C.S. (2016). Transcriptional Control by the SMADs. Cold Spring Harb Perspect Biol *8*.
Horvath, C., and Scheele, C. (2022). PRDM16 stability and metabolically healthy adipose
tissue. Nat Metab *4*, 1095-1096.
Inoue, M., Iwai, R., Tabata, H., Konno, D., Komabayashi-Suzuki, M., Watanabe, C., Iwanari,
H., Mochizuki, Y., Hamakubo, T., Matsuzaki, F.*, et al.* (2017). Prdm16 is crucial for
progression of the multipolar phase during neural differentiation of the developing
neocortex. Development *144*, 385-399.
Itasaki, N., and Hoppler, S. (2010). Crosstalk between Wnt and bone morphogenic protein
signaling: a turbulent relationship. Dev Dyn *239*, 16-33.
Johansson, P.A. (2014). The choroid plexuses and their impact on developmental
neurogenesis. Front Neurosci *8*, 340.
Kajimura, S., Seale, P., Tomaru, T., Erdjument-Bromage, H., Cooper, M.P., Ruas, J.L., Chin, S.,
Tempst, P., Lazar, M.A., and Spiegelman, B.M. (2008). Regulation of the brown and white fat
gene programs through a PRDM16/CtBP transcriptional complex. Genes Dev *22*, 1397-1409.
Kalani, M.Y., Cheshier, S.H., Cord, B.J., Bababeygy, S.R., Vogel, H., Weissman, I.L., Palmer,
T.D., and Nusse, R. (2008). Wnt-mediated self-renewal of neural stem/progenitor cells. Proc
Natl Acad Sci U S A *105*, 16970-16975.
Katada, S., Takouda, J., Nakagawa, T., Honda, M., Igarashi, K., Imamura, T., Ohkawa, Y., Sato,
S., Kurumizaka, H., and Nakashima, K. (2021). Neural stem/precursor cells dynamically
change their epigenetic landscape to differentially respond to BMP signaling for fate
switching during brain development. Genes Dev *35*, 1431-1444.
Kaya-Okur, H.S., Wu, S.J., Codomo, C.A., Pledger, E.S., Bryson, T.D., Henikoff, J.G., Ahmad, K.,
and Henikoff, S. (2019). CUT&Tag for efficient epigenomic profiling of small samples and
single cells. Nat Commun *10*, 1930.
Korinek, V., Barker, N., Moerer, P., van Donselaar, E., Huls, G., Peters, P.J., and Clevers, H.
(1998). Depletion of epithelial stem-cell compartments in the small intestine of mice lacking
Tcf-4. Nat Genet *19*, 379-383.
La Manno, G., Siletti, K., Furlan, A., Gyllborg, D., Vinsland, E., Mossi Albiach, A., Mattsson
Langseth, C., Khven, I., Lederer, A.R., Dratva, L.M.*, et al.* (2021). Molecular architecture of
the developing mouse brain. Nature *596*, 92-96.
Lehtinen, M.K., Bjornsson, C.S., Dymecki, S.M., Gilbertson, R.J., Holtzman, D.M., and Monuki,
E.S. (2013). The choroid plexus and cerebrospinal fluid: emerging roles in development,
disease, and therapy. J Neurosci *33*, 17553-17559.
Liddelow, S.A., Dziegielewska, K.M., Vandeberg, J.L., and Saunders, N.R. (2010).
Development of the lateral ventricular choroid plexus in a marsupial, Monodelphis
domestica. Cerebrospinal Fluid Res *7*, 16.
Lun, M.P., Monuki, E.S., and Lehtinen, M.K. (2015). Development and functions of the
choroid plexus-cerebrospinal fluid system. Nat Rev Neurosci *16*, 445-457.
Machanick, P., and Bailey, T.L. (2011). MEME-ChIP: motif analysis of large DNA datasets.
Bioinformatics *27*, 1696-1697.

Meers, M.P., Tenenbaum, D., and Henikoff, S. (2019). Peak calling by Sparse Enrichment
Analysis for CUT&RUN chromatin profiling. Epigenetics Chromatin *12*, 42.
Mira, H., Andreu, Z., Suh, H., Lie, D.C., Jessberger, S., Consiglio, A., San Emeterio, J.,
Hortiguela, R., Marques-Torrejon, M.A., Nakashima, K.*, et al.* (2010). Signaling through
BMPR-IA regulates quiescence and long-term activity of neural stem cells in the adult
hippocampus. Cell stem cell *7*, 78-89.
Nie, X., Luukko, K., and Kettunen, P. (2006). BMP signalling in craniofacial development. Int J
Dev Biol *50*, 511-521.
Parichha, A., Suresh, V., Chatterjee, M., Kshirsagar, A., Ben-Reuven, L., Olender, T., Taketo,
M.M., Radosevic, V., Bobic-Rasonja, M., Trnski, S.*, et al.* (2022). Constitutive activation of
canonical Wnt signaling disrupts choroid plexus epithelial fate. Nat Commun *13*, 633.
Pellegrini, L., Bonfio, C., Chadwick, J., Begum, F., Skehel, M., and Lancaster, M.A. (2020).
Human CNS barrier-forming organoids with cerebrospinal fluid production. Science *369*.
Pliner, H.A., Packer, J.S., McFaline-Figueroa, J.L., Cusanovich, D.A., Daza, R.M., Aghamirzaie,
D., Srivatsan, S., Qiu, X., Jackson, D., Minkina, A.*, et al.* (2018). Cicero Predicts cis-Regulatory
DNA Interactions from Single-Cell Chromatin Accessibility Data. Mol Cell *71*, 858-871 e858.
Pyrowolakis, G., Hartmann, B., Muller, B., Basler, K., and Affolter, M. (2004). A simple
molecular complex mediates widespread BMP-induced repression during Drosophila
development. Dev Cell *7*, 229-240.
Rhodes, C.T., Thompson, J.J., Mitra, A., Asokumar, D., Lee, D.R., Lee, D.J., Zhang, Y., Jason, E.,
Dale, R.K., Rocha, P.P.*, et al.* (2022). An epigenome atlas of neural progenitors within the
embryonic mouse forebrain. Nat Commun *13*, 4196.
Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015). limma
powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic
Acids Res *43*, e47.
Robert, B. (2007). Bone morphogenetic protein signaling in limb outgrowth and patterning.
Dev Growth Differ *49*, 455-468.
Shimada, I.S., Acar, M., Burgess, R.J., Zhao, Z., and Morrison, S.J. (2017). Prdm16 is required
for the maintenance of neural stem cells in the postnatal forebrain and their differentiation
into ependymal cells. Genes Dev *31*, 1134-1146.
Shull, L.C., Lencer, E.S., Kim, H.M., Goyama, S., Kurokawa, M., Costello, J.C., Jones, K., and
Artinger, K.B. (2022). PRDM paralogs antagonistically balance Wnt/beta-catenin activity
during craniofacial chondrocyte differentiation. Development *149*.
Sountoulidis, A., Liontos, A., Nguyen, H.P., Firsova, A.B., Fysikopoulos, A., Qian, X., Seeger,
W., Sundstrom, E., Nilsson, M., and Samakovlis, C. (2020). SCRINSHOT enables spatial
mapping of cell states in tissue sections with single-cell resolution. PLoS Biol *18*, e3000675.
Stine, R.R., Sakers, A.P., TeSlaa, T., Kissig, M., Stine, Z.E., Kwon, C.W., Cheng, L., Lim, H.W.,
Kaestner, K.H., Rabinowitz, J.D.*, et al.* (2019). PRDM16 Maintains Homeostasis of the
Intestinal Epithelium by Controlling Region-Specific Metabolism. Cell stem cell *25*, 830-845
e838.
Strassman, A., Schnutgen, F., Dai, Q., Jones, J.C., Gomez, A.C., Pitstick, L., Holton, N.E.,
Moskal, R., Leslie, E.R., von Melchner, H.*, et al.* (2017). Generation of a multipurpose Prdm16
mouse allele by targeted gene trapping. Dis Model Mech *10*, 909-922.
Takahata, M., Inoue, Y., Tsuda, H., Imoto, I., Koinuma, D., Hayashi, M., Ichikura, T., Yamori,
T., Nagasaki, K., Yoshida, M.*, et al.* (2009). SKI and MEL1 cooperate to inhibit transforming
growth factor-beta signal in gastric cancer cells. J Biol Chem *284*, 3334-3344.

Thomas, P.D., Ebert, D., Muruganujan, A., Mushayahama, T., Albou, L.P., and Mi, H. (2022).
PANTHER: Making genome-scale phylogenetics accessible to all. Protein Sci *31*, 8-22.
van Wijk, B., Moorman, A.F., and van den Hoff, M.J. (2007). Role of bone morphogenetic
proteins in cardiac differentiation. Cardiovasc Res *74*, 244-255.
Wang, Q., Li, H., Tajima, K., Verkerke, A.R.P., Taxin, Z.H., Hou, Z., Cole, J.B., Li, F., Wong, J.,
Abe, I.*, et al.* (2022). Post-translational control of beige fat biogenesis by PRDM16
stabilization. Nature *609*, 151-158.
Warner, D.R., Horn, K.H., Mudd, L., Webb, C.L., Greene, R.M., and Pisano, M.M. (2007).
PRDM16/MEL1: a novel Smad binding protein expressed in murine embryonic orofacial
tissue. Biochim Biophys Acta *1773*, 814-820.
Watanabe, M., Kang, Y.J., Davies, L.M., Meghpara, S., Lau, K., Chung, C.Y., Kathiriya, J.,
Hadjantonakis, A.K., and Monuki, E.S. (2012). BMP4 sufficiency to induce choroid plexus
epithelial fate from embryonic stem cell-derived neuroepithelial progenitors. J Neurosci *32*,
15934-15945.
Weiss, A., Charbonnier, E., Ellertsdottir, E., Tsirigos, A., Wolf, C., Schuh, R., Pyrowolakis, G.,
and Affolter, M. (2010). A conserved activation element in BMP signaling during Drosophila
development. Nat Struct Mol Biol *17*, 69-76.
Wolff, J.E., Sajedi, M., Brant, R., Coppes, M.J., and Egeler, R.M. (2002). Choroid plexus
tumours. British journal of cancer *87*, 1086-1091.
Wong, C., Rougier-Chapman, E.M., Frederick, J.P., Datto, M.B., Liberati, N.T., Li, J.M., and
Wang, X.F. (1999). Smad3-Smad4 and AP-1 complexes synergize in transcriptional activation
of the c-Jun promoter by transforming growth factor beta. Mol Cell Biol *19*, 1821-1830.
Zawel, L., Dai, J.L., Buckhaults, P., Zhou, S., Kinzler, K.W., Vogelstein, B., and Kern, S.E. (1998).
Human Smad3 and Smad4 are sequence-specific transcription activators. Mol Cell *1*, 611-
617.
Zhang, J., and Li, L. (2005). BMP signaling and stem cell regulation. Developmental biology
284, 1-11.
Zhang, Y., Feng, X.H., and Derynck, R. (1998). Smad3 and Smad4 cooperate with c-Jun/c-Fos
to mediate TGF-beta-induced transcription. Nature *394*, 909-913.

 Figure legend

Figure 1. Prdm16 mutant ChP cells fail to exit cell cycle and are still neural progenitors

(**A**) RNA *in situ* hybridization with a *Prdm16* probe, showing expression of *Prdm16* in the

E10.5 and E12.5 brains. The line and bracket indicate the developing ChP. (**B**) Staining of

PRDM16 using a PRDM16 antibody on the E13.5 control and null mutant brains. Red:

PRDM16; blue: the nucleus dye DAPI. White arrows indicate the ChP in the control. (**C-D**)

RNA in situ hybridization using a *Ttr* probe on E12.5 control and *Prdm16* null mutant brains.

Brain sections with tChP and hChP areas are shown separately. (**E**) Two-hour EdU labeling in

the E12.5 control and homozygous brains. The brain sections are co-stained with an antibody

against B-gal. Note: the lacZ gene trap is present in the heterozygous and homozygous mutants

but not in wild-type animals, and the highest expression of B-Gal is seen at the ChP indicated

by brackets. Red: B-Gal; blue: the nucleus dye DAPI; green: EdU. (**F**) DCX antibody staining

on the E12.5 control and homozygous brain slices. White arrows indicate some of the DCX

positive cells. Quantification of EdU+ (**E**) or DCX+ (**F**) cells is shown on the right. Biological

replicates: N=3. Error bars represent standard deviation (SD). Statistical significance is

calculated using unpaired t-test. ***, $p<0.001$.

Figure 2. Prdm16 is required for BMP4-induced NSC quiescence

(**A**) Illustration of the NSC culture assay. (**B**) Immunostaining of mKI67 in red and SOX2 in

green and EdU labeling in blue on indicated NSC genotypes and treatment. (**C**) Quantification

of the fraction of mKI67+ or EdU+ cells among the total cell number marked by SOX2. (**D**)

RT-qPCR measurement of *mKi67* RNA levels relative to 18S RNAs. Biological replicates: N=3.

Error bars represent standard deviation (SD). Statistical significance is calculated using

unpaired t-test. ***, $p<0.001$; **, $p< 001$; *, $p<0.05$; n.s., non-significant.

Figure 3. PRDM16 assists genomic binding of the SMAD proteins

(**A**) A Venn diagram showing the overlap of CUT&TAG peak in BMP4-treated and non-treated

Prdm16_E cells. (**B**) The mean coverages (normalized by library depths) centered at peak

summits are shown for the PRDM16 CUT&TAG signal in BMP4-treated and non-treated

Prdm16_E cells and Prdm16_KO mutant samples (4 replicates combined). (**C**) Heatmaps of

ChIP-seq coverage (normalized by library depths and input) centered at peak summits for

SMAD4, pSMAD1/5/8 and SMAD3 in Prdm16_E cells with or without BMP4. **(D)** Metaplots

of SMAD4 and pSMAD1/5/8 ChIP-seq coverage centered at SMAD or pSMAD peak summits

that have PRDM16 co-binding (top) and those without PRDM16 co-binding (bottom) in BMP4

treated *Prdm16_E* and *Prdm16_KO* cells. The average coverage is plotted in the bar plots on

the right. Only the peaks with PRDM16 co-binding show significant reduction in the KO cells

(*** P<0.001; ** P<0.01; * P<0.05; n.a. P>0.05). **(E)** The occurrence frequency of four types

of SMAD motifs is shown in five binned SMAD ChP-seq peaks scored from low to high. The

palindromic motif GTCTAGAC shows the highest occurrence frequency in peaks with

PRDM16 co-binding (solid red) and is the only one that has reduced occurrence frequency in

peaks without PRDM16 co-binding (dashed red). **(F)** A genome browser view of the *Wnt7b*

locus, showing PRDM16 CUT&TAG and SMAD ChIP-seq tracks. The co-bound site is

highlighted and shown in the enlarged view on the right. The arrows indicate ectopic SMAD

peaks. **(G)** ChIP-qPCR validation of SMAD4 binding to the highlighted region in *Prdm16_E*

and *Prdm16_KO* cells in two replicates. **(H)** A volcano plot showing differential SMAD4 ChIP-

seq read counts in BMP4 treated Prdm16_KO cells compared to BMP4 treated Prdm16_E cells.

The sites in blue are co-bound by PRDM16, while the sites in red are not bound by PRDM16

but within the 200kb of a PRDM16-cobound site. The average log2 fold-change is plotted on

the right.

Figure 4. AP1 is a co-factor that associates with SMADs in the absence of Prdm16

(**A-B**) *De novo* motif discovery from SMAD4 and pSMAD1/5/8 ChIP-seq peaks in *Prdm16_E*

cells (**A**) and *Prdm16_KO* cells (**B**). (**C**) The occurrence frequency of each indicated motif is

shown for SMAD4/pSMAD1/5/8 overlapped ChIP-seq peaks with or without PRDM16 co-

binding from BMP4 treated *Prdm16_E* cells. (**D, F**) Genome browser snapshots of the *Id3* (top)

and *Wnt7b* (bottom) gene loci, showing ectopic SMAD peaks in the regions, indicated by the

arrows. (**E, G**) ChIP-qPCR validation of increased SMAD4 and c-FOS occupancy at the

highlighted ectopic sites in *Prdm16_KO* mutant cells versus *Prdm16_E* cells. The locations of

PCR amplicons are highlighted in yellow in D and F.

Figure 5. The collaborative activity of PRDM16 and SMADs represses cell proliferation genes

(**A**) RT-qPCR measurement of *Wnt7b* mRNA levels in indicated genotypes and conditions from

three biological replicates, each with two technical replicates. Statistical significance was

calculated using unpaired t-test (***, p<0.001; **, p< 0.01; *, p<0.05; n.s., non-significant). (**B**)

Genome browser snapshots showing the H3K4me4 CUT&TAG tracks in the *Wnt7b* and

Cdkn1a loci. (**C**) A volcano plot of differential H3K4me3 read counts at TSS in Prdm16_E

cells with BMP4 versus those with no BMP4. Significant changes ($P < 0.01$) are colored. (**D**)

A volcano plot of differential H3K4me3 read counts at TSS in Prdm16_KO cells with BMP4

versus Prdm16_E cells with BMP4. Significant changes ($P < 0.01$) are colored. (**E**) A bubble

plot showing GO terms significantly enriched in SMAD-repressed genes (labelled in dark green

in C). (**F**) A heatmap of H3K4me3 normalized read counts at TSS of the 31 genes co-repressed

by PRDM16 and SMADs (overlapping genes from those in dark green in C and those in dark

pink in D). The genes with PRDM16 co-binding are in black text, while those without PRDM16

co-binding in orange. (**G**) An enhancer switch model demonstrating that SMADs change co-

factors and become into a transcriptional activator in the absence of PRDM16.

Figure 6. PRDM16 and SMADs co-regulated genes are differentially expressed in the ChP and

neural cells

(**A**) An illustration of the CH and ChP regions with opposite Wnt and BMP gradients and a

violin plot of scRNA-seq gene expression at E12.5 for the indicated genes in several selected

cell type clusters reanalyzed from (La Manno et al., 2021). Definition of the clusters is based

on marker gene expression from E9.5-E18.5 brain cells. These include the choroid plexus

clusters 1-5 (Chpl), with cluster Chpl5 identified by this study based on *Ttr* expression, Radial

glia clusters (RglCH 1-3, RglDF 6-10, and RglF7). CH: cortical hem; DF: dorsal forebrain; F:

forebrain. Per cell expression was normalized to give a log2 normalized counts. (**B**) Metaplots

and heatmaps of PRDM16 and IgG CUT&TAG coverage (normalized by library depths)

centered at peak summits from E12.5 dorsal midline tissues. (**C**) A genome browser snapshot

of the *Wnt7b* locus, showing PRDM16 CUT&TAG and SMADs ChIP-seq tracks.

Figure 7. PRDM16 suppresses Wnt signaling and cell proliferation in the ChP epithelial cells

(**A**) SCRINSHOT images of E11.5 wild-type and *Prdm16* mutant brains, with *BMP4*, *Ttr*,

Wnt7b and *Wnt2b* probes. Note: The red arrows indicate a strong unspecific signal in all

channels due to autofluorescence of blood cells. The cells selected for quantification are shown

together with the DAPI channel. (**B**) Enlarged views of SCRINSHOT images at the area

indicated in A, showing an increase of Wnt and neural gene expression levels and decrease of

Ttr and *Prdm16*. (**C**) A violin plot with the dot counts measuring expression levels of the 24

genes examined. Each gene is measured in 330 CH and 330 ChP cells from three wild-type and three Prdm16 mutant samples. (**D**) SCRINSHOT images for E12.5 wild-type and Prdm16 mutant brains, with Axin2, Ttr, Wnt 7b, Wnt2b and mKi67 probes. (**E**) Enlarged views of the areas indicated in the dash-lined rectangles in D. The ChP parts are highlighted by dashed lines. (**F**) Distribution of dot counts for indicated genes in the wild-type (*wt*) and *Prdm16* mutant (*mut*) CH and ChP areas. The X-axis shows the 110 cells in each replicate, and the Y-axis shows the mean value of the number of counts in three biological replicates. P value was calculated using area under the ROC curve in Prism Graphpad. (**G**) Multi-variant linear regression of *mKi67*, *Axin2* and *Ttr* using the dot counts for each gene in three pairs of wild-type and *Prdm16* mutant samples. All 660 cells including both CH and ChP tissues in each genotype are included. Pearson correlation and statistical difference were calculated in Prism Graphpad. (H) Illustration of the regulatory circuit, which is required for transitioning proliferative NSCs to quiescent NSCs, a prerequisite for the specification of ChP epithelial cells.

A

B

WT E12.5 ChP

Mean coverage (RPM)

PRDM16   IgG

Normalized cut&tag density

peak summits -2.0 to +2.0Kb

C

100 kb

Atxn10   16G11Rik   *Wnt7b*   AU022754

WT_E12.5_ChP_PRDM16 peak summit

WT_E12_5_ChP_PRDM16

Prdm16_E_BMP4_PRDM16

Prdm16_E_BMP4_pSMAD1/5/8

Prdm16_E_BMP4_SMAD4