Genome-wide screens identify SEL1L as an intracellular rheostat controlling collagen turnover

Michael J. Podolsky[1,*], Benjamin Kheyfets[1], Monika Pandey[1], Afaq H. Beigh[1], Christopher D.

Yang[2], Carlos Lizama Valenzuela[2], Ritwik Datta[2], Paul J. Wolters[3], Michael T. McManus[4], Ling

Qi[5,6], Kamran Atabai[2,3,7,*]

*Co-corresponding author

1- Department of Medicine, Weill Cornell Medical College, New York, NY, USA.

2- Cardiovascular Research Institute, University of California, San Francisco, CA, USA.

3- Department of Medicine, University of California, San Francisco, CA, USA.

4- Department of Microbiology and Immunology and UCSF Diabetes Center, University of

California, San Francisco, CA, USA.

5- Department of Internal Medicine, Division of Metabolism, Endocrinology and Diabetes,

University of Michigan Medical School, Ann Arbor, MI, USA.

6- Department of Molecular and Integrative Physiology, University of Michigan Medical School,

Ann Arbor, MI, USA.

7- Lung Biology Center, University of California, San Francisco, CA, USA.

The authors have declared that no conflict of interest exists.

Abstract

Accumulating evidence has implicated impaired extracellular matrix (ECM) clearance as a key

factor in fibrotic disease. Despite decades of research elucidating the effectors of ECM

clearance, relatively little is understood regarding the upstream regulation of this process.

Collagen is the most abundant constituent of normal and fibrotic ECM in mammalian tissues. Its

catabolism occurs through extracellular proteolysis and cell-mediated uptake of collagen

fragments for intracellular degradation. Given the paucity of information regarding the regulation

of this latter process, we executed unbiased genome-wide screens to understand the molecular

underpinnings of cell-mediated collagen clearance. Using this approach, we discovered a

previously unappreciated mechanism through which collagen biosynthesis is sensed by cells

internally and directly regulates clearance of extracellular collagen. The sensing mechanism is

dependent on endoplasmic reticulum-resident protein SEL1L and occurs via a noncanonical

function of SEL1L. This pathway functions as a homeostatic negative feedback loop that limits

collagen accumulation in tissues. In human fibrotic lung disease, the induction of this collagen

clearance pathway by collagen synthesis is impaired, thereby contributing to the pathological

accumulation of collagen in lung tissue. Thus cell-autonomous, rheostatic collagen clearance is

a previously unidentified pathway of tissue homeostasis.

Main Text

Fibrosis can be seen as the excess accumulation of extracellular matrix that interferes

with tissue and organ function. Collagen is the most abundant constituent of fibrotic extracellular

matrix, yet we still lack a complete understanding of the regulatory mechanisms governing

collagen homeostasis. Whereas decades of research have elucidated pathways of collagen

deposition in normal tissue and in fibrosis, relatively less is understood about collagen

clearance. Clearance of extracellular collagen from tissue occurs via two pathways that can act

in parallel and sequentially[1]: an extracellular proteolytic process and a process of intracellular

uptake and degradation. This latter process is vitally important in controlling normal tissue

homeostasis and mitigating fibrosis as demonstrated in several recent publications that show

the effects of interfering with this process in animal models of fibrosis[2–5]. Although an impaired

collagen-degradative environment is a feature of fibrotic diseases such as Idiopathic Pulmonary

Fibrosis (IPF), Scleroderma, and cirrhosis[6–13], very little is known about the role of cell-based

collagen degradation in human disease.

A key mediator of cell-based collagen clearance is MRC2, a canonical collagen

endocytic receptor that binds to and internalizes collagen via a Fibronectin-2 domain, with

subsequent lysosomal degradation[14,15]. Earlier studies of this receptor showed that it mitigates

lung fibrosis[3] as well as liver and kidney models of fibrosis[16,17]. Our recent work added to this

literature by showing that MRC2-mediated collagen clearance is essential for resolution of

fibrosis (rather than development of fibrosis) – deletion of MRC2 is sufficient to impair resolution

of fibrosis in the bleomycin mouse model of lung fibrosis[18].

We and others have been studying the process of cellular uptake and degradation of

collagen fragments because of the accumulating evidence that this is an important homeostatic

mechanism *in vivo* and controls fibrosis resolution[2,16–19]. Previous attempts at uncovering novel

mediators of this process have met with some success, for example uncovering the role of

Flotillin proteins or the regulation by MRC2 of matrix proteases[20,21]. However, experimental

approaches have either been targeted based on known genes and pathways or have been

limited to a subset of potential genes because of limited technical ability to use genome-wide

discovery approaches. The advent of high throughput CRISPR-based screens that are done in

a pooled format and leverage high throughput sequencing[22] now permits the unbiased

evaluation of all potential genes for any role in upstream regulatory pathways that could be

contributing to this process. Therefore in this work we started with complementary CRISPRi and

CRISPRa screens to address this knowledge gap, followed by mechanistic work to characterize

a previously undescribed pathway that controls cell-based collagen clearance.

Results

We performed genome wide unbiased screens using CRISPR interference (CRISPRi)

and CRISPR activation (CRISPRa) to determine genes that regulate cellular uptake of

exogenous collagen (Fig. 1). In independent pooled screens, we used CRISPRi to silence every

gene in the genome, or CRISPRa to activate every gene in the genome along with use of

10,000 non-targeting guides. We then performed a phenotypic screen for cellular uptake of

fluorescent collagen fragments. The screens were carried out independently with entirely

separate guide libraries for CRISPRi and CRISPRa. Coverage was at least 350x throughout the

screening process for each screen. The screens were performed in U937 cells, a human

monocytic leukemia cell line that readily takes up collagen[23] and grows in suspension, which

facilitates the use of a large-scale screen platform requiring culturing of hundreds of millions of

cells.

We determined multiple genetic regulators of collagen uptake (Fig. 1B-C; Supp. Table

1). Some of these were previously known to have a role in this process, most importantly

MRC2, a canonical collagen endocytic receptor. Most of the other candidates were not

previously known to play a role in collagen turnover. Importantly, we found a negative

correlation when comparing phenotype scores for individual genes that were hits in both the

CRISPRi and CRISPRa screens (Fig. 1D). This suggests that at least some of these genes we

identified are both necessary and sufficient for cellular collagen clearance and that quantitative

control of collagen clearance by these genes might be tunable based on their expression levels.

To validate our findings, we re-screened the high and low uptake bins from the original CRISPR

screens individually and compared hits in these re-screens to the corresponding original

CRISPRi or CRISPRa screen (Supp. Fig. 1A-D). We found a marked correlation between the

original and repeat screens (Supp. Fig. 1C-D). To validate the performance of our pooled guide

library, we used arrayed delivery of CRISPRi reagents for 3 sgRNAs for multiple genes

identified as hits in either positive or negative directions in the original CRISPRi screens,

followed by collagen uptake assays. Phenotypic effects of arrayed delivery of these CRISPRi

sgRNAs were 77% concordant with original screen data (Supp. Fig. 1E).

We next performed bioinformatic analyses of hits from our CRISPR screens. We noticed

that multiple gene ontology (GO) groups related to collagen metabolism were top scoring GO

terms in our analysis using GSEA (Fig. 2A). This was true in all four screens, i.e. the original

CRISRPi and CRISPRa screens as well as the re-screens (Supp. Fig. 2). Intriguingly, this did

not only include collagen catabolism GO terms, but also collagen biosynthesis or collagen

production-related GO terms (Fig. 2A). This informatic analysis implied that collagen

biosynthetic activity may positively regulate cellular uptake of extracellular collagen fragments.

We compared the enrichment of these GO terms in our collagen uptake screens with data from

previously published CRISPR screens of phagocytosis of other cargo[24] (Fig. 2B). Only our

screens for collagen uptake exhibited uniform enrichment of collagen biosynthetic GO terms

(i.e. positive regulation of collagen uptake by collagen biosynthesis), compared with uptake of

other cargo (Fig. 2B). These data suggest that there is a biological basis for positive regulation

collagen uptake by collagen biosynthesis genes, and this is specific to uptake of collagen.

To validate whether collagen biosynthesis positively regulates collagen uptake, we

decided to evaluate individual collagen biosynthesis genes. A heat map of gene-level phenotype

scores from our original screens among top-enriched genes from the Collagen Biosynthesis GO

group is shown in Figure 3A. We also note that *TRAM2*, a gene previously described to be

critically important collagen biosynthesis[25] was a top hit in the CRISPRi screen as well (Fig. 1).

We used an orthogonal technique (shRNA) to silence TRAM2 in U937 cells which resulted in

decreased collagen uptake (Fig. 3B; Supp. Fig. 3A) and decreased MRC2 expression at the

protein and mRNA levels (Fig. 3C-D). To verify whether TRAM2-silencing indeed affects

collagen synthesis, we used MRC5 human fetal lung fibroblasts which express high levels of

collagen. Silencing of TRAM2 in fibroblasts led to decreased endogenous levels of collagen

(Fig. 3E-G) and decreased collagen uptake in fibroblasts as well (Fig. 3H). We next silenced

individual collagen genes that were seen in the original screen including type I collagens and

type V collagens and determined that silencing of these genes led to decreased collagen uptake

and a decrease in *MRC2* message in U937 cells (Fig. 3I-K). Silencing of type I collagen

synthesis in MRC5 fibroblasts also caused decreased collagen uptake (Fig. 3L) as well as

decreased MRC2 protein levels (Fig. 3M-N). By contrast, stimulating collagen expression by

pharmacologic means with TGFβ1 increased collagen uptake in U937 cells (Fig. 4A). Although it

is controversial whether leukocytes or myeloid cells produce significant quantities of collagen[26–

28], they do generate collagen mRNA and recently were found to contribute to collagen content *in

vivo* under certain conditions (including types I and IV collagens in zebrafish and mouse)[29]; a

Western blot verified that U937 cells do indeed make collagen, albeit in very low quantities (Fig.

4A; note 60 μg of protein needed to be loaded to visualize collagen from U937 cells, compared

with 20 μg loaded in most other Western blots in this manuscript; additionally, a more sensitive

total type I collagen antibody was used in this subpanel rather than a procollagen antibody that

is less sensitive but is specific for newly synthesized collagen, which we used in the other

subpanels of this figure). To corroborate these findings, we additionally stimulated collagen

synthesis in MRC5 fibroblasts and measured collagen uptake and *MRC2* expression.

Stimulation with TGFβ1 led to increased collagen uptake and increased *MRC2* expression (Fig.

4B-C). Using orthogonal pharmacologic stimulators of collagen synthesis, ascorbic acid and

LPA1 respectively, led to increasing *MRC2* message as well (Fig. 4D-E). These data together

provide additional evidence for a pathway by which collagen biosynthesis regulates *MRC2*

expression and collagen uptake.

Because the screens were based on flow cytometric sorting of individual cells and each

cell had a single individual genetic perturbation, phenotypes determined by our screens are

expected to be fundamentally cell-autonomous. Therefore we presumed that collagen synthesis

in cells must be sensed by some internal cell-autonomous mechanism. In examining hits in the

screen to determine whether there could be a cell-autonomous sensor of collagen synthesis, we

were intrigued to see that *SEL1L* was a top hit in our screens (Fig. 1). The gene product of

SEL1L is a single-pass, endoplasmic reticulum-resident transmembrane protein with its luminal

face containing a Fibronectin-2 (FN2) domain (Fig. 5A). FN2 domains are generally known to

bind collagen[30–32]. However the function of the FN2 domain in SEL1L has not previously been

characterized.

To validate the effect of *SEL1L* on collagen uptake, we used an inducible knockout

mouse embryonic fibroblast (MEF) cell line characterized previously[33]. We determined that

when *Sel1L* is deleted, there is reduced collagen uptake (Fig. 5B), reduced MRC2 protein

expression (Fig. 5C-D; Supp. Fig. 3B-C), reduced *Mrc2* mRNA (Fig. 5E; Supp. Fig. 3D), and

reduced MRC2 cell surface and total protein expression (Fig. 5F-G). By contrast overexpressing

SEL1L in fibroblasts leads to increased *MRC2* message and protein (Fig. 5H-I). Importantly

although SEL1L has chiefly been described as a member of the endoplasmic reticulum-

associated protein degradation (ERAD) quality control pathway[33–35], other members of the

ERAD pathway were not hits in our CRISPR screens of collagen uptake (Fig. 1; Supp. Table 1).

This included HRD1, which acts with SEL1L and is essential for ERAD[36–39] but was not a hit in

our screens. To test if SEL1L-mediated regulation of collagen uptake is ERAD-dependent, we

used cells null for SEL1L or HRD1 and determined that only SEL1L positively regulates collagen

uptake, MRC2 expression (Fig. 6A-C; Supp. Fig. 3E), and *MRC2* message production (Fig. 6D).

For these experiments and several following cell biology experiments, HEK293T cells were

chosen for the following reasons (1) efficiency of transfection and co-transfection; (2) conserved

positive regulation of MRC2 by SEL1L in this cell type as well; and (3) decreased proliferation of

MEFs after two passages following induction of complete *Sel1L* knockout, as described

previously[33]. However, where possible we have analyzed more than one cell as described

below.

We then tested whether MRC2 can be forcibly overexpressed in wildtype (WT), SEL1L-

knockout and HRD1-knockout cells. We were able to overexpress in all three conditions to

equal levels MRC2 that is functional and can take up exogenous fluorescent collagen fragments

(Fig. 6E-G; Supp. Fig. 3F). Since HRD1 is essential for ERAD as above, these data suggest

that the ERAD pathway is not necessary for MRC2 protein expression or maturation and argues

against ERAD itself being instrumental in regulating MRC2 expression. Furthermore, our data

indicate that *MRC2* is being regulated by SEL1L at the message level based on the data in

Figures 5 and 6, rather than protein level as may be expected if the process were ERAD-

dependent. Finally, though deficiency of SEL1L or HRD1 have both been shown to predispose

to ER stress[40,41], the mechanism by which deficiency of either of these molecules predisposes

to ER stress has been shown to be ERAD-dependent. Since the data in Figures 5 and 6 show

that HRD1 deficiency does not affect MRC2 and collagen uptake regulation, this argues against

a strong contribution of ER stress from SEL1L deficiency affecting this pathway via ERAD

impairment because HRD1 deficiency would be expected to have the same effect on ER stress.

To determine whether SEL1L could be a collagen sensor, we wanted to investigate

whether it binds collagen and whether its FN2 domain is important for its function of regulating

MRC2 levels. Computational modeling predicted docking of collagen peptide in the FN2 domain

of SEL1L (Supp. Fig. 4A). In fibroblasts, we determined that SEL1L and collagen co-localize,

and this colocalization increases when collagen synthesis is induced by TGFβ1 (Fig. 7A; Supp.

Fig. 4B). Furthermore, there is a biochemical association between SEL1L and type I collagen,

as determined by co-immunoprecipitation followed by Western blotting (Fig. 7B). We next made

expression constructs of WT SEL1L vs. SEL1L with an in-frame deletion of its FN2 domain

(AA118-166; referred to as ΔFN2), each with C-terminal MYC tags, to examine the function of

this FN2 domain. Computational prediction suggested that this deletion should have minimal

overall effect on protein folding (Supp. Fig. 5A), since predicted full-length or ΔFN2 structures

had very predicted high alignment with Root Mean Squared Distance <1 Å. Transfection of

either construct led to localization of the MYC-tagged protein in the ER based on co-localization

with Calnexin (Supp. Fig. 5B). We also purified the WT and ΔFN2 proteins which migrated to the

expected sizes via SDS-PAGE (Supp. Fig. 5C) and analyzed the purified proteins with circular

dichroism (Supp. Fig. 5D); their spectra were similar, suggesting similar secondary structures.

Taken together, these data suggest the ΔFN2 mutant is fully translated, can be localized in the

correct subcellular compartment, and likely folds properly.

Next, we compared co-immunoprecipitation of the WT or ΔFN2 constructs with collagen

and show that the SEL1LΔFN2 mutant exhibited markedly diminished binding to procollagen

compared to WT SEL1L in lysates from co-transfected cells (Fig. 7C-D). To verify this is likely a

direct interaction, we incubated our purified protein constructs with rat-tail collagen and

demonstrate markedly reduced binding efficiency of the ΔFN2 construct, down to the

background levels of non-specific binding of collagen to beads (Supp. Fig. 5E). These data

indicate that the FN2 domain is essential for collagen binding by SEL1L. Importantly in both

fibroblasts and HEK293T cells, WT SEL1L drives MRC2 expression but SEL1LΔFN2 does not,

even though in these experiments these expression constructs are expressed at similar levels

(Fig. 7E-H). Furthermore, we evaluated the levels of OS9, a well-described ERAD substrate[33],

under conditions of SEL1L knockout or rescue with WT or ΔFN2 constructs (Fig. 7G). OS9

accumulated in the SEL1L knockout cells as expected, which was rescued with the WT

construct as well as the ΔFN2 mutant construct. Taken all together these data strongly suggest

that SEL1L is an internal sensor of collagen biosynthesis, and SEL1L is necessary for the effect

of collagen biosynthesis on MRC2 message levels and hence the downstream phenotype of

collagen uptake. Consistent with our data in Figure 6 suggesting this phenomenon is ERAD-

independent, the FN2 domain appears to be essential for *MRC2* regulation but not for ERAD

based on the effects on OS9 levels. This effect on OS9 also validates the stability and function

of the ΔFN2 construct since it rescues ERAD functionality. Consistent with our data suggesting

that the FN2 domain is dispensable for ERAD, a comparison of protein domain structures of

SEL1L across diverse species demonstrates that although SEL1L and its ERAD function are

evolutionarily ancient and conserved down to yeast[42], the FN2 domain is only present in

vertebrates and higher organisms, consistent with the hypothesis that the collagen-sensing

function arose later and is likely to be independent of its original ERAD function (Fig. 7I). This

occurred evolutionarily after the appearance of the earliest collagens in sponges at the dawn of

metazoan[43–45].

To determine whether this pathway is relevant in human physiology and disease, we re-

analyzed single cell data from multiple published experiments including normal and fibrotic lung.

We found that in cells that express both *MRC2* and type I collagen, there is a positive

correlation between expression levels of these two mRNA species (Fig. 8A-D). However, this

positive correlation is diminished IPF lung cells (Fig. 8D). In fact, data from multiple independent

diverse datasets examining gene expression in different compartments (whole lung,

bronchoalveolar lavage, peripheral blood mononuclear cells) from control versus IPF patients

demonstrate that in all cases there is a positive correlation between collagen and *MRC2*

expression, but the relationship is impaired in IPF (Fig. 8E-G). These findings corroborate our *in*

vitro data showing collagen biosynthesis is positively associated with *MRC2* and collagen

uptake by cells. This diminished recruitment of *MRC2* could be a potential mechanism by which

fibrotic collagenous extracellular matrix fails to be cleared in diseases of persistent and

progressive fibrosis like IPF. By contrast in a spontaneously resolving model of fibrosis (the

bleomycin lung fibrosis model in C57Bl\6 mice) there is a positive correlation between collagen

and *Mrc2* expression in mouse lung and the relationship is not impaired under conditions of

fibrosis (Fig. 8H). Given that we have previously shown that MRC2 is necessary for resolution of

fibrosis[18], these data raise the possibility that impairment of biosynthesis-induced collagen

clearance via MRC2 could be an important pathway driving non-resolving fibrosis and IPF.

Finally, we examined SEL1L levels in lungs of pulmonary fibrosis patients versus control

lungs (see Supp. Table 3 for more details). We found that in some, but not all, fibrotic human

lung specimens there is diminished full-length SEL1L (Fig. 9A-B). Immunofluorescence in

fibrosis vs. control lungs corroborated this finding (Fig. 9C). Future work will investigate how this

process is taking place. Additional mechanisms by which collagen biosynthesis-induced

collagen clearance is impaired in IPF is also an area of study in our laboratory.

Discussion

In this work we have shown that collagen synthesis directly upregulates collagen

clearance by MRC2. Our screens identified uniquely important biological pathways controlling

collagen clearance that do not regulate phagocytic uptake of other cargo. This led us to identify

SEL1L as a critical regulator of collagen synthesis-induced collagen turnover. SEL1L is an

internal sensor of collagen biosynthesis and its collagen sensing function is necessary for the

homeostatic effect of collagen biosynthesis on upregulation of MRC2. SEL1L is known to be a

key member of the ERAD pathway, but our data indicate that the collagen-sensing function of

SEL1L is independent of its ERAD function, defining a novel non-canonical function for this

protein. Consistent with this concept is our observation that although SEL1L and the ERAD

pathways are both evolutionarily ancient with conservation to yeast[42], the collagen-sensing FN2

domain of SEL1L arose more recently with vertebrates after the evolutionary origin of collagen

proteins, which appeared concurrently with metazoa[43–45].

From a teleological perspective, priming the same cells that contribute to scar formation

to upregulate machinery integral for repair and clearance of that scar would be an efficient

means of reaching a goal of tissue homeostasis and regeneration of normal tissue. It is striking

that in IPF, a disease of persistent non-resolving fibrosis, the relationship between collagen

synthesis and clearance is uncoupled, in contrast to the preserved relationship of these two

processes in the spontaneously resolving model of bleomycin-induced lung fibrosis in C57Bl\6

mice[18,46,47]. These findings suggest this pathway could be ripe for therapeutic intervention in

IPF. Until now, it has not been apparent that this rheostatic collagen turnover pathway was

inadequately engaged because previous studies have shown a modest increase in *MRC2*

expression in IPF lung compared with normal lung[48–52]. However, this belies a dramatically

inappropriate recruitment of the homeostatic turnover mechanism necessary to resolve fibrosis

which would be expected commensurate with high levels of collagen synthesis in IPF.

In short, an imbalance of collagen production and degradation is necessary for the

hallmark accumulation of collagen in pulmonary fibrosis; our work here suggests a defective

cell-mediated pathway of collagen clearance, based on expression of MRC2 which correlates

closely to functional uptake of collagen[23], is a key part of this imbalance. It is well documented

that MRC2 mitigates fibrosis in *in vivo* models[3,16,17]. Direct cellular clearance of collagen by

MRC2 is likely the key function responsible for this phenotype. Collagen fragments are also

thought to be pro-inflammatory[53,54], so it is possible that unclear collagen fragments in the

setting of inadequate MRC2 activity could add to fibrogenesis. However, MRC2 has multiple

other functions besides collagen uptake such as clearance of collectins[55] and roles in apoptosis

regulation and other cellular processes[56,57]. The inadequate upregulation of MRC2 we have

described in Pulmonary Fibrosis could have multiple downstream consequences which we

intend to explore further in future work. Despite this, our gain-of-function data showing that

overexpression of SEL1L can lead to increased MRC2 production suggests this pathway could

be ripe for therapeutic targeting (i.e. to drive increased clearance of collagen to ameliorate

fibrosis). The decrement in SEL1L seen in a subset of fibrotic lungs in both Western blotting and

immunofluorescence lends further credence to this idea, since MRC2 is pro-resolving in

fibrosis[18].

Limitations of our study include not elucidating the specific cell type(s) in which SEL1L is

apparently diminished in pulmonary fibrosis lungs; this is an area of future study in our

laboratory. SEL1L is expressed in multiple cell types in the lung and may have other roles (most

notably its function in ERAD) that are or are not relevant in fibrosis biology. Since collagen

expression can be detected in other cell types besides fibroblasts, it is possible the pathway we

have described here is operational in other cell types as well. We also do not yet understand

how SEL1L is downregulated in fibrotic human lungs. Our data suggest this is one potential

mechanism by which there is incomplete recruitment of the homeostatic collagen clearance

mechanisms in IPF. How SEL1L itself is downregulated in fibrotic tissue is also the subject of

ongoing study in our laboratory. In addition, the mechanism by which SEL1L-mediated collagen

sensing transcriptionally regulates *MRC2* is not completely determined by this work. The

luciferase activity data we have produced suggest that *MRC2* transcript production may be a

downstream effect of SEL1L expression levels. The precise transcription factors or regulators

that are responsible for this are also the subject of ongoing study. Lastly, we cannot exclude the

possibility that MRC2 can also be an ERAD target, even though the mechanism by which

SEL1L positively regulates *MRC2* mRNA we have shown here is not ERAD-dependent; in fact,

the trend of increased MRC2 protein levels in the HRD1-KO cells may suggest MRC2 protein

could also be an ERAD target.

Despite these limitations, we demonstrate here that unbiased genome-wide approaches

based on phenotypic screening can yield new insights into cell and tissue-level biology. The

relationship between matrix production and degradation is turning out to be ever more complex

than previously appreciated; we believe understanding how this core relationship is

dysregulated in pulmonary fibrosis at the cellular level has the potential to help unlock insights

about this relentlessly progressive disease.

Methods

CRISPR Screens

Single cell clones of U937 cells expressing dCas9-KRAB-BFP (CRISPRi) or dCas9-VP64-BFP

(CRISPRa) were made as previously described[58]. Subsequently, independent guide libraries for

CRISPRi and CRISPRa were used as previously described[59]. Briefly, lentivirus particles

containing these guides were transduced into U937 clones above at low MOI (0.3), followed by

subsequent selection with puromycin (1 μg/ml). Puromycin-selected cells were amplified and the

underwent fluorescent collagen uptake assay (as below) in bulk (i.e. pooled format), with sorting

into high and low bins (top or bottom 10% of collagen uptake based on mean fluorescence

intensity), simultaneously on 4x FACSAria II devices, then re-cultured and harvested for

analysis or for repeat screening. Isolation of cell DNA, followed by amplification of the guide-

region of individual cells and high-throughput amplicon sequencing on an Illumina platform were

all done as previously described[59,60].

Bioinformatics, Including Analysis of Screen Data

Guide counts were extracted using Python v3 scripts previously published[22]. Analysis of screen

data was done via MAGeCK[61] using the lfc-mean method. Bioinformatic analysis was performed

using GSEA[62,63] on phenotype scores (defined as Log-fold change multiplied by the $-\log_{10}$(p-

value) as determined by the robust rank-aggregation method in MAGeCK). For comparison

across knockdown screens for different cargo, data from Haney MS et al. were used[24] and then

re-analyzed via GSEA along with our own collagen screens. Visualization of screen data was

done with ggplot2 package in R (version 4.1.1). Visualization of domain architecture of SEL1L

protein across species was performed using the OMA Orthology Browser[64]. For re-analysis of

single cell RNAseq data from multiple experiments (GSE136831, GSE135893, GSE121611,

GSE128033, GSE132771) were integrated and re-normalized via the Seurat R package version

4; cells with non-zero expression of *COL1A1* and *MRC2* were used for analysis in Fig. 7D.

Microarray data was re-analyzed as indicated using GEO2R.

Quantitative real-time PCR

Total RNA from tissue or sorted cells was isolated via Trizol (Invitrogen) or RNeasy kit (Qiagen).

Total RNA from each sample was reverse transcribed with a first-strand cDNA synthesis kit

(Quantabio or Thermo Fisher) according to the manufacturer's instructions. Quantitative real-

time PCR reactions were performed with different sets of primers and Sensifast SYBR (Bioline,

Taunton, MA) on a CFX96 or CFX384 Real-Time PCR Detection System (Bio-Rad) or Applied

Biosystems QuantStudio 5 using 20 ng cDNA/reaction. Each quantitative real-time PCR

reaction was performed at least twice and representative results are shown relative to a control

by the standard $2^{-\Delta\Delta Ct}$ method, where Ct represents the number of cycles required to reach

threshold for the target gene subtracted from the number of cycles required to reach threshold

for a control housekeeping gene (*GAPDH* or *ACTB*). All data are shown relative to the first

control condition in the given figure panel. Primer sequences are in Supplemental Table 2.

Immunoblotting

Cells or tissues were homogenized with cold lysis buffer (RIPA buffer: 50 mM Tris-HCl pH 7.4,

150 mM NaCl, 0.1% SDS, 0.5% DOC, and 1% Nonidet P-40; coIP buffer: 20mM Tris-HCl pH 8,

137mM NaCl, 2mM EDTA, 1% Nonidet P-40) supplemented with protease and phosphatase

inhibitor cocktail (Thermo Fisher Scientific). Lysates (20 μg except where otherwise noted) were

clarified by centrifugation at 14,000g for 15 minutes at 4°C, electrophoresed under reducing

conditions on 7.5% or 4-20% gradient precast gels (Bio-Rad), and transferred to PVDF

membranes (Bio-Rad and GVS Filter Technology). Membranes were blocked for 30 minutes in

5% Bovine Serum Albumin and incubated with primary antibodies against MRC2 (1:1,200: anti-

mouse: AF4789, 1μg/mL: anti-human: AF5770; R&D Systems, Bio-Techne), SEL1L (1:2,000;

Novus Biologicals: NBP2-93746; ABCAM: ab78298), collagen I (1:2,000; Southern Biotech:

1310-01), human procollagen I alpha 1 (1:1,500; Novus Biologicals: AF6220), MYC (1:5,000;

Cell Signaling Technology, 9B11), OS9 (1:2,000; Novus Biologicals: NB100-519), GAPDH

(1:4,000; Cell Signaling Technology, 14C10), or beta-Actin (1:10,000; Cell Signaling

Technology, 13E5) overnight at 4°C. Membranes were then incubated with horseradish

peroxidase–conjugated secondary antibodies (1:4000: Cell Signaling Technology; 1:2,000: R&D

Systems, Bio-Techne; Santa Cruz Biotechnology) for 1 hour at room temperature. Blots were

developed using enhanced chemiluminescence (Cytiva) by exposure onto film or ChemiDoc

XRS+ (Bio-Rad). Where indicated densitometry was performed using ImageJ (NIH), normalizing

the protein of interest to a loading control (GAPDH or beta-Actin), and comparisons were made

to the baseline condition. For immunoblots of FN2-deleted SEL1L, the Abcam antibody was

used because the immunogen is distal to the FN2 domain, whereas the immunogen for the

Novus antibody is the FN2 domain itself.

Cell culture and molecular biology experiments

U937 and MRC5 cells were obtained from ATCC. HEK293T cells that were WT, or null for

SEL1L or HRD1 were obtained from Dr. Qi and were described previously[33,65–67]. Immortalized

MEFs from WT (*Sel1L^{f/f;}ERcre-*) or littermate inducible KO KO (*Sel1L^{f/f}; ERcre+*) mice were

provided by Dr. Qi and were described previously[33]. MEFs were used at passage 4-16 and for

experiments in which Sel1L deletion was induced, MEFs were used within 1 passage after

addition of 4-hydroxytamoxifen (at 400 nM) vs. ethanol vehicle control as described previously[33].

Cells were grown in DMEM or RPMI (Corning) with 10% FBS (Gemini Bio-Products). ORFs

containing full length murine SEL1L or GAPDH were obtained from Origene and were

maintained in pCMV6 expression vectors and included an in-frame C-terminal MYC tag. FN2

deletion was accomplished using Q5 Site-Directed Mutagenesis kit (New England Biolabs)

according to the manufacturer's directions and in-frame deletion was verified with Sanger

sequencing. ORFs (including the MYC tag) were also subcloned into pHIV lentiviral transfer

plasmids using standard techniques and lentiviral particles were packaged in HEK293T cells

with pMD2G and psPAX2. For MRC2 overexpression, pCI-Mrc2 (murine Mrc2 cloned into the

pCI-Neo expression vector, from Promega) was used as we previously described[18]. All

transfections were done using Lipofectamine 2000 (Invitrogen) according to the manufacturer's

instructions. For shRNA experiments, lentiviral particles harboring shRNA machinery within

pSicoR-puromycin vectors were obtained from Sigma-Aldrich (scramble control: SHC002V;

shRNA sequences can be found in Supplemental Table 2). Lentiviral infection was done in

these experiments at 10 multiplicity of infection (MOI) in cells overnight in complete medium

after addition of polybrene (8 µg/mL), and medium was replaced with fresh complete medium

the next day. The infected cells were allowed to grow and then selected by resistance to

puromycin (1 µg/mL Sigma-Aldrich). Puromycin-selected cells were collected for experiments as

indicated. Cells were infected at passages <10, and cells were collected for experiments

between passages 5 and 16. For the luciferase assay, a 625bp fragment around the MRC2

transcriptional start site in a Firefly luciferase vector with a minimal promoter element (Promega)

was used as previously described[18]. The Stop&Glo® (Promega) system was used to read

luciferase activity in cells on a luminometer (Synergy H1). All firefly luciferase values were

normalized to luciferase levels of a constitutive Renilla for a transfection control.

Co-immunoprecipitation

Proteins were isolated in non-denaturing coIP lysis buffer (see above). Proteins were incubated

with fast-flow protein G- or protein A- Dynabeads (Invitrogen) for 2 hours and centrifuged to

eliminate nonspecifically bound proteins. The concentrations of the precleared proteins were

estimated by Bradford's assay. Primary antibodies against Procollagen (Novus) or SEL1L

(Novus) or a non-specific IgG control antibody were incubated with protein G- or protein A-

Dynabeads for 4 hours. Then, 200-400 µg pre-cleared protein were incubated overnight with the

antibody-bound beads. After washing, attached proteins were then eluted from the beads in 1%

SDS buffer by heating at 60˚C and vortexing followed by centrifugation to collect the

supernatants as the eluates. Equal volumes of the eluates were used for immunoblotting.

Immunoprecipitation with protein isolation buffer without lysates was run as a negative control

for the beads. Proteins immunoprecipitated using non-specific IgG (NS IgG) were negative

control for specific antibodies. Immunoblotting with total pre-cleared protein lysates were used

as input samples. For purification of the WT and ∆FN2 proteins, MYC-tagged protein was

purified from transduced cells (see above) with Myc-trap agarose (Chromotek), a nanobody

based system, according to the manufacturer's instructions. For co-IP experiments with purified

collagen, purified WT or ∆FN2 proteins were incubated with rat tail collagen (EMD Millipore) and

then co-IP was carried out as described above.

Circular Dichroism

WT or ∆FN2 purified protein samples were diluted to final concentration of ~0.35 mg/mL in a

Tris-Glycine buffer. CD spectra were taken with Chirascan V100 CD Spectrometer (Applied

Photophysics) using a 0.5-mm quartz cuvette between 200 and 300 nm at a set temperature of

25 °C.

Flow cytometry and flow cytometric-based collagen uptake assay

Flow cytometric analysis was done using a FACSVerse, FACSAriaII, or Accuri c6 (BD

Biosciences) and sorting was done using FACSAriaII devices (BD Biosciences) at the UCSF

Parnassus Flow Cytometry Core. For the collagen uptake assay, cells were incubated in fresh

media with Oregon Green 488-conjugated gelatin (10 µg/ml final concentration; Life

Technologies) for 1 hour at 37°C. Cells were then harvested (for suspension cells) or trypsinized

and harvested for analysis. To assess only internalized fluorescent collagen (i.e. to ignore

bound but not internalized collagen), cells were either washed before flow cytometry or Trypan

blue was added to quench bound but not internalized fluorochromes as we have done

previously[21,58]. For cell-surface staining, cells were stained using a standard FACS staining

protocol. Antibodies included: anti-MRC2 (as above) as a primary antibody with a secondary of

Donkey anti-sheep conjugated to a fluorophore (Invitrogen). Cells were then analyzed and/or

sorted on one of the above devices, with sorting and analysis gates based on fluorescence-

minus-one controls after compensation was performed with single-stain controls. For the

collagen uptake assay. Data were analyzed with FlowJo, version 7.6.1 (Tree Star, Ashland,

OR).

Immunofluorescence Imaging

Cells plated on coverslips were fixed with 4% PFA in PBS. Coverslips were blocked in

Phosphate-buffered saline (PBS) with 0.5% Bovine serum albumin, 0.1% Triton X-100 and 5%

donkey serum. Antibodies against Type I collagen (Southern Biotech, as above), MRC2 (R&D,

as above), Calnexin (Cell Signaling Technology, C5C9), MYC (Santa Cruz Biotechnology,

9E10), or SEL1L (Novus, as above for Fig. 6A; for Fig. 6D, anti-SEL1L was provided as a gift by

Dr. Qi[33,65–67] and was used since it can recognize SEL1L without its FN2 domain) was applied at

1:100 dilution followed by secondary antibody conjugated to Alexa Fluor 488 or 594 (Invitrogen)

at 1:100. Coverslips were mounted in Vectashield (Vector Laboratories). Human lung fragments

were fixed in 4% PFA in PBS and then dehydrated in sucrose and embedded in OCT and

frozen in chilled isopropanol on dry ice. Frozen sections (30 µm) were prepared. Sections were

blocked in Phosphate-buffered saline (PBS) with 0.5% Bovine serum albumin, 0.1% Triton X-

100 and 5% donkey serum. Antibody against SEL1L (Novus, as above) and Collagen I

(Rockland, 600-401-103) were applied at 1:100 dilution followed by secondary antibody

conjugated to Alexa Fluor 488 or 594 or 647 (Invitrogen) at 1:100. Sections were mounted in

Vectashield (Vector Laboratories).

Confocal images were captured using a Leica TCS SPE confocal microscope with an ACS APO

40x or 63x Oil CS objective lens at room temperature with Leica Type F Immersion Liquid

(11513859) or with a Leica Stellaris 8 confocal microscope with a 10x air or 40x Oil objective.

Wide-field epifluorescence images were captured by an Olympus DP70 CCD using a standard

Olympus BX51 upright microscope with an Olympus UplanFL 40x objective at room

temperature or with a Nikon TE2000-U microscope and Ds-Qi2 camera. Images were

processed with ImageJ (National Institutes of Health, Bethesda, Maryland, USA). All imaging

was done on the same platform within any given experiment. Images were processed with

ImageJ, in which any brightness/contrast was changed uniformly across all conditions and

pseudocoloring was applied. Single optical slices at Airy 1 are shown for confocal images.

Human lung tissue

Normal human lung tissue from deceased donors of different ages was obtained from lungs not

used by the Northern California Transplant Donor Network according to an IRB-exempted

protocol led by Dr. Wolters; IPF lung specimens were obtained from explanted lungs removed

during lung transplantation, in an IRB-approved protocol led by Dr. Wolters (see Supp. Table 3

for more details). After harvest, lung tissue was directly snap-frozen in liquid nitrogen and

protein was extracted as above or 1cm fragments were fixed in 4% PFA in PBS and processed

as described above for imaging.

In Silico **protein structure prediction**

Docking of the FN2 domain of SEL1L (Uniprot Q9UBV2; AlphaFoldDB structural prediction) with

a collagen peptide sequence (PGASGPMGPRGPPGPPGKNG) was done using MDockPeP[68,69]

and visualized with PyMOL (version 2.3; Schrödinger). For prediction of the structures of full

length SEL1L the ΔFN2 mutant, the Robetta online platform (which relies on RoseTTAFold

prediction algorithm[70]) was used to predict the structure of both the full-length sequence

(Uniprot Q9Z2G6) and the sequence without AA118-166 (the FN2 domain). Alignment and

visualization were performed in PyMOL.

Statistics

Data were evaluated with Graphpad Prism software (San Diego, CA): for two sample

comparisons, by 2-tailed Student's *t*-test or Mann-Whitney *U* test if data did not pass a test of

normality; for multiple comparisons, by ANOVA with post-hoc Bonferroni testing if the ANOVA

was significant; or by Pearson correlation as indicated. A p-value of 0.05 or less was considered

statistically significant with *=$p<0.05$, **=$p<0.01$, ***=$p<0.001$, ****=$p<0.0001$, NS=Not

Significant. Data are presented as mean ± SEM.

Study approval

All experiments using human tissue were approved by the Weill Cornell Medicine or UCSF

Committee on Human Research. Informed consent was obtained for study participants or

waived for human tissue from deceased donors in accordance with the UCSF Committee on

Human Research.

Data Availability

Source data are provided with this paper.

Code Availability

541 The codes used in CRISPR screen analysis are described above and are available from the

542 authors upon reasonable requests.

Acknowledgments

This research was supported by US National Institutes of Health grants K08 HL145015 (MJP)

and R01 HL136377 (KA), a Grant-in-aid from The Stony Wold-Herbert Fund (MJP), and a

Sponsored Research Agreement with Celgene (KA). We acknowledge the UCSF Parnassus

Flow CoLab (RRID:SCR_018206) for assistant generating flow cytometry data, supported in

part by Grant NIH P30 DK063720 and by the NIH S10 Instrumentation Grant S10

1S10OD021822-01. We acknowledge and thank the Precision Biomolecular Characterization

Facility at Columbia University for technical support and access to the Circular Dichroism

spectrometer, which is supported by NIH Award 1S10OD025102-01. We would like to thank S.

Layer for ongoing inspiration.

Figure Legends

Figure 1. CRISPR screens identify mediators of cellular collagen uptake. (A) Schematic of

experimental approach for flow cytometry-based screening for uptake of collagen fragments. (B)

Volcano plot of gene-level statistics of CRISPRi (inhibition) screen. Most significant gene hits

are labeled and colored in red. RRA = Robust Rank Aggregation score; LFC = Log-fold change.

(C) Volcano plot of gene-level statistics of CRISPRa (activation) screen. (D) Comparison of

overall phenotype score between CRISPRi and CRISPRa screens.

Figure 2. Collagen-biosynthetic pathways are regulators of collagen uptake uniquely compared

to screens for uptake of other cargoes. (A) Ranked plot of Normalized Enrichment Scores

derived from Gene-Set Enrichment Analysis (GSEA) of CRISPRi phenotype scores, using

10,561 Gene Ontology gene sets. Collagen-related gene sets are labeled and displayed in red.

(B) Comparison of GSEA statistics for labeled collagen-related gene sets across screens for

uptake of various cargoes[24]; screen method is indicated in parentheses. KD = Knock-down.

Figure 3. Collagen biosynthesis positively regulates cell-based collagen clearance and *MRC2*

expression. (A) Heatmap of overall phenotype scores from original screens as indicated for top

genes from 'Collagen Biosynthetic Process' GO gene-set (displayed genes had most extreme

'Score' values in one or more screen result). (B) Flow-cytometry based collagen-uptake assay in

U937 cells after shRNA treatment as indicated vs. Scramble control. MFI = Mean Fluorescence

Intensity. N=4 per group. (C) Western blot of U937 cells after shRNA treatment as indicated vs.

Scramble control, representative of N=4 independent experiments. (D) Q-RT-PCR in U937 cells

after shRNA treatment as indicated vs. Scramble control. N=4 per group. (E-F) Western blot

and densitometry of MRC5 fibroblasts after shRNA treatment as indicated vs. Scramble control,

representative of N=3 independent experiments. (G) Representative confocal

immunofluorescence images of MRC5 fibroblasts after shRNA treatment as indicated vs.

Scramble control. (H) Flow-cytometry based collagen-uptake assay in MRC5 cells after shRNA

treatment as indicated vs. Scramble control. N=4 per group. (I) Flow-cytometry based collagen-

uptake assay in U937 cells after shRNA treatment as indicated vs. Scramble control. N=4-6 per

group. (J-K) Q-RT-PCR in U937 cells after shRNA treatment as indicated vs. Scramble control.

N=4-5 per group. (L) Flow-cytometry based collagen-uptake assay in MRC5 fibroblasts after

shRNA treatment as indicated vs. Scramble control. N=6 and 4 in Scramble and shCOL1A1

groups respectively, N=2 in shMRC2 group. (M-N) Western blot and densitometry of MRC5

fibroblasts after shRNA treatment as indicated vs. Scramble control, representative of N=4

independent experiments. Statistics: (B-K) ANOVA with post-hoc Bonferroni testing; (L-N)

Student's *t*-test. * $p < 0.05$, ** $p < 0.01$, **** $p < 0.0001$.

Figure 4. Pharmacological stimulation of collagen synthesis is positively associated with

collagen uptake and *MRC2* expression. (A-B) Flow-cytometry based collagen-uptake assay in

U937 (A) or MRC5 (B) cells after drug treatment as indicated vs. vehicle control. N=4 per group.

Western blot of corresponding cells and conditions shown below, each representative of N=3-4

independent experiments. (C-E) Q-RT-PCR in MRC5 cells after drug treatment as indicated vs.

vehicle control. N=3-5 per group. Western blot of corresponding cells and conditions shown

below, each representative of N=3 independent experiments. Statistics: (A-E) Student's *t*-test. *

$p < 0.05$, ** $p < 0.01$.

Figure 5. *Sel1L* positively regulates collagen uptake and *Mrc2* expression. (A) Schematic of

domain structure of SEL1L protein. FN2 = Fibronectin-2. (B) Flow-cytometry based collagen-

uptake assay in MEF cells (mouse embryonic fibroblasts) after treatment with 4-

hydroxytamoxifen (OHT) as indicated vs. vehicle control. WT = *Sel1L^{f/f;}ERcre*-, iKO =

Sel1L^{f/f};ERcre+. N=3-4 per group. (C-D) Western blot and densitometry of MEFs after OHT

treatment as indicated vs. control, representative of N=4 independent experiments. (E) Q-RT-

PCR in MEFs after OHT treatment for genes as indicated. N=3 per group. (F) Flow-cytometry

based measurement of MRC2 surface expression in MEFs after OHT treatment. N=4 per group.

(G) Representative wide-field immunofluorescence images of MEFs after OHT treatment. (H) Q-

RT-PCR of MRC5 cells transfected with overexpression vectors containing genes as indicated

vs. vehicle control (lipofectamine only). N=4 per group. (I) Western blot of MRC5 cells

transfected with overexpression vectors containing genes as indicated vs. vehicle control

(lipofectamine only). N=2. Statistics: (B, H) ANOVA with post-hoc Bonferroni testing; (D - right

panel, E, F) Student's *t*-test; (D - left panel) Mann-Whitney *U* test. * $p < 0.05$, ** $p < 0.01$, *** $p <$

0.001, **** $p < 0.0001$, ns = not significant.

Figure 6. SEL1L, but not HRD1, positively regulates MRC2. (A) Flow-cytometry based collagen-

uptake assay in 293T of different genotypes (compared with baseline control without fluorescent

collagen). N=4-5 per group. (B-C) Western blot and densitometry of 293T cells of different

genotypes, representative of N=5 independent experiments. (D) Firefly luciferase assay of

MRC2 promoter activity, normalized to constitutive Renilla luciferase in 293T cells of different

genotypes. N=3 per group. (E) Western blot of HEK293T cells of different genotypes transfected

with MRC2 overexpression vector vs. vehicle control, representative of N=6 independent

experiments. (F) Representative wide-field immunofluorescence images of 293T cells of

different genotypes transfected with MRC2 overexpression vector vs. vehicle control. (G) Flow-

cytometry based collagen-uptake assay in HEK293T of different genotypes transfected with

MRC2 overexpression vector (compared with baseline control without fluorescent collagen; note

different scale compared with panel A). N=3-4 per group. Statistics: (A, C, D, G) ANOVA with

post-hoc Bonferroni testing. * $p < 0.05$ ** $p < 0.01$, ns = not significant.

Figure 7. SEL1L binds collagen as it is being synthesized via its FN2 domain and this interaction

is necessary for regulation of *MRC2* expression. (A) Representative confocal

immunofluorescence images of MRC5 fibroblasts treated as indicated or vehicle control. (B)

Western blot of MRC5 fibroblasts after immunoprecipitation with anti-SEL1L antibody (vs.

controls as indicated), representative of N=2 independent experiments. (C) Western blot after

immunoprecipitation with anti-procollagen type I antibody (with input controls as indicated) of

HEK293T cells transfected with collagen and WT or mutant SEL1L constructs as indicated,

representative of N=3 independent experiments. (D) Representative confocal

immunofluorescence images of HEK293T cells transfected with collagen and WT or mutant

SEL1L constructs as indicated. (E) Western blot of MEFs after treatment with WT or mutant

SEL1L constructs as indicated, representative of N=3 independent experiments. (F) Flow-

cytometry based measurement of MRC2 surface expression on MEFs after treatment with WT

or mutant SEL1L constructs as indicated, vs. Empty Vector (EV). N=6 per group. (G) Western

blot of HEK293T cells either WT or KO for SEL1L as indicated after treatment with EV (empty

vector), WT or mutant SEL1L constructs as indicated, representative of N=3 independent

experiments. (H) Q-RT-PCR of SEL1L-KO HEK293T cells after transfection with WT or mutant

SEL1L as indicated vs. empty vehicle control. N=3 per group. (I) Comparison of protein domain

structures of Sel1L orthologs across phylogenetically diverse organisms; FN2 = Fibronectin-2

domain, tpr = Tetratricopeptide repeat domain. Statistics: (F-G) ANOVA with post-hoc

Bonferroni testing. * $p < 0.05$, ** $p < 0.01$, ns = not significant.

Figure 8. Collagen biosynthesis and *MRC2* expression are positively correlated in lung tissue,

but this relationship is impaired in human fibrotic lung disease. (A) UMAP plot with cell type

annotation of scRNAseq data from multiple available datasets of human lung control and IPF

tissue (GSE136831, GSE135893, GSE121611, GSE128033, GSE132771). N=184,672 single

cells. (B-C) Gene expression of individual genes as indicated in UMAP as generated in panel

(A). (D) Comparison of gene expression in all cells from data as in panel (A) that had non-zero

expression of *COL1A1* or *MRC2*. (E) Comparison of gene expression in whole human lung

microarray data from GSE110147 analyzed via GEO2R in pulmonary fibrosis lungs vs. control

specimens. N=37 vs. 11 per group. (F) Comparison of microarray gene expression in human

lung bronchoalveolar lavage specimens from GSE70867 analyzed via GEO2R from Idiopathic

Pulmonary Fibrosis (IPF) lungs vs. control specimens. N=112 vs. 20. (G) Comparison of

microarray gene expression in peripheral blood mononuclear cell (PBMC) specimens from

GSE37858 analyzed via GEO2R from Idiopathic Pulmonary Fibrosis (IPF) lungs vs. control

specimens. N=70 vs. 35. (H) Comparison of gene expression in whole mouse lung microarray

data from GSE40151 analyzed via GEO2R from Bleomycin-treated mice vs. Saline control

mice. N=56 vs. 55. Statistics: ANCOVA, p-values indicated in figures. Pearson correlations are

displayed in panel (D).

Figure 9. SEL1L protein levels are reduced in the lungs of patients with Pulmonary Fibrosis. (A-

B) Western blots of whole human lung lysates from Pulmonary Fibrosis lungs vs. control (donor

lungs deemed not suitable for transplant), with corresponding densitometry. N=13 in Control

and 11 in Pulmonary Fibrosis group. Arrowheads correspond to two lung lysates that were used

in both Western blots for normalization across blots, otherwise all other lanes are independent

samples. (C) Representative low-power wide-field immunofluorescence images of human lungs

(Pulmonary Fibrosis vs. control). Each row comes from an individual donor or Pulmonary

Fibrosis lung, with two different fields per lung (i.e. N=3 individual lungs per group). Statistics:

(B, D) Mann-Whitney U test. * p < 0.05.

Supplemental Figures

Supplemental Figure 1. Re-screens are correlated to original CRISPR screen results. (A)

Volcano plot of CRISPRi re-screen (Screen 2). (B) Volcano plot of CRISPRa re-screen (Screen

2). (C) Comparison of gene-level overall phenotype scores between CRISPRi screens 1 and 2.

(D) Comparison of gene-level overall phenotype scores between CRISPRa screens 1 and 2. (E)

Comparison of arrayed log-fold change in arrayed validation vs. original pooled screen log-fold

change; each point represents unique guide sequence. 65/84 guides demonstrated

concordance with original pooled screen data (77%). Pearson correlation coefficient is shown.

Supplemental Figure 2. Collagen-related gene ontology terms are enriched as regulators of

collagen uptake in CRISPRi and CRISPRa screens. The top twenty positive and negatively

performing gene sets are indicated with associated statistics for each screen as indicated,

among all 10,561 Gene Ontology gene sets. Collagen-related gene sets are highlighted with red

boxes (i.e. those with 'Collagen' in the GO gene set name). NES = Normalized Enrichment

Score.

Supplemental Figure 3. Validation of different tools used in this manuscript. (A) Q-RT-PCR in

U937 cells after shRNA treatment as indicated vs. Scramble control. N=4 per group. (B-C)

Additional densitometry of data from Figure 5C. N=4 per group. (D) Q-RT-PCR in MEFs after

OHT treatment for genes as indicated. N=3 per group. (E) Additional densitometry of data from

Figure 6B. N=4 per group. (F) Densitometry of the left side of Figure 6E. N=6 per group.

Statistics: (A, E, F) ANOVA with post-hoc Bonferroni testing; (B-D) Student's *t*-test. * p < 0.05, **

p < 0.01, **** p < 0.0001, ns = not significant.

Supplemental Figure 4. Additional support of SEL1L and collagen interaction. (A) Predicted

docking of collagen peptide with FN2 domain of SEL1L. (B) Additional representative confocal

immunofluorescence image of MRC5 fibroblasts at baseline (as in Figure 7A), with decreased

PMT gain in the Collagen channel, along with magnified inset to highlight co-localization.

Supplemental Figure 5. FN2 domain deletion mutant interferes with collagen binding but is

otherwise stable. (A) Alignment of predicted structures of full-length SEL1L and ΔFN2 SEL1L.

Two different views and alignments are shown. Root-mean squared distance (RMSD) is

displayed. Yellow arrow indicates the FN2 domain (in the pink structure on the left; in the green

structure on the right). (B) Representative confocal immunofluorescence images of HEK293T

cells transfected with MYC-tagged WT or ΔFN2 SEL1L constructs (at low transfection

efficiency), co-stained with ER marker Calnexin. (C) Western blot of purified WT or ΔFN2 SEL1L

proteins compared with input cell lysates, representative of N=3 independent experiments. (D)

Circular Dichroism spectra of purified WT or ΔFN2 proteins. (E) Western blot of co-incubated

purified either WT or ΔFN2 SEL1L proteins with rat tail collagen, after immunoprecipitation with

anti-MYC antibody (with input and additional controls as indicated), representative of N=3

independent experiments. 'Collagen only' indicates incubation of rat tail collagen with IP beads

without co-incubation with purified MYC-tagged protein.

Supplemental Tables

Supplemental Table 1. Gene-level data from CRISPRi and CRISPRa screens of collagen

uptake as analyzed via MAGeCK.

Supplemental Table 2. Primer and shRNA sequences used in this manuscript.

Supplemental Table 3. Demographic and clinical data on the sources of human lungs used in

this manuscript.

References

1.   McKleroy W, Lee T-H, Atabai K. Always cleave up your mess: targeting collagen
degradation to treat tissue fibrosis. *AJP: Lung Cellular and Molecular Physiology*.
2013;304(11):L709–L721.

2.   Atabai K, Jame S, Azhar N, Kuo A, Lam M, McKleroy W, DeHart G, Rahman S, Xia DD,
Melton AC, Wolters P, Emson CL, Turner SM, Werb Z, Sheppard D. Mfge8 diminishes the
severity of tissue fibrosis in mice by binding and targeting collagen for uptake by macrophages.
Journal of Clinical Investigation. 2009.
3.   Bundesmann MM, Wagner TE, Chow Y-H, Altemeier WA, Steinbach T, Schnapp LM.
Role of urokinase plasminogen activator receptor-associated protein in mouse lung. *American
journal of respiratory cell and molecular biology*. 2012;46(2):233–9.
4.   Cui H, Jiang D, Banerjee S, Xie N, Kulkarni T, Liu R-M, Duncan SR, Liu G. Monocyte-
derived alveolar macrophage apolipoprotein E participates in pulmonary fibrosis resolution. *JCI
Insight*. 2020;5(5).
5.   Fan M-H, Zhu Q, Li H-H, Ra H-J, Majumdar S, Gulick DL, Jerome JA, Madsen DH,
Christofidou-Solomidou M, Speicher DW, Bachovchin WW, Feghali-Bostwick C, Pur Eacute E.
Fibroblast Activation Protein (FAP) Accelerates Collagen Degradation and Clearance from Lung
in Mice. *The Journal of biological chemistry*. 2015.
6.   Montaño M, Ramos C, González G, Vadillo F, Pardo A, Selman M. Lung collagenase
inhibitors and spontaneous and latent collagenase activity in idiopathic pulmonary fibrosis and
hypersensitivity pneumonitis. *Chest*. 1989;96(5):1115–9.
7.   Selman M, Ruiz V, Cabrera S, Segura L, Ramírez R, Barrios R, Pardo A. TIMP-1, -2, -3,
and -4 in idiopathic pulmonary fibrosis. A prevailing nondegradative lung microenvironment?
American journal of physiology. Lung cellular and molecular physiology. 2000;279(3):L562-74.
8.   Montfort I, Pérez-Tamayo R. Collagenase in experimental carbon tetrachloride cirrhosis
of the liver. *The American journal of pathology*. 1978;92(2):411–20.
9.   Okazaki I, Maruyama K. Collagenase activity in experimental hepatic fibrosis. *Nature*.
1974;252(5478):49–50.
10.   Brady AH. Collagenase in scleroderma. *The Journal of clinical investigation*.
1975;56(5):1175–80.
11.   Mahalanobish S, Saha S, Dutta S, Sil PC. Matrix metalloproteinase: An upcoming
therapeutic approach for idiopathic pulmonary fibrosis. *Pharmacological Research*.
2020;152:104591.
12.   Costanzo L, Soto B, Meier R, Geraghty P. The Biology and Function of Tissue Inhibitor
of Metalloproteinase 2 in the Lungs. *Pulmonary Medicine*. 2022;2022:1–12.
13.   Inoue R, Yasuma T, Fridman D'Alessandro V, Toda M, Ito T, Tomaru A, D'Alessandro-
Gabazza CN, Tsuruga T, Okano T, Takeshita A, Nishihama K, Fujimoto H, Kobayashi T,
Gabazza EC. Amelioration of Pulmonary Fibrosis by Matrix Metalloproteinase-2
Overexpression. *International Journal of Molecular Sciences*. 2023;24(7):6695.
14.   Engelholm LH, List K, Netzel-Arnett S, Cukierman E, Mitola DJ, Aaronson H, Kjøller L,
Larsen JK, Yamada KM, Strickland DK, Holmbeck K, Danø K, Birkedal-Hansen H, Behrendt N,
Bugge TH. uPARAP/Endo180 is essential for cellular uptake of collagen and promotes fibroblast
collagen adhesion. *The Journal of cell biology*. 2003;160(7):1009–15.
15.   Yuan C, Jürgensen HJ, Engelholm LH, Li R, Liu M, Jiang L, Luo Z, Behrendt N, Huang
M. Crystal structures of the ligand-binding region of uPARAP: effect of calcium ion binding. *The
Biochemical journal*. 2016;473(15):2359–68.
16.   Madsen DH, Jürgensen HJ, Ingvarsen S, Melander MC, Vainer B, Egerod KL, Hald A,
Rønø B, Madsen CA, Bugge TH, Engelholm LH, Behrendt N. Endocytic collagen degradation: A
novel mechanism involved in protection against liver fibrosis. *Journal of Pathology*.
2012;227(1):94–105.
17.   López-Guisa JM, Cai X, Collins SJ, Yamaguchi I, Okamura DM, Bugge TH, Isacke CM,
Emson CL, Turner SM, Shankland SJ, Eddy AA. Mannose receptor 2 attenuates renal fibrosis.
Journal of the American Society of Nephrology : JASN. 2012;23(2):236–51.

18.   Podolsky MJ, Yang CD, Valenzuela CL, Datta R, Huang SK, Nishimura SL, Dallas SL,
Wolters PJ, Le Saux CJ, Atabai K. Age-dependent regulation of cell-mediated collagen turnover.
JCI Insight. 2020;5(10).
19.   Melander MC, Jürgensen HJ, Madsen DH, Engelholm LH, Behrendt N. The collagen
receptor uPARAP/Endo180 in tissue degradation and cancer (Review). *International Journal of*
Oncology. 2015;47(4):1177–1188.
20.   Messaritou G, East L, Roghi C, Isacke CM, Yarwood H. Membrane type-1 matrix
metalloproteinase activity is regulated by the endocytic collagen receptor Endo180. *Journal of*
cell science. 2009;122(Pt 22):4042–4048.
21.   Lee T-H, McKleroy W, Khalifeh-Soltani A, Sakuma S, Lazarev S, Riento K, Nishimura
SL, Nichols BJ, Atabai K. Functional genomic screen identifies novel mediators of collagen
uptake. *Molecular biology of the cell*. 2014;25(5):583–93.
22.   Gilbert LA, Horlbeck MA, Adamson B, Villalta JE, Chen Y, Whitehead EH, Guimaraes C,
Panning B, Ploegh HL, Bassik MC, Qi LS, Kampmann M, Weissman JS. Genome-Scale
CRISPR-Mediated Control of Gene Repression and Activation. *Cell*. 2014;159(3):647–661.
23.   Madsen DH, Ingvarsen S, Jürgensen HJ, Melander MC, Kjøller L, Moyer A, Honoré C,
Madsen CA, Garred P, Burgdorf S, Bugge TH, Behrendt N, Engelholm LH. The non-phagocytic
route of collagen uptake: A distinct degradation pathway. *Journal of Biological Chemistry*. 2011.
24.   Haney MS, Bohlen CJ, Morgens DW, Ousey JA, Barkal AA, Tsui CK, Ego BK, Levin R,
Kamber RA, Collins H, Tucker A, Li A, Vorselen D, Labitigan L, Crane E, et al. Identification of
phagocytosis regulators using magnetic genome-wide CRISPR screens. *Nature genetics*.
2018;50(12):1716–1727.
25.   Stefanovic B, Stefanovic L, Schnabl B, Bataller R, Brenner DA. TRAM2 protein interacts
with endoplasmic reticulum Ca2+ pump Serca2b and is necessary for collagen type I synthesis.
Molecular and cellular biology. 2004;24(4):1758–68.
26.   Bucala R, Spiegel LA, Chesney J, Hogan M, Cerami A. Circulating fibrocytes define a
new leukocyte subpopulation that mediates tissue repair. *Molecular medicine (Cambridge,*
Mass.). 1994;1(1):71–81.
27.   Phillips RJ, Burdick MD, Hong K, Lutz MA, Murray LA, Xue YY, Belperio JA, Keane MP,
Strieter RM. Circulating fibrocytes traffic to the lungs in response to CXCL12 and mediate
fibrosis. *The Journal of clinical investigation*. 2004;114(3):438–46.
28.   Kleaveland KR, Velikoff M, Yang J, Agarwal M, Rippe RA, Moore BB, Kim KK.
Fibrocytes are not an essential source of type I collagen during lung fibrosis. *Journal of*
immunology (Baltimore, Md. : 1950). 2014;193(10):5229–39.
29.   Simões FC, Cahill TJ, Kenyon A, Gavriouchkina D, Vieira JM, Sun X, Pezzolla D,
Ravaud C, Masmanian E, Weinberger M, Mayes S, Lemieux ME, Barnette DN, Gunadasa-
Rohling M, Williams RM, et al. Macrophages directly contribute collagen to scar formation
during zebrafish heart regeneration and mouse heart repair. *Nature Communications*.
2020;11(1):600.
30.   Ozhogina OA, Trexler M, Bányai L, Llinás M, Patthy L. Origin of fibronectin type II (FN2)
modules: structural analyses of distantly-related members of the kringle family idey the kringle
domain of neurotrypsin as a potential link between FN2 domains and kringles. *Protein science :*
a publication of the Protein Society. 2001;10(10):2114–22.
31.   Bányai L, Patthy L. Evidence for the involvement of type II domains in collagen binding
by 72 kDa type IV procollagenase. *FEBS letters*. 1991;282(1):23–5.
32.   Pickford AR, Potts JR, Bright JR, Phan I, Campbell ID. Solution structure of a type 2
module from fibronectin: implications for the structure and function of the gelatin-binding
domain. *Structure*. 1997;5(3):359–370.
33.   Sun S, Shi G, Han X, Francisco AB, Ji Y, Mendonça N, Liu X, Locasale JW, Simpson
KW, Duhamel GE, Kersten S, Yates JR, Long Q, Qi L. Sel1L is indispensable for mammalian

endoplasmic reticulum-associated degradation, endoplasmic reticulum homeostasis, and
survival. *Proceedings of the National Academy of Sciences*. 2014;111(5).
34.   Christianson JC, Olzmann JA, Shaler TA, Sowa ME, Bennett EJ, Richter CM, Tyler RE,
Greenblatt EJ, Wade Harper J, Kopito RR. Defining human ERAD networks through an
integrative mapping strategy. *Nature Cell Biology*. 2012;14(1):93–105.
35.   Mueller B, Lilley BN, Ploegh HL. SEL1L, the homologue of yeast Hrd3p, is involved in
protein dislocation from the mammalian ER. *The Journal of cell biology*. 2006;175(2):261–70.
36.   Gardner RG, Swarbrick GM, Bays NW, Cronin SR, Wilhovsky S, Seelig L, Kim C,
Hampton RY. Endoplasmic Reticulum Degradation Requires Lumen to Cytosol Signaling.
Journal of Cell Biology. 2000;151(1):69–82.
37.   Hampton RY, Gardner RG, Rine J. Role of 26S proteasome and HRD genes in the
degradation of 3-hydroxy-3-methylglutaryl-CoA reductase, an integral endoplasmic reticulum
membrane protein. *Molecular Biology of the Cell*. 1996;7(12):2029–2044.
38.   Bordallo J, Plemper RK, Finger A, Wolf DH. Der3p/Hrd1p Is Required for Endoplasmic
Reticulum-associated Degradation of Misfolded Lumenal and Integral Membrane Proteins.
Molecular Biology of the Cell. 1998;9(1):209–222.
39.   Plemper RK, Bordallo J, Deak PM, Taxis C, Hitt R, Wolf DH. Genetic interactions of
Hrd3p and Der3p/Hrd1p with Sec61p suggest a retro-translocation complex mediating protein
transport for ER degradation. *Journal of Cell Science*. 1999;112(22):4123–4134.
40.   Francisco AB, Singh R, Li S, Vani AK, Yang L, Munroe RJ, Diaferia G, Cardano M,
Biunno I, Qi L, Schimenti JC, Long Q. Deficiency of Suppressor Enhancer Lin12 1 Like (SEL1L)
in Mice Leads to Systemic Endoplasmic Reticulum Stress and Embryonic Lethality. *Journal of
Biological Chemistry*. 2010;285(18):13694–13703.
41.   Yagishita N, Ohneda K, Amano T, Yamasaki S, Sugiura A, Tsuchimochi K, Shin H,
Kawahara K, Ohneda O, Ohta T, Tanaka S, Yamamoto M, Maruyama I, Nishioka K, Fukamizu
A, et al. Essential Role of Synoviolin in Embryogenesis. *Journal of Biological Chemistry*.
2005;280(9):7909–7916.
42.   Biunno I, Castiglioni B, Rogozin IB, DeBellis G, Malferrari G, Cattaneo M. Cross-species
conservation of SEL1L, a human pancreas-specific expressing gene. *Omics : a journal of
integrative biology*. 2002;6(2):187–98.
43.   Exposito J-Y, Cluzel C, Garrone R, Lethias C. Evolution of collagens. *The Anatomical
Record*. 2002;268(3):302–316.
44.   Exposito JY, le Guellec D, Lu Q, Garrone R. Short chain collagens in sponges are
encoded by a family of closely related genes. *Journal of Biological Chemistry*.
1991;266(32):21923–21928.
45.   Fidler AL, Vanacore RM, Chetyrkin S v, Pedchenko VK, Bhave G, Yin VP, Stothers CL,
Rose KL, McDonald WH, Clark TA, Borza D-B, Steele RE, Ivy MT, Aspirnauts, Hudson JK, et al.
A unique covalent bond in basement membrane is a primordial innovation for tissue evolution.
Proceedings of the National Academy of Sciences of the United States of America.
2014;111(1):331–6.
46.   Hecker L, Logsdon NJ, Kurundkar D, Kurundkar A, Bernard K, Hock T, Meldrum E,
Sanders YY, Thannickal VJ. Reversal of persistent fibrosis in aging by targeting Nox4-Nrf2
redox imbalance. *Science Translational Medicine*. 2014;6(231).
47.   Peng R, Sridhar S, Tyagi G, Phillips JE, Garrido R, Harris P, Burns L, Renteria L, Woods
J, Chen L, Allard J, Ravindran P, Bitter H, Liang Z, Hogaboam CM, et al. Bleomycin induces
molecular changes directly relevant to idiopathic pulmonary fibrosis: a model for "active"
disease. *PloS one*. 2013;8(4):e59348.
48.   Morse C, Tabib T, Sembrat J, Buschur KL, Bittar HT, Valenzi E, Jiang Y, Kass DJ,
Gibson K, Chen W, Mora A, Benos P v, Rojas M, Lafyatis R. Proliferating SPP1/MERTK-
expressing macrophages in idiopathic pulmonary fibrosis. *The European respiratory journal*.
2019;54(2).

49.   Reyfman PA, Walter JM, Joshi N, Anekalla KR, McQuattie-Pimentel AC, Chiu S,
Fernandez R, Akbarpour M, Chen C-I, Ren Z, Verma R, Abdala-Valencia H, Nam K, Chi M, Han
S, et al. Single-Cell Transcriptomic Analysis of Human Lung Provides Insights into the
Pathobiology of Pulmonary Fibrosis. *American journal of respiratory and critical care medicine*.
2019;199(12):1517–1536.
50.   Habermann AC, Gutierrez AJ, Bui LT, Yahn SL, Winters NI, Calvi CL, Peter L, Chung M-
I, Taylor CJ, Jetter C, Raju L, Roberson J, Ding G, Wood L, Sucre JMS, et al. Single-cell RNA
sequencing reveals profibrotic roles of distinct epithelial and mesenchymal lineages in
pulmonary fibrosis. *Science advances*. 2020;6(28):eaba1972.
51.   Adams TS, Schupp JC, Poli S, Ayaub EA, Neumark N, Ahangari F, Chu SG, Raby BA,
DeIuliis G, Januszyk M, Duan Q, Arnett HA, Siddiqui A, Washko GR, Homer R, et al. Single-cell
RNA-seq reveals ectopic and aberrant lung-resident cell populations in idiopathic pulmonary
fibrosis. *Science advances*. 2020;6(28):eaba1983.
52.   Tsukui T, Sun K-H, Wetter JB, Wilson-Kanamori JR, Hazelwood LA, Henderson NC,
Adams TS, Schupp JC, Poli SD, Rosas IO, Kaminski N, Matthay MA, Wolters PJ, Sheppard D.
Collagen-producing lung cell atlas identifies multiple subsets with distinct localization and
relevance to fibrosis. *Nature communications*. 2020;11(1):1920.
53.   Weathington NM, van Houwelingen AH, Noerager BD, Jackson PL, Kraneveld AD, Galin
FS, Folkerts G, Nijkamp FP, Blalock JE. A novel peptide CXCR ligand derived from extracellular
matrix degradation during airway inflammation. *Nature Medicine*. 2006;12(3):317–323.
54.   Riley DJ, Berg RA, Soltys RA, Kerr JS, Guss HN, Curran SF, Laskin DL. Neutrophil
Response following Intratracheal Instillation of Collagen Peptides into Rat Lungs. *Experimental
Lung Research*. 1988;14(4):549–563.
55.   Jürgensen HJ, Nørregaard KS, Sibree MM, Santoni-Rugiu E, Madsen DH, Wassilew K,
Krustrup D, Garred P, Bugge TH, Engelholm LH, Behrendt N. Immune regulation by fibroblasts
in tissue injury depends on uPARAP-mediated uptake of collectins. *Journal of Cell Biology*.
2019;218(1):333–349.
56.   Li L, Chen X, Zhang H, Wang M, Lu W. MRC2 Promotes Proliferation and Inhibits
Apoptosis of Diabetic Nephropathy. *Analytical Cellular Pathology*. 2021;2021:1–10.
57.   Onursal C, Dick E, Angelidis I, Schiller HB, Staab-Weijnitz CA. Collagen Biosynthesis,
Processing, and Maturation in Lung Ageing. *Frontiers in Medicine*. 2021;8.
58.   Podolsky MJ, Gupta D, Ha A, Ta R, Khalifeh-Soltani A, McKleroy W, Datta R, Sheppard
D, Atabai K. Cell division cycle 7 kinase is a negative regulator of cell-mediated collagen
degradation. *American Journal of Physiology - Lung Cellular and Molecular Physiology*.
2018;315(3):L360–L370.
59.   Boettcher M, Tian R, Blau JA, Markegard E, Wagner RT, Wu D, Mo X, Biton A, Zaitlen
N, Fu H, McCormick F, Kampmann M, McManus MT. Dual gene activation and knockout screen
reveals directional dependencies in genetic networks. *Nature biotechnology*. 2018;36(2):170–
178.
60.   Boettcher M, Covarrubias S, Biton A, Blau J, Wang H, Zaitlen N, McManus MT. Tracing
cellular heterogeneity in pooled genetic screens via multi-level barcoding. *BMC genomics*.
2019;20(1):107.
61.   Li W, Xu H, Xiao T, Cong L, Love MI, Zhang F, Irizarry RA, Liu JS, Brown M, Liu XS.
MAGeCK enables robust identification of essential genes from genome-scale CRISPR/Cas9
knockout screens. *Genome biology*. 2014;15(12):554.
62.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich
A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP. Gene set enrichment analysis: a
knowledge-based approach for interpreting genome-wide expression profiles. *Proceedings of
the National Academy of Sciences of the United States of America*. 2005;102(43):15545–50.
63.   Mootha VK, Lindgren CM, Eriksson K-F, Subramanian A, Sihag S, Lehar J, Puigserver
P, Carlsson E, Ridderstråle M, Laurila E, Houstis N, Daly MJ, Patterson N, Mesirov JP, Golub

TR, et al. PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately
downregulated in human diabetes. *Nature genetics*. 2003;34(3):267–73.
64.   Altenhoff AM, Train C-M, Gilbert KJ, Mediratta I, Mendes de Farias T, Moi D, Nevers Y,
Radoykova H-S, Rossier V, Warwick Vesztrocy A, Glover NM, Dessimoz C. OMA orthology in
2021: website overhaul, conserved isoforms, ancestral gene order and more. *Nucleic acids*
research. 2021;49(D1):D373–D379.
65.   Shrestha N, Liu T, Ji Y, Reinert RB, Torres M, Li X, Zhang M, Tang C-HA, Hu C-CA, Liu
C, Naji A, Liu M, Lin JD, Kersten S, Arvan P, et al. Sel1L-Hrd1 ER-associated degradation
maintains β cell identity via TGF-β signaling. *The Journal of clinical investigation*.
2020;130(7):3499–3510.
66.   Zhou Z, Torres M, Sha H, Halbrook CJ, van den Bergh F, Reinert RB, Yamada T, Wang
S, Luo Y, Hunter AH, Wang C, Sanderson TH, Liu M, Taylor A, Sesaki H, et al. Endoplasmic
reticulum–associated degradation regulates mitochondrial dynamics in brown adipocytes.
Science. 2020;368(6486):54–60.
67.   Bhattacharya A, Sun S, Wang H, Liu M, Long Q, Yin L, Kersten S, Zhang K, Qi L.
Hepatic Sel1L-Hrd1 ER-associated degradation (ERAD) manages FGF21 levels and systemic
metabolism via CREBH. *The EMBO Journal*. 2018;37(22).
68.   Yan C, Xu X, Zou X. Fully Blind Docking at the Atomic Level for Protein-Peptide
Complex Structure Prediction. *Structure*. 2016;24(10):1842–1853.
69.   Xu X, Yan C, Zou X. MDockPeP: An *ab‑initio* protein–peptide docking server. *Journal of*
Computational Chemistry. 2018;39(28):2409–2413.
70.   Baek M, DiMaio F, Anishchenko I, Dauparas J, Ovchinnikov S, Lee GR, Wang J, Cong
Q, Kinch LN, Schaeffer RD, Millán C, Park H, Adams C, Glassman CR, DeGiovanni A, et al.
Accurate prediction of protein structures and interactions using a three-track neural network.
Science. 2021;373(6557):871–876.
971

Figure 1. CRISPR screens identify mediators of cellular collagen uptake. (A) Schematic of experimental approach for flow cytometry-based screening for uptake of collagen fragments. (B) Volcano plot of gene-level statistics of CRISPRi (inhibition) screen. Most significant gene hits are labeled and colored in red. RRA = Robust Rank Aggregation score; LFC = Log-fold change. (C) Volcano plot of gene-level statistics of CRISPRa (activation) screen. (D) Comparison of overall phenotype score between CRISPRi and CRISPRa screens.

Supplemental Figure 1. Re-screens are correlated to original CRISPR screen results. (A) Volcano plot of CRISPRi re-screen (Screen 2). (B) Volcano plot of CRISPRa re-screen (Screen 2). (C) Comparison of gene-level overall phenotype scores between CRISPRi screens 1 and 2. (D) Comparison of gene-level overall phenotype scores between CRISPRa screens 1 and 2. (E) Comparison of arrayed log-fold change in arrayed validation vs. original pooled screen log-fold change; each point represents unique guide sequence. 65/84 guides demonstrated concordance with original pooled screen data (77%). Pearson correlation coefficient is shown.

Figure 2. Collagen-biosynthetic pathways are regulators of collagen uptake uniquely compared to screens for uptake of other cargoes. (A) Ranked plot of Normalized Enrichment Scores derived from Gene-Set Enrichment Analysis (GSEA) of CRISPRi phenotype scores, using 10,561 Gene Ontology gene sets. Collagen-related gene sets are labeled and displayed in red. (B) Comparison of GSEA statistics for labeled collagen-related gene sets across screens for uptake of various cargoes[22]; screen method is indicated in parentheses. KD = Knock-down.

Supplemental Figure 2. Collagen-related gene ontology terms are enriched as regulators of collagen uptake in CRISPRi and CRISPRa screens. The top twenty positive and negatively performing gene sets are indicated with associated statistics for each screen as indicated, among all 10,561 Gene Ontology gene sets. Collagen-related gene sets are highlighted with red boxes (i.e. those with 'Collagen' in the GO gene set name). NES = Normalized Enrichment Score.

Figure 3. Collagen biosynthesis positively regulates cell-based collagen clearance and *MRC2* expression. (A) Heatmap of overall phenotype scores from original screens as indicated for top genes from 'Collagen Biosynthetic Process' GO gene-set (displayed genes had most extreme 'Score' values in one or more screen result). (B) Flow-cytometry based collagen-uptake assay in U937 cells after shRNA treatment as indicated vs. Scramble control. MFI = Mean Fluorescence Intensity. N=4 per group. (C) Western blot of U937 cells after shRNA treatment as indicated vs. Scramble control, representative of N=4 independent experiments. (D) Q-RT-PCR in U937 cells after shRNA treatment as indicated vs. Scramble control. N=4 per group. (E-F) Western blot and densitometry of MRC5 fibroblasts after shRNA treatment as indicated vs. Scramble control, representative of N=3 independent experiments. (G) Representative confocal immunofluorescence images of MRC5 fibroblasts after shRNA treatment as indicated vs. Scramble control. (H) Flow-cytometry based collagen-uptake assay in MRC5 cells after shRNA treatment as indicated vs. Scramble control. N=4 per group. (I) Flow-cytometry based collagen-uptake assay in U937 cells after shRNA treatment as indicated vs. Scramble control. N=4-6 per group. (J-K) Q-RT-PCR in U937 cells after shRNA treatment as indicated vs. Scramble control. N=4-5 per group. (L) Flow-cytometry based collagen-uptake assay in MRC5 fibroblasts after shRNA treatment as indicated vs. Scramble control. N=6 and 4 in Scramble and shCOL1A1 groups respectively, N=2 in shMRC2 group. (M-N) Western blot and densitometry of MRC5 fibroblasts after shRNA treatment as indicated vs. Scramble control, representative of N=4 independent experiments. Statistics: (B-K) ANOVA with post-hoc Bonferroni testing; (L-N) Student's *t*-test. * p < 0.05, ** p < 0.01, **** p < 0.0001.

Supplemental Figure 3. Validation of different tools used in this manuscript. (A) Q-RT-PCR in U937 cells after shRNA treatment as indicated vs. Scramble control. N=4 per group. (B-C) Additional densitometry of data from Figure 5C. N=4 per group. (D) Q-RT-PCR in MEFs after OHT treatment for genes as indicated. N=3 per group. (E) Additional densitometry of data from Figure 6B. N=4 per group. (F) Densitometry of the left side of Figure 6E. N=6 per group. Statistics: (A, E, F) ANOVA with post-hoc Bonferroni testing; (B-D) Student's *t*-test. * p < 0.05, ** p < 0.01, **** p < 0.0001, ns = not significant.

Figure 4. Pharmacological stimulation of collagen synthesis is positively associated with collagen uptake and *MRC2* expression. (A-B) Flow-cytometry based collagen-uptake assay in U937 (A) or MRC5 (B) cells after drug treatment as indicated vs. vehicle control. N=4 per group. Western blot of corresponding cells and conditions shown below, each representative of N=3-4 independent experiments. (C-E) Q-RT-PCR in MRC5 cells after drug treatment as indicated vs. vehicle control. N=3-5 per group. Western blot of corresponding cells and conditions shown below, each representative of N=3 independent experiments. Statistics: (A-E) Student's *t*-test. * $p < 0.05$, ** $p < 0.01$.

Figure 5. *Sel1L* positively regulates collagen uptake and *Mrc2* expression. (A) Schematic of domain structure of SEL1L protein. FN2 = Fibronectin-2. (B) Flow-cytometry based collagen-uptake assay in MEF cells (mouse embryonic fibroblasts) after treatment with 4-hydroxytamoxifen (OHT) as indicated vs. vehicle control. WT = *Sel1L^{f/f};ERcre-*, iKO = *Sel1L^{f/f};ERcre+*. N=3-4 per group. (C-D) Western blot and densitometry of MEFs after OHT treatment as indicated vs. control, representative of N=4 independent experiments. (E) Q-RT-PCR in MEFs after OHT treatment for genes as indicated. N=3 per group. (F) Flow-cytometry based measurement of MRC2 surface expression in MEFs after OHT treatment. N=4 per group. (G) Representative wide-field immunofluorescence images of MEFs after OHT treatment. (H) Q-RT-PCR of MRC5 cells transfected with overexpression vectors containing genes as indicated vs. vehicle control (lipofectamine only). N=4 per group. (I) Western blot of MRC5 cells transfected with overexpression vectors containing genes as indicated vs. vehicle control (lipofectamine only). N=2. Statistics: (B, H) ANOVA with post-hoc Bonferroni testing; (D - right panel, E, F) Student's *t*-test; (D - left panel) Mann-Whitney *U* test. * p < 0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001, ns = not significant.

Figure 6. SEL1L, but not HRD1, positively regulates MRC2. A) Flow-cytometry based collagen-uptake assay in 293T of different genotypes (compared with baseline control without fluorescent collagen). N=4-5 per group. (B-C) Western blot and densitometry of 293T cells of different genotypes, representative of N=5 independent experiments. (D) Firefly luciferase assay of *MRC2* promoter activity, normalized to constitutive Renilla luciferase in 293T cells of different genotypes. N=3 per group. (E) Western blot of HEK293T cells of different genotypes transfected with MRC2 overexpression vector vs. vehicle control, representative of N=6 independent experiments. (F) Representative wide-field immunofluorescence images of 293T cells of different genotypes transfected with MRC2 overexpression vector vs. vehicle control. (G) Flow-cytometry based collagen-uptake assay in HEK293T of different genotypes transfected with MRC2 overexpression vector (compared with baseline control without fluorescent collagen; note different scale compared with panel A). N=3-4 per group. Statistics: (A, C, D, G) ANOVA with post-hoc Bonferroni testing. * p < 0.05 ** p < 0.01, ns = not significant.

A

Green: collagen peptide; Blue: FN2 domain of SEL1L

B

Supplemental Figure 4. Additional support of SEL1L and collagen interaction. (A) Predicted docking of collagen peptide with FN2 domain of SEL1L. (B) Additional representative confocal immunofluorescence image of MRC5 fibroblasts at baseline (as in Figure 7A), with decreased PMT gain in the Collagen channel, along with magnified inset to highlight co-localization.

Figure 7. SEL1L binds collagen as it is being synthesized via its FN2 domain and this interaction is necessary for regulation of *MRC2* expression. (A) Representative confocal immunofluorescence images of MRC5 fibroblasts treated as indicated or vehicle control. (B) Western blot of MRC5 fibroblasts after immunoprecipitation with anti-SEL1L antibody (vs. controls as indicated), representative of N=2 independent experiments. (C) Western blot after immunoprecipitation with anti-procollagen type I antibody (with input controls as indicated) of HEK293T cells transfected with collagen and WT or mutant SEL1L constructs as indicated, representative of N=3 independent experiments. (D) Representative confocal immunofluorescence images of HEK293T cells transfected with collagen and WT or mutant SEL1L constructs as indicated. (E) Western blot of MEFs after treatment with WT or mutant SEL1L constructs as indicated, representative of N=3 independent experiments. (F) Flow-cytometry based measurement of MRC2 surface expression on MEFs after treatment with WT or mutant SEL1L constructs as indicated, vs. Empty Vector (EV). N=6 per group. (G) Western blot of HEK293T cells either WT or KO for SEL1L as indicated after treatment with EV (empty vector), WT or mutant SEL1L constructs as indicated, representative of N=3 independent experiments. (H) Q-RT-PCR of SEL1L-KO HEK293T cells after transfection with WT or mutant SEL1L as indicated vs. empty vehicle control. N=3 per group. (I) Comparison of protein domain structures of Sel1L orthologs across phylogenetically diverse organisms; FN2 = Fibronectin-2 domain, tpr = Tetratricopeptide repeat domain. Statistics: (F-G) ANOVA with post-hoc Bonferroni testing. * p < 0.05, ** p < 0.01, ns = not significant.

Supplemental Figure 5. FN2 domain deletion mutant interferes with collagen binding but is otherwise stable. (A) Alignment of predicted structures of full-length SEL1L and ΔFN2 SEL1L. Two different views and alignments are shown. Root-mean squared distance (RMSD) is displayed. Yellow arrow indicates the FN2 domain (in the pink structure on the left; in the green structure on the right). (B) Representative confocal immunofluorescence images of HEK293T cells transduced with MYC-tagged WT or ΔFN2 SEL1L constructs, co-stained with ER marker Calnexin. (C) Western blot of purified WT or ΔFN2 SEL1L proteins compared with input cell lysates, representative of N=3 independent experiments. (D) Circular Dichroism spectra of purified WT or ΔFN2 proteins. (E) Western blot of co-incubated purified either WT or ΔFN2 SEL1L proteins with rat tail collagen, after immunoprecipitation with anti-MYC antibody (with input and additional controls as indicated), representative of N=3 independent experiments. 'Collagen only' indicates incubation of rat tail collagen with IP beads without co-incubation with purified MYC-tagged protein.

Figure 8. Collagen biosynthesis and *MRC2* expression are positively correlated in lung tissue, but this relationship is impaired in human fibrotic lung disease. (A) UMAP plot with cell type annotation of scRNAseq data from multiple available datasets of human lung control and IPF tissue (GSE136831, GSE135893, GSE121611, GSE128033, GSE132771). N=184,672 single cells. (B-C) Gene expression of individual genes as indicated in UMAP as generated in panel (A). (D) Comparison of gene expression in all cells from data as in panel (A) that had non-zero expression of *COL1A1* or *MRC2*. (E) Comparison of gene expression in whole human lung microarray data from GSE110147 analyzed via GEO2R in pulmonary fibrosis lungs vs. control specimens. N=37 vs. 11 per group. (F) Comparison of microarray gene expression in human lung bronchoalveolar lavage specimens from GSE70867 analyzed via GEO2R from Idiopathic Pulmonary Fibrosis (IPF) lungs vs. control specimens. N=112 vs. 20. (G) Comparison of microarray gene expression in peripheral blood mononuclear cell (PBMC) specimens from GSE37858 analyzed via GEO2R from Idiopathic Pulmonary Fibrosis (IPF) lungs vs. control specimens. N=70 vs. 35. (H) Comparison of gene expression in whole mouse lung microarray data from GSE40151 analyzed via GEO2R from Bleomycin-treated mice vs. Saline control mice. N=56 vs. 55. Statistics: ANCOVA, p-values indicated in figures. Pearson correlations are displayed in panel (D).

Figure 9. SEL1L protein levels are reduced in the lungs of patients with Pulmonary Fibrosis. (A-B) Western blots of whole human lung lysates from Pulmonary Fibrosis lungs vs. control (donor lungs deemed not suitable for transplant), with corresponding densitometry. N=13 in Control and 11 in Pulmonary Fibrosis group. Arrowheads correspond to two lung lysates that were used in both Western blots for normalization across blots, otherwise all other lanes are independent samples. (C) Representative low-power wide-field immunofluorescence images of human lungs (Pulmonary Fibrosis vs. control). Each row comes from an individual donor or Pulmonary Fibrosis lung, with two different fields per lung (i.e. N=3 individual lungs per group). Statistics: (B, D) Mann-Whitney U test. * p < 0.05.