Tissue-specific chromatin binding patterns of *C. elegans* heterochromatin proteins HPL-1 and HPL-2 reveal differential roles in the regulation of gene expression.

Patricia de la Cruz Ruiz[1], Peter Askjaer[1,*] and Marta Artal-Sanz[1,2,*]

[1]Andalusian Centre for Developmental Biology, Consejo Superior de Investigaciones Científicas/Junta de Andalucía/Universidad Pablo de Olavide, Seville, Spain
[2]Department of Molecular Biology and Biochemical Engineering, Universidad Pablo de Olavide, Seville, Spain

*Correspondence to: Peter Askjaer (pask@upo.es) and Marta Artal-Sanz (martsan@upo.es)

Andalusian Centre for Developmental Biology. University Pablo de Olavide.
Ctra. De Utrera km.1. 41013 Sevilla. Spain.

Running title: Tissue-Specific HP1 Chromatin Binding

Keywords: *C. elegans*; Chromatin; DamID; Gene expression; Heterochromatin protein 1; HPL-1; HPL-2.

Abstract

Heterochromatin is characterized by an enrichment of repetitive elements and low gene density and is often maintained in a repressed state across cell division and differentiation. The silencing is mainly regulated by repressive histone marks, such as H3K9 and H3K27 methylated forms and the heterochromatin protein 1 (HP1) family. Here, we analyzed in a tissue-specific manner the binding profile of the two HP1 homologs in *C. elegans,* HPL-1 and HPL-2, at the L4 developmental stage. We identified the genome-wide binding profile of intestinal and hypodermal HPL-2 and intestinal HPL-1 and compared them to heterochromatin marks and other features. HPL-2 associated preferentially to the distal arms of autosomes and correlated positively with methylated forms of H3K9 and H3K27. HPL-1 was also enriched in regions containing H3K9me3 and H3K27me3 but exhibited a more even distribution between autosome arms and centers. HPL-2 showed a differential tissue-specific enrichment for repetitive elements, conversely with HPL-1 that exhibited a poor association. Finally, we found a significant intersection of genomic regions bound by the BLMP-1/PRDM1 transcription factor and intestinal HPL-1, suggesting a co-repression role during cell differentiation. Our study uncovers both shared and singular properties of conserved HP1 proteins, providing information about genomic binding preferences in relation to their role as heterochromatic markers.

Introduction

Heterochromatin in worms, like in other species, is tightly packaged, presents a low gene density and is usually enriched at the nuclear periphery (Ahringer and Gasser 2018; Gu and Fire 2010). Facultative heterochromatin is defined by the presence of histone H3 lysine 27 tri-methylation (H3K27me3), which can be erased by demethylation during development leading to a transcriptionally active state. The Polycomb PRC2 complex comprised by MES-2, MES-3, and MES-6 proteins is responsible for H3K27 methylation (Steffen and Ringrose 2014; Piunti and Shilatifard 2016). The constitutive form of heterochromatin is characterized by H3K9me3, which ensures homeostasis and differentiation of tissues across species (Padeken, Methot, and Gasser 2022). H3K9 methylation is performed in sequential steps by MET-2/SETDB1 enzyme that carries out mono- and di- methylation and SET-25/G9a, in charge of tri-methylation. Both enzymes are main players in the perinuclear attachment of H3K9me enriched heterochromatin (Towbin et al. 2012). These heterochromatin domains usually reside within the chromosomal distal regions ("arms"), where most meiotic recombination occurs. The arms are also enriched in repetitive DNA elements, decorated by H3K9me2 and/or H3K9me3 (Liu et al. 2011; Consortium 1998; McMurchy et al. 2017). On the contrary, genes in autosome centers are generally higher expressed and more evolutionarily conserved (Consortium 1998).

In *C. elegans*, HPL-2 is considered as a marker for heterochromatin domains in developing embryos (Grant et al. 2010). HPL-2 is part of the heterochromatin protein 1 (HP1) family, which is highly conserved across the eukaryotic kingdom. HP1 was first characterized in fruit flies, as a component responsible for variegation position effect, in which inhibition of gene expression occurs via imposition of heterochromatin-like structure to inactivate it (Eissenberg et al. 1990). The two HP1 homologues in *C. elegans,* HPL-1 and HPL-2, possess a conserved structure that consists of an N-terminal chromo domain (CD)

involved in binding to the K9 residue on histone H3 in its mono, di or tri-methylated form. The CD is followed by a flexible hinge region and a C-terminal chromo shadow domain (CSD) required for dimerization and interaction with other proteins (Schott et al. 2006; Eissenberg 2001).

HPL-1 and HPL-2 are 48% identical along their entire length, with the highest homology within the CD and CSD. Their expression is ubiquitous in nuclei of most cells throughout development (Couteau et al. 2002; Schott et al. 2006). The observation of post-embryonic defects in *hpl-2* mutants, such as sterility, defective growth and brood size, larval arrest and multivulva phenotypes demonstrated that HPL-2 functions in cell fate determination and gonad development. The similar synthetic phenotypes presented in double *hpl-1;hpl-2* mutants prompted to conclude a redundant functionality for HPL-1, which did not show obvious phenotypes when mutated alone (Schott et al. 2006; Simonet et al. 2007; Schott et al. 2009; Coustham et al. 2006; Cardoso et al. 2005; Koester-Eiserfunke and Fischle 2011). HPL-2 has also been implicated in DNA replication, stress response and small RNAs pathways, being enriched in DNA repetitive elements (Coustham et al. 2006; Schott et al. 2009; Black et al. 2010; Cardoso et al. 2005; Meister et al. 2011; Garrigues et al. 2015).

Comparatively, little is known about HPL-1 functions in *C. elegans*. A physical interaction between HPL-1 and the mono-methylated H1K14me1 histone variant has been observed to regulate innate immunity against pathogenic bacterial infection (Studencka, Konzer, et al. 2012). Like HPL-2, loss of HPL-1 results in transcriptional alteration of genes that encode nuclear hormone receptor family proteins such as NHR-60, NHR-156, transcription factors including MIZ-1, ZIP-3, KZIP-8, MADF-2, homeobox genes like CEH-82 or the homeodomain gene LIM-7, that cause male tail defects (Studencka, Wesołowski, et al. 2012). Recently, HPL-1 was related to transgenerational embryonic lethality under specific paternal DNA damage. HPL-1 depletion induced a reduction in H3K9me2 levels in the germline, which was beneficial for F2 generation survival (Wang, Meyer, and Schumacher 2022).

Despite the association of HP1 proteins with different functions, tissue-specific behavior of HPL-1 and HPL-2 has not been characterized. These types of studies have become important to investigate heterogeneity between tissues. This may help to understand, for instance, potential regulatory links between epigenetic alterations and target gene expression changes to establish an integrative analysis (Yilmaz et al. 2020). To reinforce this idea, contrasting results have been described when performing rescue experiments with HP1 proteins upon endoplasmic reticulum stress. For example, upon whole-body loss of HPL-2, intestinal-specific expression of HPL-2 restored wild type level of the stress response, while it was detrimental upon neuronal-specific expression (Kozlowski et al. 2014). However, overexpression in either of the two tissues, was sufficient to provide partial rescue of dauer exit (Meister et al. 2011). This highlights differences in tissue response to lack of HPL-2 when evaluating distinct pathways.

To contribute to the characterization of HP1 proteins, a comparative study of HPL-1 and HPL-2 homologous proteins in *C. elegans* in a tissue-specific manner is reported here. We identified specific genomic regions bound by HPL-2 in hypodermal and intestinal tissues, as well as HPL-1-associated regions in the intestine, at the L4 developmental stage. We found

that HPL-2 associated preferentially to chromosome distal arms, while HPL-1 was equally distributed between autosome arms and center. Moreover, both HP1 homologs strongly correlated with the two repressive histone marks H3K9me3 and H3K27me3 in both tissues analyzed. However, only HPL-2 binding pattern coincided with mono- and di- methylated H3K9 forms.

We confirmed a general repressive role for HP1 proteins when comparing with transcribed genes in the tissues studied. In addition, the large number of unique genes, together with gene ontology analyses, suggest tissue-specific patterns of gene silencing. Moreover, we discovered a potential co-regulation of chromatin regions by BLMP-1/PRDM1 transcription factor and intestinal HPL-1. Lastly, HPL-2 was found to be enriched at DNA repetitive elements in both tissues, conversely to intestinal HPL-1. In summary, our study uncovered both unique and common features of HP1 *C. elegans* homologous proteins in two differentiated tissues.

Results

HPL-1 and HPL-2 differentially associate to the genome

To study the function of *C. elegans* HP1 proteins, the DNA binding profiles for HPL-1 and HPL-2 were determined by tissue-specific DNA adenine methylation identification (DamID) at the L4 larval stage (de la Cruz Ruiz et al 2022). We focused on the intestine, a major metabolic tissue of *C. elegans* amenable to this technique (Cabianca et al, 2019), and included also the hypodermis for HPL-2 to compare binding profiles across two tissues. We first analyzed the results at 100 kb resolution to obtain a broad overview of the behavior of the two HP1 proteins. In agreement with previous observations in embryonic cells, HPL-2 is enriched in distal regions ("arms") of autosomes in both hypodermal and intestinal tissues and depleted from autosome centers (Figure 1A; Supplementary Figure 1A)(Garrigues et al. 2015). We found that chromosome arms with meiotic pairing center were more frequently in contact with HPL-2 in the hypodermis compared to the intestine (Figure 1A). Surprisingly, HPL-1 in the intestine is distributed more equally between autosome arms and centers, being significantly different from HPL-2 (Figure 1A). The difference between HPL-1 and HPL-2 binding profiles in the intestine is also observed at higher resolution (10 kb) for all autosomes (Figure 1B; Supplementary Figure 1A). This suggests that HPL-1 might have unique roles in *C. elegans*. In addition, both HPL-2 DamID profiles exhibit a significant overlap with previous whole-animal HPL-2 ChIP datasets (Gerstein et al. 2010)(Supplementary Figure 2 A-C; Supplementary Table 1).

HPL-1 and HPL-2 differentially correlate with H3K9me3 and H3K27me3 histone marks

As reader of methylated H3K9, HPL-2 binding sites in the genome coincide with the pattern of mono and di-methylated H3K9 (H3K9me1-2) in *C. elegans* embryos, and less so with H3K9me3 (Garrigues et al. 2015). To determine if the correlation also exists in differentiated cells and for both HP1 proteins, we compared the HPL-1 and HPL-2 binding profiles with

published histone ChIP datasets from the L3 larval stage. Indeed, a high correlation was found between both HP1 proteins and repressive H3K9me3 and H3K27me3 histone marks, clustering together in the pairwise Pearson correlation heatmap (Figure 1C). Both histone marks usually colocalize on distal autosome arms, while only H3K27me3 associates with central regions as well (Liu et al. 2011; Ahringer and Gasser 2018).

Interestingly, HPL-2 in both tissues positively correlated with mono and di-methylated H3K9, whereas HPL-1 protein failed to show any significant association (Figure 1 C-D, Supplementary Figure 1B-D). Therefore, our data indicates that only HPL-2 exhibits more flexibility to bind to the three methylated forms of H3 in both differentiated tissues, suggesting a differential role in the control of gene expression for HPL-1 and HPL-2 through development.

HP1 proteins bind to genomic regions in a tissue-specific manner

We next asked if the differences in distribution of large HP1-associated genomic regions across autosomes (Figure 1A-B) and the correlation with heterochromatin marks (Figure 1C-D) reflect a differential binding of HPL-1 and HPL-2 to smaller specific DNA regions. In addition, the different proportion of overlapping peaks between HP1 proteins and whole-animal HPL-2 ChIP data could indicate tissue-specific differences (Supplementary Figure 2A; Supplementary Table 1). To address this, we compared the regions present in each HP1 dataset. Strikingly, we found that around 70% of bound genomic regions are unique for either HPL-1 or HPL-2 (Figure 2A). The highest overlap was found between the two tissue-specific HPL-2 profiles, suggesting a conserved regulation for specific regions in both tissues (Figure 2A). However, the proportion of overlapped regions between all datasets does not reach 30%, remarking the high degree of specificity.

Annotation of genomic features for regions bound by HP1 proteins showed that more than 70% of the peaks reside inside protein coding regions (Figure 2B; see peak ratio). Most bound regions are placed within transcribed chromatin, and a notable proportion overlaps with transcription start sites (TSS) or promoter regions (Figure 2B).

In order to know whether genes bound by HP1 proteins are repressed or activated, a comparison with gene datasets from alternative methods of tissue-specific transcriptomic profiling was performed. First, for hypodermal tissue, we selected genes identified in at least three out of four transcriptomic methods (Kaletsky et al. 2018; Katsanos et al. 2021; Blazie et al. 2017; Fragoso-Luna et al. 2022). We found a significant underrepresentation of transcribed genes among HPL-2 hypodermal-bound genes (<3% overlap; $p < 0.001$), suggesting a repressive role for HPL-2 in the hypodermis (Figure 2C; Supplementary_File_1; Supplementary Table 2). We followed a similar procedure with intestinal genes, resulting in an underrepresentation of transcribed genes when compared with HPL-1-bound genes (<3% overlap; $p < 6.862e-07$) (Figure 2D; Supplementary File 1; Supplementary Table 2)(Gómez-Saldivar et al. 2020; Kaletsky et al. 2018; Katsanos et al. 2021; Blazie et al. 2017; Haenni et al. 2012). In contrast, HPL-2 in the intestine did not show any tendency (Figure 2D). These comparisons suggest a tissue-specific repressive role for HPL-2 and that HPL-1 and HPL-2 act differently in intestinal cells.

Interestingly, when comparing HP1 bound genes across the three datasets, we found that around 60% of genes are unique in each dataset, similar to the proportion of unique binding sites (Figure 3A; compare with Figure 2A; Supplementary File 2). Gene ontology (GO) enrichment analysis for biological processes with genes bound by HPL-2 in the intestine identified only a few and rather broad terms, such as development and anatomical structures (Figure 3B). Because these terms are based on genes primarily expressed in proliferative tissues, their association with HPL-2 in terminally differentiated intestinal cells is concordant with a repressive role of HPL-2. Similarly, gene sets retrieved from either HPL-2 DamID in hypodermal tissue or HPL-1 DamID in the intestine identified only a few enriched GO terms and none of them related to the tissue where the Dam fusion protein was expressed (Supplementary Figure 3A-B).

In our attempt to get more insight about new biological functions of HP1 proteins, we compared HP1-bound regions with those bound by different transcription factors at the L1 larval stage (ALR-1, MAB-5, EOR-1, PQM-1, PHA-4, ELT-3, ELT-2, EGL-5, SKN-1, UNC-130, EGL-27 and BLMP-1) (Niu et al. 2011). Interestingly, this analysis revealed a significant overlap between the transcription factor BLMP-1/PRDM1 and HPL-1 in the intestine, which is maintained at the L4 stage as well (Supplementary Figure 3C-D; Supplementary Table 3)(Stec et al. 2021). Around 13% of peaks are common to HPL-1 and BLMP-1, mainly in autosome centers in L1 (Supplementary Figure 3E). In line with this, the genome-wide average signal of HP1 binding at BLMP-1 peaks showed a strong enrichment for HPL-1 (Figure 3C). Then, we assigned the BLMP1 significant peaks to genes using the same criteria and method employed for HP1 datasets describe here. We found a significant overlap between BLMP-1 and HP1 bound genes, including more than one hundred genes in common (Supplementary Figure 3F). Collectively, those results strongly suggest that HPL-1 and BLMP-1 could potentially co-regulate specific genes or transcription factors. Knowing that PRDM1, the human BLMP-1 orthologue, was first described as a repressor of beta-interferon (beta-IFN) expression, a tissue-specific co-repression by HPL-1 and BLMP-1 may occur (Keller and Maniatis 1991).

Only HPL-2 protein is enriched at repetitive elements

It has been previously reported a role for several heterochromatin proteins, among them HPL-2, in maintaining repetitive elements and particular genes silenced, which is crucial for germline and fertility processes (McMurchy et al. 2017; Ashe et al. 2012). HPL-2 is also enriched at repetitive elements both, in embryos and young adults (Garrigues et al. 2015; McMurchy et al. 2017). These analyses were performed in whole animals and hence offered no information about tissue-specificity nor the behavior of HPL-1. Therefore, we assessed potential differences in the binding of HPL-1 and HPL-2 to repetitive elements using annotation from the most recent UCSC Genome Browser based on repeat masker that classified 61,527 individual repetitive elements (https://genome.ucsc.edu)(Tarailo-Graovac and Chen 2009). Interestingly, our analysis corroborates the high proportion of overlapping regions between HPL-2 and genomic repetitive elements in both tissues (Figure 4A; Supplementary Table 4). However, the overlap is reduced and not significant in the case of HPL-1 protein (Figure 4A; Supplementary Table 4).

Moreover, the hypodermal tissue exhibited 14% more overlapping regions compared to the intestine for HPL-2 protein, remarking tissue-specific divergence. To further study this difference, we looked at repetitive element types, finding DNA transposon as the most predominant type for both tissues (Figure 4B). Being CELE14B, CELE42 and CELE14A the most frequent DNA transposons families found for both tissues (Supplementary Table 5). However, more than a quarter of repetitive elements uniquely enriched in the intestinal tissue correspond to retrotransposons, with almost no representation in the hypodermis (Figure 4B). The strongest association reported for HPL-2 in young adults was found with DNA transposons Helitron families, being less abundant compared to other families at the L4 stage analyzed here (Supplementary Table 5)(McMurchy et al. 2017). Nonetheless, for all stages analyzed here and in other studies, DNA transposons are the most representative, suggesting the necessity to maintain these repressed in differentiated tissues.

Discussion

Here we characterized the genomic binding profile of the two *C. elegans* HP1 protein homologs HPL-1 and HPL-2 in the intestine, as well as the binding profile of HPL-2 in the hypodermis. We showed that HPL-2-bound regions are enriched in distal chromosome arms, as published for embryos and young adults (McMurchy et al. 2017; Garrigues et al. 2015). In contrast, we showed an unprecedented location of HPL-1 in the center of autosomes.

When assessing the correlation of heterochromatin histone hallmarks, we found a strong correlation of HP1 proteins with H3K9me3 and H3K27me3. On the one hand, methylated H3K9 prevents the access of transcription factors to their binding sites in differentiated cells, being crucial for restricting plasticity and maintaining tissue identity. On the other hand, H3K27me3 seems to limit plasticity, acting in a parallel manner (Patel and Hobert 2017). Our data suggest that HP1 proteins act non-redundantly in controlling tissue-specificity in differentiated somatic cells. In agreement with this, cellular plasticity is affected both in single *hpl-1* or *hpl-2* mutants and more strongly upon deletion of both HP1 proteins (Patel and Hobert 2017). Moreover, Evans et al. described some domains decorated by H3K27me marks, considered the most suceptible to be regulated during development (Evans et al. 2016). Lastly, in embryos, genes expressed in terminally differentiated tissues were marked specifically by H3K9me3 (Zeller et al. 2016). It could be that HP1 proteins contribute to the formation of these domains, as well as controlling tissue-specific genes, affecting chromatin activity and gene expression in a context-dependent manner. In other words, the ability of HP1 proteins to bind concurrently to both repressive histone modifications, reinforce the repression, and could account for developmental gene regulatory networks.

In addition, HPL-2 showed a strong correlation with H3K9me1 and H3K9me2 in both tissues analyzed, which was not observed with HPL-1. In embryos, HPL-2 closely associated with H3K9me1 and H3K9me2, and less with H3 tri-methylated forms (Garrigues et al. 2015). Moreover, more than 20% of regulatory elements were uniquely enriched for H3K9me2 in this stage, mostly tandem or simple repeats type (Zeller et al. 2016). This could be in line with the tight association between HPL-2, but not HPL-1, and repetitive elements previously reported in other stages (McMurchy et al. 2017; Garrigues et al. 2015). Those repetitive elements are mostly represented by DNA transposon families, conversely to retrotransposon families. HPL-2 could bind to H3K9 di-methylated forms to prevent de-repression of repeats

and maintain the genome instability during development. The higher proportion of repetitive elements enrichment among HPL-2 peaks in the hypodermis compared to the intestine suggests a specific regulation of genome silencing, highlighting both unique and more general role for HPL-2 during tissue differentiation.

Lastly, the significant underrepresentation of transcribed genes bound by HP1 proteins in either hypodermis or intestine strongly suggests a general repressive role for both HPL-1 and HPL-2. However, the reduced number of peaks (thus genes) shared by all HP1 datasets indicate that they control gene expression in a tissue-specific manner. In fact, in flies, the characterization of HP1a, HP1b, and HP1c protein homologs revealed a tissue-specific pattern and a reduced number of canonical target genes (Schoelz, Feng, and Riddle 2021).

Our analysis showed a significant number of genomic regions commonly bound by intestinal HPL-1 and the BLMP-1/PRDM1 transcription factor in two developmental stages. In *C. elegans* BLMP-1 is relevant for the proper development of several tissues, such as, vulva, hypodermis and gonad (Huang et al. 2014; Horn et al. 2014; Yang et al. 2015). PRDM1 was first described as responsible for B cell identity by repressing other transcription factors, thereby controlling indirectly numerous target genes (Shaffer et al. 2002). Moreover, during cell plasma differentiation, PRDM1 recruits histone deacetylase HDAC1/2 and lysine demethylase LSD1 chromatin regulators, to repress mature B-cell gene expression pattern (Su et al. 2009). In worms, the recruitment of HDACs proteins through the binding of HPL-2 to H3K27me3 represses Hox genes (Jedrusik-Bode 2013). We speculate that HPL-1 and BLMP-1 could act in parallel to maintain silent genomic regions crucial for cell identity. As occurs in plasma cells, BLMP-1 may co-regulate with HPL-1 and other epigenetic factors the repression of specific genomic regions in order to control cell fate. The specific mechanism behind remains elusive, but most recent publication proposed the increase of chromatin accessibility for transcriptional repressor by BLMP-1 binding during development (Stec et al. 2021).

In summary, this tissue-specific study of HP1 proteins revealed important differences that highlight the relevance as heterochromatic markers for cell identity during development. This potential restriction of plasticity accompanied by changes in other crucial heterochromatin marks, such as histones modifications, open new questions about the role of chromatin modifiers in this complex process. Lastly, we revealed unique features for the underexplored HPL-1 homolog in *C. elegans*, which encourage further studies to uncover its biological role.

MATERIAL AND METHODS

C. elegans maintenance

Maintenance of *C. elegans* strains were done conforming to standard protocols on Nematode Growth Medium (NGM) plates, grown in a monoxenic lawn of *Escherichia coli* OP50 at 20°C (Stiernagle 2006). For DamID experiments, a lawn of *E. coli* GM119 strain was used to grow the worms (Arraj and Marinus 1983). The complete list of strains used in this study is available in Supplementary Table 6.

Mos Single Copy Insertion (MosSCI)

Single copy integration transgenic strains were made by microinjection into strain EG4322 (ChrII) with *unc−119(+)* recombination plasmids together with individual transgenes (pBN494 and pBN495) at 25ng/µl, Mos1 transposase (P*eft-3*::transposase) at 50ng/µl and three red co-injection markers pCFJ90 (P*myo-2::mcherry*) at 2,5 ng/µl, pCFJ104 (P*myo-3::mcherry*) at 5ng/µl and pBN1 (P*lmn-1::mch::his-58*) at 10ng/µl (Dobrzynska, Askjaer, and Gruenbaum 2016; Frøkjær-Jensen et al. 2012). Plasmids were microinjected into one of the gonad arms of egg-laying and locomotion defective *unc-119* young adult worms. Injected worms were individually transferred to NGM petri dishes and placed at 25°C. Successful integrants were spotted after 7−14 days based on wild type mobility and absence of co−injection markers. Description of plasmids can be found in Supplementary Table 7.

DamID-seq protocol

DamID experiments were performed as described in (de la Cruz Ruiz, Romero-Bueno, and Askjaer 2022). Strains were synchronized by standard hypochlorite treatment and grown until the L4 stage. Optimal number of 14x cycles were selected for PCR amplification of methylated sites.

DamID-seq data processing and visualization

The quality control of the DamID technique and the calculation of chromosome association profiles at 100 kb level for all HP1 samples were performed by damid.seq.r pipeline version 0.1.3 in Rstudio (R version 3.4.3) (available at github.com/damidseq/rDamIDSeq) (Sharma, Ritler, and Meister 2016). From FastQ files, the pipeline detects and maps reads containing the DamID adapters (adapt.seq="CGCGGCCGAG") and match them to genomic regions flanked by GATCs sites (restr.seq = "GATC") to obtain Bam Files. After mapping with the *C. elegans* genome release "BSgenome.celegans.ucSc.ce11", the log2 ratio HPL::Dam/GFP::Dam across two replicas was integrated to plot the relative read counts at 100 kb bin along the genome for all chromosome autosomes. Pairwise Wilcoxon rank test values were calculated in R. Coordinates of arms as defined in (Gonzalez-Aguilera et al. 2014). Read numbers for all datasets are summarized in Supplementary Table 8.

For the visualization using UCSC Genome Browser, the Bam files obtained in the previous pipeline (damid.seq.r) were used to generate the normalized log2 HPL::Dam/GFP::Dam ratio scores per GATC genome fragments, using perl script damidseq_pipeline v1.4.5 (Marshall and Brand 2015) (available at https://github.com/owenjm/damidseq_pipeline). The pipeline integrates a Bowtie 2 v2.3.4 (Langmead and Salzberg 2012) calling to map reads on *C. elegans* bowtie indexes from

assembly WBcel235 (available in Illumina iGenomes webpage; identical to ce11). In addition, it includes also Samtools v1.9 (Li et al. 2009) for alignment guidance, and a GFF file containing all GATC sites in the *C. elegans* genome. Last GATC coordinates file was built by gatc.track.maker.pl tool provided by the same pipeline script using WBcel235 FASTA file for *C. elegans* (available at https:// ensembl.org/Caenorhabditis_elegans/Info/Index. Final bedgraph files for each replica was averaged using average_tracks perl script included in damid_misc (available at https://github.com/owenjm/damid_misc). For comparison of chromosome IV arms versus centers, we used the following border coordinates: chrIV 4,790,000 & 12,610,000.

For the heatmap depicting Pearson correlations between Histone marks datasets we did a preprocessing to average the replicas in bedgraph/wig format using WiggleTools (Zerbino et al. 2014). Then, a fixed bin size was selected to standardized all datasets by "bedgraph_to_wig.pl" tool (created by Sebastien Vigneau in Alexander Gimelbrant lab; available at https://gist.github.com/svigneau/8846527). After converting wig files to bigwig format, we lifted-over the bigwig to equivalent genome version using Crossmap tool (Zhao et al. 2014). Finally, the matrix and heatmap were generated by "multiBigwigSummary" and "plotCorrelation" tools inside Deeptools (Ramírez et al. 2016). Sources for datasets listed in Supplementary Table 9.

Peak calling and analysis

The selection of statistically significant peaks for every HP1 sample was done by find_peaks perl script (available at https://github.com/owenjm/find_ peaks) with a false discovery rate (FDR) lower than 0.05 using as input averaged bedgraph files containing normalized log2 HPL::Dam/GFP::Dam ratio scores per GATC fragment.

Intersection between peaks from different datasets and statistical calculation of Fisher exact test, were performed by "Intersect" and "Fisher" tools included in Bedtools suit (Quinlan and Hall 2010). We include r and f=0.1 options, default mode for the rest. When needed, a lift-over between genome ce10, ce6 and ce11 was performed by "liftOver" tool (available in https://genome.ucsc.edu/cgi-bin/hgLiftOver). Pairwise association for peaks comparison was performed by intervene tool, using aforementioned bedtools options (Khan and Mathelier 2017). Heatmaps and Stacked plots for peaks were generated by GraphPad Prism 8 (www.graphpad.com). The coordinates for comparison between chromosome autosome arms and center were used as described in (Garrigues et al. 2015).

A statistical method based on Montecarlo simulation was applied to plot intersected peaks in Supplementary Figure 2B and C and Supplementary Figure 3E (Ferré et al. 2019).

Gene assignment and analysis

The statistically significant peaks selected were assigned to genes using UROPA (Kondili et al. 2017) as a web tool (available at http://loosolab.mpi-bn.mpg.de/UROPA_GUI/) with Caenorhabditis_elegans.WBcel235.107.gtf (from http://ensembl.org/Caenorhabditis_elegans/Info/Index) for genome protein coding annotation. Peaks were determined to genes on any strand when their center coordinate was localized up to 3 kb upstream of a gene start site or 3 kb downstream of the gene end site. We used UROPA tool to annotate genes from significant peaks of BLMP-1 dataset. Venn diagram overlapping gene datasets was performed using "Interactivenn" webtool (available in http://www.interactivenn.net/index.html) (Heberle et al. 2015). The statistical significance

by hypergeometric distribution was done with http://nemates.org/MA/progs/ overlap_stats.html webtool.

The Gene Ontology analysis was done using unique genes from HP1 datasets, by g:Profiler webtool using default options (Raudvere et al. 2019).

Aggregation plot of signal localization

All aggregation plots were generated using Seqplots, as a GUI application (Stempor and Ahringer 2016). Plots represent the average signal for every 10 bp bins of depicted histone marks or HP1 datasets over the midpoint position of significant peaks depicted.

Data Availability

Plasmids and strains can be requested from the authors. Supplemental files are available at FigShare. Supplementary File 1 contains the overlap between HP1 bound genes with tissue-specific expressed genes. Supplementary File 2 contains the overlap between tissue-specific genes bound by HP1 proteins. Datasets generated in this paper are available at deposited at Gene Expression Omnibus (GEO) accession GSE222056 (https://www.ncbi.nlm.nih.gov/geo/).

Acknowledgements

Very special thanks to Ildefonso Cases from the CABD Bioinformatics Facility for support with data analysis.

Funding

P.dl.C.R. was supported by contract BES-2017-081183 from the Spanish FPI program. Research was funded through grants from the Ministerio de Ciencia, Innovación y Universidades, which is part of Agencia Estatal de Investigación (AEI), the Fondo Europeo de Desarrollo Regional (FEDER) and the Consejería de Transformación Económica, Industria, Conocimiento y Universidades de la Junta de Andalucía (CEX2020-001088-M, PID 2019-104145GB-I00, PID2019-105069GB-I00, FEDER 2014–2020_UPO-1260918 and P20_00873).

Author contributions

P.dl.C.R., P.A and M.A.-S. designed the experiments; P.dl.C.R., carried out experiments, P.dl.C.R. analyzed the data; P.dl.C.R., P.A and M.A.-S. interpreted results and wrote the manuscript. All authors read, commented and approved the final manuscript.

Competing Interests

The authors declare that they have no conflict of interest.

REFERENCES

Ahringer, J., and S. M. Gasser. 2018. "Repressive Chromatin in caernorhabditis elegans: Establishment, Composition and Function." *Genetics* 208 (2): 491-511. https://doi.org/10.1534/genetics.117.300386. https://www.ncbi.nlm.nih.gov/pubmed/29378810.

Arraj, J. A., and M. G. Marinus. 1983. "Phenotypic reversal in dam mutants of Escherichia coli K-12 by a recombinant plasmid containing the dam+ gene." *J Bacteriol* 153 (1): 562-5. https://doi.org/10.1128/jb.153.1.562-565.1983. https://www.ncbi.nlm.nih.gov/pubmed/6336742.

Ashe, A., A. Sapetschnig, E. M. Weick, J. Mitchell, M. P. Bagijn, A. C. Cording, A. L. Doebley, L. D. Goldstein, N. J. Lehrbach, J. Le Pen, G. Pintacuda, A. Sakaguchi, P. Sarkies, S. Ahmed, and E. A. Miska. 2012. "piRNAs can trigger a multigenerational epigenetic memory in the germline of C. elegans." *Cell* 150 (1): 88-99. https://doi.org/10.1016/j.cell.2012.06.018. https://www.ncbi.nlm.nih.gov/pubmed/22738725.

Black, J. C., A. Allen, C. Van Rechem, E. Forbes, M. Longworth, K. Tschöp, C. Rinehart, J. Quiton, R. Walsh, A. Smallwood, N. J. Dyson, and J. R. Whetstine. 2010. "Conserved antagonism between JMJD2A/KDM4A and HP1γ during cell cycle progression." *Mol Cell* 40 (5): 736-48. https://doi.org/10.1016/j.molcel.2010.11.008. https://www.ncbi.nlm.nih.gov/pubmed/21145482.

Blazie, S. M., H. C. Geissel, H. Wilky, R. Joshi, J. Newbern, and M. Mangone. 2017. "Alternative Polyadenylation Directs Tissue-Specific miRNA Targeting in." *Genetics* 206 (2): 757-774. https://doi.org/10.1534/genetics.116.196774. https://www.ncbi.nlm.nih.gov/pubmed/28348061.

Cardoso, C., C. Couillault, C. Mignon-Ravix, A. Millet, J. J. Ewbank, M. Fontés, and N. Pujol. 2005. "XNP-1/ATR-X acts with RB, HP1 and the NuRD complex during larval development in C. elegans." *Dev Biol* 278 (1): 49-59. https://doi.org/10.1016/j.ydbio.2004.10.014. https://www.ncbi.nlm.nih.gov/pubmed/15649460.

Consortium, C. elegans Sequencing. 1998. "Genome sequence of the nematode C. elegans: a platform for investigating biology." *Science* 282 (5396): 2012-8. https://doi.org/10.1126/science.282.5396.2012. https://www.ncbi.nlm.nih.gov/pubmed/9851916.

Coustham, V., C. Bedet, K. Monier, S. Schott, M. Karali, and F. Palladino. 2006. "The C. elegans HP1 homologue HPL-2 and the LIN-13 zinc finger protein form a complex implicated in vulval development." *Dev Biol* 297 (2): 308-22. https://doi.org/10.1016/j.ydbio.2006.04.474. https://www.ncbi.nlm.nih.gov/pubmed/16890929.

Couteau, F., F. Guerry, F. Muller, and F. Palladino. 2002. "A heterochromatin protein 1 homologue in Caenorhabditis elegans acts in germline and vulval development." *EMBO Rep* 3 (3): 235-41. https://doi.org/10.1093/embo-reports/kvf051. https://www.ncbi.nlm.nih.gov/pubmed/11850401.

de la Cruz Ruiz, P., R. Romero-Bueno, and P. Askjaer. 2022. "Analysis of Nuclear Pore Complexes in Caenorhabditis elegans by Live Imaging and Functional Genomics." *Methods Mol Biol* 2502: 161-182. https://doi.org/10.1007/978-1-0716-2337-4_11. https://www.ncbi.nlm.nih.gov/pubmed/35412238.

Dobrzynska, A., P. Askjaer, and Y. Gruenbaum. 2016. "Lamin-Binding Proteins in Caenorhabditis elegans." *Methods Enzymol* 569: 455-83. https://doi.org/10.1016/bs.mie.2015.08.036. https://www.ncbi.nlm.nih.gov/pubmed/26778571.

Eissenberg, J. C. 2001. "Molecular biology of the chromo domain: an ancient chromatin module comes of age." *Gene* 275 (1): 19-29. https://doi.org/10.1016/s0378-1119(01)00628-x. https://www.ncbi.nlm.nih.gov/pubmed/11574148.

Eissenberg, J. C., T. C. James, D. M. Foster-Hartnett, T. Hartnett, V. Ngan, and S. C. Elgin. 1990. "Mutation in a heterochromatin-specific chromosomal protein is associated with suppression of position-effect variegation in Drosophila melanogaster." *Proc Natl Acad Sci U S A* 87 (24): 9923-7. https://doi.org/10.1073/pnas.87.24.9923. https://www.ncbi.nlm.nih.gov/pubmed/2124708.

Evans, K. J., N. Huang, P. Stempor, M. A. Chesney, T. A. Down, and J. Ahringer. 2016. "Stable." *Proc Natl Acad Sci U S A* 113 (45): E7020-E7029. https://doi.org/10.1073/pnas.1608162113. https://www.ncbi.nlm.nih.gov/pubmed/27791097.

Ferré, Q., G. Charbonnier, N. Sadouni, F. Lopez, Y. Kermezli, S. Spicuglia, C. Capponi, B. Ghattas, and D. Puthier. 2019. "OLOGRAM: Determining significance of total overlap length between genomic regions sets." *Bioinformatics*. https://doi.org/10.1093/bioinformatics/btz810. https://www.ncbi.nlm.nih.gov/pubmed/31688931.

Frøkjær-Jensen, C., M. W. Davis, M. Ailion, and E. M. Jorgensen. 2012. "Improved Mos1-mediated transgenesis in C. elegans." *Nat Methods* 9 (2): 117-8. https://doi.org/10.1038/nmeth.1865. https://www.ncbi.nlm.nih.gov/pubmed/22290181.

Garrigues, J. M., S. Sidoli, B. A. Garcia, and S. Strome. 2015. "Defining heterochromatin in C. elegans through genome-wide analysis of the heterochromatin protein 1 homolog HPL-2." *Genome Res* 25 (1): 76-88. https://doi.org/10.1101/gr.180489.114. https://www.ncbi.nlm.nih.gov/pubmed/25467431.

Gerstein, M. B., Z. J. Lu, E. L. Van Nostrand, C. Cheng, B. I. Arshinoff, T. Liu, K. Y. Yip, R. Robilotto, A. Rechtsteiner, K. Ikegami, P. Alves, A. Chateigner, M. Perry, M. Morris, R. K. Auerbach, X. Feng, J. Leng, A. Vielle, W. Niu, K. Rhrissorrakrai, A. Agarwal, R. P. Alexander, G. Barber, C. M. Brdlik, J. Brennan, J. J. Brouillet, A. Carr, M. S. Cheung, H. Clawson, S. Contrino, L. O. Dannenberg, A. F. Dernburg, A. Desai, L. Dick, A. C. Dosé, J. Du, T. Egelhofer, S. Ercan, G. Euskirchen, B. Ewing, E. A. Feingold, R. Gassmann, P. J. Good, P. Green, F. Gullier, M. Gutwein, M. S. Guyer, L. Habegger, T. Han, J. G. Henikoff, S. R. Henz, A. Hinrichs, H. Holster, T. Hyman, A. L. Iniguez, J. Janette, M. Jensen, M. Kato, W. J. Kent, E. Kephart, V. Khivansara, E. Khurana, J. K. Kim, P. Kolasinska-Zwierz, E. C. Lai, I. Latorre, A. Leahey, S. Lewis, P. Lloyd, L. Lochovsky, R. F. Lowdon, Y. Lubling, R. Lyne, M. MacCoss, S. D. Mackowiak, M. Mangone, S. McKay, D. Mecenas, G. Merrihew, D. M. Miller, A. Muroyama, J. I. Murray, S. L. Ooi, H. Pham, T. Phippen, E. A. Preston, N. Rajewsky, G. Rätsch, H. Rosenbaum, J. Rozowsky, K. Rutherford, P. Ruzanov, M. Sarov, R. Sasidharan, A. Sboner, P. Scheid, E. Segal, H. Shin, C. Shou, F. J. Slack, C. Slightam, R. Smith, W. C. Spencer, E. O. Stinson, S. Taing, T. Takasaki, D. Vafeados, K. Voronina, G. Wang, N. L. Washington, C.

M. Whittle, B. Wu, K. K. Yan, G. Zeller, Z. Zha, M. Zhong, X. Zhou, J. Ahringer, S. Strome, K. C. Gunsalus, G. Micklem, X. S. Liu, V. Reinke, S. K. Kim, L. W. Hillier, S. Henikoff, F. Piano, M. Snyder, L. Stein, J. D. Lieb, R. H. Waterston, and modENCODE Consortium. 2010. "Integrative analysis of the Caenorhabditis elegans genome by the modENCODE project." *Science* 330 (6012): 1775-87. https://doi.org/10.1126/science.1196914. https://www.ncbi.nlm.nih.gov/pubmed/21177976.

Gonzalez-Aguilera, C., K. Ikegami, C. Ayuso, A. de Luis, M. Iniguez, J. Cabello, J. D. Lieb, and P. Askjaer. 2014. "Genome-wide analysis links emerin to neuromuscular junction activity in Caenorhabditis elegans." *Genome Biol* 15 (2): R21. https://doi.org/10.1186/gb-2014-15-2-r21. https://www.ncbi.nlm.nih.gov/pubmed/24490688.

Grant, J., C. Verrill, V. Coustham, A. Arneodo, F. Palladino, K. Monier, and A. Khalil. 2010. "Perinuclear distribution of heterochromatin in developing C. elegans embryos." *Chromosome Res* 18 (8): 873-85. https://doi.org/10.1007/s10577-010-9175-2. https://www.ncbi.nlm.nih.gov/pubmed/21116703.

Gu, S. G., and A. Fire. 2010. "Partitioning the C. elegans genome by nucleosome modification, occupancy, and positioning." *Chromosoma* 119 (1): 73-87. https://doi.org/10.1007/s00412-009-0235-3. https://www.ncbi.nlm.nih.gov/pubmed/19705140.

Gómez-Saldivar, G., J. Osuna-Luque, J. I. Semple, D. A. Glauser, S. Jarriault, and P. Meister. 2020. "Tissue-Specific Transcription Footprinting Using RNA PoI DamID (RAPID) in." *Genetics* 216 (4): 931-945. https://doi.org/10.1534/genetics.120.303774. https://www.ncbi.nlm.nih.gov/pubmed/33037050.

Haenni, S., Z. Ji, M. Hoque, N. Rust, H. Sharpe, R. Eberhard, C. Browne, M. O. Hengartner, J. Mellor, B. Tian, and A. Furger. 2012. "Analysis of C. elegans intestinal gene expression and polyadenylation by fluorescence-activated nuclei sorting and 3'-end-seq." *Nucleic Acids Res* 40 (13): 6304-18. https://doi.org/10.1093/nar/gks282. https://www.ncbi.nlm.nih.gov/pubmed/22467213.

Heberle, H., G. V. Meirelles, F. R. da Silva, G. P. Telles, and R. Minghim. 2015. "InteractiVenn: a web-based tool for the analysis of sets through Venn diagrams." *BMC Bioinformatics* 16 (1): 169. https://doi.org/10.1186/s12859-015-0611-3. https://www.ncbi.nlm.nih.gov/pubmed/25994840.

Horn, M., C. Geisen, L. Cermak, B. Becker, S. Nakamura, C. Klein, M. Pagano, and A. Antebi. 2014. "DRE-1/FBXO11-dependent degradation of BLMP-1/BLIMP-1 governs C. elegans developmental timing and maturation." *Dev Cell* 28 (6): 697-710. https://doi.org/10.1016/j.devcel.2014.01.028. https://www.ncbi.nlm.nih.gov/pubmed/24613396.

Huang, T. F., C. Y. Cho, Y. T. Cheng, J. W. Huang, Y. Z. Wu, A. Y. Yeh, K. Nishiwaki, S. C. Chang, and Y. C. Wu. 2014. "BLMP-1/Blimp-1 regulates the spatiotemporal cell migration pattern in C. elegans." *PLoS Genet* 10 (6): e1004428. https://doi.org/10.1371/journal.pgen.1004428. https://www.ncbi.nlm.nih.gov/pubmed/24968003.

Jedrusik-Bode, M. 2013. "Histone H1 and heterochromatin protein 1 (HP1) regulate specific gene expression and not global transcription." *Worm* 2 (2): e23703.

https://doi.org/10.4161/worm.23703.
https://www.ncbi.nlm.nih.gov/pubmed/24058872.

Kaletsky, R., V. Yao, A. Williams, A. M. Runnels, A. Tadych, S. Zhou, O. G. Troyanskaya, and C. T. Murphy. 2018. "Transcriptome analysis of adult Caenorhabditis elegans cells reveals tissue-specific gene and isoform expression." *PLoS Genet* 14 (8): e1007559. https://doi.org/10.1371/journal.pgen.1007559. https://www.ncbi.nlm.nih.gov/pubmed/30096138.

Katsanos, D., M. Ferrando-Marco, I. Razzaq, G. Aughey, T. D. Southall, and M. Barkoulas. 2021. "Gene expression profiling of epidermal cell types in C. elegans using Targeted DamID." *Development* 148 (17). https://doi.org/10.1242/dev.199452. https://www.ncbi.nlm.nih.gov/pubmed/34397094.

Keller, A. D., and T. Maniatis. 1991. "Identification and characterization of a novel repressor of beta-interferon gene expression." *Genes Dev* 5 (5): 868-79. https://doi.org/10.1101/gad.5.5.868. https://www.ncbi.nlm.nih.gov/pubmed/1851123.

Khan, A., and A. Mathelier. 2017. "Intervene: a tool for intersection and visualization of multiple gene or genomic region sets." *BMC Bioinformatics* 18 (1): 287. https://doi.org/10.1186/s12859-017-1708-7. https://www.ncbi.nlm.nih.gov/pubmed/28569135.

Koester-Eiserfunke, N., and W. Fischle. 2011. "H3K9me2/3 binding of the MBT domain protein LIN-61 is essential for Caenorhabditis elegans vulva development." *PLoS Genet* 7 (3): e1002017. https://doi.org/10.1371/journal.pgen.1002017. https://www.ncbi.nlm.nih.gov/pubmed/21437264.

Kondili, M., A. Fust, J. Preussner, C. Kuenne, T. Braun, and M. Looso. 2017. "UROPA: a tool for Universal RObust Peak Annotation." *Sci Rep* 7 (1): 2593. https://doi.org/10.1038/s41598-017-02464-y. https://www.ncbi.nlm.nih.gov/pubmed/28572580.

Kozlowski, L., S. Garvis, C. Bedet, and F. Palladino. 2014. "The Caenorhabditis elegans HP1 family protein HPL-2 maintains ER homeostasis through the UPR and hormesis." *Proc Natl Acad Sci U S A* 111 (16): 5956-61. https://doi.org/10.1073/pnas.1321698111. https://www.ncbi.nlm.nih.gov/pubmed/24715729.

Langmead, B., and S. L. Salzberg. 2012. "Fast gapped-read alignment with Bowtie 2." *Nat Methods* 9 (4): 357-9. https://doi.org/10.1038/nmeth.1923. https://www.ncbi.nlm.nih.gov/pubmed/22388286.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R. Durbin, and 1000 Genome Project Data Processing Subgroup. 2009. "The Sequence Alignment/Map format and SAMtools." *Bioinformatics* 25 (16): 2078-9. https://doi.org/10.1093/bioinformatics/btp352. https://www.ncbi.nlm.nih.gov/pubmed/19505943.

Liu, T., A. Rechtsteiner, T. A. Egelhofer, A. Vielle, I. Latorre, M. S. Cheung, S. Ercan, K. Ikegami, M. Jensen, P. Kolasinska-Zwierz, H. Rosenbaum, H. Shin, S. Taing, T. Takasaki, A. L. Iniguez, A. Desai, A. F. Dernburg, H. Kimura, J. D. Lieb, J. Ahringer, S. Strome, and X. S. Liu. 2011. "Broad chromosomal domains of histone modification patterns in C. elegans." *Genome Res* 21 (2): 227-36. https://doi.org/10.1101/gr.115519.110. https://www.ncbi.nlm.nih.gov/pubmed/21177964.

Marshall, O. J., and A. H. Brand. 2015. "damidseq_pipeline: an automated pipeline for processing DamID sequencing datasets." *Bioinformatics* 31 (20): 3371-3. https://doi.org/10.1093/bioinformatics/btv386. https://www.ncbi.nlm.nih.gov/pubmed/26112292.

McMurchy, A. N., P. Stempor, T. Gaarenstroom, B. Wysolmerski, Y. Dong, D. Aussianikava, A. Appert, N. Huang, P. Kolasinska-Zwierz, A. Sapetschnig, E. A. Miska, and J. Ahringer. 2017. "A team of heterochromatin factors collaborates with small RNA pathways to combat repetitive elements and germline stress." *Elife* 6. https://doi.org/10.7554/eLife.21666. https://www.ncbi.nlm.nih.gov/pubmed/28294943.

Meister, P., S. Schott, C. Bedet, Y. Xiao, S. Rohner, S. Bodennec, B. Hudry, L. Molin, F. Solari, S. M. Gasser, and F. Palladino. 2011. "Caenorhabditis elegans Heterochromatin protein 1 (HPL-2) links developmental plasticity, longevity and lipid metabolism." *Genome Biol* 12 (12): R123. https://doi.org/10.1186/gb-2011-12-12-r123. https://www.ncbi.nlm.nih.gov/pubmed/22185090.

Niu, W., Z. J. Lu, M. Zhong, M. Sarov, J. I. Murray, C. M. Brdlik, J. Janette, C. Chen, P. Alves, E. Preston, C. Slightham, L. Jiang, A. A. Hyman, S. K. Kim, R. H. Waterston, M. Gerstein, M. Snyder, and V. Reinke. 2011. "Diverse transcription factor binding features revealed by genome-wide ChIP-seq in C. elegans." *Genome Res* 21 (2): 245-54. https://doi.org/10.1101/gr.114587.110. https://www.ncbi.nlm.nih.gov/pubmed/21177963.

Padeken, J., S. P. Methot, and S. M. Gasser. 2022. "Establishment of H3K9-methylated heterochromatin and its functions in tissue differentiation and maintenance." *Nat Rev Mol Cell Biol* 23 (9): 623-640. https://doi.org/10.1038/s41580-022-00483-w. https://www.ncbi.nlm.nih.gov/pubmed/35562425.

Patel, T., and O. Hobert. 2017. "Coordinated control of terminal differentiation and restriction of cellular plasticity." *Elife* 6. https://doi.org/10.7554/eLife.24100. https://www.ncbi.nlm.nih.gov/pubmed/28422646.

Piunti, A., and A. Shilatifard. 2016. "Epigenetic balance of gene expression by Polycomb and COMPASS families." *Science* 352 (6290): aad9780. https://doi.org/10.1126/science.aad9780. https://www.ncbi.nlm.nih.gov/pubmed/27257261.

Quinlan, A. R., and I. M. Hall. 2010. "BEDTools: a flexible suite of utilities for comparing genomic features." *Bioinformatics* 26 (6): 841-2. https://doi.org/10.1093/bioinformatics/btq033. https://www.ncbi.nlm.nih.gov/pubmed/20110278.

Ramírez, F., D. P. Ryan, B. Grüning, V. Bhardwaj, F. Kilpert, A. S. Richter, S. Heyne, F. Dündar, and T. Manke. 2016. "deepTools2: a next generation web server for deep-sequencing data analysis." *Nucleic Acids Res* 44 (W1): W160-5. https://doi.org/10.1093/nar/gkw257. https://www.ncbi.nlm.nih.gov/pubmed/27079975.

Raudvere, U., L. Kolberg, I. Kuzmin, T. Arak, P. Adler, H. Peterson, and J. Vilo. 2019. "g:Profiler: a web server for functional enrichment analysis and conversions of gene lists (2019 update)." *Nucleic Acids Res* 47 (W1): W191-W198. https://doi.org/10.1093/nar/gkz369. https://www.ncbi.nlm.nih.gov/pubmed/31066453.

Schoelz, J. M., J. X. Feng, and N. C. Riddle. 2021. "The Drosophila HP1 family is associated with active gene expression across chromatin contexts." *Genetics* 219 (1). https://doi.org/10.1093/genetics/iyab108. https://www.ncbi.nlm.nih.gov/pubmed/34849911.

Schott, S., V. Coustham, T. Simonet, C. Bedet, and F. Palladino. 2006. "Unique and redundant functions of C. elegans HP1 proteins in post-embryonic development." *Dev Biol* 298 (1): 176-87. https://doi.org/10.1016/j.ydbio.2006.06.039. https://www.ncbi.nlm.nih.gov/pubmed/16905130.

Schott, S., F. Ramos, V. Coustham, and F. Palladino. 2009. "HPL-2/HP1 prevents inappropriate vulval induction in Caenorhabditis elegans by acting in both HYP7 and vulval precursor cells." *Genetics* 181 (2): 797-801. https://doi.org/10.1534/genetics.108.089276. https://www.ncbi.nlm.nih.gov/pubmed/19064713.

Shaffer, A. L., K. I. Lin, T. C. Kuo, X. Yu, E. M. Hurt, A. Rosenwald, J. M. Giltnane, L. Yang, H. Zhao, K. Calame, and L. M. Staudt. 2002. "Blimp-1 orchestrates plasma cell differentiation by extinguishing the mature B cell gene expression program." *Immunity* 17 (1): 51-62. https://doi.org/10.1016/s1074-7613(02)00335-7. https://www.ncbi.nlm.nih.gov/pubmed/12150891.

Sharma, R., D. Ritler, and P. Meister. 2016. "Tools for DNA adenine methyltransferase identification analysis of nuclear organization during C. elegans development." *Genesis* 54 (4): 151-9. https://doi.org/10.1002/dvg.22925. https://www.ncbi.nlm.nih.gov/pubmed/26845390.

Simonet, T., R. Dulermo, S. Schott, and F. Palladino. 2007. "Antagonistic functions of SET-2/SET1 and HPL/HP1 proteins in C. elegans development." *Dev Biol* 312 (1): 367-83. https://doi.org/10.1016/j.ydbio.2007.09.035. https://www.ncbi.nlm.nih.gov/pubmed/17967446.

Stec, N., K. Doerfel, K. Hills-Muckey, V. M. Ettorre, S. Ercan, W. Keil, and C. M. Hammell. 2021. "An Epigenetic Priming Mechanism Mediated by Nutrient Sensing Regulates Transcriptional Output during C. elegans Development." *Curr Biol* 31 (4): 809-826.e6. https://doi.org/10.1016/j.cub.2020.11.060. https://www.ncbi.nlm.nih.gov/pubmed/33357451.

Steffen, P. A., and L. Ringrose. 2014. "What are memories made of? How Polycomb and Trithorax proteins mediate epigenetic memory." *Nat Rev Mol Cell Biol* 15 (5): 340-56. https://doi.org/10.1038/nrm3789. https://www.ncbi.nlm.nih.gov/pubmed/24755934.

Stempor, P., and J. Ahringer. 2016. "SeqPlots - Interactive software for exploratory data analyses, pattern discovery and visualization in genomics." *Wellcome Open Res* 1: 14. https://doi.org/10.12688/wellcomeopenres.10004.1. https://www.ncbi.nlm.nih.gov/pubmed/27918597.

Stiernagle, T. 2006. "Maintenance of C. elegans." *WormBook*: 1-11. https://doi.org/10.1895/wormbook.1.101.1. https://www.ncbi.nlm.nih.gov/pubmed/18050451.

Studencka, M., A. Konzer, G. Moneron, D. Wenzel, L. Opitz, G. Salinas-Riester, C. Bedet, M. Krüger, S. W. Hell, J. R. Wisniewski, H. Schmidt, F. Palladino, E. Schulze, and M. Jedrusik-Bode. 2012. "Novel roles of Caenorhabditis elegans heterochromatin protein HP1 and linker histone in the regulation of innate immune gene expression."

Mol Cell Biol 32 (2): 251-65. https://doi.org/10.1128/MCB.05229-11. https://www.ncbi.nlm.nih.gov/pubmed/22083954.

Studencka, M., R. Wesołowski, L. Opitz, G. Salinas-Riester, J. R. Wisniewski, and M. Jedrusik-Bode. 2012. "Transcriptional repression of Hox genes by C. elegans HP1/HPL and H1/HIS-24." *PLoS Genet* 8 (9): e1002940. https://doi.org/10.1371/journal.pgen.1002940. https://www.ncbi.nlm.nih.gov/pubmed/23028351.

Su, S. T., H. Y. Ying, Y. K. Chiu, F. R. Lin, M. Y. Chen, and K. I. Lin. 2009. "Involvement of histone demethylase LSD1 in Blimp-1-mediated gene repression during plasma cell differentiation." *Mol Cell Biol* 29 (6): 1421-31. https://doi.org/10.1128/MCB.01158-08. https://www.ncbi.nlm.nih.gov/pubmed/19124609.

Tarailo-Graovac, M., and N. Chen. 2009. "Using RepeatMasker to identify repetitive elements in genomic sequences." *Curr Protoc Bioinformatics* Chapter 4: 4.10.1-4.10.14. https://doi.org/10.1002/0471250953.bi0410s25. https://www.ncbi.nlm.nih.gov/pubmed/19274634.

Towbin, B. D., C. Gonzalez-Aguilera, R. Sack, D. Gaidatzis, V. Kalck, P. Meister, P. Askjaer, and S. M. Gasser. 2012. "Step-wise methylation of histone H3K9 positions heterochromatin at the nuclear periphery." *Cell* 150 (5): 934-47. https://doi.org/10.1016/j.cell.2012.06.051. https://www.ncbi.nlm.nih.gov/pubmed/22939621.

Wang, S., D. H. Meyer, and B. Schumacher. 2022. "Inheritance of paternal DNA damage by histone-mediated repair restriction." *Nature*. https://doi.org/10.1038/s41586-022-05544-w. https://www.ncbi.nlm.nih.gov/pubmed/36544019.

Yang, J., H. T. Fong, Z. Xie, J. W. Tan, and T. Inoue. 2015. "Direct and positive regulation of Caenorhabditis elegans bed-3 by PRDM1/BLIMP1 ortholog BLMP-1." *Biochim Biophys Acta* 1849 (9): 1229-36. https://doi.org/10.1016/j.bbagrm.2015.07.012. https://www.ncbi.nlm.nih.gov/pubmed/26234645.

Yilmaz, L. S., X. Li, S. Nanda, B. Fox, F. Schroeder, and A. J. Walhout. 2020. "Modeling tissue-relevant Caenorhabditis elegans metabolism at network, pathway, reaction, and metabolite levels." *Mol Syst Biol* 16 (10): e9649. https://doi.org/10.15252/msb.20209649. https://www.ncbi.nlm.nih.gov/pubmed/33022146.

Zeller, P., J. Padeken, R. van Schendel, V. Kalck, M. Tijsterman, and S. M. Gasser. 2016. "Histone H3K9 methylation is dispensable for Caenorhabditis elegans development but suppresses RNA:DNA hybrid-associated repeat instability." *Nat Genet* 48 (11): 1385-1395. https://doi.org/10.1038/ng.3672. https://www.ncbi.nlm.nih.gov/pubmed/27668659.

Zerbino, D. R., N. Johnson, T. Juettemann, S. P. Wilder, and P. Flicek. 2014. "WiggleTools: parallel processing of large collections of genome-wide datasets for visualization and statistical analysis." *Bioinformatics* 30 (7): 1008-9. https://doi.org/10.1093/bioinformatics/btt737. https://www.ncbi.nlm.nih.gov/pubmed/24363377.

Zhao, H., Z. Sun, J. Wang, H. Huang, J. P. Kocher, and L. Wang. 2014. "CrossMap: a versatile tool for coordinate conversion between genome assemblies." *Bioinformatics* 30 (7): 1006-7. https://doi.org/10.1093/bioinformatics/btt730. https://www.ncbi.nlm.nih.gov/pubmed/24351709.

Figure 1. Tissue-specific chromatin binding of HPL-1 and HPL-2 and their association to histone marks.
(A) Boxplot showing HP1 association profile in all autosomes based on arms with and without meiotic pairing centers at 100 kb bin resolution in hypodermal and intestinal tissues. p-values from Wilcoxon rank sum test with continuity correction are indicated. PC: pairing center. (B) Example of average signal tracks over 10 kb regions in chromosome IV for HP1 proteins in hypodermal and intestinal tissues. Light blue region represents chromosome center, while flanking regions correspond to left and right arms. Snapshot from IGV viewer.
(C) Heatmap showing pairwise Pearson correlation coefficients between indicated datasets at 25 bp window size. (D) Aggregation plot depicting the genome-wide average signal tracks for H3K9me1 dataset over HPL-2 hypodermal and intestinal peaks. Center position represents the midpoint location of peaks. Regions up to 3 kb upstream and 3 kb down-stream of the midpoint in 10 bp bins are shown.

Figure 2. Tissue-specific binding of HP1 proteins and association with gene expression. (A) Venn diagram depicting the overlapping DNA regions bound by HP1 proteins in hypodermis and intestine. The intersection was performed using 10% of fraction reciprocal between genomic regions. Fisher's exact test *p*-values are represented. (B) Pie charts showing the percentages of peaks inside various genomic features. Peak assignment was done to the nearest gene whose center coordinate was within 3 kb upstream and 3 kb downstream of the TSS and TES, respectively, of a gene. Ratios above pie charts show the number of peaks assigned to genes relative to the total number of peaks. TSS: Transcription start site; TES: transcription end site. (C) Venn diagram depicting the overlap between HPL-2-bound genes in the hypodermis and genes robustly expressed in this tissue. (D) Venn diagram depicting the overlaps between HPL-1 and HPL-2-bound genes in the intestine and genes expressed in this tissue. Exact hypergeometric probability representation factor and *p*-values are shown.

A

B

C

Figure 3. Differential tissue-specific binding of HP1 proteins. (A) Venn diagram depicting the overlapping genes bound by HP1 proteins in the different tissues. Exact hypergeometric probability representation factor and p values are shown. (B) GO terms associated to Biological process category of genes bound by HPL-2 in the intestine. Number of intersected genes are depicted at the end of the bars. (C) Aggregation plot depicting the Z-score of aggregated average signal of HP1 proteins over 3 kb upstream and 3 kb downstream of BLMP-1 peaks' midpoint position.

Figure 4. Association of HP1 proteins to repetitive elements. (A) Heatmap showing the correlation between significant peaks bound by HP1 proteins and repetitive elements. (B) Stacked plot depicting the proportion of repetitive element types bound by HPL-2 datasets (See Supplementary Table 5).

Supplementary Figure 1. Tissue-specific chromatin binding of HPL-1 and HPL-2 and their association to histone marks. (A) Heatmap depicting the proportion of HP1 peaks distributed by chromosomes autosomes distinguishing between arms and center. Chromosomes autosomes coordinates based on (Garrigues et al. 2015). (B-D) Aggregation plots depicting the average signal tracks for H3K9me2 (A), H3K27me2 (B) and H3K9me3 (C) histone datasets over HP1 peaks. Center position represents the midpoint location of peaks. Regions up to 3 kb upstream and 3 kb downstream of the midpoint in 10 bp bins are shown.

A

Overlapped peaks with HPL-2 ChIP-seq data

B

Shuffled   True

Query : HPL-2_chip.bed
Total nb. of intersections per region type

p=5.1e-240
p=9.5e-135
p=3.4e-30

Number of intersections

hpl1_int   hpl2_int   hpl2_hyp

C

Shuffled   True

Query : HPL-2_chip.bed
Total overlap length per region type

p=2.1e-166
p=3.1e-249
p=2.1e-31

Nb. of overlapping base pairs

hpl1_int   hpl2_int   hpl2_hyp

Supplementary Figure 2. Overlapping of HP1 DamID and HPL-2 ChIP datasets. (A) Stacked plot depicting the number of overlapped peaks between HP1 datasets indicated and a published HPL-2 ChIP dataset. Numbers inside orange bars represent the proportion in percentage of overlapped peaks respect of total HP1 peaks number. (B-C) Statistic for overlapped peaks between HP1 and HPL-2 ChIP-seq using a Monte Carlo simulation, based on mathematical distributions of region (B) and inter-region lengths (C), according to negative binomial model of the total overlap length. Hpl-2_hyp: HPL-2 hypdermis; hpl-2_int: HPL-2 intestine; hpl-1_int: HPL-1 intestine.

Supplementary Figure 3. Differential tissue-specific binding of HP1proteins. (A-B) GO terms associated to cellular component category of unique genes bound by HPL-2 in the hypodermal tissue (A) and by HPL-1 in the intestine (B). Number of intersected genes depicted in right side bars. (C) Heatmap depicting proportion of overlapped peaks between all dataset indicated. (D) Heatmap describing chromosome autosome location of overlapped peaks between BLMP-1 and HPL-1. (E) Number of intersected peaks between HP1 datasets and BLMP-1 assessed by Montecarlo Simulation statistical test. (F) Venn diagram indicating the number of overlapping genes between HPL-1 in the intestine and BLMP-1. Exact hypergeometric probability representation factor and *p* values are shown.

Supplementary Tables

	HPL-2 ChIP-seq (23530)
HPL-2 hypodermis	# Number of query intervals: 23530 # Number of db intervals: 1572 # Number of overlaps: 1226 # Number of possible intervals (estimated): 65530 # phyper(1226 - 1. 23530. 65530 - 23530. 1572. lower.tail=F) # Contingency Table Of Counts #____ # | in -b | not in -b | # in -a | 1226 | 22304 | # not in -a | 346 | 41654 | #____ # p-values for fisher's exact test left right two-tail ratio 1 4.0067e-261 4.6837e-261 6.617
HPL-2 intestine	# Number of query intervals: 23530 # Number of db intervals: 1665 # Number of overlaps: 1409 # Number of possible intervals (estimated): 81142 # phyper(1409 - 1. 23530. 81142 - 23530. 1665. lower.tail=F) # Contingency Table Of Counts #____ # | in -b | not in -b | # in -a | 1409 | 22121 | # not in -a | 256 | 57356 | #____ # p-values for fisher's exact test left right two-tail ratio 1 0 0 14.271
HPL-1 intestine	# Number of query intervals: 23530 # Number of db intervals: 1346 # Number of overlaps: 512 # Number of possible intervals (estimated): 88730 # phyper(512 - 1. 23530. 88730 - 23530. 1346. lower.tail=F) # Contingency Table Of Counts #____ # | in -b | not in -b | # in -a | 512 | 23018 | # not in -a | 834 | 64366 | #____ # p-values for fisher's exact test left right two-tail ratio 1 9.3088e-21 1.4496e-20 1.717

Supplementary Table 1. Number of intersections with HPL-2 ChIP-seq genomic regions and fisher's exact test p-values associated.

	Hypodermal well expressed genes (2264)	Intestinal well expressed genes (5890)	HPL-2 intestine (922)	HPL-1 intestine (739)
HPL-2 hypodermis (845)	Representation factor: 0.7 p < 0.001	-		
HPL-2 intestine (922)	-	Representation factor: 1.0 p < 0.383	-	Representation factor: 5.9 p < 1.369e-100
HPL-1 intestine (739)	-	Representation factor: 0.7 p < 6.862e-07	Representation factor: 5.9 p < 1.369e-100	-

Supplementary Table 2. Hypergeometric statistical test between gene sets depicted in the table.

Samples (significant peaks)	**Number of intersection with BLMP-1 peaks (5898)**	Fisher's exact test p-values
HPL-1 intestine (1346)	169	two-tail ratio 1.8565e-09 1.701
Samples (significant peaks)	**Number of intersection with L4 BLMP-1 peaks (549)**	Fisher's exact test p-values
HPL-1 intestine (1346)	26	two-tail ratio 0.013515 1.701
HPL-2 intestine (1665)	34	two-tail ratio 0.0041704 1.726
HPL-2 hypodermis (1572)	14	two-tail ratio 0.097535 0.630

Supplementary Table 3. Number of overlapped peaks between HP1 and BLMP-1 ChIP-seq data and Fisher´s exact test p-value.

Samples (significant peaks)	**Number of intersection with Repetitive elements (61527)**	Fisher's exact test p-values
HPL-2 hypodermis (1572)	1650	two-tail ratio 2.7634e-159 inf
HPL-2 intestine (1665)	1112	two-tail ratio 0.00012955 1.222
HPL-1 intestine (1346)	567	two-tail ratio 1.1638e-23 0.575

Supplementary Table 4. Number of overlapped regions and fisher's exact test p-values between HP1 significant peaks and repetitive elements coordinates annotation from UCSC based on repeat masker (ce11).

HPL-2 hypodermis		
Family	Type	Number of repeats
CELE14B	DNA	282
CELE42	DNA	152
CELE14A	DNA	92
CELE1	DNA	75
PALTA5_CE	DNA	61
GA-rich	Low_complexity	48
A-rich	Low_complexity	46
CeRep5	Unknown	44
TIR9TA1B_CE	DNA	40
CELE2	DNA	38
TIR9TA1_CE	DNA	36
Helitron2_CE	RC	32
MINISAT1_CE	Satellite	30
HelitronY1A_CE	RC	27
HAT1_CE	DNA	25
CELE45	SINE	24
CELE4	DNA	23
RC123	Satellite	21
PALTTAA2_CE	DNA	20
CeRep3	Unknown	18
HPL-2 intestine		
CELE14B	DNA	193
CELE42	DNA	112
CELE14A	DNA	58
TIR9TA1B_CE	DNA	48
CELE45	SINE	42
TIR9TA1_CE	DNA	40
PALTA5_CE	DNA	32
GA-rich	Low_complexity	28
A-rich	Low_complexity	26
MINISAT1_CE	Satellite	26
PALTTAA3_CE	DNA	12
CeRep5	Unknown	12
CER12-I_CE	LTR	12
PALTTAA1_CE	DNA	11
Vingi-2_CE	LINE	11
LR9A	Unknown	11
HelitronY1_CE	RC	10
CER11-I_CE	LTR	9
TIR9TA1C_CE	DNA	8
CELE4	DNA	8

Supplementary Table 5. Top 20 overlapped regions between HP1 and repetitive elements.

STRAIN	DESCRIPTION	GENOTYPE	NAME	REFERENCE
BN548	FLP-Frt strain carrying P*dpy-7*::FLP transgene on chr IV.	bqSi548[pBN266(unc-119(+) P*dpy-7*::FLP)] IV; unc-119(ed3) III	-	(Muñoz-Jimenez 2019)
BN577	Visual marker in pharynx - P*myo-2*::GFP transgene on chr IV.	bqSi577[pBN306(unc-119(+) P*myo-2*::GFP)] IV; may carry unc-119(ed3) III	-	(Muñoz-Jiménez et al. 2017)
EG4322	Strain used for MosScl into chromosome II.	ttTi5605 II; unc-119(ed9) III.	EG4322	CGC
BN1103	Strain for tissue-specific HPL-1 DamID.	bqSi1103[pBN494(unc-119(+) P*hsp16.41*>mCh::*his-58*>dam::*hpl-1*)] II. May carry unc-119(ed9) III.	-	This publication. Made by injection of EG4322 and pBN494 plasmid.
BN1105	Strain for tissue-specific HPL-1 DamID co-expressing GFP in pharynx.	bqSi1103[pBN494(unc-119(+) P*hsp16.41*>mCh::*his-58*>dam::*hpl-1*)] II; bqSi577[pBN306(unc-119(+) P*myo-2*::GFP)] IV. May carry unc-119(ed9) III.	-	Made by crossing BN1103 and BN577
BN1051	Strain for control DamID in intestine.	bqSi447[pBN181(unc-119(+) hsp16.41p>mCh::his-58>gfp::dam)] II; bqSi997[pBN472(unc-119(+) nhx-2p::FLP::SL2::mNG)] IV; may carry unc-119(ed3) or unc-119(ed9) III.	-	Peter Askjaer (unpublished) Made by crossing BN448 and BN998.
BN561	Strain for control DamID in hypodermis.	bqSi447[pBN181(unc-119(+) Phsp16.41::FRT::mCh::his-58::FRT::gfp::dam)] II; bqSi548[pBN266(unc-119(+) Pdpy-7::FLP_D5)] IV. May carry unc-119(ed9) III.	-	Peter Askjaer (unpublished). Made by crossing BN550 and BN448.
BN550	Strain for control DamID in hypodermis.	FLP-Frt strain carrying Phsp16.41_FRT_mCh_his-58_FRT_gfp_his-58 on chr II and Pdpy-7::FLP-D5 on chr IV.		(Muñoz-Jimenez 2019)
BN448	Control strain for tissue-specific DamID	bqSi447[pBN181(unc−119(+).Phsp16.41::FrT::mch::his−58::FrT::gfp::dam)] II;		(Muñoz-Jimenez 2019)
BN1104	Strain for tissue-specific HPL-2 DamID.	bqSi1104[pBN495(unc-119(+) hsp16.41p>mCh::his-58>dam::hpl-2)] II. May carry unc-119(ed9) III.	-	This publication. Made by injection of EG4322 with pBN495 plasmid
BN1106	Strain for tissue-specific HPL-2 DamID co-expressing GFP in pharynx.	bqSi1104[pBN495(unc-119(+) hsp16.41p>mCh::his-58>dam::hpl-2)] II; bqSi577[pBN306(unc-119(+) myo-2p::GFP)] IV. May carry unc-119(ed9) III.	-	This thesis. Made by crossing BN1104 and BN577.

MRS578/BN1108	Strain for tissue-specific HPL-2 DamID in the hypodermis.	bqSi1104[hsp16.41p>mCh::his-58>dam::hpl-2] II; bqSi548[dpy-7p::FLP] IV	-	This publication. Made by crossing BN1104 and BN548.
BN998	FLP-Frt strain carrying P*nhx-2* controlling expression of FLP D5::SL2::mNG transgene on chr IV and visible marker on chr II (Plmn-1::mCh::his-58).	bqSi189[pBN13(unc-119(+) lmn-1p::mCherry::his-58)] II; bqSi997[pBN472(unc-119(+) nhx-2p::FLP_D5::SL2::mNG)] IV. May carry unc-119(ed3) or unc-119(ed9) III.	-	(Cabianca et al. 2019)
MRS579/BN1109	Strain for tissue-specific HPL-1 DamID in the intestine.	bqSi1104[pBN494(unc-119(+) P*hsp16.41*>mCh::*his-58*>dam::hpl-1)] II; bqSi997[pBN472(unc-119(+) P*nhx-2*::FLP::SL2::mNG)] IV. May carry unc-119(ed9) III.	-	This publication. Made by crossing BN1103 and BN998.
MRS580/BN1110	Strain for tissue-specific HPL-2 DamID in the intestine.	bqSi1103[pBN495(unc-119(+) P*hsp16.41*>mCh::*his-58*>dam::*hpl-2*)] II; bqSi997[pBN472(unc-119(+) nhx-2p::FLP::SL2::mNG)] IV. May carry *unc-119(ed9)* III.	-	This publication. Made by crossing BN1104 and BN998.
GM119	*E.coli* dam-3 dcm-6 knockout strain	(Arraj and Marinus 1983)		

Supplementary Table 6. List of *C.elegans* and bacterial strains used.

Plasmid name	Description	Reference
pBN494	*Phsp-16.41*::FRT::mCh::*his-58*::FRT::*hpl-1*::Dam Chr II	This publication
pBN495	P*hsp-16.41*::FRT:::mCh::*his-58*::FRT::hpl-2::Dam Chr II	This publication

Supplementary Table 7. List of plasmids used.

Sample	Raw reads	Cut reads	Mapped reads	GATC reads (% from Raw reads)
GFP::Dam hyp (BN561) I	28.277.140	27.192.343	11.584.959	11.548.079 (40.84%)
GFP::Dam hyp (BN561) II	27.363.912	26.009.374	12.883.562	12.859.165 (47%)
GFP::Dam int (BN1051) I	18.066.270	7.418.519	3.660.999	3.655.161 (20.23%)
GFP::Dam int (BN1051) II	4.573.067	2.080.559	1.338.144	1.336.648 (29.23%)
Dam::*hpl-2* hyp (MRS578) I	14.194.267	6.551.959	4.162.172	4.154.179 (29.27%)
Dam::*hpl-2* hyp (MRS578) II	8.507.806	3.473.687	1.594.499	1.562.681 (18.37%)
Dam::*hpl-2* int (MRS580) I	18.337.259	8.265.167	4.253.333	4.232.349 (23.08%)
Dam::*hpl-2* int (MRS580) II	1.583.607	747.426	527.174	526.083 (33.22%)
Dam::*hpl-1* int (MRS579) I	49.178.283	46.255.827	19.451.391	19.296.962 (39.24%)
Dam::*hpl-1* int (MRS579) II	8.061.225	3.861.200	696.028	694.399 (8.61%)

Supplementary Table 8. Quality controls metrics of sequencing libraries from DamID-seq data. Abbreviations hyp: hypodermis; int:intestine

Sample name	Dataset (GEO/ modENCODE)	Strain	Larval stage	Antibody	Temperature (°C)
ChIP dataset					
HPL-2 ChIP-seq	GSE49749/modENCODE_5976	N2	L3	HPL-2_SDQ2340	20
Histones datasets					
H3K9me1	GSE25353	N2	L3 (mixed Male and Hermaphrodite population)	AB9045	20
H3K9me2	GSE49728	N2	L3 (mixed Male and Hermaphrodite population)	HK00008	20
H3K9me3	GSE49719	N2	L3 (mixed Male and Hermaphrodite population)	AB8898	20
H3K27me3	GSE49724	N2	L3 (mixed Male and Hermaphrodite population)	UP07449	20
BLMP-1	GSE25803/**modENCODE_2612**	OP109	L1	GFP::3xFlag	20

Supplementary Table 9. Description and references of Dataset used for comparison.