Mechanistic insights into the activation of bacterial ExoY nucleotidyl cyclase toxins targeting actin-profilin

Magda Teixeira-Nunes[1], Pascal Retailleau[2], Dorothée Raoux-Barbot[3], Martine Comisso[1], Anani Amegan Missinou[1], Christophe Velours[1,#], Stéphane Plancqueel[1], Daniel Ladant[3], Undine Mechold[3], Louis Renault[1*]

[1] Université Paris-Saclay, CEA, CNRS, Institute for Integrative Biology of the Cell (I2BC), 91198, Gif-sur-Yvette, France

[2] ICSN, CNRS, Université Paris Saclay, 1 avenue de la terrasse, 91190, Gif-sur-Yvette, France.

[3] Institut Pasteur, Université Paris Cité, CNRS UMR 3528, Unité de Biochimie des Interactions macromoléculaires, Département de Biologie Structurale et Chimie, 25-28 rue du Docteur Roux, 75724 Paris cedex 15, France

[#] present address: Fundamental Microbiology and Pathogenicity Laboratory, UMR 5234 CNRS-University of Bordeaux, SFR TransBioMed, 33076 Bordeaux, France

[*]Correspondence: louis.renault@i2bc.paris-saclay.fr

Total characters, including spaces (from title to figure legends, including references): 79,628

Abstract

ExoY virulence factors are members of a family of bacterial nucleotidyl cyclases (NCs) that are activated by specific eukaryotic cofactors and overproduce cyclic purine and pyrimidine nucleotides in host cells. ExoY NC activity is triggered by actin. Here, we investigate ExoY-like toxins that selectively interact with monomeric actin (G-actin) using the ExoY effector domain (Vn-ExoY) of the *Vibrio nigripulchritudo* Multifunctional-Autoprocessing Repeats-in-ToXin (MARTX) toxin. Our biochemical and structural analyses suggest that Vn-ExoY uses profilin-bound G-actin as a physiological cofactor, sequestering this complex and preventing its elongation factor-mediated assembly at the barbed ends of actin filaments. A series of high-resolution crystal structures of nucleotide-free, 3'-deoxy-ATP- or 3'-deoxy-CTP-bound Vn-ExoY, activated by free or profilin-bound G-actin-ATP/-ADP, provide unprecedented mechanistic details of how the active site of NC toxins is sequentially remodelled by cofactor and substrate binding and can accommodate nucleotides of different sizes as substrates. Our structural characterisations, together with mutational data on Vn-ExoY and its distantly-related ExoY from *Pseudomonas*

aeruginosa, **elucidate important features of their catalytic reaction that will help identify how to**

inhibit NC toxins.

INTRODUCTION

ExoY-like nucleotidyl cyclase (NC) toxins are virulence factors produced by several Gram-

negative β- and γ-Proteobacteria (Teixeira-Nunes *et al*, 2022). They belong to a family of bacterial NCs

that share a structurally-related catalytic NC domain that is inactive in bacteria. When injected into

eukaryotic cells, they bind to specific eukaryotic proteins that stimulate their enzymatic activity by

several orders of magnitude (Tang & Guo, 2009; Belyy *et al*, 2016). They then become highly active

cyclases with broad substrate specificity (Beste *et al*, 2013; Raoux-Barbot *et al*, 2018) and increase the

levels of cyclic purine (cAMP, cGMP) and pyrimidine (cCMP, cUMP) nucleotides, thereby interfering

with host cell signalling (reviewed in (Ahuja *et al*, 2004; Tang & Guo, 2009; Morrow *et al*, 2017;

Seifert, 2017; Teixeira-Nunes *et al.*, 2022)). While the *Bacillus anthracis* edema factor (EF) and

Bordetella pertussis CyaA toxins are both activated by calmodulin (CaM), the ExoY-like NC toxins use

actin (Belyy *et al.*, 2016). CaM-activated toxins have been extensively studied over the last decades. In

contrast, the functional characteristics and virulence mechanisms of the ExoY-like NC toxins still need

to be clarified.

Exoenzyme Y (Pa-ExoY) is the second most prevalent type III secretion system (T3SS) exotoxin

encoded in the genomes of clinical or environmental isolates of *P. aeruginosa* (Feltman *et al*, 2001).

Pa-ExoY NC activity in host cells alters the host immune response (He *et al*, 2017; Jeon *et al*, 2017)

and leads to the release of cytotoxic amyloids from the pulmonary endothelium, impairing cellular repair

after infection (Morrow *et al.*, 2017; Renema *et al*, 2020; Voth *et al*, 2020). In *Vibrio vulnificus* biotype

3, a pathogen that causes severe foodborne and wound infections in humans, the ExoY module of its

MARTX toxin (Vv-ExoY) is essential for virulence (Ziolo *et al*, 2014). ExoY NC homologues from

different pathogens can potentially exert different cytotoxic effects in infected cells due to differences

in their protein sequence, the form of actin they require to become active, and their substrate specificities

(Raoux-Barbot *et al.*, 2018). The MARTX Vn-ExoY module of *Vibrio nigripulchritudo*, an emerging

pathogen in marine shrimp farming, is representative of the ExoY modules found in *Vibrio* strains. It

shares only 38 % sequence similarity with Pa-ExoY. Whereas Pa-ExoY requires actin filaments (F-

actin) for maximal activation (Belyy *et al.*, 2016), Vn-ExoY is selectively activated by G-actin *in vitro*.

Vn-ExoY, like EF and CyaA (Beckert *et al*, 2014), strongly prefers ATP and can use CTP as substrate,

but much less efficiently (Raoux-Barbot *et al.*, 2018). Pa-ExoY, on the other hand, synthesises a wide

range of cyclic nucleotide monophosphates (cNMPs) (Beckert *et al.*, 2014) with the following *in vitro*

substrate preference: GTP>ATP≥UTP>CTP (Raoux-Barbot *et al.*, 2018). Recently, cUMP and cCMP have emerged as novel second messengers that activate antiviral immunity in bacteria (Tal *et al*, 2021). On the other hand, very little is known about their role in eukaryotic cells (Seifert & Schirmer, 2022).

Bacterial NC toxins are defined as class II adenylate cyclases (ACs). Their catalytic domain is structurally unrelated to the widely distributed class III adenylyl or guanylyl cyclases (GCs) (Hurley, 1999; Tesmer *et al*, 1999; Steegborn, 2014; Childers & Garcin, 2018; Khannpnavar *et al*, 2020) and may represent a potential drug target against bacterial infections. In their cofactor-bound state, they are several orders of magnitude more active than activated mammalian class III ACs (Tang & Guo, 2009), which use a dimeric catalytic architecture. The molecular basis for the turnover differences between these two AC classes remains to be elucidated. The catalytic domain of NC toxins can be structurally divided into subdomains C_A and C_B, with the catalytic site located in a groove at their interface (Drum *et al*, 2002). EF and CyaA recognise CaM via two flexible regions of C_A that are called switch A (residues 502-551 in EF, 199-274 in CyaA) and switch C (630-659 in EF, 348-364 in CyaA) (Drum *et al.*, 2002; Guo *et al*, 2005; Shen *et al*, 2005). In CaM-bound EF, switch A and C stabilise another key region of C_A, called switch B (578-591 in EF, 299-311 in CyaA). This region contains residues that either directly bind ATP or stabilise catalytic residues in the groove between C_A and C_B (Drum *et al.*, 2002). In the inactive state of EF, switch B is disordered because switch A and C are distant from the catalytic site. Activation of EF or CyaA by CaM does not appear to require a large subdomain movement between C_A and C_B, but rather a CaM-induced movement of switch A and C towards the catalytic site (Tang & Guo, 2009). This in turn stabilises switch B and ATP binding to favour a catalytically competent form of the enzyme. This allosteric activation-by-stabilisation mechanism was recently endorsed for the actin-activated NC toxins. In the first structure of Pa-ExoY crystallised alone after limited in-situ proteolysis, the three switches were not visible in the electron density due to partial proteolysis and/or flexibility (Khanppnavar & Datta, 2018). Recent cryo-EM structures of actin-activated Pa-ExoY and Vv-ExoY revealed that Pa- and Vv-ExoY also use their central switch A and C-terminal switch C to recognise F- and G-actin subunits, respectively (Belyy *et al*, 2021). Molecular dynamics simulations suggested that binding of Pa-ExoY to F-actin stabilises its otherwise flexible switches A and C, thereby stabilising the entire toxin and nucleotide-binding pocket (NBP). Despite these advances, the number of ions required for maximum efficiency, the nature of the catalytic base, or the coordination and role of the residues involved in the various reaction steps are still a matter of debate (Drum *et al.*, 2002; Shen *et al*, 2004; Guo *et al.*, 2005; Shen *et al.*, 2005; Gupta *et al*, 2006; Martinez *et al*, 2009; Mones *et al*, 2013; Jara & Martinez, 2016; Belyy *et al.*, 2021; Teixeira-Nunes *et al.*, 2022). The molecular basis of the NC toxin's pyrimidinyl cyclase activity is also unknown.

Here, using *in vitro* functional studies in actin self-assembly and structural studies, we show that

Vn-ExoY targets and is potently activated by profilin-bound G-actin. This interaction also prevents the

actin-profilin complex from supporting the assembly of F-actin barbed ends. We present a series of

high-resolution structures of Vn-ExoY activated by free or profilin-bound actin-ATP/ADP, in the

absence or presence of non-cyclisable ATP or CTP analogues bound to Mg^{2+} or Mn^{2+} ions. The data

conclusively demonstrate that NC toxins employ a two-metal-ion mechanism to catalyse the cyclisation

reaction of large purine or small pyrimidine nucleotides (with double or single carbon-nitrogen ring

bases, respectively). Residues that play an important role in the purinyl and pyrimidinyl cyclase activity

of NC toxins are validated in Vn-ExoY and its distantly-related Pa-ExoY.

Results

Vn-ExoY binds to actin monomers with only modest affinity and inhibits actin self-assembly

As a model for ExoY toxins that are selectively activated by G-actin, we used a functional Vn-

ExoY$^{3412-3873}$ effector domain/module corresponding to residues 3412 to 3873 of *V. n.* MARTX (Raoux-

Barbot *et al.*, 2018). We first examined the effect of Vn-ExoY on actin-ATP/-ADP self-assembly. To

avoid the indirect effects due to ATP depletion by Vn-ExoY AC activity, we used an inactive double

mutant K3528M/K3535I, hereafter referred to as Vn-ExoY^{DM} (corresponding to Pa-ExoY$^{K81M/K88I}$)

(Raoux-Barbot *et al.*, 2018). Actin polymerisation kinetics were monitored using pyrene-labelled actin,

which shows an increase in fluorescence intensity when incorporated into filaments. While inactive Pa-

Exo^{K81M} accelerated the rate of actin polymerisation (Fig. 1A) (Belyy *et al.*, 2021), Vn-ExoY^{DM}

inhibited the polymerisation rate of both G-actin-ATP-Mg and G-actin-ADP-Mg (Fig. 1B-C). Thus,

Vn-ExoY interacts with G-actin-ATP or -ADP in a very different way than Pa-ExoY. To understand

the inhibitory effect of Vn-ExoY, we examined the association of free or Vn-ExoY^{DM}-bound G-actin

with the barbed (+) or pointed (-) ends of filaments. Spectrin-actin or gelsolin-actin seeds were used to

induce elongation at the barbed- and pointed-ends, respectively. Interaction of Vn-ExoY^{DM} with G-

actin-ATP-Mg inhibited actin assembly at both ends (Fig. 1D). The dose-dependent effects of Vn-

ExoY^{DM} on barbed- and pointed-end elongation rates were consistent with the formation of a 1:1 molar

complex between G-actin and Vn-ExoY^{DM} sequestering G-actin with an equilibrium dissociation

constant (Kd) of 21 μM (Fig. 1D). To confirm the binding affinity by direct measurement, we performed

fluorescence-detected sedimentation velocity experiments using analytical ultracentrifugation (AUC-

FDS) with G-actin-ATP labelled with the fluorescent dye Alexa-488 and unlabelled Vn-ExoY fused to

Thioredoxin (Trx-Vn-ExoY). This protein fusion, which is more soluble than Vn-ExoY, was preferred

for dose-dependent analyses at high concentrations (Fig. 1E). With free labelled G-actin (0.06 μM)

bound to Latrunculin-A, an actin-polymerisation inhibitor, the actin sedimentation coefficient

distribution gave a peak centred at an $s_{20,w}$ value of 3.3 S (Fig. 1E, left panel) consistent with a globular

actin monomer of 42 kDa. Upon incremental addition of Trx-Vn-ExoY, the sedimentation coefficient

distribution of labelled actin shifted to higher s-values (from 3.2 up to 4.7 S), demonstrating the

interaction of Vn-ExoY with actin. Peak position and shape were concentration-dependent, and the

relative proportion of the larger species in the peaks increased with increasing concentrations of Trx-

Vn-ExoY. The titration was consistent with the formation of a 1:1 Vn-ExoY:G-actin complex with a

Kd of 11.7 ± 0.9 µM (Fig. 1E, right panel). Thus, Vn-ExoY captures G-actin with only modest affinity

to form a complex that sequesters G-actin.

Vn-ExoY binds and is activated by G-actin bound to profilin

We next investigated whether Vn-ExoY could compete with G-actin-binding proteins (G-ABPs) in

host cells despite its moderate affinity for G-actin (dos Remedios *et al*, 2003). We examined the *in vitro*

AC activity of Vn-ExoY stimulated by G-actin in the presence of three regulatory G-ABPs that display

different binding interfaces on G-actin: (i) a chimeric β-thymosin domain of Thymosin-β4 and Ciboulot

proteins, which inhibits actin self-assembly by sequestering G-actin (Didry *et al*, 2012), (ii) profilin and

(iii) a profilin-like WH2 domain of the Cordon-Bleu protein (Husson *et al*, 2011). The latter two control

the unidirectional assembly of actin monomers at the barbed-ends of filaments. At saturating

concentrations of these three G-ABPs, only profilin did not interfere with G-actin-induced activation of

Vn-ExoY AC activity (Fig. 2A). We then performed AUC-FDS experiments, monitoring the

sedimentation behaviour of Alexa-488-labelled-G-actin in the presence of Vn-ExoY, profilin or both

proteins. The formation of complexes of increasing size was observed by the shifts from an $s_{20,w}$ value

of 3.5 S obtained for the peak of free Alexa488-labelled actin to 4.2 S with profilin (15 kDa), 5.0 S with

the larger Trx-Vn-ExoY protein (69.1 kDa), and to a higher value of 5.5 S with both proteins (Fig. 2B),

validating the formation of a ternary Vn-ExoY:actin:profilin complex in solution. We next investigated

whether the interaction of Vn-ExoY with G-actin was modulated by the simultaneous binding of

profilin. We measured the binding strength of Trx-Vn-ExoY[DM] to fluorescently labelled G-actin-ATP

alone or bound to profilin using microscale thermophoresis (MST). Titrations of the interaction by

measuring changes in fluorescence (Fig. 2C) or MST signal (Fig. S1) show that Vn-ExoY binds with

similar affinity to free or profilin-bound G-actin.

Assembly of actin:profilin at F-actin barbed-ends is inhibited by Vn-ExoY

To understand the effect of Vn-ExoY interaction with actin:profilin on actin self-assembly, we first

examined its effect on the spontaneous assembly of actin:profilin at F-actin barbed-ends. We measured

the normalised initial rate of barbed-end filament growth in the presence of increasing concentrations

of profilin with or without fixed concentrations of Vn-ExoY. Without Vn-ExoY, barbed-end growth

was only modestly inhibited by saturating concentrations of profilin as actin:profilin can associate with

and elongate F-actin barbed-ends (Fig. 2D). With Vn-ExoY, the dependence of barbed-end growth on

profilin concentration shifted to lower values in a Vn-ExoY dose-dependent manner. This shift was

consistent with a decrease in the G-actin pool available for barbed-end elongation due to its

sequestration by Vn-ExoY. We next examined the effect of Vn-ExoY on actin:profilin assembly at F-

actin barbed-ends regulated by VASP or formin processive barbed-end elongation factors. The latter

bind to barbed-ends via an F-actin binding domain (called Formin-Homology domain 2 (FH2) in

formins) and favour actin:profilin recruitment to barbed-ends through their adjacent profilin-binding

proline-rich motifs (PRMs, which are repeated in formin FH1 domain). Formins or VASP also

counteract the barbed-end capping and growth inhibition by capping proteins (CP). The barbed-end

growth of F-actin seeds was therefore monitored in the presence of both G-actin:profilin, CP and either

the C-terminal FH1-FH2 domains of mouse Dia1 formin (F or mDia1-FH1-FH2, Figure 2E) or full-

length human VASP (V, Figure 2F) to approach the physiological regulation of F-actin barbed-ends.

When actin (1µM) is saturated with profilin (8 µM), actin nucleation is prevented, and polymerisation

does not occur (horizontal dashed grey lines). Under these conditions, the addition of 0.35 nM of

spectrin-actin seeds induced rapid elongation of the barbed ends of actin filaments (ascending dotted

grey curves). In the presence of 2.1 nM CP, this barbed-end elongation was partially inhibited (black

dashed curves), but restored by the addition of 3.2 nM mDia1-FH1-FH2 (Fig. 2E, black curves) or 0.22

µM VASP (Fig. 2F, black curves), as both elongation factors compete with CP at the barbed ends.

Further addition of Vn-ExoY at 0.8 or 1.4 µM to mDia1-FH1-FH2 or VASP in the presence of CP

inhibited the barbed-end growth from actin:profilin induced by the two elongation factors (blue curves).

These inhibitory effects were independent of Vn-ExoY AC activity for the duration of the kinetic

experiments (Fig. S2). Thus, the interaction of Vn-ExoY with G-actin:profilin inhibits the spontaneous

or VASP-/formin-mediated assembly of this complex at the most dynamic ends of F-actin.

Crystal structures of nucleotide-free and 3'-deoxy-ATP-bound Vn-ExoY in complex with actin-

ATP/ADP:profilin or actin-ATP

Next, we investigated the structural mechanism of Vn-ExoY activation by free or profilin-bound G-

actin. To improve the crystallisation strategy (see Supplementary Information), we designed a chimeric

protein containing three fused functional domains: Vn-ExoY extended at the C-terminus (Vn-ExoY^{3455-}

^{3896}, either wild-type or with K3528M/K3535I), a short profilin-binding proline-rich motif (PRM) and

profilin (Fig. 3A, VnE/VnE^{DM}-PRM-Prof). The chimera sequestered G-actin with approximately 100-

fold higher affinity than the Vn-ExoY$^{3412-3873}$ module and exhibited strong actin-dependent AC activity

at low G-actin concentrations (Fig. 3B-C). Thus, the actin-binding regions in Vn-ExoY and profilin
appear to be fully functional in this chimera. Using this chimera (wild-type or double-mutant), we solved
two crystal structures at 1.7 Å resolution: (i) VnE^{DM}-PRM-Prof bound to actin-ADP with latrunculin B
(LatB), and (ii) VnE-PRM-Prof bound to actin-ADP without LatB (Fig. 3D, Table S1). In these
structures, profilin is bound to actin-ADP, the PRM to profilin, Vn-ExoY to actin, while residues 3863-
3895 connecting the C-terminus of Vn-ExoY to the PRM are disordered (Fig. 3D, Table S1). Vn-ExoY
contacts actin-ADP via its switch A and C, but only switch C is fully ordered. The Vn-ExoY switch A
disorder (residues 3682-3727) is close to the LatB-binding site in actin. However, solving the structure
without LatB only slightly reduces the switch A disorder. In the latter structure, the NBP of Vn-ExoY
is further stabilised by a sulphate ion (Fig. 3D, VnE-PRM-Prof-SO$_4^{2-}$:actin-ADP). On the basis of these
structural data, we designed a minimal functional Vn-ExoY module (residues 3455-3863) by removing
the C-terminal disordered residues 3864-3895 (Fig. 3A). This Vn-ExoY module (shortened by 32 C-
terminal residues) shows similar catalytic activity to our original Vn-ExoY$^{3412-3873}$ construct and
allowed the determination of two additional crystal structures: (i) nucleotide-free Vn-ExoY bound to
actin-ATP-LatB and profilin, with a sulphate ion in Vn-ExoY NBP (Fig. 3E, Vn-ExoY-SO$_4^{2-}$:actin-
ATP-LatB:profilin), and (ii) Vn-ExoY bound to actin-ATP-LatB with its NBP filled by a non-cyclisable
ATP analogue 3'-deoxy-ATP (3'dATP, Figure 3F), at 1.7 and 2.1 Å resolution, respectively. The Vn-
ExoY-SO$_4^{2-}$:actin-ATP-LatB:profilin structure is very similar to the structures of the VnE/VnE^{DM}-
PRM-Prof chimera bound to actin-ADP, with the Vn-ExoY switch A similarly disordered. In contrast,
the Vn-ExoY-3'dATP-2Mg^{2+}:actin-ATP-LatB structure co-crystallised with 3'dATP reveals a more
ordered Vn-ExoY:actin interface and a distinct conformational state of Vn-ExoY NBP that is more
competent for catalysis, as described below.

In the G-/F-actin-bound state, the switch-C of ExoYs ensures stable interactions while other regions remain dynamic

Vn-ExoY undergoes significant structural rearrangements at both the C$_A$ and C$_B$ subdomains between
its nucleotide-free conformation bound to actin-ATP/ADP and profilin, and its 3'dATP-bound
conformation in complex with actin-ATP (Fig. 4A). At the Vn-ExoY:actin interface, none of the buried
contacts of the Vn-ExoY switch C move, except for L3827, which is relocated by the remodelling of
switch A (Fig. 4A-B). Thus, switch C appears to be essential for initiating the interaction and stabilising
Vn-ExoY on actin. In the absence of 3'dATP, most of switch A is too flexible to be modelled, and the
switch A portions that are ordered in the absence of ligand undergo profound rearrangements upon
3'dATP-binding (up to 10 Å on D3682 and K3683 Cα atoms, Figure 4A). Subsequently, the letters after
the residue numbers are used to distinguish residues of different proteins or functional regions (e.g. $^{VnE-}$

^{C_A} refers to residues belonging to the Vn-ExoY subdomain C_A). The different conformations of Vn-ExoY bound to actin/actin:profilin with or without a nucleotide substrate analogue allow us to define the mobile "switch" regions as follows: Vn-ExoY switch A as residues Y3675-V3727$^{VnE-Switch-A}$ (I222-N264$^{PaE-Switch-A}$, UniProt AC Q9I1S4_PSEAE), switch B as H3753-A3770$^{VnE-Switch-B}$ (H291-A308$^{PaE-Switch-B}$), switch C as I3819-L3863$^{VnE-Switch-C}$ (K347-V378$^{PaE-Switch-C}$). Other important conformational changes near the interface occur in the parts of switch C in contact with switch A and B, and in the parts of switch B in contact with switch A and C (Fig. 4A-B). These regions show high B-factors in all the complexes with nucleotide-free Vn-ExoY (Fig. 4D-1). Near the interface, the C_B subdomain of Vn-ExoY undergoes the most drastic global rearrangement with a large 25° rotation as a rigid body around the hinge regions 3532-3533^{VnE} and 3662-3663^{VnE}. It thus acts as a lid (hereafter referred to as C_B/LID, residues K3533-M3662$^{VnE-LID}$, corresponding to K86-M209$^{PaE-LID}$ in Pa-ExoY) that closes on C_A to stabilise 3'dATP binding (Fig. 4A-B, movie S1A-B). The conformational changes induced by 3'dATP binding bring several regions of the Vn-ExoY C_A subdomain closer to actin, increasing the buried solvent-accessible area (BSA) of switch A from 338 to 707 Å^2, reaching the BSA value of switch C (810 and 766 Å^2, respectively). Vn-ExoY binds to actin over a large surface area, with switch A binding primarily to actin subdomains 2 and 4 and switch C binding to actin subdomains 1 and 3 (Fig. 4C).

To validate the importance of switch C for proper positioning of ExoY on actin, we mutated conserved buried residues of switch C that remain well anchored and stable in the complexes with or without 3'dATP. Actin D25 (D25^{Actin}) forms a buried salt bridge at the interface core with the R3824$^{VnE-switch-C}$ side chain (distances of 2.7 and 3.2 Å between N-O atom pairs, Figure 3D-F, 4B-C) and was identified as a critical residue for Pa- and Vn-ExoY activation by F- and G-actin, respectively (Belyy *et al.*, 2016). The R3824A$^{VnE-switch-C}$ mutation alone results in a 130-fold reduction in Vn-ExoY G-actin stimulated AC activity (Fig. 5A). The extreme C-terminus of Pa-ExoY is also critical for its binding to F-actin (Belyy *et al*, 2018). The corresponding region in Vn-ExoY is located at the edge of the Vn-ExoY:actin interface, where the extreme C-terminal amphipathic helix of Vn-ExoY (residues 3847-3863$^{VnE-switch-C}$) binds into the hydrophobic cleft between actin subdomains 1 and 3. Mutation of the most buried hydrophobic residues of this amphipathic helix to alanine results in a 240-fold reduction in its AC activity (Fig. 5A). Similar interactions occur in the switch C of F-actin-bound Pa-ExoY (Belyy *et al.*, 2021). Three basic amino-acids of Pa-ExoY near D25^{Actin} may play the same role as R3824$^{VnE-switch-C}$: K347$^{PaE-switch-C}$, H352$^{PaE-switch-C}$ and R364$^{PaE-switch-C}$. While the R364A mutation in Pa-ExoY switch C had almost no effect on its F-actin-stimulated GC activity, the K347A and H352A mutations caused a ~15- and ~8-fold decrease, respectively (Fig. 5C), indicating that these two amino-acids together function as R3824$^{VnE-switch-C}$. On the other hand, changing the three bulky hydrophobic side chains of the C-terminal amphipathic helix of Pa-ExoY (i.e. F367-L371-F374, Figure 5B) into alanine

caused a ~12,000-fold reduction in F-actin stimulated GC activity (Fig. 5C). These residues in the switch

C of Vn-ExoY or Pa-ExoY are conserved in most of their counterparts in other β- and γ-Proteobacteria

(Fig. 5D). Thus, during the transition of these enzymes to their catalytic conformation, their complex

with G-/F-actin is first stably docked to their cofactor at their switch-C, while other regions remain

dynamic to regulate additional functional rearrangements.

Significance of switch A and C_B/LID closure in stabilising nucleotide and Mg^{2+}

Except for a difference in the structural topology of the proteins (Fig. S3), the overall structure of

Vn-ExoY bound to actin and 3'dATP resembles *V. vulnificus* Vv-ExoY bound to actin:profilin (Belyy

et al., 2021) (Fig. S4A). In the 3.9-Å resolution cryo-EM structure, the 3'dATP ligand used for the

preparation of the complex could not be modelled in Vv-ExoY. In both structures, switch A adopts a

similar conformation between actin subdomains 2 and 4. This interaction primarily prevents Vn- and

Vv-ExoY from binding along the side of F-actin. This explains their G-actin specificity, whereas the

orientation of Pa-ExoY switch A bound to F-actin differs (Belyy *et al.*, 2021) (Fig. S4B). Furthermore,

the positioning of the Vn-ExoY switch A on G-actin prevents the association of Vn-ExoY:G-actin or

Vn-ExoY:G-actin:profilin complexes with F-actin barbed-ends (Fig. S5A) and is responsible for the

inhibition seen in Figure 1D, 2D-F. Switch A is also the region with the most divergent folding at the

toxin:cofactor interface between G/F-actin-bound ExoYs and CaM-bound EF or CyaA (Figs. S5B-C).

Thus, this region appears to be evolutionarily essential for the cofactor specificity of NC toxins.

Our different structures reveal two major consecutive conformational changes occurring at the

catalytic site of NC toxins. In the free/inactive Pa-ExoY structure (Khanppnavar & Datta, 2018), the A,

B, and C switches were not visible as partially degraded and/or flexible. In our nucleotide-free structures

of Vn-ExoY bound to G-actin-ATP/-ADP and profilin (Fig. 3D-E), the Vn-ExoY NBP is still only

partially stabilised by cofactor binding (Fig. 4D-1), showing that cofactor binding is not sufficient to

fully stabilise the NBP and catalytic residues of NC toxins, as previously suggested (Drum *et al.*, 2002;

Tang & Guo, 2009; Belyy *et al.*, 2021). 42 to 46 of the 53 residues from switch-A are still disordered.

A SO_4^{2-} ion is bound to the Vn-ExoY NBP via residues (K3528[VnE-CA], K3535[VnE-LID], S3536[VnE-LID],

K3554[VnE-LID]) that coordinate the γ- and β-phosphates of 3'dATP when this ligand is bound (Fig. 4D-2).

Binding of the ATP substrate and Mg^{2+} ion in the Vn-ExoY NBP requires rearrangement of switch B

and the supporting part of switch C (Fig. 4D-3). To reach its most favourable binding position, the

nucleotide must move ~4 Å (distance between the SO_4^{2-} ion and 3'dATP γ-phosphate) through the LID

closure (Fig. 4D-3). Structural rearrangements in the region of switch C, which supports switch B, are

accompanied by stabilisation of switch A and the substrate analogue (Fig. 4D-4,4D-5). In the final stage,

switch B and C are well-ordered, while the entire switch A is stabilised on G-actin (Fig. 4D-5). In the

Vn-ExoY-3'dATP-2Mg^{2+}:actin-ATP-LatB structure, switch A ultimately contributes to the stabilisation

of the nucleotide base through interactions with M3726$^{VnE-switch-A}$.

Catalytic mechanism of ExoY NC toxins activated by G- or F-actin with purine nucleotides

In the known cofactor:toxin structures, the nucleotide ligands adopt different positions and

conformations in the toxin catalytic pocket, variably coordinated by one or two metal ions (Fig. S6-7).

The Pa-ExoY-bound 3'dGTP is coordinated by a single metal ion in the F-actin-activated Pa-ExoY

structure (Belyy *et al.*, 2021). In the structure of Vn-ExoY bound to actin and 3'dATP, the electronic

density and coordination geometry suggest that 3'dATP is coordinated by two Mg^{2+} ions. To

unambiguously distinguish metal ions from water, we replaced Mg^{2+} with Mn^{2+} ions and collected an

anomalous data set at the manganese absorption edge (Supplementary Materials). The anomalous

difference Fourier map confirms that two Mn^{2+} ions are bound to 3'dATP at the positions occupied by

the Mg^{2+} ions, with no significant changes elsewhere (Fig. 6B). Mg$^{2+}_A$ is pentacoordinated in a square

pyramidal geometry by two aspartates D3665^{VnE-CA} and D3667^{VnE-CA}, and H3753$^{VnE-switch-B}$, which are

invariant in all NC toxins (Fig. 6A), an α-phosphate oxygen of 3'dATP, and a water molecule. Mg$^{2+}_B$ is

octahedrally hexacoordinated by the same two aspartates (D3665/D3667^{VnE-CA}), one oxygen from each

of the three phosphates of 3'dATP and a water molecule.

Figure 6A summarises the main interactions observed with 3'dATP in the Vn-ExoY-3'dATP-

2*Mg^{2+}:actin-ATP-LatB structure, extrapolated to the natural substrate ATP. The residues of Vn-ExoY

that coordinate the nucleotide and metal-ions are indicated in coloured boxes in the first inner shell. The

corresponding residues in the Pa-ExoY, EF, and CyaA sequences are in the next outer shells. 9 out of

the 14 substrate and Mg^{2+} ion binding residues belong to regions of the enzyme that are flexible and

significantly rearranged and stabilised upon binding of the cofactor, nucleotide substrate and metals.

This explains the low basal activity of the free enzyme (Raoux-Barbot *et al.*, 2018). All side-chain

contacts are provided by residues that are invariant or similar in Pa-ExoY, EF, and CyaA, and most

interactions are conserved in CaM-activated EF bound to 3'dATP with a Yb^{3+} ion and a closed LID

(PDB 1k90) (Drum *et al.*, 2002) (Fig. S7D).

To assess the relevance of the observed interactions with the ATP analogue, some of the

conserved residues involved were mutated in Vn- and Pa-ExoY and the effect on their AC and GC

activity was investigated. H351^{EF-LID} in EF has been proposed to play a role in the 3'OH deprotonation,

either by accepting the 3'OH proton (general base), or by facilitating the presence of OH$^-$ ions near the

ATP 3'OH group (Drum *et al.*, 2002; Guo *et al*, 2004; Shen *et al.*, 2005; Gupta *et al.*, 2006; Mones *et*

al., 2013; Jara & Martinez, 2016). H351^{EF-LID} is conserved in CyaA, but in ExoYs it corresponds to a

lysine: K3533$^{VnE-LID}$ or K86$^{PaE-LID}$ (Fig. 6A). In the active site of Vn-ExoY bound to actin, 3'dATP and

2 Mg^{2+} or 2 Mn^{2+}, the K3533$^{VnE-LID}$ side chain is too distant to potentially coordinate the ATP 3'OH

moiety if the latter were present. K3533Q$^{VnE-LID}$ or K86Q$^{PaE-LID}$ mutations only decreased G/F-actin-

activated AC and GC activity by 14- and 12-fold, respectively (Fig. 6C-D), further suggesting that these

residues are unlikely to act as a general base. Next, we examined K3528^{VnE-CA} and K3535$^{VnE-LID}$, which

are close to the bridging O3α and non-bridging O2α, O1β and O2γ oxygens of 3'dATP (Fig. 6A). Both

side chains are ~3.4 Å from O3α. K3528^{VnE-CA} is closer to O2α and O2γ (2.8-2.9 Å) than K3535$^{VnE-LID}$,

and K3535$^{VnE-LID}$ is closer to O1β (3.3 Å). The catalytic activity of Vn- and Pa-ExoY was essentially

abolished (at least ~23,000-fold reduction) by the K3528Q^{VnE-CA} and the corresponding K81Q^{PaE-CA}

mutations in Pa-ExoY. The K3535Q$^{VnE-LID}$ or the equivalent K88Q$^{PaE-LID}$ mutations resulted in a 250-

and 26-fold decrease in G-/F-actin-induced Vn-ExoY AC or Pa-ExoY GC activity, respectively (Fig.

6C-D). The invariant H3753$^{VnE-switch-B}$, which coordinates Mg^{2+}_A and stabilises the purine base via a π-

π stacking with its imidazole ring, is specific to NC toxins. Mg^{2+}_A may be particularly important in the

NBP of activated mammalian ACs to coordinate and polarise the 3'OH of ATP, facilitating its

deprotonation as a first step in the cyclisation reaction (Tesmer *et al.*, 1999; Steegborn, 2014; Hahn *et*

al, 2015). The H3753Q$^{VnE-switch-B}$ or equivalent H291Q$^{PaE-switch-B}$ mutations greatly reduced the catalytic

activity of the toxins (~13,000-fold reduction), suggesting a critical role for the switch-B invariant

histidine and Mg^{2+}_A in the catalysis of NC toxins.

Structural basis for the pyrimidinyl cyclase activity of ExoYs

The activity of NC toxins as pyrimidine nucleotide cyclases was identified some time ago, both

in vitro (Gottle *et al*, 2010; Raoux-Barbot *et al.*, 2018) and in eukaryotic cells (Beste *et al.*, 2013;

Beckert *et al.*, 2014), but remains poorly characterised. Among the pyrimidine nucleotides, Vn-ExoY,

EF or CyaA all primarily use CTP as a substrate, whereas Pa-ExoY prefers UTP but can also use CTP.

To determine how activated NC toxins can also use small pyrimidine nucleotides as substrates, we

solved the structure of Vn-ExoY bound to 3'dCTP in complex with actin-ADP at 2.2 Å resolution (Vn-

ExoY-3'dCTP-2*Mn^{2+}:actin-ADP-LatB). This is the first structure of an activated NC enzyme to be

determined in association with a pyrimidine nucleotide substrate analogue. The overall conformation of

Vn-ExoY in this structure is identical to those found in the complexes with actin-ATP, 3'dATP and two

Mg^{2+} or Mn^{2+}. As with 3'dATP, the LID of Vn-ExoY is closed on 3'dCTP and two metal ions coordinate

the phosphate moiety of 3'dCTP. In the NBP of Vn-ExoY, 3'dCTP is stabilised by the same residues

and types of interactions as 3'dATP (Fig. 7A). The main differences are in the interactions with the

nucleotide ribose and base moieties, which are stabilised by fewer and weaker hydrogen bonds and

framed by fewer hydrophobic interactions with 3'dCTP than with 3'dATP. With 3'dATP, the ribose is

stabilised by L3530^{VnE-CA} hydrophobic interactions and hydrogen-bonded by K3533$^{VnE-LID}$ and

N3759[VnE-switch-B] side-chains (Fig. 6A). In the activated structures of Vn-ExoY and tmAC (Tesmer *et*

al., 1999) with an NBP closed on an ATP analogue, N3759[VnE-switch-B] and N1025[tmAC-IIC2] similarly

hydrogen-bond the O4' oxygen of the nucleotide ribose ring (Fig. 6A, 6F). Just as mutations at

N1025[tmAC-IIC2] reduce tmAC kcat by 30-100 (Yan *et al*, 1997), mutations at N583[EF-switch-B] in EF

(equivalent to N3759[VnE-switch-B]) reduce EF catalytic activity by at least two orders of magnitude (Drum

et al., 2002). These data highlight the importance of hydrogen bonding with the nucleotide ribose for

correct positioning during class II and III AC catalysis. With 3'dCTP, electrostatic interactions with the

nucleotide ribose switch to a single, more labile hydrogen bond between the cytidine ribose ring O4'-

oxygen and a water molecule coordinated by D3756[VnE-switch-B] or N3759[VnE-switch-B]. The cytosine base is

hydrogen-bonded by M3726[VnE-switch-A] and A3755[VnE-switch-B] as for the adenine base. However, these

residues only coordinate the N4 atom of cytosine. For 3'dATP, they stabilise both the N1 and N6 atoms

of adenine through shorter hydrogen bonds. Similarly, in the Pa-ExoY:3'dGTP-1*Mg^{2+}:F-actin

structure, the N1, N2, and O6 guanine atoms are hydrogen-bonded by the Ser292[PaE-switch-B], H291[PaE-

switch-B], and Glu258[PaE-switch-A] side chains, respectively (Fig. S7C). The pyrimidine base is otherwise

surrounded by similar hydrophobic interactions to the purine base of 3'dATP. However, it lacks Pro3760

side-chain interaction with imidazole ring of 3'dATP purine base on solvent accessible side. On the

triphosphate moiety of 3'dCTP, the phosphates and two metal ions are tightly coordinated and less

solvent-accessible than the nucleoside moiety in Vn-ExoY NBP, which is very similar to 3'dATP.

The mutations analysed above for Vn-ExoY AC or Pa-ExoY GC activity were reconsidered and

their effect on their cytidylyl or uridylyl cyclase (CC/UC) activity was examined to determine whether

the conserved interactions with the nucleotide triphosphate moiety play a similar role in the purinyl and

pyrimidinyl cyclase activity of ExoYs. The mutations with the greatest effect on Vn-ExoY CC or Pa-

ExoY UC activity (150- to 480-fold reduction) are those that also most affect their AC or GC activity,

namely K3528Q[VnE-CA] or K81Q[PaE-CA] and H3753Q[VnE-switch-B] or H291Q[PaE-switch-B], respectively (Fig. 7B-

C). As with 3'dATP, the K3528[VnE-CA] side chain hydrogen-bonds the α, β, and γ phosphate oxygens of

3'dCTP, and the H3753[VnE-switch-B] side chain stabilises both Mg^{2+}_A via a hydrogen bond and the cytosine

base via a π-π stacking with its imidazole ring. Taken together, these results demonstrate that NC toxins

use a similar two-metal-ion catalytic mechanism with purine and pyrimidine nucleotide substrates,

relying on the close coordination of their triphosphate moiety for catalysis.

Discussion

Here, we have used the MARTX Vn-ExoY module to explore the functional specificities of

ExoY-like toxins that selectively interact with G-actin (Raoux-Barbot *et al.*, 2018). Our data show that

Vn-ExoY's modest affinity for free G-actin (Fig. 1D-E, Kd~12 μM compared to a Kd of ~0.2-20 nM

for CyaA or EF binding to CaM (Shen *et al*, 2002)) is compensated by its ability to interact with and be

efficiently activated by G-actin bound to profilin (Fig. 2A-C). Vn-ExoY is unable to bind to G-actin in

complex with other common G-actin binding proteins such as Thymosin-β4, WH2 domains (Fig. 2A)

or actin-depolymerising factor (ADF)/cofilin due to overlapping binding interfaces between actin

subdomains 1 and 3 (Fig. S9). The G-actin:profilin complex is abundant and important for supporting

filament assembly at barbed-ends in eukaryotic cells (dos Remedios *et al.*, 2003; Pimm *et al*, 2021). It

is therefore probably the major physiological form of G-actin that is recognised and used as a cofactor

by Vn-ExoY and closely-related ExoYs in eukaryotic cells. Our structural bioinformatics analysis

suggests that many ExoY-like homologues from different γ-proteobacteria are indeed structurally-

related to Vn-ExoY (Fig. 5D). Vn-ExoY interaction with G-actin:profilin inhibits both spontaneous and

formin- or VASP-mediated association of G-actin:profilin complexes with F-actin barbed-ends *in vitro*,

independent of the toxin's NC activity (Fig. 2D-F, S2). However, as shown by the multipartite

interactions of our actin-bound VnE-PRM-Prof chimera, the ternary complex does not prevent profilin

from interacting with PRM segments (Fig. 3D). Further studies are required to determine whether the

formation of such a Vn-ExoY:G-actin:profilin abortive ternary complex for barbed-end elongation may

also contribute to Vn-ExoY cytotoxicity in host cells independently of its NC activity. Several

observations suggest that Pa-ExoY may also act on eukaryotic target cells independently of its

enzymatic activity (Belyy *et al.*, 2016; Mancl *et al*, 2020; Silistre *et al*, 2021). The interaction of Vn-

ExoY with actin:profilin may be particularly important in determining subcellular localisation and

activation at sites of active actin cytoskeleton remodelling (Fig. 8). At these sites, G-actin:profilin and

elongation-promoting factors play a prominent role together in fine-tuning the coordination and

competition of F-actin networks (Suarez *et al*, 2015; Skruber *et al*, 2020). This fine-tuning of actin

dynamics is coordinated by a large number of multidomain ABPs and is tightly associated with cAMP

signalling in important cellular processes, including cell adhesion and T-cell migration (Howe, 2004;

Yamashita *et al*, 2011; Simao *et al*, 2021; Teixeira-Nunes *et al.*, 2022). VASP activity is regulated by

PKA and PKG (Aszodi *et al*, 1999) and is important for immune function (Estin *et al*, 2017).

Recruitment of the abortive ternary complex Vn-ExoY:actin:profilin by the PRMs of multidomain

ABPs may be of interest to pathogens to confine Vn-ExoY-like toxin activity to these discrete sites

where signalling and cytoskeletal remodelling are intense and highly intertwined. These elements

provide insight into why ExoY toxins target actin and not calmodulin as canonical NC toxins do.

Our structural data show that Vn-ExoY switch C interactions at the Vn-ExoY:actin interface

remain stable with different conformations of the LID/C_B subdomain and switch A (Fig. 3D-F, 4A-B),

thus firmly docking the toxin to actin. Our mutational analysis reveals two hotspots in the switch C of

Vn- and Pa-ExoY, whose mutation almost completely abolishes their G- or F-actin-dependent NC

activity (Fig. 5). These sites could be hits in the search for ways to inhibit ExoY toxins. In contrast, switch A is only barely stabilised by actin in the absence of substrate (Fig. 3D-E, 4D). EF and CyaA, on the other hand, have a switch A that is already fully folded on CaM in their nucleotide-free, cofactor-bound state (Drum *et al.*, 2002; Guo *et al.*, 2005). However, their C_B/LID subdomain can also adopt different positions in isolated CyaA (Malliavin, 2017) or between different crystal structures of CaM-bound EF (Fig. 6E, S8). Many crystal structures of CaM-activated EF or CyaA with nucleotide ligands in the active site were first crystallised without substrate analogues or reaction products and then incubated with these ligands. Crystal packing in these structures may have prevented complete rearrangement of the NBP and LID/C_B subdomain closure. As a result, and due to the intrinsic flexibility of the LID/C_B subdomain, both substrate analogues and metal ions occupy quite different positions in the NBP of CaM-bound EF or CyaA (Fig. S6-8) (Teixeira-Nunes *et al.*, 2022). These structures may correspond to intermediate stages of LID/C_B movement and substrate entry. Cofactor binding thus initiates allosteric stabilisation of switch B in NC toxins by repositioning the central switch A and the C-terminal switch C towards switch B, as previously reported (Drum *et al.*, 2002; Tang & Guo, 2009; Belyy *et al.*, 2021). In the ExoY NC toxins, however, complete stabilisation of switch A on actin also requires substrate binding and C_B/LID closure. Switch A is involved in the correct positioning of the substrate by directly coordinating the purine or pyrimidine base of the nucleotides in all cofactor-bound NC toxins (Fig. 6A, 7A, S7). In both the inactive conformation of free Pa-ExoY (Khanppnavar & Datta, 2018) and the nucleotide-free conformations of actin:profilin-bound Vn-ExoY (Fig. 3D-E), $H291^{PaE-switch-B}$ and $H3753^{VnE-switch-B}$, respectively, are stabilised away from the NBP by a salt bridge with $D3736^{VnE}$/$D274^{PaE}$, which are located near the switch A (Fig. 4D). After cofactor binding, substrate binding in ExoYs induces the flipping of $H3753^{VnE-switch-B}$/$H291^{PaE-switch-B}$ towards the NBP. This invariant histidine of NC toxin switch B assists in the correct positioning of the catalytic Mg^{2+}_A on the two invariant aspartates ($D3665D3667^{VnE-CA}$ or $D212D214^{PaE-CA}$) with purine or pyrimidine nucleotide substrate analogues (Fig. 6A-B, 7A). For all NC toxins, substrate binding therefore ultimately facilitates the C_B/LID closure.

Our data demonstrate unambiguously that Vn-ExoY uses a two-metal-ion-assisted catalytic mechanism with purine and pyrimidine nucleotides. Given the sequence similarity and structural conservation, this is most likely to be the case for all bacterial NC toxins after LID/C_B closure (Fig. 6A-B, 7A, S7). Interestingly, the coordination of metal ions and residues of the Vn-ExoY NBP with the ATP-analogue is also closer to that observed in the structures of activated class III ACs (Tesmer *et al.*, 1999) than to that previously observed in the NBPs of cofactor-activated NC toxins (Fig. 6A, 6F, S7). The active site of transmembrane or soluble class III ACs (tmAC/sAC) is located at the interface of their two homologous catalytic domains (termed C1 and C2 in tmAC, Figure 6F). It has also been

proposed to undergo an open-closed transition during catalysis. In the case of tmAC/sAC, however, the

exact role of the open-closed transition of their dimeric NBP and the identification of the dynamic active

site processes during substrate binding and turnover remain to be clarified (Steegborn, 2014). In the

closed NBP of all ACs, the correct positioning of the different ATP moieties is essential for catalysis.

This is shown by the strong effect of mutations targeting N1025$^{\text{tmAC-IIC2}}$ in tmAC (Yan *et al.*, 1997;

Tesmer *et al.*, 1999) or N583$^{\text{EF-switch-B}}$ in EF (Drum *et al.*, 2002), which corresponds to N3759$^{\text{VnE-switch-}}$

$^{\text{B}}$ in Vn-ExoY. These conserved asparagines similarly hydrogen-bond the ribose ring O4' oxygen of the

ATP analogue for catalysis (Fig. 6A, 6F). On the phosphate and 3'OH side, the most probable reaction

sequence as well as the key interactions during each step were investigated by computational studies

for tmAC with a closed ATP-binding NBP (Hahn *et al.*, 2015) and for EF with an open or nearly-closed

ATP-binding NBP (using PDB 1XFV/1SK6) (Mones *et al.*, 2013). The reaction model for tmAC better

fits the numerous and tight interactions around 3'dATP and metal ions observed in the closed NBP of

activated Vn-ExoY. The proposed mechanism for tmAC is a substrate-assisted general base catalysis,

where several residues of the enzyme appear to be particularly important with metal-ion cofactors for

lowering the two highest energy barriers of the reaction. First, K1065$^{\text{tmAC-IIC2}}$, Mg$^{2+}_{\text{A}}$, Mg$^{2+}_{\text{B}}$ and the

two conserved D396D440$^{\text{tmAC-VC1}}$ anchoring metal ions are critical for jointly facilitating the proton

transfer from the ribosyl 3′O to a γ-phosphate oxygen (Hahn *et al.*, 2015). R1029$^{\text{tmAC-IIC2}}$ is then critical

for the concerted phosphoryl transfer step in which K1065$^{\text{tmAC-IIC2}}$ is also involved. These interactions

belong to C1 and C2 structural elements that are distant in the open NBP of tmAC (Tesmer *et al.*, 1999).

Similarly, in NC toxins, the NBP closure brings together ATP/CTP- and metal-ion-coordinating

residues belonging to four distinct regions, which are distant from each other in the absence of cofactor

and substrate. Three of these ATP/CTP- and metal-ion coordination regions are intrinsically flexible.

Their positioning and stabilisation depend on both cofactor and substrate binding for switch A and B,

and substrate binding for C$_{\text{B}}$/LID (Fig. 6A, 7A). Substrate-induced C$_{\text{B}}$/LID closure after activation

ultimately reduces the degrees of freedom of the nucleotide substrate and metal ions in the NBP (Fig.

S7) to promote catalysis. Our mutational analysis in Vn-ExoY and the distantly-related Pa-ExoY

identifies the common elements of the NC toxin active sites that contribute to catalysing the limiting

steps in the cyclisation reaction with substrates of different sizes. The analysis pinpoints K3528$^{\text{VnE-}}$

$^{\text{CA}}$/K81$^{\text{PaE-CA}}$ and H3753$^{\text{VnE-switch-B}}$/H291$^{\text{PaE-switch-B}}$. Their mutations to Gln (to preserve polar interactions

with ATP or Mg^{2+}) strongly alter the AC/GC and CC/UC activity in Vn-/Pa-ExoY, respectively (Fig.

6C-D, 7B-C). These residues are invariant in EF or CyaA and the same mutations on the equivalent

K58$^{\text{CyaA-CA}}$ and H298$^{\text{CyaA-switch-B}}$ in CyaA (Glaser *et al*, 1989; Glaser *et al*, 1991) or H577$^{\text{EF-switch-B}}$ in EF

(Drum *et al.*, 2002) also largely reduce their CaM-stimulated AC activity. Our data show that the two

residues are correctly positioned and stabilised with two metal ions only after cofactor and substrate

binding and LID/C_B closure (Fig. 6A, 7A, S7). Further investigation is required to determine whether

H3753$^{VnE\text{-switch-B}}$ (or the equivalent histidine in other NC toxins) is critical for initiating 3'OH

deprotonation via Mg^{2+}_A, and K3528$^{VnE\text{-CA}}$ (or equivalent) for the phosphoryl transfer step in a manner

analogous to R1029$^{tmAC\text{-IIC2}}$ in tmAC. A key element in the ability of Vn-ExoY, and most likely other

NC toxins, to catalyse the cyclisation reaction of nucleotides of different sizes is the tight coordination

of the triphosphate moiety of purine and pyrimidine nucleotides after closure of their C_B/LID

subdomain. In the structure of Vn-ExoY bound to actin and 3'dCTP-$2Mg^{2+}$, the hydrogen bond network

around the ribose and base is weaker than for 3'dATP. This probably reduces the stability of pyrimidine

nucleoside binding in the NBP of activated Vn-ExoY and may explain why Vn-ExoY remains less

efficient as CC than AC. It will be interesting to compare these structural insights into the broad NC

activity of NC toxins with the functional specificities of newly identified bacterial PycC pyrimidine

cyclases, which appear to contain conserved class III cyclase catalytic elements (Tal *et al.*, 2021).

Despite the lack of tertiary and quaternary structural homology, class II (NC toxins) and class

III ACs may share some important determinants of catalysis. Upon activation, the collapse of their

different ATP/metal-ion coordination regions towards the active site brings together and constrains all

interactions with the various parts of the nucleotide substrate and the metal ions. This open-closed

transition of their catalytic domains may also facilitate the conformational changes of the substrate

predicted by computational studies during the cyclisation reaction (Mones *et al.*, 2013; Steegborn, 2014;

Hahn *et al.*, 2015; Jara & Martinez, 2016). For the catalytic domain of cofactor-activated NC toxins,

the substrate-induced fit mechanism is controlled by the C_B/LID subdomain, which is near the cofactor-

binding site. Its intrinsic flexibility and substrate-induced closure ensure both good substrate specificity

and efficient enzymatic turnover through repeated closure and opening of the C_B/LID for substrate

loading/processing and product release (Fig. 8). Without cofactor binding, substrate-induced NBP

closure remains unproductive in catalysing the reaction because switch A and B, and thus part of the

coordination of Mg^{2+}_A and the substrate nucleoside moiety (Fig. 6A, 7A), are not properly stabilised.

Our data suggest that the substrates enter the active site of NC toxins by binding through their

triphosphate moiety. Indeed, the base-coordinating switch A is initially unstable and incorrectly

positioned in ExoY toxins. This is consistent with the ability of NC toxins to use nucleotide substrates

with varying base sizes. Our study therefore provides a better understanding of how the NBP of NC

toxins is regulated and gradually switches to a catalytic conformation. It opens up new structural

perspectives for identifying ways to inhibit these NC toxins, which are involved in a wide range of

bacterial infections.

Materials and Methods

The SI Appendix includes the protein constructs, design of the chimeras used for crystallization and methods for protein expression and purification. Methodologies for pyrene-actin polymerization assays, analytical ultracentrifugation, microscale thermophoresis experiments, NC activity assays, protein crystallization and crystallography are also detailed. Data collection experiments were performed on the PROXIMA-1 and PROXIMA-2 beamlines at synchrotron SOLEIL (Saint-Aubin, France).

Data Availability. X-ray crystallography data have been deposited in Protein Data Bank (8BJH, 8BJI, 8BJJ, 8BR1, 8BO1, 8BR0). See Table S1 in SI Appendix for further details.

ACKNOWLEDGEMENTS. We thank SOLEIL for providing synchrotron radiation facilities, the staff of the PROXIMA-1 and PROXIMA-2 beamlines, and Pierre Legrand for technical assistance during data collection. We acknowledge the core facilities of Imagerie-Gif, (http://www.i2bc.paris-saclay.fr), supported by "France-BioImaging" (ANR-10-INBS-04-01). This work was supported by the ANR under ANR-18-CE44-0004 (to L.R., M.T.N., P.R., U.M.), by CNRS (to L.R., P.R., U.M., D.L.) and Institut Pasteur under #PTR 43-16 (to U.M.).

Author contributions: L.R. designed research. M.T.N, M.C., A.A.M., and L.R. performed the molecular biology or biochemical experiments, C.V. the AUC experiments, M.T.N, P.R., S.P., and L.R. the crystallisation and crystallographic experiments. D.R.B. and U.M. performed NC activity assays and generated the Pa-ExoY mutants. L.R. and M.T.N. designed the figures and wrote the paper. All authors reviewed and edited the article.

The authors declare no competing interest.

REFERENCES

Adasme MF, Linnemann KL, Bolz SN, Kaiser F, Salentin S, Haupt VJ, Schroeder M (2021) PLIP 2021: expanding the scope of the protein-ligand interaction profiler to DNA and RNA. *Nucleic Acids Res* 49: W530-W534

Ahuja N, Kumar P, Bhatnagar R (2004) The adenylate cyclase toxins. *Crit Rev Microbiol* 30: 187-196

Aszodi A, Pfeifer A, Ahmad M, Glauner M, Zhou XH, Ny L, Andersson KE, Kehrel B, Offermanns S, Fassler R (1999) The vasodilator-stimulated phosphoprotein (VASP) is involved in cGMP- and cAMP-mediated inhibition of agonist-induced platelet aggregation, but is dispensable for smooth muscle function. *EMBO J* 18: 37-48

Beckert U, Wolter S, Hartwig C, Bahre H, Kaever V, Ladant D, Frank DW, Seifert R (2014) ExoY from Pseudomonas aeruginosa is a nucleotidyl cyclase with preference for cGMP and cUMP formation. *Biochem Biophys Res Commun* 450: 870-874

Belyy A, Raoux-Barbot D, Saveanu C, Namane A, Ogryzko V, Worpenberg L, David V, Henriot V, Fellous S, Merrifield C *et al* (2016) Actin activates Pseudomonas aeruginosa ExoY nucleotidyl cyclase toxin and ExoY-like effector domains from MARTX toxins. *Nat Commun* 7: 13582

Belyy A, Santecchia I, Renault L, Bourigault B, Ladant D, Mechold U (2018) The extreme C terminus of the Pseudomonas aeruginosa effector ExoY is crucial for binding to its eukaryotic activator, F-actin. *J Biol Chem* 293: 19785-19796

Belyy A, Merino F, Mechold U, Raunser S (2021) Mechanism of actin-dependent activation of nucleotidyl cyclase toxins from bacterial human pathogens. *Nat Commun* 12: 6628

Beste KY, Spangler CM, Burhenne H, Koch KW, Shen Y, Tang WJ, Kaever V, Seifert R (2013) Nucleotidyl cyclase activity of particulate guanylyl cyclase A: comparison with particulate guanylyl cyclases E and F, soluble guanylyl cyclase and bacterial adenylyl cyclases CyaA and edema factor. *PLoS One* 8: e70223

Childers KC, Garcin ED (2018) Structure/function of the soluble guanylyl cyclase catalytic domain. *Nitric Oxide* 77: 53-64

Didry D, Cantrelle FX, Husson C, Roblin P, Moorthy AM, Perez J, Le Clainche C, Hertzog M, Guittet E, Carlier MF *et al* (2012) How a single residue in individual beta-thymosin/WH2 domains controls their functions in actin assembly. *EMBO J* 31: 1000-1013

dos Remedios CG, Chhabra D, Kekic M, Dedova IV, Tsubakihara M, Berry DA, Nosworthy NJ (2003) Actin binding proteins: regulation of cytoskeletal microfilaments. *Physiol Rev* 83: 433-473

Drum CL, Yan SZ, Bard J, Shen YQ, Lu D, Soelaiman S, Grabarek Z, Bohm A, Tang WJ (2002) Structural basis for the activation of anthrax adenylyl cyclase exotoxin by calmodulin. *Nature* 415: 396-402

Estin ML, Thompson SB, Traxinger B, Fisher MH, Friedman RS, Jacobelli J (2017) Ena/VASP proteins regulate activated T-cell trafficking by promoting diapedesis during transendothelial migration. *Proc Natl Acad Sci U S A* 114: E2901-E2910

Feltman H, Schulert G, Khan S, Jain M, Peterson L, Hauser AR (2001) Prevalence of type III secretion genes in clinical and environmental isolates of Pseudomonas aeruginosa. *Microbiology* 147: 2659-2669

Glaser P, Elmaoglou-Lazaridou A, Krin E, Ladant D, Barzu O, Danchin A (1989) Identification of residues essential for catalysis and binding of calmodulin in Bordetella pertussis adenylate cyclase by site-directed mutagenesis. *EMBO J* 8: 967-972

Glaser P, Munier H, Gilles AM, Krin E, Porumb T, Barzu O, Sarfati R, Pellecuer C, Danchin A (1991) Functional consequences of single amino acid substitutions in calmodulin-activated adenylate cyclase of Bordetella pertussis. *EMBO J* 10: 1683-1688

Gottle M, Dove S, Kees F, Schlossmann J, Geduhn J, Konig B, Shen Y, Tang WJ, Kaever V, Seifert R (2010) Cytidylyl and uridylyl cyclase activity of bacillus anthracis edema factor and Bordetella pertussis CyaA. *Biochemistry* 49: 5494-5503

Guo Q, Shen Y, Zhukovskaya NL, Florian J, Tang WJ (2004) Structural and kinetic analyses of the interaction of anthrax adenylyl cyclase toxin with reaction products cAMP and pyrophosphate. *J Biol Chem* 279: 29427-29435

Guo Q, Shen Y, Lee YS, Gibbs CS, Mrksich M, Tang WJ (2005) Structural basis for the interaction of Bordetella pertussis adenylyl cyclase toxin with calmodulin. *EMBO J* 24: 3190-3201

Gupta M, Alam S, Bhatnagar R (2006) Kinetic characterization and ligand binding studies of His351 mutants of
Bacillus anthracis adenylate cyclase. *Arch Biochem Biophys* 446: 28-34
Hahn DK, Tusell JR, Sprang SR, Chu X (2015) Catalytic Mechanism of Mammalian Adenylyl Cyclase: A
Computational Investigation. *Biochemistry* 54: 6252-6262
He C, Zhou Y, Liu F, Liu H, Tan H, Jin S, Wu W, Ge B (2017) Bacterial Nucleotidyl Cyclase Inhibits the Host
Innate Immune Response by Suppressing TAK1 Activation. *Infect Immun* 85
Howe AK (2004) Regulation of actin-based cell migration by cAMP/PKA. *Biochimica et Biophysica Acta -
Molecular Cell Research* 1692: 159-174
Hurley JH (1999) Structure, mechanism, and regulation of mammalian adenylyl cyclase. *J Biol Chem* 274: 7599-
7602
Husson C, Renault L, Didry D, Pantaloni D, Carlier MF (2011) Cordon-Bleu uses WH2 domains as
multifunctional dynamizers of actin filament assembly. *Mol Cell* 43: 464-477
Jara GE, Martinez L (2016) Anthrax Edema Factor: An Ion-Adaptive Mechanism of Catalysis with Increased
Transition-State Conformational Flexibility. *J Phys Chem B* 120: 6504-6514
Jeon J, Kim YJ, Shin H, Ha UH (2017) T3SS effector ExoY reduces inflammasome-related responses by
suppressing bacterial motility and delaying activation of NF-kappaB and caspase-1. *FEBS J* 284: 3392-3403
Jubb HC, Higueruelo AP, Ochoa-Montano B, Pitt WR, Ascher DB, Blundell TL (2017) Arpeggio: A Web Server for
Calculating and Visualising Interatomic Interactions in Protein Structures. *J Mol Biol* 429: 365-371
Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Zidek A,
Potapenko A *et al* (2021) Highly accurate protein structure prediction with AlphaFold. *Nature* 596: 583-
589
Khannpnavar B, Mehta V, Qi C, Korkhov V (2020) Structure and function of adenylyl cyclases, key enzymes in
cellular signaling. *Curr Opin Struct Biol* 63: 34-41
Khanppnavar B, Datta S (2018) Crystal structure and substrate specificity of ExoY, a unique T3SS mediated
secreted nucleotidyl cyclase toxin from Pseudomonas aeruginosa. *Biochim Biophys Acta Gen Subj* 1862:
2090-2103
Malliavin TE (2017) Molecular Modeling of the Catalytic Domain of CyaA Deepened the Knowledge of Its
Functional Dynamics. *Toxins (Basel)* 9
Mancl JM, Suarez C, Liang WG, Kovar DR, Tang WJ (2020) Pseudomonas aeruginosa exoenzyme Y directly
bundles actin filaments. *J Biol Chem* 295: 3506-3517
Martinez L, Laine E, Malliavin TE, Nilges M, Blondel A (2009) ATP conformations and ion binding modes in the
active site of anthrax edema factor: a computational analysis. *Proteins* 77: 971-983
Mones L, Tang WJ, Florian J (2013) Empirical valence bond simulations of the chemical mechanism of ATP to
cAMP conversion by anthrax edema factor. *Biochemistry* 52: 2672-2682
Morrow KA, Frank DW, Balczon R, Stevens T (2017) The Pseudomonas aeruginosa Exoenzyme Y: A
Promiscuous Nucleotidyl Cyclase Edema Factor and Virulence Determinant. *Handb Exp Pharmacol* 238:
67-85
Pimm ML, Liu X, Tuli F, Lojko A, Henty-Ridilla JL (2021) Visualizing functional human profilin in cells and in vitro
applications. *bioRxiv*: 2021.2009.2001.458557
Raoux-Barbot D, Belyy A, Worpenberg L, Montluc S, Deville C, Henriot V, Velours C, Ladant D, Renault L,
Mechold U (2018) Differential regulation of actin-activated nucleotidyl cyclase virulence factors by
filamentous and globular actin. *PLoS One* 13: e0206133
Renema P, Kozhukhar N, Pastukh V, Spadafora D, Paudel SS, Tambe DT, Alexeyev M, Frank DW, Stevens T
(2020) Exoenzyme Y induces extracellular active caspase-7 accumulation independent from apoptosis:
modulation of transmissible cytotoxicity. *Am J Physiol Lung Cell Mol Physiol* 319: L380-L390
Schoning-Stierand K, Diedrich K, Fahrrolfes R, Flachsenberg F, Meyder A, Nittinger E, Steinegger R, Rarey M
(2020) ProteinsPlus: interactive analysis of protein-ligand binding interfaces. *Nucleic Acids Res* 48: W48-
W53
Schrödinger L, DeLano W, 2020. PyMOL.
Seifert R (2017) cCMP and cUMP Across the Tree of Life: From cCMP and cUMP Generators to cCMP- and
cUMP-Regulated Cell Functions. *Handb Exp Pharmacol* 238: 3-23
Seifert R, Schirmer B (2022) cCMP and cUMP come into the spotlight, finally. *Trends Biochem Sci*

Shen Y, Lee YS, Soelaiman S, Bergson P, Lu D, Chen A, Beckingham K, Grabarek Z, Mrksich M, Tang WJ (2002)
Physiological calcium concentrations regulate calmodulin binding and catalysis of adenylyl cyclase
exotoxins. *EMBO J* 21: 6721-6732
Shen Y, Guo Q, Zhukovskaya NL, Drum CL, Bohm A, Tang WJ (2004) Structure of anthrax edema factor-
calmodulin-adenosine 5'-(alpha,beta-methylene)-triphosphate complex reveals an alternative mode of
ATP binding to the catalytic site. *Biochem Biophys Res Commun* 317: 309-314
Shen Y, Zhukovskaya NL, Guo Q, Florian J, Tang WJ (2005) Calcium-independent calmodulin binding and two-
metal-ion catalytic mechanism of anthrax edema factor. *EMBO J* 24: 929-941
Silistre H, Raoux-Barbot D, Mancinelli F, Sangouard F, Dupin A, Belyy A, Deruelle V, Renault L, Ladant D,
Touqui L *et al* (2021) Prevalence of ExoY Activity in Pseudomonas aeruginosa Reference Panel Strains and
Impact on Cytotoxicity in Epithelial Cells. *Front Microbiol* 12: 666097
Simao M, Regnier F, Taheraly S, Fraisse A, Tacine R, Fraudeau M, Benabid A, Feuillet V, Lambert M, Delon J *et*
al (2021) cAMP Bursts Control T Cell Directionality by Actomyosin Cytoskeleton Remodeling. *Front Cell*
Dev Biol 9: 633099
Skruber K, Warp PV, Shklyarov R, Thomas JD, Swanson MS, Henty-Ridilla JL, Read TA, Vitriol EA (2020) Arp2/3
and Mena/VASP Require Profilin 1 for Actin Network Assembly at the Leading Edge. *Curr Biol* 30: 2651-
2664 e2655
Steegborn C (2014) Structure, mechanism, and regulation of soluble adenylyl cyclases - similarities and
differences to transmembrane adenylyl cyclases. *Biochim Biophys Acta* 1842: 2535-2547
Suarez C, Carroll RT, Burke TA, Christensen JR, Bestul AJ, Sees JA, James ML, Sirotkin V, Kovar DR (2015)
Profilin regulates F-actin network homeostasis by favoring formin over Arp2/3 complex. *Dev Cell* 32: 43-
53
Tal N, Morehouse BR, Millman A, Stokar-Avihail A, Avraham C, Fedorenko T, Yirmiya E, Herbst E, Brandis A,
Mehlman T *et al* (2021) Cyclic CMP and cyclic UMP mediate bacterial immunity against phages. *Cell* 184:
5728-5739 e5716
Tang WJ, Guo Q (2009) The adenylyl cyclase activity of anthrax edema factor. *Mol Aspects Med* 30: 423-430
Teixeira-Nunes M, Retailleau P, Comisso M, Deruelle V, Mechold U, Renault L (2022) Bacterial Nucleotidyl
Cyclases Activated by Calmodulin or Actin in Host Cells: Enzyme Specificities and Cytotoxicity
Mechanisms Identified to Date. *Int J Mol Sci* 23
Tesmer JJ, Sunahara RK, Johnson RA, Gosselin G, Gilman AG, Sprang SR (1999) Two-metal-Ion catalysis in
adenylyl cyclase. *Science* 285: 756-760
Voth S, Gwin M, Francis CM, Balczon R, Frank DW, Pittet JF, Wagener BM, Moser SA, Alexeyev M, Housley N *et*
al (2020) Virulent Pseudomonas aeruginosa infection converts antimicrobial amyloids into cytotoxic
prions. *FASEB J* 34: 9156-9179
Yamashita H, Ueda K, Kioka N (2011) WAVE2 forms a complex with PKA and is involved in PKA enhancement
of membrane protrusions. *J Biol Chem* 286: 3907-3914
Yan SZ, Huang ZH, Shaw RS, Tang WJ (1997) The conserved asparagine and arginine are essential for catalysis
of mammalian adenylyl cyclase. *J Biol Chem* 272: 12342-12349
Ziolo KJ, Jeong HG, Kwak JS, Yang S, Lavker RM, Satchell KJ (2014) Vibrio vulnificus biotype 3 multifunctional
autoprocessing RTX toxin is an adenylate cyclase toxin essential for virulence in mice. *Infect Immun* 82:
2148-2157
710

Fig. 1. Vn-ExoY sequesters actin-ATP/ADP monomers in actin self-assembly by interacting with modest affinity. (A-C) Polymerisation time course of 4 μM ATP-Mg-G-actin (5 % pyrenyl-labelled, (A-B)) or 10 μM ADP-Mg-G-actin (10 % labelled, (C)) in the absence (black) and presence of Pa-ExoY^{K81M} (A) or Vn-ExoY^{DM} (B-C) at the indicated concentrations (μM). **(D)** Initial rates of barbed-end (dark blue) or pointed-end (light blue) growth from spectrin-actin or gelsolin-actin seeds, respectively, were measured with 1.5 μM G-actin (10 % pyrenyl-labelled) and increasing concentrations of Vn-ExoY^{DM}. Initial rates were normalised to the initial elongation rate with free G-actin from a linear fit to the first 120 s of each time course. Vn-ExoY^{DM} inhibits both barbed and pointed-end growth by sequestering G-actin with the indicated Kd (μM). **(E)** AUC-FDS measurement of the binding affinity of Vn-ExoY for G-actin. The sedimentation coefficient distribution c(s) (left panel) of Alexa-488-labelled-G-actin-ATP (60 nM) bound to the actin-polymerisation inhibitor Latrunculin A (2.5 μM) was determined in the absence and presence of increasing concentrations of unlabelled Trx-Vn-ExoY (from 0.5 to 50 μM). The resulting sw isotherms from c(s) integration are shown in the right panel. In the isotherm plots, the solid circles are the sw data from the dilution series. The solid line is the best-fit isotherm giving an estimated Kd of 11.7 μM for the interaction between Trx-Vn-ExoY and G-actin-ATP-Latrunculin-A.

Fig. 2. Vn-ExoY is activated by profilin-bound G-actin and inhibits spontaneous or regulated assembly of actin-profilin on F-actin barbed-ends *in vitro*. (A) cAMP synthesis by G-actin-activated Vn-ExoY in the presence of G-ABPs. Reactions containing 5 ng of Vn-ExoY and Latrunculin-A-bound actin at the indicated concentrations (μM) were initiated with 2 mM ATP and incubated at 30°C for 30 min. cAMP synthesis was measured in the absence (black) or presence of two high concentrations (10, 25 μM) of profilin (green), a sequestering β-thymosin (pink) or profilin-like WH2 (purple) domains. The Kd of their interaction with G-actin is 0.1, 0.5 and 0.5 μM, respectively (Husson *et al.*, 2011; Didry *et al.*, 2012). **(B)** Formation of the Vn-ExoY:actin:profilin complex by AUC-FDS. The c(s) distribution of Alexa-488-labelled-G-actin (60 nM) was determined alone (grey line) and in the presence of either profilin (8 μM; green line), Trx-Vn-ExoY (24 μM; yellow line) or both proteins (red line). **(C)** Measurement of Trx-Vn-ExoY^{DM} binding to free or profilin-bound actin from fluorescence changes in low ionic strength MST experiments. Titration of Alexa-488-labelled, Latrunculin-A-bound ATP-G-actin (0.075 μM) alone (cyan) or bound to 4 (red) and 7.9 (green) μM profilin by Trx-Vn-ExoY^{DM} for a 2-fold dilution series from 108 to 0.033 μM of Trx-Vn-ExoY^{DM} in MST-optimised buffer (see Methods). The inset shows the fluorescence changes in the raw data. The table shows the resulting Kd.

Error bars are s.d. (n≥3). **(D)** Vn-ExoY inhibits barbed-end elongation by actin-profilin. Initial barbed-

end growth rates from spectrin-actin seeds (0.36 nM) were measured as in figure 1D using 2 μM G-

actin (5 % pyrenyl-labelled) and increasing concentrations of profilin in the absence and presence of

Vn-ExoY at the indicated concentrations. **(E-F)** Vn-ExoY inhibits formin- (E) or VASP-mediated (F)

barbed-end elongation from actin-profilin. Barbed-end growth from spectrin-actin seeds (0.13 nM), 1

μM G-actin (5 % pyrenyl-labelled), and 8 μM profilin was measured in the absence or presence of 2.1

nM of capping protein (CP), 3.2 nM of FH1-FH2 domains of mDia1 formin (denoted by F) (E) or 0.22

μM of VASP (V) (F) at the indicated concentrations of Vn-ExoY (VnE).

Fig. 3. Crystal structures of Vn-ExoY in different conformational states bound to actin:profilin

or actin, starting with a chimeric protein containing three fused functional domains including Vn-

ExoY. (A) Domain diagrams of: (1) the VnE/VnE^{DM}-PRM-Prof chimera used to initiate the

crystallographic study containing Vn-ExoY extended at the C-terminus (ExoY$^{3455-3896}$, numbering from

Uniprot accession no. (AC) F0V1C5) fused to a short PRM/Proline-Rich-Motif capable of binding to

profilin, and human profilin-1 (Uniprot AC P07737), and (2) an optimised shorter Vn-ExoY$^{3455-3863}$

construct that remains as active as our original construct Vn-ExoY$^{3412-3873}$ (Belyy *et al.*, 2016). **(B)**

Synthesis of cAMP by actin-activated Vn-ExoY or VnE-PRM-Prof. VnE-PRM-Prof exhibits strong

actin-dependent AC activity that saturates at much lower G-actin concentrations. Reactions containing

5 ng Vn-ExoY or 2 ng VnE-PRM-Prof and actin at the indicated concentrations were initiated with 2

mM ATP and incubated at 30°C for 10 or 30 min. **(C)** VnE^{DM}-PRM-Prof sequestered G-actin-ATP with

100-fold higher affinity than Vn-ExoY^{DM} (residues 3412-3873) in the barbed-end elongation. Initial

barbed-end growth rates from spectrin-actin seeds were measured using 2 μM actin (3 % pyrenyl-

labelled) and increasing concentrations of VnE^{DM}-PRM-Prof (in red) or Vn-ExoY^{DM} (in blue). **(D)**

Structure of the nucleotide-free VnE^{DM}-PRM-Prof chimera bound to actin-ADP-LatB. In Vn-ExoY, the

C$_A$ subdomain and the flexible switch A, switch B, switch C and LID/C$_B$ subdomain are shown in yellow,

blue, cyan, purple and orange, respectively, here and in all subsequent figures. Disordered regions are

shown as dotted lines, the position of the ATP substrate in Vn-ExoY NBP by its surface in red and the

four actin subdomains by numbers in grey. Conserved interactions corresponding to hotspots are

indicated by black labels, with the Vn-ExoY and actin side chains shown as green and red sticks,

respectively. The sulphate ion bound in the VnE-PRM-Prof-SO$_4^{2-}$:actin-ADP structure is overlaid. **(E)**

Structure of nucleotide-free Vn-ExoY in complex with actin-ATP-LatB and profilin (Vn-ExoY-SO$_4^{2-}$

:actin-ATP-LatB:profilin). **(F)** Structure of 3'dATP-bound Vn-ExoY in complex with ATP-actin-LatB

(Vn-ExoY-3'dATP-2Mg^{2+}:actin-ATP-LatB).

Fig. 4. Sequence diagram of the structural rearrangements in actin-bound nucleotide-free Vn-ExoY upon substrate binding. (A) Porcupine plot illustrating the direction of Cα atom movement (Schrödinger & DeLano, 2020) between the nucleotide-free and 3'dATP-bound Vn-ExoY conformations. Red arrow length proportional to Cα atom motion amplitude. **(B)** Vn-ExoY conformations in the Vn-ExoY-SO_4^{2-}:actin-ATP-LatB:profilin and Vn-ExoY-3'dATP-2Mg^{2+}:actin-ATP-LatB structures (actin and profilin omitted) overlaid on their C_A subdomain. Residues with significant solvent-accessible areas (BSAs) buried by actin are indicated by yellow (BSA ≤ 35 Å^2), green (between 35 and 70 Å^2) and pink (≥ 70 Å^2) spheres on Cα. **(C)** Vn-ExoY-buried actin residues detailing contacts between proteins. The four numbered actin subdomains are shown in different colours. Vn-ExoY residues are boxed with the same colour code as in (A,B). **(D)** The LID/C_B, switch A, B and C regions of Vn-ExoY are coloured red, light blue, grey and violet in Vn-ExoY-SO_4^{2-}:actin-ATP-LatB:profilin, and orange, blue, cyan and purple in Vn-ExoY-3'dATP-2Mg^{2+}:actin-ATP-LatB, respectively. D-1) In the nucleotide-free state, switch B and the beginning of switch C, which contacts switch A and B, are ordered but show relatively high B-factors in Vn-ExoY. The structures are drawn in a cartoon putty representation (Schrödinger & DeLano, 2020), where the backbone thickness is proportional to the B-factor value. The black dotted lines represent intramolecular interactions. D-2) The conformations of switch B and C, and residues (L3530-F3531$^{\text{VnE}}$) near the LID hinge region V3532-K3533$^{\text{VnE}}$ are incompatible with 3'dATP binding or the final stabilised conformation of switch A induced by 3'dATP binding. To avoid steric clashes (red explosion symbols), they have to be rearranged. D-3) Structural rearrangement of switch B and LID. LID closure is stabilised by electrostatic bonds between K3535$^{\text{VnE-LID}}$ or R3559$^{\text{VnE-LID}}$ and D3766$^{\text{VnE-switch-B}}$ side chains. The red arrows indicate how the different regions move. D-4) Structural rearrangements of switch A and C. D-5) In this 3'dATP-bound state, switch B and C have B-factors of the same order as those in the overall structure, and the entire switch A stabilises on G-actin. Only the region of switch A (residues 3686-3719) that binds between actin subdomains 2 and 4 retains relatively high B-factors.

Fig. 5. Importance of the switch C at the actin:Vn-ExoY interface. (A) cAMP synthesis catalysed by wild-type or switch C mutants of Vn-ExoY (Vn-ExoY$^{3455-3863}$) activated by G-actin. Reactions containing 2.5 or 5 ng for Vn-ExoY and 5, 50 or 100 ng for Vn-ExoY mutants and 3 µM G-actin-ATP-LatB were started with 2 mM ATP and incubated at 30°C for 15 or 30 min. Values are averages of reactions with the indicated enzyme concentrations. **(B)** The amphipathic α-helix at the extreme C-terminus of Pa-ExoY (residues 365-376$^{\text{PaE-switch-C}}$) is shown in a helical wheel projection using the HeliQuest analysis (http://heliquest.ipmc.cnrs.fr/). **(C)** cGMP synthesis catalysed by wild-type Pa-ExoY or switch C mutants activated by F-actin. Reactions containing 5 ng Pa-ExoY$^{\text{wt}}$ or Pa-ExoY(R364A,

K347A, or H352A) or 1 µg Pa-ExoY(F367A-L371A-F374A) and 3 µM F-actin were started with 2 mM GTP and incubated at 30°C for 10 min. Error bars in (A,C) correspond to the s.d. of at least two experimental replicates. In F-actin:Pa-ExoY structure (Belyy *et al.*, 2021), the K347, H352 and R364 side chains are at distances of 3.1, 4.5 and 4.1 Å from the D25^{Actin} side chain, respectively. **(D)** Phylogenetic tree showing the relationships between ExoY-like proteins/effector domains produced by different β- and γ-proteobacteria. Vn- and Pa-ExoY are representative of the distant ExoYs. ExoYs are classified as Vn- or Pa-ExoY-like homologues based on both their sequence similarity and the predicted complex with actin by AlphaFold (Jumper *et al*, 2021) (see Fig. S10 for more details). A + sign indicates that the Vn- or Pa-ExoY residue/structural element is conserved at the same position in the ExoY-like homologue's sequence and predicted structure.

Fig. 6. Model of ATP-binding in activated Vn-ExoY and mammalian transmembrane AC (tmAC) and residues important for the purinylyl cyclase activities of Vn- and Pa-ExoY. (A) Model of ATP-binding in activated-Vn-ExoY based on its interactions with 3'dATP in the Vn-ExoY-3'dATP-2*Mg^{2+}:ATP-actin-LatB structure. Thresholds used for interaction detection are those of the PLIP (Adasme *et al*, 2021) and Arpeggio (Jubb *et al*, 2017) web servers and the PoseView tool in the ProteinsPlus web server (Schoning-Stierand *et al*, 2020). **(B)** The model-phased anomalous difference Fourier map from data collected at 6.55 keV, shown as a green mesh contoured at 5.0 σ, confirms the presence of two Mn^{2+} (grey spheres) bound to 3'dATP in Vn-ExoY NBP. **(C)** cAMP synthesis catalysed by wild-type or mutants of Vn-ExoY (Vn-ExoY$^{3455-3863}$) activated by G-actin. Reactions containing 10 ng Vn-ExoY, 100 ng Vn-ExoY(K3533Q) or 1 µg Vn-ExoY(K3528Q, K3535Q, or H3753Q) and 3 µM of LatB-Mg-G-actin were started with 2 mM ATP and incubated at 30°C for 10-30 min. **(D)** cGMP synthesis catalysed by wild-type or mutants of Pa-ExoY activated by F-actin. Reactions containing 10 ng Pa-ExoY^{wt}/Pa-ExoY(K86Q or K88Q) or 8 µg Pa-ExoY(K81Q or H291Q) and 3 µM of Mg-F-actin were started with 2 mM GTP and incubated at 30°C for 10-30 min. **(E)** The LID/C$_B$ subdomain of EF occupies different positions relative to the C$_A$ subdomain between the nucleotide-free (1xfz, 1xfy) (Shen *et al.*, 2005) and nucleotide-bound (1k90) (Drum *et al.*, 2002) structures of CaM-bound EF. It rotates between the 1xfz and 1k90 PDBs by 16° as a rigid body. The structures are superimposed on their C$_A$ subdomains shown in yellow. **(F)** Model of ATP-binding in activated class III tmAC based on the structure of the type V AC C1a/type II AC C2 heterodimer bound to the ATP analogue RP-ATPαS (PDB 1CJK) (Tesmer *et al.*, 1999). Activated Vn-ExoY and tmAC closed NBPs have similar interactions with ATP analogues. For example, their conserved asparagine N3759$^{VnE-switch-B}$ or N1025$^{tmAC-IIC2}$ similarly hydrogen-bonds the ribose ring O4'-oxygen of ATP analogues. In tmAC, the interaction of N1025^{tmAC-}

[IIC2] with ATP is essential for correct ribose positioning during catalysis (Yan *et al.*, 1997; Tesmer *et al.*,

1999).

Fig. 7. Model of CTP-binding in actin-activated Vn-ExoY and residues important for the CC

or UC activity of Vn- and Pa-ExoY. **(A)** Model of CTP-binding in activated-Vn-ExoY based on its

interactions with 3'dCTP in the Vn-ExoY-3'dCTP-2*Mn^{2+}:ADP-actin structure. Interaction detection

thresholds are as in Fig. 6A, F. **(B)** cCMP synthesis catalysed by wild-type or mutants of Vn-ExoY

activated by G-actin. Reactions containing 100 ng Vn-ExoYwt or 1 µg-100 ng (mean) Vn-

ExoY(K3528Q, K3533Q, K3535Q, or H3753Q) and 6 µM of Lat.B-Mg-G-actin were started with 2

mM CTP and incubated at 30° C for 60 min. **(C)** cCMP and cUMP synthesis catalysed by wild-type or

mutants of Pa-ExoY activated by F-actin. cCMP synthesis was performed by using 1µg-100 ng (mean)

Pa-ExoY[wt] or Pa-ExoY(K81Q, K86Q, K88Q, or H291Q) and 6 µM of Mg-F-actin and reactions were

started with 2 mM CTP and incubated at 30° C for 60 min. cUMP synthesis was performed by using

100-200 ng (mean) Pa-ExoY[wt] or 1000-100 ng (mean) Pa-ExoY(K81Q, K86Q, K88Q, or H291Q) and

6 µM of Mg-F-actin and reactions were started with 2 mM UTP and incubated at 30° C for 60 min. In

the absence of G-/F-actin, the basal CC and UC activity of Vn- and Pa-ExoY are 0.007 and 0.003

pmol/min/ng of the enzyme, respectively.

Fig. 8. Model of the activation mechanism and cytotoxic effects of Vn-ExoY elicited by its

interaction with actin:profilin. **(A)** Once secreted by bacterial T1SS, the MARTX toxin binds to the

target cell membrane, translocates and releases its various MARTX effector modules into host cells

(Ziolo *et al.*, 2014), including Vn-ExoY. **(B)** In isolation, Vn-ExoY's three switch regions and CB/LID

domain are flexible and unstable, and the enzyme is inactive. **(C)** The interaction of Vn-ExoY with the

actin:profilin complex is first tightly docked to actin by switch C. This first step brings together and

stabilises only part of the substrate and metal ion binding sites: it initiates stabilisation of switch B and

brings switch A closer to the NBP. This intermediate activated conformation allows the active site to

'breathe', particularly at the C_B/LID subdomain of NC toxins, which is close to the cofactor binding site.

The C_B/LID subdomain of NC toxins can therefore continue to adopt a variety of positions. **(D)**

Substrate and metal ion binding promotes NC toxin C_B/LID closure. This allows the cofactor pre-

stabilised active site of NC toxins to accommodate and ultimately constrain all substrate and metal ion

interactions. By simply closing and opening the C_B/LID to load and process nucleotide substrates of

different sizes (purine/pyrimidine nucleotides) and release the products, the large substrate-induced fit

movement of the NC toxin C_B/LID subdomain ensures both good substrate specificity and efficient

enzymatic turnover. **(E)** The formation of an abortive ternary complex, recruited by ABPs through their

PRMs (present in formin FH1) and sequestering G-actin:profilin, may act in two ways. It confines Vn-
ExoY's toxic NC activity to sites of active actin remodelling. At these sites, G-actin:profilin and
elongation-promoting factors, in close cooperation with cAMP-dependent signalling, fine-tune the
homeostasis of F-actin networks (Howe, 2004; Yamashita *et al.*, 2011; Skruber *et al.*, 2020; Simao *et*
al., 2021). It may also locally interfere with the activity or regulation of various elongation factors, such
as VASP, Arp2/3 or formin (shown here inducing membrane protrusions), if the proper functioning of
their PRMs, which deliver G-actin to the barbed ends of F-actin, is impaired.

Figure 1

Figure 2

Figure 3

Figure 4

A   Vn-ExoY

LID
sw. B
α-helix I
α-helix B
K3683^VnE
D3682^VnE
sw. A
L3827^VnE
R3824^VnE
sw. C

BSA
70 Å²
35 Å²

B   Vn-ExoY

25° LID
LID
sw. A
sw. B
sw. C
L3827^VnE
R3824^VnE
Y3850^VnE
R3854^VnE
V3857^VnE
L3861^VnE
K3860^VnE

C   Actin

P243^Actin
S3703^VnE
P3704^VnE
R62^Actin
D3701^VnE
D56^Actin
K3683^VnE
E93^Actin
D25^Actin
R3824^VnE
I345^Actin
Y3850^VnE
R3854^VnE
L349^Actin
R3854^VnE
E334^Actin
Y3675^VnE
ATP
Actin

D-1

LID
K3554
Switch B
SO₄
K3533
L3530
D3757
N3832
H3753
A3755
D3736
Switch A
Switch C
L3827
Actin

Substrate binding

D-2

K3554
3'dATP
K3533
S3723
N3832
A3755
L3827
Y3675
D3682
D3736
Switch A

D-3

LID
25°
LID
R3559
Mg_B
Mg_A
D3766
H3753
S3826
R3824
N3828
S3822
L3678
Y3675
Q3681

Switch B and LID rearrangements

Switch C and A rearrangements

D-4

R3559
D3766
Switch B
Switch C
N3832
L3827
D3682
Switch A

D-5

R3559
H3753
D3766
S3723
L3724
M3726
A3758
N3832
L3827
R3824
S3822
Y3675

Figure 5

A

cAMP (µmol/min/mg)

509.1
3.7
2.1

Vn-ExoY ^wt / mutants

□ wild-type (wt)
▨ R3824A
▦ Y3850A-R3854A-V3857A-K3860A-L3861A

B

Pa-ExoY extreme C-term. (P365-R376)

N-ter (P365)   C-ter (R376)

F373

L371

F367   F374

hydrophobic face

C

cGMP (µmol/min/mg)

599.9
465.8
40.8   79.3
0.05

Pa-ExoY ^wt / mutants

□ wild-type (wt)
▨ R364A
▧ K347A
▦ H352A
■ F367A-L371A-F374A

D

	G-/F-actin specificity (G/F)	Residues for salt bridge with Asp25^Actin				C-terminal amphipathic α-helix
		Pa-ExoY-like			**Vn-ExoY-like**	
		K347	H352	R364	R3824	
Cedecea neteri ExoY-L	F	+	+	S		+
Yersinia aldovae ExoY-L	F	+	+	N		+
Chromobacterium vaccinii ExoY-L	F	+	+	+		+
Providencia stuartii ExoY-L	F	+	+	+		+
Salmonella enterica ExoY-L	F	R	Y	S		+
Burkholderia pseudomallei ExoY-L	F	+	+	+		+
Pseudomonas reactans ExoY-L	F	+	+	+		+
Pseudomonas viridiflava ExoY-L	F	+	+	+		+
Pseudomonas sivasensis ExoY-L	F	+	+	+		+
Pseudomonas grimontii ExoY-L	F	+	Y	+		+
Pseudomonas fluorescens ExoY-L	F	+	+	+		+
Pantoea cypripedii ExoY-L	F	+	Y	S		+
Candidatus hamiltonella defensa ExoY-L	F	+	Y	S		+
Iodobacter fluviatilis ExoY-L	F	+	+	+		+
Aeromonas salmonicida ExoY-L	F	+	+	+		+
Pseudomonas aeruginosa ExoY (Pa-ExoY)	F	+	+	+		+
Vibrio anguillarum ExoY-L	G				+	+
Vibrio ordalii ExoY-L	G				+	+
Vibrio vulnificus ExoY-L (Vv-ExoY)	G				+	+
Vibrio cholerae ExoY-L	G				+	+
Vibrio nigripulchritudo ExoY-L (Vn-ExoY)	G				+	+
Vibrio neptunius ExoY-L	G				+	+
Psychromonas sp.Urea-02u-13 ExoY-L	G				+	+
Vibrio caribbeanicus ExoY-L	G				+	+
Proteus vulgaris ExoY-L	G				+	+
Providencia alcalifaciens ExoY-L	G				+	+
Providencia rettgeri ExoY-L	G				+	+
Escherichia coli ExoY-L	G				+	+

0.2

Figure 6

Figure 7

A

Flexible regions dependent on:
substrate
cofactor

C_A | LID/C_B | switch B | switch A

--- π-π interaction
⋮⋮⋮⋮ hydrophobic interaction
--- carbon-π interaction

B

Vn-ExoY^wt/mutated

cCMP (μmol/min/mg)

wt	K3528Q	K3533Q	K3535Q	H3753Q
4.8	0.01	0.03	0.01	0.01

cCMP

C

Pa-ExoY^wt/mutated

cCMP/cUMP (μmol/min/mg)

cCMP / cUMP

	wt	K81Q	K86Q	K88Q	H291Q
cCMP	0.8	0.01	0.1	0.04	0.01
cUMP	2.9	0.02	0.9	0.2	0.02

Figure 8

Host cell membrane

(+) end elongation

Vibrio nigripulchritudo (Vn) bacteria

A   T1SS MARTX

B

LID/C_B

inactive Vn-ExoY

switch A

switch B

C_A

switch C

cofactor-recruitment

(-) end

C   LID

ATP or CTP

25°

D   LID

E   sequestration

(+) end

(-) end

cAMP
cCMP

host cell intracellular signalling disruption

Vn-ExoY

G-actin

F-actin

Profilin

Formin
FH1-FH2