Title: The fitness consequences of wildlife conservation translocations: A meta-analysis

Authors: Iwo P. Gross[1, *], Alan E. Wilson[2], & Matthew E. Wolak[1]

[1]Department of Biological Sciences, Auburn University, Auburn, AL 36849, USA

[2]School of Fisheries, Aquaculture, and Aquatic Sciences, Auburn University, Auburn, AL

36849, USA

[*]Corresponding author: ipg0004@auburn.edu

Running title: Conservation translocation efficacy meta-analysis

Keywords: *Captive-breeding, captive-rearing, head-starting, animal relocation, animal*

reintroduction, ex situ reinforcement, in situ reinforcement

Abstract

Conservation translocation is a common strategy to offset mounting rates of population declines

through the transfer of captive-reared or relocated organisms to imperiled populations. However,

these programs have been criticized for low success rates and husbandry practices that produce

individuals with genetic and performance deficits. We assessed the disparity in post-release

performance of translocated organisms relative to wild-resident conspecifics through a meta-

analysis of 826 performance comparisons from 172 studies representing nine animal classes (101

species). We found that translocated organisms have 70% decreased odds of out-performing their

wild-resident counterparts, supporting published claims of systemic issues hampering

conservation translocation. Pre-release animal enrichment significantly reduced performance

disparities, whereas our results suggest no overall effects of captive generation time or the type

of fitness surrogate measured. Conservation managers should consider evaluations of post-

release performance using sympatric (resident-control) and allopatric (true-control) wild-

reference groups to ensure translocated animals contribute to recipient population growth and

viability.

Introduction

Rates of floral and faunal species loss in the new Anthropocene signal the onset of a sixth

global mass extinction [1,2]. Among vertebrates, extinction rates in the last century increased 100

times above pre-industrial levels [3] and recorded population size declines have averaged 68% in

the last 50 years [4]. Continued species loss will have irreversible consequences for ecosystem

services and resiliency. Slowing the rate of species loss will require eliminating existing threats

(e.g., disease, poaching) and extensive habitat preservation and restoration [5,6]. While

environmental threats are being addressed, supportive breeding and translocation programs have

been proposed to maintain species and their roles in the ecosystem [7,8].

Animal conservation translocations (ACTs) are a common strategy intended to directly

offset species extinction through the supplementation of threatened populations and the

repatriation of extirpated populations [9]. ACT programs supplementing extant populations in their

indigenous range are called reinforcements, and can be grouped into two categories [10,11]. First, *in*

situ reinforcements (ISRs) collect and transport wild organisms to another natural area for the

benefit of the recipient conspecific population. Second, *ex situ* reinforcements (ESRs)

incorporate a captive phase lasting from less than one generation (i.e., head-starting) up to

several generations (i.e., captive breeding) before release. Despite immense efforts across a wide

array of taxa to employ ACTs and document their efficacy in the literature, it is still unknown

whether translocated individuals are as capable as their wild conspecifics at contributing to

population viability and preventing extinctions.

To date, ACTs have had many notable successes and a quantifiable impact on global

extinction rates [12,13]. Nevertheless, ACTs are often considered a halfway technology with

logistical and systemic shortcomings that have led to generally low success rates [14,15]. In

particular, multiple studies indicate that ACTs might produce individuals with inherent genetic

and performance-related deficits that compromise ACT program goals and reduce recipient

population viability [16,17]. These deficits are most severe in ESRs involving multiple captive

generations since the capture of ESR founders acts as a bottleneck on genetic diversity. Even

with a stringent captive breeding structure, factors inherent to husbandry practices (e.g., small

effective population size, inbreeding depression, relaxed purifying selection) increase the

likelihood of fixation of undesirable alleles and a loss of allelic heterozygosity [7,18,19]. A large

influx of captive-born individuals into the wild with such genetic properties can swamp genetic

diversity of locally adapted populations and reduce effective population sizes despite an overall

increase in numbers [20]. However, such genetic deficits require multiple generations of captive

breeding to accumulate which suggests alternative factors contribute to fitness reductions

observed in ESRs lasting a generation or less [17,21].

A fundamental criticism of ESRs is that non-natural conditions present in captivity can

influence individual phenotypes, diminishing post-release performance relative to wild

conspecifics [19,22,23], where animal performance is defined as a measure of some critical life

history character (e.g., growth rate, survival, fecundity) that correlates with individual fitness [24].

In a variety of taxa, ESRs lasting a generation or less negatively influence post-release

performance metrics such as anti-predator behavior [25], dispersal [26], growth [27], survival [28], and

reproductive success [21]. Such phenotypic mismatches can arise via domestication selection,

where trait values that are selected for in captivity are maladaptive when expressed in a natural

environment [29,30]. The unintentional selection for domesticated phenotypes in artificial

environments is likely a major contributor to reduced performance of ESRs, since it can alter

phenotype distributions within a single generation [21,27]. In the absence of domestication selection,

phenotypic plasticity during less than one generation of captive development and rearing can

lead to similar deviations from the wild-type in anatomical, physiological, and behavioral traits

[29,31]. Moreover, certain plastic phenotypic deviations have an epigenetic basis, thereby

potentially influencing performance of future generations regardless of their developmental

environment [32].

By virtue of the use of wild-born organisms and lack of a captive component, ISRs are

lauded as a less intrusive and a generally more successful translocation strategy than ESRs [14,33–

35]. However, failure rates for ISRs are still notably high and potentially a consequence of similar

issues that affect ESRs. At the population-level, supplementing existing populations with

conspecifics from other wild populations can lead to maladaptive gene flow or outbreeding

depression of the recipient population and a reduction in meta-population genetic diversity [36,37].

At the individual-level, relocation can impact post-release performance either directly through

transportation-related stress [38] or indirectly through energetic costs of establishment [39], errant

dispersal and homing behaviors [40], diminished social structure [41], and phenotypic mismatches

within the novel ecosystem [31].

Exposing captive organisms to some form of enrichment is the most intuitive strategy to

counteract phenotypic mismatch, and a reasonable compromise when ISRs are not possible. We

use the term enrichment broadly to refer to a variety of supportive measures [14] employed by

ACTs to offset any maladaptive behavior, physiology, or physical injury incurred during

captivity and/or translocation [23,41]. Some examples include fostering pre-release organisms with

experienced conspecifics [42], simulating *in situ* environments with naturalistic enclosures or

antipredator training [43], implementing a soft-release phase where supplemental feeding and/or

shelter are provided [44], or simply reducing the length of the captive phase [26,45]. Literature reviews

suggest that enriched ESR cohorts are more successful post-release compared to un-enriched

cohorts [14,23], but the lack of a wild-reference group in the synthesized findings makes it difficult

to conclude whether enrichment strategies produce translocated organisms that are

indistinguishable from wild conspecifics with respect to post-release fitness [46].

Finally, the metric used to gauge relative performance can vary in its representation of

fitness, thereby shaping study conclusions [24]. The lifetime reproductive success of an individual

is the ideal measure of fitness when focusing on population dynamics and phenotypic adaptation,

but lifetime reproductive success and the component life history traits that determine it (e.g.,

survival and fecundity) are difficult to measure in natural conditions. Consequently, fitness is

often approximated using traits that typically correlate with fitness (e.g., body size, body

constitution, movement). However, the extent to which correlates and components of fitness

covary and ultimately relate to lifetime reproductive success remains uncertain [24]. For example,

Mulder et al. found that while translocated and wild-resident desert tortoises (*Gopherus*

agassizii) had similar body condition and survivorship four years post-release, translocated males

fathered none of 92 offspring for which paternity was determined [47]. This suggests that survival

is a potentially inappropriate component of fitness to gauge ACT success in adult organisms with

high adult survivorship. Conservation translocations should consider multiple relevant fitness

components during project assessment to identify potential life-history trade-offs and avoid

misleading conclusions [24].

The translocation literature has grown to a point that quantitative reviews are necessary to

summarize the high volume of papers, while also accounting for heterogeneity and bias [48].

Seminal reviews have evaluated ACT efficacy from the published literature and via mail-in

questionnaires sent to conservation program managers [11,14,33]. In both approaches, the response

variable (translocation success versus failure) reduces a highly nuanced outcome to a binary

classification based in part on unstandardized criteria (e.g., creation of a self-sustaining

population) and professional opinion. These syntheses are an invaluable resource for applied

conservation and management, but their largely narrative format is not conducive to quantitative

synthesis across broad and multifactorial datasets while accounting for bias and heterogeneity

among studies. Meta-analysis is well equipped to gauge translocation efficacy based on the post-

release performance of animal subjects [23,43,49–51]. Previous meta-analyses are limited to

comparisons among translocated cohorts and fail to include a wild-reference group (but see [22]),

therefore, they cannot address whether translocated individuals are as capable as their wild

conspecifics at contributing to population viability.

Here, we present the first comprehensive meta-analysis to assess conservation

translocation programs through the synthesis of studies involving direct, post-release

comparisons of translocated and wild-resident conspecifics. We further expand on previous

reviews by sampling the literature broadly across macroscopic animal groups and assess relative

post-release performance using multiple fitness surrogates. Our primary objective is to quantify

the disparity in post-release performance of translocated organisms relative to wild-resident

conspecifics across animal taxa and to determine whether several organismal and methodological

factors influence this disparity. Our systematic review and meta-analysis show that across a

broad array of taxa (826 effect sizes from 101 species representing 49 families), translocated

organisms have 70% decreased odds of out-performing their wild-resident counterparts. Pre-

release animal enrichment strategies significantly reduced performance disparities among

translocated and wild organisms, whereas *in situ* vs. *ex situ* reinforcement strategy, captive

generation time, and the type of fitness surrogate measured did not influence relative post-release

performance. Conservation managers should standardize the use of wild-resident controls and

multiple performance metrics in their assessments of potential conservation translocation

strategies.

Table 1. Total number of log odds ratio (LOR) estimates collected, organized by taxonomic class

and the fitness surrogate used as a metric for performance.

	Movement	Constitution	Growth	Survival	Reproduction	**Total**
Bivalvia	0	0	0	6	0	**6**
Gastropoda	0	3	5	9	6	**23**
Malacostraca	0	4	0	14	5	**23**
Echinoidea	1	1	0	0	0	**2**
Actinopterygii	38	43	26	158	81	**346**
Amphibia	4	2	0	3	0	**9**
Reptilia	101	12	26	55	9	**203**
Aves	14	12	0	59	38	**123**
Mammalia	18	13	2	46	12	**91**
Total	**176**	**90**	**59**	**350**	**151**	**826**

Results

After filtering based on the inclusion criteria, our dataset comprised 826 unique effect

sizes for relative performance of translocated organisms and their wild-resident conspecifics

(223 ESR and 603 ISR comparisons) collected from 172 peer-reviewed studies investigating 101

unique species (Table 1; Supplementary Table S1). We calculated standardized effect sizes as

log odds ratios (LORs) of performance metrics for translocated versus wild-resident cohorts such

that positive values indicate translocated individuals exhibited higher relative performance than

wild-resident conspecifics. We analyzed the LORs and subsets of these data using three linear

mixed effects meta-analytical models with fixed effect moderator variables to determine the

impact of organismal and methodological factors plus random effects to attribute sources of

variation among effect sizes.

Translocated organisms generally exhibited poorer performance following release

compared to wild-resident conspecifics. The primary model's overall log odds ratio (LOR

mode: -0.98; CrI: -1.62 to -0.36) was converted to an odds ratio by exponentiating across the

posterior distribution, indicating that translocated organisms have 69.9% decreased odds of out-

performing their wild-resident counterparts (OR mode: 0.30; CrI: 0.18 to 0.66). By quantifying

the proportion of posterior samples with LOR estimates greater than or equal to zero, we infer

that translocated organisms had an overall 0.002 probability of out-performing wild-resident

conspecifics following release (Fig. 1). Disparity in post-release performance did not vary widely

across taxonomic classes, but instead all classes generally exhibited negative LOR effect sizes,

suggesting poor performance of translocated organisms relative to wild-residents across taxa

(Supplementary Table S2). Similarly, relative post-release performance was largely unaffected

by ESR vs. ISR translocation strategy.

In studies involving mixed-sex cohorts, comparisons based on fitness components tend to

have lower LOR marginal posterior mode estimates than comparisons based on fitness

correlates, but widely overlapping credible intervals across fitness components and correlates

indicate a consensus in the interpretation of relative under-performance of translocated cohorts

across a variety of performance metrics (Supplementary Table S2). More variable effect size

outcomes observed in sex-specific comparisons might suggest the presence of unequal life-

history consequences to translocation among sexes, but low sample sizes in constitution-,

growth-, and survival-based estimates, and uneven taxonomic representation in movement-based

estimates preclude conclusive interpretations.

Due to limited reporting of key moderator variables across studies, we analyzed subsets

of the data in two additional models to assess whether the amount of captive ancestry or the

presence of enrichment, respectively, influenced relative post-release performance of

translocated cohorts. In general, LOR marginal posterior mode estimates were similar for the

overall model effects and for parameters that also appeared in the primary model 1 above

(Supplementary Tables S3-S4). Therefore, we limit the remainder of our reporting to model-

specific parameters.

Post-release performance was generally lower among translocated cohorts

compared to wild residents, regardless of the amount of captive ancestry. Post-release

performance was largely unaffected by the length of captive ancestry, except for comparisons

based on constitution and growth. In ISR studies involving constitution-based comparisons, we

observed that relocated organisms without any captive ancestry had a lower probability (0.00;

OR mode: 0.06; OR CrI: 0.03 to 0.21) of out-performing wild-resident conspecifics relative to

head-started (0.15; OR mode: 0.58; OR CrI: 0.24 to 1.33) and captive-bred cohorts (0.25; OR

mode: 0.28; OR CrI: 0.03 to 2.49; Supplementary Table S3). Comparisons with growth-based

estimates exhibited a similar pattern, but these growth-related comparisons for relocated, head-

started, and captive-bred cohorts were based on reduced datasets of only two, eight, and one

studies, respectively.

Post-release performance of translocated organisms generally benefitted as a result

of enrichment. To test the influence of enrichment strategies on relative post-release

performance among translocated and wild cohorts, we analyzed a subset of studies that gathered

data from both enriched and unenriched cohorts within the same experimental design and

compared their post-release performances relative to a common wild-control group. Unenriched

translocated organisms had a 0.07 probability of out-performing their wild-resident conspecifics

(OR mode: 0.07; CrI: 0.003 to 1.16), whereas enriched translocated organisms had a 0.58

probability of out-performing their wild counterparts (OR mode: 0.55; CrI: 0.01 to 9.95; Fig. 2;

Supplementary Table S4).

The dataset exhibited a high overall taxonomic signal but low heterogeneity at

individual taxonomic levels. Here we summarize the heterogeneity statistics from model 1,

representing the full dataset (Supplementary Table S5 for heterogeneity summary statistics of all

three models). Heterogeneity values from individual taxonomic levels were relatively small and

even in magnitude, but their combined taxonomic signal accounted for 59.9% (CrI: 46.8% to

74.4%) of total heterogeneity. Study- and group-level effects accounted for 21.8% (CrI: 12.0% to

34.3%) and 11.9% (CrI: 8.14% to 18.4%) of total heterogeneity, respectively.

Discussion

Our meta-analysis gauged the efficacy of conservation translocation programs based on

direct, *in situ* comparisons of animal performance between translocated organisms and wild-

resident conspecifics. We synthesized estimates of relative performance based on five fitness-

surrogate categories across 101 unique species spanning nine taxonomic classes. Our findings

indicate that translocated organisms tend to under-perform relative to their wild counterparts, and

that this pattern is largely independent of taxonomic class or the type of fitness surrogate

measured. This supports previous claims for the existence of systemic issues across ACTs and

suggests that translocated organisms are often at a major disadvantage following release. Under-

performance of translocated cohorts was also consistent regardless of translocation strategy or

the amount of captive ancestry, with some caveats. Finally, post-release performance was

markedly improved in translocated cohorts exposed to pre-release enrichment relative to

unenriched cohorts. Here we explore the implications of our findings and make

recommendations for future translocation program assessment and monitoring procedures.

The unexpected lack of variation in effect sizes across taxonomic classes hints at a

general influence of ACTs on animals, which is in direct odds with the large proportion of study

heterogeneity explained by taxonomic rank (Supplementary Table S5). The high taxonomic

signal (64.29%) indicates that taxonomic position explains a majority of variation in effect sizes,

which would lead us to expect large differences in relative performance values among higher

taxonomic groups. For example, Dochtermann et al. observed a consistent pattern in a

phylogenetic meta-analysis quantifying behavioral heritability: Little difference in effect size

among species and a low phylogenetic signal [52]. However, we see the proportion of

heterogeneity explained by individual taxonomic tiers is consistently low, which could explain

the minimal differentiation in effect sizes at the level of taxonomic class. This suggests that

heterogeneity incurred at different taxonomic levels is not necessarily cumulative and might be

more accurately quantified in a phylogenetic framework. Regardless, comparative studies

assessing biological patterns at the broadest taxonomic scales should adopt a mixed phylogenetic

and taxonomic approach until an appropriate phylogeny with empirically-derived branch lengths

can be resolved [53,54].

Our meta-analysis highlights a clear bias in taxonomic representation among ACTs, with

most studies involving charismatic species or those with economically important fisheries. Most

notably, there is a significant lack of ACTs involving amphibians that meet our inclusion criteria.

Amphibian populations are experiencing global declines and will likely require human

interventions in the form of ACTs to avoid extirpation [40]. Recent quantitative reviews describe a

positive trend in the success of amphibian translocations relative to earlier benchmarks [40,55].

However, these efforts fail to consider comparisons among translocated and wild-resident

conspecifics as a potential index for project feasibility. Recent amphibian studies using this

comparative approach have provided novel insights regarding the multi-generational influence of

ACTs as well as potential pitfalls of cryopreservation techniques on ACT feasibility [56,57]. Future

assessments of translocation success should include simultaneous monitoring of a wild-control

group from the donor and/or recipient populations as an additional criterion for determining

project feasibility and success.

Our primary model showed high consistency in effect sizes across fitness surrogates,

indicating that phenotypic traits considered as correlates of overall fitness have the potential to

characterize performance differences among translocated and wild-resident cohorts. We initially

predicted that fitness correlates would exhibit less consistent disparities across animal cohorts

since fitness correlates are less reliable proxies of fitness than fitness components [58]. One

explanation for the observed pattern is that fitness surrogates can vary in how well they reflect

overall individual fitness based on the relative length and positioning of the measurement

interval within the organism's lifespan [24]. Additionally, the lack of differences in fitness

surrogate effects may be due to unaccounted-for differences in life-history strategies within and

among study systems. Life-history theory suggests that the uneven allocation of finite resources

results in life-history trade-offs among fitness components such that maximizing one fitness

component can decrease overall lifetime reproductive success (e.g., increased annual fecundity

decreases annual survival)[59]. Future ACT studies and meta-analyses on the topic should

incorporate multiple fitness components at different life stages to assess relative quality of

released organisms, such that the covariance among fitness components and thus the presence of

fitness trade-offs or differing life-history strategies can be tested at the within-study level [24].

We were not able to evaluate sex-by-fitness surrogate interactions in the full dataset

because so few studies reported sex-specific effects. However, sufficient data from reptile studies

suggest that estimates of relative performance based on movement were substantially lower for

both male- and female-only cohorts relative to estimates from mixed-sex cohorts, indicating

greater disparity in post-release performance relative to wild conspecifics when males and

females were analyzed separately. Only when accounting for sex do our findings align with the

established relationship between maladaptive post-release movement patterns and reduced

survival in the herpetofaunal literature [40,60]. The influence of sex on post-release performance is

also poorly represented in the primary literature and rarely synthesized in subsequent reviews [61].

To our knowledge, no other quantitative literature review or meta-analysis examining

translocation efficacy has included sex as a moderating factor, despite well-documented and

systematic differences in life history and ecology among the sexes [62].

We observed similar levels of under-performance among translocated cohorts relative to

wild conspecifics, regardless of whether translocated organisms were ISR (0 captive

generations), head-started (ESR with < 1 captive generation), or captive-bred (ESR with > 1

captive generation). This contrasts with existing quantitative reviews describing a negative

relationship between captive ancestry and fitness, with ISRs performing relatively better than

ESRs [14,35]. One explanation for this discrepancy is the poor reporting in the literature of ESR

captive histories and animal origins which precluded us from conducting a more powerful

assessment of the effect of captive ancestry. As such, our head-starting vs. captive-rearing levels

possibly misclassified cohorts with ambiguous or mixed captive histories, which could explain

the lack of significant performance deficits commonly reported in long-term captive-breeding

populations [21]. Additionally, past reviews have used an arbitrary measure of success to

characterize performance, which could skew model outcomes. Our findings suggest that while

ISRs avoid the maladaptive selective pressures associated with extended captivity, managers

should still be wary of individual- and population-level stressors characteristic of ISRs that can

lead to disparities in post-release performance like those observed in ESRs. This underscores the

necessity of ACT programs to report the capture histories of their organisms, and to conduct

multi-generational monitoring to ensure accurate measures of meaningful life-history traits.

Disparity in post-release performance among translocated and wild individuals was

significantly reduced when translocated organisms were exposed to some form of pre-release

enrichment. This finding highlights the growing relevance of the behavioral ramifications of *in

situ* and *ex situ* reinforcements, where genetic management has historically been the only factor

informing best husbandry and breeding practices [16,46]. Our findings agree with those from other

meta-analyses showing that enriched, translocated organisms generally had higher performance

than non-enriched, translocated conspecifics in a variety of taxa [23,50]. Since few studies

incorporating pre-release enrichment also fulfilled our inclusion criteria by directly comparing

translocated cohorts to wild-resident conspecifics, we were not able to refine our categorization

of different enrichment strategies. Despite that, our study provides the broadest evidence yet—

and the first synthesized evidence to incorporate wild-control groups—that pre-release

enrichment strategies have the potential to reduce post-release performance disparities between

translocated and wild-resident organisms.

Conservation biology is in the midst of a revolutionary push for more evidence-based

decision-making to replace historically anecdotal- or experience-driven strategies [63]. Formal

meta-analysis is an important tool for synthesizing existing literature and directing new veins of

inquiry, particularly in the field of conservation, where translocation studies are numerous but

statistical power is limited by low sample sizes [64]. Our findings show that *in situ* reinforcements

(ISRs) and *ex situ* reinforcements (ESRs) produce organisms that under-perform and should

therefore be implemented only after careful consideration of the recipient population's trajectory

and the existing environmental threats at the recipient site. For example, if a wild population is

declining due to extrinsic threats, then the release of under-performing translocated conspecifics

is unlikely to reverse declines [5]. Alternatively, if environmental threats have been removed, then

the allelic diversity of a remnant wild population could benefit from the influx of translocated

cohorts, despite certain performance deficits [13,56]. Pilot studies employing the post-release

comparative techniques analyzed in this study could therefore use relative post-release

performance, population demographics, and environmental forecasts to predict the efficacy of

ACT programs. Future conservation efforts must clarify their goals and consider other

ecosystem-level restoration or mitigation efforts as alternative or complementary strategies to

ACTs.

334   Despite a number of successful efforts, animal conservation translocations have

experienced historically low success rates, without significant improvement observed over time

[65]. That said, the often-wide credible intervals in our findings suggest there are opportunities in

which ACTs can produce viable individuals and contribute to recipient populations. We

recommend that ACT managers include sympatric (resident-control) and allopatric (true-control)

wild-reference groups in their programs to properly evaluate post-release performance of

translocated organisms, and assess any negative density-dependent impacts of ACTs on resident

conspecifics [46,66]. Managers should also take advantage of next-generation sequencing

technologies and parentage assignment software to promote long-term, multi-generational

monitoring of relative reproductive success and other fitness surrogates [21]. Finally, we encourage

ACT managers to explore the natural history of their study organisms for tailored opportunities

to maximize post-release performance by implementing pre-release behavioral or environmental

enrichment.

Methods

Literature search and inclusion criteria

350   We conducted our literature review, screening, and analysis according to PRISMA

guidelines (Supplementary Fig. S1)[67]. We searched the literature using Web of Science,

including all databases (1/1/1900-8/23/2019), for peer-reviewed studies concerning conservation

translocations with some reference to a wild-resident group used as a control. Our query was:

TS = ((conservation OR reintroduc* OR repatriat* OR releas* OR supplementation OR

transloc*) AND ("captive?born" OR "captive?breed*" OR "captive?bred" OR "captive?rais*"

OR "captive?rear*" OR fisher* OR hatcher* OR "head$start*" OR relocat*) AND (native OR

resident OR wild))

We also screened studies reviewed in Stuparyk et al. for inclusion because our primary Web of

Science query did not encompass the terminology used in several ISR studies concerning

relocations intended to mitigate human-animal conflicts [22]. After screening (Supplementary Fig.

S1; Supplementary Table S1), the resulting 589 full-text papers—along with 28 additional papers

synthesized by Stuparyk et al.[22]—were assessed for our required inclusion criteria: At least one

direct, quantitative comparison of a fitness component (growth, survival, reproduction) or

fitness-correlated trait (movement, body constitution) between one translocated (either ESR or

ISR) and one wild-resident animal group. The comparison(s) must have been made in a natural

setting following the release of the translocated cohort into an area occupied by wild-resident

conspecifics of comparable age and size distributions. We excluded papers making comparisons

based on ambiguous traits without the author(s) clearly defining a relationship to fitness. These

ambiguous traits include, but are not limited to, stable isotope or itemized diet analysis [68], habitat

use [69], behavioral ethograms [70], or varying morphotypes [71]. Papers were also excluded if they did

not contain the necessary descriptive statistics to calculate effect sizes (see "effect size

extraction" below) and corresponding authors did not respond to email requests for

supplementary statistics. Finally, we excluded papers assessing the influence of translocations on

population genetic indices, as these datasets do not involve distinct animal groups and sample

sizes that are necessary to calculate effect sizes.

We quantified the disparity in performance between wild-resident and translocated

cohorts using two effect size metrics. In studies comparing animal groups based on mean

estimates of continuous traits (*e.g.,* growth rate, survivorship, clutch size), we calculated

standardized mean differences (SMD)[72]. For comparisons based on counts of dichotomous

outcomes (e.g., full vs. empty stomachs, proportion recaptured, successful vs. unsuccessful

nests), we calculated log odds ratios (LOR)[73]. Effect sizes were calculated using the function

escalc in the R package *metafor* [74]. From each paper, we extracted descriptive statistics necessary

to calculate effect sizes for all relevant fitness-related comparisons from the published and

supplemental texts and data tables, from additional materials requested from corresponding

authors, and from relevant figures and plots using the R package *digitize* [75].

We converted all SMD estimates to LORs for more intuitive interpretation of effect size

magnitudes [76]. Log odds ratios are centered at zero and were calculated such that positive values

indicate translocated individuals exhibited better relative performance than wild-resident

individuals, and negative values indicate translocated individuals under-performed relative to

wild individuals. In traits where lower values indicate better performance (e.g., rate of mortality,

parasite load), raw trait values were multiplied by negative one for consistent interpretation.

We extracted several study-level and estimate-level attributes to evaluate our predictions

in the final statistical model. Comparisons were assigned a study ID based on the paper from

which they were extracted and a group ID to distinguish unique groups of organisms within each

study. Identical group ID values were assigned to comparisons involving measures of different

traits in the same organisms [77]. We also retrieved taxonomic class, order, family, genus, and

species names of all assessed organisms from the Integrative Taxonomic Information System

database [78]. We identified the fitness-related trait under consideration in each comparison and

reclassified each into one of five broad categories: Movement, body constitution, growth,

survival, and reproduction. We also identified the predominant sex (i.e., female-only, male-only,

or mixed-sex) of organisms in each comparison. We characterized the amount of captive

ancestry of translocated individuals represented in each comparison using two separate variables.

First, we distinguished comparisons according to whether the translocation program involved a

captive phase or not (i.e., *in situ* reinforcement vs. *ex situ* reinforcement strategy). In those

studies that described the amount of captive ancestry among translocated cohorts, we further

classified captive ancestry using an ordered set of categories: 1) ISR organisms without any

captive ancestry, 2) ESR organisms with wild-origin parents not raised in captivity, and 3) ESR

organisms with one or more direct ancestral generation raised in captivity. Finally, we classified

the absence/presence (0/1) of enrichment techniques for select comparisons according to the

circumstances of each paper. Due to the arbitrary nature of enrichment treatments across our

dataset, we only assessed enrichment efficacy within studies that included post-release

performance comparisons with wild-resident conspecifics for both enriched and unenriched

translocated cohorts.

Statistical analysis

We first fit a meta-analytical taxonomic mixed model [53,79] to quantify the overall effect

size—relative post-release performance among translocated and wild-resident cohorts—across

the entire dataset, and also assess how individual-level (i.e., sex, taxonomic group) and study-

level (i.e., fitness surrogate, ESR vs. ISR strategy) factors influence this outcome (model 1;

Supplementary Table S2). We modeled the LOR responses as a function of categorical fixed

effects for sex, taxonomic class, fitness surrogate, and study strategy. We examined two-way

interactions between sex and fitness surrogate, and between sex and strategy to determine

whether these individual- and study-level factors had sex-specific relationships with relative

post-release performance. We also included a two-way interaction between fitness surrogate and

strategy to examine whether post-release performance in various fitness-related traits is

influenced by translocation strategy.

To assess the potential influence of domestication selection on the relative post-release

performance of translocated cohorts, we used a second, similar model to analyze a subset of

studies that specified the number of captive generations experienced by translocated cohorts

(model 2; Supplementary Table S3). We fit model 2 by substituting the fixed effect of strategy

from model 1 with captive ancestry, a categorical fixed effect with three levels classifying the

focal ACT cohort as either part of an ISR (without any captive ancestry), a head-starting program

(ESR with < 1 generation in captivity), or a captive-breeding program (ESR with > 1 direct

ancestral generations in captivity). We were unable to analyze captive ancestry as a continuous

variable due to a lack of reporting of exact ancestral generation times, and due to cohorts of

potentially mixed ancestry. Model 2 also included categorical fixed effects for sex, taxonomic

class, fitness surrogate, and captive ancestry, and two-way interactions between sex and fitness

surrogate, sex and captive ancestry, and between fitness surrogate and captive ancestry.

Finally, we developed model 3 to test the influence of enrichment strategies on relative

post-release performance among wild and translocated cohorts by analyzing a subset of studies

that gathered relative post-release performance data with a wild-control group from both

enriched and unenriched cohorts within the same experimental design (Supplementary Table S4).

In model 3, we included categorical fixed effects for sex, taxonomic class, fitness surrogate, ESR

vs. ISR strategy, and enrichment. We also included two-way interactions between enrichment

and sex, enrichment and surrogate, and enrichment and strategy.

In all models, we included several random effects terms to account for different sources

of non-independence among effect sizes. To account for shared taxonomy [53,79], we included all

Linnaean taxonomic ranks below the level of class as separate random effect terms in each

model. Taxonomic class was maintained as a fixed effect to quantify average differences in post-

release performance across broad, recognizable animal groups of potential relevance to

translocation managers. We chose the nested taxonomic random effects approach over the

incorporation of a phylogenetic covariance matrix because the taxonomic breadth of our dataset

precluded access to a comprehensive phylogeny with empirically-derived branch lengths [53,80] and

the goals of our analyses are not concerned with estimating the extent of correlated evolution.

We included study and group identities as two additional random effects terms in all models to

account for non-independence of effect sizes within studies and among repeated measures of

unique organisms [77]. We also included a term to model the variance in sampling errors for each

effect size (Supporting Information SI 1). To assess the overall dispersion of our effect sizes and

thus the generalizability of the findings of our models, we calculated total model heterogeneity

(Supplementary Table S5) [81]. Since measures of heterogeneity tend to be high in ecological

studies assessing diverse taxa [82], we further calculated the heterogeneity apportioned across the

aforementioned taxonomic and study-specific random effect terms to more clearly explain

patterns of dispersion in effect sizes [83].

Models were fit in a Bayesian framework using a Markov chain Monte Carlo (MCMC)

algorithm as implemented in the R package *MCMCglmm* (Supporting Information SI 1) [84]. We

address whether conservation programs produce organisms that perform similarly to or better

than wild-resident conspecifics after release by calculating the total probability that the log odds

ratio for a particular comparison is greater than or equal to zero. We also report the marginal

posterior mode and 95% highest posterior density credible interval (CrI) of the odds ratio (OR)

by exponentiating posterior distributions of log odds ratios. We used the R package *emmeans* to

report the marginal values of individual parameters and overall model effects to account for the

influence of other model parameters [85]. All derived parameters are calculated across the entire

posterior distribution to carry through all uncertainty.

We performed several meta-analysis validation tests to assess our dataset and findings for

publication bias and influential outliers (Supporting Information SI 2-4) [86]. As we found no

changes in our major findings after rerunning all models with identified outliers excluded

(Supplementary Tables S6-S8), and because we considered outlying estimates to be

representative of legitimate patterns in animal performance, we conducted the final analysis with

all outliers included [87].

Data Availability

All data and R code are available from Zenodo Digital Repository:

https://doi.org/10.5281/zenodo.7535746.

Literature Cited

1. Pimm, S. L. *et al.* The biodiversity of species and their rates of extinction, distribution,

and protection. *Science* **344**, (2014).

2. Barnosky, A. D. *et al.* Has the Earth's sixth mass extinction already arrived? *Nature* **471**,

51–57 (2011).

3. Ceballos, G. *et al.* Accelerated modern human–induced species losses: Entering the sixth

mass extinction. *Sci. Adv.* **1**, e1400253 (2015).

4. WWF/ZSL. The Living Planet Index database. (2020). Available at:

www.livingplanetindex.org.

5. Heppell, S. S., Crowder, L. B. & Crouse, D. T. Models to Evaluate Headstarting as a

Management Tool for Long-Lived Turtles Author. *Ecol. Appl.* **6**, 556–565 (1996).

6. Newmark, W. D., Jenkins, C. N., Pimm, S. L., McNeally, P. B. & Halley, J. M. Targeted

habitat restoration can reduce extinction rates in fragmented forests. *Proc. Natl. Acad. Sci.*

U. S. A. **114**, 9635–9640 (2017).

7. Ford, M. J. Selection in captivity during supportive breeding may reduce fitness in the

wild. *Conserv. Biol.* **16**, 815–825 (2002).

8. Seddon, P. J., Griffiths, C. J., Soorae, P. S. & Armstrong, D. P. Reversing defaunation:

Restoring species in a changing world. *Science* **345**, 406–411 (2014).

9. Ceballos, G., Ehrlich, P. R. & Raven, P. H. Vertebrates on the brink as indicators of

biological annihilation and the sixth mass extinction. *Proc. Natl. Acad. Sci. U. S. A.* **117**,

13596–13602 (2020).

10. IUCN/SCC. *Guidelines for Reintroductions and Other Conservation Translocations.*

Version 1.0. (2013).

11.   Seddon, P. J., Armstrong, D. P. & Maloney, R. F. Developing the science of

reintroduction biology. *Conserv. Biol.* **21**, 303–312 (2007).

12.   Hoffmann, M. *et al.* The impact of conservation on the status of the world's vertebrates.

Science **330**, 1503–1509 (2010).

13.   Jensen, E. L. *et al.* Population genomics through time provides insights into the

consequences of decline and rapid demographic recovery through head-starting in a

Galapagos giant tortoise. *Evol. Appl.* 1–11 (2018). doi:10.1111/eva.12682

14.   Fischer, J. & Lindenmayer, D. B. An assessment of the published results of animal

relocations. *Biol. Conserv.* **96**, 1–11 (2000).

15.   Pérez, I. *et al.* What is wrong with current translocations? A review and a decision-making

proposal. *Front. Ecol. Environ.* **10**, 494–501 (2012).

16.   Frankham, R. Genetic adaptation to captivity in species conservation programs. *Mol. Ecol.*

17, 325–333 (2008).

17.   Farquharson, K. A., Hogg, C. J. & Grueber, C. E. A meta-analysis of birth-origin effects

on reproduction in diverse captive environments. *Nat. Commun.* **9**, 1–10 (2018).

18.   Furlan, E. M. *et al.* Assessing the benefits and risks of translocations in depauperate

species: A theoretical framework with an empirical validation. *J. Appl. Ecol.* **57**, 831–841

(2020).

19.   Araki, H., Berejikian, B. A., Ford, M. J. & Blouin, M. S. Fitness of hatchery-reared

salmonids in the wild. *Evol. Appl.* **1**, 342–355 (2008).

20.   Ryman, N. & Laikre, L. Effects of supportive breeding on the genetically effective

population size. *Conserv. Biol.* **5**, 325–329 (1991).

21.   Araki, H., Cooper, B. & Blouin, M. S. Genetic effects of captive breeding cause a rapid,

531  cumulative fitness decline in the wild. *Science* **318**, 100–103 (2007).

532 22. Stuparyk, B., Horn, C. J., Karabatsos, S. & Arteaga, J. A meta-analysis of animal survival

533  following translocation: Comparisons between conflict and conservation efforts. *Can.*

534  *Wildl. Biol. Manag.* **7**, 3–17 (2018).

535 23. Tetzlaff, S. J., Sperry, J. H. & DeGregorio, B. A. Effects of antipredator training,

536  environmental enrichment, and soft release on wildlife translocations: A review and meta-

537  analysis. *Biol. Conserv.* **236**, 324–331 (2019).

538 24. Hendry, A. P., Schoen, D. J., Wolak, M. E. & Reid, J. M. The contemporary evolution of

539  fitness. *Annu. Rev. Ecol. Evol. Syst.* **49**, 457–476 (2018).

540 25. Kraaijeveld-Smit, F. J. L., Griffiths, R. A., Moore, R. D. & Beebee, T. J. C. Captive

541  breeding and the fitness of reintroduced species: A test of the responses to predators in a

542  threatened amphibian. *J. Appl. Ecol.* **43**, 360–365 (2006).

543 26. Degregorio, B. A., Sperry, J. H., Tuberville, T. D. & Weatherhead, P. J. Translocating

544  ratsnakes: Does enrichment offset negative effects of time in captivity? *Wildl. Res.* **44**,

545  438–448 (2017).

546 27. Horreo, J. L. *et al.* Nature versus nurture? Consequences of short captivity in early stages.

547  *Ecol. Evol.* **8**, 521–529 (2018).

548 28. McCleery, R. *et al.* Are declines of an endangered mammal predation-driven, and can a

549  captive-breeding and release program aid their recovery? *J. Zool.* **291**, 59–68 (2013).

550 29. Stringwell, R. *et al.* Maladaptation and phenotypic mismatch in hatchery-reared Atlantic

551  salmon *Salmo salar* released in the wild. *J. Fish Biol.* **85**, 1927–1945 (2014).

552 30. Heath, D. D., Heath, J. W., Bryden, C. A., Johnson, R. M. & Fox, C. W. Rapid Evolution

553  of Egg Size in Captive Salmon. *Science* **299**, 1738–1741 (2003).

554 31. Stamps, J. A. & Swaisgood, R. R. Someplace like home: Experience, habitat selection and

555   conservation biology. *Appl. Anim. Behav. Sci.* **102**, 392–409 (2007).

556 32. Evans, M. L., Wilke, N. F., O'Reilly, P. T. & Fleming, I. A. Transgenerational Effects of

557   Parental Rearing Environment Influence the Survivorship of Captive-Born Offspring in

558   the Wild. *Conserv. Lett.* **7**, 371–379 (2014).

559 33. Griffith, B., Scott, M. J., Carpenter, J. W. & Reed, C. Translocation as a Species

560   Conservation Tool: Status and Strategy. *Science* **245**, 477–480 (1989).

561 34. Wolf, C. M., Griffith, B., Reed, C. & Temple, S. A. Avian and Mammalian

562   Translocations: Update and Reanalysis of 1987 Survey Data. *Conserv. Biol.* **10**, 1142–

563   1154 (1996).

564 35. Rummel, L. *et al.* Use of wild-caught individuals as a key factor for success in vertebrate

565   translocations. *Anim. Biodivers. Conserv.* **39**, 207–219 (2016).

566 36. Ficetola, G. F. & De Bernardi, F. Supplementation or in situ conservation? Evidence of

567   local adaptation in the Italian agile frog Rana latastei and consequences for the

568   management of populations. *Anim. Conserv.* **8**, 33–40 (2005).

569 37. Lenormand, T. Gene flow and the limits to natural selection. *Trends Ecol. Evol.* **17**, 183–

570   189 (2002).

571 38. Dickens, M. J., Delehanty, D. J. & Michael Romero, L. Stress: An inevitable component

572   of animal translocation. *Biol. Conserv.* **143**, 1329–1341 (2010).

573 39. Armstrong, D. P. & Seddon, P. J. Directions in reintroduction biology. *Trends Ecol. Evol.*

574   **23**, 20–25 (2008).

575 40. Germano, J. M. & Bishop, P. J. Suitability of amphibians and reptiles for translocation.

576   *Conserv. Biol.* **23**, 7–15 (2009).

41. Goldenberg, S. Z. *et al.* Increasing conservation translocation success by building social

functionality in released populations. *Glob. Ecol. Conserv.* **18**, e00604 (2019).

42. Shier, D. M. & Owings, D. H. Effects of social learning on predator training and

postrelease survival in juvenile black-tailed prairie dogs, *Cynomys ludovicianus*. *Anim.*

Behav. **73**, 567–577 (2007).

43. Zhu, J., Gosnell, J. S., Akallal, L. & Goltsman, M. Fear changes traits and increases

survival: a meta-analysis evaluating the efficacy of antipredator training in captive-rearing

programs. *Restor. Ecol.* 1–11 (2022). doi:10.1111/rec.13674

44. Tuberville, T. D., Clark, E. E., Buhlmann, K. A. & Gibbons, J. W. Translocation as a

conservation tool: Site fidelity and movement of repatriated gopher tortoises (*Gopherus*

polyphemus). *Anim. Conserv.* **8**, 349–358 (2005).

45. DeGregorio, B. A., Weatherhead, P. J., Tuberville, T. D. & Sperry, J. H. Time in captivity

affects foraging behavior of ratsnakes: Implications for translocation. *Herpetol. Conserv.*

Biol. **8**, 581–590 (2013).

46. Mathews, F., Orros, M., McLaren, G., Gelling, M. & Foster, R. Keeping fit on the ark:

Assessing the suitability of captive-bred animals for release. *Biol. Conserv.* **121**, 569–577

(2005).

47. Mulder, K. P. *et al.* No paternal genetic integration in desert tortoises (*Gopherus agassizii*)

following translocation into an existing population. *Biol. Conserv.* **210**, 318–324 (2017).

48. Stewart, G. Meta-analysis in applied ecology. *Biol. Lett.* **6**, 78–81 (2010).

49. Skikne, S. A., Borker, A. L., Terrill, R. S. & Zavaleta, E. Predictors of past avian

translocation outcomes inform feasibility of future efforts under climate change. *Biol.*

Conserv. **247**, 108597 (2020).

50.   Zhang, Z., Gao, L. & Zhang, X. Environmental enrichment increases aquatic animal

601   welfare: A systematic review and meta-analysis. *Rev. Aquac.* **14**, 1120–1135 (2022).

602   51.   Resende, P. S., Viana-Junior, A. B., Young, R. J. & Azevedo, C. S. What is better for

603   animal conservation translocation programmes: Soft- or hard-release? A phylogenetic

604   meta-analytical approach. *J. Appl. Ecol.* **58**, 1122–1132 (2021).

605   52.   Dochtermann, N. A., Schwab, T., Anderson Berdal, M., Dalos, J. & Royauté, R. The

606   Heritability of Behavior: A Meta-analysis. *J. Hered.* **110**, 403–410 (2019).

607   53.   Hadfield, J. D. & Nakagawa, S. General quantitative genetic methods for comparative

608   biology: Phylogenies, taxonomies and multi-trait models for continuous and categorical

609   characters. *J. Evol. Biol.* **23**, 494–508 (2010).

610   54.   Lynch, M. Methods for the analysis of comparative data in evolutionary biology.

611   *Evolution* **45**, 1065–1080 (1991).

612   55.   Dodd, C. K. & Seigel, R. A. Relocation, repatriation, and translocation of amphibians and

613   reptiles: are they conservation strategies that work? *Biol. Conserv.* **62**, 230 (1992).

614   56.   Cayuela, H. *et al.* Survival cost to relocation does not reduce population self-sustainability

615   in an amphibian. *Ecol. Appl.* **29**, 1–15 (2019).

616   57.   Poo, S., Bogisich, A., Mack, M., Lynn, B. K. & Devan-Song, A. Post-release comparisons

617   of amphibian growth reveal challenges with sperm cryopreservation as a conservation

618   tool. *Conserv. Sci. Pract.* **4**, 1–9 (2022).

619   58.   Hunt, J. & Hodgson, D. What is fitness, and how do we measure it? in *Evolutionary*

620   *Behavioral Ecology* (eds. Westneat, D. & Fox, C.) (New York: Oxford University Press,

621   2010).

622   59.   Roff, D. A. *Life History Evolution*. (Sinauer Associates, Inc. Sunderland, MA, USA.,

623   2002).

624   60.   Bonnet, X., Naulleau, G. & Shine, R. The dangers of leaving home: dispersal and

625   mortality in snakes. *Biol. Conserv.* **89**, 39–50 (1999).

626   61.   Teixeira, C. P., de Azevedo, C. S., Mendl, M., Cipreste, C. F. & Young, R. J. Revisiting

627   translocation and reintroduction programmes: the importance of considering stress. *Anim.*

628   *Behav.* **73**, 1–13 (2007).

629   62.   Tarka, M., Guenther, A., Niemelä, P. T., Nakagawa, S. & Noble, D. W. A. Sex differences

630   in life history, behavior, and physiology along a slow-fast continuum: a meta-analysis.

631   *Behav. Ecol. Sociobiol.* **72**, (2018).

632   63.   Sutherland, W. J., Pullin, A. S., Dolman, P. M. & Knight, T. M. The need for evidence-

633   based conservation. *Trends Ecol. Evol.* **19**, 305–308 (2004).

634   64.   Moseby, K. E., Hill, B. M. & Lavery, T. H. Tailoring release protocols to individual

635   species and sites: One size does not fit all. *PLoS One* **9**, (2014).

636   65.   Berger-Tal, O., Blumstein, D. T. & Swaisgood, R. R. Conservation translocations: a

637   review of common difficulties and promising directions. *Anim. Conserv.* **23**, 121–131

638   (2020).

639   66.   Molony, S. E., Dowding, C. V., Baker, P. J., Cuthill, I. C. & Harris, S. The effect of

640   translocation and temporary captivity on wildlife rehabilitation success: An experimental

641   study using European hedgehogs (*Erinaceus europaeus*). *Biol. Conserv.* **130**, 530–537

642   (2006).

643   67.   Moher, D. *et al.* Preferred reporting items for systematic reviews and meta-analyses: The

644   PRISMA statement. *PLoS Med.* **6**, (2009).

68.   Quinn, T. P., Seamons, T. R. & Johnson, S. P. Stable isotopes of carbon and nitrogen

indicate differences in marine ecology between wild and hatchery-produced steelhead.

Trans. Am. Fish. Soc. **141**, 526–532 (2012).

69.   Himes, J. G., Hardy, L. M., Rudolph, D. C. & Burgdorp, S. J. Movement patterns and

habitat selection by native and repatriated Louisiana pine snakes (*Pituophis ruthveni*):

implications for conservation. *Herpetol. Nat. Hist.* **9**, 103–116 (2006).

70.   Rantanen, E. M., Buner, F., Riordan, P., Sotherton, N. & Macdonald, D. W. Vigilance,

time budgets and predation risk in reintroduced captive-bred grey partridges *Perdix*

perdix. *Appl. Anim. Behav. Sci.* **127**, 43–50 (2010).

71.   Larsen, D. A. *et al.* Early Life History Variation in Hatchery- and Natural-Origin Spring

Chinook Salmon in the Yakima River, Washington. *Trans. Am. Fish. Soc.* **142**, 540–555

(2013).

72.   Rosenthal, R. & Rubin, D. B. Comparing effect sizes of independent studies. *Psychol.*

Bull. **92**, 500–504 (1982).

73.   Haddock, C. K., Rindskopf, D. & Shadish, W. R. Using Odds Ratios as Effect Sizes for

Meta-Analysis of Dichotomous Data: A Primer on Methods and Issues. *Psychol. Methods*

3, 339–353 (1998).

74.   Viechtbauer, W. Conducting meta-analyses in R with the metafor. *J. Stat. Softw.* **36**, 1–48

(2010).

75.   Poisot, T. The digitize package: extracting numerical data from scatterplots. *R J.* **3**, 26–27

(2011).

76.   Rosenberg, M. S., Rothstein, H. R. & Gurevitch, J. Effect sizes: Conventional Choices and

Calculations. in *Handbook of meta-analysis in ecology and evolution* (eds. Kuricheva, J.,

Gurevitch, J. & Mengersen, K.) 61–71 (Princeton University Press, 2013).

77.   Noble, D. W. A., Lagisz, M., O'dea, R. E. & Nakagawa, S. Nonindependence and

670   sensitivity analyses in ecological and evolutionary meta-analyses. *Mol. Ecol.* **26**, 2410–

671   2425 (2017).

672   78.   ITIS. Retrieved 27 Sep 2021 from the Integrated Taxonomic Information System (ITIS).

673   *Integrated Taxonomic Information System (ITIS)* (2021). doi:10.5066/F7KH0KBK

674   79.   Clutton-Brock, T. H. & Harvey, P. H. Primate ecology and social organization. *J. Zool.*

675   **183**, 1–39 (1977).

676   80.   Lajeunesse, M. J., Rosenberg, M. S. & Jennions, M. Phylogenetic nonindependence and

677   meta-analysis. in *Handbook of meta-analysis in ecology and evolution* (eds. Kuricheva, J.,

678   Gurevitch, J. & Mengerson, K.) 284–299 (Princeton University Press, 2013).

679   81.   Higgins, J. P. T. & Thompson, S. G. Quantifying heterogeneity in a meta-analysis. *Stat.*

680   *Med.* **21**, 1539–1558 (2002).

681   82.   Senior, A. M. *et al.* Heterogeneity in ecological and evolutionary meta- analyses: its

682   magnitude and implications. *Ecology* **97**, 3293–3299 (2016).

683   83.   Nakagawa, S. & Santos, E. S. A. Methodological issues and advances in biological meta-

684   analysis. *Evol. Ecol.* **26**, 1253–1274 (2012).

685   84.   Hadfield, J. D. MCMCglmm: MCMC Methods for Multi-Response GLMMs in R. *J. Stat.*

686   *Softw.* **33**, 1–22 (2010).

687   85.   Lenth, R. V. emmeans: Estimated Marginal Means, aka Least-Square Means. R package

688   version 1.6.3. https://CRAN.R-project.or (2021).

689   86.   Koricheva, J., Jennions, M. D. & Lau, J. Temporal trends in effect sizes: Causes,

690   Detection, and Implications. in *Handbook of meta-analysis in ecology and evolution* (eds.

691   Koricheva, J., Gurevitch, J. & Mengersen, K.) 237–254 (Princeton University Press,

692   2013).

693   87.   Aguinis, H., Gottfredson, R. K. & Joo, H. Best-Practice Recommendations for Defining,

694   Identifying, and Handling Outliers. *Organ. Res. Methods* **16**, 270–301 (2013).

Acknowledgments

We would like to thank members of the D.A. Warner and M.E. Wolak labs, and students from

the meta-analysis course developed and taught by A.E.W. for providing comments on earlier

drafts of the manuscript.

Author contributions

Conceptualization: IPG and MEW; **Data curation:** IPG and MEW; **Formal analysis:** IPG,

with input from AEW and MEW; **Investigation:** IPG; **Project administration:** MEW;

Visualization: IPG and MEW; **Writing - original manuscript:** IPG; **Writing - review and**

editing: AEW and MEW.

Competing Interests

The authors declare no competing interests.

Figure Legends

Figure 1. Marginal posterior distributions of log odds ratios (ln odds ratio) plotted across
taxonomic classes. Values greater than zero (vertical dotted line) indicate that translocated
organisms out-performed wild-resident conspecifics, and negative values indicate translocated
organisms under-performed. Vertical lines within each density plot indicate the 95% credible
intervals (CrI). The parenthetical statements indicate the number of studies and the number of
individual comparisons comprising each density plot, respectively.

Figure 2. Marginal posterior distributions of log odds ratios (ln odds ratio) plotted across
taxonomic classes. Values greater than zero (vertical dotted line) indicate that translocated
organisms out-performed wild-resident conspecifics, and negative values indicate translocated
organisms under-performed. Blue density plots represent comparisons in which translocated
organisms received some form of pre-release enrichment, whereas tan density plots represent
comparisons where translocated organisms were unenriched. Vertical lines within each density

725 plot indicate the 95% credible intervals (CrI). The parenthetical statements indicate the number
726 of studies and the total number of comparisons between enriched/unenriched and wild cohorts
727 organized by taxonomic class.
728