Title: The fitness consequences of wildlife conservation translocations: a meta-analysis

Authors: Iwo P. Gross[1, *], Alan E. Wilson[2], & Matthew E. Wolak[1]

[1]Department of Biological Sciences, Auburn University, Auburn, AL 36849, USA

[2]School of Fisheries, Aquaculture, and Aquatic Sciences, Auburn University, Auburn, AL

36849, USA

[*]Corresponding author: ipg0004@auburn.edu

Running title: Conservation translocation efficacy meta-analysis

**ABSTRACT**

Conservation translocation is a common strategy to offset mounting rates of population declines

through the transfer of captive- or wild-origin organisms into areas where conspecific

populations are imperiled or completely extirpated. Translocations that supplement existing

populations are referred to as reinforcements, and can be conducted using captive-origin animals

(*ex situ* reinforcements [ESR]) or wild-origin animals without any captive ancestry (*in situ*

reinforcement [ISR]). These programs have been criticized for low success rates and husbandry

practices that produce individuals with genetic and performance deficits, but the post-release

performance of captive-origin or wild-origin translocated groups has not been systematically

reviewed to quantify success relative to wild-resident control groups. To assess the disparity in

post-release performance of translocated organisms relative to wild-resident conspecifics and

examine the association of performance disparity with organismal and methodological factors

across studies, we conducted a systematic review and meta-analysis of 821 performance

comparisons from 171 studies representing nine animal classes (101 species). We found that

translocated organisms have 64% decreased odds of out-performing their wild-resident

counterparts, supporting claims of systemic issues hampering conservation translocations. To

help identify translocation practices that could maximize program success in the future, we

further quantified the impact of broad organismal and methodological factors on the disparity

between translocated and wild-resident conspecific performance. Pre-release animal enrichment

significantly reduced performance disparities, whereas our results suggest no overall effects of

taxonomic group, sex, captive generation time, or the type of fitness surrogate measured. This

work is the most comprehensive systematic review to date of animal conservation translocations

in which wild conspecifics were used as comparators, thereby facilitating an evaluation of the

overall impact of this conservation strategy and identifying specific actions to increase success.

Our review highlights the need for conservation managers to include both sympatric and

allopatric wild-reference groups to ensure the post-release performance of translocated animals

can be evaluated. Further, our analyses identify pre-release animal enrichment as a particular

strategy for improving the outcomes of animal conservation translocations, and demonstrate how

meta-analysis can be used to identify implementation choices that maximize translocated animal

contributions to recipient population growth and viability.

Keywords: *Captive-breeding, captive-rearing, head-starting, animal relocation, animal*

*reintroduction, ex situ reinforcement, in situ reinforcement*

## CONTENTS

## I. INTRODUCTION

Rates of floral and faunal species loss in the new Anthropocene signal the onset of a sixth global mass extinction (Barnosky *et al.*, 2011; Pimm *et al.*, 2014). Among vertebrates, extinction rates in the last century increased 100 times above pre-industrial levels (Ceballos *et al.*, 2015) and recorded population size declines have averaged 68% in the last 50 years (WWF/ZSL, 2020). Continued species loss will have irreversible consequences for ecosystem services and resiliency. Slowing the rate of species loss will require eliminating existing threats (e.g., disease, poaching;Heppell, Crowder & Crouse, 1996; Snyder *et al.*, 1996) and extensive habitat preservation and restoration (Bouzat *et al.*, 2009; Newmark *et al.*, 2017). While environmental threats are being addressed, supportive breeding and translocation programs have been proposed to maintain species and their roles in the ecosystem (Ford, 2002; Seddon *et al.*, 2014; Hufbauer *et al.*, 2015).

Animal conservation translocations (ACTs) are a common strategy intended to directly offset species extinction or provide beneficial ecosystem functions through the supplementation of threatened populations and the repatriation of extirpated populations (Ceballos, Ehrlich & Dirzo, 2017; Ceballos, Ehrlich & Raven, 2020). ACT programs supplementing extant populations in their indigenous range are called reinforcements, and can be grouped into two categories (Seddon, Armstrong & Maloney, 2007; IUCN/SCC, 2013). First, *in situ* reinforcements (ISRs) collect and transport wild organisms to another natural area for the benefit of the recipient conspecific population. Second, *ex situ* reinforcements (ESRs) incorporate a captive phase lasting from less than one generation (i.e., head-starting) up to several generations (i.e., captive breeding) before release. Despite immense efforts across a wide array of taxa to employ ACTs and document their efficacy in the literature, a major unresolved question is

whether translocated individuals are as capable as their wild conspecifics at contributing to

population viability and preventing extinctions.

To date, ACTs have had many celebrated results and a quantifiable impact on global

extinction rates (Hoffmann *et al.*, 2010; Jachowski *et al.*, 2011; Hess *et al.*, 2012; Jensen *et al.*,

2018; Bolam *et al.*, 2021). Nevertheless, ACTs are often considered a halfway technology with

logistical and systemic shortcomings that result in failure to achieve pre-defined conservation

goals (Snyder *et al.*, 1996; Fischer & Lindenmayer, 2000; Pérez *et al.*, 2012; Germano *et al.*,

2015; Sullivan, Nowak & Kwiatkowski, 2015). In particular, multiple studies indicate that ACTs

might produce individuals with inherent genetic and performance-related deficits that

compromise ACT program goals and reduce recipient population viability (Frankham, 2008;

Satake & Araki, 2012; Farquharson, Hogg & Grueber, 2018, 2021). These deficits are most

severe in ESRs involving multiple captive generations since the capture of ESR founders acts as

a bottleneck on genetic diversity. Even with a stringent captive breeding structure, factors

inherent to husbandry practices (e.g., small effective population size, inbreeding depression,

relaxed purifying selection) increase the likelihood of fixation of undesirable alleles and a loss of

allelic heterozygosity (Lynch & O'Hely, 2001; Ford, 2002; Araki *et al.*, 2008; Furlan *et al.*,

2020). A large influx of captive-born individuals into the wild with such genetic properties can

swamp genetic diversity of locally adapted populations and reduce effective population sizes

despite an overall increase in numbers (Ryman & Laikre, 1991; Alleaume-Benharira, Pen &

Ronce, 2006; Laikre *et al.*, 2010; Ford, Murdoch & Howard, 2012; Waples *et al.*, 2016; Pinter,

Epifanio & Unfer, 2019). For example, Klütsch *et al.,* (2019) examined how stocking of captive-

bred brown trout (*Salmo trutta*) influenced the genetic diversity of wild sub-populations isolated

by hydroelectric dams. They found that historically stocked sections of the system exhibited

genetic signs of bottleneck events indicative of genetic swamping by the released captive-bred

fish. However, such genetic deficits require multiple generations of captive breeding to

accumulate which suggests alternative factors contribute to fitness reductions observed in ESRs

lasting a generation or less (Araki, Cooper & Blouin, 2007; Farquharson *et al.*, 2018).

A fundamental criticism of ESRs is that non-natural conditions present in captivity can

influence individual phenotypes, diminishing post-release performance relative to wild

conspecifics (Araki *et al.*, 2008; Stuparyk *et al.*, 2018; Tetzlaff, Sperry & DeGregorio, 2019a;

Crates, Stojanovic & Heinsohn, 2022), where animal performance is defined as a measure of

some critical life history character (e.g., growth rate, survival, fecundity) that correlates with

individual fitness (Thoday, 1953; Hendry *et al.*, 2018). In a variety of taxa, ESRs lasting a

generation or less negatively influence post-release performance metrics such as anti-predator

behavior (Kraaijeveld-Smit *et al.*, 2006; Melstrom, Salau & Shanafelt, 2019), dispersal (Lehrer

*et al.*, 2016; Degregorio *et al.*, 2017; McCallen *et al.*, 2018), growth (Horreo *et al.*, 2018),

survival (McCleery *et al.*, 2013; Vanderwerf *et al.*, 2014), and reproductive success (Araki *et al.*,

2007; Knudsen *et al.*, 2008; Ford *et al.*, 2016). Such phenotypic mismatches can arise *via*

domestication selection, where trait values that are selected for in captivity are maladaptive when

expressed in a natural environment (Heath *et al.*, 2003; Stringwell *et al.*, 2014).

The unintentional selection for domesticated phenotypes in artificial environments is

likely a major contributor to reduced performance of ESRs, since it can alter phenotype

distributions within a single generation (Araki *et al.*, 2007; Horreo *et al.*, 2018). For example,

Araki *et al.,* (2007) discovered that captive-bred steelhead trout (*Oncorhynchus mykiss*)

exhibited a 40% reduction in fecundity relative to their wild counterparts for every additional

generation of captive ancestry. In the absence of domestication selection, phenotypic plasticity

during less than one generation of captive development and rearing can lead to similar deviations

from the wild-type in anatomical, physiological, and behavioral traits (Huntingford, 2004;

Stamps & Swaisgood, 2007; Stringwell *et al.*, 2014). Moreover, certain plastic phenotypic

deviations have an epigenetic basis, thereby potentially influencing performance of future

generations regardless of their developmental environment (Day & Bonduriansky, 2011; Evans

*et al.*, 2014).

By virtue of the use of wild-born organisms and lack of a captive component, ISRs are

lauded as a less intrusive and a generally more successful translocation strategy than ESRs

(Griffith *et al.*, 1989; Wolf *et al.*, 1996; Fischer & Lindenmayer, 2000; Kingsbury & Attum,

2009; Rummel *et al.*, 2016). However, failure rates for ISRs are still notably high and potentially

a consequence of similar issues that affect ESRs. At the population-level, supplementing existing

populations with conspecifics from other wild populations can lead to maladaptive gene flow or

outbreeding depression of the recipient population and a reduction in meta-population genetic

diversity (Lenormand, 2002; Moritz, 2002; Ficetola & De Bernardi, 2005). At the individual-

level, ISRs can impact post-release performance either directly through transportation-related

stress (Dickens, Delehanty & Michael Romero, 2010) or indirectly through energetic costs of

establishment (Armstrong & Seddon, 2008), errant dispersal and homing behaviors (Germano &

Bishop, 2009), diminished social structure (Goldenberg *et al.*, 2019), and phenotypic mismatches

within the novel ecosystem (Stamps & Swaisgood, 2007; Turlure *et al.*, 2013).

Exposing captive organisms to some form of enrichment is the most intuitive strategy to

counteract phenotypic mismatch, and a reasonable compromise when ISRs are not possible. We

use the term enrichment broadly to refer to a variety of supportive measures (Fischer &

Lindenmayer, 2000) employed by ACTs to offset any maladaptive behavior, physiology, or

physical injury incurred during captivity and/or translocation (Reading, Miller & Shepherdson,

2013; Goldenberg *et al.*, 2019; Tetzlaff *et al.*, 2019a). Some examples include fostering pre-

release organisms with experienced conspecifics (Lumsden & Drever, 2002; Shier & Owings,

2007), simulating *in situ* environments with naturalistic enclosures or antipredator training

(Brown, Davidson & Laland, 2003; Nazar & Marin, 2011; Roe, Frank & Kingsbury, 2015;

Tetzlaff, Sperry & DeGregorio, 2018; Zhu *et al.*, 2022), implementing a soft-release phase where

supplemental feeding and/or shelter are provided (Tuberville *et al.*, 2005; Brown *et al.*, 2006), or

simply reducing the length of the captive phase (Degregorio et al., 2017; DeGregorio,

Weatherhead, Tuberville, & Sperry, 2013; but see Tetzlaff, Sperry, Kingsbury, & DeGregorio,

2019). Literature reviews suggest that enriched ESR cohorts are more successful post-release

compared to un-enriched cohorts (Fischer & Lindenmayer, 2000; Tetzlaff *et al.*, 2019a), but the

lack of a wild-reference group in the synthesized findings makes it difficult to conclude whether

enrichment strategies produce translocated organisms that are indistinguishable from wild

conspecifics with respect to post-release fitness (Mathews *et al.*, 2005).

Finally, the metric used to gauge relative performance can vary in its representation of

fitness, thereby shaping study conclusions (Moyes *et al.*, 2009; Wilson & Nussey, 2010; Hendry

*et al.*, 2018). The lifetime reproductive success of an individual is the ideal measure of fitness

when focusing on population dynamics and phenotypic adaptation, but lifetime reproductive

success and the component life history traits that determine it (e.g., survival and fecundity) are

difficult to measure in natural conditions (Coulson *et al.,* 2006; Fairbairn & Reeve, 2001; Hendry

*et al*., 2018; Kozłowski, 1993). Consequently, fitness is often approximated using traits that

typically correlate with fitness (e.g., body size, body constitution, movement). However, the

extent to which correlates and components of fitness (hereafter collectively referred to as fitness

surrogates) covary and ultimately relate to lifetime reproductive success remains uncertain

(Moyes *et al.*, 2009; Hendry *et al.*, 2018). For example, Mulder *et al.,* (2017) found that while

translocated and wild-resident desert tortoises (*Gopherus agassizii*) had similar body condition

and survivorship four years post-release, translocated males fathered none of 92 offspring for

which paternity was determined. This suggests that survival is a potentially inappropriate

component of fitness to gauge ACT success in adult organisms with high adult survivorship.

Conservation translocations should consider multiple relevant fitness components during project

assessment to identify potential life-history trade-offs and avoid misleading conclusions (Fincke

& Hadrys, 2001; Wilson & Nussey, 2010; Pekkala, Kotiaho & Puurtinen, 2011; Hendry *et al.*,

2018).

The translocation literature has grown to a point that quantitative reviews are necessary to

summarize the high volume of papers, while also accounting for heterogeneity and bias (Bajomi

*et al.*, 2010; Stewart, 2010). Seminal reviews have evaluated ACT efficacy from the published

literature and *via* mail-in questionnaires sent to conservation program managers (e.g., Griffith *et*

*al.,* 1989; Wolf *et al.,* 1996b; Fischer & Lindenmayer, 2000; Kraaijeveld-Smit *et al.,* 2006;

Seddon *et al.*, 2007; Rummel *et al.,* 2016; Brichieri-Colombi *et al.,* 2019). In these approaches,

the response variable (translocation success *versus* failure) reduces a highly nuanced outcome to

a binary classification based in part on unstandardized criteria (e.g., creation of a self-sustaining

population) and professional opinion. These syntheses are an invaluable resource for applied

conservation and management (Nason *et al.*, 2021), but their largely narrative format is not

conducive to quantitative synthesis across broad and multifactorial datasets while accounting for

bias and heterogeneity among studies.

Meta-analysis is a quantitative tool that integrates the findings of published papers to

examine broader patterns and draw more powerful conclusions about a focal topic (Glass, 1976;

Stewart, 2010; Gurevitch *et al.*, 2018). Meta-analyses standardize and pool effect sizes from

multiple studies while preserving the magnitude and variance of the original effects, thus

enabling predictions about particular driving variables to be modeled as covariates that account

for unexplained variation among studies (Glass, 1976; Bennett *et al.*, 2017). A properly executed

meta-analysis of the conservation literature will synthesize existing trends, assess knowledge

gaps, and develop recommendations for future research and management actions.

Meta-analysis is well-equipped to gauge translocation efficacy for establishing viable

populations based on the post-release performance of animal subjects (Ducatez & Shine, 2019;

Jule, Leaver, & Lea, 2008; Linklater *et al.,* 2011; Skikne *et al.,* 2020; Tetzlaff, Sperry, &

DeGregorio, 2019; Zhang, Gao, & Zhang, 2022; Zhu *et al.,* 2022; also see Resende *et al.,* 2021).

Previous meta-analyses fail to include a wild-reference group (but see Stuparyk *et al.,* 2018) and

therefore cannot address the way by which translocated individuals may contribute to the

population viability goals of ACTs. For instance, translocated animals that are on average less

likely to survive, reproduce, or perform relative to wild conspecifics in the recipient population

(i.e., a sympatric control group comparison) will have limited contributions to genetic or

evolutionary mechanisms increasing population persistence (i.e., genetic and evolutionary

rescue; Carlson *et al.,* 2014). The performance of wild members of the recipient population

should be considered by conservation managers as an adaptive baseline shaped by local

environmental and social pressures that translocation programs should strive to emulate

(Mathews *et al.*, 2005). Even though translocated cohorts might exhibit lower survival or

reproductive success compared to wild control groups, translocations may positively affect other

demographic and ecological processes (e.g., demographic rescue and Allee effects) that fulfill

primary goals of ACTs (IUCN/SSC, 2013). Distinguishing these beneficial effects on the wild

resident recipient population requires comparison with wild allopatric control groups to separate

the effect of translocation from the measurement of performance in both the translocated cohort

as well as the wild sympatric recipient group. However, use of both wild allopatric and sympatric

control groups has rarely been reported (e.g., Molony *et al.,* 2006; Sah *et al.,* 2016).

Here, we present the first comprehensive meta-analysis to assess conservation

translocation programs through the synthesis of studies involving direct, post-release

comparisons of translocated and wild-resident conspecifics. We further expand on previous

reviews by sampling the literature broadly across macroscopic animal groups and assessing

relative post-release performance using multiple fitness surrogates. Our primary objective is to

quantify the disparity in post-release performance of translocated organisms relative to wild-

resident conspecifics across animal taxa and ultimately gauge the potential contribution of

translocated individuals to genetic or evolutionary rescue strategies. We performed a systematic

review and meta-analysis to quantify this performance disparity and also sought to determine

whether organismal and methodological factors influence this disparity by extracting information

about (*i*) sex and (*ii*) taxonomy as well as (*iii*) what strategies were implemented, if at all,

regarding pre-release animal enrichment and categorizing (*iv*) translocation reinforcements as *in*

*situ* vs. *ex situ*, (*v*) captive generation time, and (*vi*) the type of fitness surrogate measured. Our

systematic review and meta-analysis were guided by how we predict the above factors will

influence the relative performance among translocated and wild cohorts:

(1) Animals incorporated into conservation translocation programs represent a wide

taxonomic breadth in which natural history, sensory capabilities, and native environments

will vary substantially. These factors will also affect the capacities of conservation

managers to recreate natural conditions in captivity and in turn influence organismal

response. We predict that taxonomic groups will broadly vary in their relative post-

release performance.

(2) Disparity in performance among translocated and wild-resident organisms will increase

as a function of whether a trait is considered an indirect correlate (e.g., movement and

body constitution) or a direct component of an organism's fitness (e.g., growth, survival,

reproductive success).

(3) Within species, sexes can vary in physical form and physiology, seasonal and annual

energetic investments, optimal life-history strategy, spatial ecology, and behavior

(Trivers, 1972; Parker, 1979; Ruckstuhl & Neuhaus, 2005; Bondurianski & Chenoweth,

2009; Fairbairn, 2013). We predict significant intraspecific variation in post-release

relative performance among male and female cohorts.

(4) We predict varying amounts of captive ancestry will explain variation in relative post-

release performance of translocated cohorts. We predict that ESRs lasting > 1 generation

will sustain greater disparity in relative performance than ESRs lasting < 1 generation

(e.g., head-starting programs). We also predict that ISRs involving zero captive

generations will have more similar post-release performances to wild-reference groups

compared to ESRs.

(5) Translocated organisms exposed to one or more forms of enrichment will perform better

post-release relative to wild conspecifics compared to unenriched translocated

conspecifics.

## II. METHODS

### (1) Literature search and inclusion criteria

We conducted our literature review, screening, and analysis according to Preferred Reporting

Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (Fig. 1; Moher *et al.*,

2009) and following a PRISMA EcoEvo checklist (Appendix S1; O'Dea *et al.*, 2021). We

searched the literature using Web of Science, including all databases (01/01/1900-08/23/2019),

for peer-reviewed studies concerning animal conservation translocations with some reference to

a wild-resident group used as a control. Our query was:

TS = ((conservation OR reintroduc* OR repatriat* OR releas* OR supplementation OR

transloc*) AND ("captive born" OR "captive breed*" OR "captive bred" OR "captive rais*" OR

"captive rear*" OR fisher* OR hatcher* OR "headstart*" OR "head start*" OR relocat*) AND

(native OR resident OR wild) AND (animal OR invertebrate OR fish* OR amphib* OR reptil*

OR mammal* OR bird* OR avian*))

We also screened studies reviewed in Stuparyk *et al.,* (2018) for inclusion because our primary

Web of Science query did not encompass the terminology used in several ISR studies concerning

ISRs intended to mitigate human-animal conflicts. After screening (Fig. 1; Table S1 in Appendix

S2), the resulting 589 full-text papers—along with 28 additional papers synthesized by Stuparyk

*et al.,* (2018)—were assessed for our required inclusion criteria: At least one direct, quantitative

comparison of a fitness component (growth, survival, reproduction) or fitness-correlated trait

(movement, body constitution) between one translocated (either ESR or ISR) and one wild-

resident animal group. The comparison(s) must have been made in a natural setting following the

release of the translocated cohort into an area occupied by wild-resident conspecifics of

comparable age and size distributions. We excluded papers making comparisons based on

ambiguous traits without the author(s) clearly defining a relationship to fitness. These ambiguous

traits include, but are not limited to, stable isotope or itemized diet analysis (e.g., Bourass &

Hingrat, 2015; Quinn, Seamons, & Johnson, 2012), habitat use (e.g., Himes, Hardy, Rudolph, &

Burgdorp, 2006; Mann, Marty Holtgren, Auer, & Ogren, 2011), behavioral ethograms (e.g.,

Rantanen, Buner, Riordan, Sotherton, & Macdonald, 2010), or varying morphotypes (Larsen *et*

*al.*, 2013). Papers were also excluded if they did not contain the necessary descriptive statistics to

calculate effect sizes (see "effect size extraction" below) and corresponding authors did not

respond to email requests for supplementary statistics. Finally, we excluded papers assessing the

influence of translocations on population genetic indices, as these datasets do not involve distinct

animal groups and sample sizes that are necessary to calculate effect sizes.

**(2) Effect size extraction and calculation**

An effect size is a statistic used to estimate the magnitude of a relationship between two

experimental groups or variables (Nakagawa & Cuthill, 2007). We quantified the disparity in

performance between wild-resident and translocated cohorts using two effect size metrics. In

studies comparing animal groups based on mean estimates of continuous traits (*e.g.,* growth rate,

survivorship, clutch size), we calculated standardized mean differences (SMD; Rosenthal &

Rubin, 1982). For comparisons based on counts of dichotomous outcomes (e.g., full vs. empty

stomachs, proportion recaptured, successful vs. unsuccessful nests), we calculated log odds ratios

(LOR; Haddock, Rindskopf, & Shadish, 1998). Effect sizes were calculated using the function

*escalc* in the R package *metafor* (Viechtbauer, 2010). From each paper, we extracted descriptive

statistics necessary to calculate effect sizes for all relevant fitness-related comparisons from the

published and supplemental texts and data tables, from additional materials requested from

corresponding authors, and from relevant figures and plots using the R package *digitize* (Poisot,

2011).

We converted all SMD estimates to LORs for more intuitive interpretation of effect size

magnitudes (Chinn, 2000; Rosenberg, Rothstein & Gurevitch, 2013). Log odds ratios are

centered at zero and were calculated such that positive values indicate translocated individuals

exhibited better relative performance than wild-resident individuals, and negative values indicate

translocated individuals under-performed relative to wild individuals. In traits where lower

values indicate better performance (e.g., rate of mortality, parasite load), raw trait values were

multiplied by negative one for consistent interpretation.

**(3) Moderators**

We extracted several study-level and estimate-level attributes to evaluate our predictions in the

final statistical model. Comparisons were assigned a study ID based on the paper from which

they were extracted and a group ID to distinguish unique groups of organisms within each study.

Identical group ID values were assigned to comparisons involving measures of different traits in

the same organisms (Noble *et al.*, 2017). We also retrieved taxonomic class, order, family,

genus, and species names of all assessed organisms from the Integrative Taxonomic Information

System database (ITIS, 2021). We identified the fitness-related trait under consideration in each

comparison and reclassified each into one of five broad categories: Movement, body

constitution, growth, survival, and reproduction. We also identified the predominant sex (i.e.,

female-only, male-only, or mixed-sex) of organisms in each comparison. We characterized the

amount of captive ancestry of translocated individuals represented in each comparison using two

separate variables. First, we distinguished comparisons according to whether the translocation

program involved a captive phase or not (i.e., *in situ* reinforcement vs. *ex situ* reinforcement

strategy). In those studies that described the amount of captive ancestry among translocated

cohorts, we further classified captive ancestry using an ordered set of categories: 1) ISR

organisms without any captive ancestry, 2) ESR organisms with wild-origin parents not raised in

captivity, and 3) ESR organisms with one or more direct ancestral generation raised in captivity.

Finally, we classified the absence/presence (0/1) of enrichment techniques for select comparisons

according to the circumstances of each paper. Due to the arbitrary nature of enrichment

treatments across our dataset, we only assessed enrichment efficacy within studies that included

post-release performance comparisons with wild-resident conspecifics for both enriched and

unenriched translocated cohorts.

**(4) Statistical analysis**

*(a) Model structure*

We first fit a meta-analytical taxonomic mixed model (Clutton-Brock & Harvey, 1977; Hadfield

& Nakagawa, 2010) to quantify the overall effect size—relative post-release performance among

translocated and wild-resident cohorts—across the entire dataset, and also assess how individual-

level (i.e., sex, taxonomic group) and study-level (i.e., fitness surrogate, ESR vs. ISR strategy)

factors influence this outcome (model 1; Table S2 in Appendix S2). We modeled the LOR

responses as a function of categorical fixed effects for sex, taxonomic class, fitness surrogate,

and study strategy. We examined two-way interactions between sex and fitness surrogate, and

between sex and strategy to determine whether these individual- and study-level factors had sex-

specific relationships with relative post-release performance. We also included a two-way

interaction between fitness surrogate and strategy to examine whether post-release performance

in various fitness-related traits is influenced by translocation strategy.

*(b) Quantification of study heterogeneity*

We included several random effects terms to account for different sources of non-independence

among effect sizes. To account for shared taxonomy (Clutton-Brock & Harvey, 1977; Hadfield

& Nakagawa, 2010), we included all Linnaean taxonomic ranks below the level of class as

separate random effect terms in each model. Taxonomic class was maintained as a fixed effect to

quantify average differences in post-release performance across broad, recognizable animal

groups of potential relevance to translocation managers. We chose the nested taxonomic random

effects approach over the incorporation of a phylogenetic covariance matrix because the

taxonomic breadth of our dataset precluded access to a comprehensive phylogeny with

empirically-derived branch lengths (Hadfield & Nakagawa, 2010; Lajeunesse, Rosenberg &

Jennions, 2013) and the goals of our analysis are not concerned with estimating the extent of

correlated evolution. We included study and group identities as two additional random effects

terms to account for non-independence of effect sizes within studies and among repeated

measures of unique organisms (Noble *et al.*, 2017). We also included a term to model the

variance in sampling errors for each effect size, which incorporates study-level precision into the

model (see "*Meta-analysis model specifications*" below; Hadfield, 2010; Mengersen *et al*.,

2013). To assess the overall dispersion of our effect sizes and thus the generalizability of the

findings of our models, we calculated total model heterogeneity (Higgins & Thompson, 2002;

Table S5 in Appendix S2). Since measures of heterogeneity tend to be high in ecological studies

assessing diverse taxa (Senior *et al.*, 2016), we further calculated the heterogeneity apportioned

across the aforementioned taxonomic and study-specific random effect terms to more clearly

explain patterns of dispersion in effect sizes (Nakagawa & Santos, 2012).

*(c) Meta-analysis model specifications*

The model was fit in a Bayesian framework using a Markov chain Monte Carlo (MCMC)

algorithm as implemented in the R package *MCMCglmm* (Hadfield, 2010). We specified diffuse

normal prior distributions for all fixed effects (mean=0, variance=$10^{10}$) and parameter expanded

priors on all variance components to yield scaled non-central F-distributions with numerator and

denominator degrees of freedom equal to one and a scale parameter of 1000 (Gelman, 2006;

Hadfield, 2010). The model was constructed as a random-effect meta-analysis by including a

term that contained the estimated sampling variance for each effect size and fixing the variance

associated with this term to one (in the prior). The MCMC chain for each model was run for an

initial burn-in of 3000 iterations with samples from the chain being saved every $150^{th}$ iteration

until 3000 samples were obtained. All models were checked for autocorrelation values $< |0.1|$,

and model convergence was confirmed using a Heidelberg stationarity test and through visual

assessment of posterior distributions.

*(d) Influence of captive ancestry and enrichment*

To assess the potential influence of domestication selection on the relative post-release

performance of translocated cohorts, we used a second, similar model to analyze a subset of

studies from the model 1 dataset that specified the number of captive generations experienced by

translocated cohorts (model 2; Table S3 in Appendix S2). We fit model 2 by substituting the

fixed effect of ESR vs. ISR strategy from model 1 with captive ancestry, a categorical fixed

effect with three levels classifying the focal ACT cohort as either part of an ISR (without any

captive ancestry), a head-starting program (ESR with < 1 generation in captivity), or a captive-

breeding program (ESR with > 1 direct ancestral generations in captivity). We were unable to

analyze captive ancestry as a continuous variable due to a lack of reporting of exact ancestral

generation times, and due to cohorts of potentially mixed ancestry. Model 2 also included

categorical fixed effects for sex, taxonomic class, fitness surrogate, and captive ancestry, and

two-way interactions between sex and fitness surrogate, sex and captive ancestry, and between

fitness surrogate and captive ancestry.

Finally, we developed model 3 to test the influence of enrichment strategies on relative

post-release performance among wild and translocated cohorts by analyzing a subset of studies

from the model 1 dataset that gathered relative post-release performance data with a wild-control

group from both enriched and unenriched cohorts within the same experimental design (Table S4

in Appendix S2). In model 3, we included categorical fixed effects for sex, taxonomic class,

fitness surrogate, ESR vs. ISR strategy, and enrichment. We also included two-way interactions

between enrichment and sex, enrichment and surrogate, enrichment and strategy.

*(e) Results reporting*

We addressed whether conservation programs produce organisms that perform similarly to or

better than wild-resident conspecifics after release by calculating the total probability that the log

odds ratio for a particular comparison is greater than or equal to zero. We also reported the

marginal posterior mode and 95% highest posterior density credible interval (CrI) of the odds

ratio (OR) by exponentiating posterior distributions of log odds ratios. We further interpreted

each effect size according to established threshold values delimiting small, medium, and large

magnitudes of effect (see Table 1; Cohen, 1992; Olivier *et al*., 2013). While these values provide

potential useful benchmarks for the assessment of biological importance in effect sizes

(Nakagawa & Cuthill, 2007), we focused our interpretations on the true effect size values and

associated credible intervals. We used the R package *emmeans* to report the marginal values of

individual parameters and overall model effects to account for the influence of other model

parameters (Lenth, 2021). All derived parameters are calculated across the entire posterior

distribution to carry through all uncertainty.

**(5) Meta-analysis validation**

*(a) Outlier identification*

We performed several meta-analysis validation tests to assess our dataset and findings for

influential outliers and publication bias (Koricheva, Jennions & Lau, 2013). We identified

outliers by calculating studentized deleted residuals for each comparison (Viechtbauer &

Cheung, 2010). Comparisons with estimates over 1.96 standard deviations from the overall mean

effect were considered outliers. We reran all models with outliers excluded to examine whether

our findings were significantly affected.

Table 1. Established recommendations for the interpretation of log odds ratio (LOR) and odds

ratio (OR) effect size magnitude and direction (Cohen, 1992; Olivier *et al*., 2013). Effect size

values in gray describe ACT comparisons where translocated cohorts have decreased odds of

out-performing their wild-resident counterparts.

| Magnitude, direction of effect | Log odds ratio (LOR) | Odds ratio (OR) |
| --- | --- | --- |
| Large, negative | -1.10 | 0.33 |
| Medium, negative | -0.62 | 0.54 |
| Small, negative | -0.20 | 0.82 |
| No effect | 0 | 1.00 |
| Small, positive | 0.20 | 1.22 |
| Medium, positive | 0.62 | 1.86 |
| Large, positive | 1.10 | 3.00 |

*(b) Publication bias*

To test for temporal patterns in effect sizes and potential time-lag bias (Jennions & Møller, 2002;

Koricheva *et al.*, 2013), we fit a linear model with our dataset's calculated effect sizes as the

response variable and year of publication as the predictor. We included study ID, group ID, and

all Linnaean taxonomic ranks below the level of phylum as random effects. To assess the dataset

for evidence of publication bias, we used Egger's regression to model the relationship of the

meta-analytic residuals of each effect size against their precision (Egger *et al.*, 1997; Nakagawa

& Santos, 2012). Funnel plots were also created to visualize this relationship and identify any

systematic absence of values, which could also indicate publication bias. To directly test the

influence of publication bias on our overall effect, we conducted a trim-and-fill analysis using

the *trimfill* function in the R package *metafor*, which quantifies funnel plot asymmetry and

estimates the number of missing studies on either side of the overall mean effect (Duval &

Tweedie, 2000; Viechtbauer, 2010).

**Table 2.** Total number of log odds ratio (LOR) estimates collected, organized by taxonomic

class and the fitness surrogate used as a metric for performance.

|  | Movement | Constitution | Growth | Survival | Reproduction | **Total** |
|---|---|---|---|---|---|---|
| Bivalvia | 0 | 0 | 0 | 6 | 0 | **6** |
| Gastropoda | 0 | 3 | 5 | 9 | 6 | **23** |
| Malacostraca | 0 | 4 | 0 | 14 | 5 | **23** |
| Echinoidea | 1 | 1 | 0 | 0 | 0 | **2** |
| Actinopterygii | 38 | 43 | 26 | 158 | 77 | **342** |
| Amphibia | 4 | 2 | 0 | 3 | 0 | **9** |
| Reptilia | 101 | 12 | 26 | 54 | 9 | **202** |
| Aves | 14 | 12 | 0 | 59 | 38 | **123** |
| Mammalia | 18 | 13 | 2 | 46 | 12 | **91** |
| **Total** | **176** | **90** | **59** | **349** | **147** | **821** |

**III. RESULTS**

**(1) Description of data**

Our initial Web of Science query returned 6906 unique articles. After filtering based on the inclusion criteria, our dataset comprised 821 unique effect sizes for relative performance of translocated organisms and their wild-resident conspecifics collected from 171 peer-reviewed studies (Table S1 in Appendix S2). The studies investigated 101 unique species in either *in situ* reinforcements (ISR; 599 effect sizes) or *ex situ* reinforcements (ESR; 222 effect sizes; Table 2). We calculated standardized effect sizes as log odds ratios (LORs) of performance metrics for translocated *versus* wild-resident cohorts such that positive values indicate translocated individuals exhibited higher relative performance than wild-resident conspecifics.

**(2) Meta-analysis**

We analyzed the LORs and subsets of these data using three linear mixed effects meta-analytical models with fixed effect moderator variables to determine the impact of organismal and methodological factors plus random effects to attribute sources of variation among effect sizes.

*(a) Model 1: Overall model*

The primary model's overall log odds ratio (LOR mode: -1.02; 95% CrI: -1.59 to -0.32) was converted to an odds ratio by exponentiating across the posterior distribution, indicating that translocated organisms have 63.9% decreased odds of out-performing their wild-resident counterparts (OR mode: 0.36; 95% CrI: 0.16 to 0.66). These effect size values correspond to a large magnitude of effect (Cohen, 1992; Olivier *et al.*, 2013). By quantifying the proportion of posterior samples with LOR estimates greater than or equal to zero, we infer that translocated

organisms had an overall 0.001 probability of out-performing wild-resident conspecifics

following release (Fig. 2). Disparity in post-release performance did not vary widely across

taxonomic classes, but instead all classes generally exhibited negative LOR effect sizes of

moderate-to-high magnitude, suggesting poor performance of translocated organisms relative to

wild-residents across taxa (Table S2 in Appendix S2). Similarly, relative post-release

performance was largely unaffected by ESR vs. ISR translocation strategy.

In studies involving mixed-sex cohorts, comparisons based on fitness components tend to

have lower LOR marginal posterior mode estimates than comparisons based on fitness

correlates, but widely overlapping credible intervals across fitness components and correlates

indicate a consensus in the interpretation of relative under-performance of translocated cohorts

across a variety of performance metrics (Table S2 in Appendix S2). More variable effect size

outcomes observed in sex-specific comparisons might suggest the presence of unequal life-

history consequences to translocation among sexes, but low sample sizes in constitution-,

growth-, and survival-based estimates, and uneven taxonomic representation in movement-based

estimates preclude conclusive interpretations.

*(b) Dataset heterogeneity*

Here we summarize the heterogeneity statistics from model 1, representing the full dataset (see

Table S5 in Appendix S2 for heterogeneity summary statistics of all three models).

Heterogeneity values from individual taxonomic levels were relatively small and even in

magnitude, but their combined taxonomic signal accounted for 60.6% (95% CrI: 47.0% to

74.7%) of total heterogeneity. Study- and group-level effects accounted for 23.9% (95% CrI:

12.0% to 34.3%) and 13.3% (95% CrI: 8.67% to 19.7%) of total heterogeneity, respectively.

*(c) Subset analyses*

Due to limited reporting of key moderator variables across studies, we analyzed subsets of the

data in two additional models to assess whether the amount of captive ancestry or the presence of

enrichment, respectively, influenced relative post-release performance of translocated cohorts. In

general, LOR marginal posterior mode estimates were similar for the overall model effects and

for parameters that also appeared in the primary model 1 above (Tables S3-S4 in Appendix S2).

Therefore, we limit the remainder of our reporting to model-specific parameters.

*(i) Model 2: Influence of captive ancestry*

Post-release performance was largely unaffected by the length of captive ancestry, except for

comparisons based on constitution and growth. In ISR studies involving constitution-based

comparisons, we observed that relocated organisms without any captive ancestry had a lower

probability (0.00; OR mode: 0.07; OR 95% CrI: 0.03 to 0.21) of out-performing wild-resident

conspecifics relative to head-started (0.15; OR mode: 0.55; OR 95% CrI: 0.22 to 1.28) and

captive-bred cohorts (0.25; OR mode: 0.21; OR 95% CrI: 0.02 to 2.51; Table S3 in Appendix

S2). Comparisons with growth-based estimates exhibited a similar pattern, but these growth-

related comparisons for relocated, head-started, and captive-bred cohorts were based on reduced

datasets of only two, eight, and one studies, respectively.

*(ii) Model 3: Influence of pre-release enrichment*

To test the influence of enrichment strategies on relative post-release performance among

translocated and wild cohorts, we analyzed a subset of studies that gathered data from both

enriched and unenriched cohorts within the same experimental design and compared their post-

release performances relative to a common wild-control group. Unenriched translocated

organisms had a 0.06 probability of out-performing their wild-resident conspecifics (OR mode:

0.06; 95% CrI: 0.002 to 1.06), whereas enriched translocated organisms had a 0.58 probability of

out-performing their wild counterparts (OR mode: 0.75; 95% CrI: 0.008 to 9.36; Fig. 3; Table S4

in Appendix S2).

*(d) Validation*

*(i) Outlier identification*

We identified 56 outlying effect sizes based on studentized deleted residuals greater in

magnitude than 1.96. Removal of outliers generally reduced effect sizes across all parameters

and caused the 95% credible intervals for the overall effects of model 1 and model 2 to overlap

with zero (Table S6-S8 in Appendix S2). However, outlier exclusion did not qualitatively affect

our overall findings nor the indices of publication bias. Moreover, the examination of all outlying

effect sizes revealed these effect sizes were calculated correctly and reflect legitimate patterns in

relative post-release performance (Aguinis, Gottfredson & Joo, 2013). Therefore, we completed

the final analysis with outliers included.

*(ii) Publication bias*

Visual examination of the marginal posterior distributions across years shows consistent effect

sizes, indicating no time-lag bias in our dataset (Fig. 4). Removing outliers did not affect patterns

of effect size magnitude across years (Fig. S1 in Appendix S3).

The Egger's regression slopes were significantly different from zero after being run with

both the dataset including (slope = 0.18; SE: 0.04; p < .001) and excluding (slope = 0.19; SE:

0.03; p < 0.001) outliers, which suggests the presence of publication bias. The resulting funnel

plots revealed the presence of some effect size asymmetry in the dataset including outliers (Fig.

5), and to a lesser degree in the dataset excluding outliers (Fig. S2 in Appendix S3). However,

the significant Egger's regression slopes for both datasets could be an artifact of the high model

heterogeneity and not publication bias (Higgins & Thompson, 2002).

The trim-and-fill analysis estimated that the overall model including outliers was missing

zero estimates to the left side of the mean and 3 estimates from the right side (SE: 2.83; p =

0.06), whereas the model excluding outliers had zero missing estimates to either side of the mean

(SE: 1.41; p = 0.5).

In conclusion, our validation tests indicate negligible amounts of publication bias.

Moreover, we consider our estimates to be highly conservative given the fact that conservation

translocations describing unsuccessful or uncertain outcomes are less likely to be published than

those with successful outcomes (Fischer and Lindenmayer, 2000; Miller *et al*., 2014; Reading *et*

*al*., 1997).

**IV. DISCUSSION**

Our meta-analysis gauged the efficacy of conservation translocation programs for benefitting

population and species persistence based on direct, *in situ* comparisons of animal performance

between translocated organisms and wild-resident conspecifics. We synthesized estimates of

relative performance based on five fitness-surrogate categories across 101 unique species

spanning nine taxonomic classes. Our findings indicate that translocated organisms tend to

under-perform relative to their wild counterparts, and that this pattern is largely independent of

taxonomic class or the type of fitness surrogate measured. This supports previous claims for the

existence of general issues across ACTs and suggests that translocated organisms are often at a

major disadvantage following release (see the Introduction; e.g., Snyder *et al.,* 1996; Fischer &

Lindenmayer, 2000; Pérez *et al.,* 2012; Germano *et al.,* 2015; Sullivan, Nowak & Kwiatkowski,

2015). Under-performance of translocated cohorts was also consistent regardless of translocation

strategy or the amount of captive ancestry, with some caveats. Finally, post-release performance

was markedly improved in translocated cohorts exposed to pre-release enrichment relative to

unenriched cohorts. Here we explore the implications of our findings and make

recommendations for future translocation program assessment and monitoring procedures.

**(1) Overall model**

The unexpected lack of variation in effect sizes across taxonomic classes hints at a general

influence of ACTs on animals, which is in direct odds with the large proportion of study

heterogeneity explained by taxonomic rank (Table S5 in Appendix S2). The high taxonomic

signal (64.29%) indicates that taxonomic position explains a majority of variation in effect sizes,

which would lead us to expect large differences in relative performance values among higher

taxonomic groups. For example, Dochtermann *et al*., (2019) observed a consistent pattern in a

phylogenetic meta-analysis quantifying behavioral heritability: little difference in effect size

among species and a low phylogenetic signal. However, we see the proportion of heterogeneity

explained by individual taxonomic tiers is consistently low, which could explain the minimal

differentiation in effect sizes at the level of taxonomic class. Equal variances estimated at each

taxonomic level is consistent with an explanation of constant phylogenetic inertia across the

phylogeny, in which equal branch lengths of the phylogeny coincide with equal taxonomic

branch lengths. Alternatively, if phylogenetic and taxonomic branching patterns do not coincide,

then the observed results would imply variation in taxonomic inertia across evolutionary time

(Hadfield & Nakagawa, 2010). To further explore these alternative explanations, and investigate

patterns of correlated evolution with respect to performance of translocated animals, comparative

studies focusing on biological patterns at the broadest taxonomic scales could adopt a mixed

phylogenetic and taxonomic approach until an appropriate phylogeny with empirically-derived

branch lengths can be resolved for all species in the data (Lynch, 1991; Hadfield & Nakagawa,

2010; Johnson *et al.*, 2018).

Our meta-analysis highlights a clear bias in taxonomic representation among ACTs, with

most studies involving charismatic species or those with economically important fisheries. Most

notably, there is a significant lack of ACTs involving amphibians that meet our inclusion criteria.

Amphibian populations are experiencing global declines and will likely require human

interventions in the form of ACTs to avoid extirpation (Wake & Vredenburg, 2008; Germano &

Bishop, 2009; Scheele *et al.*, 2021). Recent quantitative reviews describe a positive trend in the

success of amphibian translocations relative to earlier benchmarks (Dodd & Seigel, 1992;

Germano & Bishop, 2009). However, these efforts fail to consider comparisons among

translocated and wild-resident conspecifics as a potential index for project feasibility. Recent

amphibian studies using this comparative approach have provided novel insights regarding the

multi-generational influence of ACTs as well as potential pitfalls of cryopreservation techniques

on ACT feasibility (Cayuela *et al.*, 2019; Poo *et al.*, 2022). Future assessments of translocation

success should include simultaneous monitoring of a wild-control group from the donor and/or

recipient populations as an additional criterion for determining project feasibility and success.

Our primary model showed high consistency in effect sizes across fitness surrogates,

indicating that phenotypic traits considered as correlates of overall fitness have the potential to

characterize performance differences among translocated and wild-resident cohorts. We initially

predicted that fitness correlates would exhibit less consistent disparities across animal cohorts

since fitness correlates are less reliable proxies of fitness than fitness components (Hunt &

Hodgson, 2010). One explanation for the observed pattern is that fitness surrogates can vary in

how well they reflect overall individual fitness based on the relative length and positioning of the

measurement interval within the organism's lifespan (Pekkala *et al.*, 2011; Hendry *et al.*, 2018).

Additionally, the lack of differences in fitness surrogate effects may be due to unaccounted-for

differences in life-history strategies within and among study systems. Life-history theory

suggests that the uneven allocation of finite resources results in life-history trade-offs among

fitness components such that maximizing one fitness component can decrease overall lifetime

reproductive success (e.g., increased annual fecundity decreases annual survival; Haave-Audet,

Besson, Nakagawa, & Mathot, 2022; Moyes *et al.*, 2009; Roff, 2002; van Noordwijk & de Jong,

1986). Future ACT studies and meta-analyses on the topic should incorporate multiple fitness

components at different life stages to assess relative quality of released organisms, such that the

covariance among fitness components and thus the presence of fitness trade-offs or differing life-

history strategies can be tested at the within-study level (Fincke & Hadrys, 2001; Hendry *et al.*,

2018).

We were not able to evaluate sex-by-fitness surrogate interactions in the full dataset

because so few studies reported sex-specific effects. However, sufficient data from reptile studies

suggest that estimates of relative performance based on movement were substantially lower for

both male- and female-only cohorts relative to estimates from mixed-sex cohorts, indicating

greater disparity in post-release performance relative to wild conspecifics when males and

females were analyzed separately. Only when accounting for sex do our findings align with the

established relationship between maladaptive post-release movement patterns and reduced

survival in the herpetofaunal literature (Bonnet, Naulleau & Shine, 1999; Germano & Bishop,

2009). The influence of sex on post-release performance is also poorly represented in the

primary literature and rarely synthesized in subsequent reviews (Teixeira *et al.*, 2007). To our

knowledge, no other quantitative literature review or meta-analysis examining translocation

efficacy has included sex as a moderating factor, despite well-documented and systematic

differences in life history and ecology among the sexes (Immonen *et al.*, 2018; Tarka *et al.*,

2018).

**(2) Influence of captive ancestry**

We observed similar levels of under-performance among translocated cohorts relative to wild

conspecifics, regardless of whether translocated organisms were ISR (0 captive generations),

head-started (ESR with < 1 captive generation), or captive-bred (ESR with > 1 captive

generation). This contrasts with existing quantitative reviews describing a negative relationship

between captive ancestry and fitness, with ISRs performing relatively better than ESRs (Fischer

& Lindenmayer, 2000; Rummel *et al.*, 2016). One explanation for this discrepancy is the poor

reporting in the literature of ESR captive histories and animal origins which precluded us from

conducting a more powerful assessment of the effect of captive ancestry. As such, our head-

starting vs. captive-rearing levels possibly misclassified cohorts with ambiguous or mixed

captive histories, which could explain the lack of significant performance deficits commonly

reported in long-term captive-breeding populations (Araki *et al.*, 2007). It should also be noted

692 that the ACT strategies specific to certain taxonomic groups might be more prone to reporting

693 ambiguous captive ancestry information (Table S9). For example, reptilian, mammalian, and

694 avian studies regularly describe captive ancestries in great detail (Jones *et al*., 2002), whereas

695 tracking individual captive-ancestries in fisheries systems is impossible without genetic

696 parentage assignment (Lutz *et al*., 2021). This can be especially problematic if there is captive

697 ancestry in the purported wild-resident group, which would deflate expected fitness differences

698 between translocated and wild-resident conspecifics (Araki *et al*., 2007). Our findings suggest

699 that while ISRs avoid the maladaptive selective pressures associated with extended captivity,

700 managers should still be wary of individual- and population-level stressors characteristic of ISRs

701 that can lead to disparities in post-release performance like those observed in ESRs. This

702 underscores the necessity of ACT programs to report the capture histories of their organisms, and

703 to conduct multi-generational monitoring to ensure accurate measures of meaningful life-history

704 traits.

706 **(3) Influence of pre-release enrichment**

707 Disparity in post-release performance among translocated and wild individuals was significantly

708 reduced when translocated organisms were exposed to some form of pre-release enrichment.

709 This finding highlights the growing relevance of the behavioral ramifications of *in situ* and *ex*

710 *situ* reinforcements, where patterns of allelic heterozygosity and population growth rate have

711 historically been the primary factors used to assess translocation efficacy (Frankham, 2008;

712 Greggor *et al.,* 2016; Mathews *et al.*, 2005; Morris *et al.,* 2021). Our findings agree with those

713 from other meta-analyses showing that enriched, translocated organisms generally had higher

714 performance than non-enriched, translocated conspecifics in a variety of taxa (Tetzlaff *et al.*,

2019a; Zhang *et al.*, 2022). Since few studies incorporating pre-release enrichment also fulfilled

our inclusion criteria by directly comparing translocated cohorts to wild-resident conspecifics,

we were not able to refine our categorization of different enrichment strategies. Despite that, our

study provides the broadest evidence yet—and the first synthesized evidence to incorporate wild-

control groups—that pre-release enrichment strategies have the potential to reduce post-release

performance disparities between translocated and wild-resident organisms.

**(4) Future directions**

Despite notable positive outcomes, animal conservation translocations have experienced

historically low success rates without significant improvement observed over time (Berger-Tal,

Blumstein & Swaisgood, 2020). Ensuring future success of ACTs requires improved alignment

between the specific conservation benefits sought in the translocation (e.g., as defined in Annex

2 of IUCN/SSC, 2013) coupled with objective evaluation of success achieving the desired

benefits. Conservation biology is in the midst of a revolutionary push for more evidence-based

decision-making to replace historically anecdotal- or experience-driven strategies (Pullin &

Knight, 2001; Sutherland *et al.*, 2004). Formal meta-analysis is an important tool for

synthesizing existing literature and directing new veins of inquiry, particularly in the field of

conservation, where translocation studies are numerous but statistical power is limited by low

sample sizes (Moseby, Hill & Lavery, 2014). Here we discuss several aspects of ACT practices

and conservation goals that are relevant to increasing insight from future meta-analyses.

The conservation benefits of many ACTs (*sensu* IUCN/SSN, 2013) target population

persistence and preventing outright species extinction. Short-term strategies for achieving these

goals can be generalized according to three different ways to 'rescue' populations (Carlson *et al.*,

2014): increase population density and buffer against ecological effects and demographic

stochasticity (i.e., demographic rescue), increase genetic heterozygosity and reduce the

accumulation of deleterious recessive alleles (i.e., genetic rescue), or increase adaptive potential

(i.e., evolutionary rescue). Ultimately, to achieve these goals, translocation activities must

contribute to populations in ways that enable recipient populations to adapt to their environment,

increase average individual fitness, and thereby bolster population growth rate (Lande 1988;

Carlson *et al.,* 2014; Hendry *et al.,* 2018; Shaw, 2019). To evaluate the density-dependent

ecological dynamics associated with demographic rescue, the recipient population cannot serve

as the control group and so an allopatric population should be used instead. In our systematic

review, we found only one study that included both sympatric and allopatric control groups

(Molony *et al.*, 2006). Thus, our results can only be interpreted in light of how successfully

ACTs contribute to increasing mean fitness of the recipient population.

Our findings show that *in situ* reinforcements (ISRs) and *ex situ* reinforcements (ESRs)

produce organisms that under-perform and should therefore be implemented only after careful

consideration of the recipient population's trajectory and the existing environmental threats at the

recipient site (IUCN/SSC, 2013). For example, if a wild population is declining due to extrinsic

threats, then the release of under-performing translocated conspecifics is unlikely to reverse

declines (Heppell *et al.*, 1996). Alternatively, if environmental threats have been removed, then

the allelic diversity of a remnant wild population could benefit from the influx of translocated

cohorts, despite certain performance deficits (Jensen *et al.*, 2018; Cayuela *et al.*, 2019). Pilot

studies employing the post-release comparative techniques analyzed in this study could therefore

use relative post-release performance, population demographics, and environmental forecasts to

predict the efficacy of ACT programs.

The initial prevention and reversal of invasive species spread is a major aspect of global

habitat restoration which shares a conceptual basis with conservation translocations (Bright &

Smithson, 2001; Armstrong & Seddon, 2008) and thus stands to benefit from *in situ* comparisons

with wild-control groups. The potential for non-invasive species establishment and persistence

can be assessed through sympatric comparisons with non-conspecifics of overlapping ecological

niche, or through allopatric comparisons with conspecifics from the original donor population. If

an invasion is ongoing, direct comparisons between recent invaders and the offspring of

previous-generation invaders could also promote the investigation of the genetic basis of

invasive adaptations. Similar investigations could act as the prelude to assisted colonization or

rewilding efforts to ensure that species introduced to areas outside of their historic range are

capable of colonizing successfully without risk of disrupting existing ecosystems (Corlett, 2016;

Hunter-Ayad *et al.*, 2021).

The often-wide credible intervals in our findings suggest there are opportunities in which

ACTs can produce viable individuals and contribute to recipient populations. We recommend

that ACT managers include sympatric and allopatric wild-reference groups in their programs to

properly evaluate post-release performance of translocated organisms, and assess any negative

density-dependent impacts of ACTs on resident conspecifics (Mathews *et al.*, 2005; Molony *et*

*al.*, 2006). Managers should also take advantage of next-generation sequencing technologies and

parentage assignment software to promote long-term, multi-generational monitoring of relative

reproductive success and other fitness surrogates (Araki *et al.*, 2007).

**V. CONCLUSIONS**

(1) Translocated organisms generally exhibited poorer performance following release compared

to wild-resident conspecifics. The translocated organisms showed an overall 64% decreased odds

of out-performing their wild-resident counterparts (odds ratio posterior mode: 0.30; 95% CrI:

0.18 to 0.66) and this degree of disparity was consistent across taxonomic classes as well as

whether an *in situ versus ex situ* reinforcement (ISR vs. ESR) strategy had been implemented.

(2) The overall disparity between translocated and wild-resident cohorts was consistent across

studies regardless of the amount of captive ancestry in any particular implementation.

(3) Post-release performance of translocated organisms generally benefitted as a result of some

level of enrichment activity incorporated into the translocation program.

(4) The dataset exhibited a high overall taxonomic signal but low heterogeneity at higher

individual taxonomic levels. Variation across taxonomic levels suggests a potential influence of

shared evolutionary history on relative post-release performance, but more comprehensive

taxonomic sampling coupled with comparative analyses in a phylogenetic framework is

necessary to draw accurate conclusions.

(5) Overall, the disparity in post-release performance between translocated organisms and their

wild-resident counterparts was not affected by sex, taxonomic group, type of fitness surrogate

measured, or ESR vs. ISR translocation strategy.

(6) Only 2% of animal conservation translocation studies returned during the initial literature

search of our systematic review report or design translocations that use wild-resident control

groups. We suggest both sympatric and allopatric wild-reference groups will further enable post-

release performance of translocated organisms to be evaluated for any negative density-

dependent impacts of ACTs on resident conspecifics.

(7) Conservation managers should standardize the use of wild-resident controls and multiple

performance metrics that capture several different aspects of the life history in their assessments

of potential conservation translocation strategies to ensure population or species persistence.

**VI. ACKNOWLEDGEMENTS**

We would like to thank members of the D.A. Warner and M.E. Wolak labs, and students from

the Auburn University meta-analysis course (taught by A.E.W.) for providing comments on

earlier drafts of the manuscript.

**VII. AUTHOR CONTRIBUTIONS**

**Conceptualization:** IPG and MEW; **Data curation:** IPG and MEW; **Formal analysis:** IPG,

with input from AEW and MEW; **Investigation:** IPG; **Project administration:** MEW;

**Visualization:** IPG and MEW; **Writing - original manuscript:** IPG; **Writing - review and**

**editing:** AEW and MEW.

**VIII. DATA ACCESSIBILITY**

All data and R code are available from the Zenodo Digital Repository

https://doi.org/10.5281/zenodo.7535746

**IX. REFERENCES**

References included only in the meta-analysis dataset are identified with an asterisk (*).

*AALTONEN, K., BRYANT, A.A., HOSTETLER, J.A. & OLI, M.K. (2009) Reintroducing endangered

Vancouver Island marmots: Survival and cause-specific mortality rates of captive-born

versus wild-born individuals. *Biological Conservation* **142**, 2181–2190.

*AARESTRUP, K., BAKTOFT, H., KOED, A., DEL VILLAR-GUERRA, D. & THORSTAD, E. (2014)

Comparison of the riverine and early marine migration behaviour and survival of wild and

hatchery-reared sea trout *Salmo trutta* smolts. *Marine Ecology Progress Series* **496**, 197–

206.

*AGNALT, A.L., KRISTIANSEN, T.S. & JØRSTAD, K.E. (2007) Growth, reproductive cycle, and

movement of berried European lobsters (*Homarus gammarus*) in a local stock off

southwestern Norway. *ICES Journal of Marine Science* **64**, 288–297.

AGUINIS, H., GOTTFREDSON, R.K. & JOO, H. (2013) Best-practice recommendations for defining,

identifying, and handling outliers. *Organizational Research Methods* **16**, 270–301.

ALLEAUME-BENHARIRA, M., PEN, I.R. & RONCE, O. (2006) Geographical patterns of adaptation

within a species' range: Interactions between drift and gene flow. *Journal of Evolutionary*

*Biology* **19**, 203–215.

*ANDERSON, J.H., FAULDS, P.L., ATLAS, W.I. & QUINN, T.P. (2013) Reproductive success of

captively bred and naturally spawned Chinook salmon colonizing newly accessible habitat.

*Evolutionary Applications* **6**, 165–179.

ARAKI, H., BEREJIKIAN, B.A., FORD, M.J. & BLOUIN, M.S. (2008) Fitness of hatchery-reared

salmonids in the wild. *Evolutionary Applications* **1**, 342–355.

ARAKI, H., COOPER, B. & BLOUIN, M.S. (2007) Genetic effects of captive breeding cause a rapid,

cumulative fitness decline in the wild. *Science* **318**, 100–103.

ARMSTRONG, D.P. & SEDDON, P.J. (2008) Directions in reintroduction biology. *Trends in*

*Ecology and Evolution* **23**, 20–25.

*ATTUM, O. & CUTSHALL, C.D. (2015) Movement of translocated turtles according to

translocation method and habitat structure. *Restoration Ecology* **23**, 588–594.

*BACON, L., ROBERT, A. & HINGRAT, Y. (2019) Long lasting breeding performance differences

between wild-born and released females in a reinforced North African houbara bustard

(*Chlamydotis undulata undulata*) population: a matter of release strategy. *Biodiversity and*

*Conservation* **28**, 553–570.

BAJOMI, B., PULLIN, A.S., STEWART, G.B. & TAKÁCS-SÁNTA, A. (2010) Bias and dispersal in the

animal reintroduction literature. *Oryx* **44**, 358–365.

BARNOSKY, A.D., MATZKE, N., TOMIYA, S., WOGAN, G.O.U., SWARTZ, B., QUENTAL, T.B.,

MARSHALL, C., MCGUIRE, J.L., LINDSEY, E.L., MAGUIRE, K.C., MERSEY, B. & FERRER,

E.A. (2011) Has the Earth's sixth mass extinction already arrived? *Nature* **471**, 51–57.

*BAUDER, J.M., CASTELLANO, C., JENSEN, J.B., STEVENSON, D.J. & JENKINS, C.L. (2014)

Comparison of movements, body weight, and habitat selection between translocated and

resident gopher tortoises. *The Journal of Wildlife Management* **78**, 1444–1455.

BENNETT, A.M., STEINER, J., CARSTAIRS, S., GIELENS, A. & DE, C.M.D. (2017) A question of

scale: replication and the effective evaluation of conservation interventions. *Facets* **2**, 892–

909.

BERGER-TAL, O., BLUMSTEIN, D.T. & SWAISGOOD, R.R. (2020) Conservation translocations: a

review of common difficulties and promising directions. *Animal Conservation* **23**, 121–131.

*BERINGER, J., HANSEN, L.P., DEMAND, J.A., SARTWELL, J., WALLENDORF, M. & MANGE, R.

(2002) Efficacy of translocation to control urban deer in Missouri: costs, efficiency, and

outcome. *Wildlife Society Bulletin* **30**, 767–774.

*BLANCHARD, B.M. & KNIGHT, R.R. (1995) Biological consequences of relocating grizzly bears

in the Yellowstone ecosystem. *The Journal of Wildlife Management* **59**, 560–565.

*BLYTHE, R.M., SMYSER, T.J., JOHNSON, S.A. & SWIHART, R.K. (2015) Post-release survival of

captive-reared Allegheny woodrats. *Animal Conservation* **18**, 186–195.BOLAM, F.C., MAIR,

L., ANGELICO, M., BROOKS, T.M., BURGMAN, M., HERMES, C., HOFFMANN, M., MARTIN,

R.W., MCGOWAN, P.J.K., RODRIGUES, A.S.L., RONDININI, C., WESTRIP, J.R.S., WHEATLEY,

H., BEDOLLA-GUZMÁN, Y., CALZADA, J., ET AL. (2021) How many bird and mammal

extinctions has recent conservation action prevented? *Conservation Letters* **14**, e12762.

*BOLLAND, J.D., COWX, I.G. & LUCAS, M.C. (2009) Dispersal and survival of stocked cyprinids

in a small English river: comparison with wild fishes using a multi-method approach.

*Journal of Fish Biology* **74**, 2313–2328.

BONDURIANSKY, R. & CHENOWETH, S.F. (2009) Intralocus sexual conflict. *Trends in Ecology*

*and Evolution* **24**, 280–288.

BONNET, X., NAULLEAU, G. & SHINE, R. (1999) The dangers of leaving home: dispersal and

mortality in snakes. *Biological Conservation* **89**, 39–50.

*BORGSTROM, R., SKAALA, Ø. & AASTVEIT, A.H. (2002) High mortality in introduced brown

trout depressed potential gene flow to a wild population. *Journal of Fish Biology* **61**, 1085–

1097.

*BOSE, A.P.H., ZAYONC, D., AVRANTINIS, N., FICZYCZ, N., FISCHER-RUSH, J., FRANCIS, F.T.,

GRAY, S., MANNING, F., ROBB, H., SCHMIDT, C., SPICE, C., UMEDALY, A., WARDEN, J. &

CÔTÉ, I.M. (2019) Effects of handling and short-term captivity: a multi-behaviour approach

using red sea urchins, *Mesocentrotus franciscanus*. *PeerJ* **7**, e6556.

*BOSSON, C.O., PALME, R. & BOONSTRA, R. (2013) Assessing the impact of live-capture,

confinement, and translocation on stress and fate in eastern gray squirrels. *Journal of*

*Mammalogy* **94**, 1401–1411.

BOURASS, K. & HINGRAT, Y. (2015) Diet of released captive-bred North-African houbara

bustards. *European Journal of Wildlife Research* **61**, 563–574.

BOUZAT, J.L., JOHNSON, J.A., TOEPFER, J.E., SIMPSON, S.A., ESKER, T.L. & WESTEMEIER, R.L.

(2009) Beyond the beneficial effects of translocations as an effective tool for the genetic

restoration of isolated populations. *Conservation Genetics* **10**, 191–201.

*BRADLEY, E.H., PLETSCHER, D.H., BANGS, E.E., KUNKEL, K.E., SMITH, D.W., MACK, C.M.,

MEIER, T.J., FONTAINE, J.A., NIEMEYER, C.C. & JIMENEZ, M.D. (2005) Evaluating

translocation as a nonlethal method to reduce livestock conflicts in the northwestern United

States. *Conservation Biology* **19**, 1498–1508.

*BRAND, L.A., FARNSWORTH, M.L., MEYERS, J., DICKSON, B.G., GROUIOS, C., SCHEIB, A.F. &

SCHERER, R.D. (2016) Mitigation-driven translocation effects on temperature, condition,

growth, and mortality of Mojave desert tortoise (*Gopherus agassizii*) in the face of solar

energy development. *Biological Conservation* **200**, 104–111.

*BRANNELLY, L.A., HUNTER, D.A., SKERRATT, L.F., SCHEELE, B.C., LENGER, D., MCFADDEN,

M.S., HARLOW, P.S. & BERGER, L. (2016) Chytrid infection and post-release fitness in the

reintroduction of an endangered alpine tree frog. *Animal Conservation* **19**, 153–162.

BRICHIERI-COLOMBI, T.A., LLOYD, N.A., MCPHERSON, J.M. & MOEHRENSCHLAGER, A. (2019)

Limited contributions of released animals from zoos to North American conservation

translocations. *Conservation Biology* **33**, 33–39.

BRIGHT, P.W. & SMITHSON, T.J. (2001) Biological invasions provide a framework for

reintroductions: Selecting areas in England for pine marten releases. *Biodiversity and*

*Conservation* **10**, 1247–1265.

*BRITTAS, R., MARCSTRÖM, V., KENWARD, R.E. & KARLBOM, M. (1992) Survival and breeding

success of reared and wild ring-necked pheasants in Sweden. *The Journal of Wildlife*

*Management* **56**, 368–376.

BROWN, C., DAVIDSON, T. & LALAND, K. (2003) Environmental enrichment and prior experience

of live prey improve foraging behaviour in hatchery-reared Atlantic salmon. *Journal of Fish*

*Biology* **63**, 187–196.

*BROWN, J.R., BISHOP, C.A. & BROOKS, R.J. (2009) Effectiveness of short-distance translocation

and its effects on western rattlesnakes. *The Journal of Wildlife Management* **73**, 419–425.

BROWN, J.L., COLLOPY, M.W., GOTT, E.J., JUERGENS, P.W., MONTOYA, A.B. & HUNT, W.G.

(2006) Wild-reared aplomado falcons survive and recruit at higher rates than hacked falcons

in a common environment. *Biological Conservation* **131**, 453–458.

*BURNSIDE, R.J., COLLAR, N.J. & DOLMAN, P.M. (2017) Comparative migration strategies of

wild and captive-bred Asian houbara *Chlamydotis macqueenii*. *Ibis* **159**, 374–389.

*BURNSIDE, R.J., COLLAR, N.J. & DOLMAN, P.M. (2018) Dataset on the numbers and proportion

of mortality attributable to hunting, trapping, and powerlines in wild and captive-bred

migratory Asian houbara *Chlamydotis macqueenii*. *Data in Brief* **21**, 1848–1852.

*BUTLER, H., MALONE, B. & CLEMANN, N. (2005) The effects of translocation on the spatial

ecology of tiger snakes (*Notechis scutatus*) in a suburban landscape. *Wildlife Research* **32**,

165.

CARLSON, S.M., CUNNINGHAM, C.J. & WESTLEY, P.A.H. (2014) Evolutionary rescue in a

changing world. *Trends in Ecology & Evolution* **29**, 521–530.

CAYUELA, H., GILLET, L., LAUDELOUT, A., BESNARD, A., BONNAIRE, E., LEVIONNOIS, P.,

MUTHS, E., DUFRÊNE, M. & KINET, T. (2019) Survival cost to relocation does not reduce

population self-sustainability in an amphibian. *Ecological Applications* **29**, 1–15.

CEBALLOS, G., EHRLICH, P.R., BARNOSKY, A.D., GARCÍA, A., PRINGLE, R.M. & PALMER, T.M.

(2015) Accelerated modern human–induced species losses: entering the sixth mass

extinction. *Science Advances* **1**, e1400253.

CEBALLOS, G., EHRLICH, P.R. & DIRZO, R. (2017) Biological annihilation via the ongoing sixth

mass extinction signaled by vertebrate population losses and declines. *Proceedings of the*

*National Academy of Sciences of the United States of America* **114**, E6089–E6096.

CEBALLOS, G., EHRLICH, P.R. & RAVEN, P.H. (2020) Vertebrates on the brink as indicators of

biological annihilation and the sixth mass extinction. *Proceedings of the National Academy*

*of Sciences of the United States of America* **117**, 13596–13602.

*CHAMPAGNON, J., GUILLEMAIN, M., ELMBERG, J., MASSEZ, G., CAVALLO, F. & GAUTHIER-

CLERC, M. (2012) Low survival after release into the wild: assessing "the burden of

captivity" on mallard physiology and behaviour. *European Journal of Wildlife Research* **58**,

255–267.

CHINN, S. (2000) A simple method for converting an odds ratio to effect size for use in

meta‐analysis. *Statistics in Medicine* **19**, 3127–3131.

*CHITTARO, P., JOHNSON, L., TEEL, D., MORAN, P., SOL, S., MACNEALE, K. & ZABEL, R. (2018)

Variability in the performance of juvenile Chinook salmon is explained primarily by when

and where they resided in estuarine habitats. *Ecology of Freshwater Fish* **27**, 857–873.

*CHITTENDEN, C.M., SURA, S., BUTTERWORTH, K.G., CUBITT, K.F., PLANTALECH MANEL-LA, N.,

BALFRY, S., ØKLAND, F. & MCKINLEY, R.S. (2008) Riverine, estuarine and marine

migratory behaviour and physiology of wild and hatchery-reared coho salmon

*Oncorhynchus kisutch* (Walbaum) smolts descending the Campbell River, BC, Canada.

*Journal of Fish Biology* **72**, 614–628.

*CLIFFORD, D.L., WOODROFFE, R., GARCELON, D.K., TIMM, S.F. & MAZET, J. A. K. (2007) Using

pregnancy rates and perinatal mortality to evaluate the success of recovery strategies for

endangered island foxes. *Animal Conservation* **10**, 442–451.

CLUTTON-BROCK, T.H. & HARVEY, P.H. (1977) Primate ecology and social organization.

*Journal of Zoology* **183**, 1–39.

COHEN, J. (1992) Quantitative methods in psychology: a power primer. *Psychological Bulletin*

**112**, 1155–1159.

*COLLIS, K., ROBY, D.D., CRAIG, D.P., RYAN, B.A. & LEDGERWOOD, R.D. (2001) Colonial

waterbird predation on juvenile salmonids tagged with passive integrated transponders in

the Columbia River estuary: vulnerability of different salmonid species, stocks, and rearing

types. *Transactions of the American Fisheries Society* **130**, 385–396.

CORLETT, R.T. (2016) Restoration, reintroduction, and rewilding in a changing world. *Trends in*

*Ecology and Evolution* **31**, 453–462.

COULSON, T., BENTON, T.G., LUNDBERG, P., DALL, S.R.X., KENDALL, B.E. & GAILLARD, J.M.

(2006) Estimating individual contributions to population growth: evolutionary fitness in

ecological time. *Proceedings of the Royal Society B: Biological Sciences* **273**, 547–555.

CRATES, R., STOJANOVIC, D. & HEINSOHN, R. (2022) The phenotypic costs of captivity.

*Biological Reviews*, 434–449.

\*CROMWELL, J.A., WARREN, R.J. & HENDERSON, D.W. (1999) Live-capture and small-scale

relocation of urban deer on Hilton Head Island, South Carolina. *Wildlife Society Bulletin*

*(1973-2006)* **27**, 1025–1031.

\*DANNEWITZ, J., PETERSSON, E., DAHL, J., PRESTEGAARD, T., LÖF, A.-C. & JÄRVI, T. (2004)

Reproductive success of hatchery-produced and wild-born brown trout in an experimental

stream. *Journal of Applied Ecology* **41**, 355–364.

\*DAVIS, J.L.D., ECKERT-MILLS, M.G., YOUNG-WILLIAMS, A.C., HINES, A.H. & ZOHAR, Y.

(2005) Morphological conditioning of a hatchery-raised invertebrate, *Callinectes sapidus*, to

improve field survivorship after release. *Aquaculture* **243**, 147–158.

\*DAVIS, J.L.D., YOUNG-WILLIAMS, A.C., AGUILAR, R., CARSWELL, B.L., GOODISON, M.R.,

HINES, A.H., KRAMER, M.A., ZOHAR, Y. & ZMORA, O. (2004) Differences between

hatchery-raised and wild blue crabs: implications for stock enhancement potential.

*Transactions of the American Fisheries Society* **133**, 1–14.

\*DAVIS, M.J., WOO, I., ELLINGS, C.S., HODGSON, S., BEAUCHAMP, D.A., NAKAI, G. & DE LA

CRUZ, S.E.W. (2018) Integrated diet analyses reveal contrasting trophic niches for wild and

hatchery juvenile chinook salmon in a large river delta. *Transactions of the American*

*Fisheries Society* **147**, 818–841.

\*DAVIS, M.J., WOO, I., ELLINGS, C.S., HODGSON, S., BEAUCHAMP, D.A., NAKAI, G. & DE LA

CRUZ, S.E.W. (2019) Freshwater tidal forests and estuarine wetlands may confer early life

growth advantages for delta-reared chinook salmon. *Transactions of the American Fisheries*

*Society* **148**, 289–307.

DAY, T. & BONDURIANSKY, R. (2011) A unified approach to the evolutionary consequences of

genetic and nongenetic inheritance. *American Naturalist* **178**.

DEGREGORIO, B.A., SPERRY, J.H., TUBERVILLE, T.D. & WEATHERHEAD, P.J. (2017)

Translocating ratsnakes: Does enrichment offset negative effects of time in captivity?

*Wildlife Research* **44**, 438–448.

DEGREGORIO, B.A., WEATHERHEAD, P.J., TUBERVILLE, T.D. & SPERRY, J.H. (2013) Time in

captivity affects foraging behavior of ratsnakes: implications for translocation.

*Herpetological Conservation and Biology* **8**, 581–590.

*DEL REAL, S.C., WORKMAN, M. & MERZ, J. (2012) Migration characteristics of hatchery and

natural-origin *Oncorhynchus mykiss* from the lower Mokelumne River, California.

*Environmental Biology of Fishes* **94**, 363–375.

*DELGADO, G.A., BARTELS, C.T., GLAZER, R.A., BROWN-PETERSON, N.J. & MCCARTHY, K.J.

(2004) Translocation as a strategy to rehabilitate the queen conch (*Strombus gigas*)

Population in the Florida Keys. *Fishery Bulletin* **102**, 278–288.

*DEVAN-SONG, A., MARTELLI, P., DUDGEON, D., CROW, P., ADES, G. & KARRAKER, N.E. (2016)

Is long-distance translocation an effective mitigation tool for white-lipped pit vipers

(*Trimeresurus albolabris*) in South China? *Biological Conservation* **204**, 212–220.

DICKENS, M.J., DELEHANTY, D.J. & MICHAEL ROMERO, L. (2010) Stress: An inevitable

component of animal translocation. *Biological Conservation* **143**, 1329–1341.

DOCHTERMANN, N.A., SCHWAB, T., ANDERSON BERDAL, M., DALOS, J. & ROYAUTÉ, R. (2019)

The heritability of behavior: a meta-analysis. *Journal of Heredity* **110**, 403–410.

DODD, C.K. & SEIGEL, R.A. (1992) Relocation, repatriation, and translocation of amphibians and

reptiles: are they conservation strategies that work? *Biological Conservation* **62**, 230.

*DRAKE, K.K., NUSSEAR, K.E., ESQUE, T.C., BARBER, A.M., VITTUM, K.M., MEDICA, P.A.,

TRACY, C.R. & HUNTER JR, K.W. (2012) Does translocation influence physiological stress

in the desert tortoise? *Animal Conservation* **15**, 560–570.

DUCATEZ, S. & SHINE, R. (2019) Life-history traits and the fate of translocated populations.

*Conservation Biology* **33**, 853–860.

DUVAL, S. & TWEEDIE, R. (2000) Trim and fill: a simple funnel-plot-based method. *Biometrics*

**56**, 455–463.

EGGER, M., SMITH, G.D., SCHNEIDER, M. & MINDER, C. (1997) Bias in meta-analysis detected by

a simple, graphical test. *British Medical Journal* **315**, 629–634.

*ELSEY, R.M., JOANEN, T., MCNEASE, L. & KINLER, N. (1992) Growth rates and body condition

factors of *Alligator mississippiensis* in coastal Louisiana wetlands: A comparison of wild

and farm-released juveniles. *Comparative biochemistry and physiology. Part A, Molecular*

*& integrative physiology* **103A**, 667–672.

*ELSEY, R.M., MCNEASE, L. & JOANEN, T. (2000) Louisiana's alligator ranching programme: a

review and analysis of releases of captive-raised juveniles. In *Crocodilian Biology and*

*Evolution* pp. 426–441. Surrey Beatty & Sons, Chipping Norton.

*ESCOBAR, R.A., BESIER, E. & HAYES, W.K. (2010) Evaluating headstarting as a management

tool: post- release success of green iguanas (*Iguana iguana*) in Costa Rica. *International*

*Journal of Biodiversity and Conservation* **2**, 204–214.

*ESQUE, T., NUSSEAR, K., DRAKE, K., WALDE, A., BERRY, K., AVERILL-MURRAY, R.,

WOODMAN, A., BOARMAN, W., MEDICA, P., MACK, J. & HEATON, J. (2010) Effects of

subsidized predators, resource variability, and human population density on desert tortoise

populations in the Mojave Desert, USA. *Endangered Species Research* **12**, 167–177.

*EVANS, A.F., HOSTETTER, N.J., COLLIS, K., ROBY, D.D. & LOGE, F.J. (2014) Relationship

between Juvenile Fish Condition and Survival to Adulthood in Steelhead. *Transactions of*

*the American Fisheries Society* **143**, 899–909.

*EVANS, M.L., JOHNSON, M.A., JACOBSON, D., WANG, J., HOGANSEN, M. & O'MALLEY, K.G.

(2016) Evaluating a multi-generational reintroduction program for threatened salmon using

genetic parentage analysis. *Canadian Journal of Fisheries and Aquatic Sciences* **73**, 844–

852.

EVANS, M.L., WILKE, N.F., O'REILLY, P.T. & FLEMING, I.A. (2014) Transgenerational effects of

parental rearing environment influence the survivorship of captive-born offspring in the

wild. *Conservation Letters* **7**, 371–379.

FAIRBAIRN, D.J. (2013) *Odd couples: Extraordinary differences between the sexes in the animal*

*kingdom*. Princeton University Press.

FAIRBAIRN, D.J. & REEVE, J.P. (2001) Natural selection: Measuring selection in natural

populations. In *Evolutionary Ecology: Concepts and case studies* (eds C. FOX, D.A. ROFF &

D.J. FAIRBAIRN), Oxford University Press, NY.

*FARNSWORTH, M.L., DICKSON, B.G., ZACHMANN, L.J., HEGEMAN, E.E., CANGELOSI, A.R.,

JACKSON, T.G. & SCHEIB, A.F. (2015) Short-term space-use patterns of translocated Mojave

Desert tortoise in Southern California. *PLOS ONE* **10**, e0134250.

*FARNSWORTH, S.D. & SEIGEL, G., RICHARD A. (2013) Responses, movements, and survival of

relocated box turtles during the construction of the Inter-County Connector Highway in

Maryland. *Transportation Research Record Journal of the Transportation Research Board*

**2362**, 1–8.

FARQUHARSON, K.A., HOGG, C.J. & GRUEBER, C.E. (2018) A meta-analysis of birth-origin

effects on reproduction in diverse captive environments. *Nature Communications* **9**, 1–10.

FARQUHARSON, K.A., HOGG, C.J. & GRUEBER, C.E. (2021) Offspring survival changes over

generations of captive breeding. *Nature Communications* **12**, 1–9.

*FERNANDO, P., LEIMGRUBER, P., PRASAD, T. & PASTORINI, J. (2012) Problem-elephant

translocation: translocating the problem and the elephant? *PLoS ONE* **7**, e50917.

*FERRETTI, M., PACI, G., PORRINI, S., GALARDI, L. & BAGLIACCA, M. (2010) Habitat use and

home range traits of resident and relocated hares (*Lepus europaeus*, Pallas). *Italian Journal*

*of Animal Science* **9**, e54.

FICETOLA, G.F. & DE BERNARDI, F. (2005) Supplementation or *in situ* conservation? Evidence of

local adaptation in the Italian agile frog *Rana latastei* and consequences for the management

of populations. *Animal Conservation* **8**, 33–40.

*FINSTAD, B. & HEGGBERGET, T.G. (1993) Migration, growth and survival of wild and hatchery-

reared anadromous Arctic charr (*Salvelinus alpinus*) in Finnmark, Northern Norway.

*Journal of Fish Biology* **43**, 303–312.

FINCKE, O.M. & HADRYS, H. (2001) Unpredictable offspring survivorship in the damselfly,

*megaloprepus coerulatus*, shapes parental behavior, constrains sexual selection, and

challenges traditional fitness estimates. *Evolution* **55**, 762–772.

FISCHER, J. & LINDENMAYER, D.B. (2000) An assessment of the published results of animal

relocations. *Biological Conservation* **96**, 1–11.

*FJELLHEIM, A., RADDUM, G.G. & BARLAUP, B.T. (1995) Dispersal, growth and mortality of

brown trout (*Salmo trutta L.*) stocked in a regulated west Norwegian river. *Regulated*

*Rivers: Research & Management* **10**, 137–145.

*FLÁVIO, H., AARESTRUP, K., JEPSEN, N. & KOED, A. (2019) Naturalised Atlantic salmon smolts

1096   are more likely to reach the sea than wild smolts in a lowland fjord. *River Research and*

1097   *Applications* **35**, 216–223.

1098 *FOLEY, A.M., PIERCE, B., HEWITT, D.G., DEYOUNG, R.W., CAMPBELL, T.A., HELLICKSON,

1099   M.W., RANCH, K., BOX, P.O., FEILD, J., RANCH, K., BOX, P.O., MITCHELL, S., PARKS, T.,

1100   LOCKWOOD, M.A., PARKS, T., ET AL. (2008) Survival and movements of translocated white-

1101   tailed deer in South Texas.

1102 FORD, M.J. (2002) Selection in captivity during supportive breeding may reduce fitness in the

1103   wild. *Conservation Biology* **16**, 815–825.

1104 FORD, M.J., MURDOCH, A. & HOWARD, S. (2012) Early male maturity explains a negative

1105   correlation in reproductive success between hatchery-spawned salmon and their naturally

1106   spawning progeny. *Conservation Letters* **5**, 450–458.

1107 FORD, M.J., MURDOCH, A.R., HUGHES, M.S., SEAMONS, T.R. & LAHOOD, E.S. (2016)

1108   Broodstock history strongly influences natural spawning success in hatchery steelhead

1109   (*Oncorhynchus mykiss*). *PLoS ONE*, 1–20.

1110 FRANKHAM, R. (2008) Genetic adaptation to captivity in species conservation programs.

1111   *Molecular Ecology* **17**, 325–333.

1112 FURLAN, E.M., GRUBER, B., ATTARD, C.R.M., WAGER, R.N.E., KEREZSY, A., FAULKS, L.K.,

1113   BEHEREGARAY, L.B. & UNMACK, P.J. (2020) Assessing the benefits and risks of

1114   translocations in depauperate species: A theoretical framework with an empirical validation.

1115   *Journal of Applied Ecology* **57**, 831–841.

1116 GELMAN, A. (2006) Prior distributions for variance parameters in hierarchical models (comment

1117   on article by Browne and Draper). *Bayesian Analysis* **1**, 515–534.

GERMANO, J.M. & BISHOP, P.J. (2009) Suitability of amphibians and reptiles for translocation.

*Conservation Biology* **23**, 7–15.

GERMANO, J.M., FIELD, K.J., GRIFFITHS, R.A., CLULOW, S., FOSTER, J., HARDING, G. &

SWAISGOOD, R.R. (2015) Mitigation-driven translocations: are we moving wildlife in the

right direction? *Frontiers in Ecology and the Environment* **13**, 100–105.

*GILMARTIN, W.G., SLOAN, A.C., HARTING, A.L., JOHANOS, T.C., BAKER, J.D., BREESE, M. &

RAGEN, T.J. (2011) Rehabilitation and relocation of young Hawaiian monk seals (*Monachus*

*schauinslandi*). *Aquatic Mammals* **37**, 332–341.

GLASS, G. V. (1976) Primary, secondary, and meta-analysis of research. *Educational Researcher*

**5**, 3–8.

*GOETZ, F.A., JEANES, E., MOORE, M.E. & QUINN, T.P. (2015) Comparative migratory behavior

and survival of wild and hatchery steelhead (*Oncorhynchus mykiss*) smolts in riverine,

estuarine, and marine habitats of Puget Sound, Washington. *Environmental Biology of*

*Fishes* **98**, 357–375.

GOLDENBERG, S.Z., OWEN, M.A., BROWN, J.L., WITTEMYER, G., OO, Z.M. & LEIMGRUBER, P.

(2019) Increasing conservation translocation success by building social functionality in

released populations. *Global Ecology and Conservation* **18**, e00604.

*GOOSSEN, J.P., VANDER LEE, R., KRUSE, C., GRATTO-TREVOR, C.L. & WESTWORTH, S.M.

(2011) Resightings of captive-reared and wild piping plovers from Saskatchewan, Canada.

*Wader Study Group Bulletin* **118**, 118–122.

GREGGOR, A.L., BERGER-TAL, O., BLUMSTEIN, D.T., ANGELONI, L., BESSA-GOMES, C.,

BLACKWELL, B.F., ST CLAIR, C.C., CROOKS, K., DE SILVA, S., FERNÁNDEZ-JURICIC, E.,

GOLDENBERG, S.Z., MESNICK, S.L., OWEN, M., PRICE, C.J., SALTZ, D., ET AL. (2016)

Research priorities from animal behaviour for maximising conservation progress. *Trends in Ecology and Evolution* **31**, 953–964.

GRIFFITH, B., SCOTT, M.J., CARPENTER, J.W. & REED, C. (1989) Translocation as a species conservation tool: status and strategy. *Science* **245**, 477–480.

GUREVITCH, J., KORICHEVA, J., NAKAGAWA, S. & STEWART, G. (2018) Meta-analysis and the science of research synthesis. *Nature* **555**, 175–182.

HAAVE-AUDET, E., BESSON, A.A., NAKAGAWA, S. & MATHOT, K.J. (2022) Differences in resource acquisition, not allocation, mediate the relationship between behaviour and fitness: a systematic review and meta-analysis. *Biological Reviews* **97**, 708–731.

HADDOCK, C.K., RINDSKOPF, D. & SHADISH, W.R. (1998) Using odds ratios as effect sizes for meta-analysis of dichotomous data: a primer on methods and issues. *Psychological Methods* **3**, 339–353.

HADFIELD, J.D. (2010) MCMCglmm: MCMC methods for multi-response GLMMs in R. *Journal of Statistical Software* **33**, 1–22.

HADFIELD, J.D. & NAKAGAWA, S. (2010) General quantitative genetic methods for comparative biology: phylogenies, taxonomies and multi-trait models for continuous and categorical characters. *Journal of Evolutionary Biology* **23**, 494–508.

*HADJISTERKOTIS, E. (1999) The survival of captive bred chukar *Alectoris chukar cypriotes*, released for restocking in Cyprus. Zeitschrift Fur Jagdwissenshaft **45**, 238–249.

*HAGELIN, A., CALLES, O., GREENBERG, L., PICCOLO, J. & BERGMAN, E. (2016) Spawning migration of wild and supplementary stocked landlocked Atlantic salmon (*Salmo Salar*). *River Research and Applications* **32**, 383–389.

*HAMEAU, O. & MILLON, A. (2019) Assessing the effectiveness of bird rehabilitation:

temporarily captive-reared little owls (*Athene noctua*) experience a similar recruitment rate

as wild birds. *Journal of Ornithology* **160**, 581–585.

*HANSEN, S.C. & GOSSELIN, L.A. (2016) Are hatchery-reared abalone naïve of predators?

Comparing the behaviours of wild and hatchery-reared northern abalone, *Haliotis*

*kamtschatkana* (Jonas, 1845). *Aquaculture Research* **47**, 1727–1736.

*HARING, M.W., JOHNSTON, T.A., WIEGAND, M.D., FISK, A.T. & PITCHER, T.E. (2016)

Differences in egg quantity and quality among hatchery- and wild-origin Chinook salmon

(*Oncorhynchus tshawytscha*). *Canadian Journal of Fisheries and Aquatic Sciences* **73**, 737–

746.

*HEATH, S.R., KERSHNER, E.L., COOPER, D.M., LYNN, S., TURNER, J.M., WARNOCK, N.,

FARABAUGH, S., BROCK, K. & GARCELON, D.K. (2008) Rodent control and food

supplementation increase productivity of endangered San Clemente loggerhead shrikes

(*Lanius ludovicianus mearnsi*). *Biological Conservation* **141**, 2506–2515.

HEATH, D.D., HEATH, J.W., BRYDEN, C.A., JOHNSON, R.M. & FOX, C.W. (2003) Rapid evolution

of egg size in captive Salmon. *Science* **299**, 1738–1741.

*HELLSTEDT, P. & KALLIO, E.R. (2005) Survival and behaviour of captive-born weasels

(*Mustela nivalis nivalis*) released in nature. *Journal of Zoology* **266**, 37–44.

HENDRY, A.P., SCHOEN, D.J., WOLAK, M.E. & REID, J.M. (2018) The contemporary evolution of

fitness. *Annual Review of Ecology, Evolution, and Systematics* **49**, 457–476.

*HENRIQUEZ, M.C., MACEY, S.K., BAKER, E.E., KELLY, L.B., BETTS, R.L., RUBBO, M.J. &

CLARK, J.A. (2017) Translocated and resident eastern box turtles (*Terrapene c. carolina*) in

New York: movement patterns and habitat use. *Northeastern Naturalist* **24**, 249–266.

HEPPELL, S.S., CROWDER, L.B. & CROUSE, D.T. (1996) Models to evaluate headstarting as a

management tool for long-lived turtles. *Ecological Applications* **6**, 556–565.

HESS, M.A., RABE, C.D., VOGEL, J.L., STEPHENSON, J.J., NELSON, D.D. & NARUM, S.R. (2012)

Supportive breeding boosts natural population abundance with minimal negative impacts on

fitness of a wild population of Chinook salmon. *Molecular Ecology* **21**, 5236–5250.

*HESTER, J.M., PRICE, S.J. & DORCAS, M.E. (2008) Effects of relocation on movements and

home ranges of eastern box turtles. *The Journal of Wildlife Management* **72**, 772–777.

*HIGGINS, J.P.T. & THOMPSON, S.G. (2002) Quantifying heterogeneity in a meta-analysis.

*Statistics in Medicine* **21**, 1539–1558.

*HILL, D. & ROBERTSON, P. (1988) Breeding success of wild and hand-reared ring-necked

pheasants. *The Journal of Wildlife Management* **52**, 446–450.

HIMES, J.G., HARDY, L.M., RUDOLPH, D.C. & BURGDORP, S.J. (2006) Movement patterns and

habitat selection by native and repatriated Louisiana pine snakes (*Pituophis ruthveni*):

implications for conservation. *Herpetological Natural History* **9**, 103–116.

HOFFMANN, M., HILTON-TAYLOR, C., ANGULO, A., BÖHM, M., BROOKS, T.M., BUTCHART,

S.H.M., CARPENTER, K.E., CHANSON, J., COLLEN, B., COX, N.A., DARWALL, W.R.T.,

DULVY, N.K., HARRISON, L.R., KATARIYA, V., POLLOCK, C.M., ET AL. (2010) The impact of

conservation on the status of the world's vertebrates. *Science* **330**, 1503–1509.

*HOLDING, M.L., FRAZIER, J.A., TAYLOR, E.N. & STRAND, C.R. (2012) Experimentally altered

navigational demands induce changes in the cortical forebrain of free-ranging Northern

Pacific rattlesnakes *(Crotalus o. oreganus)*. *Brain, Behavior and Evolution* **79**, 144–154.

HORREO, J.L., VALIENTE, A.G., ARDURA, A., BLANCO, A., GARCIA-VAZQUEZ, C. & GARCIA-

GONZALEZ, E. (2018) Nature versus nurture? Consequences of short captivity in early stages.

*Ecology and Evolution* **8**, 521–529.

HUFBAUER, R.A., SZUCS, M., KASYON, E., YOUNGBERG, C., KOONTZ, M.J., RICHARDS, C., TUFF,

T. & MELBOURNE, B.A. (2015) Three types of rescue can avert extinction in a changing

environment. *Proceedings of the National Academy of Sciences* **112**, 10557–10562.

HUNT, J. & HODGSON, D. (2010) What is fitness, and how do we measure it? In *Evolutionary*

*Behavioral Ecology* (eds D. WESTNEAT & C. FOX), p. New York: Oxford University Press.

HUNTER-AYAD, J., JARVIE, S., GREAVES, G., DIGBY, A., OHLEMÜLLER, R., RECIO, M.R. &

SEDDON, P.J. (2021) Novel conditions in conservation translocations: a conservative-

extrapolative strategic framework. *Frontiers in Conservation Science* **2**, 1–14.

HUNTINGFORD, F.A. (2004) Implications of domestication and rearing conditions for the

behaviour of cultivated fishes. *Journal of Fish Biology* **65,** 122–142.

IMMONEN, E., HÄMÄLÄINEN, A., SCHUETT, W. & TARKA, M. (2018) Evolution of sex-specific

pace-of-life syndromes: genetic architecture and physiological mechanisms. *Behavioral*

*Ecology and Sociobiology* **72**, 60.

*ISLAM, M.Z., SINGH, A., BASHEER, M.P., JUDAS, J. & BOUG, A. (2013) Differences in space use

and habitat selection between captive-bred and wild-born houbara bustards in Saudi Arabia:

results from a long-term reintroduction program. *Journal of Zoology* **289**, 251–261.

ITIS (2021) Retrieved 27 Sep 2021 from the Integrated Taxonomic Information System (ITIS).

*Integrated Taxonomic Information System (ITIS)*. Www.itis.gov [accessed 27 September

2021].

IUCN/SCC (2013) Guidelines for reintroductions and other cnservation translocations. Version

1.0. Gland, Switzerland: IUCN Species Survival Commission.

JACHOWSKI, D.S., GITZEN, R.A., GRENIER, M.B., HOLMES, B. & MILLSPAUGH, J.J. (2011) The

1232  importance of thinking big: large-scale prey conservation drives black-footed ferret

1233  reintroduction success. *Biological Conservation* **144**, 1560–1566.

1234 JENNIONS, M.D. & MØLLER, A.P. (2002) Publication bias in ecology and evolution: An empirical

1235  assessment using the 'trim and fill' method. *Biological Reviews of the Cambridge*

1236  *Philosophical Society* **77**, 211–222.

1237 JENSEN, E.L., EDWARDS, D.L., GARRICK, R.C., MILLER, J.M., GIBBS, J.P., CAYOT, L.J., TAPIA,

1238  W., CACCONE, A. & RUSSELLO, M.A. (2018) Population genomics through time provides

1239  insights into the consequences of decline and rapid demographic recovery through head-

1240  starting in a Galapagos giant tortoise. *Evolutionary Applications*, 1–11.

1241 JOHNSON, M.A., FRANCIS, C.D., MILLER, E.T., DOWNS, C.J. & VITOUSEK, M.N. (2018) Detecting

1242  bias in large-scale comparative analyses: Methods for expanding the scope of hypothesis-

1243  testing with hormonebase. *Integrative and Comparative Biology* **58**, 720–728.

1244 *JOHNSON, O., NEELY, K. & WAPLES, R. (2004) Lopsided fish in the Snake River Basin –

1245  fluctuating asymmetry as a way of assessing impact of hatchery supplementation in Chinook

1246  salmon, *Oncorhynchus tshawytscha*. *Environmental Biology of Fishes* **69**, 379–393.

1247 *JOHNSON, S.L., POWER, J.H., WILSON, D.R. & RAY, J. (2010) A comparison of the survival and

1248  migratory behavior of hatchery-reared and naturally-reared steelhead smolts in the Alsea

1249  River and Estuary, Oregon, using acoustic telemetry. *North American Journal of Fisheries*

1250  *Management* **30**, 55–71.

1251 *JOKIKOKKO, E., KALLIO-NYBERG, I., SALONIEMI, I. & JUTILA, E. (2006) The survival of semi-

1252  wild, wild and hatchery-reared Atlantic salmon smolts of the Simojoki River in the Baltic

1253  Sea. *Journal of Fish Biology* **68**, 430–442.

*JONES, J.M. & WITHAM, J.H. (1990) Post-translocation survival and movements of metropolitan

white-tailed deer. *Wildlife Society Bulletin* **18**, 434–441.

JONES, K.L., GLENN, T.C., LACY, R.C., PIERCE, J.R., UNRUH, N., MIRANDE, C.M. & CHAVEZ-

RAMIREZ, F. (2002) Refining the whooping crane studbook by incorporating microsatellite

DNA and leg-banding analyses. *Conservation Biology* **16**, 789–799.

*JONSSON, B., JONSSON, N. & HANSEN, L.P. (1990) Does juvenile experience affect migration

and spawning of adult Atlantic salmon? *Behavioral Ecology and Sociobiology* **26**, 225–230.

*JONSSON, N., JONSSON, B. & FLEMING, I.A. (1996) Does early growth cause a phenotypically

plastic response in egg production of Atlantic salmon? *Functional Ecology* **10**, 89–96.

JULE, K.R., LEAVER, L.A. & LEA, S.E.G. (2008) The effects of captive experience on

reintroduction survival in carnivores: a review and analysis. *Biological Conservation* **141**,

355–363.

*JUTILA, E., JOKIKOKKO, E., KALLIO-NYBERG, I., SALONIEMI, I. & PASANEN, P. (2003)

Differences in sea migration between wild and reared Atlantic salmon (*Salmo salar L.*) in

the Baltic Sea. *Fisheries Research* **60**, 333–343.

*KÅLÅS, J.A., HEGGBERGET, T.G., BJØRN, P.A. & REITAN, O. (1993) Feeding behaviour and diet

of goosanders *(Mergus merganser)* in relation to salmonid seaward migration. *Aquatic*

*Living Resources* **6**, 31–38.

*KALLIO-NYBERG, I., ROMAKKANIEMI, A., JOKIKOKKO, E., SALONIEMI, I. & JUTILA, E. (2015)

Differences between wild and reared *Salmo salar* stocks of two northern Baltic Sea rivers.

*Fisheries Research* **165**, 85–95.

*KELLISON, G., EGGLESTON, D., TAYLOR, J., BURKE, J. & OSBORNE, J. (2003) Pilot evaluation of

summer flounder stock enhancement potential using experimental ecology. *Marine Ecology*

*Progress Series* **250**, 263–278.

*KING, R.B. & STANFORD, K.M. (2006) Headstarting as a management tool: A case study of the

plains gartersnake. *Herpetologica* **62**, 282–292.

KINGSBURY, B.A. & ATTUM, O. (2009) Conservation strategies: captive rearing, translocation,

and repatriation. In *Snakes: Ecology and Conservation* (eds S.J. MULLIN & SEIGEL), pp.

202–220. Cornell University Press.

KLÜTSCH, C.F.C., MADUNA, S.N., POLIKARPOVA, N., FORFANG, K., ASPHOLM, P.E., NYMAN, T.,

EIKEN, H.G., AMUNDSEN, P.A. & HAGEN, S.B. (2019) Genetic changes caused by restocking

and hydroelectric dams in demographically bottlenecked brown trout in a transnational

subarctic riverine system. *Ecology and Evolution* **9**, 6068–6081.

*KNOX, C.D., JARVIE, S., EASTON, L.J. & MONKS, J.M. (2017) Soft-Release, but not cool winter

temperatures, reduces post-translocation dispersal of jewelled geckos. *Journal of*

*Herpetology* **51**, 490–496.

KNUDSEN, C.M., SHRODER, S.L., BUSACK, C., JOHNSTON, M. V., PEARSONS, T.N. & STROM, C.R.

(2008) Comparison of female reproductive traits and progeny of first-generation hatchery

and wild upper Yakima River spring Chinook salmon. *Transactions of the American*

*Fisheries Society* **137**, 1433–1445.

*KOCK, T.J., PERRY, R.W., POPE, A.C., SERL, J.D., KOHN, M. & LIEDTKE, T.L. (2018) Responses

of hatchery- and natural-origin adult spring Chinook salmon to a trap-and-haul

reintroduction program. *North American Journal of Fisheries Management* **38**, 1004–1016.

*KOŁODZIEJ-SOBOCIŃSKA, M., DEMIASZKIEWICZ, A.W., PYZIEL, A.M. & KOWALCZYK, R. (2018)

Increased parasitic load in captive-released European bison (*Bison bonasus*) has important

implications for reintroduction programs. *EcoHealth* **15**, 467–471.

KORICHEVA, J., JENNIONS, M.D. & LAU, J. (2013) Temporal trends in effect sizes: causes,

detection, and implications. In *Handbook of meta-analysis in ecology and evolution* (eds J.

KORICHEVA, J. GUREVITCH & K. MENGERSEN), pp. 237–254. Princeton University Press.

KOZŁOWSKI, J. (1993) Measuring fitness in life-history studies. *Trends in Ecology and Evolution*

**8**, 84–85.

KRAAIJEVELD-SMIT, F.J.L., GRIFFITHS, R.A., MOORE, R.D. & BEEBEE, T.J.C. (2006) Captive

breeding and the fitness of reintroduced species: a test of the responses to predators in a

threatened amphibian. *Journal of Applied Ecology* **43**, 360–365.

*KRAUS, B.T., MCCALLEN, E.B. & WILLIAMS, R.N. (2017) Evaluating the survival of

translocated adult and captive-reared, juvenile eastern hellbenders (*Cryptobranchus*

*alleganiensis alleganiensis*). *Herpetologica* **73**, 271–276.

*KROCHMAL, A.R., ROTH, T.C. & O'MALLEY, H. (2018) An empirical test of the role of learning

in translocation. *Animal Conservation* **21**, 36–44.

LAIKRE, L., SCHWARTZ, M.K., WAPLES, R.S. & RYMAN, N. (2010) Compromising genetic

diversity in the wild: unmonitored large-scale release of plants and animals. *Trends in*

*Ecology and Evolution* **25**, 520–529.

LAJEUNESSE, M.J., ROSENBERG, M.S. & JENNIONS, M. (2013) Phylogenetic nonindependence and

meta-analysis. In *Handbook of meta-analysis in ecology and evolution* (eds J. KURICHEVA,

J. GUREVITCH & K. MENGERSON), pp. 284–299. Princeton University Press.

LANDE, R. (1998) Anthropogenic, ecological and genetic factors in extinction and conservation.

*Population Ecology* **40**, 259–269.

*LAPIDGE, S.J. (2005) Reintroduction increased vitamin E and condition in captive-bred yellow-

footed rock wallabies *Petrogale xanthopus*. *Oryx* **39**, 56–64.

LARSEN, D.A., HARSTAD, D.L., STROM, C.R., JOHNSTON, M. V., KNUDSEN, C.M., FAST, D.E.,

PEARSONS, T.N. & BECKMAN, B.R. (2013) Early life history variation in hatchery- and

natural-origin spring Chinook salmon in the Yakima River, Washington. *Transactions of the*

*American Fisheries Society* **142**, 540–555.

*LARSSON, S., LINNANSAARI, T., VATANEN, S., SERRANO, I. & HAIKONEN, A. (2011) Feeding of

wild and hatchery reared Atlantic salmon (Salmo salar L.) smolts during downstream

migration. *Environmental Biology of Fishes* **92**, 361–369.

*LARSSON, S., SERRANO, I., ERIKSSON, L.-O. & FLEMING, I. (2012) Effects of muscle lipid

concentration on wild and hatchery brown trout (*Salmo trutta*) smolt migration. *Canadian*

*Journal of Fisheries & Aquatic Sciences* **69**, 1–12.

*LEBATA, MA.J.H.L., LE VAY, L., WALTON, M.E., BIÑAS, J.B., QUINITIO, E.T., RODRIGUEZ, E.M.

& PRIMAVERA, J.H. (2009) Evaluation of hatchery-based enhancement of the mud crab,

*Scylla spp.*, fisheries in mangroves: comparison of species and release strategies. *Marine*

*and Freshwater Research* **60**, 58.

*LEBATA-RAMOS, MA.J.H., DOYOLA-SOLIS, E.F.C., ABROGUEÑA, J.B.R., OGATA, H., SUMBING,

J.G. & SIBONGA, R.C. (2013) Evaluation of post-release behavior, recapture, and growth

rates of hatchery-reared abalone *Haliotis asinina* released in Sagay Marine Reserve,

Philippines. *Reviews in Fisheries Science* **21**, 433–440.

*LEE, J.-H. & PARK, D.-S. (2011) Spatial ecology of translocated and resident Amur ratsnakes

(*Elaphe schrenckii*) in two mountain valleys of South Korea. *Asian Herpetological*

*Research* **2**, 223–229.

LEHRER, E.W., SCHOOLEY, R.L., NEVIS, J.M., KILGOUR, R.J., WOLFF, P.J. & MAGLE, S.B. (2016)

Happily ever after? Fates of translocated nuisance woodchucks in the Chicago metropolitan

area. *Urban Ecosystems* **19**, 1389–1403.

LENORMAND, T. (2002) Gene flow and the limits to natural selection. *Trends in Ecology &*

*Evolution* **17**, 183–189.

LENTH, R. V. (2021) emmeans: Estimated marginal means, aka least-square means.

*LEPEIGNEUL, O., BALLOUARD, J.M., BONNET, X., BECK, E., BARBIER, M., EKORI, A., BUISSON,

E. & CARON, S. (2014) Immediate response to translocation without acclimation from

captivity to the wild in Hermann's tortoise. *European Journal of Wildlife Research* **60**, 897–

907.

*LETTINK, M. (2007) Detectability, movements and apparent lack of homing in *Hoplodactylus*

*maculatus* (Reptilia: Diplodactylidae) following translocation. *New Zealand Journal of*

*Ecology* **31**, 111–116.

LINKLATER, W.L., ADCOCK, K., DU PREEZ, P., SWAISGOOD, R.R., LAW, P.R., KNIGHT, M.H.,

GEDIR, J. V. & KERLEY, G.I.H. (2011) Guidelines for large herbivore translocation

simplified: black rhinoceros case study. *Journal of Applied Ecology* **48**, 493–502.

*LINO, P.G., BENTES, L., ABECASIS, D., SANTOS, M.N.D. & ERZINI, K. (2009) Comparative

behavior of wild and hatchery reared white sea bream (*Diplodus sargus*) released on

artificial reefs off the Algarve (Southern Portugal). In *Tagging and Tracking of Marine*

*Animals with Electronic Devices* (eds J.L. NIELSEN, H. ARRIZABALAGA, N. FRAGOSO, A.

HOBDAY, M. LUTCAVAGE & J. SIBERT), pp. 23–34. Springer Netherlands, Dordrecht.

LUMSDEN, H.G. & DREVER, M.C. (2002) Overview of the trumpeter swan reintroduction

program in Ontario, 1982-2000. *Waterbirds: The International Journal of Waterbird*

*Biology* **25**, 301–312.

LUTZ, M.L., TONKIN, Z., YEN, J.D.L., JOHNSON, G., INGRAM, B.A., SHARLEY, J., LYON, J.,

CHAPPLE, D.G., SUNNUCKS, P. & PAVLOVA, A. (2021) Using multiple sources during

reintroduction of a locally extinct population benefits survival and reproduction of an

endangered freshwater fish. *Evolutionary Applications* **14**, 950–964.

LYNCH, M. (1991) Methods for the analysis of comparative data in evolutionary biology.

*Evolution* **45**, 1065–1080.

LYNCH, M. & O'HELY, M. (2001) Captive breeding and the genetic fitness of natural

populations. *Conservation Genetics* **2**, 363–378.

MANN, K.A., MARTY HOLTGREN, J., AUER, N.A. & OGREN, S.A. (2011) Comparing size,

movement, and habitat selection of wild and streamside-reared lake sturgeon. *North*

*American Journal of Fisheries Management* **31**, 305–314.

MATHEWS, F., ORROS, M., MCLAREN, G., GELLING, M. & FOSTER, R. (2005) Keeping fit on the

ark: assessing the suitability of captive-bred animals for release. *Biological Conservation*

**121**, 569–577.

MCCALLEN, E.B., KRAUS, B.T., BURGMEIER, N.G., FEI, S. & WILLIAMS, R.N. (2018) Movement

and habitat use of eastern hellbenders (*Cryptobranchus alleganiensis alleganiensis*)

following population augmentation. *Herpetologica* **74**, 283–293.

MCCLEERY, R., OLI, M.K., HOSTETLER, J.A., KARMACHARYA, B., GREENE, D., WINCHESTER, C.,

GORE, J., SNECKENBERGER, S., CASTLEBERRY, S.B. & MENGAK, M.T. (2013) Are declines of an endangered mammal predation-driven, and can a captive-breeding and release program aid their recovery? *Journal of Zoology* **291**, 59–68.

*MELNYCHUK, M.C., KORMAN, J., HAUSCH, S., WELCH, D.W., MCCUBBING, D.J.F. & WALTERS, C.J. (2014) Marine survival difference between wild and hatchery-reared steelhead trout determined during early downstream migration. *Canadian Journal of Fisheries and Aquatic Sciences* **71**, 831–846.

MELSTROM, R.T., SALAU, K.R. & SHANAFELT, D.W. (2019) The optimal timing of reintroducing captive populations into the wild. *Ecological Economics* **156**, 174–184.

MENGERSEN, K., SCHMID, C.H., JENNIONS, M.D., & GUREVITCH, J. (2013) Statistical models and approaches to inference. In *Handbook of meta-analysis in ecology and evolution* (eds J. KORICHEVA, J. GUREVITCH & K. MENGERSEN), pp. 89-107. Princeton University Press.

MILLER, K.A., BELL, T.P., & GERMANO, J.M. (2014) Understanding publication bias in reintroduction biology by assessing translocations of New Zealand's herpetofauna. *Conservation Biology* 28, 1045-1056.

*MILLS, D.J., GARDNER, C. & JOHNSON, C.R. (2006) Experimental reseeding of juvenile spiny lobsters (*Jasus edwardsii*): Comparing survival and movement of wild and naïve lobsters at multiple sites. *Aquaculture* **254**, 256–268.

MOHER, D., LIBERATI, A., TETZLAFF, J., ALTMAN, D.G., ALTMAN, D., ANTES, G., ATKINS, D., BARBOUR, V., BARROWMAN, N., BERLIN, J.A., CLARK, J., CLARKE, M., COOK, D., D'AMICO, R., DEEKS, J.J., ET AL. (2009) Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Medicine* **6**.

MOLONY, S.E., DOWDING, C. V., BAKER, P.J., CUTHILL, I.C. & HARRIS, S. (2006) The effect of

1409 translocation and temporary captivity on wildlife rehabilitation success: An experimental

1410 study using European hedgehogs (*Erinaceus europaeus*). *Biological Conservation* **130**,

1411 530–537.

1412 *MONZYK, F.R., JONASSON, B.C., HOFFNAGLE, T.L., KENIRY, P.J., CARMICHAEL, R.W. &

1413 CLEARY, P.J. (2009) Migration characteristics of hatchery and natural spring Chinook

1414 salmon smolts from the Grande Ronde River Basin, Oregon, to Lower Granite Dam on the

1415 Snake River. *Transactions of the American Fisheries Society* **138**, 1093–1108.

1416 *MOORE, M., BEREJIKIAN, B.A. & TEZAK, E.P. (2012) Variation in the early marine survival and

1417 behavior of natural and hatchery-reared Hood Canal steelhead. *PLoS ONE* **7**, e49645.

1418 *MOORE, S.J. & BATTLEY, P.F. (2006) Differences in the digestive organ morphology of captive

1419 and wild brown teal *Anas chlorotis* and implications for releases. *Bird Conservation*

1420 *International* **16**, 253–264.

1421 MORITZ, C. (2002) Strategies to protect biological diversity and the evolutionary processes that

1422 sustain it. *Systematic Biology* **51**, 238–254.

1423 MORRIS, S.D., BROOK, B.W., MOSEBY, K.E., & JOHNSON, C.N. (2021) Factors affecting success

1424 of conservation translocations of terrestrial vertebrates: a global systematic review. *Global*

1425 *Ecology and Conservation* **28,** e01630.

1426 MOSEBY, K.E., HILL, B.M. & LAVERY, T.H. (2014) Tailoring release protocols to individual

1427 species and sites: one size does not fit all. *PLoS ONE* **9**.

1428 MOYES, K., MORGAN, B.J.T., MORRIS, A., MORRIS, S.J., CLUTTON-BROCK, T.H. & COULSON, T.

1429 (2009) Exploring individual quality in a wild population of red deer. *Journal of Animal*

1430 *Ecology* **78**, 406–413.

1431 MULDER, K.P., WALDE, A.D., BOARMAN, W.I., WOODMAN, A.P., LATCH, E.K. & FLEISCHER,

R.C. (2017) No paternal genetic integration in desert tortoises (*Gopherus agassizii*)

following translocation into an existing population. *Biological Conservation* **210**, 318–324.

NAKAGAWA, S. & SANTOS, E.S.A. (2012) Methodological issues and advances in biological

meta-analysis. *Evolutionary Ecology* **26**, 1253–1274.

NAKAGAWA, S. & CUTHILL, I. C. (2007) Effect size, confidence interval and statistical

significant: a practical guide for biologists. *Biological Reviews* **82**: 591–605.

*NASH, D.J. & GRIFFITHS, R.A. (2018) Ranging behaviour of adders (*Vipera berus*) translocated

from a development site. *Herpetological Journal* **28**, 155–159.

NASON, S.E., LLOYD, N., KELLY, C.D., BRICHIERI-COLOMBI, T.A., DALRYMPLE, S.E. &

MOEHRENSCHLAGER, A. (2021) Maximizing the effectiveness of qualitative systematic

reviews: A case study on terrestrial arthropod conservation translocations. *Biological*

*Conservation* **254**, 108948.

NAZAR, F.N. & MARIN, R.H. (2011) Chronic stress and environmental enrichment as opposite

factors affecting the immune response in Japanese quail (*Coturnix coturnix japonica*). *Stress*

**14**, 166–173.

*NELSON, T.C., ROSENAU, M.L. & JOHNSTON, N.T. (2005) Behavior and survival of wild and

hatchery-origin winter steelhead spawners caught and released in a recreational fishery.

*North American Journal of Fisheries Management* **25**, 931–943.

*NEUMAN, K., STENZEL, L., WARRINER, J., PAGE, G., ERBES, J., EYSTER, C., MILLER, E. &

HENKEL, L. (2013) Success of captive-rearing for a threatened shorebird. *Endangered*

*Species Research* **22**, 85–94.

NEWMARK, W.D., JENKINS, C.N., PIMM, S.L., MCNEALLY, P.B. & HALLEY, J.M. (2017) Targeted

habitat restoration can reduce extinction rates in fragmented forests. *Proceedings of the*

*National Academy of Sciences of the United States of America* **114**, 9635–9640.

*Nicoll, M.A.C., Jones, C.G. & Norris, K. (2004) Comparison of survival rates of captive-reared and wild-bred Mauritius kestrels (*Falco punctatus*) in a re-introduced population. *Biological Conservation* **118**, 539–548.

Noble, D.W.A., Lagisz, M., O'dea, R.E. & Nakagawa, S. (2017) Nonindependence and sensitivity analyses in ecological and evolutionary meta-analyses. *Molecular Ecology* **26**, 2410–2425.

van Noordwijk, A.J. & de Jong, G. (1986) Acquisition and allocation of resources: their influence on variation in life history tactics. *The American Naturalist* **128**, 137–142.

*Norris, T.A., Littnan, C.L. & Gulland, F.M.D. (2011) Evaluation of the captive care and post-release behavior and survival of seven juvenile female Hawaiian monk seals (*Monachus schauinslandi*). *Aquatic Mammals* **37**, 342–353.

*Nussear, K.E., Tracy, C.R., Medica, P.A., Wilson, D.S., Marlow, R.W. & Corn, P.S. (2012) Translocation as a conservation tool for Agassiz's desert tortoises: survivorship, reproduction, and movements. *The Journal of Wildlife Management* **76**, 1341–1353.

*Nyqvist, D., McCormick, S.D., Greenberg, L., Ardren, W.R., Bergman, E., Calles, O. & Castro-Santos, T. (2017) Downstream migration and multiple dam passage by Atlantic salmon smolts. *North American Journal of Fisheries Management* **37**, 816–828.

O'Dea, R.E., Lagisz, M., Jennions, M.D., Koricheva, J., Noble, D.W.A., Parker, T.H., Gurevitch, J., Page, M.J., Stewart, G., Moher, D. & Nakagawa, S. (2021) Preferred reporting items for systematic reviews and meta-analyses in ecology and evolutionary biology: a PRISMA extension. *Biological Reviews* **96**, 1695–1722.

*ØKLAND, F., HEGGBERGET, T.G. & JONSSON, B. (1995) Migratory behaviour of wild and

farmed Atlantic salmon (*Salmo salar*) during spawning. *Journal of Fish Biology* **46**, 1–7.

OLIVIER, J., & BELL, M. L. (2013) Effect sizes for 2 x 2 contingency tables. PSoS ONE **8**, e5877.

*OSTERBACK, A.-M.K., FRECHETTE, D.M., HAYES, S.A., BOND, M.H., SHAFFER, S.A. & MOORE,

J.W. (2014) Linking individual size and wild and hatchery ancestry to survival and

predation risk of threatened steelhead (*Oncorhynchus mykiss*). *Canadian Journal of*

*Fisheries and Aquatic Sciences* **71**, 1877–1887.

*OSTERMANN, S.D., DEFORGE, J.R. & EDGE, W.D. (2001) Captive breeding and reintroduction

evaluation criteria: a case study of peninsular bighorn sheep. *Conservation Biology* **15**, 749–

760.

*PARISH, D.M.B. & SOTHERTON, N.W. (2007) The fate of released captive-reared grey partridges

*Perdix perdix*: implications for reintroduction programmes. *Wildlife Biology* **13**, 140–149.

PARKER, G. (1979) Sexual selection and sexual conflict. In *Sexual Selection and Reproductive*

*Competition in Insects* (ed M. BLUM), pp. 123–166. New York Academic Press.

*PECK, A.J., HARRIS, J.L., FARRIS, J.L. & CHRISTIAN, A.D. (2014) Survival and horizontal

movement of the freshwater mussel *Potamilus capax* (Green, 1832) following relocation

within a Mississippi Delta stream system. *American Midland Naturalist* **172**, 76–90.

*PEDRONO, M. & SAROVY, A. (2000) Trial release of the world's rarest tortoise *Geochelone*

*yniphora* in Madagascar. *Biological Conservation* **95**, 333–342.

PEKKALA, N., KOTIAHO, J.S. & PUURTINEN, M. (2011) Laboratory relationships between adult

lifetime reproductive success and fitness surrogates in a *Drosophila littoralis* population.

*PLoS ONE* **6**, e24560.

PÉREZ, I., ANADÓN, J.D., DÍAZ, M., NICOLA, G.G., TELLA, J.L. & GIMÉNEZ, A. (2012) What is

1500 wrong with current translocations? A review and a decision-making proposal. *Frontiers in*

1501  *Ecology and the Environment* **10**, 494–501.

1502 *PÉREZ-BUITRAGO, N., GARCÍA, M., SABAT, A., DELGADO, J., ÁLVAREZ, A., MCMILLAN, O. &

1503  FUNK, S. (2008) Do headstart programs work? Survival and body condition in headstarted

1504  Mona Island iguanas *Cyclura cornuta stejnegeri*. *Endangered Species Research* **6**, 55–65.

1505 *PIGGINS, D.J. & MILLS, C.P.R. (1985) Comparative aspects of the biology of naturally produced

1506  and hatchery-reared Atlantic salmon smolts (*Salmo salar L.*). *Aquaculture* **45**, 321–333.

1507 *PILLE, F., CARON, S., BONNET, X., DELEUZE, S., BUSSON, D., ETIEN, T., GIRARD, F. &

1508  BALLOUARD, J.-M. (2018) Settlement pattern of tortoises translocated into the wild: a key to

1509  evaluate population reinforcement success. *Biodiversity and Conservation* **27**, 437–457.

1510 PIMM, S.L., JENKINS, C.N., ABELL, R., BROOKS, T.M., GITTLEMAN, J.L., JOPPA, L.N., RAVEN,

1511  P.H., ROBERTS, C.M. & SEXTON, J.O. (2014) The biodiversity of species and their rates of

1512  extinction, distribution, and protection. *Science* **344**, 987.

1513 PINTER, K., EPIFANIO, J. & UNFER, G. (2019) Release of hatchery-reared brown trout (*Salmo*

1514  *trutta*) as a threat to wild populations? A case study from Austria. *Fisheries Research* **219**,

1515  105296.

1516 *PLUMMER, M.V. & MILLS, N.E. (2000) Spatial ecology and survivorship of resident and

1517  translocated hognose snakes (*Heterodon platirhinos*). *Journal of Herpetology* **34**, 565–575.

1518 *POIRIER, M.-A. & FESTA-BIANCHET, M. (2018) Social integration and acclimation of

1519  translocated bighorn sheep (*Ovis canadensis*). *Biological Conservation* **218**, 1–9.

1520 POISOT, T. (2011) The digitize package: extracting numerical data from scatterplots. *The R*

1521  *Journal* **3**, 26–27.

1522 POO, S., BOGISICH, A., MACK, M., LYNN, B.K. & DEVAN-SONG, A. (2022) Post-release

comparisons of amphibian growth reveal challenges with sperm cryopreservation as a
conservation tool. *Conservation Science and Practice* **4**, 1–9.

*POULIN, R.G., TODD, L.D., WELLICOME, T.I. & BRIGHAM, R.M. (2006) Assessing the feasibility
of release techniques for captive-bred burrowing owls. *Journal of Raptor Research* **40**, 142–
150.

*POWELL, M.S., HARDY, R.W., FLAGG, T.A. & KLINE, P.A. (2010) Proximate composition and
fatty acid differences in hatchery-reared and wild Snake River Sockeye salmon
overwintering in nursery lakes. *North American Journal of Fisheries Management* **30**, 530–
537.

, A.S. & KNIGHT, T.M. (2001) Effectiveness in conservation practice: pointers from medicine
and public health. *Conservation Biology* **15**, 50–54.

QUINN, T.P., SEAMONS, T.R. & JOHNSON, S.P. (2012) Stable isotopes of carbon and nitrogen
indicate differences in marine ecology between wild and hatchery-produced steelhead.
*Transactions of the American Fisheries Society* **141**, 526–532.

RANTANEN, E.M., BUNER, F., RIORDAN, P., SOTHERTON, N. & MACDONALD, D.W. (2010)
Vigilance, time budgets and predation risk in reintroduced captive-bred grey partridges
*Perdix perdix*. *Applied Animal Behaviour Science* **127**, 43–50.

*RAY, M., STONER, A. & O'CONNELL, S. (1994) Size-specific predation of juvenile queen conch
*Strombus gigas*: implications for stock enhancement. *Aquaculture* **128**, 79–88.

READING, R.P., CLARK, T.W., & GRIFFITH, B. (1997) The influence of valuational and
organization consideration on the succes of rare species translocations. *Biological
Conservation* **79**, 217-225.

READING, R.P., MILLER, B. & SHEPHERDSON, D. (2013) The value of enrichment to

reintroduction success. *Zoo Biology* **32**, 332–341.

REID, J.M., ARCESE, P., NIETLISBACH, P., WOLAK, M.E., MUFF, S., DICKEL, L., & KELLER, L.F.

(2021) Immigration counter-acts local micro-evolution of a major fitness component:

migration-selection balance in free-living song sparrows. *Evolution Letters* **5**, 48-60.

*REINERT, H.K. & RUPERT, R.R. (1999) Impacts of translocation on behavior and survival of

timber rattlesnakes, *Crotalus horridus*. *Journal of Herpetology* **33**, 45–61.

RESENDE, P.S., VIANA-JUNIOR, A.B., YOUNG, R.J. & AZEVEDO, C.S. (2021) What is better for

animal conservation translocation programmes: soft- or hard-release? A phylogenetic meta-

analytical approach. *Journal of Applied Ecology* **58**, 1122–1132.

*RITTENHOUSE, C.D., MILLSPAUGH, J.J., HUBBARD, M.W. & SHERIFF, S.L. (2007) Movements of

translocated and resident three-toed box turtles. *Journal of Herpetology* **41**, 115–121.

*ROCHE, E.A., CUTHBERT, F.J. & ARNOLD, T.W. (2008) Relative fitness of wild and captive-

reared piping plovers: does egg salvage contribute to recovery of the endangered Great

Lakes population? *Biological Conservation* **141**, 3079–3088.

ROE, J.H., FRANK, M.R. & KINGSBURY, B.A. (2015) Experimental evaluation of captive-rearing

practices to improve success of snake reintroductions. *Herpetological Conservation and*

*Biology* **10**, 711–722.

ROFF, D.A. (2002) *Life History Evolution*. Sinauer Associates, Inc. Sunderland, MA, USA.

ROSENBERG, M.S., ROTHSTEIN, H.R. & GUREVITCH, J. (2013) Effect sizes: Conventional choices

and calculations. In *Handbook of meta-analysis in ecology and evolution* (eds J.

KURICHEVA, J. GUREVITCH & K. MENGERSEN), pp. 61–71. Princeton University Press.

ROSENTHAL, R. & RUBIN, D.B. (1982) Comparing effect sizes of independent studies.

*Psychological Bulletin* **92**, 500–504.

RUCKSTUHL, K.E. & NEUHAUS, P. (2005) *Sexual Segregation in Vertebrates: Ecology of the two*

*sexes*. Cambridge University Press.

RUMMEL, L., MARTÍNEZ-ABRAÍN, A., MAYOL, J., RUIZ-OLMO, J., MAÑAS, F., JIMÉNEZ, J.,

GÓMEZ, J.A. & ORO, D. (2016) Use of wild-caught individuals as a key factor for success in

vertebrate translocations. *Animal Biodiversity and Conservation* **39**, 207–219.

*RUTH, T.K., LOGAN, K.A., SWEANOR, L.L., HORNOCKER, M.G. & TEMPLE, L.J. (1998)

Evaluating Cougar Translocation in New Mexico. *The Journal of Wildlife Management* **62**,

1264–1275.

*RYAN, B.A., SMITH, S.G., BUTZERIN, J.A.M. & FERGUSON, J.W. (2003) Relative vulnerability to

avian predation of juvenile salmonids tagged with passive integrated transponders in the

Columbia River estuary, 1998–2000. *Transactions of the American Fisheries Society* **132**,

275–288.

RYMAN, N. & LAIKRE, L. (1991) Effects of supportive breeding on the genetically effective

population size. *Conservation Biology* **5**, 325–329.

SAH, P., NUSSEAR, K.E., ESQUE, T.C., AIELLO, C.M., HUDSON, P.J. & BANSAL, S. (2016)

Inferring social structure and its drivers from refuge use in the desert tortoise, a relatively

solitary species. *Behavioral Ecology and Sociobiology* **70**, 1277–1289.

*SALONIEMI, I., JOKIKOKKO, E., KALLIO-NYBERG, I., JUTILA, E. & PASANEN, P. (2004) Survival

of reared and wild Atlantic salmon smolts: size matters more in bad years. *ICES Journal of

Marine Science* **61**, 782–787.

*SANTOS, T., PÉREZ-TRIS, J., CARBONELL, R., TELLERÍA, J.L. & DÍAZ, J.A. (2009) Monitoring the

performance of wild-born and introduced lizards in a fragmented landscape: implications for

ex situ conservation programmes. *Biological Conservation* **142**, 2923–2930.

SATAKE, A. & ARAKI, H. (2012) Stocking of captive-bred fish can cause long-term population

decline and gene pool replacement: predictions from a population dynamics model

incorporating density-dependent mortality. *Theoretical Ecology* **5**, 283–296.

SCHEELE, B.C., HOLLANDERS, M., HOFFMANN, E.P., NEWELL, D.A., LINDENMAYER, D.B.,

MCFADDEN, M., GILBERT, D.J. & GROGAN, L.F. (2021) Conservation translocations for

amphibian species threatened by chytrid fungus: a review, conceptual framework, and

recommendations. *Conservation Science and Practice* **3**, 1–15.

*SCILLITANI, L., STURARO, E., MENZANO, A., ROSSI, L., VIALE, C. & RAMANZIN, M. (2012) Post-

release spatial and social behaviour of translocated male Alpine ibexes (*Capra ibex ibex*) in

the eastern Italian Alps. *European Journal of Wildlife Research* **58**, 461–472.

SEDDON, P.J., ARMSTRONG, D.P. & MALONEY, R.F. (2007) Developing the science of

reintroduction biology. *Conservation Biology* **21**, 303–312.

SEDDON, P.J., GRIFFITHS, C.J., SOORAE, P.S. & ARMSTRONG, D.P. (2014) Reversing defaunation:

Restoring species in a changing world. *Science* **345**, 406–411.

SENIOR, A.M., GRUEBER, C.E., KAMIYA, T., LAGISZ, M., O'DWYER, K., SANTOS, E.S.A. &

NAKAGAWA, S. (2016) Heterogeneity in ecological and evolutionary meta- analyses: its

magnitude and implications. *Ecology* **97**, 3293–3299.

*SERRANO, I., LARSSON, S. & ERIKSSON, L.-O. (2009) Migration performance of wild and

hatchery sea trout (*Salmo trutta L.*) smolts—implications for compensatory hatchery

programs. *Fisheries Research* **99**, 210–215.

SHAW, R.G. (2019) From the past to the future: considering the value and limits of evolutionary

Prediction. *The American Naturalist* **193**, 1–10.

SHIER, D.M. & OWINGS, D.H. (2007) Effects of social learning on predator training and

postrelease survival in juvenile black-tailed prairie dogs, *Cynomys ludovicianus*. *Animal*

*Behaviour* **73**, 567–577.

*SINN, D.L., CAWTHEN, L., JONES, S.M., PUKK, C. & JONES, M.E. (2014) Boldness towards

novelty and translocation success in captive-raised, orphaned Tasmanian devils. *Zoo*

*Biology* **33**, 36–48.

*SKAALA, O., JØRSTAD, K. & BORGSTRØM, R. (2011) Genetic impact on two wild brown trout

(*Salmo trutta*) populations after release of non-indigenous hatchery spawners. *Canadian*

*Journal of Fisheries and Aquatic Sciences* **53**, 2027–2035.

SKIKNE, S.A., BORKER, A.L., TERRILL, R.S. & ZAVALETA, E. (2020) Predictors of past avian

translocation outcomes inform feasibility of future efforts under climate change. *Biological*

*Conservation* **247**, 108597.

*SLAUGH, B.T., FLINDERS, J.T., ROBERSON, J.A. & JOHNSTON, N.P. (1992) Effect of rearing

method on chukar survival. *The Great Basin Naturalist* **52**, 25–28.

SNYDER, N.F.R., DERRICKSON, S.R., BEISSINGER, S.R., WILEY, J.W., SMITH, T.B., TOONE, W.D.

& MILLER, B. (1996) Limitations of captive breeding in endangered species recovery.

*Conservation Biology* **10**, 338–348.

STAMPS, J.A. & SWAISGOOD, R.R. (2007) Someplace like home: experience, habitat selection

and conservation biology. *Applied Animal Behaviour Science* **102**, 392–409.

*STARK, E.J., VIDERGAR, D.T., KOZFKAY, C.C. & KLINE, P.A. (2018) Egg viability and egg-to-

fry survival of captive-reared Chinook salmon released to spawn naturally. *Transactions of*

*the American Fisheries Society* **147**, 128–138.

*STEWART, D.C. (2015) Ranching to the rod: an evaluation of adult returns from hatchery-reared

Atlantic salmon smolts released in Scottish rivers. *Scottish Marine and Freshwater Science*

**6,** 1-13.

STEWART, G. (2010) Meta-analysis in applied ecology. *Biology Letters* **6**, 78–81.

*STONER, A. & DAVIS, M. (1994) Experimental outplanting of juvenile queen conch, Strombus

gigas: comparison of wild and hatchery-reared stocks. *Fishery Bulletin- National Oceanic*

*and Atmospheric Administration* **92**, 390–411.

STRINGWELL, R., LOCK, A., STUTCHBURY, C.J., BAGGETT, E., TAYLOR, J., GOUGH, P.J. &

LEANIZ, C.G. DE (2014) Maladaptation and phenotypic mismatch in hatchery-reared

Atlantic salmon *Salmo salar* released in the wild. *Journal of Fish Biology* **85**, 1927–1945.

STUPARYK, B., HORN, C.J., KARABATSOS, S. & ARTEAGA, J. (2018) A meta-analysis of animal

survival following translocation: comparisons between conflict and conservation efforts.

*Canadian Wildlife Biology and Managment* **7**, 3–17.

*STURDEVANT, M.V., FERGUSSON, E., HILLGRUBER, N., REESE, C., ORSI, J., FOCHT, R.,

WERTHEIMER, A. & SMOKER, B. (2012) Lack of trophic competition among wild and

hatchery juvenile chum salmon during early marine residence in Taku Inlet, Southeast

Alaska. *Environmental Biology of Fishes* **94**, 101–116.

*SULAK, K.J., RANDALL, M.T. & CLUGSTON, J.P. (2014) Survival of hatchery Gulf sturgeon

(*Acipenser oxyrinchus desotoi* Mitchill, 1815) in the Suwannee River, Florida: a 19-year

evaluation. *Journal of Applied Ichthyology* **30**, 1428–1440.

SULLIVAN, B.K., NOWAK, E.M. & KWIATKOWSKI, M.A. (2015) Problems with mitigation

translocation of herpetofauna. *Conservation Biology* **29**, 12–18.

SUTHERLAND, W.J., PULLIN, A.S., DOLMAN, P.M. & KNIGHT, T.M. (2004) The need for

evidence-based conservation. *Trends in Ecology and Evolution* **19**, 305–308.

TARKA, M., GUENTHER, A., NIEMELÄ, P.T., NAKAGAWA, S. & NOBLE, D.W.A. (2018) Sex

differences in life history, behavior, and physiology along a slow-fast continuum: a meta-

analysis. *Behavioral Ecology and Sociobiology* **72**. Behavioral Ecology and Sociobiology.

*TAYLOR, M.D., LAFFAN, S.W., FAIRFAX, A.V. & PAYNE, N.L. (2017) Finding their way in the

world: Using acoustic telemetry to evaluate relative movement patterns of hatchery-reared

fish in the period following release. *Fisheries Research* **186**, 538–543.

TEIXEIRA, C.P., DE AZEVEDO, C.S., MENDL, M., CIPRESTE, C.F. & YOUNG, R.J. (2007) Revisiting

translocation and reintroduction programmes: the importance of considering stress. *Animal*

*Behaviour* **73**, 1–13.

*TEMSIRIPONG, Y., WOODWARD, A.R., ROSS, J.P., KUBILIS, P.S. & PERCIVAL, H.F. (2006)

Survival and growth of american alligator (*Alligator mississippiensis*) hatchlings after

artificial incubation and repatriation. *Journal of Herpetology* **40**, 415–423.

*TERHUNE, T.M., SISSON, D.C. & STRIBLING, H.L. (2006) The efficacy of relocating wild

northern bobwhites prior to breeding season. *The Journal of Wildlife Management* **70**, 914–

921.

TETZLAFF, S.J., SPERRY, J.H. & DEGREGORIO, B.A. (2018) Captive-reared juvenile box turtles

innately prefer naturalistic habitat: Implications for translocation. *Applied Animal Behaviour*

*Science* **204**, 128–133.

TETZLAFF, S.J., SPERRY, J.H. & DEGREGORIO, B.A. (2019a) Effects of antipredator training,

environmental enrichment, and soft release on wildlife translocations: a review and meta-

analysis. *Biological Conservation* **236**, 324–331.

TETZLAFF, S.J., SPERRY, J.H., KINGSBURY, B.A. & DEGREGORIO, B.A. (2019b) Captive-rearing

duration may be more important than environmental enrichment for enhancing turtle head-

starting success. *Global Ecology and Conservation* **20**, e00797.

THODAY, J.M. (1953) Components of fitness. *Symposium of the Society for Experimental Biology*

**8**, 96–113.

*THOMPSON, B.C., PORAK, W.F., LEONE, E.H. & ALLEN, M.S. (2016) Using radiotelemetry to

compare the initial behavior and mortality of hatchery-reared and wild juvenile Florida bass.

*Transactions of the American Fisheries Society* **145**, 374–385.

*THORSTAD, E.B., HEGGBERGET, T.G. & ØKLAND, F. (1998) Migratory behaviour of adult wild

and escaped farmed Atlantic salmon, *Salmo salar L.*, before, during and after spawning in a

Norwegian river. *Aquaculture Research* **29**, 419–428.

*THORSTAD, E.B., ØKLAND, F., FINSTAD, B., SIVERTSGÅRD, R., PLANTALECH, N., BJØRN, P.A. &

MCKINLEY, R.S. (2007) Fjord migration and survival of wild and hatchery-reared Atlantic

salmon and wild brown trout post-smolts. *Hydrobiologia* **582**, 99–107.

*TRESTED, D.G., WARE, K., BAKAL, R. & ISELY, J.J. (2011) Microhabitat use and seasonal

movements of hatchery-reared and wild shortnose sturgeon in the Savannah River, South

Carolina – Georgia. *Journal of Applied Ichthyology* **27**, 454–461.

TRIVERS, R.L. (1972) Parental investment and sexual selection. In *Sexual Selection and the

Descent of Man, 1871-1971* pp. 136–179. Aldine, Chicago, IL.

*TSAKIRIS, E.T., RANDKLEV, C.R., BLAIR, A., FISHER, M. & CONWAY, K.W. (2017) Effects of

translocation on survival and growth of freshwater mussels within a West Gulf Coastal Plain

river system. *Aquatic Conservation: Marine and Freshwater Ecosystems* **27**, 1240–1250.

TUBERVILLE, T.D., CLARK, E.E., BUHLMANN, K.A. & GIBBONS, J.W. (2005) Translocation as a

conservation tool: Site fidelity and movement of repatriated gopher tortoises (*Gopherus*

1705  *polyphemus). Animal Conservation* **8**, 349–358.

1706  *Turek, J., Horký, P., Velíšek, J., Slavík, O., Hanák, R. & Randák, T. (2010a) Recapture

1707  rate and growth of hatchery-reared brown trout (*Salmo trutta v. fario, L.*) in Blanice River

1708  and the effect of stocking on wild brown trout and grayling (*Thymallus thymallus, L.*).

1709  *Journal of Applied Ichthyology* **26**, 881–885.

1710  *Turek, J., Randák, T., Horký, P., Z☐Lábek, V., Velíšek, J., Slavík, O. & Hanák, R.

1711  (2010b) Post-release growth and dispersal of pond and hatchery-reared European grayling

1712  Thymallus thymallus compared with their wild conspecifics in a small stream. *Journal of

1713  Fish Biology* **76**, 684–693.

1714  Turlure, C., Radchuk, V., Baguette, M., Meijrink, M., van den Burg, A., De Vries, M.W.

1715  & van Duinen, G.J. (2013) Plant quality and local adaptation undermine relocation in a bog

1716  specialist butterfly. *Ecology and Evolution* **3**, 244–254.

1717  *Tweed, E.J., Foster, J.T., Woodworth, B.L., Monahan, W.B., Kellerman, J.L. &

1718  Lieberman, A. (2006) Breeding biology and success of a reintroduced population of the

1719  critically endangered Puaiohi (*Myadestes Palmeri*). *The Auk* **123**, 753–763.

1720  *Unger, C. & Klaus, Siegfried (2009) Lebenserwartung und Verlustursachen umgesiedelter

1721  Auerhühner *Tetrao urogallus* in Thüringen. *Ornithologischer Anzeiger* **48**, 50–55.

1722  *Urke, H.A., Kristensen, T., Ulvund, J.B. & Alfredsen, J.A. (2013) Riverine and fjord

1723  migration of wild and hatchery-reared Atlantic salmon smolts. *Fisheries Management and

1724  Ecology* **20**, 544–552.

1725  Vanderwerf, E.A., Crampton, L.H., Diegmann, J.S., Atkinson, C.T. & Leonard, D.L.

1726  (2014) Survival estimates of wild and captive-bred released Puaiohi, an endangered

1727  Hawaiian thrush. *The Condor: Ornithological Applications* **116**, 609–618.

VIECHTBAUER, W. (2010) Conducting meta-analyses in R with the metafor. *Journal of Statistical*

*Software* **36**, 1–48.

VIECHTBAUER, W. & CHEUNG, M.W.-L. (2010) Outlier and influence diagnostics for meta-

analysis. *Research Synthesis Methods* **1**, 112–125.

WAKE, D.B. & VREDENBURG, V.T. (2008) Are we in the midst of the sixth mass extinction? A

view from the world of amphibians. *Proceedings of the National Academy of Sciences of the*

*United States of America* **105**, 11466–11473.

WAPLES, R.S., HINDAR, K., KARLSSON, S. & HARD, J.J. (2016) Evaluating the Ryman–Laikre

effect for marine stock enhancement and aquaculture. *Current Zoology* **62**, 617–627.

*WEIGEL, D.E., CONNOLLY, P.J. & POWELL, M.S. (2014) Fluvial rainbow trout contribute to the

colonization of steelhead (*Oncorhynchus mykiss*) in a small stream. *Environmental Biology*

*of Fishes* **97**, 1149–1159.

*WEISE, F.J., LEMERIS, J., STRATFORD, K.J., VAN VUUREN, R.J., MUNRO, S.J., CRAWFORD, S.J.,

MARKER, L.L. & STEIN, A.B. (2015) A home away from home: insights from successful

leopard (*Panthera pardus*) translocations. *Biodiversity and Conservation* **24**, 1755–1774.

*WELCH, D.W., WARD, B.R. & BATTEN, S.D. (2004) Early ocean survival and marine

movements of hatchery and wild steelhead trout (*Oncorhynchus mykiss*) determined by an

acoustic array: Queen Charlotte Strait, British Columbia. *Deep Sea Research Part II:*

*Topical Studies in Oceanography* **51**, 897–909.

*WILLIAMSON, K.S., MURDOCH, A.R., PEARSONS, T.N., WARD, E.J. & FORD, M.J. (2010) Factors

influencing the relative fitness of hatchery and wild spring Chinook salmon (*Oncorhynchus*

*tshawytscha*) in the Wenatchee River, Washington, USA. *Canadian Journal of Fisheries*

*and Aquatic Sciences* **67**, 1840–1851.

WILSON, A.J. & NUSSEY, D.H. (2010) What is individual quality? An evolutionary perspective.

*Trends in Ecology and Evolution* **25**, 207–214.

*WISELY, S.M., HOWARD, J., WILLIAMS, S.A., BAIN, O., SANTYMIRE, R.M., BARDSLEY, K.D. &

WILLIAMS, E.S. (2008) An unidentified filiarial species and its impact on fitness in wild

population of the black-footed ferret (*Mustela nigripes*). *Journal of Wildlife Diseases* **44**,

53–64.

WOLF, C.M., GRIFFITH, B., REED, C. & TEMPLE, S.A. (1996) Avian and mammalian

translocations: update and reanalysis of 1987 survey data. *Conservation Biology* **10**, 1142–

1154.

*WOLFE, A.K., FLEMING, P.A., BATEMAN, P.W., WOLFE, A.K., FLEMING, P.A. & BATEMAN,

P.W. (2018) Impacts of translocation on a large urban-adapted venomous snake. *Wildlife

Research* **45**, 316–324.

*WOOD, C., WELCH, D., GODBOUT, L. & CAMERON, J. (2012) Marine migratory behavior of

hatchery-reared anadromous and wild non-anadromous sockeye salmon revealed by acoustic

tags. *American Fisheries Society Symposium* **76**, 289–311.

WWF/ZSL (2020) The Living Planet Index database. Www.livingplanetindex.org.

ZHANG, Z., GAO, L. & ZHANG, X. (2022) Environmental enrichment increases aquatic animal

welfare: A systematic review and meta-analysis. *Reviews in Aquaculture* **14**, 1120–1135.

ZHU, J., GOSNELL, J.S., AKALLAL, L. & GOLTSMAN, M. (2022) Fear changes traits and increases

survival: a meta☐analysis evaluating the efficacy of antipredator training in captive☐rearing

programs. *Restoration Ecology* **31**, 1–11.

 **X. SUPPORTING INFORMATION**

Additional supporting information may be found online in the Supporting Information section at

the end of the article.

**Appendix S1:** PRISMA EcoEvo checklist.

**Appendix S2:** Supplemental tables summarizing full-text screening outcomes, meta-

analysis model summary tables, and heterogeneity statistics.

**Appendix S3:** Supplemental figures.

1779

**Figure Captions**

**Figure 1.** A flowchart describing the dataset accession and screening processes as prescribed by
Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines
(Moher *et al.*, 2009; O'Dea *et al.*, 2021). †We supplemented the query-returned papers with 28
additional papers synthesized by Stuparyk *et al.*, (2018) focused on relative post-release
performance of organisms translocated to alleviate human-animal conflicts.
1786
**Figure 2.** Marginal posterior distributions of log odds ratios (ln odds ratio) across taxonomic
classes. Values greater than zero (vertical dashed line) indicate that translocated organisms out-
performed wild-resident conspecifics, and negative values indicate translocated organisms under-
performed. Vertical lines within each density plot indicate the 95% highest posterior density
credible intervals (95% CrI). The parenthetical statements indicate the number of studies and the
number of individual comparisons comprising each density plot, respectively.
1793
**Figure 3.** Marginal posterior distributions of log odds ratios (ln odds ratio) across enriched vs.
unenriched translocated cohorts and taxonomic classes. Values greater than zero (vertical dashed
line) indicate that translocated organisms out-performed wild-resident conspecifics, and negative
values indicate translocated organisms under-performed. Blue density plots represent
comparisons in which translocated organisms received some form of pre-release enrichment,
whereas tan density plots represent comparisons where translocated organisms were unenriched.
Vertical lines within each density plot indicate the 95% highest posterior density credible
intervals (95% CrI). The parenthetical statements indicate the number of studies and the total
number of comparisons between enriched/unenriched and wild cohorts organized by taxonomic
class.
1804
1805    **Figure 4.** Time-lag bias. The relationship between effect size and publication year. Effect sizes
1806    are marginal posterior distributions, with 95% credible intervals indicated by black vertical lines.
1807    The dashed vertical line indicates an effect size of zero.
1808

1809 **Figure 5.** Egger's regression. Funnel plots describing patterns of publication bias for the full
1810 dataset. (A) Relationship between effect size (log odds ratio) and precision. The solid line
1811 indicates the meta-analytic mean. (B) Relationship between the meta-analytic residuals and
1812 precision. The solid vertical line indicates zero.