Insm1 regulates the development of mTECs and immune tolerance

Wehuai Tao[1,2,3], Yiqiu Wei[1,2,3], Zhihuan Ye[1,2,3], Jianxue Wang[1], Weixin Yang[1],

Guoxing Yu[1,2,3], Jieyi Xiong[*4,5], Shiqi Jia[*1,2,3]

1 The First Affiliated Hospital of Jinan University, Guangzhou, China

2 The Guangdong-Hong Kong-Macao Joint University Laboratory of Metabolic

and Molecular Medicine, Jinan University, Guangzhou, China

3 The Institute of Clinical Medicine, Jinan University, Guangzhou, China

4 Max Delbrück Center for Molecular Medicine in the Helmholtz Association,

Berlin, Germany

5 VIB-KU Leuven Center for Cancer Biology, Leuven, Belgium

#These authors contribute equally to this work

* Corresponding author:
Shiqi Jia, Phone: +86-20-3868 0563, Email: shiqijia@jnu.edu.cn
ORCID for Shiqi Jia: 0000-0001-8595-9314

Jieyi Xiong, Phone: +32 478 11 49 46, Email: jieyi.xiong@kuleuven.be
ORCID for Jieyi Xiong: 0000-0002-4302-3515

Abstract

The *Insm1* gene encodes a zinc finger protein with known functions in neuroendocrine cells and neurons. Here we characterized the expression and function of *Insm1* in medullary thymic epithelial cells (mTECs). *Insm1* is co-expressed with Aire in majority of Insm1- or Aire- positive cells, while a few Insm1-positive cells did not express Aire. Mutation of *Insm1* impair the expression of *Aire* and the generation of normal numbers of Aire-expressing mTECs during development. We detected downregulation of genes that expressed specifically in Aire-expressing mTEC and mimetic cells in *Insm1* mutant mTECs. Conversely, when *Insm1* was overexpressed in thymic epithelial cells *in vivo*, the size of the mTECs compartment was enlarged and the expression of *Aire* and genes expressed specifically in the neuroendocrine mimetic cells were increased. Mechanistically, Insm1 bound DNA in mTECs and the majority of the Insm1 binding sites were co-occupied by Aire. These Insm1 binding sites were enriched on super-enhancer regions and thus may contributed to remoted regulation. Both, mice with a thymus-specific mutation in *Insm1* or nude mice transplanted with *Insm1* mutant thymus, displayed autoimmune responses in multiple peripheral tissues. Together, our data demonstrate a role of Insm1 in development of mTECs and immune tolerance.

Introduction

The thymus is a primary lymphoid organ where T cell progenitors undergo maturation and selection to become functional T cells. The medullary thymic epithelial cells (mTECs) control negative selection which eliminates self-antigen-recognizing T cells, and promote differentiation of Treg (Foxp3+CD25+) cells (1, 2). Expression of tissue-restricted antigens in mTECs, also called promiscuous gene expression, is believed to be a key for negative selection and the establishment of self-tolerance (3). The autoimmune regulator (Aire) is the essential transcription factor that directly regulate TRA expression and autoimmunity (4, 5). In the last two decades, studies showed that the Aire-expressing cells defined the functional mTEC populations and have provided key molecular insights into how the TRAs expression is regulated by Aire (6-12). Particularly, Aire was identified as a chromatin looping regulator in mTECs, where Aire binds to super-enhancers and promotes the interaction between enhancers and the TRA genes (13). Studying on Aire also revealed that correct TRAs expressing in the perinatal period but not at the adult stages is essential for the establishment of self-resistance (14, 15).

Recently, lineage specific transcription factor defined mTECs were believed appeared in post-Aire populations, which is called mimetic cells (16). Aire is partially and variably required for the development of mimetic cells and the expression of the mimetic cells expressing TRAs. For each lineage of mimetic cells, specific transcription factors were required for the driving of the lineage development and accumulation. Therefore, the Aire-dependent TRAs

and the mimetic cells expressing TRAs establishment the cellular and

molecular bases of the central tolerance.

Herzig *et al*. screened potential regulators of Aire-expression, and

identified Insulinoma-associated protein 1 (Insm1) as a transcription factor

expressed in mature mTECs and a candidate regulator of Aire. However, no

functional analysis of Insm1 was performed in mTECs (17). We present here

a genetic and molecular analysis of Insm1 in the thymus, and show that

Insm1 is a novel regulator in both Aire-expressing mTECs and mimetic

neuroendocrine cells.

Results

***Insm1* is expressed in Aire-expressing and post-Aire mTECs**

To investigate the expression of *Insm1* in the thymus, we performed

immunofluorescence analysis. For this we used *Insm1$^{+/lacZ}$* and *Insm1$^{lacZ/lacZ}$*

mice in which one or two alleles of *Insm1* codon sequence were replaced by

lacZ that encodes beta-galactosidase (β-gal) (18, 19). Therefore, the

expression of *Insm1* can be monitored using β-gal. We compared signals

obtained using antibodies against β-gal and Insm1, and observed that β-gal

was expressed in the medulla of the thymus in both *Insm1$^{+/lacZ}$* and

Insm1$^{lacZ/lacZ}$ mice (Fig. S1A). Insm1 immunoreactivity overlapped with β-gal in

the thymus of *Insm1$^{+/lacZ}$* but was absent in *Insm1$^{lacZ/lacZ}$* mice (Fig. 1A and

Fig.S1A). Thus, the Insm1 antibody specific detected the endogenous Insm1

expression. Unlike the exclusive nuclear location in neuroendocrine cells (20,

21), Insm1 protein was detected in both, nuclei and cytoplasm of thymic cells

(Fig 1B).

The thymic epithelial compartment is formed by cTECs and mTECs, and

contains also lymphocytes and dendritic cells (22). To define the cell-type that

expresses *Insm1*, we performed immunofluorescence using antibodies

against Insm1 and cell type specific markers in both, the fetal and adult

thymus. We did not observe co-expression of Insm1 with the lymphocyte

specific marker CD45, or the dendritic cell marker Cd11b and Cd11c in

thymuses of E18.5 and 6-week old mice (Fig. S1B,C). However, Insm1 was

co-expressed with the mTECs marker keratin 5 (Krt5) but not the cTECs

marker keratin 8 (Krt8) (Fig. 1C,D). Using Insm1/Aire double antibodies

staining, we further observed that 70% of the Insm1-positive cells co-

expressed Aire and vice versa (Fig. 1E). The similar co-expression pattern of

Insm1 and Aire was also observed in thymuses of adult mice (Fig S1D).

Moreover, the Aire-dependent TRA insulin and somatostatin can be detected

co-expressing with Insm1 (Fig S1E and S2A). Using the published single cell

RNAseq data (16), we investigated the sub-populations of mTEC that

expressing *Insm1*. We identified that *Insm1* is expressed in Aire-expressing,

transit-amplifying and neuroendocrine mTECs, while Aire is expressed mainly

in Aire-expressing and transit-amplifying mTECs (Fig 1F,G and Fig S2B,C).

This is consistent with the observation in immunofluorescence analysis that

most of the of Insm1 expressing cells co-expressed Aire (Fig 1D). We further

verified the co-expressing of Insm1 with neuroendocrine mimetic cell markers

Foxa2 and chromogranin A in adult thymus (Fig 2A,B). In summary, *Insm1* is

expressed in Aire-expressing mTECs and neuroendocrine mimetic cells.

Tissue morphology and cell populations in *Insm1* mutant thymus

There was no difference in the appearance of size and shapes of thymus

at E18.5 in mutants (Fig S2D). The distribution and relative amounts of

mTECs and cTECs identified by Krt5 and Krt8, respectively, were comparable

between wildtype and *Insm1* mutant mice (Fig S2E,F). We used flow

cytometry to further investigated subpopulation of mTECs. The number of

EpCAM labeled thymic epithelial cells (CD45$^-$EpCAM$^+$) and UEA1 labeled

mTECs (CD45$^-$EpCAM$^+$UEA1$^+$) were comparable in wildtype controls and

Insm1 mutants both at E18.5 and adults (Fig.2C,D). However, the proportions

of Aire-positive cells were significantly decreased in populations of mTECs

(CD45$^-$EpCAM$^+$Ly51$^-$) in *Insm1* mutants at E18.5 (Fig.2E). We did not observe

significant alteration of Aire-positive cell proportions in mTECs in adults

(Fig.2E). Thus, *Insm1* mutation affected the development of Aire-positive

mTECs.

Using qRT-PCR, we examined the expression of genes that were

identified specifically expressed in each of the mimetic cells(16) (Fig 2F upper

panel). We detected genes that are specifically expressed in neuroendocrine,

entero-hepatocyte and Ptf1a+pancreatic mimetic cells were downregulated in

Insm1 mutant adult mice (Fig 2F lower panel). Thus, the neuroendocrine,

entero-hepatocyte and Ptf1a+pancreatic mimetic cells were affected by *Insm1*

mutation.

Decreased expression of *Aire* and mTEC genes in *Insm1* mutants

As we observed a decreased number of Aire-positive cells in the *Insm1*

mutant thymus, we investigated the expression levels of Aire. The

immunofluorescence intensity which indicated the Aire protein level was

moderately but significantly decreased in mTECs of *Insm1* mutant mice

(Fig.3A). Using real-time quantitative reverse transcription PCR (qRT-PCR),

we detected decreased transcript levels of *Aire* in mTECs analyzing either in

Insm1 mutant E18.5 mice or thymic specific mutant adult mice

(*Foxn1Cre;Insm1^{flox/flox}*, subsequently called *Thy-cKO*) (Fig. 2B,C). Thus, the

Insm1 mutation resulted in decreased expression of *Aire*.

To analyze the global gene expression changes in *Insm1* mutant thymus,

we performed RNA-seq using mTECs isolated from both *Insm1* mutant E18.5

and *thy-cKO* adult mice. To our surprising, differential gene expression test

by DESeq2 under cutoff of FDR<0.1 and fold change>1.5, identified 63 and

97 dysregulated genes in E18.5 and adult mice, respectively (Fig 3D,E and

Table S1,S2). The numbers of the dysregulated genes were far less than that

of Aire-regulated genes reported in previous study (23). Nevertheless, *Aire*

expression was detected down-regulated in both E18.5 and adults in the

RNAseq data (Fig 3D,E), which was consistent with the observation in qRT-

PCR and immunofluorescence analysis (Fig 3A-C). Recent studies using

single cell technology illustrated the sub-lineages of thymic epithelial cells (16,

24). To characterize the *Insm1* effects on different mimetic cell types, we

analyzed the expression pattern of our dysregulated genes in the published

thymus single-cell data (16). The down-regulated genes distributed both in

Aire-expressing mTECs and in mimetic cells (Fig 3F upper and middle panels).

However, the cTECs specific genes were up-regulated in mTECs of *Insm1*

mutant adult mice (Fig 3F lower panel). Thus, although moderate numbers of

the dysregulated genes were identified in *Insm1* mutants, Insm1 regulates the

gene expression in Aire-expressing mTECs and mimetic cells.

177   Considering each of the TRA typically expressed in only 1-5% of mTECs

178 (25), to clarify the TRA expression trends, we used a loose cutoff (p value

179 <0.05 and fold change>1.5) (Fig 3D,E) and distinguished 78 and 81

180 downregulated TRA genes at E18.5 and adults, respectively (Fig S3A, Fig 3G

181 and Table S3,S4). Using qRT-PCR, we verified some of the dysregulated

182 TRAs genes that were identified with the this cutoff (Fig 3B,C). We compared

183 the downregulated genes observed in *Insm1* mutant mTECs with the ones

184 identified in *Aire* mutant mTECs (23). Among the dysregulated TRA genes,

185 we found both, Aire-dependent and -independent TRAs to be downregulated

186 (Fig 3G and Fig S3A). Among the 81 downregulated TRA genes in adults,

187 tween-six was downregulated in both, *Aire* and *Insm1* mutants. However, fifty-

188 five (68% of all TRAs) TRA genes were unique to *Insm1* mutant mTECs

189 (Fig.3G). Thus, Insm1 regulates the expression of TRA genes in mTECs, and

190 particularly both Aire-dependent and -independent TRAs.

191   We characterized the expression patterns of downregulated TRAs in

192 mimetic cells using the published thymus single-cell data (16). Although

193 *Insm1* expression was restricted in neuroendocrine cells in mimetic cells (Fig

194 1F), the downregulated TRAs were not restricted in neuroendocrine cells,

195 instead, the distribution of these TRAs was found in Aire-expressing mTECs

196 and multiple mimetic cell types, i.e., enterocytes/hepatocytes and microfolds,

197 both in E18.5 and in adult (Fig 3H). We investigated the tissue types where

198 TRA genes are dominantly expressed using public microarray data (26). We

199 noticed that Insm1-induced TRAs were expressed in all the tested peripheral

200 tissues and with relative higher frequency in the gastrointestinal tract and

201 neuroendocrine tissues (Fig.S3B).

Insm1 promotes *Aire* and TRA expression

We induced over-/ectopic-expression of *Insm1* (*Insm1OE*) *in vivo* using

genetic tools (Fig S4), i.e., a mouse line that harbors a *loxp-STOP-loxp-Insm1*

cassette in the *Rosa26* locus, and the F*oxn1Cre* transgenic allele that allows

thymus specific expression. We observed increased numbers of Insm1-

positive cells and enlarged Krt5-positive mTECs areas in the *Insm1OE*

thymus (Fig 4A). Moreover, the Aire-positive cells were increased in the

thymic sections, although the increase of such cells was less obvious as the

one of Insm1-positive cells (Fig 4A). Increased expression levels of *Aire* were

detected by qRT-PCR using RNA isolated from the whole thymus or from

mTECs (Fig 4B,C). Interestingly, four of the nine TRA genes that were

downregulated in *Insm1* mutant thymuses showed significantly increased

expression upon *Insm1* overexpression (Fig 4B,C). We investigated the

expression of mimetic cell marker genes that was downregulated in *Insm1*

mutant mTEC. *Insm1* overexpressing promoted the neuroendocrine cells

specific genes expression (Fig 4B,C). Thus, Insm1 promoted mTECs fate and

the expression mTEC specific genes such as *Aire,* TRAs and neuroendocrine-

specific markers.

Insm1 binds on promoters and super-enhancers in mTECs

Insm1 was identified as a transcriptional factor in endocrine cells (20). To

investigate the mechanism of Insm1 function, we examined the genome wide

DNA-binding of Insm1 using CUT & Tag analysis. We detected 5,206 Insm1

binding peaks (union of 3 repeats, q<1e-5) in mTEC of E18.5 and 1,458

(union of 2 repeats, q<1e-5) peaks in mTEC of adults (Fig 5A,B). Although a

less numbers of Insm1 binding peaks were identified in adults, 1,382 (94.7%)

peaks identified in adult mTEC were overlapped with the peaks detected in

E18.5 mTEC, indicating a significant amount of consensus Insm1 binding on

DNA in mTEC during development and in adults.

Although the majority of Insm1 binding sites were located on promoter

regions in both E18.5 (50%) and adults (65.8%) (Fig 5A,B), only two (Fig 5C,

E18.5) and three (Fig 5D, adult) of these Insm1 bound genes were

dysregulated in Insm1 mutation (Fig 5C,D). Furthermore, we found the

dysregulating genes, which were identified by a looser cutoff (p-value<0.05,

FC>1.5) in the RNAseq data, were not enriched but depleted for Insm1

binding on the promoter (Fisher test, p<0.0001) (Fig 5E). These evidences

suggest that Insm1 binds on promoters but did not contribute to the correlated

gene expression.

Insm1 is co-expressed with Aire in mTECs (Fig1E, Fig S1D), we

therefore compared the DNA binding of Insm1 with that of Aire identified in

published data sets (23, 27). The majority of Insm1 binding sites (72-78%)

were co-occupied by Aire (Fig S5A,B). Since Aire was reported binding on

super-enhancers and performing regulatory roles in mTEC (13), we

investigated the Insm1 binding sites and found significantly enriched binding

of Insm1 on super-enhancers (Fig.5F-H). We further found that the

dysregulated genes were significantly over-represented within 500kb of the

Insm1-binding super-enhancers at E18.5 mTECs (Fig. 5I). These results imply

that Insm1 could bind on super-enhancers and participate in the gene

expression regulation in the developing mTECs.

Autoimmune responses in nude mice transplanted with *Insm1* mutant thymuses

We used thymus transplantation in nude mice to study autoimmune reactions. For this, T cells depleted thymuses that were isolated from fetal control (*Insm1*$^{+/lacZ}$) and mutant (*Insm1*$^{lacZ/lacZ}$) mice were transplanted into the renal capsule of six-week old nude mice (28) (Fig.6A,B). Eight weeks after transplantation, the size and structure of the transplanted thymuses were similar regardless whether they derived from control or *Insm1* mutant animals (Fig.6C and Fig.S6). Further, the numbers of CD4+ and CD8+ cells detected by flow cytometry were comparable in the transplanted control and mutant thymuses (Fig.6D). However, the spleens of nude mice transplanted with a mutant thymus (*KO/nu* mice) were increased in size and weight compared to the one of animals that received a control transplant (*Het/nu* mice) (Fig. 6E). Furthermore, the structure of the spleen was altered (Fig 6F), but the lymph nodes were unchanged (Fig S6). In particular, spleen follicles of *KO/nu* mice were smaller than that those of *Het/nu* mice (Fig. 6F). We investigated lymphocyte infiltration of various organs using hematoxylin eosin (H&E) staining, that detects the accumulation of lymphocyte nuclei. Increased infiltration was observed in pancreatic islets, lungs, kidneys and salivary glands of *KO/nu* mice (Fig.6G,H), but not in other investigated tissues (Fig.S6).

Next, we analyzed autoimmune reactions by the detection of autoimmune antibodies. Among the 13 investigated tissues, we observed autoimmune antibody reactions in pancreatic islets, kidney, and testis by staining these tissues with the serum obtained from the *KO/nu* mice (Fig 6I and Fig S7). In

particular, serum of *KO/nu* mice detected subsets of pancreatic polypeptide

(Pp) positive cells, glucagon (Gcg) positive alpha cells, and somatostatin (Sst)

positive delta cells, whereas insulin (Ins) positive beta cells were rarely

detected (Fig 6J). Thus, autoimmune antibody against PP-, alpha- and delta-

cells were present in *KO/nu* mice. To sum up, the transplantation of *Insm1*

mutant thymuses into nude mice results in autoimmune responses in multiple

tissues.

Thymus specific *Insm1* mutant mice show autoimmune responses

We employed thymus-specific *Insm1* mutant mice, *Foxn1Cre;Insm1^{flox/flox}*

(*Thy-cKO*), and littermates *Foxn1Cre* or *Insm1^{flox/flox}* controls to further

examine the autoimmune phenotype. The *Thy-cKO* mice were born healthy

and displayed no obvious abnormalities in adulthood. We observed

decreased expression levels of *Insm1, Aire* and the Aire-dependent and -

independent TRAs in mTECs of *Thy-cKO* mice (Fig 3C), which is similarly to

the changes observed in *Insm1* null mutant mTECs (Fig 3B). CD4$^+$ and CD8$^+$

T cells was properly generated in the thymus (Fig 7A) and developed in lymph

nodes (Fig 7B) of *Thy-cKO* mice, as analyzed by flow cytometry. However, we

detected significantly decreased numbers of Treg cells (CD8$^-$

CD4$^+$Foxp3$^+$CD25$^+$) in the thymus of *Thy-cKO* mice (Fig 7C), and we

observed a trend towards a reduced number of Treg cells in lymph nodes that

did not reach statistical significance (Fig 7D).

We investigated the infiltration of lymphocytes into various tissues at 6-

week, 6-, 12- and 18-month-old *Thy-cKO* animals. Generally, we observed

pronounced lymphocyte infiltration in multiple organs that increased in

frequency with the age of the animals. Infiltrated lymphocytes were observed

in salivary glands of 6-week-old and in multiple organs of 1.5-year-old *Thy-*

cKO mice, the infiltrated organs including the pancreas, the slavery gland, the

liver, the lung and the brown fat (Table 1 and Fig 7E). Further, we detected

autoimmune antibodies against multiple tissues including the pancreas, the

salivary gland and the ovary when serum of *Thy-cKO* mice was used for

immunostaining (Table 2 and Fig 7F). Thus, mutation of *Insm1* in the thymus

resulted in autoimmune responses in periphery organs in a subset of the *thy-*

cKO mice.

Discussion:

Insm1 was previously described as a factor functioning in neuroendocrine

cells and neurons (19-21, 29-31). Herzig, Y *et al* reported the first evidence

indicating Insm1 is a candidate transcriptional regulator of *Aire* (17). However,

to our knowledge functional data on a role of Insm1 in thymus have not been

systematic analyzed. We used *Insm1* null mutant mice, thymus-specific *Insm1*

mutants and transplanted the thymus of *Insm1* mutants for an analysis of a

role of Insm1 in autoimmunity.

We detected that *Insm1* is expressed in mTECs. Majority of Insm1 is

expressed in Aire positive cells while a minor population of Insm1 positive

cells did not express Aire. Using the published single cell RNAseq data (16),

we found that *Insm1* is expressed in Aire-expressing and transit-amplifying

TECs, which is similar as the expression pattern of *Aire*. In addition, *Insm1* but

not *Aire* is expressed in neuroendocrine cells, a mimetic cell type in post-Aire

population. In *Insm1* mutant mTECs, the expression of *Aire*, the Aire-

dependent and -independent TRAs was decreased. In consistent with the

gene expression, *Insm1* mutation results in less Aire-positive cells in

developmental mTECs. To identify the regulatory role of Insm1 in mimetic

cells, we examined the mimetic cell-type-specific gene expression and found

only those genes expressed in the neuroendocrine, enterohepatic, and

pancreatic cells was downregulated in *Insm1* mutation. When *Insm1*

overexpression was induced in the thymus, enlarged areas of Krt5 expression

were observed, and the expression level of *Aire* was increased. Further, TRAs

and neuroendocrine cell marker genes was induced. Currently, the molecular

and cellular model of TRA expression showed that Aire-expressing mTECs,

which developed from transit-amplifying TECs, express a selected set of

TRAs that directedly regulated by Aire. Aire-expressing mTEC subsequently

differentiate into diverse mimetic cell types, which produce mimetic cell

specific TRAs. The maturing thymocytes then undergo negative or agonist

selection against the full ranges of TRAs expressed by Aire-expressing

mTECs and mimetic cells (24). In this model, Aire deficiency impaired the

function of Aire-expressing mTECs and the development of mimetic cells,

which results in diminished expression of numerous TRAs. In our study, we

showed that *Insm1* is expressed in transit-amplifying TECs, Aire-expressing

mTECs and neuroendocrine mTECs, indicating Insm1 exists in various

developmental stages of mTECs. In addition, Insm1 regulates not only the

expression of Aire and Aire-dependent TRAs but also neuroendocrine mimetic

cell specific gene. Thus, Insm1 contributes to the development and function of

Aire-expressing mTECs and post-Aire neuroendocrine mimetic cells.

Interestingly, we observed that Insm1 binding on the super-enhancer

regions in mTECs. These super-enhancers contribute to the expression of

Insm1-dependent genes. In addition, the majority of Insm1 binding sites

overlapped with those of Aire, pointing towards the functional importance of

the set of cis-elements binding by both, Insm1 and Aire in mTEC. Although a

relative high proportion of common Insm1 binding sites were observed in both

developing and adult mTECs, the dysregulated genes were not conserved

between E18.5 and adult in *Insm1* mutants. Rather, more up-regulated genes

were identified in *Insm1* mutant adult mTECs than that of E18.5. These up-

regulated genes are cTEC-specific genes, indicating a cTEC gene expression

feature appeared in *Insm1* mutant adult mTECs. In addition, the Insm1-

binding super-enhancers were significant correlated with the Insm1

dependent gene expression at E18.5 but not at adult stages, one explanation

is that the dysregulated genes which are not directly regulated by Insm1 may

have been accumulated in adults. As *Insm1* is expressed in several

subpopulations of TECs, Further analysis on the mechanism of Insm1

function in the individual sub-population is needed, particularly in Aire-

expressing mTECs and/or neuroendocrine mimetic cells.

Mutation of *Insm1* results in about 50% decrease of *Aire* expression. A

dosage effect of Aire in TRAs regulation was discussed controversially. *Herzig*

et al showed that 50% decreased Aire expression in *Aire*[+/-] mice did not alter

the Aire-dependent TRAs expression (17). In contrast, others demonstrated

an 80-90% reduction in the expression of a particular TRAs (*Ins2, Tff3, Mup1*

and *Spt1*) in *Aire*[+/-] compared to wild-type littermate control mice (32).

Additional studies also showed that TRAs levels were regulated by dosage of

Aire expression (33, 34). In the *Insm1* mutant thymus, we observed

downregulated expression of Aire-dependent TRA genes, possibly due to the

decreased Aire expression.

In nude mice transplanted with *Insm1* mutant thymuses or in thymus

specific *Insm1* mutant mice, we observed lymphocyte infiltration and

autoimmune antibodies. These autoimmune responses were directed against

various tissues, among them are pancreatic islets, the salivary gland, the lung

and the kidney. However, not all these were equally attacked by immune

reactions. Whereas pancreatic islets and the salivary gland were affected

frequently in both, nude mice transplanted with *Insm1* mutant thymuses or

thymus specific *Insm1* mutant mice, immune reactions in other tissues were

observed only in occasionally. In addition, autoimmune responses in brown

adipose tissues were observed in *Insm1* mutants, and have not been reported

for *Aire* mutant mice (4, 5, 9, 10, 12, 14). In conclusion, out data show that

Insm1 is a novel regulator of mTEC development, gene expression and

particularly of TRAs expression in mTECs. Mutation of *Insm1* can cause

autoimmune reactions.

Materials and Methods:

Animals

Insm1$^{lacZ/lacZ}$ mice were described (18). E18.5 embryos were collected,

genotyped and analyzed. A thymus specific mutation of *Insm1* was

introduced by crossing *Insm1$^{flox/flox}$* (20) and *Foxn1Cre* mice (JAX No.

018448). Tissues collected from *Rag1$^{-/-}$* mice (GemPharmatech

No.T004753) were used for autoimmune antibody test. The *Insm1KI*

mouse strain was generated by introducing Insm1 coding sequences

preceded 5' by a stop cassette, that can be removed by Cre, into the

Rosa26 locus (Cyagen, Suzhou, China) (Fig S8). The littermate wildtype or

heterozygous animals were used for control animals in the analysis. All

animal experiments were approved by the institutional animal care and use

committee of the Jinan University (IACUC-20211123-02).

Thymus transplantation

Thymuses were isolated from E18.5 *Insm1$^{+/lacZ}$* and *Insm1$^{lacZ/lacZ}$* mice

and cultured for eight days in RPMI-1640 containing 10% FBS and 1.25

mM 2'-deoxyguanosine. One thymus was transplanted into renal capsule

of a nude mouse (6-week-old, CD1 background) using the procedure

described previously (28). Eight weeks after transplantation, the thymus

and newly generated lymphocytes were collected from the nude mice,

various tissues of the animals were also collected and analyzed.

Thymic cell dissociation and mTEC isolation

The fetal and adult thymic cells were dissociated using Liberase (35).

In brief, the thymuses were separated and connective tissues and fat were

removed. For E18.5 animals, 2-3 thymuses with of same genotype were

pooled, and thymic lobes were cut into small pieces in RPMI to release the

lymphocytes. Lymphocytes were collected in those cases when the entire

thymic cell population was analyzed. Otherwise, the thymuses were

digested in 100ul RPMI1640 containing 0.1mg/ml Liberase (Roche,

Germany) and 20U/ml DNase I at 37°C for 4 minutes. The digestion

procedure was repeated twice, and cells were collected after each round

of digestion.

mTECs isolation was performed using a magnetic beads-based

method as previously described (36). In brief, 1μl biotinylated UEA1

(Vector Laboratories, B-1065-2) was incubated with 25μl Biotin Binder

Dynabeads (Thermo Fisher Scientific,11047) for 30 minutes at room

temperature. Unbound UEA1 was removed by washing in 1ml FACS buffer

(PBS with 0.02%BSA and 5mM EDTA). Thymuses from 2-3 E18.5 fetal or

one adult mice were incubated with above prepared 25μl UEA1-coated

beads at 4°C for 30 minutes. Unbound cells were removed by three

washes with FACS buffer and by additional incubation with CD45 S-

pluriBeads (pluriSelect Life Science, SKU#70-50010-11), before mTECs

was collected for analysis. The isolated mTECs were used for RNAseq,

Cut&Tag and qRT-PCR analysis.

Flow cytometry analysis

For analysis of lymphocytes in the transplanted thymuses, thymic

lobes were gentle cut, released lymphocytes were collected and directly

stained with CD4 and CD8 antibodies in FACS buffer for flow cytometric

analysis. For analysis of thymic cells, stroma cells were dissociated using

Liberase as described above. Staining with antibodies recognizing cell

surface proteins were performed directly in FACS buffer, whereas staining

with antibodies recognizing nuclear antigens was performed after fixation

for 10 minutes in 4% PFA. Foxp3 staining was performed in

Foxp3/Transcription Factor Staining Buffer (Invitrogen, 00-5523-00).

Immunostaining procedures were performed as described (37). The

antibodies used were listed in the supplementary table 5.

BD FACSCanto II or FACSAria II were used to analyze and collect

cells (Beckton Dickenson, Franklin Lakes, USA). Cell quantification was

determined using FloJo 7.6.5 software.

Immunohistochemistry, western blot analysis and hematoxylin and

eosin (H&E) staining

Immunohistochemistry and western blot analysis were performed as

described (20). Fluorescence was imaged on a Zeiss LSM 700 confocal or

a Leica DMi8 microscope, and the images were processed using Adobe

Photoshop software. The antibodies used are listed in supplementary table

5. For the autoimmune antibody tests, serum of the *KO/nu* or *Thy-cKO*

mice was used at the concentration of 1:25 on tissue sections obtained

from *Rag1^{-/-}* mice.

For H&E staining, the Lillie-Mayer method was used. In brief, 5%

aluminum ammonium sulphate and 0.5% hematoxylin were used to stain

the nuclei, followed by 0.3% acid alcohol incubation for the differentiation

of nuclear staining. Acetified eosin was used for counterstain to reveal

cellular detail.

Insm1 **mutants RNA-seq analysis**

For E18.5 mice, six to nine thymuses of each genotype (*Insm1* mutant

and wildtype) were collected, and mTECs were isolated after pooling two

to three thymuses of the same genotype. For 6-8 weeks mice, mTECs

were isolated from each of the thymus for each genotype. Total RNAs

were isolated using TRIZOL reagents. Three independent sequencing

libraries for each genotype were generated using the NEBNext Ultra RNA

Library Prep Kit for Illumina (NEB, USA). Illumina NovaSeq 6000 was used

for sequencing, and 150-nt paired-end reads were generated. Sequencing

data (6 Gb or more) with more than 94% of bases scoring above Q30

(accuracy rate 99.9%) were produced from each library sample. RNAseq

fastq files were aligned to the mouse genome (mm10) using STAR

(v2.5.2a) (38) with parameter --outFilterMismatchNmax 2. The proportions

of uniquely mapped reads were 79.4%-85.9%. Gene read numbers were

counted using HTSeq (v0.11.0) (39) based on mouse gene annotation

Ensembl v79, and the value of fragment per kilobase of exon per million

reads mapped (FPKM) was calculated. Next, we used DESeq2 (v1.34.0)

(40) to detect dysregulated genes under cutoff FDR<0.1 and fold

change>1.5. To analyze downregulated TRAs and dysregulating-trend

genes for down-steam analysis, a looser cutoff (P value<0.05 and fold

change>1.5) was performed. All the RNA-seq data are accessible on Gene

Expression Omnibus (GSE193929).

For comparison of genes dysregulated after *Aire* mutation, published

data were used (23). Gene read counts were downloaded from GEO

(ID:GSE144877). Differential expressed genes were analyzed according to

the method used in the referred paper (23).

GO enrichment analysis was performed by R package topGO

(v2.38.1) (41) with "classic" algorithm and "Fisher exact test", followed by

the Benjamini-Hochberg multiple test correction using FDR=0.05 as a

cutoff.

qRT-PCR analysis

Cells were lysed and total RNA was isolated using TRIZOL reagent

(Invitrogen). For qRT-PCR analysis, RNA was isolated from thymuses or

mTECs of E18.5 fetal mice. cDNA was synthetized using HiScript II Q RT

SuperMix for qPCR (+gDNA wiper) (R223-01, Vazyme, China) and

analyzed using SYBR Green I based real-time quantitative PCR method

on a CFX96 RT-PCR system (Bio-Rad). Expression levels were

determined using the $2^{-\Delta\Delta Ct}$ method and *ActB* or *Krt5* as internal

standards, and displayed as proportion of control. Primers used for

quantitative analysis are listed in supplemental table S6.

Insm1 CUT&Tag-seq analysis

CUT&Tag (Cleavage Under Targets and Tagmentation) analysis was

performed according to the instruction of the manufacturer (Novoprotein,

N259-YH01) and previously published protocols (42). In brief, mTECs

were isolated using UEA1 coupled Dynabeads from wildtype mice. 0.5µl of

Insm1 antibody or 1µg IgG antibody was incubated overnight at 4°C with

5×10^{4} mTEC cells in a volume of 100µl first antibody buffer. Donkey anti-

guinea pig (Jackson ImmunoResearch,706-005-148) secondary antibody

was used and incubated with the mTEC for 45 minutes in 100µl secondary

antibody buffer at room temperature. After washing to remove unbound

antibodies, the ChiTag transposon was incubated with mTECs for one

hour at room temperature. Fragmentation, DNA purification and library

construction were performed according to the protocol of the manufacturer.

At least three biological replicas were obtained.

Illumina NovaSeq 6000 was used for sequencing, and 150-nt paired-

end reads were generated. At least twelve Gigabyte pass filtering

sequencing data was generated. Raw CUT&Tag sequencing data was

trimmed to remove adaptor sequences using fastp (v0.12.0) (43) and

aligned to mouse genome (mm10) using Bowtie2 (v2.2.9) (44) with the

parameters "--local --very-sensitive-local --no-unal --no-mixed --no-

discordant --phred33 -X 700". Peaks were called using MACS2 (v2.2.6)

(45) with the parameter "--keep-dup auto". IgG data were used as a

control. Insm1-binding peaks identified in three replicates with q-

value<10e-5 were merged for subsequent analysis.

To calculate the proportions of Insm1 binding on different genomic

regions, peaks were assigned to different genomic regions based on their

summit positions on the Ensembl transcriptome annotation according to

the priority order of CDS, 5'-UTR, 3'-UTR, promoter, ncRNA, intron, and

intergenic, where the promoter were defined as the regions of -2000 bp to

+500 bp of transcription start site (TSS). The CUT&Tag sequencing data

are accessible on Gene Expression Omnibus (GSE193929).

Single cell RNA-seq data analysis

Preprocessed transcript-by-cell matrix of scRNA-seq data of Pdpn⁻CD104⁻

mTEC^{lo} from perinatal and adult mice was downloaded from GEO

(ID:GSM5831744), and read by scanpy (46) for downstream analysis. First,

the valid 8,236 mTECs used in the study (16) were selected according to the

barcode list provided. The gene expression matrix was normalized to counts

per 10,000 counts, followed by log-transfer and scaling to unit variance and

zero mean. Next, the top 2,000 highly variable genes were identified and

selected, and PCA was performed for dimensionality reduction. According to

the elbow point of the PC contribution curve, the top 30 PCs were used for the

neighborhood graph construction, and UMAP was created for visualization.

Cell clustering were preformed using the Leiden method (resolution=1.9). The

scores of gene sets were calculated using the scanpy function tl.score_genes.

TRA gene analysis

the list of TRA genes were identified in an earlier study (47) and

provided by the authors (Prof. Perreault C, Université de Montréal,

personal communication). Aire-dependent TRAs were identified by the

overlaps of downregulated genes from the public Aire mutant RNA-seq

data (23). The TRA expression profiles of various mouse tissues were

generated in pervious microarray studies (26) and downloaded from GEO

(GSE10246). To visualize TRA expressions in different tissues (Figure S3),

samples of each tissue were merged by mean expression values, followed

by z-score normalization on each gene.

Super enhancer analysis

Super enhancer regions were identified in the published study (13) and

were provided by the author via email (Prof. Diane Mathis, Harvard

Medical School, personal communication). The control regions used in Fig

4B were the sequences with the same length as the super enhancer and

with an offset of the distance of the super enhancer length plus 200kb.

Acknowledgments

The authors thank Prof. Carmen Birchmeier (Max-Delbrück-Center for

Molecular Medicine, Germany) for helpful discussions at the initiation of

the project, for providing the *Insm1$^{+/lacZ}$ Insm1$^{+/flox}$* animals and for critical

reading of the manuscript. The authors thank Dr. St-Pierre, C (University of

Montreal) and Prof. Perreault, C (University of Montreal) for kindly sharing

the detailed information of TRAs. The authors thank Prof. Diane Mathis

(Harvard Medical School) for providing the super-enhancer coordinates.

The authors thank Prof. Yuanzhi Lu (Department of Pathology, The First

affiliated hospital of Jinan University) for the help in analyzing lymphocytes

infiltration.

Funding

This work was supported by the National Natural Science Foundation of China (31970856) and the Clinical Frontier Technology Program of the First Affiliated Hospital of Jinan University, China (JNU1AF-CFTP-2022-a01236).

Author Contributions

W.T. designed the study and performed the molecular experiments. Y.W. and J.W. contributed to the animal experiments, tissues analysis and flow cytometry analysis. Y.W., Z.Y., W.Y. and G.Y. contributed to the molecular experiments, cells and tissues experiments. J.X. performed and managed the bioinformatic analysis and participated manuscript preparation. S.J. supervised the project, analyzed the data and wrote the manuscript. S.J. is the guarantor of this work and takes responsibility for the integrity of the data and the accuracy of the data analysis.

References

1. Takahama Y (2006) Journey through the thymus: stromal guides for T-cell development and selection. *Nat Rev Immunol* 6(2):127-135.
2. Alves NL*, et al.* (2014) Serial progression of cortical and medullary thymic epithelial microenvironments. *Eur J Immunol* 44(1):16-22.
3. Kyewski B & Derbinski J (2004) Self-representation in the thymus: an extended view. *Nat Rev Immunol* 4(9):688-698.

4. Ramsey C, *et al.* (2002) Aire deficient mice develop multiple features of APECED phenotype and show altered immune response. *Hum Mol Genet* 11(4):397-409.

5. Anderson MS, *et al.* (2002) Projection of an immunological self shadow within the thymus by the aire protein. *Science* 298(5597):1395-1401.

6. Abramson J, Giraud M, Benoist C, & Mathis D (2010) Aire's partners in the molecular control of immunological tolerance. *Cell* 140(1):123-135.

7. Anderson AO & Shaw S (2005) Conduit for privileged communications in the lymph node. *Immunity* 22(1):3-5.

8. Chuprin A, *et al.* (2015) The deacetylase Sirt1 is an essential regulator of Aire-mediated induction of central immunological tolerance. *Nat Immunol* 16(7):737-745.

9. Danso-Abeam D, Humblet-Baron S, Dooley J, & Liston A (2011) Models of aire-dependent gene regulation for thymic negative selection. *Front Immunol* 2:14.

10. Giraud M, *et al.* (2012) Aire unleashes stalled RNA polymerase to induce ectopic gene expression in thymic epithelial cells. *Proc Natl Acad Sci U S A* 109(2):535-540.

11. Matsumoto M, *et al.* (2013) Which model better fits the role of aire in the establishment of self-tolerance: the transcription model or the maturation model? *Front Immunol* 4:210.

12. Oven I, *et al.* (2007) AIRE recruits P-TEFb for transcriptional elongation of target genes in medullary thymic epithelial cells. *Mol Cell Biol* 27(24):8815-8823.

13. Bansal K, Yoshida H, Benoist C, & Mathis D (2017) The transcriptional regulator Aire binds to and activates super-enhancers. *Nat Immunol* 18(3):263-273.

14. Guerau-de-Arellano M, Martinic M, Benoist C, & Mathis D (2009) Neonatal tolerance revisited: a perinatal window for Aire control of autoimmunity. *J Exp Med* 206(6):1245-1252.

15. Gabler J, Arnold J, & Kyewski B (2007) Promiscuous gene expression and the developmental dynamics of medullary thymic epithelial cells. *Eur J Immunol* 37(12):3363-3372.

16. Michelson DA, Hase K, Kaisho T, Benoist C, & Mathis D (2022) Thymic epithelial cells co-opt lineage-defining transcription factors to eliminate autoreactive T cells. *Cell* 185(14):2542-2558 e2518.

17. Herzig Y, *et al.* (2017) Transcriptional programs that control expression of the autoimmune regulator gene Aire. *Nat Immunol* 18(2):161-172.

18. Gierl MS, Karoulias N, Wende H, Strehle M, & Birchmeier C (2006) The zinc-finger factor Insm1 (IA-1) is essential for the development of pancreatic beta cells and intestinal endocrine cells. *Genes Dev* 20(17):2465-2478.

19. Jia S, Wildner H, & Birchmeier C (2015) Insm1 controls the differentiation of pulmonary neuroendocrine cells by repressing Hes1. *Dev Biol* 408(1):90-98.

20. Jia S, *et al.* (2015) Insm1 cooperates with Neurod1 and Foxa2 to maintain mature pancreatic beta-cell function. *EMBO J* 34(10):1417-1433.

21. Welcker JE, *et al.* (2013) Insm1 controls development of pituitary endocrine cells and requires a SNAG domain for function and for recruitment of histone-modifying factors. *Development* 140(24):4947-4958.

22. Anderson G, *et al.* (2014) Mechanisms of thymus medulla development and function. *Curr Top Microbiol Immunol* 373:19-47.

23. Tomofuji Y, *et al.* (2020) Chd4 choreographs self-antigen expression for central immune tolerance. *Nat Immunol* 21(8):892-901.

24. Michelson DA & Mathis D (2022) Thymic mimetic cells: tolerogenic masqueraders. *Trends Immunol* 43(10):782-791.

25. Derbinski J, Schulte A, Kyewski B, & Klein L (2001) Promiscuous gene expression in medullary thymic epithelial cells mirrors the peripheral self. *Nat Immunol* 2(11):1032-1039.

26. Lattin JE, *et al.* (2008) Expression analysis of G Protein-Coupled Receptors in mouse macrophages. *Immunome Res* 4:5.

27. Goldfarb Y, *et al.* (2021) Mechanistic dissection of dominant AIRE mutations in mouse models reveals AIRE autoregulation. *J Exp Med* 218(11).

28. Wang J, *et al.* (2019) Renal Subcapsular Transplantation of 2'-Deoxyguanosine-Treated Murine Embryonic Thymus in Nude Mice. *J Vis Exp* (149).

29. Jacob J, *et al.* (2009) Insm1 (IA-1) is an essential component of the regulatory network that specifies monoaminergic neuronal phenotypes in the vertebrate hindbrain. *Development* 136(14):2477-2485.

30. Wildner H, Gierl MS, Strehle M, Pla P, & Birchmeier C (2008) Insm1 (IA-1) is a crucial component of the transcriptional network that controls differentiation of the sympatho-adrenal lineage. *Development* 135(3):473-481.

31. Wiwatpanit T, *et al.* (2018) Trans-differentiation of outer hair cells into inner hair cells in the absence of INSM1. *Nature* 563(7733):691-695.

32. Kont V, *et al.* (2008) Modulation of Aire regulates the expression of tissue-restricted antigens. *Mol Immunol* 45(1):25-33.

33. Fornari TA, *et al.* (2010) Age-related deregulation of Aire and peripheral tissue antigen genes in the thymic stroma of non-obese diabetic (NOD) mice is associated with autoimmune type 1 diabetes mellitus (DM-1). *Mol Cell Biochem* 342(1-2):21-28.

34. Oliveira EH, *et al.* (2016) Aire Downregulation Is Associated with Changes in the Posttranscriptional Control of Peripheral Tissue Antigens in Medullary Thymic Epithelial Cells. *Front Immunol* 7:526.

35. Seach N, Wong K, Hammett M, Boyd RL, & Chidgey AP (2012) Purified enzymes improve isolation and characterization of the adult thymic epithelium. *J Immunol Methods* 385(1-2):23-34.

36. Dohr D, Engelmann R, & Muller-Hilke B (2019) A novel method to efficiently isolate medullary thymic epithelial cells from murine thymi based on UEA-1 MicroBeads. *J Immunol Methods* 467:12-18.

37. Griger J, *et al.* (2017) Loss of Ptpn11 (Shp2) drives satellite cells into quiescence. *Elife* 6.

38. Dobin A, *et al.* (2013) STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29(1):15-21.

39. Anders S, Pyl PT, & Huber W (2015) HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinformatics* 31(2):166-169.

40. Love MI, Huber W, & Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology* 15(12).

41. Rahnenfuhrer AAaJ (2016) topGO: Enrichment Analysis for Gene Ontology.

42. Kaya-Okur HS, *et al.* (2019) CUT&Tag for efficient epigenomic profiling of small samples and single cells. *Nat Commun* 10(1):1930.

43. Chen S, Zhou Y, Chen Y, & Gu J (2018) fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* 34(17):i884-i890.

44.   Langmead B & Salzberg SL (2012) Fast gapped-read alignment with Bowtie
2. *Nat Methods* 9(4):357-359.
45.   Zhang Y, *et al.* (2008) Model-based analysis of ChIP-Seq (MACS). *Genome*
Biol 9(9):R137.
46.   Wolf FA, Angerer P, & Theis FJ (2018) SCANPY: large-scale single-cell
gene expression data analysis. *Genome Biol* 19(1):15.
47.   St-Pierre C, Trofimov A, Brochu S, Lemieux S, & Perreault C (2015)
Differential Features of AIRE-Induced and AIRE-Independent Promiscuous
Gene Expression in Thymic Epithelial Cells. *J Immunol* 195(2):498-506.
48.   Szot GL, Koudria P, & Bluestone JA (2007) Transplantation of pancreatic
islets into the kidney capsule of diabetic mice. *J Vis Exp* (9):404.
714

Figure legends

Figure 1. *Insm1* is expressed in thymic epithelial cells

(A) Immunofluorescence analysis using antibodies against β-gal (red) and

Insm1 (green); analyzed were thymuses of E18.5 *Insm1$^{+/lacZ}$* and

Insm1$^{lacZ/lacZ}$ mice. DAPI (48) was used as a counterstain. Scale bar =

40µm. (B) Immunofluorescence analysis of thymuses at E18.5 using

antibodies against Insm1 (red), EpCAM (green) and DAPI (48). EpCAM is

expressed in thymic epithelial cells and labels the cell membrane and

cytoplasm. Yellow arrowheads indicated the cytoplasmic and nuclear

location of Insm1, and the magenta arrowhead indicated a nucleus. Scale

bar=10µm.

(C,D) Immunofluorescence analysis of thymuses of E18.5 (C) and 6 weeks

old (D) mice using antibodies against Insm1 (red), Krt5 (green) and Krt8

(48). Magnifications are shown in the panels on the right. Scale bar =

100µm; and 20µm in magnified panels. (E) Immunofluorescence analysis

detected Insm1 (green) and Aire (red) in thymuses of E18.5 mice. Insm1 is

co-localized with Aire. Yellow arrowheads indicated the cell expressing

only Insm1, magenta arrowheads indicated cells expressing only Aire.

Scale bar = 40µm. Quantification of single and double positive cells is

shown on the right (animal number n=3, around 300 cells/animal/antibody

were counted). (F) Insm1 expressing in subpopulations of Pdpn⁻CD104⁻

mTEC^{lo} cells analyzed using the published data (16). (G) Aire expressing

in subpopulations of Pdpn⁻CD104⁻mTEC^{lo} cells analyzed using the

published data (16).

Figure 2. Cellular phenotypes observed in the *Insm1* mutant thymus

(A,B) Immunofluorescence analysis of thymuses of 6-week-old mice using

antibodies against (A) Foxa2 (green) and Insm1 (red), and (B) ChgA

(green) and Insm1 (red). Scale bar = 25µm and 10µm in the main and

magnified panels, respectively. (C) Flow cytometry analysis of CD45⁻

/EpCAM⁺ epithelial cells (the upper panel) and CD45⁻/EpCAM⁺/UEA1⁺

mTECs (the lower panel) in thymuses of *Insm1* mutant and wildtype mice

(n=11). (D) Flow cytometry analysis of CD45⁻/EpCAM⁺ epithelial cells (the

upper panel) and CD45⁻/EpCAM⁺/UEA1⁺ mTECs (the lower panel) in

thymuses of *Thy-cKO* and wildtype mice (n=6). (E) Flow cytometry

analysis of Aire⁺ cells in *Insm1* mutant and wildtype control animals at the

age of E18.5 and adult (animal number n=4 for each genotype). Isotype

control staining for Aire was showed on the left, quantifications were

showed on the right of the panel. (F) Quantitative RT-PCR analysis of the

expression of the mimetic cells specific genes. Upper panel showed the

genes specific expressed in the mimetic cell types refereed to the

published data (16). Red labeled the downregulated genes in *Insm1*

mutants. Lower panel showed the qRT-PCR results. Significant

downregulated genes were labeled red in the upper panel. Data in the

figure panel are presented as means ± SD, statistical significance was

assessed by 2-tailed unpaired Student's t-test. Statistical significant was

defined as *P* < 0.05.

Figure 3. Changes in gene expression in the *Insm1* mutant thymus

(A) Immunofluorescence analysis of the expression of β-galactosidase (β-

gal, red) and Aire (green) in thymuses of E18.5 mice. DAPI (48) was used

as a counterstain. Quantifications of the fluorescence intensity are shown

on the right (animal number n=4 for each genotype, 80-100 Aire+/β-gal+

cells were counted). Scale bar=20µm. (B,C) Quantitative RT-PCR analysis

of the expression of *Aire* and TRA genes in mTECs of E18.5 (B) and at

adults (C) (n=4). Data in the figure panel are presented as means ± SD,

statistical significance was assessed by 2-tailed unpaired Student's t-test.

ns: *P*>0.05; *: *P* < 0.05, **: *P* < 0.01, ***: *P* < 0.001. (D,E) Volcano blots of

the RNAseq data in *Insm1* mutants versus wildtype controls at E18.5 (D)

and adult stages (E). FDR≦0.1 or p-value≦0.05 combined with Fold

change ≧ 1.5 were used for identifying the dysregulated genes. The

numbers of the dysregulated genes were showed in figures accordingly.

(F) Expression distribution of the down- and up- regulated genes identified

in the sub-populations of mTECs, presented by the violin plots of gene

score for each cell. Upper panel, the down-regulated genes in *Insm1*

mutants at E18.5; middle panel, down-regulated genes in *Insm1* mutants

at adult; lower panel, up-regulated genes in *Insm1* mutants at adult. (G)

Heatmap of dysregulated TRA genes expressed in mTEC of *cKO* mice.

(H) Distribution of the gene scores for each cells showing the dysregulated

TRA expression in the sub-populations of mTECs. The dysregulated TRAs

genes in *Insm1* mutants of E18.5 (the upper panel) and of the adult (the

lower panel) were showed respectively.

Figure 4. Insm1 promotes thymic gene expression in the thymus

(A) Immunofluorescence analysis of the expression of Insm1 (green), and

Krt5 (red, upper panels), Aire (red, lower panels) in *Insm1*-overexpression

(*Insm1OE*) thymuses of postnatal 2-day animals. DAPI (48) was used as a

counterstain. (B-C) Comparison of gene expression in *Insm1OE* and

control thymuses using postnatal 2-day animals (animal number n=5-6) (B)

and mTECs (animal number n=3-4) (C) using qRT-PCR. Data are

presented as means ± SD, statistical significance was assessed by 2-

tailed unpaired Student's t-test. *: $P < 0.05$, **: $P < 0.01$, ***: $P < 0.001$.

Figure 5. Insm1 binds to super-enhancer loci

(A, B) Distribution of Insm1 binding sites in mTEC genome at the E18.5

stage (A) and at the adult stage (B). (C,D) the Insm1 binding traces on

dysregulated genes at the E18.5 stage (C) and at the adult stage (D). (E)

Proportions of genes with Insm1 binding sites in the promoter regions (-

2000bp to +500bp of TSS) at the E18.5 stage (left) and adult stage (right).

Insm1 binding sites are significantly depleted on the promoters of

dysregulated genes (F) Density curves of detected loci Inms1 and Aire co-

binding (48), Insm1-only binding (orange), and Aire-only binding (green),

aligned to the center (0 position) of the super-enhancers. (G,H) Proportion

of Insm1 binding peaks overlapping with super-enhancer and control loci

at the E18.5 stage (G) and the adult stage (H). An identical length

sequence located 200kb plus the length of super-enhancer away from

each super-enhancer was selected as the control sequence. (I)

Proportions of super-enhancers located within ±500kb of genes that were

dysregulated (p-value $\leqq$ 0.05,FC $\geqq$ 1.5) in *Insm1* mutant mTECs at the

E18.5 stage. Stratified by the super-enhancers bound or unbound by

Insm1. In (E), (G), (H) and (I), numerators and denominators of each

proportion are given above the bars. The bars show 95% confidential

intervals. The *p*-values of Fisher test are also given.

Figure 6. Transplantation of *Insm1* mutant thymuses in nude mice

(A) Schematic outline for transplantation experiments. Thymuses were

isolated, depleted for lymphocytes by 8-day culture in 2-deoxygranosine,

transplanted into nude mice, and the mice were analyzed 8 weeks after

transplantation. (B) Transplantation of the thymus under the kidney

capsule of the nude mouse. Enlarged picture shows the kidney with the

transplanted thymus. (C) Isolated kidneys showing the thymuses under the

kidney capsule 8-week after transplantation. *Het/nu*: the thymus was

isolated from an *Insm1$^{+/lacZ}$* mouse and transplanted into a nude mouse;

KO/nu: the thymus was isolated from an *Insm1$^{lacZ/lacZ}$* mouse and

transplanted into a nude mouse. (D) CD4 and CD8α staining and flow

cytometry analysis of thymocytes isolated from the transplanted thymuses

of *Het/nu* and *KO/nu* mice (n=3). (E) Appearance (upper panel) and

quantification of the weight (lower panel) of spleens isolated from *Het/nu*

and *KO/nu* mice (n=5-6). (F) H&E staining shows the structures of spleens

isolated from *Het/nu* and *KO/nu* mice. (G) Summary of lymphocytes

infiltration in multiple tissues of *Het/nu* and *KO/nu* mice (n=6). (H) H&E

staining of the pancreas, the lung, the kidney and the salivary gland

isolated from *Het/nu* and *KO/nu* mice. 6-8 sections from the non-serial

sections were used for H&E staining for each tissue. Red arrows indicate

sites of lymphocytes infiltration. (I) Immunostaining using serum (green)

isolated from *Het/nu* and *KO/nu* mice on sections that were prepared from

Rag1$^{-/-}$ mice. 3-4 sections collected from the non-serial sections of two

Rag1$^{-/-}$ mice were used for the serum staining for each tissue. (J) Shown is

the co-staining of the *KO/nu* serum (green) with pancreatic islet specific

hormones PP, Gcg, Sst and Insulin (red, as indicated in the panels).

Statistical data are presented as means ± SD, significance was assessed

by 2-tailed unpaired Student's t-test. *: p<0.05.

Figure 7. Autoimmune phenotype in thymus specific *Insm1* mutant

mice

(A) CD4 and CD8α staining and flow cytometry analysis of thymocytes

from thymuses of *Thy-cKO* and wildtype mice (mean and SD showed in

figures, n=10-12). (B) CD4 and CD8α staining and flow cytometry analysis

of thymocytes from axillary lymph nodes of T*hy-cKO* and wildtype mice

(mean and SD showed in figures,n=9). (C, D) Flow cytometry analysis of

CD4$^+$/CD25$^+$/Foxp3$^+$ Treg cells from the thymus (C, n=8) and the axillary

lymph node (D, n=9) of *Thy-cKO* and wildtype mice. (E) H&E staining of

brown fat, pancreas, salivary gland, liver and lung in *Thy-cKO* and *Insm1*$^{f/f}$

or *Foxn1Cre* control mice. 6-8 sections from the non-serial sections were

used for H&E staining for each tissue. Animal numbers were list in Table 1.

(F) Immunostaining using serum isolated from *Thy-Cko* and *Insm1*$^{f/f}$ or

Foxn1Cre mice on sections of pancreas, salivary gland and ovary

prepared from *Rag1*$^{-/-}$ mice. 3-4 sections collected from the non-serial

sections of two *Rag1*$^{-/-}$ mice were used for the serum staining for each

tissue. The animals that the serum was collected were list in Table 2.

Statistical data are presented as means ± SD, significance was assessed

by 2-tailed unpaired Student's t-test. $p > 0.05$ in figure A and B, p values

showed in figures C and D.

Table 1. Lymphocytes infiltration in tissues

	17-month									6-month								6-week							
Genotypes	*thy-cKO*					*Ctl*				*thy-cKO*				*Ctl*				*thy-cKO*				*Ctl*			
Mice	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25
Gender	F	F	F	M	M	F	F	M	M	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F
Pancreas	+	-	-	+	++	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Salivary gland	++	+/-	-	++	++	+/-	+/-	+/-	+/-	+/-	+/ -	+	+	nt	+/-	+/-	-	+	-	+	-	-	-	-	-
Liver	++	+/-	+/-	-	nt	+/-	+/-	nt	-	+/-	++	++	+/-	-	+/-	+/-	-	+	-	-	-	-	-	-	-
Kidney	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Spleen	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Brain	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Ovary	-	-	-			-	-			-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Intestine	-	-	-	nt	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	nt	-	-	-
Colon	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Stomach	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Eye	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Ganglia	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Heart	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Lung	++	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Skeletal muscle	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	nt	-	nt	-	-	nt	-	-
Brown fat	-	+	+-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Testis				-	-			-	-																

881   Severe infiltration: ++; frequently observed infiltration: +; occasionally observed infiltration: +/-; Negtive: -;   nt: not test. F: female; M: male.

Table 2. Autoimmune antibody reaction in tissues

Genotypes	17-month									12-month						6-month											
	thy-cKO					Ctl				thy-cKO			Ctl			thy-cKO				Ctl				thy-cKO		Ctl	
Mice	1	2	3	4	5	6	7	8	9	26	27	28	29	30	31	10	11	12	13	14	15	16	17	32	33	34	35
Gender	F	F	F	M	M	F	F	M	M	F	F	M	F	F	M	F	F	F	F	F	F	F	F	M	M	M	M
Pancreas	+	-	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	+	+	-	-	-	-
Salivary gland	-	-	-	++	-	-	-	-	-	+	-	-	-	-	-	+	-	-	+	-	-	-	-	-	-	-	-
Liver	-	-	-	-	-	-	-	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Kidney	-	-	-	-	-	-	-	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Brain	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Ovary	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	+	+	+	-	-	-	-	-	-	-	-
Intestine	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Colon	-	-	-	+	-	-	-	-	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Stomach	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Eye	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Ganglia	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Nerve fiber	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Heart	-	-	-	-	-	-	-	-	-	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Lung	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Skeletal muscle	-	-	-	-	-	-	-	-	-	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Brown fat	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Testis	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Severe positive: ++; Positive: +; Negtive: -. F: female; M: male.

Fig.1

Fig. 2

Fig 3

Fig. 5

Fig. 5

A E18.5 thymic organ culture

F *Het/nu*   *KO/nu* — Spleen

B

C *Het/nu*   *KO/nu*

E *Het/nu*   *KO/nu*

D *Het/nu*   *KO/nu*

CD4 vs CD8α
Het/nu: 31.5%, 49.4%, 12.8%, 6.27%
KO/nu: 43.1%, 40.1%, 11.6%, 5.20%

Weight of spleen (mg)
Het/nu   *KO/nu*   *

G *Het/nu*   *KO/nu*

Pancreas   Lung   Kidney   Salivary gland   Liver   Intestine

H *Het/nu*   *KO/nu*
Pancreas — 200μm
Lung — 200μm
Kidney — 200μm
Salivary — 200μm

I Pancreatic islets   Kidney   Testis
Het/nu
KO/nu

J
PP KO/nu serum   Gcg KO/nu serum
Sst KO/nu serum   Ins KO/nu serum

Fig. 6

Fig. 7