Genomes and transcriptomes help unravel the complex life cycle of the blastoclad fungus,

Coelomomyces lativittatus, an obligate parasite of mosquitoes and microcrustaceans

Cassandra L. Ettinger[a,#], Talieh Ostovar[a,b,$], Mark Yacoub[a,$], Steven Ahrendt[a,*], Robert H. Hice[c],

Brian A. Federici[c,d], Jason E. Stajich[a,d,#]

[a]Department of Microbiology and Plant Pathology, University of California, Riverside, Riverside,

California, USA.

[b]UCR/SDSU Joint Doctoral Program in Evolutionary Biology, San Diego State University, San

Diego, CA, USA.

[c]Department of Entomology, University of California, Riverside, Riverside, California, USA.

[d]Institute for Integrative Genome Biology, University of California, Riverside, Riverside, CA, USA

[*]**Present address** U.S. Department of Energy Joint Genome Institute, Lawrence Berkeley

National Laboratory, Berkeley, California, USA

[$]**These authors contributed equally to this work.**

[#]**Address correspondence to**: Cassandra L. Ettinger, cassande@ucr.edu; Jason E. Stajich,

jason.stajich@ucr.edu

KEYWORDS: Blastocladiomycota, RNAseq, development, genome assembly, genome

annotation, host-associated, alternations of generations

ABSTRACT:

Species of the phylum Blastocladiomycota, early diverging zoosporic (flagellated) lineages of fungi, are vastly understudied. This phylum includes the genus *Coelomomyces* which consists of more than 80 fungal species that are obligate parasites of arthropods. Known *Coelomomyces* species lack a complete asexual life cycle, instead surviving through an obligate heteroecious alternation of generations life cycle. Despite their global distribution and interesting life cycle, little is known about the genomics of any *Coelomomyces* species. To address this, we generated three draft-level genomes and annotations for *C. lativittatus* representing its haploid meiospore, orange gamete, and amber gamete life stages. These draft genome assemblies ranged in size from 5002 to 5799 contigs with a total length of 19.8-22.8 Mb and a mean of 7416 protein-coding genes. We then demonstrated the utility of these genomes by combining the draft annotations as a reference for analysis of *C. lativittatus* transcriptomes. We analyzed transcriptomes from across host-associated life stages including infection of larva and excised mature sporangia from the mosquito, *Anopheles quadrimaculatus*. We identified differentially expressed genes and enriched GO terms both across and within life stages and used these to make hypotheses about *C. lativittatus* biology. Generally, we found the *C. lativittatus* transcriptome to be a complex and dynamic expression landscape; GO terms related to metabolism and transport processes were enriched during infection and terms related to dispersal were enriched during sporulation. We further identified five HMG box genes in *C. lativittatus*, three belonging to clades with mating type (MAT) loci from other fungi, as well as four ortholog expansions in *C. lativittatus* compared to other fungi. The *C. lativittatus* genomes and transcriptomes reported here are a valuable resource and may be leveraged toward furthering understanding of the biology of these and other early diverging fungal lineages.

INTRODUCTION:

Fungi contribute to critical roles in the global ecosystem, yet knowledge of their biology,

genetics and biochemistry largely stems from observations of only two phyla, the Ascomycota

and Basidiomycota (i.e., the Dikarya). Zoosporic (flagellated) lineages of fungi make up

additional fungal phyla (including the Blastocladiomycota and Chytridiomycota), but are

generally understudied (James et al. 2020). The phylogenetic placement of these early-

diverging zoosporic lineages is controversial and under constant revision as new genomic data

becomes available (James et al. 2020; Li et al. 2021). For example, fungi belonging to the

Blastocladiomycota were originally placed together with lineages in the Chytridiomycota, but

now Blastocladiomycota is its own phylum (James et al. 2006; Porter et al. 2011). In addition,

recent phylogenetic efforts have suggested that Blastocladiomycota may be more closely

related to the Dikarya than the Chytridiomycota (Amses et al. 2022).

Within the Blastocladiomycota, the genus *Coelomomyces* (Blastocladiales;

Coelomomycetaceae) consists of more than 80 highly fastidious fungal species that are obligate

fatal parasites, primarily of mosquitoes and microcrustaceans (Whisler et al. 1975; Couch and

Bland 1985; Powell 2017; Shen et al. 2020). These fungi have a worldwide distribution and over

the last hundred years have been reported infecting all major genera of mosquitoes, i.e., *Aedes*,

Culex, and *Anopheles*, each of which contain many species that transmit pathogens that cause

medically important diseases such as malaria, filariasis, and various viral encephalitides.

Moreover, because it is difficult to detect *Coelomomyces* infections in larval and adult

mosquitoes, it is estimated that there are more than several hundred species worldwide yet to

be described (Couch and Bland 1985), making these fungi a very large group for which we

know virtually nothing about their genomes and biochemistry. This lack of knowledge is due to

the failure, despite numerous attempts, to culture any species of *Coelomomyces in vitro*. One

major difficulty is the lack of a cell wall on hyphae growing in the vegetative stages of their

mosquito and copepod hosts. As far as is known, *Coelomomyces* species lack a complete

asexual life cycle, instead surviving through an obligate alternation of generations in which a

sporophytic phase parasitizes mosquitoes (e.g., larva) and a gametopytic phase parasitizes

microcrustaceans (e.g., copepods) (Figure 1) (Whisler et al. 1975; Federici and Chapman 1977;

Couch and Bland 1985). This type of lifecycle is uncommon in fungi, though a similar

heteroecious life cycle is observed in the rust fungi (Duplessis et al. 2021).

The *Coelomomyces* life cycle begins when a biflagellate zygote encounters a mosquito larva.

The motile spore encysts on the intersegmental membrane of the mosquito cuticle, a process

facilitated by the secretion of adhesion vesicles (Travland 1979). The encysted spore develops

an appressorium and penetration tube, which penetrates through the host cuticle (Zebold et al.

1979). Once inside the mosquito larva, the so-called hyphagens grow into coenocytic hyphae

that ramify within the hemocoel over a period of seven to ten days, which then form sporangia at

their tips (Federici and Chapman 1977; Couch and Bland 1985). The mosquito larva

subsequently die and putrefy, liberating the sporangia. Meiosis then occurs in the sporangia

resulting in haploid uniflagellate meiospores, which after sporangial dehiscence seek out and

infect a crustacean host (typically copepods, though ostracods can serve as hosts as in some

Coelomomyces species (Whisler et al. 2009)).

The penetration of copepods is thought to occur in a manner similar to that of the mosquito larva

(Federici and Chapman 1977; Zebold et al. 1979), after which hyphae of the gametophyte form

a holocarpic gametangium that cleaves into gametes. The meiospores that infect the copepods

are of opposite mating types, thus forming gametangia in the copepod host that can generate

gametes of opposite mating types. When gametogenesis is complete, the gametangium bursts,

killing the copepod host and allowing the gametes to escape through fissures in the

intersegmental membranes. If a copepod is infected by meiospores of each mating type, the

gametangia burst simultaneously, and gametes of opposite mating types mate forming

biflagellate zygotes within the dead copepod, which then seek out a mosquito larval host after

release, thereby completing the alternations of generations life cycle (Whisler et al. 1975). If

only a single mating type gametangium develops within a copepod, the gametes swim to the

surface where they seek a mate in the water in which the mosquito larvae are breeding. In

some species, such as *C. punctatus* and *C. dodgei*, the gametangia and gametes of different

mating types, much like those of *Blastocladiella emersonii*, are different colors, apparently due

to different levels of β-carotene, with one isoforms being bright orange, and the other light

amber (Federici 1977; Federici and Thompson 1979).

As noted above, despite their worldwide distribution and relatively unique life cycle, little is

known about the biology, biochemistry, or genomic landscape of *Coelomomyces* species.

Modern molecular and genomic techniques allow us to circumvent the need for *in vitro* culturing

and to expand foundational knowledge of this enigmatic fungal genus. Toward this goal, we

have established an *in vivo* culture of *C. lativittatus*, a close relative of *C. dodgei* and *C.*

punctatus (Federici 1979; Couch and Bland 1985), which we maintain using the mosquito,

Anopheles quadrimaculatus, and the copepod, *Acanthocyclops vernalis*. The research

presented here represents the first exploratory investigation of *Coelomomyces* genomics and

the *C. lativittatus* transcriptome.

To begin to answer questions related to *Coelomomyces* biology, we generated draft genomes

and annotations for *C. lativittatus* from three life stages: (i) meiospores, (ii) orange gametes, and

(iii) amber gametes. We generated transcriptomes from across life stages including infection of

An. quadrimaculatus hosts and sporangia excised from *An. quadrimaculatus*, to elucidate genes

involved in the unique biology and alternation of generations life cycle of this fungus. We then

searched for mating type loci in *C. lativittatus*, as well as looked at expansions of orthologous

genes compared to close relatives in the Blastocladiomycota and Chytridiomycota. The *C.*

lativittatus genomes and transcriptomes reported here provide an invaluable foundational

resource for understanding the biology of this and other unique and important understudied

fungal lineages in various worldwide aquatic ecosystems.

MATERIALS AND METHODS:

Study system

Larvae and copepods used for maintenance of the *in vivo* culture of *C. lativittatus* were,

respectively, *Anopheles quadrimaculatus* and *Acanthocyclops vernalis*. These were maintained

in culture as described previously (Federici 1983).

DNA extraction methods & sequencing

We sought to generate genomes from three haploid life stages (i) meiospores, (ii) orange

gametes, and (iii) amber gametes. To obtain *C. lativittatus* meiospores, mosquito larvae with

advanced infections were collected when full of sporangia, within a day or two of death. To

induce germination of sporangia, each larva was surface-sterilized by rinsing it in 70% ethanol

for 20 seconds, after which each was placed in 1 mL of double-distilled water in a 22 mm plastic

Petri dish at room temperature. The larvae were dissected with jeweler's forceps and most of

the cuticle and midgut were removed from the water. Typically, the sporangia dehisced,

releasing meiospores, 48-72 hours after incubation at room temperature. Meiospore samples

were collected using a 1 mL pipette and centrifuged using a table-top mini-fuge for 3 seconds to

sediment any sporangia in the sample. The meiospores were then pelleted by centrifugation at

16,000x*g* for 2 minutes. To obtain *C. lativittatus* gametes, infected copepods containing the

orange and amber mating types were separated prior to copepod lysis. Liquid was removed and

then copepods were surface-sterilized by rinsing in 70% ethanol and then washed with double-

distilled water to reduce contaminants. After the gametes were released from the copepods, the

copepod carcasses were removed by allowing them to settle in the microfuge tube and the

supernatant containing the gametes was transferred to a new tube. Samples were spun for 3

minutes at 6000x*g* to pellet the gametes, and the supernatant was removed.

DNA was extracted from the resulting meiospore and gamete pellets using a Qiagen genomic

DNA purification kit with Qiagen genomic-20/G tips following the standard manufacturer

protocol. DNA was then amplified using the Qiagen repli-G Whole Genome Amplification Kit

according to the standard manufacturer protocol. Illumina Libraries were prepared with the NEB

DNA Library Prep Kit. Libraries were sequenced on an Illumina HiSeq 2500 with 100 bp PE

sequencing by the UC Riverside Genomics Core Facilities.

RNA extraction and library preparation

For transcriptome analysis of the sporophyte, hyphae were excised from infected fourth instar

larvae of *An. quadrimaculatus* during either early, middle, or late stages of fungal development.

For the purpose of this study, we define early, middle, and late infection stages as follows.

Typically, the early stage of obvious infection appears as a few unpigmented, i.e., white,

sporangia at the tips of hyphae about six days after molting to the fourth instar. The fat body in

these larvae is quite well developed, and hyphae can be seen adhering to this tissue in each of

the larval abdominal segments and in the thorax. The middle stage occurs over days seven and

eight, during which the number of sporangia increases significantly, with most being mature,

meaning rusty brown in color. The late stage occurs during days nine and ten, by which time

many larvae are full of sporangia and die, although many other larvae survive another four to

five days before dying. These larvae continue to grow, being at least twice the size at which

healthy larvae pupate. Two replicate samples were collected from each time point.

For the sporangia transcriptomes, mosquito larvae with advanced infections were collected

when full of sporangia, within a day or two of death. To induce sporangia to undergo meiosis

and germinate, each larva was surface-sterilized by rinsing it in 70% ethanol for 20 seconds,

after which each was placed in 1 mL of double-distilled water in a 22 mm plastic Petri dish at

room temperature. The sporangia were excised from the larvae using jeweler's forceps, after

which the cuticle and midgut were removed from the water. Typically, the sporangia dehisced,

releasing meiospores, 48-72 hours after incubation at room temperature. Sporangia were

collected at 0, 24, 36, and 48 hour time points starting from the period the sporangia were

excised from the mosquito larva (0 hour) through dehiscence (48 hour), when the uniflagellate

meiospores burst out of the sporangia. Samples were collected using a 1 mL pipette and

centrifuged using a table-top mini-fuge for 3 seconds to obtain the sporangia in the sample. Two

replicate samples were collected from each time point.

RNA from all samples was extracted with Trizol (Life Technologies, Grand Island, NY) as per

the manufacturer protocol. 1.2 µg of RNA was used as the starting material for the NEBNext

Ultra Directional RNA Library Prep Kit for Illumina (New England BioLabs, Ipswich, MA). Poly-A

RNA was purified as per instructions and converted to adapter-ligated, size-selected cDNA. An

aliquot of the library was cloned into pJet1.2 (Thermo Fisher Scientific, Waltham, MA) and

clones sequenced with standard methods to check library quality. An aliquot was also run on a

Bioanalyzer 2100 (Agilent Technologies, Santa Clara, CA). The resulting sequencing libraries

were sequenced by the Institute for Integrative Genome Biology Core facility at the University of

California at Riverside using the MiSeq instrument with 100 bp paired-end reads (Illumina, San

Diego, CA).

Genome assembly

Genomes for the meiospore (MEIOSPORE), orange gamete (ORANGE) and amber gamete

(AMBER) libraries were assembled using the Automatic Assembly of the Fungi (AAFTF)

pipeline v. 0.2.3 (Stajich and Palmer 2019). Briefly, this involved first trimming and filtering reads

using bbduk.sh from BBTools v. 37.76 (Bushnell 2014). Next, assemblies were produced using

the 'assemble' command in AAFTF which relies on SPAdes v. 3.12.0 (Prjibelski et al. 2020) run

with default parameters to select optimal *k*-mer size and screened for contaminant vectors with

AAFTF vecscreen step using NCBI BLAST (Camacho et al. 2009). The AAFTF sourpurge step

was run which utilizes sourmash v. 3.5.0 (Brown et al. 2016) to further purge any remaining

contaminant contigs and AAFTF rmdup step using Minimap2 v. 2.17 (Li 2018) was run to

identify duplicate contigs for removal. Finally, AAFTF polish step ran Pilon v. 1.22 (Walker et al.

2014) to polish the resulting contigs in each assembly to remove potentially mis-called

consensus nucleotides or indels by SPAdes.

Assembly evaluation for each genome was performed using QUAST v. 5.0.0 (Gurevich et al.

2013) and BUSCO v. 5.0.0 (Simão et al. 2015) against both the eukaryote_odb10 and

fungi_odb10 gene sets. BUSCO assessment was also performed for reference genomes from

other fungal lineages for comparison (for lineages see Table S1). We performed telomere

searches against the *Coelomomyces* assemblies using find_telomeres.py

(https://github.com/markhilt/genome_analysis_tools) to test for telomeric repeats at the ends of

the scaffolds and determine chromosome completeness (Hiltunen et al. 2021).

Contamination screen and removal

Given the obligate nature of *C. lativittaus* with its hosts, microbial contamination was assessed

in each assembly using the BlobTools2 pipeline (Figure S1) (Challis et al. 2020). This involved

first assigning taxonomy against the UniProt Reference Proteomes database (v. 2020_10) to

each contig using diamond (v. 2.0.4) and command-line BLAST v. 2.2.30+ (Camacho et al.

2009; Buchfink et al. 2021). Next, read coverage was calculated by aligning the reads from the

MEIOSPORE, AMBER and ORANGE libraries to their respective genome assemblies with BWA

v.0.7.17 (Li and Durbin 2009) and sorted using samtools v. 1.11 (Li et al. 2009). Finally, the

BlobToolKit Viewer was used to visualize the GC content, read coverage, and predicted

taxonomies of contigs to identify contaminants. This approach flagged 1969, 11, and 24 contigs

as putative contaminants in the AMBER, MEIOSPORE and ORANGE assemblies respectively.

Microbial contamination was further assessed with the anvi'o v.7 pipeline (Eren et al. 2015,

2021), a complementary method, for the AMBER assembly given the large number of

contaminants predicted by BlobTools2 (Figure S1). This involved first obtaining read coverage

from each of the three genomic samples (AMBER, ORANGE, and MEIOSPORE) to the AMBER

assembly with bowtie2 v. 2.4.2 (Langmead et al. 2009) and samtools v. 1.11 (Li et al. 2009). A

contig database for the AMBER assembly was then generated using 'anvi-gen-contigs-

database' which uses Prodigal v. 2.6.3 (Hyatt et al. 2010) to predict open-reading frames. This

command also identifies single-copy bacterial (Lee 2019), archaeal (Lee 2019) and protista

(Delmont 2018) genes using HMMER v. 3.2.1 (Eddy 2011) and ribosomal RNA genes using

barrnap (Seemann 2018). We predicted taxonomy for each predicted ORF using Kaiju v. 1.7.2

(Menzel et al. 2016) with the NCBI BLAST non-redundant protein database nr including fungi

and microbial eukaryotes v. 2020-05-25. We then constructed anvi'o profiles for each sample

(AMBER, ORANGE, and MEIOSPORE) using 'anvi-profile' with the '--cluster-contigs' option and

a contig length cut-off of >2.5 kbp. These profiles were then merged using 'anvi-merge'.

Contaminant contigs in the AMBER assembly were then identified through 'anvi-interactive'

using a combination of hierarchical clustering based on coverage and tetranucleotide frequency,

taxonomic identity, and GC content. This second method identified 1127 contaminant contigs in

the AMBER assembly.

Contaminant contigs (e.g., any contig identified by the BlobTools2 pipeline as assigned to

bacteria, archaea or viruses and any contig identified using the anvi'o pipeline) were

subsequently removed from the draft assemblies. For the AMBER assembly, this meant

conservatively removing a total of 2091 contaminant contigs (1005 identified by both methods,

964 contigs identified by BlobTool2 only, and 122 identified by anvi'o only).

Genome annotation

Genome annotation was performed using the Funannotate pipeline v.1.7.4 (Palmer and Stajich

2020). This first involved using RepeatModeler v.1.0.11 (Flynn et al. 2020) and RepeatMasker

v.4.0.7 (Smit et al. 2013-2015) to generate a library of predicted repetitive elements and then

soft mask these elements in the draft genomes. Next the RNASeq data was assembled using

Trinity v. 2.10.0 in Genome-Guided mode and aligned with PASA v.2.3.3 to train the *ab initio*

gene prediction algorithms augustus and SNAP (Haas et al. 2003; Grabherr et al. 2011).

Consensus gene models were generated using EVidenceModeler v.1.1.1 (Haas et al. 2008) on

predicted protein-coding gene models from a combination of algorithms including CodingQuarry

v. 2.0, Augustus v. 3.3.3, GeneMark-ETS v. 4.38, GlimmerHMM, and SNAP v 2013_11_29 (Korf

2004; Majoros et al. 2004; Stanke et al. 2006; Ter-Hovhannisyan et al. 2008; Testa et al. 2015).

Transfer rRNA genes were predicted using tRNAscan-SE v. 1.3.1 (Lowe and Eddy 1997).

Protein annotations were predicted for the consensus gene models based on similarity to Pfam

(Finn et al. 2014), CAZyme domains (Lombard et al. 2014; Huang et al. 2018), MEROPS

(Rawlings et al. 2014), eggNOG v. 1.0.3 (Huerta-Cepas et al. 2016), InterProScan v. 5 (Jones et

al. 2014), and Swiss-Prot (Boutet et al. 2016) using HMMER v.3 (Eddy 2011) or diamond

BLASTP (Buchfink et al. 2015). Phobius (Käll et al. 2004) and SignalP v. 4.1c (Armenteros et al.

2019) were also run to predict transmembrane proteins and secreted proteins respectively. Any

problematic gene models that were flagged by Funannotate were manually curated as needed.

The annotation results were summarized in custom code written in R v. 4.0.3 using the tidyverse

v. 1.3.0 package (Wickham et al. 2019; R Core Team 2020). The annotated genomes of the

MEIOSPORE, ORANGE, and AMBER assemblies were aligned and mapped to the RNA

sequencing reads using HISAT2 (Kim et al. 2019).

Transcriptome analysis

Given the obligate relationship of *Coelomomyces* with its hosts, we chose a reference-based

transcriptome approach as initial results from *de-novo* approaches revealed host contamination

even after removal using a reference host transcriptome. To provide a more comprehensive

gene set to use as a reference for transcriptome analysis, we combined the predicted transcript

sets from all three genome annotations. We then used CD-HIT-EST to cluster transcripts at

90% sequence identity and evaluated this combined set (AOM90) with BUSCO v. 5.0.0 (Fu et

al. 2012; Simão et al. 2015). For comparative purposes, a protein alignment of *C. lativittatus*

with other fungal lineages (for lineages see Table S1), was constructed using the

PHYling_unified (https://github.com/stajichlab/PHYling_unified) pipeline, which uses HMMER

v.3 (Eddy 2011) and ClipKIT (Steenwyk et al. 2020) to search for markers in the protein

sequences, build, and trim an alignment based on BUSCO fungi_odb10 HMMs. A maximum

likelihood phylogeny was built from this alignment using IQ-TREE2 v.2.2.0 (Minh et al. 2020).

BUSCO fungi_odb10 gene partitions were provided to IQ-TREE2 using -p (Chernomor et al.

2016) and ModelFinder Plus was run using -m MFP to ensure use of the best evolutionary

model for each partition based on BIC (Kalyaanamoorthy et al. 2017).

Mosquito host transcripts were removed from the transcriptome data using BBMap against an

An. quadrimaculatus (Accession: GBTE00000000) reference transcriptome prior to read

quantification (Bushnell 2014; Desjardins et al. 2015). Host-filtered transcriptome read counts

were quantified against the AOM90 transcript set using kallisto (Bray et al. 2016). The count

data were then imported into R for analysis with the DESeq2, ggplot2 and GSEABase packages

(Wickham 2009; Love et al. 2014; R Core Team 2020; Morgan et al. 2022).

General expression trends across all samples were visualized using variance stabilized count

data. We then used DESeq2 to identify differentially expressed transcripts between life stages

(e.g., sporangial vs. infection). Significant genes were defined as transcripts with a false

discovery rate adjusted *p*-value of < 0.01 and a |log2 fold change| > 2. Gene Ontology (GO)

enrichment analysis was performed to assess if the differentially expressed transcripts were

significantly enriched in any particular functions ($p < 0.05$). This analysis was performed at each

of three GO classes: biological processes (BP), molecular functions (MF), and cellular

components (CC). Significantly enriched GO terms were simplified using Revigo with the default

settings (Supek et al. 2011).

In order to identify transcripts that show change in expression across the development time

course conditions within each sporangium (e.g., 0 hr vs. 24 hr vs. 32 hr vs. 48 hr) and infection

stages (e.g., early vs. middle vs. late infection), we performed a Likelihood Ratio Test (LRT).

Significant transcripts from LRT were filtered with a false discovery rate adjusted p-value of <

0.01 and a |log2 fold change| > 2. The "DEGreport" package was used to cluster genes with

similar expression profiles based on the LRT results across different time series conditions

(Pantano 2022). A GO enrichment analysis was performed to identify enriched GO terms in

each of the clusters with similar expression patterns ($p < 0.05$). GO terms were simplified using

Revigo with default settings (Supek et al. 2011).

Identification of mating type (MAT) loci in C. lativattus

To identify high mobility group box (HMG-box) genes putatively involved in mating in the *C.*

lativittatus genomes, we used HMMsearch v. 3.3.2 for PFAM PF00505 with an e-value of 1e-15

(Eddy 2011). Given that genes neighboring HMG-boxes are thought to be involved in mating in

other fungi (Vossenberg et al. 2019), we used Clinker (Gilchrist and Chooi 2021) and Cblaster

(Gilchrist et al. 2021) to assess the syntenic regions surrounding the HMG-boxes in the

Coelomomyces assemblies to identify conserved regions neighboring HMG-boxes. To confirm

phylogenetic placement of the identified HMG-box genes, we aligned the candidate genes from

C. lativattatus with those of other fungi (for fungi used see Table S1) using HMMalign (Eddy

2011). We then constructed a maximum likelihood phylogenetic tree of the HMG-box genes

using IQ-TREE using the VT+R6 model which was selected by ModelFinder Plus (Minh et al.

2020). Finally, to compare expression of HMG-box genes across *C. lativittatus* life stages, the

variance stabilized expression levels of the HMG-box genes were plotted using ggplot2

(Wickham 2009)*.* Pairwise t-tests were performed to assess differential expression between life

stages*.*

Identification of orthologous gene expansions relative to other fungal lineages

Orthofinder v. 2.5.4 was used to identify whether any differentially expressed transcripts

represented genes expanded in copy number in *C. lativittatus* compared to other fungi (for fungi

used see Table S1) (Emms and Kelly 2019). We filtered the Orthofinder results to orthogroups

containing genes with differentially expressed transcripts in the RNAseq data. These results

were subsetted by orthogroups that were at least |log2 fold change| > 2 higher in copy number

in *C. lativittatus* compared to the other fungi. Orthogroup expansions were confirmed through

phylogenetic analyses. Briefly, a nucleotide alignment of all genes in an orthogroup of interest

was produced using MUSCLE v. 5.1 and then a phylogenetic tree was built with IQ-TREE2

using -m MFP which runs ModelFinder Plus (Kalyaanamoorthy et al. 2017).

RESULTS:

Over half of the genomic landscape is represented in C. lativittatus assemblies and annotations

After successful contaminant removal, the draft genome assemblies ranged in size from 5002 to

5799 contigs with a total length between 19.8 Mb and 22.8 Mb (Table 1). Although the

assemblies were fragmented with an average N50 of 6128 bp, BUSCO assessment found that

the draft assemblies were halfway complete (Table S2). Mean completeness in 'genome' mode

was 43.1% and 56.3% using the fungi_odb10 and eukaryota_odb10 sets, respectively. While,

these values were lower than those of other blastoclads on average (fungi_odb10: 75.5%,

eukaryota_odb10: 83.8%), including a recent long-read genome from *B. emersonii*

(fungi_odb10: 81.8%) (Leonard et al. 2022), it is important to note that BUSCO sets are biased

against early diverging fungal lineages. Nonetheless, these draft assemblies provide a valuable

starting point for further improvement and refinement moving forward.

Annotation of the three assemblies with Funannotate identified on average 7416 protein-coding

genes and 59 tRNA genes, with 63.2% of these having a hit to at least one functional database

(Table 1). We combined the three predicted transcript sets together at 90% identity using CD-

HIT-EST to generate a comprehensive final gene set (AOM90) of 8645 transcripts and leading

to improved BUSCO 'protein' scores of 62.5% and 82.8% using the fungi_odb10 and

eukaryota_odb10 sets, respectively (Table S2; Figure S2). Despite being slightly lower than the

average scores for other blastoclads (Figure 2; fungi_odb10: 84.1, eukaryota_odb10: 90.1%),

the AOM90 transcript set represents a promising and robust reference for beginning to elucidate

C. lativittatus biology.

Differential expression analysis reveals distinct expression profiles between life stages

Initial analysis of the transcriptome profiles supported a distinct divide between infection and

sporangial life stages with each stage clustering separately (Figure 3A) and clear differences in

expression in the most abundant differentially expressed transcripts (Figure 3B). We followed

this analysis with differential transcript expression analysis using DESeq2 to identify specific

transcripts responsible for these patterns.

We found 1262 differentially expressed transcripts between life stages, with 395 transcripts

enriched during infection compared to 867 enriched during sporangial life stages ($p < 0.01$,

log2foldchange > 2). Of these, 575 (45.6%) had no matches to any of the databases used for

annotation. Interestingly, while more transcripts were enriched in sporangial stages, many of the

most abundant transcripts were representative of transcripts enriched in the infection stages

and many of these transcripts were unannotated. For example, of the top 25 most abundant

differentially expressed transcripts, 22 were upregulated during infection relative to sporangial

stages (Figure 3B). Further, 18 of these 25 transcripts had no significant similarity to any

features in the annotation databases, and further only one transcript was fully annotated,

HMI54_014395 (*ERG10*), an acetyl-CoA C-acetyltransferase.

We performed GO enrichment analysis on the differentially expressed transcripts to identify

enriched GO terms of interest (Table S3, $p < 0.05$). For the infection stages, GO terms from 38

biological processes (BP), 9 cellular compartments (CC), and 46 molecular functions (MF) were

identified. Of these, seven of the top ten significantly enriched BP were metabolic processes

(e.g., GO:0006082: organic acid metabolic process, GO:0046394: carboxylic acid biosynthetic

process) and three were transport-related (e.g., GO:0006848: pyruvate transport, GO:1905039:

carboxylic acid transmembrane transport). For the sporangial stages, GO terms from 35 BP, 26

CC, and 41 MF were identified. Of these, the most significantly enriched BP was related to

reproduction (GO:0000003: reproduction), with four of the top ten significantly enriched

biological processes related to metabolic and biosynthetic processes (e.g., GO:0005975:

carbohydrate metabolic process, GO:0006183: GTP biosynthetic process, GO:0006228: UTP

biosynthetic process) and three related to movement or organization of cellular machinery (e.g.,

GO:0006928: movement of cell or subcellular component, GO:0007010: cytoskeleton

organization, GO:0007017: microtubule-based process).

Differential expression analysis reveals complex pattern of expression clusters within life stages

Identification of differentially expressed transcripts across the development time course within

sporangial and infection stages was done using a Likelihood Ratio Test (LRT). We found 380

transcripts that were significantly differentially expressed between infection stages ($p < 0.01$). Of

these, we found two clusters of differentially expressed transcripts with similar expression

patterns in each, group 1 and group 2 with 167 and 213 transcripts, respectively (Figure 4A).

The same analysis with the sporangial stages indicated 3701 differentially expressed transcripts

($p < 0.01$) and out of which, we identified 7 clusters, each including differentially expressed

transcripts with similar patterns of expression. There were 1083, 965, 859, 219, 355, 156 and 64

genes in groups 1 to 7, respectively (Figure 4B).

In order to identify enriched GO terms of interest, we performed GO enrichment analysis on the

differentially expressed transcripts across the developmental conditions within infection and

sporangial stages (Figure S3, $p < 0.05$). Within the infection stages, we found 30 BP, 12 CC,

and 33 MF enriched GO terms. Of these, the top significantly enriched GO terms were in the CC

category (GO:0000786: nucleosome, GO:0043228: non-membrane-bounded organelle,

GO:0043232: intracellular non-membrane-bounded organelle, GO:0044815: DNA packaging

complex, GO:0032993: protein-DNA complex), and were also seen in group 2 with decreasing

expression across infection stages (Figure 4A).

Furthermore, we found 19 BP, 10 CC and 28 MF enriched GO terms represented within the

sporangial stages and of them, the most significantly enriched GO terms were related to

structural molecule activity and binding (GO:0005488: binding, GO:0003735: structural

constituent of ribosome, GO:0004100: chitin synthase activity, GO:0005198: structural molecule

activity, GO:0003779: actin binding) as well as the GO terms related to peptide biological

processes (GO:0006412: translation, GO:0043043: peptide biosynthetic process, GO:0006518:

peptide metabolic process). Terms related to structural molecule activity and binding were

generally seen in expression pattern groups 1 and 5 both of which generally increased in

expression over time (Figure 4B).

HMG-box loci were identified with differential expression across life stages

A total of five unique HMG-box genes were identified, with all five HMG-box genes present in

the MEIOSPORE assembly and three in each of the AMBER and ORANGE assemblies. The

identified HMG-box genes were found on small fragmented contigs (average: 8300 bp) which

contained only a few neighboring genes (average: 3 genes). Despite their small size, synteny

analysis across the three assemblies found that the AMBER and ORANGE assemblies share

two HMG-box loci with each other. The third HMG-box genes in the ORANGE and AMBER

assemblies were only shared with the MEIOSPORE assembly, which has four syntenic

orthologous HMG-boxes (Figure S4). We tested the five HMG-box loci in *C. lativittatus* for

synteny against *Allomyces macrogynus* to determine whether neighboring genes around HMG-

boxes are conserved in sexually reproducing chytrids (Lee et al. 2010). We were unable to

determine synteny of neighboring genes around HMG-box loci, possibly due to the fragmented

scaffolds where these genes are found in our *C. lativittatus* assemblies. Phylogenetic analyses

of the HMG-box genes showed that four of the HMG-box genes generally fell in clades with

other blastoclads or chytrids, while HMI54_015288 fell into a clade with Dikarya. Further, three

of the five HMG-box orthologs from *C. lativitattus* (HMI56_006544, HMI55_007199, and

HMI54_004920) were present in a clade containing known mating-related HMG-boxes (Figure

S5). HMI56_006544 and HMI54_004920 were present in all three *C. lativitattus* assemblies

while HMI55_007199 was only present in the ORANGE assembly. Interestingly HMI56_006544

is closely related to SexM in *Phycomyces blakesleeanus,* however we were unable to identify a

SexP homolog in *C. lativitattus.* The number of identified HMG-box genes here is in line with

that of other blastoclads (i.e., *Blastocladiella britannica* and *Catenaria anguillulae*; Figure 5). Of

the five identified HMG-box genes, HMI54_015288 was significantly over-expressed in the

infection stages compared to the sporangial stages. (Figure S6; $p < 0.01$).

C. lativattatus may have expanded gene families of orthologs related to its unique life history

In order to determine if gene families expanded in *C. lativattatus* we performed Orthofinder

analysis comparing *C. lativattatus* with other fungal taxa. In total we identified 37 398

orthogroups among the fungal taxa. Of these, 182 orthogroups were exclusively present in *C.*

lativattatus and absent in all other fungal taxa.

We identified 10 orthogroups that both contained genes with differentially expressed transcripts

and were expanded in copy number compared to other fungi. Of these 10 orthogroups, only four

had predicted functional annotations (Figure 5). We tested the four orthogroups for duplication

errors and removed any sequences that appeared truncated or had 100% sequence similarity to

an ortholog from a different *C. lativittatus* assembly. Most of the copy number expansions for

these four orthogroups appeared to occur on clades exclusive to *C. lativittatus.* The putative

functions for the four validated orthogroups were replication timing regulatory factor 1 (*RIF1*),

chitin deacetylase, adenosine monophosphate (AMP)-activated protein kinase 1 (*AMPK1*), and

Egh16-like virulence factor. Within these orthogroups, only the *RIF1* orthogroup contained a

gene that had higher expression in infection stages; the other three orthogroups had genes with

higher expression during sporangial stages.

DISCUSSION

C. lativittatus annotated genomes are an important community resource

The genomes assembled and annotated here, while partial, are a promising and critical

community resource as little genomic data exists for members of the Blastocladiomycota. The

smallest public genome is *B. britannica* with a genome size of 19 Mb with 9431 predicted gene

models and the largest is *Allomyces macrogynus* with a genome size of 47 Mb and 19 446

predicted genes (Grigoriev et al. 2014). *C. lativittatus* falls on the shorter end of this range with

an average genome length of 21.5 Mb and average 7475 predicted gene models, possibly due

to the incomplete nature of the draft genomes reported here. While partial, we think that the

three *C. lativattaus* genomes assembled and annotated in this study provide much needed

community resources for study of these obligate fungi. For example, the phylogenetic placement

of Blastocladiomycota has been disputed, but inclusion of additional genomes like those

reported here can help elucidate these ancient phylogenetic relationships (Amses et al. 2022).

Transcriptomic landscape of C. lativittatus life stages provides insight into Blastocladiomycota

biology

The transcriptome of *C. lativittatus* is a complex, dynamic, and underexplored landscape. The

results of this study highlight a need for future refinement and exploration of gene annotation in

this species, as evidenced by the 45.6% of differentially expressed transcripts with no

annotation, and the majority of the top 25 abundant differentially expressed transcripts lacking

functional annotation. In spite of these shortcomings, we were able to make generalizable

insights about *C. lativittatus* biology from the subset of transcripts that were annotated with GO

terms. Overall, during infection, GO terms were enriched related to metabolism and transport

processes. While during sporangial stages, GO terms were enriched related to dispersal (i.e.,

cell signaling, locomotion and transport machinery). Looking at the expression patterns within

life stages, we can begin to see more complicated trends emerge.

Within infection stages, we identified two different patterns of gene expression. In the first, gene

expression increased over the course of development time (early, middle and late infection),

with enriched GO terms related to membrane transport and metabolic processes. In the second,

gene expression decreased over time with GO terms related to DNA replication, nucleic acid

and amino acid biosynthetic processes.

The enrichment of metabolism and membrane transport processes in the transcriptomes during

infection stages is similar to reports from other early diverging fungal lineages. Upregulation of

transport-related pathways has been reported in chytrid infection of frog hosts, which the

authors suggest might be related to nutrient availability and proliferation related to host

association (Ellison et al. 2017). Further, in *Vavraia culicis* (Microsporidia) enriched GO terms

for growth, metabolism and replication were identified and posited to be related to its generalist

lifestyle and ability to infect multiple types of hosts (Desjardins et al. 2015). Here, we observed

an enrichment of metabolism terms as part of a pattern of increasing expression (group 1,

Figure 4A), and an enrichment of replication-related terms as part of a pattern of decreasing

expression (group 2, Figure 4A). Thus, we may be observing a shift in priorities during infection,

with early infection stages marked by increased replication as hyphagens grow into coenocytic

hyphae inside the host and later infection stages marked by increased metabolism as the

fungus proliferates and begins preparing to make sporangia.

Within sporangial stages, we identified seven expression patterns with two patterns displaying

higher expression over these developmental stages, two patterns displaying lower expression,

and three patterns with a relatively higher gene expression at the second time point (24 hrs),

followed by stable or decreasing expression. In general, expression patterns with higher

expression were enriched in GO terms related to chitin activity as well as terms related to

dispersal and microtubule-based processes. In decreasing expression patterns the enriched GO

terms were mostly related to metabolism and transcription. The other three expression patterns,

which displayed the highest expression at the second developmental time point, were

functionally different from each other. One group was enriched in GO terms related to dispersal

and structural machinery, another in terms related to oxidative stress responses, and the third in

terms related to lipid metabolism. Similarly, time series clustering of the transcriptome profiles of

differentially expressed genes during the sporulation of the blastoclad, *Blastocladiella emersonii,*

showed eight different patterns (Vieira and Gomes 2010).

The enrichment of reproduction and dispersal machinery-related mechanisms during the

sporangial stages likely relates to the production of meiospores. For example, signal

transduction and microtubule and cytoskeleton biogenesis were similarly reported to be

enriched during sporulation in *B. emersonii (Vieira and Gomes 2010)*. These authors also

observed a decrease in transcription and metabolism during sporulation which they attributed to

the nutritional starvation required in order to sporulate. Additionally, previous investigations into

protein synthesis in chytrids (Léjohn and Lovett 1966) and blastoclads (Lovett 1968; Schmoyer

and Lovett 1969) suggest that translation does not occur until zoospore germination and that

zoospores are likely partially dependent on maternal mRNA and ribosomes for initial protein

production. Laundon *et. al*. (2022) posited that the chytrid zoospore life stage is optimized for

dispersal to new hosts rather than general metabolism. The authors also reported complex lipid

dynamics throughout the lifecycle of the chytrid, *Rhizoclosmatium globosum.* Of particular note,

they observed increased expression of genes related to lipid transport and metabolism in

zoospores, which often have large amounts of intracellular storage lipids. In this study across

sporangial stages, we observed an enrichment of dispersal and microtubule-based terms as

part of patterns with increasing expression, and transcription terms as part of patterns with

decreasing expression (Figure 4B). We also observed one pattern of increasing and then

decreasing expression related to lipid metabolism (group 7, Figure 4B). Therefore, here we may

be discerning the metabolic preparation and production of meiospores for optimal dispersal,

survival and host identification.

Later stages of host association are likely characterized by increased immune response in the

host and countered by increased stress response by the fungus in order to continue to evade

the host immune system. Upregulation of stress response genes has been reported in the plant

pathogen *Zymoseptoria tritici* during late stages of sporulation in its host, which the authors

posited might be protective (Keon et al. 2005). Similarly, we observed an enrichment of terms

related to oxidative stress responses in the later stages of sporulation (group 6, Figure 4B),

which we speculate may assist *Coelomomyces* evade host defenses during meiospore

production.

Mating-type genes may be useful for future work on evolution of sex in fungi

Unlike in animals and plants which have sex-specific chromosomes, sex determination in fungi

is regulated by only a handful of genes. These mating-type (*MAT*) loci include HMG-box genes

(Benkhali et al. 2013). While *MAT* loci in Dikarya have been widely studied (Wallen and Perlin

2018), the *MAT* loci of early diverging lineages of fungi have received relatively less attention

(Idnurm et al. 2008). Given its obligate sexual two-host alternation of generations life cycle and

the ability to separate sexed haploid gametes by color (orange or amber), *C. lativattatus*

provides an intriguing system for investigating the evolution of sex in early diverging fungi. Using

the genomes generated here, we identified five putative HMG-box genes, including one gene,

HMI54_015288, which was differentially expressed between life stages, and three genes which

were in a clade with known mating-related genes. Additionally, HMI56_006544, which was

highly expressed during the sporangial stage, is homologous to the SexM gene of *Phycomyces*

blakesleeanus. Interestingly, the three *C. lativitattus* HMG-boxes within the clade containing

mating-related genes were up-regulated in the sporangial stage. Future work should tease apart

the role in mating of these putative HMG-box genes in *C. lativattatus* and also place these

genes in a comparative framework in order to further investigate the evolution of sex

determination in fungi.

Ortholog expansions in C. lativittatus may relate to host-association

Compared to other fungal lineages, *C. lativattatus* genomes had an enriched number of gene

copies in four orthologs with functional annotations representing chitin deacetylase, *AMPK1*,

Egh16-like virulence factor, and *RIF1* (Figure 5). We believe these expanded orthogroups may

be good candidates for future investigations elucidating mechanisms behind *C. lativattatus*-host

interactions.

In *C. lativattatus,* we found that members of the chitin deacetylase orthogroup were upregulated

during sporangial stages and that "chitin synthase activity" was also an enriched GO term in

group 1, which is a group represented by increasing expression (Figure 4B). Chitin

deacetylases catalyze the deacetylation of chitin, an important structural component of fungal

cell walls and insect cuticles, and have been previously reported in many fungal species (Zhao

et al. 2010). Chitin-binding proteins and chitin deacetylation are thought to protect fungal

pathogens against plant chitinases during infection (Gueddari et al. 2002; van den Burg et al.

2006), and have also been shown to be upregulated during infection of amphibian hosts by the

chytrid pathogen, *Batrachochytrium dendrobatidis (Ellison et al. 2017)*. Further, the *B.*

dendrobatidis genome has gene expansions of chitin-binding proteins which can confer

protection against chintanse activity when expressed in the fungus, *Trichoderma reeseii*

(Abramyan and Stajich 2012; Liu and Stajich 2015). Thus, it is possible that the expansion in the

chitin deacetylase orthogroup here may be related to *C. lativattatus* defense against its two

hosts.

AMPK1 genes are sensors that modulate energy metabolism and homeostasis, and can be

important for regulating stress responses (Hardie et al. 2012). These genes can also be used to

alter host energy metabolism by microbial pathogens (Mesquita et al. 2016). Increased counts

of *AMPK1* orthologs and higher expression in sporangial stages in *C. lativattatus* may be related

to regulation of increased stress during its two-host life cycle or to evasion of host immune

detection.

In *C. lativattatus,* we found that members of the *Egh16*-like expanded orthogroup displayed

higher expression during sporangial stages. *Egh16*-like virulence factors are proteins related to

appressorium formation and pathogenesis which are present in pathogenic fungi including

insect pathogens such as *Metarhizium acridum (Grell et al. 2003; Cao et al. 2012). Egh16*-like

factors have been postulated to aid in the penetration of insect cuticles (Cao et al. 2012). Thus,

it is possible that the expanded *Egh16*-like virulence factor orthogroup is contributing to *C.*

lativattatus virulence.

RIF1 is important for telomere length control and subtelomeric silencing in fungi and other

eukaryotes (Sreesankar et al. 2012). Subtelomeric regions have increased variability, caused

by duplications and rearrangements that can result in functional novelty, including secondary

metabolites related to pathogenicity and virulence. Silencing of subtelomeric regions is one way

pathogens can control secondary metabolite expression (Wyatt et al. 2020; Diotti et al. 2021).

Given the increased expression of members of the expanded *RIF1* ortholog during infection,

these genes may have roles in silencing subtelomeric regions and may be another tool in the *C.*

lativattatus toolbox for interacting with its hosts.

CONCLUSION

We generated three draft genomes and annotations for *C. lativittatus* and characterized the *C.*

lativittatus transcriptome landscape across infection and sporangial life stages. Little is known

about the genomic landscapes of blastoclads or zoosporic fungi and thus, the genomic and

transcriptomic data described here represent a valuable foundational resource. In the

transcriptome investigation, we identified differentially expressed transcripts and enriched GO

terms that provide insight into the blastoclad life cycle, with infection stages marked by an

enrichment of metabolism and transport processes and sporangial stages by dispersal-related

processes. Further, *C. lativittatus* has an interesting obligate alternation of generations life cycle

with two hosts and here we found several ortholog expansions in virulence related genes which

may have roles in modulating its host-associated lifecycle. As additional genomic data from

other blastoclads and zoosporic fungi are generated, more powerful comparative approaches

can be used to assess the evolutionary relationships of these lineages to other fungi, as well as

better understand their complex life histories. We hope that this work sets the stage for this

future studies by providing some foundational knowledge of these unique fungi.

DATA AVAILABILITY

The decontaminated *C. lativittatus* genome assemblies and annotation are deposited at

DDBJ/ENA/GenBank under the accessions JADGJU000000000, JADGJV000000000 and

JADHYY000000000. The versions described in this paper are JADGJU010000000,

JADGJV010000000 and JADHYY010000000. The raw sequence reads for the genome

assemblies are available through BioProjects PRJNA631428, PRJNA631429, and

PRJNA631430. The raw sequence reads for the transcriptome work are available under GEO

accession GSE222209. Analysis scripts for this work are available on GitHub

(https://github.com/stajichlab/Chytrid_Coelomomyces_Project) and archived in Zenodo

(https://doi.org/10.5281/zenodo.7435008).

ACKNOWLEDGEMENTS

Funding to support sequencing work was provided by a seed grant from the UC Riverside Office

of Research and Chancellor's Strategic Investment Funds project "*Coelomomyces Genomics*

for Mosquito Vector Control". CLE was supported by funds from California Department of Food

and Agriculture agreement # 20–0267. MY was supported by funds from Gordon and Betty

Moore Foundation (award #9337). J.E.S. is a CIFAR Fellow in the program Fungal Kingdom:

Threats and Opportunities and partially supported by USDA Agriculture Experimental Station at

the University of California, Riverside and NIFA Hatch projects CA-R-PPA-5062-H. Data

analyses performed at the High-Performance Computing Cluster at the University of California

Riverside in the Institute of Integrative Genome Biology were supported by NSF grant DBI-

1429826 and NIH grant S10-OD016290. We thank Jericho Ortañez for assistance in

experiments for generating genomic DNA and RNA.

LITERATURE CITED

Abramyan J, Stajich JE. 2012. Species-specific chitin-binding module 18 expansion in the
amphibian pathogen *Batrachochytrium dendrobatidis*. mBio 3:e00150–12.

Amses KR, Rabern Simmons D, Longcore JE, Mondo SJ, Seto K, Jerônimo GH, Bonds AE,
Alisha Quandt C, Davis WJ, Chang Y, Federici BA, Kuo A, LaButti K, Pangilinan J,
Andreopoulos W, Tritt A, Riley R, Hundley H, Johnson J, Lipzen A, Barry K, Franz Lang B,
Cuomo CA, Buchler NE, Grigoriev IV, Spatafora JW, Stajich JE, James TY. 2022. Diploid-
dominant life cycles characterize the early evolution of Fungi. Proceedings of the National
Academy of Sciences 119:e2116841119.

Armenteros JJA, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Brunak S, Heijne G von,
Nielsen H. 2019. SignalP 5.0 improves signal peptide predictions using deep neural networks.
Nature Biotechnology 37:420–423.

Benkhali JA, Coppin E, Brun S, Peraza-Reyes L, Martin T, Dixelius C, Lazar N, Tilbeurgh H van,
Debuchy R. 2013. A network of HMG-box transcription factors regulates sexual cycle in the
fungus *Podospora anserina*. PLoS Genetics 9:e1003642.

Boutet E, Lieberherr D, Tognolli M, Schneider M, Bansal P, Bridge AJ, Poux S, Bougueleret L,
Xenarios I. 2016. UniProtKB/Swiss-Prot, the manually annotated section of the UniProt
KnowledgeBase: how to use the entry view. Methods in Molecular Biology 1374:23–54.

Bray NL, Pimentel H, Melsted P, Pachter L. 2016. Near-optimal probabilistic RNA-seq
quantification. Nature Biotechnology 34:525–527.

Brown CT, Titus Brown C, Irber L. 2016. sourmash: a library for MinHash sketching of DNA. The
Journal of Open Source Software 1:27.

Buchfink B, Reuter K, Drost H-G. 2021. Sensitive protein alignments at tree-of-life scale using
DIAMOND. Nature Methods 18:366–368.

Buchfink B, Xie C, Huson DH. 2015. Fast and sensitive protein alignment using DIAMOND.
Nature Methods 12:59–60.

Burg HA van den, Harrison SJ, Joosten MHAJ, Vervoort J, Wit PJGM de. 2006. *Cladosporium*
fulvum Avr4 protects fungal cell walls against hydrolysis by plant chitinases accumulating during
infection. Molecular Plant-Microbe Interactions: MPMI 19:1420–1430.

Bushnell B. 2014. BBMap: Short read aligner for DNA and RNA-seq data. Available online at
https://sourceforge.net/projects/bbmap/.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.
BLAST+ : architecture and applications. BMC Bioinformatics 10:1–9.

Cao Y, Zhu X, Jiao R, Xia Y. 2012. The Magas1 gene is involved in pathogenesis by affecting
penetration in *Metarhizium acridum*. Journal of Microbiology and Biotechnology 22:889–893.

Challis R, Richards E, Rajan J, Cochrane G, Blaxter M. 2020. BlobToolKit – interactive quality
assessment of genome assemblies. G3 Genes|Genomes|Genetics 10:1361–1374.

Chernomor O, Haeseler A von, Minh BQ. 2016. Terrace aware data structure for phylogenomic
inference from supermatrices. Systematic Biology 65:997–1008.

Couch JN, Bland CE. 1985. The genus *Coelomomyces*. Orlando, FL 32887: Academic Press,
Inc.

Delmont T. 2018. Assessing the completion of eukaryotic bins with anvi'o. Available online at
http://merenlab.org/delmont-euk-scgs.

Desjardins CA, Sanscrainte ND, Goldberg JM, Heiman D, Young S, Zeng Q, Madhani HD,
Becnel JJ, Cuomo CA. 2015. Contrasting host–pathogen interactions and genome evolution in
two generalist and specialist microsporidian pathogens of mosquitoes. Nature Communications
6:7121.

Diotti R, Esposito M, Shen CH. 2021. Telomeric and sub-telomeric structure and implications in
fungal opportunistic pathogens. Microorganisms 9.

Duplessis S, Lorrain C, Petre B, Figueroa M, Dodds PN, Aime MC. 2021. Host adaptation and
virulence in heteroecious rust fungi. Annual Review of Phytopathology 59:403–422.

Eddy SR. 2011. Accelerated Profile HMM Searches. PLoS Computational Biology 7:e1002195.

Ellison AR, DiRenzo GV, McDonald CA, Lips KR, Zamudio KR. 2017. First in vivo
Batrachochytrium dendrobatidis transcriptomes reveal mechanisms of host exploitation, host-
specific gene expression, and expressed genotype shifts. G3 7:269–278.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative
genomics. Genome Biology 20:238.

Eren AM, Murat Eren A, Esen ÖC, Quince C, Vineis JH, Morrison HG, Sogin ML, Delmont TO.
2015. Anvi'o: an advanced analysis and visualization platform for 'omics data. PeerJ 3:e1319.

Eren AM, Murat Eren A, Kiefl E, Shaiber A, Veseli I, Miller SE, Schechter MS, Fink I, Pan JN,
Yousef M, Fogarty EC, Trigodet F, Watson AR, Esen ÖC, Moore RM, Clayssen Q, Lee MD,
Kivenson V, Graham ED, Merrill BD, Karkman A, Blankenberg D, Eppley JM, Sjödin A, Scott JJ,
Vázquez-Campos X, McKay LJ, McDaniel EA, Stevens SLR, et al. 2021. Community-led,
integrated, reproducible multi-omics with anvi'o. Nature Microbiology 6:3–6.

Federici BA. 1977. Differential pigmentation in the sexual phase of *Coelomomyces*. Nature
267:514–515.

Federici BA. 1979. Experimental hybridization of *Coelomomyces dodgei* and *Coelomomyces*
punctatus. Proceedings of the National Academy of Sciences of the United States of America
76:4425–4428.

Federici BA. 1983. Species-specific gating of gametangial dehiscence as a temporal
reproductive isolating mechanism in *Coelomomyces*. Proceedings of the National Academy of
Sciences of the United States of America 80:604–607.

Federici BA, Chapman HC. 1977. *Coelomomyces dodgei*: establishment of an in vivo laboratory
culture. Journal of Invertebrate Pathology 30:288–297.

Federici BA, Thompson SN. 1979. β-Carotene in the gametophytic phase of *Coelomomyces*
dodgei. Experimental Mycology 3:281–284.

Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY, Eddy SR, Heger A, Hetherington K,
Holm L, Mistry J, Sonnhammer ELL, Tate J, Punta M. 2014. Pfam: the protein families
database. Nucleic Acids Research 42:D222–30.

Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, Smit AF. 2020.
RepeatModeler2 for automated genomic discovery of transposable element families.
Proceedings of the National Academy of Sciences of the United States of America 117:9451–
9457.

Fu L, Niu B, Zhu Z, Wu S, Li W. 2012. CD-HIT: accelerated for clustering the next-generation
sequencing data. Bioinformatics 28:3150–3152.

Gilchrist CLM, Booth TJ, Wersch B van, Grieken L van, Medema MH, Chooi Y-H. 2021.
cblaster: a remote search tool for rapid identification and visualization of homologous gene
clusters. Bioinformatics Advances 1:vbab016.

Gilchrist CLM, Chooi Y-H. 2021. Clinker & clustermap.js: Automatic generation of gene cluster
comparison figures. Bioinformatics 37:2473–2475.

Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L,
Raychowdhury R, Zeng Q, Chen Z, Mauceli E, Hacohen N, Gnirke A, Rhind N, Palma F di,
Birren BW, Nusbaum C, Lindblad-Toh K, Friedman N, Regev A. 2011. Full-length transcriptome
assembly from RNA-Seq data without a reference genome. Nature Biotechnology 29:644–652.

Grell MN, Mouritzen P, Giese H. 2003. A *Blumeria graminis* gene family encoding proteins with
a C-terminal variable region with homologues in pathogenic fungi. Gene 311:181–192.

Grigoriev IV, Nikitin R, Haridas S, Kuo A, Ohm R, Otillar R, Riley R, Salamov A, Zhao X,
Korzeniewski F, Smirnova T, Nordberg H, Dubchak I, Shabalov I. 2014. MycoCosm portal:
gearing up for 1000 fungal genomes. Nucleic Acids Research 42:D699–704.

Gueddari NEE, El Gueddari NE, Rauchhaus U, Moerschbacher BM, Deising HB. 2002.
Developmentally regulated conversion of surface-exposed chitin to chitosan in cell walls of plant
pathogenic fungi. New Phytologist 156:103–112.

Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome
assemblies. Bioinformatics 29:1072–1075.

Haas BJ, Delcher AL, Mount SM, Wortman JR, Smith RK Jr, Hannick LI, Maiti R, Ronning CM,
Rusch DB, Town CD, Salzberg SL, White O. 2003. Improving the *Arabidopsis* genome
annotation using maximal transcript alignment assemblies. Nucleic Acids Research 31:5654–
5666.

Haas BJ, Salzberg SL, Zhu W, Pertea M, Allen JE, Orvis J, White O, Buell CR, Wortman JR.
2008. Automated eukaryotic gene structure annotation using EVidenceModeler and the
Program to Assemble Spliced Alignments. Genome Biology 9:R7.

Hardie DG, Ross FA, Hawley SA. 2012. AMPK: a nutrient and energy sensor that maintains
energy homeostasis. Nature Reviews Molecular Cell Biology 13:251–262.

Hiltunen M, Ament-Velásquez SL, Johannesson H. 2021. The Assembled and Annotated
Genome of the Fairy-Ring Fungus *Marasmius oreades*. Genome Biology and Evolution
13:evab126.

Huang L, Zhang H, Wu P, Entwistle S, Li X, Yohe T, Yi H, Yang Z, Yin Y. 2018. dbCAN-seq: a
database of carbohydrate-active enzyme (CAZyme) sequence and annotation. Nucleic Acids
Research 46:D516–D521.

Huerta-Cepas J, Szklarczyk D, Forslund K, Cook H, Heller D, Walter MC, Rattei T, Mende DR,
Sunagawa S, Kuhn M, Jensen LJ, Mering C von, Bork P. 2016. eggNOG 4.5: a hierarchical
orthology framework with improved functional annotations for eukaryotic, prokaryotic and viral
sequences. Nucleic Acids Research 44:D286–93.

Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic
gene recognition and translation initiation site identification. BMC Bioinformatics 11:119.

Idnurm A, Walton FJ, Floyd A, Heitman J. 2008. Identification of the sex genes in an early
diverged fungus. Nature 451:193–196.

James TY, Letcher PM, Longcore JE, Mozley-Standridge SE, Porter D, Powell MJ, Griffith GW,
Vilgalys R. 2006. A molecular phylogeny of the flagellated fungi (Chytridiomycota) and
description of a new phylum (Blastocladiomycota). Mycologia 98:860–871.

James TY, Stajich JE, Hittinger CT, Rokas A. 2020. Toward a fully resolved fungal tree of life.
Annual Review of Microbiology 74:291–313.

Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell A,
Nuka G, Pesseat S, Quinn AF, Sangrador-Vegas A, Scheremetjew M, Yong S-Y, Lopez R,
Hunter S. 2014. InterProScan 5: genome-scale protein function classification. Bioinformatics

30:1236–1240.

Käll L, Krogh A, Sonnhammer ELL. 2004. A combined transmembrane topology and signal
peptide prediction method. Journal of Molecular Biology 338:1027–1036.

Kalyaanamoorthy S, Minh BQ, Wong TKF, Haeseler A von, Jermiin LS. 2017. ModelFinder: fast
model selection for accurate phylogenetic estimates. Nature Methods 14:587–589.

Keon J, Rudd JJ, Antoniw J, Skinner W, Hargreaves J, Hammond-Kosack K. 2005. Metabolic
and stress adaptation by *Mycosphaerella graminicola* during sporulation in its host revealed
through microarray transcription profiling. Molecular Plant Pathology 6:527–540.

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. Nature Biotechnology 37:907–915.

Korf I. 2004. Gene finding in novel genomes. BMC Bioinformatics 5:59.

Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient alignment
of short DNA sequences to the human genome. Genome Biology 10:R25.

Laundon D, Chrismas N, Bird K, Thomas S, Mock T, Cunliffe M. 2022. A cellular and molecular
atlas reveals the basis of chytrid development. eLife 11:e73933.

Lee MD. 2019. GToTree: a user-friendly workflow for phylogenomics. Bioinformatics 35:4162–
4164.

Lee SC, Ni M, Li W, Shertz C, Heitman J. 2010. The evolution of sex: a perspective from the
fungal kingdom. Microbiology and Molecular Biology Reviews: MMBR 74:298–340.

Léjohn HB, Lovett JS. 1966. Ribonucleic acid and protein synthesis in *Rhizophlyctis rosea*
zoospores. Journal of Bacteriology 91:709–717.

Leonard G, Galindo LJ, Milner DS, Avelar GM, Gomes-Vieira AL, Gomes SL, Richards TA.
2022. A genome sequence assembly of the phototactic and optogenetic model fungus
Blastocladiella emersonii reveals a diversified nucleotide-cyclase repertoire. Genome Biology
and Evolution 14:evac157.

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics 34:3094–
3100.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics 25:1754–1760.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,
1000 Genome Project Data Processing Subgroup. 2009. The sequence alignment/map format
and SAMtools. Bioinformatics 25:2078–2079.

Liu P, Stajich JE. 2015. Characterization of the Carbohydrate Binding Module 18 gene family in
the amphibian pathogen *Batrachochytrium dendrobatidis*. Fungal Genetics and Biology: FG & B
77:31–39.

Li Y, Steenwyk JL, Chang Y, Wang Y, James TY, Stajich JE, Spatafora JW, Groenewald M,
Dunn CW, Hittinger CT, Shen X-X, Rokas A. 2021. A genome-scale phylogeny of the kingdom

Fungi. Current Biology: CB 31:1653–1665.e5.

Lombard V, Golaconda Ramulu H, Drula E, Coutinho PM, Henrissat B. 2014. The carbohydrate-
active enzymes database (CAZy) in 2013. Nucleic Acids Research 42:D490–5.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biology 15:550.

Lovett JS. 1968. Reactivation of ribonucleic acid and protein synthesis during germination of
Blastocladiella zoospores and the role of the ribosomal nuclear cap. Journal of Bacteriology
96:962–969.

Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA
genes in genomic sequence. Nucleic Acids Research 25:955–964.

Majoros WH, Pertea M, Salzberg SL. 2004. TigrScan and GlimmerHMM: two open source ab
initio eukaryotic gene-finders. Bioinformatics 20:2878–2879.

Menzel P, Ng KL, Krogh A. 2016. Fast and sensitive taxonomic classification for metagenomics
with Kaiju. Nature Communications 7:11257.

Mesquita I, Moreira D, Sampaio-Marques B, Laforge M, Cordeiro-da-Silva A, Ludovico P,
Estaquier J, Silvestre R. 2016. AMPK in Pathogens. In: M. Cordero, and B. Viollet, eds. AMP-
Activated Protein Kinase. Springer, Cham. p. 287–323.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Haeseler A von, Lanfear R.
2020. IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic
era. Molecular Biology and Evolution 37:1530–1534.

Morgan M, Falcon S, Gentleman R. 2022. GSEABase: Gene set enrichment data structures and
methods. Available online at http://bioconductor.org/packages/GSEABase/.

Palmer JM, Stajich J. 2020. Funannotate v1.8.1: Eukaryotic genome annotation. Available
online at http://doi.org/10.5281/zenodo.4054262.

Pantano L. 2022. DEGreport: Report of DEG analysis. Available online at
http://lpantano.github.io/DEGreport/.

Porter TM, Martin W, James TY, Longcore JE, Gleason FH, Adler PH, Letcher PM, Vilgalys R.
2011. Molecular phylogeny of the Blastocladiomycota (Fungi) based on nuclear ribosomal DNA.
Fungal Biology 115:381–392.

Powell MJ. 2017. Blastocladiomycota. In: J. Archibald, A. Simpson, and C. Slamovits, eds.
Handbook of the Protists. p. 1497–1521.

Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. 2020. Using SPAdes de novo
assembler. Current Protocols in Bioinformatics 70:e102.

Rawlings ND, Barrett AJ, Bateman A. 2014. Using the MEROPS database for proteolytic
enzymes and their inhibitors and substrates. Current Protocols in Bioinformatics 48:1.25.1–33.

R Core Team. 2020. R: A language and environment for statistical computing. Available online
at https://www.R-project.org/. Vienna, Austria.

Schmoyer IR, Lovett JS. 1969. Regulation of protein synthesis in zoospores of *Blastocladiella*.
Journal of Bacteriology 100:854–864.

Seemann T. 2018. barrnap: BAsic Rapid Ribosomal RNA Predictor. Available online at
https://github.com/tseemann/barrnap.

Shen D, Nyawira KT, Xia A. 2020. New discoveries and applications of mosquito fungal
pathogens. Current Opinion in Insect Science 40:111–116.

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics 31:3210–3212.

Smit AFA, Hubley R, Green P. 2013-2015. RepeatMasker. Available online at
http://www.repeatmasker.org.

Sreesankar E, Senthilkumar R, Bharathi V, Mishra RK, Mishra K. 2012. Functional
diversification of yeast telomere associated protein, Rif1, in higher eukaryotes. BMC Genomics
13:255.

Stajich J, Palmer J. 2019. stajichlab/AAFTF: v0.2.3. Available online at
http://dx.doi.org/10.5281/zenodo.3437300.

Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B. 2006. AUGUSTUS: ab initio
prediction of alternative transcripts. Nucleic Acids Research 34:W435–9.

Steenwyk JL, Buida TJ 3rd, Li Y, Shen X-X, Rokas A. 2020. ClipKIT: A multiple sequence
alignment trimming software for accurate phylogenomic inference. PLoS Biology 18:e3001007.

Supek F, Bošnjak M, Škunca N, Šmuc T. 2011. REVIGO summarizes and visualizes long lists
of gene ontology terms. PloS One 6:e21800.

Ter-Hovhannisyan V, Lomsadze A, Chernoff YO, Borodovsky M. 2008. Gene prediction in novel
fungal genomes using an *ab initio* algorithm with unsupervised training. Genome Research
18:1979–1990.

Testa AC, Hane JK, Ellwood SR, Oliver RP. 2015. CodingQuarry: highly accurate hidden
Markov model gene prediction in fungal genomes using RNA-seq transcripts. BMC Genomics
16:170.

Travland LB. 1979. Initiation of infection of mosquito larvae (*Culiseta inornata*) by
Coelomomyces psorophorae. Journal of Invertebrate Pathology 33:95–105.

Vieira ALG, Gomes SL. 2010. Global gene expression analysis during sporulation of the aquatic
fungus *Blastocladiella emersonii*. Eukaryotic Cell 9:415–423.

Vossenberg BTLH van de, Vossenberg BTLH van de, Warris S, Nguyen HDT, Gent-Pelzer MPE
van, Joly DL, Geest HC van de, Bonants PJM, Smith DS, André Lévesque C, Lee TAJ van der.
2019. Comparative genomics of chytrid fungi reveal insights into the obligate biotrophic and
pathogenic lifestyle of *Synchytrium endobioticum*. Scientific Reports 9:1–14.

Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q,

Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated tool for comprehensive microbial
variant detection and genome assembly improvement. PloS One 9:e112963.

Wallen RM, Perlin MH. 2018. An overview of the function and maintenance of sexual
reproduction in dikaryotic fungi. Frontiers in Microbiology 9:503.

Whisler HC, Karanja DMS, Shemanchuk JA, Zebold SL, Romney SV, Nielsen LT. 2009. The life
history and *in vivo* culture of *Coelomomyces utahensis* (Blastocladiomycetes). Journal of
Invertebrate Pathology 100:40–43.

Whisler HC, Zebold SL, Shemanchuk JA. 1975. Life history of *Coelomomyces psorophorae*.
Proceedings of the National Academy of Sciences of the United States of America 72:693–696.

Wickham H. 2009. ggplot2: Elegant graphics for data analysis. Springer-Verlag New York.

Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, Grolemund G, Hayes A,
Henry L, Hester J, Kuhn M, Pedersen TL, Miller E, Bache SM, Müller K, Ooms J, Robinson D,
Seidel DP, Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K, Yutani H. 2019. Welcome to the
tidyverse. Journal of Open Source Software 4:1686.

Wyatt NA, Richards JK, Brueggeman RS, Friesen TL. 2020. A comparative genomic analysis of
the barley pathogen *Pyrenophora teres f. teres* identifies subtelomeric regions as drivers of
virulence. Molecular Plant-Microbe Interactions: MPMI 33:173–188.

Zebold SL, Whisler HC, Shemanchuk JA, Travland LB. 1979. Host specificity and penetration in
the mosquito pathogen *Coelomomyces psorophorae*. Canadian Journal of Botany Journal
Canadien de Botanique 57:2766–2770.

Zhao Y, Park R-D, Muzzarelli RAA. 2010. Chitin deacetylases: properties and applications.
Marine Drugs 8:24–46.

Figures/Tables:

Table 1. Genome assembly and annotation statistics. Various statistics calculated by

QUAST for each of the assemblies are provided here including the total number of contigs in the

assembly, the total assembly length, percent GC content, the N50 and the L50. All statistics

from QUAST are based on contigs of size >= 500 bp, unless specifically noted (e.g., "# contigs

(>= 0 bp)" and "Total length (>= 0 bp)" include all contigs in each assembly). We also report the

percentage of RNAseq reads that were aligned to each assembly using HISAT, the percentage

of each assembly masked by RepeatMasker (this value includes total interspersed repeats,

simple repeats and low complexity regions), and the number of predicted telomeres. A summary

of genome annotation results is reported here including the total number of gene models,

mRNAs, tRNAs, the mean exon length (bp), tRNA length (bp), mRNA length (bp), and the

percentage of proteins with a PFAM domain, InterProScan, EggNog, COG, or GO term

matches.

Software	Statistic	Amber	Orange	Meiospore
QUAST	# contigs (>= 0 bp)	5799	5413	5002
	Largest contig	56469	30249	37025
	Total length (>= 0 bp)	22882397	19891335	21857322
	GC (%)	33.43	32.32	32.33
	N50	6067	5623	6695
	N90	1782	1603	1996
	L50	1089	1030	988
	L90	3884	3610	3254

	# N's per 100 kbp	480.4	362.27	101.38
Telomeres	Telomeres TOTAL	23	18	22
	Telomeres FRWRD	8	9	10
	Telomeres RVRS	15	9	12
HISAT2	Overall RNAseq alignment to genome	55.40%	55.17%	59.01%
RepeatMasker	Repetitive regions (%)	10.77%	12.57%	14.04%
Funannotate	Total number of gene models	7404	7344	7677
	Number of mRNA genes	7352	7289	7608
	Number of tRNA genes	52	55	69
	Mean exon length	367.89	358.18	372.63
	Mean tRNA length	74.58	74.62	74.91
	Mean mRNA length	1501.03	1309.02	1465.69
	Genes with PFAM hit (%)	48.74	44.91	48.04
	Genes with InterProScan hit (%)	65.11	63.82	64.91
	Genes with EggNog hit (%)	54.16	52.78	53.82
	Genes with COG hit (%)	53.28	52.03	52.98
	Genes with GO Term (%)	46.83	45.68	46.72

Figure 1. Alternation of generations life cycle of *Coelomomyces lativittatus*.

Diagram showing the general alternation of generations life cycle of *C. lativittatus* between

copepod and mosquito hosts. We have circled and highlighted the different life stages that were

used in this study. Genomic sequencing was performed on haploid stages: including orange

gametes (orange), amber gametes (yellow) and meiospores (purple). RNA sequencing was

performed on diploid stages including mosquito larval infection stages (green) as well as

sporangial stages excised from mosquito larva (blue).

Figure 2. *C. lativittatus* protein set is comparable to those of other fungal taxa. Moving

from left to right, here we show a maximum likelihood phylogeny which shows the relationship of

C. lativittatus to other fungal lineages. This tree was generated using IQ-TREE2 on an

alignment of BUSCO fungi_odb10 HMMs constructed using the PHYling_unified pipeline. Taxa

labels in the phylogeny are shown colored by assigned fungal phylum. In association with this

phylogeny, we then depict a barchart of the draft genome size (Mbp) for each taxon with

genome size colored by fungal lifestyle (saprobe = purple, algal associated = yellow, animal

associated = pink), followed by a barchart of predicted gene counts for each taxon with counts

colored by fungal lifestyle. Next, we show barcharts of BUSCO 'protein' completion status for

the eukaryota_odb10 and fungal_odb10 sets. Bars show the percent of genes found in each

genome annotation as a percentage of the total gene set and are colored by BUSCO status

(missing = grey, fragmented = yellow, complete and duplicated = green, complete and single-

copy = blue). The values depicted here for *C. lativittatus* gene counts and BUSCO scores are

based on the combined clustered transcript set (AOM90), and the genome size is the average

size across all three assemblies. The BUSCO scores for individual *C. lativittatus* assemblies can

be seen in Figure S2.

Figure 3. Transcript expression differs between *C. lativittatus* life stages. (A) Principal

Component Analysis (PCA) of variance stabilized transcriptomic count data. Samples are

colored by time points while shapes are used to broadly represent life stages (circle = infection,

triangle = sporangial). Ellipses represent the 95% confidence interval around the centroid of

each life stage. Replicate samples may be overlapping. (B) Heatmap showing the variance

stabilized counts of the 25 most expressed transcripts with differential expression across life

stages. Replicates are indicated by 'A' or 'B'.

1038

1039

1040

1041

1042

1043

Figure 4. Transcript expression differs across the development time course conditions in

each sporangium and infection stage. (A) The plots on the left are the two possible groups

with specific transcript expression patterns across early, middle, and late infection timepoints

and number of transcripts in each group. The table on the right shows the top five enriched GO

terms for each of the groups of transcripts. (B) The plots on the left are the seven groups with

specific transcript expression patterns across 0 hr, 24 hr, 36 hr and 48 hr timepoints within the

sporangial life stages. The table on the right shows the top five enriched GO terms for each of

the seven groups of transcripts.

1052

1053 **(A)**

1054

1055 **(B)**

Figure 5. *C. lativittatus* copy number for HMG box, *RIF1*, chitin deacetylase, *Egh16-like*,

and *AMPK1* gene orthogroups compared to other fungi. Here we depict a heatmap,

organized by fungal phylogenetic relationships, depicting the copy number of orthologous

genes, representing differentially expressed transcripts, with expanded gene counts in *C.*

lativittatus relative to other fungi. The colors of the gene counts are normalized per gene family

with red indicating a high normalized count and blue indicating a low count. The phylum for each

taxon is indicated by the colored row labels at the left. The dendrogram above the heatmap

clusters the columns by similarity in counts between the different gene families. Validated HMG

boxes were found in two orthogroups which are shown combined here for simplicity.

1073

Group	Over-represented GO term	GO ID
1 167 transcripts	aspartate family amino acid metabolic process	GO:0009066
	methionine metabolic process	GO:0006555
	alpha-amino acid metabolic process	GO:1901605
	active transmembrane transporter activity	GO:0022804
	sulfur amino acid metabolic process	GO:0000096
2 213 transcripts	cellular amide metabolic process	GO:0043603
	organonitrogen compound biosynthetic process	GO:1901566
	non-membrane-bounded organelle	GO:0043228
	nucleosome	GO:0000786
	DNA packaging complex	GO:0044815

Group	Over-represented GO term	GO ID
1 1083 transcripts	structural constituent of ribosome	GO:0003735
	structural molecule activity	GO:0005198
	chitin synthase activity	GO:0004100
	translation	GO:0006412
	cellular amide metabolic process	GO:0043603
2 965 transcripts	RNA metabolic process	GO:0016070
	nucleic acid metabolic process	GO:0090304
	nucleobase-containing compound metabolic process	GO:0006139
	gene expression	GO:0010467
	heterocycle metabolic process	GO:0046483
3 859 transcripts	nucleic acid binding	GO:0003676
	binding	GO:0005488
	RNA binding	GO:0003723
	ribosome biogenesis	GO:0042254
	RNA metabolic process	GO:0016070
4 219 transcripts	microtubule-based movement	GO:0007018
	movement of cell or subcellular component	GO:0006928
	microtubule-based process	GO:0007017
	microtubule motor activity	GO:0003777
	dynein complex	GO:0030286
5 355 transcripts	structural constituent of ribosome	GO:0003735
	structural molecule activity	GO:0005198
	7S RNA binding	GO:0008312
	branched-chain-amino-acid transaminase activity	GO:0004084
	glucose-6-phosphate dehydrogenase activity	GO:0004345
6 156 transcripts	oxidoreductase activity, acting on the aldehyde or oxo group of donors, NAD or NADP as	GO:0016620
	oxidoreductase activity, acting on the aldehyde or oxo group of donors	GO:0016903
	response to oxidative stress	GO:0006979
	proteasome assembly	GO:0043248
	proline metabolic process	GO:0006560
7 64 transcripts	reactive oxygen species metabolic process	GO:0072593
	superoxide metabolic process	GO:0006801
	lipid metabolic process	GO:0006629
	methionyl-tRNA aminoacylation	GO:0006431
	cellular lipid metabolic process	GO:0044255