If it's there, could it be a bear?

Floe Foxon[1]

[1] Department of Data Management and Statistical Analysis, PinneyAssociates – Pittsburgh, USA

Correspondence: ffoxon@pinneyassociates.com

Abstract

It has been suggested that the American black bear (*Ursus americanus*) may be responsible for a significant number of purported sightings of an alleged unknown species of hominid in North America. Previous analyses have identified correlation between 'sasquatch' or 'bigfoot' sightings and black bear populations in the Pacific Northwest using ecological niche models and simple models of expected animal sightings. The present study expands the analysis to the entire US and Canada by regressing sasquatch sightings on bear populations in each state/province while adjusting for human population and land area in a generalized linear model. Sasquatch sightings were statistically significantly associated with bear populations such that, on the average, every $1,000$ bear increase in the bear population is associated with a 4% (95% CI: 1%–7%) increase in sasquatch sightings. Thus, as black bear populations increase, sasquatch sightings are expected to increase also. On the average, across all states and provinces in 2006, after controlling for human population and forest area, there were approximately $5,000$ bears per sasquatch sighting. Based on statistical considerations, it is likely that many supposed sasquatch are really misidentified known forms. If bigfoot is there, it may be a bear.

Keywords: Black bear; *Ursus americanus*; Sasquatch; Bigfoot; Regression

Introduction

The United States and Canada feature nearly 20 million square kilometers of land, hosting hundreds of mammal species in its woodlands, prairies, boreal forests, and along its coasts (Kays and Wilson, 2009). Due to this size and diversity, it is unlikely that the North American faunal catalogue is complete, with numerous insects and possibly small mammals remaining to be discovered. While most scientists believe that the existence of a large hominid species in North America, as-yet unrecognised in conventional science, is not likely (King and Greenwell, 1983), some authors have entertained this possibility in the field of research called 'hominology'.

Hominology, as formulated by Boris Porshnev and Dimitri Bayanov, alleges that hairy, bipedal primates not recognised in conventional zoology *do* exist and are relicts of *Homo neanderthalensis*, banished to the wild through competition with *H. sapiens* (Bayanov, 2012). One such mystery hominid reported in North America is variously dubbed 'sasquatch' after West Coast First Nations tradition (Heuvelmans, 1986), 'bigfoot' to Westerners, and the *nomen dubium Gigantopithecus canadensis* to cryptozoologists who believe that rather than a hominin, this animal is a relict Pleistocene pongid (Heaney, 1990). Purported sasquatch sightings number in the thousands (Bigfoot Field Researchers Organization, 2023), making bigfoot a prominent anthrozoological phenomenon.

Hominologists describe six lines of evidence to support the above claims (Bayanov, 2012): (1) descriptions of wild men in the natural history texts of ancient Roman and Arab philosophers and medieval Europeans; (2) folklore and mythology; (3) ancient and medieval art; (4) footprints, tracks, etc.; (5) photographs; and (6) eyewitness testimony. These are all categories of indirect evidence (i.e., testimonial and circumstantial).

Concerning (1), while some legendary animals in ancient texts were likely inspired by encounters with real forms (e.g., the Greco-Roman griffin and its probable inspiration in fossil ceratopsians (Mayor, 1991)), the existence of other beings catalogued in ancient and medieval bestiaries have not been treated with such conviction. Similarly concerning (2), while various cross-culture traditions, myths, and legends have been identified in association with mystery hominids including sasquatch (Bayanov, 2011), many characters in folklore such as magical witches and sorcerers do not correspond to counterparts in reality (Colarusso, 1983). In hominology, folklore is interpreted as the emotional expression of lived experiences (Bayanov and Bourtsev, 1976), therefore sasquatch traditions are seen as evidence for sasquatch existence. However, many of these folkloric references are extremely heterogeneous in their descriptions of mystery hominids.

Concerning (3), 'hairy man' pretroglyphs in the Tule River Tribe Reservation, approximately 1,000 years old, have been interpreted as pictorial representations of bigfoot (Strain, 2012). However, other Native American petroglyphs depicting animals such as humanoid frogs have not received such literal interpretations, which may indicate a kind of confirmation bias in hominology research.

Concerning (4), numerous casts and measurements of tracks and footprints attributed to sasquatch have been presented (Napier, 1976, 'Tables'), some apparently featuring primate-like dermal ridge patterns, sweat pores, and sole pads (Cachel, 1985; Krantz, 1983). These have been criticised as hoaxes constructed with modelling clay and the 'latex-and-kerosene expansion method' of preserving details of [human] footprints while greatly increasing their size (Baird, 1989; Bodley, 1988). It has been suggested that features such as 'sweat pores' are casting artefacts such as air bubbles (Freeland and Rowe, 1989). Genetic and microscopic analyses of supposed hairs, faeces, and other specimens attributed to sasquatch have been variously identified as synthetic fiber (Somer, 1989; Winn, 1991), or material from known forms such as cervids, bovines, and ursids (Bryant and Trevor-Deutsch, 1980; Coltman and Davis, 2005; Federal Bureau of Investigation, 2019; Hart,

2016a,b; Sykes et al., 2014).

Concerning (5), the 'Patterson-Gimlin film' is perhaps the 'best' photographic evidence for sasquatch. This notorious 16 mm motion picture purportedly depicts an unknown hominid over six feet tall in California (Discovery, 2022; Kelsey, 2022; W Munns, 2014). The film apparently was not spliced or edited (B Munns and Meldrum, 2013), but many have noted the imposing likelihood that the film subject is a suited actor.

This leaves (6), the testimony of thousands of eyewitnesses. The aim of the present study is to better understand what thousands of people see when they report a sasquatch encounter. Besides hoaxes, the leading hypothesis among skeptics is that when eyewitnesses report seeing bigfoot, they are actually seeing bears (Nickell, 2013). To understand how this can be the case, it is necessary in the following to describe the characteristics of bears in North America, and how these are similar to purported characteristics of sasquatch.

North America features three bear species, chiefly among them being the American black bear (*Ursus americanus*). The black bear is a large tetrapod (lengths up to 6 feet or more and masses up to 300 kg), with a pelage of various shades of black, brown, red, blue, blond, and white (Burton, 1998). Morphologically, the black bear is stocky with strong musculature around its thick neck, shoulders, and legs (Clark et al., 2021). The tail is short and therefore may be difficult to notice.

The geographic distribution of the black bear is widespread, ranging all across Canada and Alaska, the East and West coasts of the United States, south into Mexico, and with various isolated populations in-between (Beer and Morris, 2004). Although the present distribution is much diminished from the historical range, the black bear has experienced a notable expansion since the 1990s (Scheick and Mccown, 2014). Black bears may inhabit scrub, swamps, and mountains, but are primarily found in forest and woodland (Beer and Morris, 2004; Burton, 1998).

Black bears have opportunistic forraging habits; being omnivores, they consume various plant material (including berries, fruits, nuts, and grasses), other mammals (neonate ungulates), fish, invertebrates, and carrion (Beer and Morris, 2004). This opportunism means habituated and food-conditioned bears can and will approach people and buildings in search of anthropogenic food sources such as agriculture and garbage (Herrero, 2018, Chapter 4). Although this dietary range adaptation has allowed black bears to remain widespread (Clark et al., 2021), it is also the primary cause of confrontation between humans and bears in areas such as national parks (Herrero, 2018, Chapter 4). Black bears are easily food conditioned (Brown, 1993, Chapter 4), and because of this they are considered pests in much of North America (Burton, 1998).

The presence of the black bear is indicated by such signs as overturned logs and broken twigs and branches (Burton, 1998), overturned surface stones, tracks, hairs, and claw and tooth marks left on trees (Herrero, 2018, Chapter 4). These bears are solitary and active at night, and may be heard vocalising through grunts, growls, woofs, and howls (Beer and Morris, 2004), as well as snorts, roars, coughs, and squeaks, all indicative of emotion (Brown, 1993, Chapter 4). Black bears also have a pronounced odor (Brown, 1993, Chapter 3). They are inquisitive (Brown, 1993, Chapter 4) and intelligent animals (Vonk et al., 2012), capable of forming natural concepts at concrete, intermediate and abstract levels. Bears in general are plantigrade and are known to stand upright and walk on their hind legs; some bear species are known to ambulate over 400 meters bipedally (Brown, 1993, Chapter 4). Black bears are also strong swimmers, outstanding climbers, and can run at speeds of up to 13 meters per second (Brown, 1993, Chapter 4).

It is precisely the above characteristics of the black bear that make this animal a likely candidate for many purported sasquatch sightings (Nickell, 2013). Indeed, Brown (1993, Chapter 5) lists numerous lines of evi-

dence connecting sasquatch traditions with bears, including reported size, strength, and speed; bipedal loco- 113
motion; leaving human- or ape-like footprint outlines; tracks found and encounters occuring in bear habitats; 114
and having the same colouration. Cryptozoologist Ivan Sanderson (1967, Chapter 18) also described sasquatch 115
as "forest dwellers", sightings of which fall within the distribution of the Earth's vegetational belts (i.e., where 116
bears and bear foods are found). Other reported characteristics of sasquatch include distinctive smell and 117
various vocalisations; eyes "like a bear's"; "reddish", "dark brown", "light brown", and other fur colours; con- 118
sumption of berries and other vegetation; rolling boulders; and raiding of houses for food (Sanderson, 1967, 119
Chapters 2–4), are all consistent with black bears in North America, as described above. Of course, an alter- 120
native interpretation of these correlates is that the sasquatch is an animal distinct from the black bear but is 121
similar to it in many ways, including habitat and the characteristics described previously. 122
123

Existing studies have explored the possible link between the American black bear and bigfoot. Blight (2005) 124
used probabilistic models to examine the relationship between sasquatch sightings and black bear popula- 125
tions in the Pacific Northwest (PNW; including Alaska, Montana, Oregon, Washington, Northern California, and 126
Idaho) and the rest of the US. The method used was a simple calculation of the bivariate correlation coeffi- 127
cient between sasquatch sighting frequency and black bear population density. That study identified positive 128
correlation between black bear populations and sasquatch sightings. One criticism of this study is that the 129
test used assumed a Gaussian distribution for count data. Futhermore, only the US was considered, despite 130
sasquatch sightings being considerably more widespread across the entire US and Canada. 131
132

In another study, Lozier et al. (2009) identified a high degree of overlap in predicted geographic distribu- 133
tions for black bears and sasquatch. The method used was to compare results from ecological niche models 134
(ENMs) for each of sasquatch and black bears in the PNW. ENMs are tools used in conservation to predict ge- 135
ographic ranges of animals; they take as input georeferenced data (i.e., events with geographic coordinates) 136
and environmental data layers constructed from bioclimatic variables (e.g. annual mean temperature), and 137
output a predicted geographic distribution for the animal (Sillero et al., 2021). Although the purpose of the 138
study by Lozier et al. was the warn of the potential pitfalls of these models when errors are introduced by 139
inaccurate specimen identities and georeferencing, the analysis is supportive of a statistical association be- 140
tween sasquatch and black bears; the authors state explicitly that the predicted sasquatch distribution may 141
be seriously biased if many (or all) of the sightings represent misidentified black bears, hence the overlap. The 142
Lozier et al. analysis considered only data from the PNW (Washington, Oregon and California). 143
144

The present study builds upon the works of Blight (2005) and Lozier et al. (2009) by using more recent data, 145
by expanding the analysis to the entire US and Canada, and more appropriately assuming a count data dis- 146
tribution as opposed to a Gaussian distribution. Statistical methods (regression modelling) are used to test 147
the hypothesis that sasquatch sightings and populations of black bears are associated, while controlling for 148
potential confounding by human population and forest area. 149
150

Methods

Data

The number of black bears in each US state and Canadian province were sourced from Table 1. of Spencer et al. (2007). To the author's knowledge, this represents the latest peer-reviewed, publicly available collection of black bear population data, providing population estimates in each state/province for the year 2006. Because Delaware, Hawaii, Illinois, Indiana, Iowa, Kansas, Nebraska, North Dakota, and South Dakota have no known breeding populations of black bear, the black bear populations were assumed to be zero in these states. Other species of bear were not considered, as their impact on the result is likely to be negligible; black bears are by far the most abundant species, numbering over two times that of all other bear species combined (Garshelis et al., 2021).

The number of sasquatch sighting reports in each US state and Canadian province were sourced from the Bigfoot Field Researchers Organization (2023) Geographic Database of Bigfoot/Sasquatch Sightings & Reports. These data consist of eyewitness testimonials, mostly from the second half of the 20th century to the present. To maximise the validity of the analysis, only sightings in the year 2006 were considered such that bear populations and sasquatch sightings were date-matched in these analyses.

Human population statistics for each US state and Canadian province in the year 2006 were obtained from the United States Census Bureau (2021) and Statistics Canada (2022), respectively. Forest area estimates (in km^2) for each US state in 2006 were sourced from the US Department of Agriculture Forest Service (USDA Forest Service, 2006), and estimates for Canadian provinces were sourced from World Atlas (Sawe, 2017). For choropleth map plotting, a geojson map of the US and Canada was sourced from Cartograhy Vectors (2022).

Four US states (Rhode Island, Texas, Wisconsin, and Wyoming) and four Canadian provinces (Alberta, Newfoundland and Labrador, Northwest Territories, and Nova Scotia) were missing data on bear population. These were necessarily excluded from analyses.

Analysis

To test the hypothesis that sasquatch sightings and populations of black bears are associated, a regression model was implemented. This model regressed the number of sasquatch sighting reports in each state/province in 2006 (the dependent/response variable) on the black bear population in each state/province in 2006 (the independent/predictor variable). An intercept term was included to model the possibility of sasqautch sightings in states with no bears (i.e., to model the value of the response variable when the predictor equals zero).

Two more variables were included in the model to reduce confounding and increase the internal validity. These variables were the human population and forest area in each state/province. The more people in a given state/province, the more likely a sighting is to occur. To account for this, the human population in each state/province in 2006 was included as an independent variable in the model. Similarly, the area of forest land in each state/province is likely to impact the number of sasquatch sightings due to the relationship between area and population density, and because black bears are primarily found in forest and woodland (Beer and Morris, 2004; Burton, 1998). Consequently, the total amount of forest area (in square kilometers) in each state/province was also included as an independent variable in the model.

Because these data (number of sightings) are count data (i.e., countable quantities), it was necessary to use a model appropriate for count data. The model used was a generalized linear model assuming a Negative

Binomial distribution and the log link function, which are appropriate for count data. Variations on this model design (e.g., instead assuming Gaussian and Poisson distributions; including interaction terms and random effects; and excluding the intercept term) were investigated in previous exploratory analyses (Foxon, 2023a,b). In those exploratory analyses, model fits were assessed by the root mean square error (RMSE; lower is better) and log-likelihood (higher is better). Gaussian models provided the lowest RMSE but it is generally known that Guassian approximations do not correctly describe count data (Lass et al., 2021). Interaction terms were not statistically significant and/or provided poorer model fits to the data, and so were not included in the final model. The inclusion of random intercepts for each state/province overfit the model because only one timepoint is used (i.e., just 2006). The final model in the present study (Negative Binomial, log link, and intercept) provided the highest log-likelihood and is arguably the most appropriate model for these data for the reasons described above. Indeed, statistical models using negative binomial regression techniques describing the abundance of various animals species have been successfully developed in previous studies in the context of conservation biology, with high predictive power for some species (Acevedo et al., 2014; Pearce and Ferrier, 2001; Pradhan and Leung, 2006).

All analyses were performed in Python 3.8.16 with the packages Numpy 1.21.5, Pandas 1.5.2, Scipy 1.7.3, Statsmodels 0.13.5, and Plotly 5.9.0. All code and data are available in the online Supplementary Information (Foxon, 2023c). Statsmodels uses the iteratively reweighted least squares (IRLS) method for generalized linear models to reduce the impact of outliers.

Results

Figure 1 shows choropleth maps for the number of sasquatch sightings, black bear populations, human populations, and forest area in the United States and Canada in 2006. At first glance, there are locations such as Florida with very many sasquatch sightings but relatively few bears (at least compared to some other states/provinces). This appears to be inconsistent with the hypothesis that sasquatch sightings and populations of black bears are associated. However, this simple comparison of bears to bigfoot is confounded by the human population and forest area in each state/province. When considering the three maps of bears, people, and forest area simultaneously, the possible associations become more apparent. For example, Florida has relatively few bears and relatively little forest area, but also a relatively large number of people to make sightings. It may be that the combination of fewer bears and less forest area but more people explains the higher number of sightings. Thus, the only way to investigate the possible association between sasquatch sightings and populations of black bears is by considering the three predictors (bears, people, and forest area) simultaneously, which is the purpose of the model.

The results of the model regressing bears, people, and forest area on sasquatch sightings in 2006 are described in Table 1. In this model, black bear population was associated with sasquatch sightings such that, on the average, every $1,000$ bear increase in the bear population is associated with a 4% (95% CI: 1%–7%) increase in sasquatch sightings. Thus, as black bear populations increase, sasquatch sightings are expected to increase also. On the average, across all states and provinces in 2006, after controlling for human population and forest area, there were approximately $5,000$ bears per sasquatch sighting (note that being an average, this estimate is not accurate for each individual state/province). The RMSE of the model (19) was high relative to the number of bigfoot sightings in 2006 in each state/province (range: 0–21).

Discussion

In the present study, a model was used to investigate the hypothesis that sasquatch sightings are associated with black bear (*Ursus americanus*) populations across the US and Canada. From this model, a positive association was identified between sasquatch sightings and black bears such that, after adjusting for human population and forest area, one sasquatch sighting per year is expected for every $5,000$ bears in North America.

While correlation does not equal causation, perhaps the most parsimonious interpretation of these findings is that many supposed sasquatch sightings in North America may be explained as misidentified black bears. This is logical, because as noted by Nickell (2013) and Brown (1993, Chapter 5), bears and sasquatch share many characteristics in habitat, appearance (e.g., size, hair/fur coverage, and colouration), attributes (e.g., speed and strength), and behaviour (e.g., bipedal locomotion and curiosity). Nickel also notes that poor viewing conditions, non-expert observation, and expectant attention (i.e. 'seeing' what one anticipates) could explain why some people might confuse bears for mystery hominids. Of course, hominologists may point to the possibility that sasquatch are animals distinct from bears but which live in the same places and look and act similarly. However, in the absence of autoptical evidence (i.e., not testimonial or circumstantial), there is not at present strong scientific support for the existence of sasquatch as a mystery hominid.

Although preliminary, these findings suggest that sasquatch sightings may have some utility in bear conservation efforts as a form of citizen science. Indeed, data from amateurs/non-professionals are already in use in conservation biology/ecology with great success while also being low in cost (MacPhail and Colla, 2020; McKinley et al., 2017; Sumner et al., 2019). It may be that collaboration between bear conservationist and bigfoot field researchers could lead to more recent and accurate estimates of bear population and geographic distribution across North America.

The primary limitation of this study is the potential for residual confounding by factors other than bear population, human population, and forest area. For example, people living, hunting, or hiking in forest areas may also be misidentified as sasquatch. Furthermore, it is important to note that sasquatch sightings have been reported in states with no known breeding black bear populations. Indeed, the intercept term of the model was positive, implying that there are sasquatch sightings even when the number of bears equals zero. Although this may be interpreted as evidence for the possible existence of an unknown hominid in North America, it may also explained by misidentifiation of other animals (including humans), among other possibilities. Still, the present study does not prove that all sasquatch sightings are bear sightings (nor was this study designed for that purpose). Due to these and other limitations specified in the data sources, the findings of the present study must be interpreted as only approximate and not exact (Indeed, as noted above the RMSE was high). Further research and more data are necessary to improve these estimates.

Strengths of this study include the use of a quantitative model that controlled for possible confounding by human population size and forest area in each US state and Canadian province. Furthermore, positive associations between bear populations and bigfoot sightings were also identified in previous exploratory analyses with variations on the model design and data (Foxon, 2023a,b), which may suggest that these findings are robust. These findings are also in agreement with the results of previous studies by Blight (2005) and Lozier et al. (2009), which similarly indentified positive associations between bears and bigfoot using different methods. Therefore, the association is likely to be real. Blight reported only weak positive correlation between bears and bigfoot, whereas the association in the present study is rather more strong. This may be because Blight assumed a continuous probability distribution (as opposed to a count distribution), did not consider Canada, did not date-match the data, and did not control for forest area. The findings of the present study were closer

to those of Lozier et al., who reported a high ecological niche model overlap statistic (i.e., a strong associa- 286
tion between bear and sasquatch geographic distributions). Another stength is that in the present study, the 287
analyses considered the entire US and Canada, whereas the previously-published works only considered the 288
US or Pacific Northwest. Furthermore, the data in the present study were date-matched (in 2006) to provide 289
more accurate estimates of the association. 290
291
In conclusion, if bigfoot is there, it may be a bear. 292
293

Acknowledgements 294

The author thanks Julie Sheldon and Rahul Raveendran for helpful comments and suggestions. 295

Fundings 296

This work was not supported by any specific grant from funding agencies in the public, commercial, or 297
not-for-profit sectors. 298

Conflict of interest disclosure 299

The author declares that they have no financial conflicts of interest in relation to the content of the article. 300

Data, script, code, and supplementary information availability 301

Data and code are available online: https://doi.org/10.17605/OSF.IO/AV3G2 302

References 303

Acevedo P, F Quirós-Fernández, J Casal, and J Vicente (2014). Spatial distribution of wild boar population abun- 304
dance: Basic information for spatial epidemiology and wildlife management. *Ecological Indicators* 36, 594– 305
600. ISSN: 1470-160X. https://doi.org/10.1016/j.ecolind.2013.09.019. 306
Baird D (1989). Sasquatch Footprints: A Proposed Method of Fabrication. *Cryptozoology* 8, 43–46. 307
Bayanov D and I Bourtsev (1976). The Mysterious Biped. *Science and Religion* 39. 308
Bayanov D (2011). *Learning from Folklore* in *Russian Hominology*. Hancock House. Originally published on the 309
Bigfoot Encounters website. 310
— (2012). *Historical Evidence for the Existence of Relict Hominoids* in *Russian Hominology*. Hancock House. Orig- 311
inally published in *The Relict Hominoid Inquiry*, Idaho State University. 312
Beer AJ and P Morris (2004). *Encyclopedia of North American Mammals*. Thunder Bay Press. 313
Bigfoot Field Researchers Organization (2023). Geographic Database of Bigfoot/Sasquatch Sightings & Reports. 314
https://bfro.net/GDB/. 315
Blight EA (2005). Relationships between the American Black Bear Population and the Bigfoot Phenomenon. 316
blightinvestigations.freeservers.com/bigfootblackbearpaper.pdf. *Blight Investigations*. 317
Bodley JH (1988). Sasquatch Footprints: Can Dermal Ridges be Faked? *Northwest science.* 62. 318
Brown G (1993). *Great bear almanac*. Lyons & Burford. 319
Bryant VM and B Trevor-Deutsch (1980). Analysis of Feces and Hair Suspected to Be of Sasquatch Origin. In: 320
Manlike Monsters on Trial: Early Records and Modern Evidence. Ed. by Halpin MM and Ames MM. Vancouver 321
and London: University of British Columbia Press. Chap. 21, pp. 291–300. 322

Burton JA (1998). *Guide to Mammals of North America*. Parkgate Books. 323

Cachel S (1985). Sole Pads and Dermatoglyphics of the Elk Wallow Footprints. *Cryptozoology* 4, 45–54. 324

Cartograhy Vectors (2022). Combined US & Canada with States & Provinces. https://web.archive.org/web/ 325
20221128141742/https://cartographyvectors.com/map/793-combined-us-canada-with-states-provinces. 326

Clark JD, JP Beckmann, MS Boyce, BD Leopold, AE Loosen, and MR Pelton (2021). American Black Bear (*Ursus* 327
americanus). 328

Colarusso J (1983). Further Notes on the Role of Folklore in Hominology. *Cryptozoology* 2, 90–97. 329

Coltman D and C Davis (2005). Molecular cryptozoology meets the Sasquatch. *Trends in Ecology & Evolution* 21, 330
60–61. https://doi.org/10.1016/j.tree.2005.11.010. 331

Discovery (2022). Expedition Bigfoot Season 3 Episode 13 - A Massive Discovery. https://web.archive.org/web/ 332
20221007103940/https://www.travelchannel.com/shows/expedition-bigfoot. 333

Federal Bureau of Investigation (2019). FBI Records: The Vault - Bigfoot Part 01 of 01. https://web.archive.org/ 334
web/20191220161650/https://vault.fbi.gov/bigfoot/bigfoot-part-01-of-01/view. 335

Foxon F (2023a). If it's there, could it be a bear? (Version 1). https://www.biorxiv.org/content/10.1101/2023. 336
01.14.524058v1. *bioRxiv*. https://doi.org/10.1101/2023.01.14.524058. 337

— (2023b). If it's there, could it be a bear? (Version 2). https://www.biorxiv.org/content/10.1101/2023.01.14. 338
524058v2. *bioRxiv*. 339

— (2023c). Supplemental materials for paper: If it's there, could it be a bear? https://doi.org/10.17605/OSF. 340
IO/AV3G2. 341

Freeland D and W Rowe (1989). Alleged pore structure in Sasquatch (Bigfoot) footprints. *Skeptical Inquirer* 13, 342
273–276. 343

Garshelis DL, KV Noyce, MA Ditmer, PL Coy, AN Tri, TG Laske, and PA Iaizzo (2021). Remarkable adaptations of 344
the American Black Bear help explain why it is the most common bear: a long-term study from the center 345
of its range. 346

Hart HV (2016a). DNA as Evidence for the Existence of Relict Hominoids. *Relict Hominoid Inquiry* 5, 8–31. 347

— (2016b). Not Finding Bigfoot in DNA. *The Journal of Cryptozoology* 4, 39–51. 348

Heaney M (1990). A More Appropriate Procedure for Naming Sasquatch. *Cryptozoology* 9, 52–56. 349

Herrero S (2018). *Bear attacks: their causes and avoidance*. Rowman & Littlefield. 350

Heuvelmans B (1986). Annotated Checklist of Apparently Unknown Animals With Which Cryptozoology is Con- 351
cerned. *Cryptozoology* 5, 1–26. 352

Kays RW and DE Wilson (2009). *Mammals of North America*. Princeton University Press. https://doi.org/10. 353
1515/9781400833504. 354

Kelsey P (2022). A Forensic Analysis of the Patterson-Gimlin Film. https://web.archive.org/web/20221211054745/ 355
https://adventuresinrediscovery.com/2022/06/13/a-forensic-analysis-of-the-patterson-gimlin-film/. 356

King JE and JR Greenwell (1983). Attitudes of Biological Limnologists and Oceanographers toward Supposed 357
Unknown Animals in Loch Ness. *Cryptozoology* 2, 98–102. 358

Krantz GS (1983). Anatomy and Dermatoglyphics of Three Sasquatch Footprints. *Cryptozoology* 2, 53–81. 359

Lass J, ME Bøggild, P Hedegård, and K Lefmann (2021). Multinomial, Poisson and Gaussian statistics in count 360
data analysis. *Journal of Neutron Research* 23, 69–92. 361

Lozier J, P Aniello, and M Hickerson (2009). Predicting the distribution of Sasquatch in western North America: 362
anything goes with ecological niche modelling. https://doi.org/10.1111/j.1365-2699.2009.02152.x. 363

MacPhail VJ and SR Colla (2020). Power of the people: A review of citizen science programs for conservation. 364
Biological Conservation 249, 108739. ISSN: 0006-3207. https://doi.org/10.1016/j.biocon.2020.108739. 365

Mayor A (1991). Griffin Bones: Ancient Folklore and Paleontology. *Cryptozoology* 10, 16–41. 366

McKinley DC, AJ Miller-Rushing, HL Ballard, R Bonney, H Brown, SC Cook-Patton, DM Evans, RA French, JK Par- 367
rish, TB Phillips, SF Ryan, LA Shanley, JL Shirk, KF Stepenuck, JF Weltzin, A Wiggins, OD Boyle, RD Briggs, SF 368
Chapin, DA Hewitt, PW Preuss, and MA Soukup (2017). Citizen science can improve conservation science, 369
natural resource management, and environmental protection. *Biological Conservation* 208. The role of citi- 370

zen science in biological conservation, 15–28. ISSN: 0006-3207. https://doi.org/10.1016/j.biocon.2016.05.015.

Munns B and J Meldrum (2013). Analysis Integrity of the Patterson-Gimlin Film Image. *The Relict Hominoid Inquiry* 2, 41–80.

Munns W (2014). *When Roger Met Patty*. Self-published: CreateSpace.

Napier J (1976). *Bigfoot - The Yeti and Sasquatch in Myth and Reality*. Sphere Books Ltd.

Nickell J (2013). Bigfoot Lookalikes: Tracking Hairy Man-Beasts. *Skeptical Inquirer* 37.

Pearce J and S Ferrier (2001). The practical value of modelling relative abundance of species for regional conservation planning: a case study. *Biological Conservation* 98, 33–43. ISSN: 0006-3207. https://doi.org/10.1016/S0006-3207(00)00139-7.

Pradhan NC and P Leung (2006). A Poisson and negative binomial regression model of sea turtle interactions in Hawaii's longline fishery. *Fisheries Research* 78, 309–322. ISSN: 0165-7836. https://doi.org/10.1016/j.fishres.2005.12.013.

Sanderson IT (1967). *Abominable Snowmen: Legend Come to Life*. Chilton.

Sawe BE (2017). Forest Land By Canadian Province And Territory. https://web.archive.org/web/20221206040420/https://www.worldatlas.com/articles/forest-land-by-canadian-province-and-territory.html.

Scheick B and J Mccown (2014). Geographic distribution of American black bears in North America. *Ursus* 25, 24–33. https://doi.org/10.2192/URSUS-D-12-00020.1.

Sillero N, S Arenas-Castro, U Enriquez-Urzelai, CG Vale, D Sousa-Guedes, F Martínez-Freiría, R Real, and A Barbosa (2021). Want to model a species niche? A step-by-step guideline on correlative ecological niche modelling. *Ecological Modelling* 456, 109671. https://doi.org/10.1016/j.ecolmodel.2021.109671.

Somer L (1989). Morphological Analysis of Possible Sasquatch Hair Recovered from the Blue Mountains of Washington State. *Sasquatch Evidence: Scientific and Social Implications Symposium. Eight Annual Membership Meeting, International Society of Cryptozoology, June 24-25, 1989, Washington State University, Pullman, Washington*.

Spencer RD, RA Beausoleil, and DA Martorello (2007). How agencies respond to human–black bear conflicts: a survey of wildlife agencies in North America. *Ursus* 18, 217–229.

Statistics Canada (2022). Population and private dwellings occupied by usual residents and intercensal growth, 1851 to 2021 censuses. https://web.archive.org/web/20221128125153/https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=9810001701.

Strain KM (2012). Mayak Datat: The Hairy Man Pictographs. *Relict Hominoid Inquiry* 1, 1–12.

Sumner S, P Bevan, AG Hart, and NJ Isaac (2019). Mapping species distributions in 2 weeks using citizen science. *Insect Conservation and Diversity* 12, 382–388. https://doi.org/10.1111/icad.12345.

Sykes BC, RA Mullis, C Hagenmuller, TW Melton, and M Sartori (2014). Genetic analysis of hair samples attributed to yeti, bigfoot and other anomalous primates. *Proceedings of the Royal Society B: Biological Sciences* 281, 20140161. https://doi.org/10.1098/rspb.2014.0161.

United States Census Bureau (2021). State Intercensal Tables: 2000-2010. https://web.archive.org/web/20230327112644/https://www.census.gov/data/tables/time-series/demo/popest/intercensal-2000-2010-state.html.

USDA Forest Service (2006). Forest Inventory and Analysis Fiscal Year 2006 Business Report. https://web.archive.org/web/20230314130101/https://www.fia.fs.usda.gov/library/bus-org-documents/index.php.

Vonk J, SE Jett, and KW Mosteller (2012). Concept formation in American black bears, Ursus americanus. *Animal Behaviour* 84, 953–964.

Winn EB (1991). Physical and Morphological Analysis of Samples of Fiber Purported to be Sasquatch Hair. *Cryptozoology* 10, 55–65.

Figure 1. Choropleth maps for sasquatch reports, black bear (*Ursus americanus*) populations, human populations, and forest area across the United States and Canada.

Table 1. Regression Model Parameters

Variable	Regression Coefficient $\pm$ Standard Error
Intercept	0.5 ± 0.2 *
Black Bear Population	$(4 \pm 2) \times 10^{-5}$ *
Human Population	$(1.2 \pm 0.2) \times 10^{-7}$ **
Forest Area (km^2)	$(-7 \pm 3) \times 10^{-6}$ *

* $p < 0.05$, ** $p < 0.0001$