Improved cognitive-motor processing speed and decreased functional connectivity after
high intensity aerobic exercise in individuals with chronic stroke.

Justin W. Andrushko[1,2], Shie Rinat[1,3], Brian Greeley[1,2], Beverley C. Larssen[1,2], Christina B.
Jones[1,3], Cristina Rubino[1,3], Ronan Denyer[1,4], Jennifer Ferris[1,3], Kristin L. Campbell[2], Jason L.
Neva[5,6], Lara A. Boyd[1,2]

1. Brain Behaviour Laboratory, Vancouver British Columbia, Canada, V6T 1Z3
2. Faculty of Medicine, Department of Physical Therapy, University of British Columbia –
Vancouver, British Columbia, Canada, V6T 1Z3
3. Faculty of Medicine, Graduate Program in Rehabilitation Sciences, University of British
Columbia, Vancouver, British Columbia, Canada, V6T 1Z3
4. Faculty of Medicine, Graduate Program in Neuroscience, University of British Columbia,
Vancouver, British Columbia, Canada, V6T 1Z3
5. Faculty of Medicine, School of Kinesiology and Physical Activity Sciences, University
of Montreal, Montreal, Quebec, Canada, H3T 1J4
6. Research Center of the Montreal Geriatrics Institute (CRIUGM), Montreal, QC, Canada

Correspondance

Lara A. Boyd, Ph.D.
Email: lara.boyd@ubc.ca
ORCID: https://orcid.org/0000-0002-2828-4549
Department of Physical Therapy
The University of British Columbia
22

Number of pages: 37

Number of figures: 7

Number of tables: 1

Number of words:

• **Abstract:** 250
• **Introduction:** 688
• **Discussion:** 1236

Conflict of interest statement

The authors have no conflicts to declare.

Data sharing

All data will be available from the corresponding author upon reasonable request.

Keywords

Chronic stroke, functional connectivity, high intensity aerobic exercise, trail making test,
cognitive function

Significance statement

After stroke, impaired motor performance is linked to an increased demand for cognitive

resources. In our work we show that high intensity aerobic exercise paired with an implicit motor

learning task improves cognitive-motor processing speed and reduces resting-state functional

connectivity between the dorsolateral prefrontal cortex and the sensorimotor network in

individuals living with chronic stroke. These data likely reflect a reduction in cognitive resource

dependence during a cognitive-motor task after stroke and a shift towards cognitive-motor

automaticity.

Abstract

After stroke, impaired motor performance is linked to an increased demand for cognitive

resources. Aerobic exercise improves cognitive function in healthy populations and may be

effective in altering cognitive function post-stroke. We sought to determine if high intensity

aerobic exercise paired with motor training in individuals with chronic stroke alters cognitive-

motor function and functional connectivity between the dorsolateral prefrontal cortex (DLPFC),

a key region for cognitive-motor processes, and the sensorimotor network. Twenty-five

participants with chronic stroke were randomly assigned to exercise (n = 14; 66 ± 11 years; 4

females), or control (n = 11; 68 ± 8 years; 2 females) groups. Both groups performed five-days

of paretic upper limb motor training after either high intensity aerobic exercise (3 intervals of 3

minutes each, total exercise duration of 23-minutes) or watching a documentary (control).

Resting-state fMRI, and TMT-A and B were recorded pre- and post-intervention. Both groups

showed implicit motor sequence learning ($p < .001$), but there was no added benefit of exercise

($p = .738$). Regardless of group, the changes in task score ($p = .025$), and dwell time ($p = .043$)

were correlated with a decrease in DLPFC-sensorimotor network functional connectivity ($p =$

.024), which is thought to reflect a reduction in the cognitive demand and increased automaticity.

The exercise group experienced greater overall cognitive-motor improvements measured with

the trail making test part A (TMT-A: task score: $p = .012$; dwell time: $p = .024$; movement time:

$p = .567$). Aerobic exercise may improve cognitive-motor processing speed post-stroke.

Introduction

Roughly 15 million people experience a stroke each year (Mittmann *et al.*, 2012). Stroke

is the second most common cause of death globally and one of the leading causes of severe, adult

disability (Katan & Luft, 2018). Due to advancements in preventive care, rates of stroke declined

between 1990 and 2016, yet the number of individuals that survive and live with severe disability

nearly doubled during that same timeframe (Lindsay *et al.*, 2019). Identifying methods to

enhance recovery from stroke to improve or maintain independence of living is an important and

persistent research objective.

After stroke, cognitive impairment may interact with or influence motor recovery.

Greater cognitive resources are needed to successfully plan and execute voluntary movements

after stroke (Puh *et al.*, 2007). This impaired cognitive demand is observed through an increase

in cortical activity in prefrontal areas including the dorsolateral prefrontal cortex (DLPFC) (Puh

et al., 2007; Meehan *et al.*, 2011; Li *et al.*, 2014), and is often associated with worse motor

function (Meehan *et al.*, 2011; Li *et al.*, 2014; Lin *et al.*, 2021; Hall *et al.*, 2021). The DLPFC is

an important brain region involved in several cognitive-motor processes (i.e., cognitive processes

involved in cognitively demanding motor tasks) including processing speed (Hillary *et al.*, 2006;

Kaller *et al.*, 2011), response selection (Boyd *et al.*, 2009*b*), and task switching (Badre &

Wagner, 2004; Hart *et al.*, 2013; Brunoni & Vanderhasselt, 2014). Importantly, past work from

our lab showed that despite an equal dose of motor practice and subsequent learning, individuals

with chronic stroke were unable to shift cortical activity away from the prefrontal cortex during a

motor learning task (rather, an increase was observed), while age matched healthy controls did

(Meehan *et al.*, 2011). This shift in cognitive resources away from the DLPFC during cognitive-

motor tasks has been observed in healthy cohorts with enhanced automaticity coinciding with a

decrease in functional connectivity between the DLPFC and the sensorimotor network (Mazzoni,

2008; Wu *et al.*, 2008). Therefore, interventions that can reduce the cognitive demand and

DLPFC activity in individuals with stroke may improve motor performance.

Aerobic exercise can alter patterns of brain activity, including resting-state functional

connectivity between brain networks as measured by functional magnetic resonance imaging

(fMRI) (Weng *et al.*, 2017; Greeley *et al.*, 2021), and is known to improve cognitive function in

healthy older adults (Barnes, 2015), and in individuals with stroke (Zheng *et al.*, 2016). High

intensity aerobic exercise in particular can also improve motor skill acquisition in neurologically

intact people (Skriver *et al.*, 2014; Stavrinos & Coxon, 2017; Dal Maso *et al.*, 2018; Kendall *et*

al., 2020), and individuals with chronic stroke (Nepveu *et al.*, 2017). These improvements are

linked to a decrease in the neurotransmitter gamma-aminobutyric acid (Singh *et al.*, 2014; Singh

& Staines, 2015; Stavrinos & Coxon, 2017; Hendy *et al.*, 2022), and an increase in the protein

brain-derived neurotrophic factor (Sleiman *et al.*, 2016), both of which play important roles in

neuroplasticity (Stagg *et al.*, 2011; Mang *et al.*, 2013; Andreska *et al.*, 2020). Therefore, beyond

the potential cognitive benefits associated with high intensity aerobic exercise, pairing it with

motor rehabilitation is a promising method to enhance cognitive-motor function.

The purpose of the current study was to determine if high intensity aerobic exercise

paired with skilled motor training alters cognitive-motor function and DLPFC-sensorimotor

network functional connectivity in individuals living with chronic stroke. To address these

questions participants were tested using the cognitive-motor assessments Trail Making Test parts

A (TMT-A) and B (TMT-B). The Trail Making Tests are commonly used to assess executive

function and can measure processing speed with TMT-A, and mental flexibility and task

switching with TMT-B (Kortte *et al.*, 2002; Bowie & Harvey, 2006; Gläscher *et al.*, 2012;

MacPherson *et al.*, 2017). Additionally, resting-state fMRI scans were acquired to measure

change in functional connectivity. We hypothesized that high intensity aerobic exercise paired

with skilled motor training would enhance cognitive-motor performance as measured with TMT-

A and TMT-B in individuals living with chronic stroke, and these changes would be correlated

with a decrease in DLPFC-sensorimotor network functional connectivity, reflecting a reduction

in cognitive resources needed to perform the tasks.

Methods

Ethical approval

This study conformed to the standards set by the Declaration of Helsinki and was

approved by the University of British Columbia Clinical Research Ethics Board: #H16-01945.

Participants

Individuals living with chronic stroke (> 6 months post-stroke) were recruited to

participate in this study if they had an ischemic or hemorrhagic stroke and were between the ages

of 21-85 years old. In addition, to participate individuals had to score > 23 on the Montréal

Cognitive Assessment (MoCA) (Nasreddine *et al.*, 2005), and be cleared by a cardiologist for

safe participation in an exercise protocol after performing a supervised stress-test. Eligible

participants were randomly allocated to either an exercise or control group. Both groups

performed the same motor training intervention immediately following either aerobic exercise or

rest. The data in this manuscript come from a large study on the impact of exercise on behaviour,

brain function, and physiology in individuals with stroke. The data reported here are a subset of

the larger study (Greeley *et al.*, 2021, 2023; Neva *et al.*, 2022).

Experimental design

After obtaining informed and written consent, participants performed a graded maximal

exercise stress test to determine their eligibility to participate in the intervention aspects of this

study. Testing was done before (pre-testing) and after (24h-post) five-days of skilled motor

training using the paretic arm paired with either high intensity aerobic exercise (exercise group),

or rest (control group). Resting-state fMRI, TMT-A, TMT-B, Fugl-Meyer Assessment of Motor

Recovery (FMA) (Fugl Meyer *et al.*, 1975), and the Wolf Motor Function Test (WMFT) (Wolf

et al., 2001) were completed at pre-testing and 24h-post testing time points to determine brain -

behaviour relationships (Figure 1).

Stress test

The graded maximal exercise stress test was administered by a cardiology technician.

Electrocardiogram electrodes were used for continuous heart rate monitoring throughout the

exercise protocol. Prior to the test participant lay supine for approximately three minutes, after

which resting heart rate (HR), blood pressure (BP) and rating of perceived exertion (RPE) were

recorded. Next, the participant was seated in an upright recumbent bike (SCIFIT, Tulsa,

Oklahoma, USA), that was adjusted to fit the participant. During the stress test, HR and RPE

were recorded every minute and BP was recorded every two minutes. Participants were

instructed to maintain a cadence between 50-80 revolutions per minute (RPM) and that dropping

below 50 RPM would terminate the test. The stress test began with a two-minute warm-up at 10

Watts (W) of resistance. Following the warm up, the wattage was increased by 5, 10, or 15 W

depending on subjective observation of performance (Beltz *et al.*, 2016). The resistance was

increased every minute until the participant was unable to maintain a 50 or greater RPM cadence

or when volitional fatigue was reached. Once the participant reached the termination criteria, the

resistance was dropped back to 10 W for approximately three minutes as a cool down. Next, the

participant moved back to the supinated resting position until their HR, BP, and RPE recovered

back to baseline levels.

Groups

High intensity aerobic exercise

Each exercise session was completed on an upright recumbent bike (SCIFIT, Tulsa,

Oklahoma, USA). Each session started with a five-minute warm-up at 10 watts. Following the

warm-up, the participants performed three, three-minute intervals of cycling at 75% of their

maximum power output, based on the maximum power output achieved during the final fully

completed minute during the exercise stress test. Each interval was separated by three-minutes of

low intensity cycling against 10 watts of resistance. The total duration of each aerobic exercise

session lasted 23-minutes. BP, HR, and RPE were recorded every three-minutes until the end of

the protocol. Immediately following the exercise protocol, participants proceeded with motor

training.

Control

Participants allocated to the control group watched a Planet Earth documentary for 23-

minutes immediately prior to engaging in motor training each session. Heart rate was recorded

every three-minutes throughout the video.

Motor training

The serial targeting task was employed as the motor training task (Brodie *et al.*, 2014;

Mang *et al.*, 2016; Greeley *et al.*, 2021). Participants used their paretic arm to control a

frictionless manipulandum to move a cursor between a start position and an end target projected

by the Kinarm end-point robot (Kinarm, Kingston, ON, Canada). Targets appeared one at a time;

as soon as participants finished movement to one target, they were required to hold that

positioning for 500 ms for the next appeared. Participants had 10,000 ms to reach each target. A

repeated six-element sequence of movements was embedded between seven-element random

sequences. The inclusion of both sequences allows us to separate improvements in motor

learning (repeated sequence) from those associated with motor control (random sequences)

(Boyd *et al.*, 2009*a*). In each of the five-training-sessions, participants practiced four-blocks of

the serial targeting task (444 movements per session).

Data were analyzed using an exponential curve-fitting algorithm that enables

parameterization of motor data across practice (Wadden *et al.*, 2017, 2019). Motor learning

related change was characterized by fitting behavioural data to an exponential equation:

$$E(RT_N) = A + Be^{-\alpha N}$$

211   where **RT** is reaction time, **A** is predicted asymptote in performance, **B** is the performance

212   change score to predicted asymptote, **α** is rate of change and **N** is number of practice trials

213   (Brown & Heathcote, 2003). Our dependent measure of motor learning was a change score (**B**)

214   extracted from individual learning curves for each individual's practice sessions by group

215   (exercise, control).

Cognitive-motor testing

To assess cognitive-motor performance TMT-A and TMT-B were performed by the

participants with their less-affected arm, or in scenarios where there were bilateral lesions

participants were instructed to use their preferred arm on a Kinarm end-point robot. The use of

the less affected limb allowed us to assess cognitive performance without stroke related motor

impairment affecting responses. TMT-A involves connecting dispersed numbered targets in

ascending numerical order from '1' to '25' with linear reaching movements. The objective is to

locate the next number and reach to it without touching any other numbers or connected lines

from the already completed reaches. This test assesses cognitive-motor processing speed, in

which the participant must visually navigate, identify the appropriate target, and execute a

movement as fast and accurately as possible (Corrigan & Hinkeldey, 1987; Bowie & Harvey,

2006).

TMT-B is similar to TMT-A in that it also involves searching and connecting targets in

ascending order. However, TMT-B differs in that the task must be completed in an alternating

numeric and alphabetic sequence, where the number '1' must connect to the letter 'A', then 'A'

must connect to the number '2', until the final target number '13' is reached, still equaling 25

total targets. The added complexity stresses the cognitive system and requires mental flexibility

and task switching to complete. For TMT A and B, different sequences were used at pre-testing

and 24h-post to avoid any possible learning effects.

Three metrics of task performance were assessed for TMT-A and TMT-B. First, to assess

overall performance the task score at each time point was used. This metric provides a global

measure of an individual's performance. Specifically, the task score measures deviations from an

individual's best performance. Task scores are always positive with zero representing best

performance and deviations from zero reflecting poorer performance (See Dexterit-E Explorer

manual for more information - https://kinarm.com/support/user-guides-documentation/).

Additionally, to separate the cognitive and the motor aspects of these assessments the total dwell

time (i.e., the time that the participant remains on a target while they visually search and plan

their next movement) was subtracted from the total task time to isolate the movement time. Then

statistical analyses were carried out on the movement time and dwell time separately to

determine the impact of the intervention on the cognitive and motor aspects of task performance.

Clinical assessments

Paretic arm impairment was quantified using the upper extremity portion of the Fugl-

Meyer Assessment (FMA; 0-66; higher scores indicate less paretic arm impairment) (Lin *et al.*,

2004). The 17-item version of WMFT was used to characterize arm motor function (Wolf *et al.*,

2001). The WMFT contains 15 timed movement tasks. For each WMFT task, the rate

(repetitions/60 seconds, with a rate of zero recorded if no repetitions were completed within 120

seconds) was calculated to characterize functional impairment (Hodics *et al.*, 2012); higher

scores reflect a faster movement rate and thus greater motor function. All assessors for FMA and

WMFT were trained physical or occupational therapists, or a clinical student in training.

Magnetic resonance imaging

Magnetic resonance imaging acquisition

Participants received structural and functional brain scans on a Philips Achieva 3 tesla or

a Philips Elition 3 tesla MRI. At both testing time points a T1-weighted (T1w) structural brain

scan (TR = 8.1 ms, TE = 3.61 ms, flip angle = 8°, 1mm^3 isotropic voxels, field of view = 256 $\times$

256 $\times$ 165mm field of view, total scan time = 6.4 minutes), and a resting-state fMRI scan (TR =

2.000 ms, TE = 30 ms, flip angle = 90°, 120 volumes, voxel dimensions = 3 $\times$ 3 $\times$ 3 mm with a 1

mm gap, total scan time = 4 minutes) were acquired. During resting-state fMRI scans

participants were asked to look at a fixation cross, to think of nothing and stay awake.

Anatomical data preprocessing

Anatomical T1w images were preprocessed using the fMRIprep pipeline (v22.0.0)

(Esteban *et al.*, 2018). Briefly, for each participant, the two T1w images from pre-testing and

24h-post were first corrected for intensity inhomogeneity using N4BiasFieldCorrection (Tustison

et al., 2010) as part of ANTs (v2.3.1). Next, both T1w images were used to create a participant

specific average template using mri_robust_template (Reuter *et al.*, 2010) from FreeSurfer

(v6.0.1), followed by skull-stripping with antsBrainExtraction.sh using OASIS as a target

template.

Lesion masking

For lesions, a binary mask was created by manually evaluating the T1w images from

session one and drawing a mask over the lesioned tissue in 3D space using ITK-Snap (v3.8.0).

These binary lesion masks were used by fMRIprep to assist in the registration steps (Figure 2 for

lesion mask overlap).

Resting-state fMRI data preprocessing

Resting-state fMRI data were initially minimally preprocessed in native space using

fMRIprep (v22.0.0) (Esteban *et al.*, 2018) to carry out fieldmap-less susceptibility distortion

correction (Wang *et al.*, 2017). Next, MELODIC (v3.15) as part of the FMRIB Software Library

(FSL v6.0.3; (Jenkinson *et al.*, 2012)) was used to carry out motion correction (MCFLIRT;

(Jenkinson *et al.*, 2002)), high-pass temporal filtering at 0.01Hz, and the decomposition of the

functional runs into independent components for denoising. Following MELODIC

preprocessing, FMRIB's Independent Component Analysis (ICA)-based Xnoiseifier (ICA-FIX)

was used to automatically denoise the data (Griffanti *et al.*, 2014; Salimi-Khorshidi *et al.*, 2014).

A custom training-weight was created on a subset of the present study's data (20 runs total: 10

runs from pre-testing and 10 runs from 24h-post testing) and used with a threshold value of 20 to

denoise the data. Finally, after the automated denoising, data were smoothed with a 5 mm full-

width half maximum (FWHM) kernel and registered to the MNI152_T1_1mm standard space

template included in FSL.

Resting-state fMRI data analysis

After preprocessing, a group-level functional connectivity analysis was carried out in

FSL's melodic command-line tool with a dimensionality constraint of 11 group-level

components. The constraint of 11 components was selected after multiple cross-correlation

analyses between the spatial maps of several different ICA constraints ranging from 5-15

components with the BrainMap 10-ICA template (Smith *et al.*, 2009). Dual regression was then

performed to estimate a version of the group-level spatial maps for each participant and run

(Nickerson *et al.*, 2017). The component that best represented the sensorimotor network was then

used to constrain a seed-based functional connectivity analysis. A right DLPFC mask was

extracted from the Sallet dorsal frontal connectivity-based parcellation (Sallet *et al.*, 2013), and

used as a seed region for this analysis.

Statistical inference was determined with a four-contrast general linear model (GLM) that

compared pre- and post-testing rs-fMRI scans for the exercise group (pre > post, and post > pre),

and the control group (pre > post, and post > pre). Non-parametric permutation testing with 5000

permutations for each contrast was carried out using Permutation Analysis of Linear Models

(PALM) with family-wise-error-rate corrected contrasts and cluster-extent based thresholding

with a z-score of 3.1 (Winkler *et al.*, 2014).

Statistical analysis

To test our hypotheses that high intensity aerobic exercise would enhance less-affected

upper limb cognitive-motor performance in individuals living with chronic stroke, as measured

with TMT-A and TMT-B, separate mixed repeated measures analysis of variance (RM-ANOVA)

tests were performed when data met parametric assumptions. For RM-ANOVA tests, group was

entered as a between factor, and time was entered as a within factor variable. When an

interaction was significant, a priori pairwise comparisons were performed. For these pairwise

comparisons, independent samples t-tests, and paired-sampled t-tests were used to investigate

differences between groups at each time point, and between time points within each group

respectively. Additionally, where significant effects were observed, secondary exploratory

analyses were performed to assess sex differences with no a-priori hypotheses. Partial eta^2 (η_p^2)

effect sizes were reported for all interactions and main effects. Data normality was assessed with

Shapiro-Wilk's tests, and homogeneity of variance was assessed with Levene's tests.

When data did not meet the appropriate parametric testing assumptions of normality or

heterogeneity of variance, Mann-Whitney U tests were used to assess between group differences,

and Wilcoxon signed-rank tests were used to analyze paired-sample within group data. Rank-

Biserial Correlations were reported as the effect sizes for non-parametric tests.

For all parametric and non-parametric tests, a manual Bonferroni adjusted alpha-level (α

= .0125) was then used to reduce family-wise-error rates. To improve clarity in statistical

reporting, the uncorrected p-values from the individual pairwise tests from the parametric and

non-parametric comparisons were Bonferroni adjusted by multiplying the uncorrected p-value by

the number of comparisons within a given variable (4 tests; 2 within group, and 2 between group

comparisons). This is a mathematically equivalent approach to adjusting the alpha-level

threshold by dividing it by the number of tests (i.e., α = .0125; α = .05/4 tests), and is the same

approach used for Bonferroni post-hoc testing in statistical software packages

(https://www.ibm.com/support/pages/calculation-bonferroni-adjusted-p-values). This approach

allows significance interpretation to remain at α = .05. Finally, Spearman's Rho (ρ) correlations

were used to test our hypothesis that behavioural changes would relate to a reduction in

functional connectivity between the DLPFC and the sensorimotor network. Statistical analyses

were performed in JASP (v0.16.4.0) (Love *et al.*, 2019).

Results

Participants

A total of 41 individuals with stroke consented to participate in this study, however, the

exercise stress test revealed abnormalities in three individuals, while two others were ineligible

due to low MoCA scores. Eligible participants were randomly assigned to an exercise or control

group. Of the remaining 36 individuals, two had missing MRI data, and nine had excessive head

motion exceeding a mean framewise displacement greater than .5 mm during MRI scans, thereby

rendering at least one of their time points unusable. Therefore, a total of 25 participants were

included in this study (Table 1).

Table 1. Demographics

Group	Age	Sex	Affected hemisphere	Stress test Max watts	MoCA	FMA		WMFT	
						Pre-testing	24h-post	Pre-testing	24h-post
Exercise n = 14	66 ± 11	F: n = 4 M: n = 10	Left: n = 5 Right: n = 9	83 ± 32	26 ± 2	52 ± 16	52 ± 16	48 ± 27	52 ± 36
Control n = 11	68 ± 8	F: n = 2 M: n = 9	Left: n = 6 Right: n = 5	73 ± 31	25 ± 2	53 ± 12	54 ± 11	37 ± 17	42 ± 20

MoCA = Montréal Cognitive Assessment; FMA = Fugl-Meyer Assessment; WMFT = Wolf Motor Function Test

Clinical assessments

Separate group × time RM-ANOVA tests for FMA and WMFT were assessed. The

ANOVA failed to detect a significant group × time interaction in FMA score [$F(1, 22) = .286$, p

$= .598$, $\eta_p^2 = .013$], nor main effects of time [$F(1, 22) = .750$, $p = .396$, $\eta_p^2 = .033$], or group

$[F(1, 22) = .132, p = .719, \eta_p^2 = .006]$. Similarly, a group × time interaction $[F(1, 20) = .017, p =$

$.897, \eta_p^2 < .001]$, and main effects of time $[F(1, 20) = 3.391, p = .080, \eta_p^2 = .145]$, and group

$[F(1, 20) = 1.052, p = .317, \eta_p^2 = .050]$ were not significant for the mean rate of performance

from the WMFT.

Serial targeting task

For the B values there was a violation of normality for the exercise group as assessed

with Shapiro-Wilk's test $(W = .513, p < .001)$. Therefore, a non-parametric Mann-Whitney U

test was used to assess between group differences. There was no statistically significant

difference between exercise (Mean ± SD; .364 ± .352 B value) and control (.315 ± .122 B value)

groups $(W = 71, p = .738$, Rank-Biserial Correlation = .092). However, both groups learned the

task, and improved their motor performance throughout the intervention, as evidenced by a one-

sample Wilcoxon signed-rank test, which indicates that the B values were significantly different

from zero (V = 276, $p < .001$, Rank-Biserial Correlation = 1.000; Figure 3).

Trail making test part A

Task score

For TMT-A, a RM-ANOVA test revealed a significant group × time interaction $[F(1, 22)$

$= 9.257, p = .006, \eta_p^2 = .296]$, and a significant main effect for time $[F(1, 22) = 9.269, p = .006,$

$\eta_p^2 = .296]$, but the main effect of group was not significant $[F(1, 22) = .761, p = .393, \eta_p^2 =$

$.033]$. Bonferroni adjusted post-hoc testing revealed that the interaction was influenced by a

significant pre- to 24h-post testing difference for the exercise group only (-.673 ± .150, $t = 4.496$,

$p = .012$). No other post-hoc tests were statistically significant (all $p > .180$; Figure 4).

Since an effect was only observed for the exercise group, only it was used to investigate

sex differences with a sex (male, female) × time RM-ANOVA. For this analysis, only the sex ×

time interaction and the main effect of group were reported given the main effect of time is

redundant with what has already been reported in the previous analyses. The RM-ANOVA test

did not detect a sex × time interaction [$F(1, 11) = .104$, $p = .753$, $\eta_p^2 = .009$], or a main effect of

group [$F(1, 11) = .052$, $p = .824$, $\eta_p^2 = .005$].

Dwell time

For TMT-A dwell time, there was a homogeneity of variance violation (24h-post, $p =$

.05), and therefore the higher order RM-ANOVA was not assessed. Mann-Whitney U tests for

between group comparisons at pre-testing ($W = 48$, $p = .744$, Rank-Biserial Correlation = -.329)

and 24h-post ($W = 93$, $p = .912$, Rank-Biserial Correlation = .301) were not significant. A

Wilcoxon signed-rank test revealed a significant decrease in dwell time for the exercise group (W

$= 83$, $z = 2.621$, $p = .024$, Rank-Biserial Correlation = .824), but not for the control group ($W =$

17, $z = -1.423$, $p = .700$, Rank-Biserial Correlation = -.485; Figure 4).

Once again, only an effect was observed for the exercise group, and therefore only the

exercise group data were examined for sex differences. Separate Mann-Whitney U tests at pre-

testing ($W = 16$, $p > .999$, Rank-Biserial Correlation = .067) and 24h-post ($W = 17$, $p > .999$,

Rank-Biserial Correlation = .133) failed to detect any differences between sexes. Although

separate Wilcoxon signed rank tests for males ($W = 53$, $z = 2.599$, $p = .024$, Rank-Biserial

Correlation = .927) and females ($W = 4.00$, $z = .535$, $p > .999$, Rank-Biserial Correlation = .333)

revealed that only males improved their processing speed after exercise.

Movement time

For TMT-A, a RM-ANOVA revealed a significant main effect of time [$F(1, 22) = 4.629$,

$p = .043$, $\eta_p^2 = .174$] indicating that both groups decreased their movement speed over the course

of the intervention. However, the group × time interaction [$F(1, 22) = .339$, $p = .567$, $\eta_p^2 = .015$]

and the main effect of group [$F(1, 22) = 1.563$, $p = .224$, $\eta_p^2 = .066$] were not statistically

significant. These findings suggest that there were no differences between groups in change in

movement speed after five-days of motor training (Figure 4).

Trail making test part B

Task score

For TMT-B Task score, normality ($W = .871$, $p = .005$), and homogeneity of variance

(24h-post, $p = .004$) were violated. Therefore, non-parametric Mann-Whitney U tests were used

to determine there were no between group differences at pre-testing ($W = 95.00$, $p = .744$, Rank-

Biserial Correlation $= .329$), and 24h-post ($W = 78.00$, $p > .999$, Rank-Biserial Correlation $=$

.091) timepoints. Separate Wilcoxon signed-rank tests were used to determine that there were

also no within group differences between pre-testing and 24h-post testing for the exercise group

($W = 33.00$, $z = -.874$, $p > .999$, Rank-Biserial Correlation $= -.275$) or the control group ($W =$

52.00, $z = 1.689$, $p = .408$, Rank-Biserial Correlation $= .576$; Figure 5).

Dwell time

For TMT-B, the change values from pre-testing to 24h-post were not normally distributed

based on Shapiro-Wilk's test of normality ($W = .905$, $p = .027$). Therefore, non-parametric

Mann-Whitney U tests were performed on pre- and 24h-post testing time points between groups,

and these analyses failed to detect differences at pre-testing ($W = 97.00$, $p = .600$, Rank-Biserial

Correlation $= .357$), and 24h-post ($W = 63.00$, $p > .999$, Rank-Biserial Correlation $= -.119$).

Finally, separate Wilcoxon signed-rank tests suggest that both the exercise ($W = 33.00$, $z = -.874$,

$p > .999$, Rank-Biserial Correlation = -.275) and control ($W = 58.00$, $z = 2.223$, $p = .096$, Rank-

Biserial Correlation = .758) groups did not reduce their dwell time from pre-testing to 24h-post

timepoints (Figure 5).

Movement time

For TMT-B, the change values from pre-testing to 24h-post were not normally distributed

based on Shapiro-Wilk's test of normality ($W = .762$, $p = .003$), and Levene's homogeneity of

variance tests were violated at pre-testing ($p = .019$) and 24h-post ($p = .050$). Therefore, non-

parametric Mann-Whitney U tests were carried out to assess between group differences at pre-

testing ($W = 107.00$, $p = .164$, Rank-Biserial Correlation = .497), and 24h-post ($W = 111.00$, $p =$

.088, Rank-Biserial Correlation = .552) timepoints. These analyses did not show any differences

between groups in TMT-B movement time. Additionally, no differences between time points

were observed for either the exercise group ($W = 49.00$, z $= .245$, $p > .999$, Rank-Biserial

Correlation = .077), or the control group ($W = 44.00$, z $= .978$, $p > .999$, Rank-Biserial

Correlation = .333; Figure 5).

Resting-state functional connectivity

Since different MRI scanners were used in this study (Philips Achieva: n = 14; exercise =

12; control = 2; Philips Elition: n = 11; exercise = 2; control = 9) and there were clear differences

between groups for which scanner was used, we first assessed whether scanner type was a

significant covariate in a group × time RM-ANOVA model for DLPFC-sensorimotor network

functional connectivity. Scanner type was not a significant covariate [$F(1,22)= .197$, $p = .661$,

$\eta_p^2= .009$] and therefore no adjustments to the statistical models were made. A group-level seed-

to-network functional connectivity analysis revealed a decrease in functional connectivity

between the right DLPFC (seed) and the sensorimotor network in the exercise group only

(family-wise error rate corrected $p = .024$). Specifically, within the sensorimotor network, a

significant cluster of decreased functional connectivity was observed over the left inferior

parietal lobule [IPL, 232 voxels, CoG MNI coordinates: X = -58.91, Y = -25.05, Z = 20.58;

Figure 6]. We did observe a significant pre-testing difference between groups in the DLPFC-

sensorimotor network functional connectivity [$t(23) = -4.589$, $p < .001$, Cohen's $d = -1.849$]. To

determine if this pre-testing difference impacted our results, an additional between groups

analysis of covariance test was run with pre-testing DLPFC-sensorimotor network functional

connectivity serving as the covariate. This analysis revealed that a significant difference

remained between groups (exercise: -5.841 ± 10.844; control: -2.817 ± 8.474) after covarying

pre-testing differences [$F(1,22) = 5.623$, $p = .027$, $\eta_p^2 = .204$]. To visualize the participant-level

change in functional connectivity that gave rise to this significant cluster, participant-level

connectivity values at each time point were extracted using *fslmeants.* These functional

connectivity values were then used in correlation analyses to determine if the change in

functional connectivity was related to the changes in TMT-A performance.

455   To determine if the significant decrease in DLPFC-sensorimotor network functional

456   connectivity was a by-product of global changes in functional connectivity, we investigated the

457   intra-network functional connectivity of the sensorimotor network. This analysis did not show

458   any statistically significant differences between groups or time points.

459   Since an effect was only observed for the exercise group, only it was used to investigate

sex differences with a sex (male, female) $\times$ time RM-ANOVA. For this analysis, only the sex $\times$

time interaction and the main effect of group were examined given the main effect of time is

redundant with what has already been reported in the previous analyses. The RM-ANOVA failed

to detect a significant sex $\times$ time interaction [$F(1, 12) = 1.480$, $p = .247$, $\eta_p^2 = .110$], or a main

effect of sex [$F(1, 12) = 1.121$, $p = .311$, $\eta_p^2 = .085$].

Relationship between functional connectivity and motor learning

To determine if motor learning of the serial targeting task correlated with the change in

DLPFC-sensorimotor network functional connectivity, a Spearman's Rho correlation was

performed. The Spearman's rho correlation failed to detect a significant relationship between

these two variables ($\rho = .389$, $p = .067$), indicating that the improvement in motor learning was

not related to the change in functional connectivity between the DLPFC and the sensorimotor

network.

Relationship between functional connectivity and improved processing speed

After observing significant effects for TMT-A task score and dwell time, in addition to a

significant cluster of decreased functional connectivity between the DLPFC-sensorimotor

network for the exercise group, we sought to determine if a change in functional connectivity

was related to a change in cognitive-motor performance for both groups. With both groups data

pooled together, Spearman's rho correlations between the change in functional connectivity and

the change in TMT-A task score ($\rho = .458$, $p = .025$) and dwell time ($\rho = .418$, $p = .043$) were

both positively correlated. These relationships suggest that the individuals that experienced a

greater reduction in DLPFC-sensorimotor network functional connectivity also improved their

overall task performance, and reduced the time needed to visually scan for the target and plan

their next arm movement (Boyd *et al.*, 2009*b*) (Figure 7).

Discussion

In the present study, we investigated the impact of high intensity aerobic exercise paired

with motor training on cognitive-motor function in individuals living with chronic stroke. After a

five-day intervention pairing either exercise or rest before a paretic arm implicit motor learning

task, we observed significant improvements in cognitive processing speed with TMT-A, but not

with TMT-B with the less-affected arm. We also used resting-state functional brain imaging to

determine if changes in functional connectivity between the DLPFC and the sensorimotor

network would be observed after our intervention and whether these changes would relate to

behavioural changes. We observed a decrease in DLPFC-sensorimotor network functional

connectivity that was correlated with a change in overall TMT-A task performance ($\rho = .453$),

and the TMT-A dwell time ($\rho = .418$) regardless of group.

Cognitive-motor performance is enhanced after exercise

In the present study, we saw an improvement in TMT-A performance for the exercise

group but not the control group, using the TMT-A task score. After separating the cognitive and

motor components of TMT-A, it was evident that pairing high intensity aerobic exercise with

motor training had a positive impact on performance in the cognitive domain, this was supported

by a significant decrease in dwell time for the exercise group only, with no between group

differences for movement time.

The decreased dwell time for TMT-A and not TMT-B hints at how exercise differentially

impacts the neurocognitive processes involved in these tasks. TMT-A is a measure of cognitive-

motor processing speed (Bowie & Harvey, 2006), whereas TMT-B assesses more complex

cognitive processes like task switching and mental flexibility (Bowie & Harvey, 2006). Our data

suggest that high intensity aerobic exercise had a specific impact on processing speed rather than

on complex cognitive processes such as task switching and mental flexibility.

Altered resting-state functional connectivity related to cognitive-motor performance

In the present study, an expected decrease in functional connectivity was observed

between the right DLPFC, which is known for its involvement in processing speed (Hillary *et al.*,

2006; Kaller *et al.*, 2011) and response selection (Boyd *et al.*, 2009*b*), and the sensorimotor

network in the exercise group only. Specifically, the cluster of decreased functional connectivity

within the sensorimotor network was found over the left inferior parietal lobule, which is

involved in action planning and prediction (Wolpert & Ghahramani, 2000; Kilner *et al.*, 2007;

Elk, 2014). The decrease in functional connectivity between these regions after exercise may

reflect a beneficial effect of exercise on cognitive-motor processing speed in individuals with

stroke, whereby the decreased coupling reflects a shift towards automaticity of perception and

action, and illustrates a reduced dependence on cognitive resources to complete a cognitively

demanding motor task (Mazzoni, 2008; Wu *et al.*, 2008). This notion is supported by previous

research that found a decrease in DLPFC BOLD signal after learning a cognitively challenging

repeated sequence continuous target tracking task with healthy controls, but not in individuals

with stroke (Meehan *et al.*, 2011). These findings likely coincide with "slow" or "late" phases of

motor learning (Dayan & Cohen, 2011). In these stages of learning the attentional demand and

executive resources are no longer required for effective task execution (Schneider & Shiffrin,

1977; Doyon & Benali, 2005; Ashby *et al.*, 2010; Wu *et al.*, 2015). However, past work

suggested that motor practice alone was insufficient to stimulate automaticity of motor plans

after stroke; importantly in this previous work, the same dose of practice enabled age matched

healthy controls to reduce their reliance on DLPFC suggesting that they automated learned

movements (Meehan *et al.*, 2011). Critically, the current study suggests that motor training

paired with high intensity aerobic exercise facilitates the acquisition of TMT-A after stroke,

which is a cognitive-motor task that specifically relies on processing speed.

In the present study, we observed significant motor learning improvements in both

groups (one sample Wilcoxon signed-rank test: $p < .001$). However, there were no differences

between groups ($p = .738$), and the improvements were not related to the change in functional

connectivity. Previous research from our lab using the same experimental paradigm in a healthy

aging cohort (Greeley *et al.*, 2021) and individuals with chronic stroke (Greeley *et al.*, 2023) also

failed to see a preferential advantage of high intensity aerobic exercise for enhancing implicit

motor sequence learning on the serial targeting task compared to controls. These findings may

suggest implicit motor sequence learning tasks, which do not rely heavily on the prefrontal

cortex, do not need the benefits conferred by acute bouts of high intensity aerobic exercise to be

learned; instead, they potentially rely on plasticity within motor networks. In contrast, our data

show that tasks that require cognitive-motor interactions appear to benefit greatly from an

intervention that amplifies plasticity in the prefrontal cortex.

Limitations and future directions

In the context of the present study, high intensity aerobic exercise paired with motor

training failed to alter cognitive-motor performance in TMT-B, which depends on mental

flexibility and task switching. Future work may explore alternative manipulations to various

exercise variables such as duration or intensity of exercise bouts, frequency of exercise sessions,

the timing of exercise sessions in proximity to motor training or even explore anaerobic or

resistance exercise training modalities to determine their efficacy for improving not only

processing speed but other more complex neuro-cognitive processes. It is also currently unclear

how long the exercise-related effects on TMT-A would be retained, and future work should

consider investigating this phenomenon with an additional delayed retention test after the

intervention. In addition, there was no differential impact of high intensity aerobic exercise on

motor learning as characterized by B scores. Importantly this shows that for an implicit motor

sequence learning task (the serial targeting task) practice alone enabled both groups to learn.

Future work should consider more complex, cognitive-motor learning tasks to understand what

types of skills benefit from being paired with high intensity aerobic exercise. We also paired

exercise with motor practice in this study, and it is unclear whether exercise alone would produce

similar outcomes. Therefore, future research should explore the effects of exercise on cognitive-

motor processing speed without skilled motor practice. Finally, the participant sample in the

exercise group was predominantly male (n =19) compared to female (n = 6). This biological sex

imbalance limits our ability to accurately interpret sex differences and future work should

consider larger sample sizes with a more balanced sex distribution to be able to adequately

explore if our findings translate equally or differ between sex.

Conclusions

Five-sessions of high intensity aerobic exercise paired with skilled motor training

improved cognitive-motor performance on a processing speed dependent task. Interestingly, this

effect was not observed for a more complex cognitive-motor task that depended on task

switching and mental flexibility. We also observed a relationship between the amount of change

in DLPFC-sensorimotor network functional connectivity and the change in overall task

performance, and processing speed during TMT-A. Regardless of group, the individuals that had

greater reductions in functional connectivity performed better on TMT-A. These findings suggest

that in individuals with chronic stroke, high intensity aerobic exercise may lead to brain changes

that enable a beneficial decrease in cognitive resources dedicated to task execution, thereby

correcting the high cognitive demand of complex motor tasks often seen after stroke (Puh *et al.*,

2007; Li *et al.*, 2014; Hall *et al.*, 2021). This intervention allowed a restoration of cognitive-

motor function that may have meaningful effects on complex motor task performance in

individuals with chronic stroke.

Acknowledgements

The authors would like to acknowledge Daniela J. Palombo Ph.D. from the Department

of Psychology at the University of British Columbia for her valuable insight and feedback with

analyses, and interpretation of this work.

Funding sources

This work was funded by the Canadian Institutes of Health Research (PI L.A.B., PJT-

148535). JWA, JF, and BCL are funded by Canadian Institutes of Health Research (CIHR)

fellowships. CR received funding from the Natural Sciences and Engineering Research Council

(NSERC) of Canada. CBJ is funded by MITACS (Canada). JN and JWA were also funded by

the Michael Smith Foundation for Health Research.

References

Andreska T, Rauskolb S, Schukraft N, Lüningschrör P, Sasi M, Signoret-Genest J, Behringer M,
Blum R, Sauer M, Tovote P & Sendtner M (2020). Induction of BDNF expression in layer
II/III and layer V neurons of the motor cortex Is essential for motor learning. *J Neurosci* **40,**
6289–6308.

Ashby FG, Turner BO & Horvitz JC (2010). Cortical and basal ganglia contributions to habit
learning and automaticity. *Trends Cogn Sci* **14,** 208–215.

Badre D & Wagner AD (2004). Selection, integration, and conflict monitoring: Assessing the
nature and generality of prefrontal cognitive control mechanisms. *Neuron* **41,** 473–487.

Barnes JN (2015). Exercise, cognitive function, and aging. *Adv Physiol Educ* **39,** 55–62.

Beltz NM, Gibson AL, Janot JM, Kravitz L, Mermier CM & Dalleck LC (2016). Graded
exercise testing protocols for the determination of VO2max: Historical perspectives,
progress, and future considerations. *J Sports Med* **2016,** 1–12.

Bowie CR & Harvey PD (2006). Administration and interpretation of the Trail Making Test. *Nat*
Protoc 2006 15 **1,** 2277–2281.

Boyd LA, Edwards JD, Siengsukon CS, Vidoni ED, Wessel BD & Linsdell MA (2009*a*). Motor
sequence chunking is impaired by basal ganglia stroke. *Neurobiol Learn Mem* **92,** 35–44.

Boyd LA, Vidoni ED, Siengsukon CF & Wessel BD (2009*b*). Manipulating time-to-plan alters
patterns of brain activation during the Fitts' task. *Exp Brain Res* **194,** 527–539.

Brodie SM, Meehan S, Borich MR & Boyd LA (2014). 5 Hz repetitive transcranial magnetic
stimulation over the ipsilesional sensory cortex enhances motor learning after stroke. *Front*
Hum Neurosci; DOI: 10.3389/FNHUM.2014.00143.

Brown S & Heathcote A (2003). Averaging learning curves across and within participants.
Behav Res Methods Instrum Comput **35,** 11–21.

Brunoni AR & Vanderhasselt MA (2014). Working memory improvement with non-invasive
brain stimulation of the dorsolateral prefrontal cortex: a systematic review and meta-
analysis. *Brain Cogn* **86,** 1–9.

Corrigan JD & Hinkeldey NS (1987). Relationships between parts A and B of the Trail Making
Test. *J Clin Psychol*; DOI: 10.1002/1097-4679(198707)43:4<402::AID-
JCLP2270430411>3.0.CO;2-E.

Dal Maso F, Desormeau B, Boudrias MH & Roig M (2018). Acute cardiovascular exercise
promotes functional changes in cortico-motor networks during the early stages of motor
memory consolidation. *Neuroimage* **174,** 380–392.

Dayan E & Cohen LG (2011). Neuroplasticity subserving motor skill learning. *Neuron* **72,** 443–
454.

Doyon J & Benali H (2005). Reorganization and plasticity in the adult brain during learning of
motor skills. *Curr Opin Neurobiol* **15,** 161–167.

Elk M van (2014). The left inferior parietal lobe represents stored hand-postures for object use and action prediction. *Front Psychol* **5,** 333.

Esteban O, Markiewicz CJ, Blair RW, Moodie CA, Isik AI, Erramuzpe A, Kent JD, Goncalves M, DuPre E, Snyder M, Oya H, Ghosh SS, Wright J, Durnez J, Poldrack RA & Gorgolewski KJ (2018). fMRIPrep: a robust preprocessing pipeline for functional MRI. *Nat Methods 2018 161* **16,** 111–116.

Fugl Meyer AR, Jaasko L & Leyman I (1975). The post stroke hemiplegic patient. I. A method for evaluation of physical performance. *Scand J Rehabil Med*.

Gläscher J, Adolphs R, Damasio H, Bechara A, Rudrauf D, Calamia M, Paul LK & Tranel D (2012). Lesion mapping of cognitive control and value-based decision making in the prefrontal cortex. *Proc Natl Acad Sci U S A* **109,** 14681–14686.

Greeley B, Chau B, Jones CB, Neva JL, Kraeutner SN, Campbell KL & Boyd LA (2021). Multiple bouts of high-intensity interval exercise reverse age-related functional connectivity disruptions without affecting motor learning in older adults. *Sci Rep* **11,** 1–16.

Greeley B, Larssen BC, Ferris J, Yeganeh NM, Andrushko JW, Chau B, Jones CB, Campbell KL, Neva JL, Boyd LA & Boyd L (2023). High-intensity exercise paired with motor practice benefits cognitive performance in stroke and older adults. *medRxiv*; DOI: 10.1101/2023.02.09.23285669.

Griffanti L, Salimi-Khorshidi G, Beckmann CF, Auerbach EJ, Douaud G, Sexton CE, Zsoldos E, Ebmeier KP, Filippini N, Mackay CE, Moeller S, Xu J, Yacoub E, Baselli G, Ugurbil K, Miller KL & Smith SM (2014). ICA-based artefact removal and accelerated fMRI acquisition for improved resting state network imaging. *Neuroimage*; DOI: 10.1016/j.neuroimage.2014.03.034.

Hall GR, Kaiser M & Farr TD (2021). Functional connectivity change in response to stroke Is comparable across species: From mouse to man. *Stroke* **52,** 2961–2963.

Hart H, Radua J, Nakao T, Mataix-Cols D & Rubia K (2013). Meta-analysis of functional magnetic resonance imaging studies of inhibition and attention in attention-deficit/hyperactivity disorder: exploring task-specific, stimulant medication, and age effects. *JAMA psychiatry* **70,** 185–198.

Hendy AM, Andrushko JW, Della Gatta PA & Teo WP (2022). Acute effects of high-intensity aerobic exercise on motor cortical excitability and inhibition in sedentary adults. *Front Psychol*; DOI: 10.3389/FPSYG.2022.814633/BIBTEX.

Hillary FG, Genova HM, Chiaravalloti ND, Rypma B & DeLuca J (2006). Prefrontal modulation of working memory performance in brain injury and disease. *Hum Brain Mapp* **27,** 837.

Hodics TM, Nakatsuka K, Upreti B, Alex A, Smith PS & Pezzullo JC (2012). Wolf motor function test for characterizing moderate to severe hemiparesis in stroke patients. *Arch Phys Med Rehabil* **93,** 1963–1967.

Jenkinson M, Bannister P, Brady M & Smith S (2002). Improved optimization for the robust and accurate linear registration and motion correction of brain images. *Neuroimage* **17,** 825–841.

Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW & Smith SM (2012). FSL.
Neuroimage **62,** 782–790.

Kaller CP, Rahm B, Spreer J, Weiller C & Unterrainer JM (2011). Dissociable contributions of
left and right dorsolateral prefrontal cortex in planning. *Cereb Cortex* **21,** 307–317.

Katan M & Luft A (2018). Global burden of stroke. *Semin Neurol* **38,** 208–211.

Kendall BJ, Siekirk NJ & Lai Q (2020). Acute high-intensity interval training improves motor
skill acquisition. *J Sports Med Phys Fitness* **60,** 1065–1071.

Kilner JM, Friston KJ & Frith CD (2007). Predictive coding: An account of the mirror neuron
system. *Cogn Process* **8,** 159–166.

Kortte KB, Horner MD & Windham WK (2002). The trail making test, part B: Cognitive
flexibility or ability to maintain set? *Appl Neuropsychol* **9,** 106–109.

Li W, Li Y, Zhu W & Chen X (2014). Changes in brain functional network connectivity after
stroke. *Neural Regen Res* **9,** 51.

Lin DJ, Erler KS, Snider SB, Bonkhoff AK, DiCarlo JA, Lam N, Ranford J, Parlman K, Cohen
A, Freeburn J, Finklestein SP, Schwamm LH, Hochberg LR & Cramer SC (2021).
Cognitive demands influence upper extremity motor performance during recovery from
acute stroke. *Neurology* **96,** e2576–e2586.

Lin JH, Hsueh IP, Sheu CF & Hsieh CL (2004). Psychometric properties of the sensory scale of
the Fugl-Meyer Assessment in stroke patients. *Clin Rehabil* **18,** 391–397.

Lindsay MP, Norrving B, Sacco RL, Brainin M, Hacke W, Martins S, Pandian J & Feigin V
(2019). World stroke organization (WSO): Global stroke fact sheet 2019. *Int J Stroke* **14,**
806–817.

Love J, Selker R, Marsman M, Jamil T, Dropmann D, Verhagen J, Ly A, Gronau QF, Šmíra M,
Epskamp S, Matzke D, Wild A, Knight P, Rouder JN, Morey RD & Wagenmakers EJ
(2019). JASP: Graphical statistical software for common statistical designs. *J Stat Softw*;
DOI: 10.18637/jss.v088.i02.

MacPherson SE, Cox SR, Dickie DA, Karama S, Starr JM, Evans AC, Bastin ME, Wardlaw JM
& Deary IJ (2017). Processing speed and the relationship between Trail Making Test-B
performance, cortical thinning and white matter microstructure in older adults. *Cortex* **95,**
92.

Mang CS, Campbell KL, Ross CJD & Boyd LA (2013). Promoting neuroplasticity for motor
rehabilitation after stroke: Considering the effects of aerobic exercise and genetic variation
on brain-derived neurotrophic factor. *Phys Ther* **93,** 1707–1716.

Mang CS, Snow NJ, Wadden KP, Campbell KL & Boyd LA (2016). High-intensity aerobic
exercise eenhances motor memory retrieval. *Med Sci Sports Exerc* **48,** 2477–2486.

Mazzoni P (2008). Efficient motor control: how can less be more? *J Physiol* **586,** 4031.

Meehan SK, Randhawa B, Wessel B & Boyd LA (2011). Implicit sequence-specific motor
learning after subcortical stroke is associated with increased prefrontal brain activations: An

fMRI Study. *Hum Brain Mapp* **32,** 290–303.

Mittmann N, Seung SJ, Hill MD, Phillips SJ, Hachinski V, Coté R, Buck BH, Mackey A,
714   Gladstone DJ, Howse DC, Shuaib A & Sharma M (2012). Impact of disability status on
715   ischemic stroke costs in Canada in the first year. *Can J Neurol Sci / J Can des Sci Neurol*
716   **39,** 793–800.

717   Nasreddine ZS, Phillips NA, Bédirian V, Charbonneau S, Whitehead V, Collin I, Cummings JL
718   & Chertkow H (2005). The montreal cognitive assessment, MoCA: A brief screening tool
719   for mild cognitive impairment. *J Am Geriatr Soc* **53,** 695–699.

720   Nepveu JF, Thiel A, Tang A, Fung J, Lundbye-Jensen J, Boyd LA & Roig M (2017). A single
721   bout of high-intensity interval training improves motor skill retention in individuals with
722   stroke. *Neurorehabil Neural Repair* **31,** 726–735.

723   Neva JL, Greeley B, Chau B, Ferris JK, Jones CB, Denyer R, Hayward KS, Campbell KL &
724   Boyd LA (2022). Acute high-intensity interval exercise modulates corticospinal excitability
725   in older adults. *Med Sci Sports Exerc* **54,** 673–682.

726   Nickerson LD, Smith SM, Öngür D & Beckmann CF (2017). Using Dual Regression to
727   Investigate Network Shape and Amplitude in Functional Connectivity Analyses. *Front*
728   *Neurosci* **11,** 115.

729   Puh U, Vovk A, Sevšek F & Šuput D (2007). Increased cognitive load during simple and
730   complex motor tasks in acute stage after stroke. *Int J Psychophysiol* **63,** 173–180.

731   Reuter M, Rosas HD & Fischl B (2010). Highly accurate inverse consistent registration: a robust
732   approach. *Neuroimage* **53,** 1181–1196.

733   Salimi-Khorshidi G, Douaud G, Beckmann CF, Glasser MF, Griffanti L & Smith SM (2014).
734   Automatic denoising of functional MRI data: Combining independent component analysis
735   and hierarchical fusion of classifiers. *Neuroimage* **90,** 449–468.

736   Sallet J, Mars RB, Noonan MP, Neubert FX, Jbabdi S, O'Reilly JX, Filippini N, Thomas AG &
737   Rushworth MF (2013). The organization of dorsal frontal cortex in humans and macaques. *J*
738   *Neurosci* **33,** 12255.

739   Schneider W & Shiffrin RM (1977). Controlled and automatic human information processing: I.
740   Detection, search, and attention. *Psychol Rev* **84,** 1–66.

741   Singh AM, Duncan RE, Neva JL & Staines WR (2014). Aerobic exercise modulates intracortical
742   inhibition and facilitation in a nonexercised upper limb muscle. *BMC Sports Sci Med*
743   *Rehabil* **6,** 1–10.

744   Singh AM & Staines WR (2015). The effects of acute aerobic exercise on the primary motor
745   cortex. *J Mot Behav* **47,** 328–339.

746   Skriver K, Roig M, Lundbye-Jensen J, Pingel J, Helge JW, Kiens B & Nielsen JB (2014). Acute
747   exercise improves motor memory: exploring potential biomarkers. *Neurobiol Learn Mem*
748   **116,** 46–58.

Sleiman SF, Henry J, Al-Haddad R, El Hayek L, Haidar EA, Stringer T, Ulja D,

Karuppagounder SS, Holson EB, Ratan RR, Ninan I & Chao M V. (2016). Exercise
promotes the expression of brain derived neurotrophic factor (BDNF) through the action of
the ketone body β-hydroxybutyrate. *Elife*; DOI: 10.7554/ELIFE.15092.

Smith SM, Fox PT, Miller KL, Glahn DC, Fox PM, Mackay CE, Filippini N, Watkins KE, Toro
R, Laird AR & Beckmann CF (2009). Correspondence of the brain's functional architecture
during activation and rest. *Proc Natl Acad Sci U S A* **106,** 13040–13045.

Stagg CJ, Bachtiar V & Johansen-Berg H (2011). The role of GABA in human motor learning.
Curr Biol **21,** 480–484.

Stavrinos EL & Coxon JP (2017). High-intensity interval exercise promotes motor cortex
disinhibition and early motor skill consolidation. *J Cogn Neurosci* **29,** 593–604.

Tustison NJ, Avants BB, Cook PA, Zheng Y, Egan A, Yushkevich PA & Gee JC (2010).
N4ITK: improved N3 bias correction. *IEEE Trans Med Imaging* **29,** 1310–1320.

Wadden KP, Asis K De, Mang CS, Neva JL, Peters S, Lakhani B & Boyd LA (2017). Predicting
motor sequence learning in individuals with chronic stroke. *Neurorehabil Neural Repair* **31,**
95–104.

Wadden KP, Hodges NJ, De Asis KL, Neva JL & Boyd LA (2019). Individualized challenge
point practice as a method to aid motor sequence learning. *J Mot Behav* **51,** 467–485.

Wang S, Peterson DJ, Gatenby JC, Li W, Grabowski TJ & Madhyastha TM (2017). Evaluation
of field map and nonlinear registration methods for correction of susceptibility artifacts in
diffusion MRI. *Front Neuroinform* **11,** 17.

Weng TB, Pierce GL, Darling WG, Falk D, Magnotta VA & Voss MW (2017). The acute effects
of aerobic exercise on the functional connectivity of human brain networks. *Brain Plast* **2,**
171–190.

Winkler AM, Ridgway GR, Webster MA, Smith SM & Nichols TE (2014). Permutation
inference for the general linear model. *Neuroimage* **92,** 381–397.

Wolf SL, Catlin PA, Ellis M, Archer AL, Morgan B & Piacentino A (2001). Assessing Wolf
motor function test as outcome measure for research in patients after stroke. *Stroke* **32,**
1635–1639.

Wolpert DM & Ghahramani Z (2000). Computational principles of movement neuroscience. *Nat
Neurosci 2000 311* **3,** 1212–1217.

Wu T, Chan P & Hallett M (2008). Modifications of the interactions in the motor networks when
a movement becomes automatic. *J Physiol* **586,** 4295.

Wu T, Hallett M & Chan P (2015). Motor automaticity in Parkinson's disease. *Neurobiol Dis* **82,**
226–234.

Zheng G, Zhou W, Xia R, Tao J & Chen L (2016). Aerobic exercises for cognition rehabilitation
following stroke: A systematic review. *J Stroke Cerebrovasc Dis* **25,** 2780–2789.

Figures

Figure 1. *Experimental design timeline. MRI = magnetic resonance imaging, TMT-A & B = Trail Making Test Part A and B, Clinical*
tests = Fugl-Meyer Assessment, and Wolf Motor Function Test at pre-testing and 24h-post timepoints, and Montréal cognitive
assessment at pre-testing only.

Figure 2. *Lesion overlap. Colour bar represents the number of participants that have a lesion in a given location (i.e., lesion voxel*
overlap; A value of 4 means 4 participants have a lesion in the same location). Figure is in radiological view with the right side of
the brain on the left, and the left side of the brain on the right.

Figure 3. Serial targeting task B values. There were no differences between groups (p = .738), but data were significantly
different from zero (p < .001). Black bars represent the group means. Grey circles represent individual data points.

Figure 4. *Trail Making Test Part A (TMT-A). Pre-testing and 24h-post testing participant values for the Exercise group (left) and control group (right) for **A)** Task score, **C)** Total dwell time, **E)** Total movement time. Panels **B), D),** and **F)** represent change scores for each of the measures respectively. group × time RM-ANOVA Interaction results are represented with the line and p-values over the change scores on the right. Total dwell time was run with non-parametric testing and therefore no group × time interaction was assessed. All p-values are Bonferroni adjusted. Blue shaded bars represent standard deviation. For panels B, D, and F Black bars represent the group means. Grey circles represent individual data points*

Figure 5. *Trail Making Test Part B (TMT-B). Pre-testing and 24h-post testing participant values for the Exercise group (left) and control group (right) for **A)** Task score, **C)** Total dwell time, **E)** Total movement time. Panels **B), D),** and **F)** represent change scores for each of the measures respectively. All tests were run with non-parametric testing and therefore no group × time interactions were assessed. All p-values are Bonferroni adjusted. Blue shaded bars represent standard deviation. For panels B, D, and F Black bars represent the group means. Grey circles represent individual data points*

Figure Error! No text of specified style in document.**6.** Resting-state functional connectivity results from the seed-to-network
analysis. **A)** A glass brain showing the right dorsolateral prefrontal cortex (DLPFC) seed in green, the sensorimotor network
component from the group level ICA (grey), and the significant cluster of decreased functional connectivity over the left inferior
parietal lobule (IPL) in red for the exercise group. **B)** individual functional connectivity scores at pre-testing and 24h-post time
points for the exercise group (left) and the control group (right). **C)** Functional connectivity change scores (24h-post – pre-
testing) for the exercise group (left) and control group (right). Blue shaded bars represent standard deviation. For panel C, black
bars represent the group means. Grey circles represent individual data points ANCOVA = Analysis of Covariance adjusted for pre-
test differences.

Figure 7. *Spearman's Rho (ρ) correlations between the change in DLPFC-sensorimotor network functional connectivity (x-axes), and the change in Trail Making Test Part A (TMT-A) **A)** Task score, and **B)** Total dwell time. The Grey shaded areas represent the 95% confidence interval around the regression line.*

Trail Making Test Part A

Trail Making Test Part B

A

DLPFC

Sensorimotor Network

Sensorimotor Network

L   R

IPL

Cluster information:
Size: 232 voxels
Mean value: 0.9786 (i.e., FWE-corrected p = 0.024)
CoG: -58.91 -35.05 20.58 (MNI coordinates)

B

Functional Connectivity (Arbitrary Units)

p = .024

ANCOVA adjusted p = .027

Pre-Test   24h-Post   Pre-Test   24h-Post

Exercise   Control

C

Δ Functional Connectivity (Arbitrary Units)

Exercise   Control