PML mutants resistant to arsenic induced degradation fail to generate the

appropriate SUMO and ubiquitin signals required for RNF4 and p97

recruitment.

Ellis G. Jaffray, Michael H. Tatham, Alejandro Rojas-Fernandez, Adel Ibrahim,

Graeme Ball and Ronald T. Hay*

Centre for Gene Regulation and Expression, School of Life Sciences, University

of Dundee, Dundee, UK

***Corresponding author**

Email: r.t.hay@dundee.ac.uk

ORCID: 0000-0001-7113-9024

ABSTRACT

Arsenic is an effective treatment for Acute Promyelocytic Leukaemia as it induces

degradation of the Promyelocytic Leukaemia (PML) – retinoic acid receptor alpha

(RARA) oncogenic fusion protein. Some patients relapse with arsenic resistant

disease because of missense mutations in PML. To determine the mechanistic basis

for arsenic resistance we reconstituted PML-/- cells with YFP fusions of wild type

(WT) and two mutant forms of PMLV found in patients refractory to arsenic, A216T

and L217F. Both mutants formed PML bodies that were larger, but less numerous

than WT and neither responded to arsenic by degradation. Analysis of

immunoprecipitated PML bodies indicated that while WT PML experiences increased

SUMO1, SUMO2/3 and ubiquitin conjugation, A216T PML is almost completely

unresponsive and therefore does not recruit the SUMO targeted ubiquitin E3 ligase

RNF4. Compared to WT PML, L217F PML was modified to a similar extent by

SUMO2 but not SUMO1 and although it recruited RNF4, it failed to develop the

appropriate poly-ubiquitin conjugates required to recruit the segregase p97, which is

essential for PML degradation.

INTRODUCTION

Acute Promyelocytic Leukaemia (APL) is caused by a reciprocal chromosomal

translocation t(15;17) that fuses the genes encoding the Promyelocytic Leukaemia

(PML) protein and the retinoic acid receptor alpha (RARA) (de The, Lavau et al.

1991, Kakizuka, Miller et al. 1991). This generates the PML-RARA oncoprotein that

deregulates transcriptional programmes required for the differentiation of

haematopoetic progenitor cells. Therefore, differentiation is blocked and

promyelocytes accumulate causing leukaemia (Grignani, Ferrucci et al. 1993, Kwok,

Zeisig et al. 2006, Martens, Brinkman et al. 2010, Mikesch, Gronemeyer et al. 2010,

Tan, Wang et al. 2021). In healthy cells the PML gene is alternatively spliced to

generate multiple mRNAs that encode seven protein isoforms (I-VII) that differ in

their C-terminal regions.

PML is a member of the Tripartite Motif (TRIM) family of proteins and is also known

as TRIM19. Located in the N-terminal region of PML and present in all PML isoforms

and PML-RARA fusions, the highly conserved TRIM is composed of a RING domain,

two zinc co-ordinating B-boxes and a coiled-coil that mediates dimerization (Bernardi

and Pandolfi 2007). In cells with wild type (WT) PML genes, the protein is associated

with non-membranous nuclear structures known as PML nuclear bodies, that also

accumulate a variety of other proteins including Small Ubiquitin-like Modifiers

(SUMOs), p53, Daxx and SP100. In cells containing a PML-RARA fusion protein,

PML and PML-RARA are co-associated in a large number of tiny nuclear speckles

(Dyck, Maul et al. 1994, Weis, Rambaud et al. 1994). In the past, treatment options

were limited and APL was a disease with a very poor prognosis. However, the

majority of patients are now cured by a combination therapy consisting of arsenic

trioxide (referred to herein as As) and *all trans* retinoic acid (ATRA) (Wang and Chen

2008, Lo-Coco, Avvisati et al. 2013, Mi, Chen et al. 2015). These induce degradation

of the PML-RARA oncogene and allows the WT version of RARA to initiate a

transcriptional programme that drives the accumulated promyelocytes down a

pathway of terminal differentiation that ultimately cures the disease. Arsenic alone

induces degradation of both the unfused PML and PML-RARA, identifying PML as

the target of As. Treatment with As leads to rapid multisite modification of PML and

PML-RARA with SUMO (Muller, Matunis et al. 1998). SUMO modified PML and

PML-RARA then serves to recruit the SUMO Targeted Ubiquitin E3 ligase (STUbL)

RING Finger Protein 4 (RNF4) (Perry, Tainer et al. 2008, Geoffroy and Hay 2009).

Recruitment of RNF4 to SUMO modified PML and PML-RARA present in PML

nuclear bodies is mediated by multiple SUMO Interaction Motifs (SIMs) present in

the N-terminal region of the protein. The high local concentration of RNF4 leads to

dimerization of the C-terminal RING domains and activation of the ubiquitin E3 ligase

activity of the protein (Rojas-Fernandez, Plechanovova et al. 2014). RNF4 mediated

ubiquitination of SUMO modified PML and PML-RARA ultimately leads to their

proteolytic degradation by the proteasome (Lallemand-Breitenbach, Jeanne et al.

2008, Tatham, Geoffroy et al. 2008). Recently, an additional step required for arsenic

induced degradation of PML and PML-RARA has been uncovered that involves the

p97/VCP segregase (Jaffray, Tatham et al. 2023). RNF4 mediated ubiquitination of

PML leads to recruitment of UFD1-NPLOC4-p97 which extracts ubiquitinated PML

and presumably feeds the modified protein to the proteasome where it is

proteolytically degraded.

While the majority of APL patients treated with As are cured, a small proportion of

these patients relapse and present with arsenic resistant disease. In some cases

resistance may be a consequence of metabolic reprogramming linked to expression

of additional known oncogenes (Madan, Shyamsunder et al. 2016, Iaccarino, Ottone

et al. 2019), but in many patients, arsenic resistance is a result of missense

mutations in PML clustered between residues L211 and S220 in B-box 2. These

mutations are not only found in the PML-RARA oncogene (Goto, Tomita et al. 2011,

Chendamarai, Ganesan et al. 2015, Lou, Ma et al. 2015, Iaccarino, Ottone et al.

2016, Liu, Zhu et al. 2016) but also in the unfused version of PML (Lehmann-Che,

Bally et al. 2014, Zhu, Qin et al. 2014, Iaccarino, Ottone et al. 2016). These

mutations are undetectable at initial diagnosis, suggesting that cells expressing

these mutations are selected for by arsenic treatment (Alfonso, Iaccarino et al. 2019,

Balasundaram, Ganesan et al. 2022).

To establish the mechanistic basis for the arsenic resistance we investigated the

properties of two mutations in PML that responded differently to As. While both

mutations were resistant to As-induced degradation, the A216T mutant was severely

compromised for SUMO modification, failed to recruit RNF4 and as a result was not

ubiquitinated. In contrast, like WT PML the L217F mutant was modified by

SUMO2/3, recruited RNF4, and was ubiquitinated in response to As. However,

SUMO1 modification was compromised and the ubiquitin signal was not sufficient to

recruit the p97 segregase that is required to extract PML from nuclear bodies. Thus,

while two mutations that are adjacent in the PML B-box both inhibit As-induced PML

degradation, mechanistic failure is at different steps of the degradation pathway.

RESULTS

PML mutants A216T and L217F form atypical PML bodies in U2OS cells.

Mutations in PML and PML-RARA that arise in APL patients with arsenic resistant

disease are a unique resource as these mutations are selected in vivo to be resistant

to arsenic while still retaining the biological functions of PML that contribute to the

disease phenotype. The majority of these mutations are located to a 10 amino acid

region (L211-S220) in B-box 2 of PML that encompasses, but never includes, the

zinc co-ordinating residues C212 and C215 (Fig. 1A). As PML is the direct target of

arsenic and mutations from patients with arsenic resistant disease can also be

detected in the non-rearranged allele of PML (Lehmann-Che, Bally et al. 2014,

Iaccarino, Ottone et al. 2016), we chose to reconstitute U2OS PML-/- cells with wild-

type (WT) or mutant versions of PMLV N-terminally fused to YFP (Jaffray, Tatham et

al. 2023). In testing a number of PML variants, we found that they fell into two main

groups exemplified by the most frequently identified mutations in patients; A216T

and L217F. U2OS cells expressing YFP fused forms of WT, A216T and L217F

PMLV were isolated by FACS and the derived cell lines analysed by high content

microscopy. Images of these cells show that while the total YFP fluorescence signal

was similar for WT, A216T and L217F, the average size and number of PML bodies

per cell varied (Fig. 1B). Quantitative analysis of thousands of cells showed that the

average expression levels, measured by YFP-PML intensity per cell, were similar for

all three PML forms (Fig. 1C). However, cells expressing A216T and L217F PMLV

had on average approximately half the number of PML bodies per cell (median = 4)

compared to WT (median = 8) (Fig. 1C). This was contrasted by both mutants

showing larger than WT PML bodies according to their measured cross-sectional

area (Fig. 1C). This ~20-25% larger area equates to almost twice the volume when

considering a spherical PML body. These results show that A216T and L217F

mutations to PML reduce PML body number and increase PML body size in

untreated cells.

PML mutants A216T and L217F are not degraded in response to arsenic in

U2OS cells

Arsenic exposure is responsible for dramatic changes to the number, morphology

and content of PML bodies (Geoffroy et al., 2010; Jeanne et al., 2010). To determine

how these parameters were influenced by mutations in PML, the cells expressing

WT, A216T or L217F YFP-PMLV were either untreated or treated with 1μM arsenic

and monitored by time-lapse fluorescence microscopy over a 15 hr period. At least 9

movies were collected for each condition and an automated, quantitative pipeline

(Jaffray, Tatham et al. 2023) was used to follow PML body size, number and total

YFP-PMLV fluorescence intensity over time (see Supplementary movies 1-6 for

representative time-course data). Representative images from the cells at 0 and 15

hr (Fig. 2A) showed the expected reduction in the number and total intensity of

nuclear bodies in WT PMLV expressing cells, but not in cells expressing A216T or

L217F PMLV (Fig. 2A). Averaged values of these metrics across all time courses

indicated that PML body size did not significantly change over time after arsenic

treatment for any of the PML forms studied (Fig. 2B). Furthermore, the number of

bodies per cell and normalized YFP-PML intensity was reduced during arsenic

exposure for WT PML but not for either mutant (Fig. 2B). More specifically, after 15

hr As there was a >4 fold reduction in the number of PML bodies per cell and >3 fold

reduction in YFP-PMLV intensity for the WT PMLV, while little or no change was

observed for A216T and L217F mutants (Fig. 2C). Western-blot analysis of extracts

gathered from cells exposed to arsenic over a 24 hr time-course showed that as

expected, WT PML rapidly shifts to a higher molecular weight species and is

completely degraded by 24 hr (Fig. 2D). In contrast, A216T PML does not appear to

shift to a higher molecular weight form and its levels do not change over the 24 hr

period. Unlike A216T, L217F PML does shift to higher molecular weight species, but

it is not degraded in response to arsenic after 24 hr (Fig. 2D).

In response to arsenic, WT and L217F PML recruit RNF4 but the A216T mutant

does not

Although A216T does not appear to respond to arsenic by altered post-translational

modification, both the WT and L217F forms display reduced electrophoretic mobility

upon exposure of cells to arsenic (Fig. 2D). To better understand the nature of the

modification status of all PML variants, nuclear bodies were purified from WT, A216T

and L217F YFP-PMLV expressing cells (Jaffray, Tatham et al. 2023) both before and

after treatment with $1\mu M$ As for 2h and analysed by Western blot for PML, SUMO1

or SUMO2/3. Consistent with the analysis of PML from whole cell extracts (Fig. 2D),

purified YFP-PMLV WT and L217F shifted to higher molecular weight species in

response to arsenic treatment for 2 hrs, whereas this response was almost

completely blunted for A216T (Fig. 3A). SUMO1 modification of all three PML forms

in untreated cells appears to be similar, but in response to arsenic, WT YFP-PMLV

undergoes a much greater increase in SUMO1 modification than either A216T or

L217F mutants. As with SUMO1, in untreated cells conjugation of all three PML

forms to SUMO2/3 is broadly comparable (Fig. 3A). However, there are some

notable differences in the SUMO2/3 response to arsenic in comparison with SUMO1.

Firstly, the scale of the As-induced increase in SUMO2/3 conjugation to WT-PML is

not as large as the relative increase in SUMO1. Secondly, both mutants show a

SUMO2/3 conjugation response to As treatment, albeit weakest for the A216T

mutant (Fig. 3A).

Enhanced SUMO modification of PML is responsible for the recruitment of the

STUbL RNF4 that mediates PML ubiquitination prior to degradation by the

proteasome. To study RNF4 recruitment in response to arsenic, nuclear bodies from

WT, A216T and L217F YFP-PMLV expressing cells were again purified and the

associated RNF4 determined by Western blotting (Fig. 3B). RNF4 was not stably

associated with nuclear bodies in the absence of arsenic, but was detected in

purifications containing WT and L217F YFP-PMLV forms after arsenic treatment

(Fig. 3B). By this method RNF4 was not detectable in A216T YFP-PMLV

purifications in response to arsenic (Fig. 3B). The YFP-PMLV purifications were then

analysed for ubiquitin modification by Western blotting with an antibody that detects

ubiquitin chains. This revealed that WT YFP-PMLV was extensively ubiquitinated in

response to arsenic and, consistent with the failure to recruit RNF4, increase in the

ubiquitination of A216T YFP-PMLV was not detected (Fig. 3B). Interestingly,

although L217F YFP-PMLV recruited RNF4, the increase in ubiquitin chains was not

apparently as great as for the WT protein, and the modified species had a lower

molecular weight (Fig. 3B).

The arsenic-induced recruitment of SUMO1, SUMO2/3 and RNF4 to PML bodies

was also analysed by fluorescence microscopy for the three different cell lines. This

confirmed that colocalization of SUMO1, SUMO2/3 and RNF4 with PML was

increased for the WT protein in response to arsenic (Supplementary Figs. 1-3).

Consistent with the Western blot data (Fig. 3A+B), the A216T PML mutant showed

little response to arsenic in terms of recruitment of SUMO1, SUMO2/3 or RNF4. The

L217F PML variant showed lower than WT recruitment of SUMO1 and SUMO2/3 in

response to arsenic (Supplementary Fig. 1&2), although RNF4 levels appeared

similar (Supplementary Fig. 3). In this experiment, RNF4 appears to colocalise with

PML nuclear bodies prior to arsenic treatment, but this increases in response to

arsenic for both WT and L217F and less so for A216T (Supplementary Fig. 3). Thus,

both Western blotting and fluorescence microscopy indicate that the inability of

A216T PML to be degraded is the consequence of insufficient SUMO conjugation

which fails to recruit RNF4, and therefore is not ubiquitinated. This was not the case

for L217F which could recruit RNF4 and induce ubiquitination, but was still resistant

to degradation.

220   In response to arsenic L217F PML fails to recruit the p97 segregase

Recently we determined that the p97 segregase was required to extract ubiquitinated

PML from nuclear bodies prior to degradation by the proteasome (Jaffray, Tatham et

al. 2023). To determine if the mutants were compromised for p97 extraction we

analysed WT, A216T and L217F YFP-PMLV expressing cells by fluorescence

microscopy 0, 2, 4 and 6 hr after arsenic treatment. p97 recruitment was determined

by evaluating the extent of co-localisation between antibody stained p97 and PML

marked by YFP fluorescence. As p97 is a highly abundant protein we employed a

pre-extraction procedure to release the bulk of the soluble p97 from the cells before

fixation. In WT YFP-PMLV untreated cells p97 was associated with only a small

proportion of PML bodies, whereas after 6 hr arsenic treatment a larger fraction of

PML bodies had associated p97 (Supplementary Fig. 4). In cells expressing A216T

and L217F YFP-PMLV, arsenic treatment did not appear to increase the association

of p97 with PML bodies (Fig. 4A, Supplementary Fig. 4). To quantify this, the

percentage of PML bodies co-localised with p97 in WT, A216T and L217F YFP-

PMLV expressing cells was determined at 0, 2, 4 and 6 hr after arsenic treatment

(Fig. 4B). In untreated WT YFP-PMLV expressing cells about 20% of PML bodies

were associated with p97, which increased to over 60% at 2, 4 and 6 hr after arsenic

treatment. In untreated cells expressing A216T and L217F YFP-PMLV, p97

association was less than 10% and this did not increase in response to arsenic (Fig.

4B). Statistical analysis of the 0 hr and 6 hr time-points data revealed that there was

a substantial and statistically significant recruitment of p97 to PML bodies in cells

expressing WT YFP-PMLV, but not in cells expressing A216T or L217F YFP-PMLV

(Fig. 4C). Changes in the expression levels of p97 in the different cell lines, or upon

arsenic treatment could not explain these differences (Supplementary Fig. 5). Thus,

although L217F YFP-PMLV recruits SUMO and RNF4 to generate ubiquitin

conjugates, this signal is somehow not sufficient to engage p97, and therefore

proteasomal degradation fails.

Differential modification with SUMO and ubiquitin of PML mutants

To provide more detail on the post-translational modifications that accumulate on

PML, we repeated the purification of YFP-PML nuclear bodies with a view to release

SUMO or ubiquitin from resin-bound material by treatment with proteases specific for

each (Supplementary Fig. 6A). The material released by protease treatment, and that

remaining on the beads can be analysed by Western blotting, which should provide

information not only on the covalent modification status of PML, but also that of the

SUMO1, SUMO2/3 and ubiquitin species attached to PML (Supplementary Fig. 6B).

Prior to purification, nuclear extracts for WT, A216T and L217F YFP-PMLV displayed

the expected changes in electrophoretic mobility (Supplementary Fig. 6C), and were

almost completely depleted from nuclear extracts after YFP-PML purification.

Importantly, the modifications to PML were preserved during purification (Fig. 5A "No

protease"). For all PML variants treatment of the beads with SENP1 returned most of

the heavier PML forms back to more rapidly migrating species consistent with

unmodified YFP-PMLV (Fig. 5 "SENP1" lanes). Treatment with USP2 had little impact

on the distribution of PML species, indicating only a relatively small proportion of the

modified PML is ubiquitin conjugated (Fig. 5A "USP2"). SUMO1 blots show that in

untreated cells there are relatively low levels of SUMO1 associated with PML, while

after arsenic treatment this increases dramatically for WT YFP-PMLV, modestly for

L217F YFP-PMLV, and not at all for A216T YFP-PMLV (Fig. 5B). As expected, these

species disappear upon SENP1 treatment, but are largely unaltered by treatment with

USP2 (Fig. 5B). After arsenic exposure, the relative increase to SUMO2/3 modification

is not as large as for SUMO1. Notably all three PML forms undergo an increase in

SUMO2/3 conjugation, although A216T PML is modified to a much lesser degree than

either WT or L217F (Fig. 5C). SUMO2/3 is also completely removed by SENP1, but

USP2 treatment does not result in significant changes to SUMO2/3 antibody reactive

species (Fig. 5C). Blotting with an antibody to ubiquitin reveals that it is present at low

levels on WT, A216T and L217F YFP-PMLV in untreated cells but is dramatically

increased on WT, modestly increased on L217F but any increase on A216T is

undetectable in this assay (Fig. 5D). For both WT PML and the L217F mutant, SENP1

treatment not only reduces the amount of ubiquitin remaining on the beads, it also

causes the ubiquitin-reactive species to run at a reduced molecular weight (Fig. 5D).

This ubiquitin released by SENP1 is probably conjugated directly to SUMO, while the

material remaining is directly conjugated to PML. As expected, ubiquitin is removed

from purifications by treatment with USP2 (Fig. 5D). Although ubiquitination of L217F

in response to arsenic is less than that of WT YFP-PMLV, it is still far more substantial

than the A216T mutant (Fig. 5D).

The material released from the bead-associated YFP-PMLV by treatment with the

proteases was also analysed by Western blotting. As expected, none of the treatments

released YFP-PMLV (Supplementary Fig. 7A). SENP1 treatment released low levels

of monomeric SUMO1 from untreated WT and L217F YFP-PMLV samples, with none

detected from A216T purifications. After arsenic treatment the amount of SUMO1

released by SENP1 from WT samples increased for both WT and L217F PML forms,

albeit less extensively for the mutant. Compared to these two, the amount of SUMO1

conjugated to A216T after arsenic exposure was negligible although it was at least

detectable compared to the sample derived from untreated cells (Supp. Fig. 7B).

SENP1 treatment of WT-PML purifications also released forms of SUMO1 shifted up

around 10kDa from the unmodified SUMO1 species, which are presumed to be

SUMO1 conjugated by ubiquitin, as they are sensitive to USP2 (Supplementary Fig.

7B). However, these species were not apparent in SENP1 elutions from L217F-PML

purifications according to this assay, implying ubiquitination of SUMO1 is most

extensive for in WT-PML. SENP1 digestion released substantial amounts of

monomeric SUMO2/3 from preparations derived from untreated cells for all PML

variants including A216T (Supp. Fig. 7C). Arsenic treatment induced a modest

increase in SUMO2/3 conjugates on all three PML forms, although a ubiquitinated

form of SUMO2/3 was only apparent for WT and L217F variants (Supp. Fig. 7C).

These SUMO-ubiquitin hybrid species released by SENP1 were also detected using

a ubiquitin antibody (Supplementary Fig. 7D). Although data shown in Fig. 5C suggest

SUMO2/3 modification of PML is low prior to arsenic and increases after arsenic

treatment, this is not consistent with analysis of the material released by SENP1

treatment (Supplementary Fig. 6C) which shows similar SUMO2/3 amounts both

before and after arsenic. A possible explanation for this apparent discrepancy is that

in untreated cells the SUMO2/3 modified PML are dispersed throughout the gel,

whereas after SENP1 treatment all the SUMO2/3 runs as a single species.

In summary, A216T YFP-PMLV is defective for arsenic induced SUMO1, SUMO2/3

and ubiquitin modification. L217F YFP-PMLV appears to be modestly compromised

for SUMO2/3 conjugation but shows much less than WT-PML levels of SUMO1

conjugation. Furthermore, WT PML shows higher levels of direct ubiquitination than

L217F, whereas L217F ubiquitination appears to mostly take place on SUMO2/3

linked to PML. Finally, a significant proportion of SUMO1 conjugated to WT-PML is

covalently conjugated by ubiquitin, while this was not detected for L217F PML.

Quantitative proteomic analysis of PML post-translational modifications in

response to arsenic

Early descriptions of PML modification by SUMO identified three major acceptors at

lysines 65, 160 and 490 (Kamitani, Kito et al. 1998). More recently, the single largest

SUMO site proteomic analysis to date (Hendriks, Lyon et al. 2017) described a total

of 15 acceptor lysines in PML (Supplementary figure 8A). To provide quantitative

data at the amino-acid site level for the post-translational modifications present in

PML and SUMO, a peptide-level quantitative proteomics analysis was undertaken.

Trypsin digestion leaves long SUMO adducts attached to modified lysines

(Supplementary Fig 8B), that are challenging to identify by mass-spectrometry. Thus,

a strategy combining GluC digestion and trypsin digestion was employed to give

shorter peptide fragments. As GluC is less specific and less stable than trypsin,

multiple GluC adducts from SUMOs remain on lysines after digestion

(Supplementary Fig. 8B), which has the advantage of providing multiple peptide

evidences for a single modification.

Cultures of U2OS PML-/- +YFP-PML cells were grown for the WT, A216T and L217F

PML variants, and were either treated or not with 1μM arsenic for 2 hr

(Supplementary Fig. 9A). Four experimental replicates for each condition were

prepared and YFP-PML purified by the same method as described above.

Purifications were analysed by SDS-PAGE (Supplementary Fig. 9B) and the

uppermost section of the gel containing proteins larger than 100kDa, including YFP-

PML, was processed for mass-spectrometry analysis. Anti-PML Western blot of a

fraction of the inputs revealed the YFP-PML proteins responded as expected to

arsenic treatment (Supplementary Fig. 9C). 21 phosphorylation, 14 SUMOylation

and 11 ubiquitination sites were identified in PML (Supplementary datafile 1 and

Supplementary Fig. 10A). Principal component analysis using intensity data for all

modified and unmodified peptides from YFP-PML, SUMO1, SUMO2, SUMO3 and

ubiquitin shows replicates cluster by experimental condition and are clearly

separated from one-another (Supplementary Fig. 10B), indicating consistent

differences among the conditions. Intensity data for all peptides were normalized by

total YFP peptide intensity (Supplementary Fig. 10C), to minimize any systematic

errors relating to PML expression or loading (Supplementary Fig. 10D). Using the

total intensity of the unmodified peptides from SUMO1, SUMO2/3 and ubiquitin as a

proxy for their overall protein abundance (Supplementary Fig. 10E-G), a remarkably

similar picture to that revealed in the Western blot experiments above (Fig. 3) was

found. Namely, prior to exposure to arsenic all three PML forms are associated with

broadly similar amounts of SUMO1, SUMO2/3 and ubiquitin. Arsenic treatment has

little effect on the modification of the A216T mutant, but triggers increased

conjugation by SUMO1, SUMO2/3 and ubiquitin for WT and L217F PML, with the

mutant showing a weaker response in SUMO1 and ubiquitin conjugation. Also

consistent with the Western blot data is the finding that SUMO1 has a larger relative

increase (5 fold) than SUMO2/3 (1.8 fold) in conjugation to WT PML after arsenic

treatment. To more easily compare multiple protein/peptide intensities, each

normalized intensity can be represented relative to the average intensity across all

samples (Supplementary Fig. 10H), which can then be plotted in a single compact

radial chart to show which differ the most between cell types and treatments

(compare with Fig. 6A).

Site-specific differences in SUMO modification to PML mutants

10 SUMO1 modifications to PML had reliable enough data for comparisons among conditions (Fig. 6B). After arsenic treatment, WT PML was much more extensively modified with SUMO1 than either mutant at all PML sites except K160 in B-Box1, which appeared to be more occupied by SUMO1 in the L217F mutant (Fig. 6B). Site-level analysis confirmed A216T PML was relatively unresponsive in SUMO1 conjugation. The results for SUMO2/3 were less clear-cut (Fig. 6C). Firstly, the scale of the difference among the three PML forms in the SUMO2 conjugation is much smaller than for SUMO1. Surprisingly, two of the 12 sites with good enough data for quantitative comparisons were most highly modified in the apparently inert A216T mutant (Fig. 6C). These were K226 close to the mutation hotspot and K487 in the NLS (Fig. 6C). Several other sites in or around the NLS, K400, K476, and K497, showed greater modification for L217F than WT PML (Fig. 6C). The WT protein only showed the most extensive SUMO2/3 modification at K380 (between the coiled-coil and NLS) and marginally higher conjugation at three sites; K65 (RING domain), K426 and K460 (both between the coiled-coil and the NLS). Site-level analysis of SUMO-SUMO branched peptides showed again that A216T PML was almost completely unresponsive (Fig. 6D). SUMO polymers did accumulate on L217F PML, but not to the same extent as for WT PML (Fig. 6D). Notably all bar one of the identified SUMO-SUMO linages contained SUMO1 as conjugate donor, acceptor, or both. Together these results show the higher levels of SUMO1 recruitment to WT PML compared to the mutants was via direct conjugation to PML, as well as SUMO1-SUMO polymer formation. Differences among the PML variants for SUMO2 conjuagation were more subtle and site-specific.

Neither A216T nor L217F PML accumulate polyubiquitin chains in response to arsenic

Of the 11 ubiquitination sites identified in PML, the majority were sites also identified as SUMO acceptors, except K183 between B-Boxes 1 and 2 and K337 in the coiled-coil domain. 9 sites had high quality quantitative data which showed that A216T did not respond to arsenic with increased ubiquitination at any site (Fig. 6E). In general WT and L217F PML showed similar global increases in ubiquitination on PML although there were differences at the site level. Notably both A216T and L217F PML were much more highly modified at lysines 160 and 490 than WT (Fig. 6E). These are two of the three sites originally identified as major SUMO acceptors, and likely to already have relatively high occupancy with SUMO. K183 and K337 were more occupied by ubiquitin in WT PML than either mutant (Fig. 6E). There is a striking difference between WT and L217F in the ubiquitination of the associated SUMO (Fig. 6F). 4 of the 5 ubiquitination sites in SUMO 1 and SUMO2/3 quantified were more extensively ubiquitinated in L217F PML purifications then the WT equivalent (Fig. 6F). Conversely, ubiquitin-ubiquitin conjugations were generally more abundant in WT PML purifications than for L217F PML (Fig. 6G). These results show that compared with WT PML, arsenic-induced ubiquitination of L217F PML is skewed towards SUMO ubiquitination, and away from ubiquitin-ubiquitin polymer formation.

SUMO1 is essential for arsenic-induced PML degradation

The finding that the stable L217F PML mutant is more defective in SUMO1 conjugation than SUMO2 is unexpected. RNF4 has been shown to have specificity

for polymeric SUMO conjugates, with a chain of 4 or more SUMOs having higher

affinity for RNF4 than shorter polymers or monomeric SUMO conjugates (Tatham,

Geoffroy et al. 2008). Due to the presence of an internal SUMO conjugation

consensus motif in SUMO2/3 but not SUMO1, it is assumed that SUMO2/3 form

branched conjugates more readily than SUMO1, and therefore represent the more

common RNF4 substrate. However, SUMO1 can still participate in polySUMO

conjugates as a SUMO donor, and proteomics studies have also found SUMO1 to

be an acceptor for SUMOylation, although the salience of this is poorly understood.

To investigate the role of SUMO1 in arsenic-induced PML degradation, a U2OS cell

line lacking SUMO1 (SUMO1 -/- U2OS) was generated by CRISPR-cas9 genome

editing. Immunofluorescence confirmed the loss of SUMO1 expression in these cells

(Fig. 7A). Anti-PML Western-blot analysis of crude cell extracts of cells exposed to

1µM As over 24 hr showed that in cells lacking SUMO1 there is a molecular weight

shift in PML in a manner similar to SUMO1 +/+ cells (Fig. 7B), but the modified forms

of PML appear to be more resistant to degradation. These persist up to 24 hr after

addition of As to the medium, when PML from WT cells is almost completely

degraded (Fig. 7B). Furthermore, after these extended periods of As exposure,

unconjugated PML begins to re-accumulate in SUMO1 - / cells (Fig. 7C), almost

returning the cellular pool of PML to the initial state (Compare 0h and 24h lanes in

Fig. 7C). This confirms SUMO1 is essential for PML degradation and insufficient

conjugation of the L217F PML mutant by SUMO1 potentially explains its stability

during arsenic exposure.

DISCUSSION

The overarching cellular mechanisms governing arsenic-induced degradation of PML

and the oncogenic fusion PML-RARA were broadly characterised more than ten

years ago (Lallemand-Breitenbach, Jeanne et al. 2008, Tatham, Geoffroy et al.

2008). Arsenic triggers the SUMOylation of PML, followed by its ubiquitination, then

degradation by the proteasome. However, the precise molecular signals and effector

proteins required to progress through each step are not fully understood. The

functional characterisation of mutant forms of PML and PML-RAR found in APL

patients refractory to arsenic treatment, provides an opportunity to better understand

the biochemical details of arsenic-induced PML degradation by understanding their

loss of function. We have used a combination of high content, time-lapse

microscopy, and peptide-level proteomic analysis to study two common patient-

derived PML mutants; A216T and L217F. We generated three model U2OS cell lines

lacking endogenous PML but expressing YFP fusions of WT PML, A216T PML or

L217F PML. In untreated cells the two mutants of PML formed atypical nuclear

bodies, fewer in number and larger in size than WT bodies, and both mutants were

stable when cells were exposed to $1\mu M$ arsenic for 24 hours. Importantly, during

arsenic treatment the modification status of each was quite different, and both

differed from WT PML.

After 2 hours exposure to $1\mu M$ arsenic, A216T PML appeared almost inert and

showed little increase in either SUMOylation or ubiquitination. Thus, A216T PML

stability is explained by a failed SUMOylation response early in the degradation

pathway. In contrast, arsenic did trigger SUMO and ubiquitin conjugation to L217F

PML, and yet it was not degraded. Site level proteomics revealed that while

ubiquitination did increase for L217F PML, it differed from WT PML in the precise

lysine sites to which it conjugated. The ubiquitin associated with L217F appeared to

favour attachment to SUMO molecules, while on WT-PML ubiquitin accumulated

more in polyubiquitin chains. This lack of polyubiquitin chain formation potentially

explains the observed lack of recruitment to L217F-PML of the segregase p97, which

recognises ubiquitin chains and is essential for PML degradation (Jaffray, Tatham et

al. 2023). p97 recognises polyubiquitin chains containing 5 or more copies of

ubiquitin (Twomey, Ji et al. 2019), and both K11 and K48 linkages are implicated in

p97 function (Locke, Toth et al. 2014, Meyer and Rape 2014, Yau, Doerner et al.

2017). Crucially, WT-PML showed statistically significant increases in ubiquitin-

ubiquitin linkages via K11, K33 and K48 (Supp. Fig. 11A). Although these increases

were modest, we expect an appropriate degradation signal for PML to be short-lived

in cells, making their detection difficult in studies like these.

Arsenic exposure triggered a much greater SUMO1 response on WT PML than it did

for SUMO2/3. From these studies we cannot determine whether more SUMO1 than

SUMO2/3 was conjugated to PML, but relative to initial conjugation levels, the

SUMO1 response was much greater than SUMO2/3. Furthermore, the L217F mutant

displayed lower levels of SUMO1 conjugation upon arsenic exposure than WT PML,

while total amounts of SUMO2/3 associated with both PML forms were comparable.

The importance of SUMO1 was further evidenced by the finding that WT PML is

stable in cells lacking SUMO1, suggesting that SUMO2/3 cannot compensate for its

loss. This raises the possibility that sub-optimal SUMO1 conjugation may lead to the

failure of L217F to generate the appropriate ubiquitin signal for p97 binding.

Based on the current model for PML degradation, critical requirement for SUMO1

seems counter intuitive. The E3 ligase responsible for PML ubiquitination, RNF4 is

poly-SUMO specific (Tatham, Geoffroy et al. 2008), and by lacking an internal

SUMO conjugation consensus, SUMO1 is thought not to form polySUMO chains as

readily as SUMO2/3. Incorporation of SUMO1 into SUMO2/3 polymers could

suppress further polymer formation by blocking lysines in SUMO2/3, and so SUMO1

may be expected to be inhibitory to PML degradation via RNF4. One potential

explanation is enough SUMOs in close proximity could substitute for a SUMO

polymer, and engage and activate RNF4 (Aguilar-Martinez and Sharrocks 2016).

This may be more likely for a protein like PML which forms multimeric complexes

and contains numerous SUMOylation sites. It is also conceivable that independent of

RNF4, SUMO1 provides a unique signal recognised by an as-yet unidentified protein

effector of the PML degradation pathway that specifically recognises SUMO1 and

not SUMO2/3. These prospects will require further investigation.

Alternatively, rather than generating a new signal, SUMO-1 may function to suppress

the formation of unproductive signals. For example SUMO1 may compete with

ubiquitin for conjugation to lysine residues on proteins other than ubiquitin itself, and

so promote polyubiquitin chain formation. In this way SUMO1 modification of lysines

from SUMOs attached to WT-PML or PML itself may block spurious ubiquitination

and effectively drive the synthesis of ubiquitin-ubiquitin linkages required for

degradation. L217F PML which is less SUMO1 modified than WT-PML might then

have more available non-ubiquitin lysines and so does not synthesise ubiquitin

chains as readily. Consistent with this, ubiquitination of SUMOs was found to be

generally higher for L217F PML than WT (Fig. 6F), and SUMO1-SUMO linkages

were more abundant in WT-PML preps than L217F (Fig. 6D).

In summary, two patient-derived mutant forms of PML, A216T and L217F result in

similar clinical aetiologies but have subtly different biochemical responses to arsenic

resulting in blocked degradation (See Supplementary figure 12). A216T is almost

completely unresponsive to arsenic, experiencing only very modest changes in

SUMOylation and ubiquitination, most likely because it does not engage the

appropriate SUMO conjugation machinery. L217F PML does become modified with

both SUMO1 and SUMO2/3, but while ubiquitination is stimulated, it does not create

the appropriate signal required to recruit p97 prior to degradation by the proteasome.

These results reveal the subtle intricacies of the PML degradation pathway, and

propose a higher level of complexity than previously considered.

MATERIALS AND METHODS

Preparation of anti-GFP nanobody magnetic beads. 300mg of Dynabeads M-270

Epoxy (ThermoFisher Scientific Catalogue No. 14302D) were resuspended in 10 ml

of 0.1 M sodium phosphate buffer pH 7.5 and washed twice using a DynaMag-15

magnet (ThermoFisher Scientific Catalogue No. 12301D) with 10 ml of the same

buffer. 4 mg of the LaG16 nanobody that recognises GFP (Fridy, Li et al. 2014) in 10

ml of 1 M $(NH_4)_2SO_4$, 0.1 M sodium phosphate buffer pH 8.0 was added to the beads

which were incubated overnight at 27ºC with end over end rotation. After coupling was

complete the beads were washed 6 times with phosphate buffered saline (PBS) and

stored in PBS, 0.1% NaN_3 at 4^0C.

Purification of YFP-PML from U2OS PML-/- YFP-PML cells. For each 'replicate' of

the proteomics experiment five 15 cm diameter plates of approximately 80% confluent

cells were used. Prior to harvesting, growth medium was removed and cells washed

in situ twice with 5mL PBS/100mM iodoacetamide. For each dish, cells were scraped

into 5mL PBS+100mM iodoacetamide and transferred to a 50mL tube. A second 2mL

volume of PBS+100mM iodoacetamide was used to clean the plate of all cells and

pooled with the first 5mL. This was repeated for all dishes for each replicate and

pooled in the same 50mL tube (~35mL total). Cells were pelleted by centrifugation at

400g for 5 minutes at 22ºC and resuspended in 10 mL 4ºC hypotonic buffer including

iodoacetamide (10mM HEPES pH 7.9, 1.5mM MgCl2, 10mM KCl, 0.08% NP-40, 1x

EDTA-free protease inhibitor cocktail (Roche), + 100mM Iodoacetamide). A sample of

this may be retained as a "Whole cell extract". These were snap-frozen in liquid

nitrogen and stored at -80ºC until required. Cells were thawed by tube rotation at 4ºC

for 30 minutes. Tubes were then centrifuged at 2000g, 4ºC for 10 minutes to pellet

nuclei. Supernatants were discarded but a sample of this can be taken as a "Cytosolic

fraction". Each pellet of nuclei was resuspended in 5mL ice-cold hypotonic buffer

containing 20mM DTT but without iodoacetamide, transferred to a 15mL tube and

centrifuged at 2000g, 4ºC for 10 minutes. The nuclei were washed once more in 5mL

ice-cold hypotonic buffer (without iodoacetamide or DTT) and the supernatant

discarded. To digest DNA, nuclei were resuspended in 5mL "Buffer A" (50mM Tris pH

7.5, 150mM NaCl, 0.2mM DTT, 1µg/ml benzonase, 1x EDTA-free protease inhibitor

cocktail (Roche)) and incubated on ice for 15 minutes. Samples were sonicated in

15mL tubes on ice (Branson Sonifier) using the small probe, 6 cycles of 30 seconds

at approximately 50% amplitude with at least 2 minutes cooling on ice in between

cycles. PML bodies should remain intact at this point. Insoluble debris was removed

by centrifugation at 1200g for 20 minutes at 4ºC, and the supernatant containing

soluble PML bodies, transferred to a new 15ml centrifuge tube. A sample of this can

be taken as a "Nuclear fraction". To each 5mL supernatant 6mL "Buffer B" was added

(50mM Tris pH 7.5, 1% Triton-X-100, 0.1% Deoxycholic acid, 1x EDTA-free protease

inhibitor cocktail (Roche)). To each 11mL solution, 1mL volume of a 5% beads:buffer

slurry (v:v) of magnetic anti-GFP nanobody beads, pre-equilibrated in 1:1.4 Buffer A:B

mix (v:v), was added and mixed on a tube roller at 4ºC for 16h (50μL beads per

sample). Using the DynaMag rack (Thermofisher scientific) the beads were extracted

from suspension and the supernatant removed. A sample of this can be analysed to

ensure YFP-PML depletion. Beads were then washed with 5mL "Buffer C" (50mM Tris

pH 7.5, 500mM NaCl, 0.2mM DTT) and the beads extracted. These beads were then

resuspended in 1mL "Buffer D" (50mM Tris pH 7.5, 150mM NaCl, 0.2mM DTT) and

transferred to a new 1.5mL protein LoBind tube (Eppendorf) and after bead separation,

washed once more with 1mL Buffer D. One half of this final sample of beads (25μL

beads) was eluted for proteomic analysis by adding 35μl 1.2X NuPAGE LDS sample

buffer (Thermofisher scientific), and incubated at 70ºC for 15 minutes with agitation,

followed by fractionation by SDS-PAGE. If Western blot analysis was also required,

the second half of the resin (25μL) would be resuspended in 250μl 1.2X NuPAGE LDS

sample buffer (Thermo) and incubated at 70ºC for 15 minutes with agitation.

Approximately 30μl would be used per gel lane. If treatment with proteases was

required to monitor PML modification status, then the second half volume of beads

(25μL) was divided into four and treated as follows: 1-Add 150 μl Buffer D. 2-Add 150

μl 250nM SENP1 in Buffer D. 3- Add 150 μl 500nM USP2 in Buffer D. 4- Add 150 μl

250nM SENP1/500nM USP2 in Buffer D. All tubes were agitated for 1-2 hrs at 22ºC,

supernatants removed (and can be kept for analysis) and beads washed twice with

1mL buffer D before elution with 250 μl 1.2X LDS, 70ºC, 15 minutes with agitation.

Proteomic analysis of YFP-PML purifications. Elutions from anti-GFP nanobody

beads in 1.2x LDS/60mM DTT protein sample buffer were fractionated on 4-12%

polyacrylamide Bis-Tris NuPAGE gels (Thermofisher Scientific) using MOPS running

buffer. After Coomassie staining and destaining gels were excised into 4 slices per

lane and tryptic peptides extracted from top slice (>100kDa) as previously described

(Shevchenko, Tomas et al. 2006). Peptides were resuspended in 40μL 0.1% TFA

0.5% acetic acid and half STAGE tip purified prior to digestion with GluC. This gave

two fractions per replicate: Trypsin digested, and Trypsin+GluC digested. 6μL of each

was analysed by LC-MS/MS. This was performed using a Q Exactive mass

spectrometer (Thermofisher Scientific) coupled to an EASY-nLC 1000 liquid

chromatography system (Thermofisher Scientific), using an EASY-Spray ion source

(Thermofisher Scientific) running a 75 μm x 500 mm EASY-Spray column at 45ºC. A

150 minute elution gradient with a top 8 data-dependent method was applied. Full

scan spectra (m/z 300–1800) were acquired with resolution R = 70,000 at m/z 200

(after accumulation to a target value of 1,000,000 ions with maximum injection time of

20 ms). The 8 most intense ions were fragmented by HCD and measured with a

resolution of R = 17,500 at m/z 200 (target value of 500,000 ions and maximum

injection time of 120 ms) and intensity threshold of 2.1×10^4. Peptide match was set to

'preferred', a 40 second dynamic exclusion list was applied and ions were ignored if

they had unassigned charge state 1, 8 or >8.

MS data were processed in MaxQuant version 1.6.1.0 (Cox and Mann 2008, Cox,

Neuhauser et al. 2011). Multiple MaxQuant runs were performed considering multiple

SUMO1 and SUMO2/3 derived adducts as variable modifications (see Supplementary

Datafile 1 for details), as well as those typical for ubiquitin and phosphorylation to Ser

and Thr residues. A MaxQuant search for SUMO adducts was also conducted using

a concatenated database approach using linear peptide fusions of SUMO C-termini to

substrate peptides as search library (Matic, van Hagen et al. 2008). Oxidised

methionine and acetylated protein N-termini were also selected as variable

modifications and carbamidomethyl-C was the only fixed modification. Maximum

number of variable modifications was set to 4. The match between runs option was

enabled, which matches identified peaks among slices from the same position in the

gel as well as one slice higher or lower. The uniport human proteome database

(downloaded 19/4/2019 - 73920 entries) digested with Trypsin/P (maximum missed

cleavages was 5) was used as search space. LFQ intensities were required for each

slice. All FDR filtering was set to 1%. For peptides derived from GluC digestions 8

maximum missed cleavages were considered. Raw peptide intensities were manually

normalized by the total YFP peptide intensity for each sample. Mass spectrometry

data from samples derived from the same replicate were numerically summed to

provide a final normalized peptide intensity for each identification in each sample.

Downstream data processing used Perseus v1.6.1.1 (Tyanova, Temu et al. 2016).

Zero intensity values were replaced from $\log_2$ transformed data (width 0.3 and

downshift 1.8).

Pre-extraction of cells. Cells were grown on coverslips in 24 well plates. Coverslips

were washed twice with PBS before being exposed to pre-extraction buffer (25 mM

HEPES pH 7.4, 50 mM NaCl, 3 mM MgCl2, 0.5% Triton-X-100, 0.3M Sucrose) for 10

min. Cells were then processed through the procedure below for immunofluorescence.

Immunofluorescence analysis. Adherent cells were washed three times with PBS,

fixed with 4% Formaldehyde in PBS for 15mins, then washed again three times with

PBS before blocking for 30 min in 5% BSA, 0.1% TWEEN 20 in PBS. Fixed cells were

washed once with 1% BSA, 0.1% TWEEN 20 in PBS before incubation for 1-2 hr with

primary antibody in 1% BSA, 0.1% TWEEN 20 in PBS. These were chicken anti-PML

(in house) and sheep anti-p97 (MRC Dundee). After 3 washes in 1% BSA, 0.1%

TWEEN 20 in PBS cells were incubated with secondary antibody in 1% BSA, 0.1%

TWEEN 20 in PBS for 60 min., washed three times in 1% BSA, 0.1% TWEEN 20 in

PBS and stained with 0.1µg/ml DAPI for 2 min. These were Alexa Fluor (Invitrogen)

Donkey anti-chicken 594, donkey anti-sheep 594 and donkey anti-chicken 488. Cover

slips were rinsed twice with PBS, twice with water and dried before mounting in

MOWIOL. Images were collected using a DeltaVision DV3 widefield microscope and

processed using Softworx (both Applied Precision, Issaquah, WA). Images are

presented as maximal intensity projections using Omero software. For quantitative

p97-PML colocalization analysis the same image settings were applied across all

fields and PML bodes per cell were counted for between 20 and 40 cells per condition.

P97 colocalisation was defined as any p97 signal above background within the same

region as a PML body. Percent colocalization was then calculated for each cell.

Live-Cell Imaging in Real Time and Measurement of Fluorescence Intensities.

For all live-cell imaging experiments, cells were seeded onto µ-slide 8 well glass

bottom (Ibidi 80827). Immediately prior to imaging cell medium was replaced with

DMEM minus phenol red (gibco 31053-038) supplemented with 2mM glutamine and

10% fetal bovine serum. Cells were untreated or treated with 1µM arsenic. Time-laspe

microscopy was performed on a DeltaVision Elite restoration microscope (Cytiva)

fitted with an incubation chamber (Solent Scientific) set to 37 ºC and 5% CO2 and a

cooled charge-coupled device camera (CoolSnap HQ; Roper). SoftWorx software was

used for image collection. Datasets were deconvolved using the constrained iterative

algorithm (Swedlow, Sedat et al. 1996, Wallace, Schaefer et al. 2001) using SoftWorx

software. Using a YFP specific filter set (Ex510/10nm, Em537/26nm), 5 z-planes

spaced at 2 µm were taken every 15 min. A single brightfield reference image was

taken at each time point to monitor cell health. Time courses were presented as

maximum intensity projections of deconvolved 3D datasets. Movies were created in

OMERO using the deconvolved images.

Quantitative analysis of PML body composition during Live-Cell imaging

Batch analysis of time-lapse movies was carried out using ImageJ (Schneider,

Rasband et al. 2012) macros in Fiji (Schindelin, Arganda-Carreras et al. 2012)

(Supplementary Materials). Briefly: 1) datasets (deconvolved with SoftWoRx) were

cropped to remove border artefacts, 2) for each timepoint the maximum focus slice

was selected according to intensity variance (radius 2 pixels), 3) auto-thresholding

was performed using a threshold of mean intensity + 3 standard deviations, and 4)

ImageJ's built-in "Analyze Particles" function was used to measure PML body count,

size, total area, and intensity. Results were aggregated and summarized using R

scripts as described previously (Jaffray, Tatham et al. 2023).

Preparation of cell extracts and Western blotting. Adherent cells were washed with

PBS and lysed by the addition of 1.2X LDS buffer and heated to 70ºC for 10 min. Cell

lysates were sonicated using a probe sonicator for two bursts of 30 sec. Samples were

run on NuPage 3-8% or 4-12% Bis-Tris precast gels (ThermoFisher Scientific) and

transferred to nitrocellulose. Membranes were blocked in PBS with 5% non-fat milk

and 0.1% Tween-20. Primary antibody incubations were performed in PBS with 3%

BSA and 0.1% Tween-20. For the HRP-coupled secondary antibody incubations 5%

milk was used instead of the BSA. The signal was detected by Pierce enhanced

chemiluminescence (ThermoFisher scientific 32106) and X-ray films. Antibodies.

Chicken anti-PML (homemade), sheep anti-SUMO-1 (homemade), rabbit anti-SUMO-

2 (Cell signaling 4971S), rabbit FK2 ubiquitin (Enzo BML-PW8810-0500), rabbit anti-

p97 (Abcam ab109240). HRP-coupled secondary anti-mouse, anti-rabbit anti-chicken

and anti-sheep were purchased from Sigma.

FIGURE LEGENDS

Figure 1. Characterization of PML bodies in U2OS PML-/- cell lines expressing

YFP-fusions of WT, A216T and L217F forms of PML-V. A. Schematic depiction of

PML-V primary structure indicating the major domains, regions of common mutations

associated with arsenic treatment resistance, and the common breakpoints leading

to PML-RARA fusions. B. Representative images of the indicated PML-/- U2OS +

YFP-PML-V cell lines C. High content imaging data summarizing the calculated

values for total PML intensity per cell, PML body number per cell and area per PML

body for each cell line. Scatter plots are shown with median (solid line) and quartiles

(error bars). n=38411 (wt), 35210 (A216T) and 41288 (L217F).

Figure 2. A216T and L217F mutants of PML are not rapidly degraded in

response to arsenic treatment in vitro. A. YFP fluorescence image stills from one

live cell analysis of each YFP-PML cell type at both 0h and 15h treatment with 1μM

arsenic. B. Averaged data of 9 or 10 fields of view (median 12 cells per field)

followed over a 15 hour time course of treatment or not with 1μM arsenic (15 minute

intervals). Intensity measurements are normalized by the t=0hr values. C. Summary

statistics (n=9 or 10) for 15hr data comparing 1μM arsenic treated cells with

untreated cells. P-values are unpaired, two-tailed student's T-tests using Welch's

correction for unequal variances where appropriate (§). See Supplementary movie

files 1-6 for real-time data of a single representative field. D. Anti-PML Western-blots

of fractionated whole cell extracts showing time-courses of the response of YFP-

PML-V variants to 1μM arsenic exposure over 24 hours.

Figure 3. In response to arsenic A216T PML fails to recruit RNF4. Two separate

purifications of YFP-PMLV WT, A216T and L217F (A+B): PML bodies were enriched

from U2OS PML-/- + YFP-PMLV WT, A216T or L217F cells, either nontreated (NT)

or treated with 1μM arsenic (1μM As 2h) for 2 hr. Bound material was eluted and

analysed by Western blotting using antibodies to (A) PML, SUMO1, SUMO2/3 and

(B) PML, RNF4 and ubiquitin.

Figure 4. A216T and L217F mutants of PML do not recruit p97 in response to

arsenic treatment. A. Immunofluorescence images showing YFP-PMLV and p97 in

the indicated cell lines after 6 hours arsenic treatment. Data showing merged images

for these and the 0h time-point data can be seen in supplementary figure 4. B.

Quantitative summary of p97 colocalization with PML bodies in the indicated cell

lines during 0h, 2h, 4h and 6h arsenic exposure. Markers show average values and

error bars are standard error of the mean. C. Statistical analysis of the comparison

between 0h and 6h arsenic exposure for PML-p97 colocalization. Columns represent

average % colocalization, error bars are standard error of the mean, cell count is

shown in brackets below each column and student's two-tailed t-test p-values are

shown above.

Figure 5. Failure of arsenic resistant mutants to generate the SUMO and

ubiquitin modified intermediates required for degradation. PML bodies were

enriched from U2OS PML-/- + YFP-PMLV WT, A216T or L217F cells, either

untreated (-) or treated with 1μM arsenic (+) for 2 hr. Purified PML bodies bound to

anti-GFP nanobody magnetic beads were either untreated (No protease), treated

with SUMO specific protease (SENP1), treated with ubiquitin specific protease

(USP2) or a combination of both (USP2 + SENP1). Material remaining bound to the

beads after treatment was eluted and analysed by Western blotting using antibodies

to A. PML, B. SUMO1, C. SUMO2/3 and D. ubiquitin.

Figure 6. Differential site-specific modifications of WT, A216T and L217F PML

forms before and after arsenic. PML bodies were enriched from U2OS PML-/- +

YFP-PMLV WT, A216T or L217F cells and quantitatively analysed by mass

spectrometry for protein content and site-specific modifications by SUMO1,

SUMO2/3 and ubiquitin. A-G. Radial plots showing relative intensity for each of the

indicated proteins or peptides. A. is derived from the total of the unmodified peptides

for YFP, PML, SUMO1, SUMO2/3 and ubiquitin and is a proxy for of total protein

abundance. Circles indicate the upper limit of the range, with the maximum relative

intensity in brackets.

Figure 7. Effect of SUMO1 deletion on arsenic-induced WT PML degradation. A.

Wild-type (SUMO1 +/+) and SUMO1 knockout (SUMO1 -/-) U2OS cells were

exposed to 1μM As for 2 hours to allow SUMO1 recruitment to PML bodies, fixed

and double stained with DAPI and anti-SUMO1 (FITC) before image acquisition

using InCell analyser 2200 high-content imaging microscope and differential image

visualization by ImageJ software. B. SUMO1 +/+ and SUMO1 -/- U2OS cells were

exposed to 1μM As for the indicated periods before lysis and analysis by Western

blot for PML. C. As in B except only for SUMO -/- cells and with more time-points.

Supplementary Figure 1. Arsenic-induced colocalisation of SUMO1 with WT, A216T and L217F forms of YFP-PML. Colocalisation of SUMO1 with WT, A216T and L217F PML variants after 1μM arsenic treatment for 2h studied by fluorescence microscopy. Scale bar is 10μm.

Supplementary Figure 2. Arsenic-induced SUMO2/3 colocalisation with WT, A216T and L217F forms of YFP-PML. Colocalisation of SUMO2/3 with WT, A216T and L217F PML variants after 1μM arsenic treatment for 2h studied by fluorescence microscopy. Scale bar is 10μm.

Supplementary Figure 3. Arsenic-induced RNF4 colocalisation with WT, A216T and L217F forms of YFP-PML. Colocalisation of RNF4 with WT, A216T and L217F PML variants after 1μM arsenic treatment for 2h studied by fluorescence microscopy. Scale bar is 10μm.

Supplementary Figure 4. Arsenic-induced p97 colocalisation with WT, A216T and L217F forms of YFP-PML. Representative immunofluorescence images for 1μM arsenic treated YFP-PMLV cells at 0 hr and 6 hr. Monochrome images for YFP-PML and p97 channels are shown along with the merge of YFP-PMLV (green) and p97 (red) with DAPI (blue).

Supplementary Figure 5. p97 expression levels do not differ between cell types or upon arsenic treatment. Western blots for p97 and PML of whole cell extracts from the indicated cell lines either treated or not with arsenic.

Supplementary Figure 6. Treatment of purified PML bodies with recombinant SUMO and Ubiquitin proteases reveals the nature of PML post-translational modifications. A. Overview of the experimental design for detection of ubiquitin and SUMO conjugated to PML. B. Anti-PML Western blot for nuclear extracts from the indicated cell lines before and after purification of YFP-PML with GFP nanobody beads. C. Schematic depiction of the contents of YFP-PML bound to beads and resulting supernatants after treatment with specific proteases.

Supplementary Figure 7. Western blot analysis of the material eluted by SENP1 and USP2 from the purified YFP-PML forms indicated. Associated with Figure 5. Western blots using antibodies specific for PML (A), SUMO1 (B), SUMO2/3 (C) and ubiquitin (D). 2ng standards of recombinant proteins (Rec.) were included with the gels. The positions of recombinant PML (Rec.PML), unmodified forms of YFP-PML, SUMO1, SUMO2 and ubiquitin are indicated, along with ubiquitinated species (*).

Supplementary Figure 8. Known sites of PML SUMOylation and potential peptide adducts remaining on substrates after trypsin and GluC digestion. A. Positions of previously identified SUMO conjugation sites relative to the structural domains in PML. Isoform V is shown although identifications could be from other isoforms. Data taken from Hendriks et al 2017 (Hendriks, Lyon et al. 2017). Common breakpoints found in the PML-RARA and mutational hotspots found in arsenic insensitive forms are indicated. B. Overview of the potential remnants of SUMO C-termini attached to substrate lysine residues (K) when conjugates are cleaved by Trypsin, or GluC considering multiple missed cleavages.

Supplementary Figure 9. A proteomic study to quantitatively analyse arsenic-induced changes to PML post-translational modifications at the site level. A. Overview of the proteomic experimental design. B. Coomassie-stained gel showing the elutions from anti-GFP nanobody purifications for each replicate. Gel slices are numbered. C. Anti-PML western blot of a fraction of the anti-GFP nanobody purifications for each replicate.

Supplementary Figure 10. Overview of the proteomic data for the post-translational modification of PML WT, A216T and L217F

A. Summary of SUMO and ubiquitin conjugation sites identified in this study. Region of mutation hotspots is boxed. B. Principal component analysis of the quantitative proteomic data using only peptides from YFP-PML, SUMOs or ubiquitin. These included all unmodified peptides, phospho peptides, Ubiquitin branches and SUMO branches on YFP-PML, SUMO1, SUMO2/3 and Ubiquitin. Total is 343 peptides and uses $\log_2$ transformed and zero-replaced intensity values. C-G. Total unmodified peptide intensity data for the indicted proteins. Bars are average intensities with standard deviation error bars. p values are for two tailed student's t-test comparing untreated with arsenic treated samples. n=4. H. Summary of the data shown in C-G normalized by the average intensity for each protein.

Supplementary Figure 11. Column plots of relative peptide intensity data for selected peptide types. A. Quantitative data for peptides indicative of Ubiquitin-Ubiquitin linkages detected in the proteomics study. Columns are average values with SD error bars. Student's t-test p-values are indicated.

Supplementary Figure 12. Schematic summary of the arsenic-induced

responses of WT, A216T and L217F forms of PML. WT, A216T and L217F forms

of PML are modified by broadly similar amounts of SUMO1 and SUMO2/3 in

untreated cells. Exposure of cells to arsenic triggers an increase in SUMO2/3

conjugation for WT PML and L217F PML, and a much smaller increase for A216T

PML. A large increase in SUMO1 conjugation is triggered for WT PML, with a

smaller increase for L217F PML, and no significant SUMO1 change for A216T.

Arsenic does not induce increased ubiquitination of A216T, while L217F experiences

increased ubiquitination to PML and SUMO proteins. WT PML also becomes

ubiquitinated on PML and SUMOs, but a fraction also forms ubiquitin-ubiquitin

polymers. Neither A216T PML nor L217F PML develop the requisite SUMO and/or

ubiquitin signals required to bind p97 and target them for degradation.

REFERENCES

Aguilar-Martinez, E. and A. D. Sharrocks (2016). "The Use of Multimeric Protein Scaffolds for Identifying Multi-SUMO Binding Proteins." Methods Mol Biol **1475**: 195-204.

Alfonso, V., L. Iaccarino, T. Ottone, L. Cicconi, S. Lavorgna, M. Divona, R. Cairoli, A. Cristiano, C. Ciardi, S. Travaglini, G. Falconi, S. K. Hasan, A. Venditti, W. Arcese, M. T. Voso and F. Lo-Coco (2019). "Early and sensitive detection of PML-A216V mutation by droplet digital PCR in ATO-resistant acute promyelocytic leukemia." Leukemia **33**(6): 1527-1530.

Balasundaram, N., S. Ganesan, E. Chendamarai, H. K. Palani, A. Venkatraman, A. A. Alex, S. David, S. P. Kumar, N. R. Radhakrishnan, M. Yasar, S. Krishna, A. Korula, U. Kulkarni, N. B. Janet, P. Balasubramanian and V. Mathews (2022). "Metabolic adaptation drives arsenic trioxide resistance in acute promyelocytic leukemia." Blood Adv **6**(2): 652-663.

Bernardi, R. and P. P. Pandolfi (2007). "Structure, dynamics and functions of promyelocytic leukaemia nuclear bodies." Nat Rev Mol Cell Biol **8**(12): 1006-1016.

Chendamarai, E., S. Ganesan, A. A. Alex, V. Kamath, S. C. Nair, A. J. Nellickal, N. B. Janet, V. Srivastava, K. M. Lakshmi, A. Viswabandya, A. Abraham, M. Aiyaz, N. Mullapudi, R. Mugasimangalam, R. A. Padua, C. Chomienne, M. Chandy, A. Srivastava, B. George, P. Balasubramanian and V. Mathews (2015). "Comparison of newly diagnosed and relapsed patients with acute promyelocytic leukemia treated with arsenic trioxide: insight into mechanisms of resistance." PLoS One **10**(3): e0121912.

Cox, J. and M. Mann (2008). "MaxQuant enables high peptide identification rates,
individualized p.p.b.-range mass accuracies and proteome-wide protein quantification." <u>Nat</u>
<u>Biotechnol</u> **26**(12): 1367-1372.

Cox, J., N. Neuhauser, A. Michalski, R. A. Scheltema, J. V. Olsen and M. Mann (2011).
"Andromeda: a peptide search engine integrated into the MaxQuant environment." <u>J</u>
<u>Proteome Res</u> **10**(4): 1794-1805.

de The, H., C. Lavau, A. Marchio, C. Chomienne, L. Degos and A. Dejean (1991). "The PML-
RAR alpha fusion mRNA generated by the t(15;17) translocation in acute promyelocytic
leukemia encodes a functionally altered RAR." <u>Cell</u> **66**(4): 675-684.

Dyck, J. A., G. G. Maul, W. H. Miller, Jr., J. D. Chen, A. Kakizuka and R. M. Evans (1994). "A
novel macromolecular structure is a target of the promyelocyte-retinoic acid receptor
oncoprotein." <u>Cell</u> **76**(2): 333-343.

Fridy, P. C., Y. Li, S. Keegan, M. K. Thompson, I. Nudelman, J. F. Scheid, M. Oeffinger, M. C.
Nussenzweig, D. Fenyo, B. T. Chait and M. P. Rout (2014). "A robust pipeline for rapid
production of versatile nanobody repertoires." <u>Nat Methods</u> **11**(12): 1253-1260.

Geoffroy, M. C. and R. T. Hay (2009). "An additional role for SUMO in ubiquitin-mediated
proteolysis." <u>Nat Rev Mol Cell Biol</u> **10**(8): 564-568.

Goto, E., A. Tomita, F. Hayakawa, A. Atsumi, H. Kiyoi and T. Naoe (2011). "Missense
mutations in PML-RARA are critical for the lack of responsiveness to arsenic trioxide
treatment." <u>Blood</u> **118**(6): 1600-1609.

Grignani, F., P. F. Ferrucci, U. Testa, G. Talamo, M. Fagioli, M. Alcalay, A. Mencarelli, F.
Grignani, C. Peschle, I. Nicoletti and et al. (1993). "The acute promyelocytic leukemia-
specific PML-RAR alpha fusion protein inhibits differentiation and promotes survival of
myeloid precursor cells." <u>Cell</u> **74**(3): 423-431.

Hendriks, I. A., D. Lyon, C. Young, L. J. Jensen, A. C. Vertegaal and M. L. Nielsen (2017). "Site-
specific mapping of the human SUMO proteome reveals co-modification with
phosphorylation." <u>Nat Struct Mol Biol</u> **24**(3): 325-336.

Iaccarino, L., T. Ottone, V. Alfonso, L. Cicconi, M. Divona, S. Lavorgna, S. Travaglini, A.
Ferrantini, G. Falconi, C. Baer, M. Usai, F. Forghieri, A. Venditti, M. I. Del Principe, W. Arcese,
M. T. Voso, T. Haferlach and F. Lo-Coco (2019). "Mutational landscape of patients with acute
promyelocytic leukemia at diagnosis and relapse." <u>Am J Hematol</u> **94**(10): 1091-1097.

Iaccarino, L., T. Ottone, M. Divona, L. Cicconi, R. Cairoli, M. T. Voso and F. Lo-Coco (2016).
"Mutations affecting both the rearranged and the unrearranged PML alleles in refractory
acute promyelocytic leukaemia." <u>Br J Haematol</u> **172**(6): 909-913.

Jaffray, E. G., M. H. Tatham, B. Mojsa, M. Liczmanska, A. Rojas-Fernandez, Y. Yin, G. Ball and
R. T. Hay (2023). "The p97/VCP segregase is essential for arsenic-induced degradation of
PML and PML-RARA." <u>J Cell Biol</u> **222**(4).

Kakizuka, A., W. H. Miller, Jr., K. Umesono, R. P. Warrell, Jr., S. R. Frankel, V. V. Murty, E.
Dmitrovsky and R. M. Evans (1991). "Chromosomal translocation t(15;17) in human acute
promyelocytic leukemia fuses RAR alpha with a novel putative transcription factor, PML."
<u>Cell</u> **66**(4): 663-674.

Kamitani, T., K. Kito, H. P. Nguyen, H. Wada, T. Fukuda-Kamitani and E. T. Yeh (1998).
"Identification of three major sentrinization sites in PML." <u>J Biol Chem</u> **273**(41): 26675-
26682.

Kwok, C., B. B. Zeisig, S. Dong and C. W. So (2006). "Forced homo-oligomerization of
RARalpha leads to transformation of primary hematopoietic cells." <u>Cancer Cell</u> **9**(2): 95-108.

Lallemand-Breitenbach, V., M. Jeanne, S. Benhenda, R. Nasr, M. Lei, L. Peres, J. Zhou, J. Zhu,
B. Raught and H. de The (2008). "Arsenic degrades PML or PML-RARalpha through a SUMO-
triggered RNF4/ubiquitin-mediated pathway." Nat Cell Biol **10**(5): 547-555.
Lehmann-Che, J., C. Bally and H. de The (2014). "Resistance to therapy in acute
promyelocytic leukemia." N Engl J Med **371**(12): 1170-1172.
Liu, J., H. H. Zhu, H. Jiang, Q. Jiang and X. J. Huang (2016). "Varying responses of PML-RARA
with different genetic mutations to arsenic trioxide." Blood **127**(2): 243-250.
Lo-Coco, F., G. Avvisati, M. Vignetti, C. Thiede, S. M. Orlando, S. Iacobelli, F. Ferrara, P. Fazi,
L. Cicconi, E. Di Bona, G. Specchia, S. Sica, M. Divona, A. Levis, W. Fiedler, E. Cerqui, M.
Breccia, G. Fioritoni, H. R. Salih, M. Cazzola, L. Melillo, A. M. Carella, C. H. Brandts, E. Morra,
M. von Lilienfeld-Toal, B. Hertenstein, M. Wattad, M. Lubbert, M. Hanel, N. Schmitz, H. Link,
M. G. Kropp, A. Rambaldi, G. La Nasa, M. Luppi, F. Ciceri, O. Finizio, A. Venditti, F. Fabbiano,
K. Dohner, M. Sauer, A. Ganser, S. Amadori, F. Mandelli, H. Dohner, G. Ehninger, R. F.
Schlenk, U. Platzbecker, d. A. Gruppo Italiano Malattie Ematologiche, G. German-Austrian
Acute Myeloid Leukemia Study and L. Study Alliance (2013). "Retinoic acid and arsenic
trioxide for acute promyelocytic leukemia." N Engl J Med **369**(2): 111-121.
Locke, M., J. I. Toth and M. D. Petroski (2014). "Lys11- and Lys48-linked ubiquitin chains
interact with p97 during endoplasmic-reticulum-associated degradation." Biochem J **459**(1):
205-216.
Lou, Y., Y. Ma, J. Sun, X. Ye, H. Pan, Y. Wang, W. Qian, H. Meng, W. Mai, J. He, H. Tong and J.
Jin (2015). "Evaluating frequency of PML-RARA mutations and conferring resistance to
arsenic trioxide-based therapy in relapsed acute promyelocytic leukemia patients." Ann
Hematol **94**(11): 1829-1837.
Madan, V., P. Shyamsunder, L. Han, A. Mayakonda, Y. Nagata, J. Sundaresan, D. Kanojia, K.
Yoshida, S. Ganesan, N. Hattori, N. Fulton, K. T. Tan, T. Alpermann, M. C. Kuo, S. Rostami, J.
Matthews, M. Sanada, L. Z. Liu, Y. Shiraishi, S. Miyano, E. Chendamarai, H. A. Hou, G.
Malnassy, T. Ma, M. Garg, L. W. Ding, Q. Y. Sun, W. Chien, T. Ikezoe, M. Lill, A. Biondi, R. A.
Larson, B. L. Powell, M. Lubbert, W. J. Chng, H. F. Tien, M. Heuser, A. Ganser, M. Koren-
Michowitz, S. M. Kornblau, H. M. Kantarjian, D. Nowak, W. K. Hofmann, H. Yang, W. Stock,
A. Ghavamzadeh, K. Alimoghaddam, T. Haferlach, S. Ogawa, L. Y. Shih, V. Mathews and H. P.
Koeffler (2016). "Comprehensive mutational analysis of primary and relapse acute
promyelocytic leukemia." Leukemia **30**(12): 2430.
Martens, J. H., A. B. Brinkman, F. Simmer, K. J. Francoijs, A. Nebbioso, F. Ferrara, L. Altucci
and H. G. Stunnenberg (2010). "PML-RARalpha/RXR Alters the Epigenetic Landscape in
Acute Promyelocytic Leukemia." Cancer Cell **17**(2): 173-185.
Matic, I., M. van Hagen, J. Schimmel, B. Macek, S. C. Ogg, M. H. Tatham, R. T. Hay, A. I.
Lamond, M. Mann and A. C. Vertegaal (2008). "In vivo identification of human small
ubiquitin-like modifier polymerization sites by high accuracy mass spectrometry and an in
vitro to in vivo strategy." Mol Cell Proteomics **7**(1): 132-144.
Meyer, H. J. and M. Rape (2014). "Enhanced protein degradation by branched ubiquitin
chains." Cell **157**(4): 910-921.
Mi, J. Q., S. J. Chen, G. B. Zhou, X. J. Yan and Z. Chen (2015). "Synergistic targeted therapy
for acute promyelocytic leukaemia: a model of translational research in human cancer." J
Intern Med **278**(6): 627-642.
Mikesch, J. H., H. Gronemeyer and C. W. So (2010). "Discovery of novel transcriptional and
epigenetic targets in APL by global ChIP analyses: Emerging opportunity and challenge."
Cancer Cell **17**(2): 112-114.

Muller, S., M. J. Matunis and A. Dejean (1998). "Conjugation with the ubiquitin-related
modifier SUMO-1 regulates the partitioning of PML within the nucleus." <u>Embo J</u> **17**(1): 61-
70.
Perry, J. J., J. A. Tainer and M. N. Boddy (2008). "A SIM-ultaneous role for SUMO and
ubiquitin." <u>Trends Biochem Sci</u> **33**(5): 201-208.
Rojas-Fernandez, A., A. Plechanovova, N. Hattersley, E. Jaffray, M. H. Tatham and R. T. Hay
(2014). "SUMO chain-induced dimerization activates RNF4." <u>Mol Cell</u> **53**(6): 880-892.
Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C.
Rueden, S. Saalfeld, B. Schmid, J. Y. Tinevez, D. J. White, V. Hartenstein, K. Eliceiri, P.
Tomancak and A. Cardona (2012). "Fiji: an open-source platform for biological-image
analysis." <u>Nat Methods</u> **9**(7): 676-682.
Schneider, C. A., W. S. Rasband and K. W. Eliceiri (2012). "NIH Image to ImageJ: 25 years of
image analysis." <u>Nat Methods</u> **9**(7): 671-675.
Shevchenko, A., H. Tomas, J. Havlis, J. V. Olsen and M. Mann (2006). "In-gel digestion for
mass spectrometric characterization of proteins and proteomes." <u>Nat Protoc</u> **1**(6): 2856-
2860.
Swedlow, J. R., J. W. Sedat and D. A. Agard (1996). Deconvolution in optical microscopy
<u>Deconvolution of Images and Spectra (2nd Ed)</u>. USA, Academic Press, Inc.**:** 284-309.
Tan, Y., X. Wang, H. Song, Y. Zhang, R. Zhang, S. Li, W. Jin, S. Chen, H. Fang, Z. Chen and K.
Wang (2021). "A PML/RARalpha direct target atlas redefines transcriptional deregulation in
acute promyelocytic leukemia." <u>Blood</u> **137**(11): 1503-1516.
Tatham, M. H., M. C. Geoffroy, L. Shen, A. Plechanovova, N. Hattersley, E. G. Jaffray, J. J.
Palvimo and R. T. Hay (2008). "RNF4 is a poly-SUMO-specific E3 ubiquitin ligase required for
arsenic-induced PML degradation." <u>Nat Cell Biol</u> **10**(5): 538-546.
Twomey, E. C., Z. Ji, T. E. Wales, N. O. Bodnar, S. B. Ficarro, J. A. Marto, J. R. Engen and T. A.
Rapoport (2019). "Substrate processing by the Cdc48 ATPase complex is initiated by
ubiquitin unfolding." <u>Science</u> **365**(6452).
Tyanova, S., T. Temu, P. Sinitcyn, A. Carlson, M. Y. Hein, T. Geiger, M. Mann and J. Cox
(2016). "The Perseus computational platform for comprehensive analysis of (prote)omics
data." <u>Nat Methods</u> **13**(9): 731-740.
Wallace, W., L. H. Schaefer and J. R. Swedlow (2001). "A workingperson's guide to
deconvolution in light microscopy." <u>Biotechniques</u> **31**(5): 1076-1078, 1080, 1082 passim.
Wang, Z. Y. and Z. Chen (2008). "Acute promyelocytic leukemia: from highly fatal to highly
curable." <u>Blood</u> **111**(5): 2505-2515.
Weis, K., S. Rambaud, C. Lavau, J. Jansen, T. Carvalho, M. Carmo-Fonseca, A. Lamond and A.
Dejean (1994). "Retinoic acid regulates aberrant nuclear localization of PML-RAR alpha in
acute promyelocytic leukemia cells." <u>Cell</u> **76**(2): 345-356.
Yau, R. G., K. Doerner, E. R. Castellanos, D. L. Haakonsen, A. Werner, N. Wang, X. W. Yang, N.
Martinez-Martin, M. L. Matsumoto, V. M. Dixit and M. Rape (2017). "Assembly and Function
of Heterotypic Ubiquitin Chains in Cell-Cycle and Protein Quality Control." <u>Cell</u> **171**(4): 918-
933 e920.
Zhu, H. H., Y. Z. Qin and X. J. Huang (2014). "Resistance to arsenic therapy in acute
promyelocytic leukemia." <u>N Engl J Med</u> **370**(19): 1864-1866.
1009

Figure 1. Jaffray et al 2023

Figure 2. Jaffray et al 2023

Figure 3. Jaffray et al 2023

Figure 4. Jaffray et al 2023

Figure 5. Jaffray et al 2023

Figure 6. Jaffray et al 2023

Figure 7. Jaffray et al 2023

Supplementary Figure 1. Jaffray et al 2023

Supplementary Figure 2. Jaffray et al 2023

Supplementary Figure 3. Jaffray et al 2023

YFP-PML	p97	Merge+DAPI	
			YFP-WT-PMLV 0h ATO
			YFP-WT-PMLV 6h ATO
			YFP-A216T-PMLV 0h ATO
			YFP-A216T-PMLV 6h ATO
			YFP-L217F-PMLV 0h ATO
			YFP-L217F-PMLV 6h ATO

Supplementary Figure 4. Jaffray et al 2023

Supplementary Figure 5. Jaffray et al 2023

Supplementary Figure 6. Jaffray et al 2023

Supplementary Figure 7. Jaffray et al 2023

Supplementary Figure 8. Jaffray et al 2023

A

PML-/- U2-OS
+YFP-PMLV-WT

PML-/- U2-OS
+YFP-PML-A216T

PML-/- U2-OS
+YFP-PML-L217F

1μM 2h   -As   +As   -As   +As   -As   +As

Anti-GFP nanobody purifications (24)

SDS-PAGE, In Gel digestion Trypsin (96) Trypsin+GluC (120)

LC-MS/MS, MaxQuant & Perseus data processing

B

YFP-WT-PML-V   YFP-A216T-PML-V   YFP-L217F-PML-V

-As   +As   -As   +As   -As   +As

Coomassie SDS-PAGE

kDa

1
2
3
4

C

-As   +As   -As   +As   -As   +As

250-
150-
100-

250-
150-
100-

250-
150-
100-

Anti-PML Western Blot

Supplementary Figure 9. Jaffray et al 2023

A

```
  1  MEPAPARSPRPQQDPARPQEPTMPPPETPSEGRQPSPSPSPTERAPASEE
 51  EFQFLRCQQCQAEAKCPKLLPCLHTLCSGCLEASGMQCPICQAPWPLGAD
                   65  68
101  TPALDNVFFESLQRRLSVYRQIVDAQAVCTRCKESADFWCFECEQLLCAK
                                  84                   101
151  CFEAHQWFLKHEARPLAELRNQSVREFLDGTRKTNNIFCSNPNHRTPTLT
                160                      183
201  SIYCRGCSKPLCCSCALLDSSHSELKCDISAEIQQRQEELDAMTQALQEQ
                209                 226
251  DSAFGAVHAQMHAAVGQLGRARAETEELIRERVRQVVAHVRAQERELLEA
301  VDARYQRDYEEMASRLGRLDAVLQRIRTGSALVQRMKCYASDQEVLDMHG
                                              337
351  FLRQALCRLRQEEPQSLQAAVRTDGFDEFKVRLQDLSSCITQGKDAAVSK
                                   380             394      400
401  KASPEAASTPRDPIDVDLPEEAERVKAQVQALGLAEAQPMAVVQSVPGAH
     401                      426
451  PVPVYAFSIKGPSYGEDVSNTTTAQKRKCSQTQCPRKVIKMESEEGKEAR
              460              476 478          487 490      497
501  LARSSPEQPRPSTSKAVSPPHLDGPPSPRSPVIGSEVFLPNSNHVASGAG
                   515
551  EAEERVVVISSSEDSDAENSVSGPEVQPRTPASPHFRSQGAQPQQVTLRL
601  ALRLGNFPVRH
```

RING (49-104)
B-BOX1 (127-165)
B-BOX2 (187-228)
COILED-COIL (229-339)
NLS (476-490)
SIM (556-562)
K Ubiquitinated
K SUMOylated
K SUMOylated and ubiquitinated
K Other lysine

B

C YFP

D PML

E SUMO1

F SUMO2+3

G Ubiquitin

H

Supplementary Figure 10. Jaffray et al 2023

A. Ubiquitin-Ubiquitin branched peptides

Untreated cells   **+Arsenic (1μM 2 hr)**

WT PML

Increase in SUMO modification
    Large - SUMO1 increase
    Modest - SUMO2/3 increase
RNF4 engaged
Ubiquitinated appropriately
(p97 recruited)
(PML degraded)

A216T PML

Small increase in SUMO modification
    Very small - SUMO1
    Small - SUMO2
RNF4 **not** engaged
Little increase in ubiquitination
(p97 **not** recruited)
(PML **not** degraded)

L217F PML

Increase in SUMO modification
    Modest - SUMO1 increase
    Modest - SUMO2/3 increase
RNF4 engaged
Ubiquitinated **in**appropriately
(p97 **not** recruited)
(PML **not** degraded)

Supplementary Figure 12. Jaffray et al 2023