Impaired polyamine metabolism causes behavioral and neuroanatomical defects in a novel mouse model of Snyder-Robinson Syndrome

Oluwaseun Akinyele[1], Anushe Munir[1,2], Marie Johnson[1], Megan E. Perez[1,2], Yuan Gao[3], Jackson R. Foley[4], Yijen Wu[5], Tracy Murray-Stewart[4], Robert Casero Jr.[4], Hulya Bayir[3,], Dwi U. Kemaladewi[1,2]

[1]Div. of Genetic and Genomic Medicine, Dept. of Pediatrics, University of Pittsburgh School of Medicine, Pittsburgh, USA.

[2]Dept. of Human Genetics, School of Public Health, University of Pittsburgh, Pittsburgh, USA.

[3]Children's Neuroscience Institute, Dept. of Pediatrics, University of Pittsburgh School of Medicine, Pittsburgh, USA.

[4]Sidney Kimmel Comprehensive Cancer Center, Johns Hopkins School of Medicine, Baltimore, Maryland, USA.

[5]Dept. of Developmental Biology, University of Pittsburgh School of Medicine, Pittsburgh, USA.

Corresponding author:

Dwi U. Kemaladewi, Ph.D.

Department of Pediatrics, University of Pittsburgh School of Medicine

UPMC Children's Hospital of Pittsburgh

4401 Penn Ave, Pittsburgh, PA 15224 USA

Dwi.kemaladewi@chp.edu

Abstract

Polyamines (putrescine, spermidine, and spermine) are essential for cellular growth and are subject to strict metabolic regulation. Mutations in the gene encoding spermine synthase (SMS) lead to the accumulation of spermidine and an X-linked recessive disorder known as Snyder-Robinson syndrome (SRS). There are no treatments available for this rare disease that manifests as mental retardation, thin habitus, and low muscle tone. The development of therapeutic interventions for SRS will require a suitable disease-specific animal model that recapitulates many of the abnormalities observed in these patients.

Here, we characterize the molecular, behavioral, and neuroanatomical features of a mouse model with a missense mutation in *Sms* that results in a glycine-to-serine substitution at position 56 (G56S) of the SMS protein. Mice harboring this mutation exhibited a complete loss of SMS protein and elevated spermidine/spermine ratios in skeletal muscles and the brain. In addition, the G56S mice demonstrated increased anxiety, impaired learning, and decreased explorative behavior in fear conditioning, Morris water maze, and open field tests, respectively. Furthermore, the mice undergo significant reductions in body weight over time, as well as abnormalities in brain structure and bone density. Transcriptomic analysis of the cerebral cortex revealed downregulation of genes associated with mitochondrial oxidative phosphorylation and synthesis of ribosomal proteins. Our findings also revealed impaired mitochondrial bioenergetics in fibroblasts isolated from the G56S mice, which provides a link between these processes and the pathogenesis of SRS. Collectively, our findings establish the first in-depth characterization of SRS-associated mechanisms in a preclinical animal model and identify cellular processes that can be targeted for future therapeutic development.

Keywords: Polyamines, Spermine synthase, spermine, neurological, disease, pathogenesis.

Introduction

Putrescine, spermidine, and spermine are ubiquitous polyamines that are essential for normal cellular functions. As a group, these polycationic molecules are responsible for maintaining chromatin structure and regulating gene expression and signaling pathways, as well as cell growth and death. Polyamines are also critical contributors to the many processes underlying immune cell activation, wound healing, and general tissue growth and development (1–6). Given their central role in cellular metabolism, the levels of intracellular polyamines are tightly regulated via *de novo* synthesis, interconversion, and transport into and out of cells (Figure 1).

Most of the intracellular polyamine pool arises from *de novo* synthesis. The diamine putrescine, which is the main precursor in this pathway, is synthesized from ornithine via the actions of the enzyme ornithine decarboxylase (ODC). Spermidine and spermine are higher-order polyamines derived from putrescine. Spermidine synthase (SPDS) catalyzes the reaction between putrescine and decarboxylated S-adenosylmethionine (dc-SAM) that generates spermidine. Spermine is then synthesized from spermidine and dc-SAM by spermine synthase (SMS). Spermidine and spermine can be converted back to their respective precursors via the actions of spermidine/spermine N^1-acetyltransferase (SSAT) and acetylpolyamine oxidase (APAO). Dysregulation or absence of any of these enzymes alters the homeostatic levels of cellular polyamines. Aberrancies in polyamine metabolism have been implicated in numerous diseases, including cancers, Alzheimer's disease, and Snyder-Robinson Syndrome (SRS) (7–10).

SRS (OMIM: 309583) is a rare X-linked intellectual disability syndrome associated with several known deleterious mutations in the *SMS* gene that lead to the loss of spermine synthase enzymatic activity and thus reductions in the intracellular spermine concentration (10). Loss of SMS activity also leads to an accumulation of spermidine and a high spermidine/spermine ratio, which is a biomarker used clinically to diagnose this disorder (10). SRS is characterized by intellectual disability, seizures, and general developmental delay (10–12). Individuals diagnosed with SRS exhibit thin body habitus and low muscle tone (hypotonia); these clinical findings may be apparent in infancy and worsen over time (10,13). Some SRS patients have difficulty walking and may never become ambulatory(11).

While a specific mutation in the *SMS* was first identified as the cause of SRS as early as 2003 (14), there are still no suitable animal models available to study disease pathophysiology and support the development of effective therapeutic interventions. Previous reports featured a *Drosophila* model in which an appropriate mutation was introduced into the *sms* gene (15) (16). While *Drosophila* models are readily available, many pathogenic factors that are critical to our understanding of vertebrate diseases are not conserved in this species and thus key aspects and

mechanisms may be overlooked (17). To address this concern, several groups used *Gy* (*Gyro*)

mice to explore the pathogenesis of SRS (17,18). The *Gy* mouse strain features a complete

deletion of the *Sms* gene and exhibits several of the abnormalities described in SRS patients,

including altered polyamine content, cognitive impairment, and bone abnormalities (19,20).

However, in addition to *Sms* deletion, the *Gy* mice also have deletion of the *Phex* gene. *Phex*

encodes the phosphate-regulating endopeptidase homolog, which is a protein involved in

phosphate transport that has been implicated in the pathogenesis of X-linked hypophosphatemia

(19,20). These attributes (i.e., the complete deletion of both *Sms* and *Phex*) complicate the

interpretation of many of the abnormalities observed in the *Gy* mice. Thus, this strain is not an

ideal model of SRS.

90
Figure 1. The polyamine metabolic pathway. Abbreviations: ODC, ornithine decarboxylase; SPDS, spermidine
synthase; SMS, spermine synthase; s-Adomet, s-adenosylmethionine; AMD-1, adenosylmethionine decarboxylase; dc-
SAM, decarboxylated s-adenosylmethionine; APAO, acetylpolyamine oxidase; SSAT, spermidine/spermine
acetyltransferase; PAs, polyamines.

Here, we explore disease pathogenesis in a recently generated mouse model of human SRS (21)

that carries a disease-specific missense mutation in *Sms* gene. This mouse strain harbors a

glycine to serine substitution in the amino acid sequence at position 56 of the SMS protein (G56S),

analogous to a mutation reported in individuals diagnosed with severe SRS (11). In this study. we

characterized the disease presentation in the G56S mice and explored potential mechanisms

underlying some of the observed abnormalities. Our findings revealed that G56S mice exhibit
many of the phenotypic (both behavioral and neuroanatomical) abnormalities reported in SRS
individuals (11,22,23). Furthermore, a transcriptomic analysis identified critical changes in gene
expression patterns that might contribute to these observed aberrancies.

Materials and methods

Mice: All animals used in this study were housed at the University of Pittsburgh Division of
Laboratory Animal Resources, Rangos Research Building, following the IACUC protocol number
2206137, which was approved by the University of Pittsburgh's Institutional Animal Care and Use
Committee. The colony of mutant mice was established by breeding female heterozygous *Sms*
mutation carriers (C57BL/6J-Sms[em2Lutzy]/J; Jackson Laboratory stock # 031170) and male WT
C57BL/6J mice (Jackson Laboratory stock # 000664). The male offspring of this cross that
harbored the X-linked G56S *Sms* mutation and WT littermate controls were used in the
experiments described in this study. To ensure that only male mice harboring the desired mutation
was used, pups were genotype at Transnetyx.com using the following probes: forward primer
ACCTGGCAGGACCATGGATATTTA, reverser primer GTGTTCACATCTAAAGCCCATGAGA,
reporter 1 AACAAGAATGGCAGGTAAG and reporter 2 ACGAACAAGAATTCCAGG.

Open field activity assay: The open field chamber is a hollow square field box equipped with
tracking software (ACTITRACK, Panlab/Harvard Apparatus, USA) connected to an infrared
tracking system that monitors animal movement. The walls of the box were opacified (covered
with aluminum foil) to prevent the environment from influencing the behavior of the mice that were
undergoing testing. The chamber was divided into two imaginary zones: an outer zone (45 x 45
cm) and an inner or center zone (18.5 cm x 18.5 cm, centered at 22.5 cm from the wall on each
side). Experiments were undertaken under constant room temperature ($22 - 25^{O}C$) and light
levels. Each mouse was released at the same location near the wall of the box and movement
was evaluated for 15 minutes using the infrared tracking system. The positions recorded for each
mouse were used to generate tracking plots and to determine the distance traveled, speed, and
time spent in each zone (i.e., within the entire apparatus and specifically in the center zone). The
total amount of time spent and the type of body motion (i.e., rearing, leaning, and vertical activity)
detected in the center zone were used as relative measurements of explorative behavior and
anxiety-related responses, respectively

Auditory-cued fear conditioning: The conditioning procedure was carried out using a
specifically designed chamber (model *H10-11M-TC-SF* Coulbourn Instruments, Whitehall, PA,
USA). The conditioning chamber (25 × 25 × 25 cm) had three grey methacrylate walls, a grid floor
connected to a shock scrambler to deliver foot shock as the unconditioned stimulus (US), and a

speaker mounted on the chamber ceiling to deliver audible tones as the conditioned stimulus (CS). The conditioning chamber was fitted with a high sensitivity camera system that monitored animal movement. The chamber was confined in a ventilated, soundproof enclosure (78 × 53 × 50 cm) on an anti-vibration table in a quiet room. The door to the room remained closed throughout the conditioning and testing periods.

On the first day (fear acquisition), the animals were habituated for 120 sec in the chamber before the delivery of CS-US pairs (i.e., a 75 dB tone [CS] for 20 sec followed by a 15-sec trace and then foot shocks [US] of 0.6 mA for 2 sec) with variable and pseudo-randomly distributed intervals between pairs of stimuli (90 – 203 sec). On the second day (fear retention), the session started with the mice placed in the same environment. During this phase, the mice were provided with no stimulation that might elicit contextual fear responses. Freezing responses in this otherwise familiar environment were monitored.

For the third session, the mice were placed in a different environmental setting (i.e., a chamber with a covered floor and white walls) to assess the retention of cued fear in a novel context. Baseline fear responses were monitored for 90 sec followed by the delivery of three CS (75 dB and 20 s) separated by variable inter-trial intervals (ITIs). The movement of the animal was sampled at a frequency of 50 Hz for quantitative analysis (Freezeframe, Coulbourn Instruments, USA). Freezing was analyzed during the delivery of the CS (20 sec periods) as well as during the 15 sec trace period that would ordinarily precede the US (not delivered) to monitor the associative fear response. The animals were gently handled before, during, and after the test to avoid introducing any additional potential stress before or during each test that could influenced the measured responses.

Morris Water Maze (MWM) Task: The MWM task was performed in a circular pool containing water using the procedure described by Tsien et al. (24) with slight modifications. The animals were trained to find an escape platform that was submerged in the water. The animals were not habituated in the pool before the training. The training protocol (hidden platform, used to evaluate spatial learning) included five sessions with 4 trials per session per day. Navigation was tracked by a video camera and the escape latency (i.e., the time required to locate the platform) was recorded. An animal that failed to locate the platform within 90 sec was guided to the platform. We then performed visible (to measure spatial memory) and probe (to measure non-spatial memory) tests on day six. In the visible test, colored tape was placed at the top of the platform. For the probe test, the platform was removed; the mice were allowed to swim in the pool for 60 s, and the time spent in each quadrant of the pool was recorded. The acquire data was analyzed using the ANY-maze software.

In vivo Magnetic Resonance Imaging (MRI) scans: All mice were subjected to *in vivo* brain
imaging while under isoflurane anesthesia. The mice were placed in a clear plexiglass anesthesia
induction box that permitted unimpeded visual monitoring. Induction was achieved by the
administration of 3% isoflurane in oxygen for several minutes. The depth of anesthesia was
monitored by toe reflex (extension of limbs, spine positioning) and respiration rate. Once
established, the appropriate level of anesthesia was maintained by continuous administration of
1-2 % isoflurane in oxygen via a nose cone. The mice were then transferred to the designated
animal bed for imaging. Respiration was monitored using a pneumatic sensor placed between
the animal bed and the mouse's abdomen. Rectal temperature was measured with a fiber optic
sensor and maintained with a feedback-controlled source of warm air (SA Instruments, Stony
Brook, NY, USA).

In vivo brain MRI was carried out on a Bruker BioSpec 70/30 USR spectrometer (Bruker BioSpin
MRI, Billerica, MA, USA) operating at 7-Tesla field strength and equipped with an actively shielded
gradient system and a quadrature radio-frequency volume coil with an inner diameter of 35 mm.
Multi-planar T_2-weighted anatomical images were acquired with a Rapid Imaging with Refocused
Echoes (RARE) pulse sequence with the following parameters: field of view (FOV) = 2 cm, matrix
= 256 X 256, slice thickness = 1 mm, in-plane resolution = 78 μm X 78 μm, echo time (TE) = 12
msec, RARE factor = 8, effective echo time (ETE) = 48 msec, repetition time (TR) = 1800 msec,
and flip angle = 180^0. Multi-planar diffusion MRI was performed using the following parameters:
field of view (FOV) = 2.0 cm, matrix = 128 X 128, slice thickness = 1.5 mm, in-plane resolution =
156 μm X 156 μm, TE = 16.31 msec, TR = 1500 msec, diffusion preparation with the spin echo
sequence, diffusion gradient duration = 4 msec, diffusion gradient separation = 8 msec, diffusion
direction = 30, number of A_0 images = 1, and b value = 1500 s/mm^2.

The MRI data were exported to a DICOM format and analyzed using the open source ITK-SNAP
(http://www.itksnap.org) brain segmentation software by 2 independent observers who were
blinded to the experimental conditions. The volumes of each region of interest (ROI), including
the amygdala, corpus callosum, thalamus, ventricles, hippocampus, and cortex were manually
drawn by blinded observers based on the information obtained from the Allen mouse brain atlas
(https://mouse.brain-map.org/static/atlas). To account for potential difference in the sizes of
brains in G56S and WT mice, volumes from each brain region were normalized to the total brain
volume of each mouse.

Diffusion MRI was analyzed by the open source DSI studio (http://dsi-studio.labsolver.org/) to
obtain fractional anisotropy (FA). ROIs contributing to quantitative and statistical analyses,
including the cortex, hippocampus, thalamus, corpus callosum, and ventricles with cerebrospinal
fluid (CSF) were manually segmented and defined by blinded independent observers.

***In vivo* micro-Computed Tomography (micro-CT) scans**: All mice undergoing *in vivo* micro-
CT imaging were maintained under general inhalation anesthesia with isoflurane as described for
MRI scans above. Once established, anesthesia was maintained with 1.5% isoflurane in oxygen
administered using a nose cone and the mouse was transferred to the designated animal bed for
imaging. Respiration was monitored as described above. Respiration gating was performed using
BioVet system that was triggered by maximal inhalation with a 500 ms trigger delay.

Respiration-gated *in vivo* micro-CT imaging was performed with Siemens Inveon Multimodality
micro-CT-SPECT-PET system with the following parameters: full rotation, 360^0 projections; settle
time 1000 msec; 4X4 binning; effective pixel size of 76.75 µm; trans axial field of view (FOV) 78.6
mm with 4096 pixels; axial FOV 76.1 mm with 3968 pixels 80 kV of voltage; current of 500 µA;
exposure time of 410 ms. The three-dimensional (3D) micro-CT images were reconstructed using
the Feldkamp algorithm and were calibrated in Hounsfield Units (HU). Double distilled water was
set at a readout of 0 and air at 1000 HU.

The 3D micro-CT image stacks were analyzed using the Inveon Research Workplace (IRW). The
ROI analysis function was used with a thresholding tool that created several ROIs with different
Hounsfield Unit (HU). A cylindrical 3D ROI was drawn around the body that encompassed the
entire body. All external air around the mouse was excluded from the ROI and a custom threshold
was set between 400 – 5700 HU to capture the bones. The mean HU values obtained from each
ROI were used to quantify bone density.

Body composition measurements: Body composition (percentage lean and fat weight) of the
mice was measured by quantitative MRI (EchoMRI, Echo Medical Systems, Houston, TX).
Animals were placed in thin-walled plastic cylinders with a plastic restraining inserts. Each animal
was briefly subjected to a low-intensity electromagnetic field that measured total body
composition. Percentages of fat and lean body weights were determined based on total body
weight.

Primary cell isolation: Primary fibroblast cultures from the ears of G56S and WT mice were
established using the protocol described by Khan and Gasser(25).

RNA isolation and quantitative polymerase chain reaction (qPCR): Total RNA was isolated
from mouse tissues using the Nucleospin RNA Plus kit (Macherey-Nagel, cat# 740984.50),
following the manufacturer's instructions. cDNA synthesis was performed using the iScript
Reverse Transcriptase Supermix kit (BioRad, Cat# 1708841) according to the manufacturer's
instructions. qPCR was performed using 2X SYBR Green Fast qPCR Mix kit (ABclonal, cat#
RM21203) in a C1000 Touch Thermal Cycler (BioRad, USA). The primer sequences used to

amplify target sequences of interest are listed in Supplementary **Table 1**. Expression of endogenous *Gapdh* was used as an internal control to measure the relative expression of genes of interest. The $2^{\Delta\Delta Ct}$ was used to assess relative fold change in gene expression in tissue samples from WT and G56S mice samples. Values are presented as the percentage change in fold expression.

RNA-sequencing (RNAseq) and pathway enrichment analysis: After completing the extraction procedure described above, RNA samples were submitted to the Health Sciences Genomic Core at the UPMC Children's Hospital of Pittsburgh. RNA quality was determined using the Agilent Bioanalyzer 2100 (Agilent Technologies). cDNA libraries were prepared using a 3′-Tag-RNA-Seq library kit (Illumina). Sequencing was performed using one lane of a Hi-Seq 4000 platform with pair-end 40 bp reads. Analysis of sequence reads, including quality control, mapping, and generation of tables of differentially expressed genes (DEGs), heatmaps, and volcano plots) were performed using the Qiagen licensed CLC Genomic Workbench software version 22.0.1. Pathway enrichment analysis of the DEGs was performed using the Qiagen-licensed Ingenuity Pathway Analysis (IPA) software. The gene expression profile identified by RNA-seq was validated by qPCR as described above.

***In vivo* analysis of mitochondria respiration:** Oxygen consumption rates (OCRs) were determined with a Seahorse XFe96 Extracellular Flux Bioanalyzer (Agilent Technologies, Santa Clara, California, USA). Primary fibroblasts plated in 96-well assay plate at a density of 40,000 cells/well were cultured overnight and then equilibrated with Seahorse XF base medium (Agilent Technologies) supplemented with glucose, sodium pyruvate, and L-glutamine at 37°C in a CO_2-free incubator for 1 hour prior to the assay. Mitochondrial function was assessed by sequential addition of 1.5 μM oligomycin, 1 μM FCCP (carbonyl cyanide-4-[trifluoromethoxy] phenylhydrazone), and 0.5 μM rotenone/antimycin A by the Seahorse Bioanalyzer. Data was normalized by the total protein content of the cells.

Protein isolation, quantification, and western blotting. Total proteins were extracted from tissues isolated from G56S and WT mice tissues using RIPA homogenizing buffer (150 μL of 50 mM Tris HCl pH 7.4, 150 nM NaCl, 1 mM EDTA) followed by homogenization using a bullet blender. After homogenization, 150 μL of RIPA double-detergent buffer (2% deoxycholate, 2% NP-40, 2% Triton X-100 in RIPA homogenizing buffer) supplemented with protease inhibitor cocktail (Roche, cat# A32953) was added to the tissue homogenate followed by incubation on a shaker for 1 h at 4^0C. The tissue homogenate was then centrifuged at 11,000 *g* for 10 min at 4^0C. The resulting supernatant was used to quantify total protein using the Pierce BCA protein assay kit (Thermo Scientific, cat# 23225) according to the manufacturer's protocol. Twenty microgram

of total protein was fractionated on 4 – 12% gradient gel (Thermo Scientific, cat# NP0336BOX).
After proteins had separated on the gel, they were transferred by electroblotting onto a
Polyvinylidene fluoride (PVDF) membrane and blocked with 5% non-fat milk in TBS-Tween-20.
The membrane was then incubated overnight with rabbit anti-spermine synthase (Abcam, cat#
ab156879 [EPR9252B]) or rabbit anti-vinculin (Abcam, cat# ab129002 [EPR8185]). After
incubation with the primary antibody, the membranes were washed and then incubated with the
secondary antibody (Goat Anti-Rabbit IgG – HRP conjugate, Bio-Rad cat# 1706515) for one hour
at room temperature. Specific protein bands were detected using SuperSignal^{TM} West Femto
Maximum Sensitivity Substrate (Thermo Scientific, cat# 34095). Bands corresponding to
immunoreactive SMS and Vinculin were identified and quantified using the ChemiDoc Imaging
System (BioRad).

Polyamine measurement: The polyamine content in isolated tissues was measured by the
precolumn dansylation, high-performance liquid chromatography method described by Kabra et
al. using 1,7-diaminoheptane as the internal standard (26).

Statistical analysis: Statistical analysis was performed using Graphpad Prism software vs 9.0.
Each variable was statistically compared between the WT and G56S mice using unpaired
students' t-test unless otherwise stated. A *p* value less than 0.05 was considered statistically
significant.

288
RESULTS
Sms **mutation altered the tissue polyamine content in mouse tissues**
To determine whether the substitution of two nucleotides (GGC to TCC) in exon two of the *Sms*
gene altered its expression profile and tissue polyamine levels, we measured *Sms* mRNA and
protein in tissues isolated from bothG56S and WT control mice. While there were no significant
changes in the level of *Sms* mRNA (Figure 2a), we observed a near-complete loss of SMS protein
in both the brain and skeletal muscles of G56S mice (Figure 2b). Similarly, an LC-MS analysis of
tissue polyamines revealed elevated levels of both putrescine and spermidine and a significant
decrease in spermine in brain tissue from G56S mice. The spermidine to spermine ratio was 4-
times higher in brain tissues from G56S mice compared to WT controls (Figure 2c). Our findings
also revealed a significant increase in spermidine but no changes in the putrescine levels in
skeletal muscles of the G56S strain (Figure 2d and inset). The spermine concentration was below
the limits of detection in the mutant strain. Figure 2e presents the two-dimensional (2D) crystal
structure of the dimerized and fully functional SMS protein including the C-terminal (catalytic) and
N-terminal (dimerization) domains; the amino acid within the latter domain at position 56 (WT,

glycine) is highlighted in Figure 2f. Figure 2g presents the mutant SMS protein with a serine at position 56; the extended side chain characteristic of this amino acid may interfere with monomer dimerization and may lead to the loss of SMS protein despite normal transcript level.

Figure 2. Tissue polyamine content and a potential mechanism underlying SMS protein loss. (a) Mice brain *Sms* gene transcript level as quantified by RT-qPCR. The level of gene transcript in the mutant was compared to the WT mice set at 100% reference value (b) Expression of SMS protein in isolated brain and skeletal muscles (triceps and gastrocnemius) from WT and G56S mice. (c) LC-MS determination of brain polyamine content and SPD/SPM ratios in WT and G56S mice quantified by LC-MS. (d) Skeletal muscle polyamine content in both G56S and WT mice. Note: spermine levels were below the limit of detection in the G56S skeletal muscle. (e) 2D-crystal crystal structure of the SMS protein with glycine at position 56 in the N-terminal region (circled). The 2D-crystal structure SMS was modeled from protein data bank ID: 3C6M. (f) The atomic structure of glycine at position 56 in the N-terminal region of SMS protein. (g) Serine in place of glycine at position 56 of SMS protein; the extended serine sidechain is highlighted in yellow. Panels a-d, values shown are mean ± standard error of the mean (S.E.M.); n = 3 – 5 mice per group; *$p < 0.05$, **$p < 0.01$, ns = not significant.

Biometric parameters are significantly altered in G56S mice

Patients diagnosed with SRS typically exhibit an asthenic physique with thin body build and short stature. Many of these patients also exhibit significant bone abnormalities and low muscle tone

(10). To understand the mechanisms contributing to these physical aberrations, we determined

the impact of altered polyamine content on the growth and development of G56S mice. Routine

measurements revealed that the G56S mice have significantly lower body weight than age-

matched WT counterparts and that they gained little to no weight throughout the duration of the

study period (Figure 3a). Similarly, the G56S mice exhibit significantly reduced body lengths

(Figure 3b). Collectively, these results suggest that G56S mice may experience failure to thrive

compared to their WT littermates. Also, an analysis of the body composition of mutant mice

revealed higher percentage of lean weight and a lower percentage of fat weight compared to their

WT counterparts (Figure 3c). This finding may reflect the diminished body build typically observed

in patients diagnosed with SRS. Similarly, quantification of bone density by micro-CT revealed

that the G56S mice exhibited decreased bone density compared to their WT littermates (Figure

3d). Taken together, these data suggest that changes in tissue polyamine content observed in

the G56S mice may have an impact on tissue development and result in general failure to thrive.

337
338 **Figure 3. Biometric analysis of male G56S and WT mice.** (a) Total body weight was measured in 3-week intervals.
339 (b) Body length of 24-week-old mice. (c) Body composition of 15-week-old mice (% lean and % fat weight) determined
340 by by Echo-MRI scan. (d) Bone mineral density (BMD) of 20-week-old mice measured by micro-CT scan. Values shown
341 are mean ± S.E.M., n = 7 mice per group,** *p < 0.01*, *** *p < 0.001*, **** *p < 0.0001*.

G56S mice exhibit signs of cognitive impairment

Mild to severe mental retardation is one the major clinical consequences of SRS. Two patients diagnosed with the G56S mutation were reported to have severe cognitive disabilities (11). To explore the impact of the G56S mutation on cognition, we evaluated the performance of G56S and WT mice using the MWM test. For this test, the mice were placed in a pool of water and were provided with five days of training in which they learned to locate an escape platform that was submerged and thus hidden under the water (Figure 4a, panel 1). The time taken to find the platform on each day was recorded.

While our findings revealed no significant differences in the time required to find the escape platform, there was a trend suggesting that the WT were somewhat more effective than the G56S mice at performing this task (Figure 4b). Our findings from the probe test in which the platform was completely removed from the MWM (day 6; see Figure 4a, panel 2) revealed no significant differences in the time spent in the escape quadrant (Figure 4c). However, the G56S mice required significantly more time to locate a visible platform (see Figure 4a, panel 3) compared to the WT mice (Figure 4d). This result suggests that the WT mice learned and retained the information needed to complete this task more effectively than the G56S mice. Overall, the results of the MWM test suggest that the G56S mice exhibit relatively mild learning impairments compared to their WT littermates; however, this trait may become more apparent and severe as the mice age.

Figure 4. Performance of G56S and WT mice in a Morris Water Maze (MWM) test. (a) Description of the three components of the MWM test. (b) Time required to locate a hidden escape platform on each day of a five-day training period. (c) Time spent in the escape quadrant (probe test without the platform) given a limit of 60 sec on day 6. (d) Time required to locate a visible escape platform on day 6. Data are mean ± s.e.m, n=7 mice per group; two-way ANOVA for repeated measures (a) and unpaired *t*-tests for (b) and (c), not significant (ns); *p < 0.05.

G56S mice exhibit diminished explorative behavior in an open field test

Anxiety-related responses are among the major symptoms of many neurological and neurodevelopmental disorders, including SRS. Tests designed to evaluate these responses have been used to characterize disease-specific mouse strains. Spontaneous locomotor activity was used to evaluate anxiety-related responses of both G56S and WT mice using a test that provided

them with the opportunity to explore an open field arena for 10 min. For this test, the open field

arena was divided into an outer or peripheral zone and an inner or center zone (Figure 5a). We

assessed the time spent in the center zone, the resting time, and the number of times each mouse

entered the center zone. The test was performed once every three weeks to characterize changes

in behavior with advancing age.

Among our results, we found that the G56S mice were less active compared to their WT

counterparts (Figure 5a). While no statistically significant differences were detected between the

two strains when the mice were less than 18 weeks old, the G56S mice became significantly less

active with increasing age compared to their WT littermates (Figure 5b). Similarly, G56S mice

were much less likely to enter the center zone of the open field arena than their WT counterparts

beginning at 18 weeks of age (Figure 5c). These results suggest the possibility that the G56S

mice experience slow but steady disease progression.

Older G56S mice also spent significantly more time resting in the outer zone (Figure 5d) and less

time resting in the inner zone of the open field compared to the WT control (Figure 5e).

Collectively, these findings suggest that G56S mice exhibit less explorative behavior than their

WT counterparts and that these responses may represent higher anxiety or fear that increases

as the mice age.

Figure 5. Anxiety-related response monitoring in an open field test. (a) Movement pattern of the animals for 20
min in the open field test at 24 weeks of the age. (b) Total activity of the animals in the open field test measured as the
number of total numbers of movement, rearing, and other bodily activities. (c) Number of entries to the center zone of
the open field test. (d and e) Resting time of the animals in the outer and center zones of the open field respectively.
Data are mean ± s.e.m., *$p < 0.05$; **$p < 0.01$; ns, not significant.

G56S mice exhibit increased fear-related responses

Fear-related responses exhibited by both WT and G56S mice were assessed via measurements of stress-induced freezing. This response is an innate anti-predator fear-related behavior that is characterized by a complete tonic immobilization while sparing respiration. This test uses auditory-cued fear conditioning, which requires the mouse to associate an aversive outcome with an otherwise unrelated cued stimulation.

In this experiment, anxiety or fear responses are expressed as the percentage of time spent in a freezing position after sound stimulation (an auditory cue) followed by foot shock (aversive condition).

On the first day of the test, the mice were trained to associate the sound (conditioned stimulus, CS) with the foot shock (unconditioned stimulus, US). The percentage of time spent in stress-induced freezing was recorded during the CS and after delivery of the US. We observed a consistent increase in the freezing responses following the CS in both groups of mice; this response reached a plateau after the delivery of the fourth sound stimulation and decreased after the fifth (Figure 6a). These results suggest similar rates of fear acquisition by both groups of mice. However, we observed significant differences in the freezing responses displayed during the inter-trial intervals (i.e., the period between the last foot shock and the next sound stimulation). The G56S mice exhibited significantly longer, and more frequent freezing responses compared to the WT mice (Figure 6b). Similarly, WT mice recovered ambulation more rapidly than the G56S mice following administration of the CS and US. This result indicates that the G56S mice exhibit more profound fear responses following stimulations than their WT counterparts.

We then compared the freezing responses of G56S and WT mice in a contextual test. This test was performed 24 hours after the CS-US training and involved no stimulation; the mice were placed in the same experimental chamber and their freezing responses in this environment were measured. While no statistically significant differences were observed, we detected a pattern that suggested that the G56S might exhibit increased freezing responses compared to the WT mice (Figure 6c). This trend suggests that the innate fear responses exhibited by the G56S mice may be more profound when compared to those of their WT littermates.

In the cued test, we evaluated the freezing responses of both WT and G56S mice to the CS only in a different environmental setting. After measuring baseline freezing responses during an initial 60 sec habituation period, the mice were subjected to 20 sec of auditory stimulation (CS). Their fear response during the CS and various intertrial intervals were then monitored. While the G56S mice exhibited comparatively higher baseline freezing responses compared to their WT counterparts in the new environmental setting, the differences did not achieve statistical

significance. By contrast, the G56S mice exhibited significantly higher fear responses compared to the WT controls both during the CS as well as the ITIs (Figure 6d). Of note, the percentage freezing during the ITIs exhibited by the G56S mice was not only elevated, it remained at a level that was similar to that observed during the CS. Taken together, these data suggest that the G56S mice exhibit higher anxiety-related fear responses than their WT counterparts. This finding may represent a specific neurological dysfunction similar to those observed in patients diagnosed with SRS.

Figure 6. Auditory-cued fear responses following a conditioned stimulus (CS). (a) Fear learning and fear response levels after each conditioned stimulus (CS; inter-trial interval, ITI) were assessed in 4–5-month-old G56S and WT mice (n=7 per group). The fear response was expressed as the percentage of time spent in a stereotypical freezing state during the presentation of the CS repeated five times during fear acquisition (CS 1-5; a 75 dB tone lasting 20 sec followed by a foot shock). This was repeated four times with staggered inter-trial intervals. The inset documents the experimental sequences on day 1, which include habituation (hab), tone presentation (CS, loud-speaker symbol), shock delivery (lightning symbol), and ITIs of varying durations. (b) WT and G56S mice demonstrated significantly different responses during fear acquisition (genotype effects and genotype trial interactions were evaluated by two-way ANOVA for repeated measures). (c) Fear response in contextual setting, without sound and shock stimulation, performed 24 hours after fear acquisition. (d) Cue-fear response (with a changed environment) induced by CS alone (tone: 20 sec, repeated three times with no foot shock) with variable ITIs. The Baseline value represents the percentage spent in the freezing state during habituation before the CS (sound); the CS response represents the average of three CS trials; the bars labeled ITI represent the average of the three it is for each mouse. The inset documents the experimental sequences used on day 2, including habituation (hab), tone presentation (CS, loud-speaker symbol), and ITIs of varying durations. Results obtained from WT and G56S mice were compared using unpaired t-tests. Values shown are mean ± S.E.M. *$p < 0.05$; ** $p < 0.01$; ***$p < 0.001$; ****$p < 0.0001$.

G56S mice exhibit neuroanatomical changes

We explored brain anatomical structures to determine whether the G56S mutation and resulting alterations in polyamine metabolism were associated with major structural changes. We also determined how altered brain structures might correlate with the behavioral defects described in previous tests. For these experiments, brain volumes were assessed using T2-weighted anatomical scans and diffusion tensor imaging (DTI) (Figure 7a). Other DTI parameters collected included fractional anisotropy (fa), mean diffusivity (md), axial diffusivity (ad), and radial diffusivity (rd). The results of whole-brain imaging showed that the G56S mice have significantly smaller brain volumes than WT mice (Figure 7b). Similarly, analysis of several specific brain regions that were selected based on reports describing SRS patients (22,23) and our behavioral data revealed that the volumes of the amygdala (involved in fear learning and emotional responses), the hippocampus (involved in cognitive functions), and the corpus callosum were all significantly lower in the G56S mouse strain (Figure 7c). Also, DTI analysis of the various brain regions revealed that fa, which is a measure of the microstructural integrity of the white matter of the amygdala and the corpus callosum, was significantly lower in the G56S mice compared to the WT controls. The other regions, including the hippocampus and the cortex, exhibited decreasing trends, although did not achieve statistically significant (Figure 7d).

Figure 7. *In vivo* MRI structural analyses. (a) Representative MRI images of coronal sections of brains from WT and G56S mice. Annotations of different regions of the brain were based on the Allen Mouse Brain Atlas. (b and c) MRI volumetric analyses of the total brain volume and volumes of annotated regions highlighted in panel (a). Quantification of the volume of each region was normalized to the total brain volume for each sample. Volumes of each brain region were quantified using ITK snap software to assess RER8 MRI scan images. (d) Fractional anisotropy (fa) was quantified using DSI studio software. Comparisons of single variables between WT and G56S mice were performed using unpaired *t*-tests. Data shown are mean ± S.E.M. *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$; ns, not significant. **Note**: The amygdala region was not visible in the image shown in panel (a) above.

Mutation in *Sms* gene alters transcriptomic profile of the G56S mouse brain cortex

To elucidate the molecular mechanisms underlying some of the observed phenotypic and behavioral differences, we isolated RNA and performed a transcriptomic analysis of the brain cortex tissue of WT and G56S mice. We focused on the cortex because of its role in directing higher complex tasks, including learning, memory, and consciousness. Furthermore, results from previous studies suggest that spermine may have protective role specifically within the cerebral cortex (22,27). The results of our transcriptomic analysis of brain cortex tissue from WT and G56S mice revealed more than 1,000 differentially expressed genes (DEGs) (Figure 8a and Supplementary **Table 2**). A heatmap revealed differential expression of genes involved in several

key cellular and metabolic processes (Figure 8b). Importantly we found differential expression of

genes that contribute to mitochondria function and ribosomal protein synthesis. We performed

gene enrichment pathway analysis to identify possible metabolic pathways that might be altered

in the G56S mice. Our results revealed that downregulation of genes involved in mitochondrial

oxidative phosphorylation and ribosomal protein synthesis (i.e., eukaryotic initiation factor 2 [eIF2]

signaling) in cortical tissue from the G56S mice (Figure 8c). In addition, pathways involved in

Huntington's disease, sirtuin, and synaptogenesis signaling pathways were all upregulated in the

G56S mice brain cortex compared to the WT (Figure 8c). The expression pattern of genes involve

in these metabolic processes were further confirmed in the G56S mice relative to the WT by a

Volcano plot (Figure 8d). Quantitative polymerase chain reaction (qPCR) performed to validate

the differential expression of genes associated with mitochondrial oxidative phosphorylation

confirmed decreased expression of *ATP5e, Uqcr10, Cox6B1, Cox4i1, Cox7b, Ndufa4, and*

Ndufa7 as well as decreased expression of *Rpl17* and *Rsp14* (both implicated in ribosome protein

synthesis via eIF2 signaling) in the brain cortex of G56S compared to WT mice (Figure 8e).

qPCR analysis also confirmed the observed upregulation of Huntington-associated protein 1,

Hap1, and the disease-related ionotropic NMDA receptor subunit 2b, *Grin2b* (Figure 8e).

Collectively, these data suggest that the G56S mutation in the *Sms* gene and alterations in tissue

polyamine levels resulted in specific changes in the expression profile of genes involved in central

cellular and metabolic processes of the brain. These observations may explain one or more of

the phenotypic abnormalities observed in G56S mice.

Figure 8. **Transcriptomic analysis of isolated brain cortical tissues**. (a) Differentially expressed genes (DEGs)
were identified using Qiagen CLC Genomic Workshop software. (b) Heatmap of selected genes that exhibit statistically
significant differences in expression (*p < 0.05*) and absolute values of log₂-fold change (LFC) greater than or equal to
1. (c) Gene enrichment analysis of up- and downregulated transcripts with *p < 0.05* and absolute LFC ≥ 0.5 in the brain
tissue from G56S compared to WT mice. Gene enrichment analysis was done using the Qiagen Ingenuity Pathway
Analysis (IPA) software. (d) A volcano plot showing the relative expression of selected genes involved in oxidative
phosphorylation and other pathways based on the results of the enrichment analysis. (e) qPCR validation of RNA-seq
results for selected genes implicated in oxidative phosphorylation, eukaryotic initiation factor 2 (eIF2) signaling, and
Huntington's disease. Data shown are mean ± S.E.M, (n=3 for both WT and G56S mice). *$p < 0.05$; **$p < 0.01$, ns, not
significant.

G56S mice exhibit impaired mitochondrial respiration

To confirm the downregulation of mitochondrial oxidative phosphorylation predicted by gene

expression analysis, we isolated primary fibroblasts from both the WT and G56S mice and

evaluated mitochondrial respiration using the XFe96 Seahorse bioanalyzer. Our results revealed

that both basal and oligomycin-sensitive respiration rates were significantly diminished in

fibroblasts isolated from G56S mice (Figures 9a and 9b). Similarly, maximum respiration and rates

of ATP synthesis were also significantly reduced in fibroblasts from the G56S mice compared to

the WT (Figures 9c and 9d). Taken together, these data suggest that SMS deficiency and

impaired polyamine metabolism in G56S fibroblasts will lead to impaired mitochondrial
bioenergetics and functions.

536
Figure 9. **Mitochondrial respiration in isolated fibroblasts**. (a) Respiratory profiles of primary fibroblasts isolated
from WT (blue line) and G56S (red). Oligomycin (ATP synthase inhibitor), FCCP (H$^+$ ionophore), and
rotenone/antimycin (mitochondria complex I/III inhibitors) were added at the times indicated. (b) Basal respiration, (c)
maximal respiration, and (d) ATP production in fibroblasts from WT and G56S mice were assessed using a Seahorse
XFe96 analyzer. Data shown are mean ± s.e.m. of average n=8 technical replicates of two independent experiments.
**p < 0.01; ***p < 0.001.

Discussion

One of the main challenges of preclinical research intended to elucidate the mechanisms

underlying rare brain-related disorders and other diseases associated with cognitive impairment

is the identification of robust and translationally-relevant behavioral phenotypes using animal

models. While the G56S mouse features an *Sms* gene-specific missense mutation that replicates

a genetic lesion identified in patients with severe form of SRS, however it is not known whether

this mouse represents a useful tool for therapeutic development. Therefore, the goal of this study

is to provide a detailed behavioral and neuroanatomical assessment of these mice. We also

studied the transcriptomic landscape of the brain cortex to identify gene expression patterns that

may contribute to the abnormalities observed in the mice and conceivable in SRS patients.

SRS is a disease characterized by abnormal somatic features and a general failure to thrive.
Patients are described as having asthenic body builds, mild short stature, and diminished body
mass. These patients also exhibit abnormal bone structure and sustain frequent fractures (10). In
the current study, we report that the G56S mice exhibit lower body weights as well as reduced
overall length, bone mineral density, and body fat composition compared to their WT counterparts.
These findings suggest that the G56S mouse model reproduces many of the abnormal clinical
features described for SRS. However, the impact of the resulting disruptions in polyamine
metabolism and their contributions to disease-specific symptomatology remain unknown. A
previous study focused on bone marrow-derived multipotent stromal cells (MSCs) revealed that
the mRNA silencing of SMS resulted in impaired cell proliferation and a reduced capacity for
osteogenesis (28). These results suggest a possible link between the *Sms* mutation, dysregulated
polyamine metabolism, and thus decreased bone mineral density in both patients and the G56S
mice. While most SRS patients, including those with the G56S mutation (11) are eventually
diagnosed with kyphoscoliosis, no scoliosis was detected in micro-CT scans of these mice.
However, we cannot rule out the possibility that abnormal spines may develop in older mice.

The decreased body fat composition might be attributed to the increase in tissue spermidine, as
this polyamine has been implicated in promoting lipolysis (29). However, it is also possible that
the absence of SMS can result in impaired mitochondrial functions (15,28). In this case, the mice
will depend more heavily on glycolysis as a means of energy generation; this will result in the
utilization of greater amounts of food with less available to be converted to and stored as body
fat. The decreased body weight seen in these mice supports the general idea that disturbances
in polyamine homeostasis impair cell growth and tissue development (4) and lead to general
growth failure.

Similar to what has been reported in many SRS patients, the G56S mouse displays signs of
cognitive impairment and exhibits significant reductions in exploratory behavior when evaluated
in an open field test. Collectively, these findings suggest that these mice exhibit amplified anxiety-
related behaviors. This hypothesis was further confirmed using the fear conditioning test. Mice
displayed significantly heightened fear responses when presented with an auditory conditioned
stimulation (CS) followed by unconditioned stimulus (US; foot shock). The increase in fear
responses in both the contextual and cued tests further confirm the observed heightened anxiety-
related behavior in the G56S mouse strain. Overall, these data provide some evidence suggesting
the existence of neurological abnormalities in these mice. Since polyamines are important
contributors to the development of the nervous system (30), several regions of the brain might be
contributing to these behavioral defects. Both the amygdala and the hippocampus in the G56S

mice, which are brain regions involved in fear-associated memory and learning, respectively, were decreased in volume compared to the WT, similar to that reported in SRS patients (22). Thus, these results suggest that impaired polyamine metabolism and toxic spermidine accumulation may lead to atrophy and neuronal loss in these regions (22) as well as the defects in behavioral and learning outcomes observed in these mice. The possibility that disrupted polyamine metabolism might lead to brain atrophy was further confirmed by the decrease in the fa value found in G56S mice. Although we did not measure the polyamine content in specific regions of the brain, we believe that the total brain polyamine content most likely reflects the overall content in the different regions, given that G56S mice lack SMS in all tissues and that this mutation is not tissue or brain-region specific.

Another potential mechanism that might explain the behavioral defects observed in G56S mice is spermidine-mediated disruption of receptor signaling. In an earlier study, Rubin *et al.* (31) reported that intra-amygdala administration of spermidine in an experimental rat model resulted in a dose-dependent increase in freezing responses. These results suggested that the accumulation of spermidine in the brains of G56S mice might contribute to the observed increase in anxiety-related behaviors. The mechanisms underlying spermidine-mediated increases in fear responses have not yet been clarified.

Spermidine may regulate the function of the amygdala via interactions with and modulation of the ion channel receptor for N-methyl-D-aspartate (NMDA); earlier report detail polyamine-mediated negative regulation of this receptor (32). Administration of arcaine, a putative competitive antagonist at the polyamine binding site of the NMDA receptor, resulted in a decrease in spermidine-induced fear responses in rats (31). Collectively, these results suggest that spermidine levels may have an impact on amygdala function and that accumulation of spermidine may induce a fear response as well as other behavioral abnormalities seen in the G56S mice. It is important to note that anxiety has been identified as one of the main symptoms of numerous neurological disorders (33). Although there are no clinical reports that document this specific behavior in SRS patients, family members have confirmed anxiety and the prevalence of fear-related behaviors in some patients (Personal communication, Snyder-Robinson Foundation conference 2022)

In addition to the neuroanatomic defects, other potential mechanisms contributing to the phenotypic abnormalities observed in the G56S mice were revealed by the transcriptomic analysis. These include impaired mitochondrial function, alterations in ribosomal protein synthesis signaling pathways, and upregulation of genes implicated in the pathogenesis of Huntington's disease. The mitochondria are important energy-generating cellular organelles; mitochondrial

dysfunction has been implicated in a variety of different neurological or neurodegenerative
diseases (34). Results from several previous studies have suggested that impaired mitochondrial
function might contribute to the pathogenesis of SRS (15,28) via one of several potential
mechanisms:

(i)   Increased spermidine levels that accumulate in cells that lack SMS may promote the
synthesis and release of reactive oxygen species (ROS) secondary to increased
catabolism. Elevated ROS results in mitochondrial oxidative stress and impaired
mitochondrial function (15).

(ii)   Polyamines play essential roles in modulating gene expression. Earlier reports
suggest that spermine modulates mammalian mitochondrial translation initiation
processes (35,36). Thus, the lack of SMS or spermine could inhibit synthesis of
mitochondrial proteins and result in impaired mitochondrial functions.

(iii)   Normal mitochondrial metabolism can result in the accumulation of potentially
damaging levels of by-products including ROS and Ca^{2+} (37). As a polycationic
molecule, spermine can scavenge mitochondrial ROS (38,39), and reduce the levels
of mitochondrial permeability transition pore (mPTP) generated in response to Ca^{2+}
accumulation (40). Thus, the lack of SMS or an observed decrease in cellular spermine
content may result in mitochondrial damage.

(iv)   Another potential mechanism of mitochondrial impairment in SRS may relate to the
decreased expression of nuclear genes encoding mitochondrial proteins reported in
this study. Although we do not yet understand how SMS mutations and/or altered
spermine concentrations result in the changes in gene expression observed, either
factor may be involved in direct or indirect interactions with critical transcription factors.
It will be important to identify relevant transcription factors as this may improve our
understanding of how spermine and/or SMS modulate mitochondrial function.

In conclusion, efforts to develop effective therapies for SRS will require a better understanding of
the disease pathophysiology as well as suitable disease-specific animal models that recapitulate
many of the critical abnormalities diagnosed in these patients. The findings presented in this study
suggest that the G56S mouse is a good model that can be used to study SRS pathogenesis and
may be an important tool for therapeutic development. This study also provides parameters that
may be used to assess the effectiveness of therapy for SRS in a murine model. Finally, the data
shown here offers some insights that could be used to improve the current clinical management
of SRS.

Acknowledgements:

We thank the Snyder-Robinson Foundation, C. Lutz, and A. Zuberi for generating and providing the G56S mouse model; A. Schmidt, E. Goetzman, and K. Schwab for the advice and assistance with animal imaging; W. MacDonald, R. Elbakri, and A. Chattopadhyay for the help on transcriptomic and pathway analyses; C. van 't Land, A. Karunanidhi, and G. Vockley for SeaHorse experiment; O. Tomisin for help with the 2D protein crystal structure. Members of the Kemaladewi laboratory are acknowledged for technical support and input in this study. This work was supported by the RK Mellon Institute for Pediatric Research Postdoctoral Fellowship (to O.A.), Children's Trust of UPMC Children's Hospital of Pittsburgh Foundation, Dept. of Pediatrics, Univ. of Pittsburgh School of Medicine, Research Advisory Committee Grant, NIH Director's New Innovator Award DP2-AR081047, NIH R01-AR078872 (to D.U.K).

Author contributions:

Conceptualization: O.A. and D.U.K; Investigation: O.A., A.M., M.J., M.E.P., Y.G., Y.W., J.F., T.M.S, R.A.C.; Formal analysis and visualization: O.A., A.M., M.E.P., Y.G., Y.W., H.B., D.U.K; Writing: O.A. and D.U.K.; Supervision, D.U.K.

References:

1. Bachrach U, Wang YC, Tabib A. Polyamines: New cues in cellular signal transduction. News in Physiological Sciences. 2001;16(3).

2. Landau G, Bercovich Z, Park MH, Kahana C. The role of polyamines in supporting growth of mammalian cells is mediated through their requirement for translation initiation and elongation. Journal of Biological Chemistry. 2010;285(17).

3. Hesterberg R, Cleveland J, Epling-Burnette P. Role of Polyamines in Immune Cell Functions. Medical Sciences. 2018;6(1).

4. Kemaladewi DU, Benjamin JS, Hyatt E, Ivakine EA, Cohn RD. Increased polyamines as protective disease modifiers in congenital muscular dystrophy. Hum Mol Genet. 2018;27(11).

5. Akinyele O, Wallace HM. Understanding the Polyamine and mTOR Pathway Interaction in Breast Cancer Cell Growth. Medical Sciences. 2022 Sep 10;10(3):51.

6. Arruabarrena-Aristorena A, Zabala-Letona A, Carracedo A. Oil for the cancer engine: The cross-talk between oncogenic signaling and polyamine metabolism. Vol. 4, Science Advances. 2018.

7. Lewandowski NM, Ju S, Verbitsky M, Ross B, Geddie ML, Rockenstein E, et al. Polyamine pathway contributes to the pathogenesis of Parkinson disease. Proc Natl Acad Sci U S A. 2010;107(39).

8. Soda K. The mechanisms by which polyamines accelerate tumor spread. Journal of Experimental and Clinical Cancer Research. 2011;30(1).

688   9.   Milovic V, Turchanowa L. Polyamines and colon cancer. In: Biochemical Society Transactions.
689   2003.

690   10.   Schwartz CE, Peron A, Kutler MJ, Snyder-Robinson Syndrome ;, Adam MP, Ardinger HH, et al.
691   Snyder-Robinson Syndrome Synonym: Spermine Synthase Deficiency. 2013.

692   11.   de Alencastro G, McCloskey DE, Kliemann SE, Maranduba CMC, Pegg AE, Wang X, et al. New SMS
693   mutation leads to a striking reduction in spermine synthase protein function and a severe form of
694   Snyder-Robinson X-linked recessive mental retardation syndrome. J Med Genet. 2008;45(8).

695   12.   Larcher L, Norris JW, Lejeune E, Buratti J, Mignot C, Garel C, et al. The complete loss of function
696   of the SMS gene results in a severe form of Snyder-Robinson syndrome. Eur J Med Genet.
697   2020;63(4).

698   13.   Valera Ribera C, Martinez-Ferrer, Flores Fernández E, Vázquez Gómez I, Orenes Vera A, Valls
699   Pascual E, et al. Snyder-Robinson syndrome: differential diagnosis of osteogenesis imperfecta.
700   Osteoporosis International. 2022;33(5).

701   14.   Cason AL, Ikeguchi Y, Skinner C, Wood TC, Holden KR, Lubs HA, et al. X-linked spermine synthase
702   gene (SMS) defect: The first polyamine deficiency syndrome. European Journal of Human
703   Genetics. 2003;11(12).

704   15.   Li C, Brazill JM, Liu S, Bello C, Zhu Y, Morimoto M, et al. Spermine synthase deficiency causes
705   lysosomal dysfunction and oxidative stress in models of Snyder-Robinson syndrome. Nat
706   Commun. 2017;8(1).

707   16.   Tao X, Zhu Y, Diaz-Perez Z, Yu SH, Foley JR, Stewart TM, et al. Phenylbutyrate modulates
708   polyamine acetylase and ameliorates Snyder-Robinson syndrome in a Drosophila model and
709   patient cells. 2022; Available from: https://doi.org/10.1172/jci.

710   17.   MacKintosh CA, Pegg AE. Effect of spermine synthase deficiency on polyamine biosynthesis and
711   content in mice and embryonic fibroblasts, and the sensitivity of fibroblasts to 1,3-bis-(2-
712   chloroethyl)-N-nitrosourea. Biochemical Journal. 2000;351(2).

713   18.   Wang X, Ikeguchi Y, McCloskey DE, Nelson P, Pegg AE. Spermine synthesis is required for normal
714   viability, growth, and fertility in the mouse. Journal of Biological Chemistry. 2004;279(49).

715   19.   Meyer RA, Henley CM, Meyer MH, Morgan PL, McDonald AG, Mills C, et al. Partial deletion of
716   both the spermine synthase gene and the Pex gene in the X-linked hypophosphatemic, Gyro (Gy)
717   mouse. Genomics. 1998;48(3).

718   20.   Lyon MF, Scriver CR, Baker LR, Tenenhouse HS, Kronick J, Mandla S. The Gy mutation: Another
719   cause of X-linked hypophosphatemia. Proc Natl Acad Sci U S A. 1986;83(13).

720   21.   Laskowski S. THE JACKSON LABORATORY AND SNYDER-ROBINSON FOUNDATION ANNOUNCE
721   NEW RESEARCH MODELS FOR SNYDER-ROBINSON SYNDROME.

722   22.   Kesler SR, Schwartz C, Stevenson RE, Reiss AL. The impact of spermine synthase (SMS) mutations
723   on brain morphology. Neurogenetics. 2009;10(4).

724   23.   Starks RD, Kirby PA, Ciliberto MA, Hefti MM. Snyder-Robinson Syndrome: An Autopsy Case
725   Report. Autops Case Rep. 2018;8(3).

726   24.   Tsien JZ, Huerta PT, Tonegawa S. The essential role of hippocampal CA1 NMDA receptor-
727   dependent synaptic plasticity in spatial memory. Cell. 1996;87(7).

728   25.   Khan M, Gasser S. Generating primary fibroblast cultures from mouse ear and tail tissues. Journal
729   of Visualized Experiments. 2016;2016(107).

730   26.   Kabra PM, Lee HK, Lubich WP, Marton LJ. Solid-phase extraction and determination of dansyl
731   derivatives of unconjugated and acetylated polyamines by reversed-phase liquid
732   chromatography: Improved separation systems for polyamines in cerebrospinal fluid, urine and
733   tissue. J Chromatogr B Biomed Sci Appl. 1986;380(C).

734   27.   Clarkson AN, Liu H, Pearson L, Kapoor M, Harrison JC, Sammut IA, et al. Neuroprotective effects
735   of spermine following hypoxia-ischemia-induced brain damage: A mechanistic study. The FASEB
736   Journal. 2004 Jul 7;18(10):1114–6.

737   28.   Ramsay AL, Alonso-Garcia V, Chaboya C, Radut B, Le B, Florez J, et al. Modeling Snyder-Robinson
738   Syndrome in multipotent stromal cells reveals impaired mitochondrial function as a potential
739   cause for deficient osteogenesis. Sci Rep. 2019;9(1).

740   29.   Monelli E, Villacampa P, Zabala-Letona A, Martinez-Romero A, Llena J, Beiroa D, et al. Angiocrine
741   polyamine production regulates adiposity. Nat Metab. 2022;4(3).

742   30.   Slotkin TA, Bartolome J. Role of ornithine decarboxylase and the polyamines in nervous system
743   development: A review. Brain Res Bull. 1986;17(3).

744   31.   Rubin MA, Berlese DB, Stiegemeier JA, Volkweis MA, Oliveira DM, dos Santos TLB, et al. Intra-
745   Amygdala Administration of Polyamines Modulates Fear Conditioning in Rats. Journal of
746   Neuroscience. 2004;24(9).

747   32.   Subramaniam S, O'connor MJ, Masukawa LM, McGonigle P. Polyamine effects on the NMDA
748   receptor in human brain. Exp Neurol. 1994;130(2).

749   33.   Davies RD, Gabbert SL, Riggs PD. Anxiety disorders in neurologic illness. Curr Treat Options
750   Neurol. 2001;3(4).

751   34.   Guo CY, Sun L, Chen XP, Zhang DS. Oxidative stress, mitochondrial damage and
752   neurodegenerative diseases. Neural Regen Res. 2013;8(21).

753   35.   Lee M, Matsunaga N, Akabane S, Yasuda I, Ueda T, Takeuchi-Tomita N. Reconstitution of
754   mammalian mitochondrial translation system capable of correct initiation and long polypeptide
755   synthesis from leaderless mRNA. Nucleic Acids Res. 2021;49(1).

756   36.   Christian BE, Haque ME, Spremulli LL. The effect of spermine on the initiation of mitochondrial
757   protein synthesis. Biochem Biophys Res Commun. 2010;391(1).

758   37.   Zorov DB, Juhaszova M, Sollott SJ. Mitochondrial reactive oxygen species (ROS) and ROS-induced
759   ROS release. Vol. 94, Physiological Reviews. 2014.

760   38.   Ha HC, Sirisoma NS, Kuppusamy P, Zweier JL, Woster PM, Casero RA. The natural polyamine
761   spermine functions directly as a free radical scavenger. Proc Natl Acad Sci U S A. 1998;95(19).

762   39.   Sava IG, Battaglia V, Rossi CA, Salvi M, Toninello A. Free radical scavenging action of the natural
763   polyamine spermine in rat liver mitochondria. Free Radic Biol Med. 2006;41(8).

764   40.   Elustondo PA, Negoda A, Kane CL, Kane DA, Pavlov E v. Spermine selectively inhibits high-
765   conductance, but not low-conductance calcium-induced permeability transition pore. Biochim
766   Biophys Acta Bioenerg. 2015;1847(2).

768