Title: SARS-CoV-2 variant natural evolution in K18-ACE2 mice increases virulence and

produces treatment resistance-linked variant alleles.

Authors: Julian Daniel Sunday Willett[1,2,3], Annie Gravel[4], Isabelle Dubuc[4], Leslie Gudimard[4],

Paul Fortin[4,5,6], Ju-Ling Liu[2,8], Jose Avila Cervantes[9], Jose Hector Galvez[7], Melissa Zwaig[2,8],

Anne-Marie Roy[2,8], Sally Lee[2,8], Shu-Huang Chen[2,8], Jiannis Ragoussis[2,8]*, Louis Flamand[4,10]*

Affiliations:

1. Quantitative Life Sciences Ph.D. Program, McGill University, Montreal, QC, Canada

2. McGill University Genome Centre, Montreal, QC, Canada

3. Lady Davis Institute, Jewish General Hospital, Montreal, QC, Canada

4. Axe maladies infectieuses et immunitaires, Centre de Recherche du Centre Hospitalier

Universitaire de Québec- Université Laval, Canada

5. Centre de Recherche ARThrite-Arthrite, Recherche et Traitements, Université Laval,

Québec, QC, Canada.

6. Division of Rheumatology, Department of Medicine, CHU de Québec-Université Laval,

Québec, QC, Canada

7. Canadian Centre for Computational Genomics, McGill University, Montreal, QC, Canada

8. Department of Human Genetics, McGill University, Montreal, QC, Canada

9. Advanced Genomic Technologies Laboratory, McGill Genome Centre, McGill University,

Montreal, QC, Canada.

10. Département de microbiologie-infectiologie et d'immunologie, Université Laval, QC,

Canada.

*Co-corresponding authors

Abstract:

The continuing COVID-19 pandemic is partially due to viral evolution reducing vaccine and treatment efficacy. Others have observed serial infections in mice with reference SARS-CoV-2 lineage IME-BJ05 produce adapted strains with significantly greater infectivity and mortality. We investigated the impact of long-term B.1.351 (Beta) and B.1.617.2 (Delta) lineage viral evolution in K18-ACE2 mice, which express human ACE2 receptor. We infected mice with these unmodified variants in a BSL-3 laboratory and studied viral evolution across 20 passages without selective pressure, sequencing virus at key points. Our IRB determined the research was not gain-of-function. We observed variant alleles with documented contributions towards evading vaccine-induced immunity arise *de-novo*, including Omicron-characteristic mutation S371F. Passage 20 (P20) viruses were more virulent, and P20 Delta was significantly more resistant to antibody neutralization. These developments occurred rapidly, emulating pandemic progression in non-immune mammals. Our model could be rapidly adapted to include selective pressures to study the evolution of viruses resistant to antivirals or vaccine-induced immunity.

Keywords: SARS-CoV-2, COVID-19, pandemic modelling, treatment resistance, viral evolution.

Introduction:

Part of the difficulty in responding to the COVID-19 pandemic has been predicting viral evolution and its impact on clinically relevant traits, such as infectivity, treatment resistance, and disease severity[1]. The basic biology of the severe acute respiratory syndrome Coronavirus 2 (SARS-CoV-2), the virus that causes COVID-19, is well understood. However, knowledge of SARS-CoV-2 evolution is limited. Many RNA viruses, such as SARS-CoV-2, exist as quasi-species meaning that within a given viral population, a multitude of alleles (major and minor) are present. These alleles give plasticity to the viral population allowing for rapid adaptation/selection to a variety of circumstances[2]. Viral evolution is also complex and affected by several factors, including host genetic background[3], host immune status[4], the organs targeted by the virus[5], and a population's collective immunity[6]. As a result, it has been difficult to predict the emergence of new variants with altered virulence.

Several variants of concern (VOC) have emerged, each with mutations providing dissemination advantages[1]. For example, the SARS-CoV-2 alpha variant (B.1.1.7) gained a growth advantage and rapidly spread globally due to the spike protein N501Y mutation that enhances affinity for the cellular entry receptor, angiotensin-converting enzyme 2 (ACE2)[7]. Other VOC, such as the Delta (B.1.617.2) variant, gained advantages with additional substitutions in spike, including L452R[8], T478K[9], and P681R[10] that affect viral transmissibility and antibody neutralization for naturally or artificially vaccinated individuals[11]. To better predict changes in SARS-CoV-2 and reduce the impact of a future pandemic, time effective models must be defined for studying viral changes over successive generations.

In 2020, Gu *et al.* studied short-term viral evolution by infecting aged mice with the SARS-CoV-2 reference strain (IME-BJ05) and collecting lung tissue at set timepoints following infection, using this isolate to infect successive generations of mice[12]. While wild-type mice are thought less

susceptible to the virus as mouse Ace2 is not used by SARS-CoV-2 as its receptor, Gu *et al.*

demonstrated that serial passaging of lung homogenate across mice produced a mouse-

adapted strain (MASCp6) that causes pulmonary symptoms, matching adaptations observed by

others who previously applied the technique to influenza[12,13]. After six passages, they found

several clinically relevant genetic changes, including de-novo appearance of A23063T that

changes spike to N501Y[12]. This mutation is associated with enhanced viral infection and

transmission[7]. Subsequently, the group passaged the MASCp6 isolate an additional 30 times

and characterized the MASCp36 isolate[14]. MASCp36 contained several mutations (i.e. K417N,

Q493H and N501Y mutations) in spike conferring greater Ace2 receptor binding affinity,

increased infectivity and greater virulence[14]. Others replicated this study using the same viral

variant, observing similar development of clinically relevant variant alleles[15].

Since these studies, new lineages arisen, with the present being Omicron, negatively impacting

treatment efficacy[1]. Others have studied viral dynamics using recent lineages in serial passages

in cells, observing similar development of directional selection and clinically concerning

mutations[16,17]. Given the pandemic's prolonged nature, understanding longitudinal viral changes

in a mammalian host is pressing. We adapted Gu *et al.*'s approach to evaluate how viral

lineages more closely related to the present lineages, B.1.351 (Beta) and B.1.617.2 (Delta),

changed over twenty passages[12] in transgenic mice with human ACE2 receptors, instead of

murine. Given clinically relevant mutations arising in Gu *et al.*'s studies[12,14], we sought to

determine if impactful variant alleles would develop in more recent viral variants. We report on

the *de novo* appearance and persistence of variant alleles associated with treatment resistance

with serially passaged virus demonstrating increased virulence and antibody resistance.

Results:

Variant allele frequency of alleles across passages. To ensure effective comparisons between key viral passages, we downsampled sequencing reads for each period of the study (**Figure 1**). The minimum number of reads per sample in the first part of the study across passage 0 (P0) and P10 was 8,812,604. The minimum number of reads per sample in the second part of the study across P13, P17, and P20 was 34,253,784. In Beta lineage, there was a significantly lower variant allele frequency (VAF) in P17 and P20 versus other passages (adjusted p-value < 0.05), with a significant difference in Delta comparing P10 and P20 (**Figure 2a-b**). As there was no significant change in allele depth across the study, these VAF differences could be due to growing allelic diversity. Such significant changes, less consistent with random allelic drift, could align with suggested mechanisms of within-host evolution of the variant allele pool, per Tonkin-Hill *et al*.[18]. Sample contamination is less likely as there was no significant difference in VAF or variant allele depth between samples for either viral lineage (**Figure 2c-d**).

Gene involvement and variant allele functional consequence severity did not significantly vary across passages. To evaluate for large-scale variant allele shifts across the study, we used chi-squared tests, adjusted for multiple-testing, to compare the net proportion change in variant allele subcategories by passage. There was no statistically significant change in the proportion of variant alleles involving a specific gene or severity of variant allele predicted functional consequence across the study (adjusted p-value > 0.05) (**Figure 3**), suggesting that the overall allele population did not significantly change in our model without experimental selective pressures [6].

Development of variant alleles associated with clinically relevant traits. To study viral evolution in our model at the single variant allele level, we identified those variant alleles that changed in frequency across passages, including those that arose *de novo* and disappeared, and annotated them for clinically relevant traits. We observed 82 unique variant alleles across our

study. There were 41 whose VAF changed with 32 having clinically-relevant annotations,

including interactions with human proteins, mutations that may confer drug resistance, variant

alleles associated with the Omicron lineage, mutations associated with antibody escape, and

mutations at vaccine targets (**Table 1, Supplemental Table 1**) [19,20]. A notable novel variant

included C22674T, or Spike S371F, that arose *de-novo* in two animals in P20 for Beta virus and

two animals in P13 for Delta virus (**Figure 4**). The allele is associated with protein interactions,

antibody escape, affects a vaccine target, and is an Omicron-specific mutation (**Table 1**) [20,21].

We thus observed an Omicron specific mutation in a study not involving Omicron viral lineages.

Several other variants changing in frequency also conferred possible drug resistance and

targeted the vaccine site (**Table 1**) [20].

Most alleles in the P0 Beta isolate, passaged several times *in vitro* in Vero cells, encoded a

Spike protein with a tryptophan (W) at position 682 (C23606T), destroying the furin cleavage

site (**Table 1**). We observed a rapid selection of viruses having the reference allele encoding an

arginine (R) at position 682 and functional furin site in all animals infected with Beta variant

starting in P10 (**Table 1**). This suggests that the furin site provides a growth advantage under *in*

vivo conditions and supports a previous report indicating the Spike furin site being important for

SARS-CoV-2 virulence in K18-ACE2 mice [22].

Later passage viruses cause greater weight loss and viral loads and demonstrate reduced

neutralization by immune sera. To study our model's viral evolutionary impact on the organism

level, we compared clinical scores of mice infected with early and late passaged virus. Unlike

Gu *et al.*'s study that required adaptation of SARS-CoV-2 to a murine Ace2 receptor for

infectivity, our study made use of transgenic mice expressing the human ACE2 receptor [12]. We

determined whether repeated passaging of SARS-CoV-2 in K18-ACE2 mice affected virulence

by infecting animals with P0 and P20 viruses with clinical scores and viral loads determined. On

day 3 post-infection, no differences in viral load were noted between Beta P0 and Beta P20

viruses with mice infected with P20 Delta virus having a significant increase (6.3x) in lung viral loads (**Figure 5A**). Both P20 Beta and P20 Delta viruses caused more rapid and greater weight loss than P0 viruses (**Figures 5B-C**) with similar survival curves (data not shown).

We then assessed whether repeated passages in K18-ACE2 mice would result in viruses capable of infecting non transgenic mice. C57BL6 mice were infected intranasally with P0 and P20 viruses and lung infectious viral loads determined 3 days later. Both P0 and P20 Beta viruses successfully replicated in mouse lungs with both viruses producing similar viral loads (**Supplementary Figure 2**). None of the P0 or the P20 Delta viruses generated viral loads in lungs of B6 mice (**Supplementary Figure 2**).

Our results indicate that Delta P20 viruses generate a greater viral load that P0. To determine if this was related to growth kinetic differences between P0 and P20 viruses, we performed *in vitro* growth curve analysis. We observed two independently evolved P20 Delta viruses replicating more efficiently after 24 h versus P0, yielding 4-5X more infectious virus after 24h of infection (**Figure 6**). This difference in viral titers was no longer observed after 48 h (**Figure 6**).

We next determined the ability of sera from vaccinated subjects to neutralize passaged viruses, comparing it to parental P0 viruses. Consistent with previous reports, sera from vaccinated subjects had varied neutralizing potential depending on the variant, with neutralization titers being greatest against the original Wuhan-like strain and lowest for the Beta isolate (**Figure 7A**) [23,24]. The same sera were analyzed for their ability to neutralize P20 viruses. P20 Beta was neutralized equally (13 out of 24) or more efficiently (8 of 24) that the P0 virus (**Figure 7B**). In contrast, P20 Delta virus was significantly less sensitive to antibody neutralization by vaccinated subject sera than P0 ($p < 0.0002$) where 15 of 24 sera showed decreased neutralizing potential toward the P20 Delta virus relative to P0 (**Figure 7C**).

Discussion:

In this study, we serially infected mice with Beta or Delta lineage virus for twenty passages without experimental selective pressures. The evolved viruses developed and maintained variant alleles associated with treatment resistance and later lineages like Omicron (**Figure 4, Table 1**), were more virulent (**Figure 5**), grew faster and to higher levels (Delta) (**Figure 6**) and were less sensitive to antibody neutralization (Delta) (**Figure 7**), demonstrating immune escape potential. These developments occurred within roughly two months, illustrating the model's potential in rapidly emulating pandemic progression that could facilitate treatment/vaccine development less sensitive to viral evolution. The model can readily be adapted to mimic viral adaptation in the context of immune or anti-viral selection pressures, affording rapid results in a system often seen in real world conditions.

Our model lacked experimental selective pressures, mimicking the evolution of SARS-CoV-2 in a naïve population. Using K18-ACE2 transgenic mice eliminated the selective pressure of adapting virus to the murine Ace2 receptor. The Beta P0 virus harbored the N501Y mutation previously reported to enable adaptation of reference SARS-CoV-2 in non transgenic mice [12]. This mutation was maintained throughout passages, suggesting that this virus can infect cells through the murine Ace2 receptor as well (**Figure 4**). Our data supports this, as non transgenic C57BL6 mice were equally susceptible to infection by Beta P0 and P20 viruses. This capacity of Beta to use human and murine receptors might explain, at least in part, the greater virulence of the Beta isolate in ACE2 transgenic mice relative to other viral isolates, such as Wuhan and Delta lacking the N501Y mutation in transgenic mouse studies [25].

The Delta P0 virus lacked the N501Y mutation and never acquired it following repeated passages in ACE2 transgenic mice. This highlights that when lacking selective pressure, such as neutralizing antibodies, the appearance of N501Y is not favored (**Figure 4**). Acquisition of

N501Y in the human population was first reported almost a year after the initial report of the

SARS-CoV-2 epidemic, suggesting that this mutation arose during the development of herd

immunity [26].

Of the 41 variant alleles that arose *de-novo* or changed in frequency during the study, several

are of clinical interest (**Figure 4, Table 1**). A4981G, which arose *de-novo* and persisted in Beta

lineage virus in P17 (**Figure 4**), was predicted to possibly confer drug resistance to papain-like

protease inhibitors [27], such as ritonavir used clinically in combination with nirmatrelvir that

protects patients against severe COVID-19 [28,29]. C22674T, or spike S371F, which arose *de-*

novo and persisted in two animals in P20 for Beta and P13 for Delta virus (**Figure 4**), represents

an Omicron-specific mutation arising in a study that did not use Omicron. This last mutation has

been previously reported to facilitate antibody escape (**Table 1**) [21]. Considering that the

mutation S371F is reported to bear a significant fitness cost [30], this might explain why isolated

S371L/F mutations are not viable or have not yet been detected in isolation. We would argue

that mutations of residues E484 present in Delta and K417/E484/N501 present in Beta Spike

proteins, also found together with the S371F mutation in Omicron BA.2, represent such

compensatory mutations. G23593T, or spike Q677H, which disappeared from Beta virus

samples after P0, has been determined to enhance treatment resistance and increase viral

infectivity [31]. The reversion of Spike amino acid 677 to Q after passage in mice could explain the

increased neutralization activity for some of the sera against Beta P20 relative to P0 (**Figure 7**).

Variants that could impair treatment efficacy appeared and disappeared in this study,

showcasing the model's potential in evaluating diverse variants.

This study has limitations. Here we used the K18-ACE2 transgenic mice but the mechanisms of

new viral variants arising in humans could significantly differ. This is relevant as host factors are

believed to contribute to viral variant allele diversity [32]. Our model lacked experimental selective

pressures, while the COVID-19 pandemic has been accompanied by contact limitations and

vaccines with the dominant SARS-CoV-2 evolutionary mechanism believed to be natural

selection [6]. Development or dominance of new variants likely varies with these pressures,

necessitating reproducing this model with them. However, a baseline comparison model is

required to get the most from these supplemental models, hence the relevance of our model.

We studied Beta and Delta lineage virus, which are no longer dominant. Patterns of genetic

changes observed in these lineages could vary from Omicron. There was no statistical change

in the variant allele population's involvement of specific genes or predicted functional

consequence when comparing Beta and Delta lineages across passages, which is reassuring

for generalizability. However, virulence significantly increased across the study, despite this lack

of population-level changes (**Figure 5**). This could suggest greater value in studying changes in

single variant alleles over allele populations, as done in this study, as they are associated with

fitness-related traits and can capture evolutionary changes of more subtle mechanisms, such as

increasing virus adaptation for cellular infection. Single or combinations of variant allele changes

could explain the change in virulence and antibody neutralization observed in this study (**Figure**

4).

Conclusions:

The model presented rapidly emulated pandemic progression in a naïve population, with virus

accumulating variant alleles that could limit therapies for severe disease, becoming more

virulent, and ultimately becoming treatment resistant in mice harboring human ACE2 receptor.

These results support that serial viral passaging in mice is a useful method to rapidly study viral

evolution enabling development of targeted anti-viral therapies.

Methods:

Viruses. SARS-CoV-2 Wuhan-like strain (LSPQ, B1 lineage) was obtained from the Laboratoire de Santé Publique du Québec ([LSPQ] Sainte-Anne-de-Bellevue, QC, Canada). SARS-CoV-2 Beta strain was obtained from BEI resources and SARS-CoV-2 Delta strain from the BC Centers for Disease Control. Unmodified SARS-CoV-2 strains were propagated on Vero cells (American Type Culture Collection, Manassas, Virginia, USA).

Determination of the viral titer. Vero cells were plated in a 96-well plate ($2x10^4$/well) and infected with 200µl of serial dilution of the viral preparation or lung homogenate in the M199 media supplemented with 10mM HEPES pH 7.2, 1mM of sodium pyruvate, 2.5g/L of glucose, 5µg/mL *Plasmocin*® and 2% FBS. Three days post-infection plates were analyzed using a EVOS M5000 microscope (ThermoFisher Scientific, Waltham, MA, USA) and the viral titer was determined using the Karber method [33].

Mouse models. B6.Cg-Tg(K18-hACE2)2Prlmn/J (stock#3034860) and B6 mice were purchased from the Jackson Laboratories (Bar Harbor, ME). All mouse studies were conducted in a BSL-3 laboratory. One K18-ACE2 mouse was intranasally infected with either B.1.351 (Beta) or B.1.617.2 (Delta) SARS-CoV-2 lineages. Following 3-4 days, the animals were sacrificed and lung homogenate collected. The homogenate was used to intranasally infect sequential animals, defined as passage. This was repeated for 10 passages. After passage 10 (P10), virus in lung homogenate was sequenced. K18-ACE2 mice were then intranasally infected with Beta P10 or Delta P10 with viruses passaged in three mice at a time for an additional 10 times (total of 20 passages) with virus in lung homogenate sequenced following passages 13, 17, and 20 (Figure 1). Viral stocks were then made from lung homogenates of Beta and Delta P20 viruses. K18-ACE2 mice were infected intranasally with 500 $TCID_{50}$ of P0 or P20 viruses. Lung viral loads were determined on day 3 post-infection with weight loss and survival monitored daily for up to 9 days.

Infectivity of P0 and P20 viruses to non-transgenic mice. C57BL6 mice were infected

intranasally with 3000 TCID$_{50}$ of P0 and P20 Beta and Delta viruses. Three days later, mice

were euthanized, and lungs collected for viral load (titer) determination.

RNA extraction: Up to 30 mg of lung tissue were used for RNA extraction using the Bead Mill

Tissue RNA Purification Kit and the Bead Mill Homogenizer, following manufacturer's

instructions.

Viral sequencing. RNA extracts were processed for reverse transcription (step a) and targeted

SARS-CoV-2 amplification using the ARTIC V3 or V4.1 primer scheme (https://github.com/artic-

network/primer-schemes/tree/master/nCoV-2019) (step b). Samples were purified (step b) and

Nextera DNA Flex library preparation (step c) was performed for Illumina PE150 paired-end

amplicon sequencing on a NovaSeq or MiSeq instrument at the McGill Genome Centre using

best practices. Each sample was sequenced twice on different days to obtain a target of

minimum 10 million reads per sample. The detailed protocols can be accessed with the

following links.

Step a: dx.doi.org/10.17504/protocols.io.bjgekjte.

Step b: dx.doi.org/10.17504/protocols.io.ewov18e4ygr2/v2.

Step c: dx.doi.org/10.17504/protocols.io.bjgnkjve.

Genome data processing. Following sequencing, the reads from each sample were used to

call variants using Freebayes v1.3.6 [34] and the results were saved as a VCF file which was used

to compare genomic variation across passages. Others have called variants with other tools

such as DeepSNV [18] and VarScan [35]. DeepSNV was not used as it only investigates SNVs [36],

when indels have been found to confer selective advantages [37]. Freebayes has been observed

to perform slightly superiorly than VarScan when calling variants in wastewater samples [38] and

in internal benchmarking (data not shown). The process is described briefly as follows: first, the

raw reads from each sample were aligned with BWA MEM v0.7.17 [39] to the reference strain

genome sequence (NC_045512.2). The resulting alignments were sorted with duplicate reads

flagged. The minimum number of aligned reads for each was logged, and each sample was

again aligned and processed, then randomly down sampled to match this minimum number of

reads. All variants with a quality of less than 20 or a depth below 10 were removed from the

analysis. Any variant alleles that overlapped with ARTIC sequencing primers were also

removed. Merged VCF files, representing the combination of reads across two sequencing

batches for each sample, were used for the paper's main results.

Variant annotation. Every variant allele was inputted into COVID-19 Ensembl variant effect

predictor (VEP) (https://covid-19.ensembl.org/index.html) to determine functional consequences

and the involved gene and amino acid change [40]. Variants whose VAF changed at least once

across passages during the study were logged and input to the UCSC Genome Browser

(https://genome.ucsc.edu/), checking for annotations of antibody escape, CD8 escape, vaccine

targets, drug resistance, or variants of concern [20].

Antibody neutralization studies. Sera from twenty-four healthy vaccinated (3x) individuals

were analyzed for neutralizing activity against Wuhan-like, and Beta and Delta P0 and P20

viruses. Serially diluted sera (in quadruplicate) were incubated with 100 $TCID_{50}$ of virus for 1 h

at 37˚C before addition to Vero E6 cells. Three days later, the 96-well plates were observed

using an inverted microscope for signs of infection. Neutralization titers were determined and

defined as the highest serum dilution preventing infection.

Ethics. The study was conducted in accordance with the Declaration of Helsinki, and mouse

protocols were approved by the Comité de protection des animaux de l'Université Laval. Our

IRB determined that our work did not constitute gain-of-function research as virus was

unmodified and represented circulating variants with the study conducted without experimental

selective pressures. Sera obtained from humans were procured following ERB approval (ERB

#2022-6204).

Statistics and reproducibility. All statistical analyses were conducted using R/4.2.1. All

statistical comparisons between passages were corrected for multiple testing using a Benjamini-

Hochberg p-value adjustment. Reproducibility is described in each method.

Source code. All code used in this study, from sample processing to data analysis, are publicly

available on Github at the following link:

https://github.com/juliandwillett/SARS_CoV_2_Serial_Passaging_Study.

Data availability. VCF files used to complete computational analyses are available by

reasonable request.

Declarations:

Competing interest.

All authors declare no competing interests.

Authors contributions. Julian Daniel Sunday Willett conceived of the computational analyses

needed for this study, completed all computational analysis of viral sequencing data, and wrote

the majority of the manuscript. Louis Flamand and Jiannis Ragoussis conceived of the study.

Louis Flamand contributed to manuscript revisions, wrote sections relevant to animal models

and antibody neutralization studies and performed analysis of results. Jiannis Ragoussis

supervised sequencing data generation and analysis. Isabelle Dubuc and Leslie Gudimard

performed all works involving live SARS-CoV-2 viruses. Annie Gravel participated in the

sequence analysis and methodological design of experiments. Paul Fortin and Ines Colmegna

provided key reagents. Jose Hector Galvez contributed bioinformatics analysis review and

manuscript revisions. Ju-Ling Liu, Jose Avila Cervantes, Anne-Marie Roy, Sally Lee and Shu-

Huang Chen supported the SARS-CoV-2 viral sequencing.

Acknowledgments.

This research was enabled in part by support provided by Calcul Québec (calculquebec.ca) and

the Digital Research Alliance of Canada (alliancecan.ca). This study was supported by the

Canadian Institutes for Health Research (CIHR 476575) to JDSW, and an operating grant to the

Coronavirus Variants Rapid Response Network (CoVaRR-Net) to LF and IR and CIHR grant

(GA1-117694) to LF. The funding body had no role in the design of the study or collection,

analysis, interpretation, or reporting of the data. We thank Ines Colmegna for providing

vaccinated patient sera. We thank Silvia Vidal for her guidance regarding SARS-CoV-2

evolutionary mechanisms. ChatGPT was used to decrease the word count in the abstract with

output carefully examined for representing our results.

Figures:

Figure 1. **Study design**. First, SARS-CoV-2 ACE2-adapted mice were inoculated with either

B.1.351 (Beta) or B.1.617.2 (Delta) virus and left for three days for virus to proliferate. They

were then sacrificed with the lung homogenate used to infect another mouse. This was repeated

for ten passages. Second, P10 virus for each lineage was used to infect three additional mice

and the process was repeated for an additional ten passages. Third, P0 and P20 virus for each

lineage was used to infect additional mice to compare weight loss and survival between early

and late passage virus, and test non-adapted mouse susceptibility to the virus. Antibody

neutralization of virus was also compared.

Figure 2. Boxplots of variant allele frequency by (a-b) passage and (c-d) sample. Each N

refers to the number of variant alleles for each MAF and in total against the NC_045512.2

reference sequence. Single samples were sequenced for P0 and P10 with three samples from

P13, P17, and P20. Box fill in (c-d) signifies sample number. For significance, n.s.: adjusted p-

value > 0.05, *: adjusted p-value > 0.01, **: adjusted p-value ≤ 0.01.

 Figure 3. **Gene involvement and predicted variant allele functional consequence severity**

 (VEP) in variant allele population did not significantly change (adjusted p-value > 0.05)

 with passage.

364 **Figure 4. Variants of clinical relevance arose de novo and grew to dominance**. VAF:

365 variant allele frequency. P refers to passage.

366   **Table 1. Annotations of select variants that changed in frequency across the study.**

Variant	Amino Acid Change	Annotation
A4981G	ORF1ab T1572T	Mutations that may confer drug resistance[20]
C22674T	Spike S371F	Protein interacting, antibody escape, vaccine target, Omicron-associated variant allele[20]
G23593T	Spike Q677H	Protein interacting, vaccine target[20], enhances treatment resistance[31], increases infectivity[31]

Figure 5. **Pathogenesis of P0 and P20 viruses in K18-ACE2 mice**. Mice (n = 15/group) were

infected with 500 $TCID_{50}$ of P0 or P20 Beta and Delta viruses. (A) On day 3 post-infection, mice

(n = 5) were euthanized and lungs collected for infectious viral load determination. Results are

expressed as mean ± SD SARS-CoV-2 $TCID_{50}$/mg of lung tissue. (B-C) The weight of each

mouse (n = 10/group) was monitored daily throughout the experiment and reported as mean

percentage ± SD of weight relative to day 0. **: $p \leq 0.008$ as determined using non-parametric

Mann-Whitney test.

Figure 6. **Growth kinetics of P0 and P20 Delta viruses**. P0 and two independently evolved

P20 Delta viruses were used to infect Vero cells at a multiplicity of infection of 0.005. Cell-free

supernatants were collected at 24 h and 48 h and used for viral titration. Results are expressed

as mean ± SD $TCID_{50}$/mL. *: p=0.02 as determined using an unpaired t-test.

Figure 7. **Neutralization of P0 and P20 viruses with sera from vaccinated subjects**. (A)

Neutralization assay of Wuhan-like, Beta, and Delta viruses using sera from 24 vaccinated

subjects. Individual results and mean ± SD neutralization titer against different SARS-CoV-2

isolates are presented. ****: p<0.0001 determined using one way-ANOVA. ns: not statistically

significant. (B) Pair-wise comparison of the sera used in A in neutralization assay against P0

and P20 Beta viruses. (C) Pair-wise comparison of the sera used in A in neutralization assay

against P0 and P20 Delta viruses. ***: p>0.002 determined using paired t-test.

391   **Supplemental Figures**

392   **Supplemental Table 1. Table of variant alleles that changed MAF during the study.**

393   **Available in "Supplemental Tables.csv".**

Supplemental Figure 1. **Viral load comparison between P0 and P20 virus in K18-ACE2**

mice.

Supplementary Figure 2. Infection of C57BL6 mice with P0 and P20 Beta and Delta

viruses. Mice (n = 5/group) were infected intranasally with 3000 $TCID_{50}$ of P0 or three

independently evolved Beta P20 or two independently evolved Delta viruses. On the third day

post infection, mice were euthanized and lungs collected for infectious viral load determination.

Individual results and mean ± SD $TCID_{50}$/mg of lung tissue are presented.

References:

1. Jacobs, J.L., Haidar, G. & Mellors, J.W. COVID-19: Challenges of Viral Variants. *Annu Rev Med* (2022).

2. Sun, F. *et al.* SARS-CoV-2 Quasispecies Provides an Advantage Mutation Pool for the Epidemic Variants. *Microbiol Spectr* **9**, e0026121 (2021).

3. Velavan, T.P. *et al.* Host genetic factors determining COVID-19 susceptibility and severity. *EBioMedicine* **72**, 103629 (2021).

4.   Belsky, J.A. *et al.* COVID-19 in immunocompromised patients: A systematic review of

cancer, hematopoietic cell and solid organ transplant patients. *J Infect* **82**, 329-338

(2021).

5.   Van Cleemput, J. *et al.* Organ-specific genome diversity of replication-competent SARS-

CoV-2. *Nat Commun* **12**, 6612 (2021).

6.   Wang, R., Chen, J. & Wei, G.W. Mechanisms of SARS-CoV-2 Evolution Revealing

Vaccine-Resistant Mutations in Europe and America. *J Phys Chem Lett* **12**, 11850-

11857 (2021).

7.   Liu, Y. *et al.* The N501Y spike substitution enhances SARS-CoV-2 infection and

transmission. *Nature* **602**, 294-299 (2022).

8.   Motozono, C. *et al.* SARS-CoV-2 spike L452R variant evades cellular immunity and

increases infectivity. *Cell Host Microbe* **29**, 1124-1136 e11 (2021).

9.   Cherian, S. *et al.* SARS-CoV-2 Spike Mutations, L452R, T478K, E484Q and P681R, in

the Second Wave of COVID-19 in Maharashtra, India. *Microorganisms* **9**(2021).

10.   Liu, Y. *et al.* Delta spike P681R mutation enhances SARS-CoV-2 fitness over Alpha

variant. *Cell Rep* **39**, 110829 (2022).

11.   Planas, D. *et al.* Reduced sensitivity of SARS-CoV-2 variant Delta to antibody

neutralization. *Nature* **596**, 276-280 (2021).

12.   Gu, H. *et al.* Adaptation of SARS-CoV-2 in BALB/c mice for testing vaccine efficacy.

Science **369**, 1603-1607 (2020).

13.   Wasik, B.R. *et al.* Influenza Viruses in Mice: Deep Sequencing Analysis of Serial

Passage and Effects of Sialic Acid Structural Variation. *J Virol* **93**(2019).

14.   Sun, S. *et al.* Characterization and structural basis of a lethal mouse-adapted SARS-

CoV-2. *Nat Commun* **12**, 5654 (2021).

15.   Huang, K. *et al.* Q493K and Q498H substitutions in Spike promote adaptation of SARS-

CoV-2 in mice. *EBioMedicine* **67**, 103381 (2021).

16. Chung, H., Noh, J.Y., Koo, B.S., Hong, J.J. & Kim, H.K. SARS-CoV-2 mutations
acquired during serial passage in human cell lines are consistent with several of those
found in recent natural SARS-CoV-2 variants. *Comput Struct Biotechnol J* **20**, 1925-
1934 (2022).

17. Sonnleitner, S.T. *et al.* The mutational dynamics of the SARS-CoV-2 virus in serial
passages in vitro. *Virol Sin* **37**, 198-207 (2022).

18. Tonkin-Hill, G. *et al.* Patterns of within-host genetic diversity in SARS-CoV-2. *Elife*
10(2021).

19. Turakhia, Y. *et al.* Pandemic-scale phylogenomics reveals the SARS-CoV-2
recombination landscape. *Nature* **609**, 994-997 (2022).

20. Kent, W.J. *et al.* The human genome browser at UCSC. *Genome Res* **12**, 996-1006
(2002).

21. Cao, Y. *et al.* BA.2.12.1, BA.4 and BA.5 escape antibodies elicited by Omicron infection.
Nature **608**, 593-602 (2022).

22. Johnson, B.A. *et al.* Loss of furin cleavage site attenuates SARS-CoV-2 pathogenesis.
Nature **591**, 293-299 (2021).

23. Tada, T. *et al.* Convalescent-Phase Sera and Vaccine-Elicited Antibodies Largely
Maintain Neutralizing Titer against Global SARS-CoV-2 Variant Spikes. *mBio* **12**,
e0069621 (2021).

24. Dupont, L. *et al.* Neutralizing antibody activity in convalescent sera from infection in
humans with SARS-CoV-2 and variants of concern. *Nat Microbiol* **6**, 1433-1442 (2021).

25. Tarres-Freixas, F. *et al.* Heterogeneous Infectivity and Pathogenesis of SARS-CoV-2
Variants Beta, Delta and Omicron in Transgenic K18-hACE2 and Wildtype Mice. *Front*
Microbiol **13**, 840757 (2022).

26. Rahimi, F. & Talebi Bezmin Abadi, A. Implications of the Emergence of a New Variant of
SARS-CoV-2, VUI-202012/01. *Arch Med Res* **52**, 569-571 (2021).

27.   Sedova, M., Jaroszewski, L., Iyer, M. & Godzik, A. Monitoring for SARS-CoV-2 drug

resistance mutations in broad viral populations. *bioRxiv* (2022).

28.   Kandeel, M. *et al.* Repurposing of FDA-approved antivirals, antibiotics, anthelmintics,

antioxidants, and cell protectives against SARS-CoV-2 papain-like protease. *J Biomol*

Struct Dyn **39**, 5129-5136 (2021).

29.   Hammond, J. *et al.* Oral Nirmatrelvir for High-Risk, Nonhospitalized Adults with Covid-

19. *N Engl J Med* **386**, 1397-1408 (2022).

30.   Pastorio, C. *et al.* Determinants of Spike infectivity, processing, and neutralization in

SARS-CoV-2 Omicron subvariants BA.1 and BA.2. *Cell Host Microbe* **30**, 1255-1268 e5

(2022).

31.   Zeng, C. *et al.* Neutralization of SARS-CoV-2 Variants of Concern Harboring Q677H.

mBio **12**, e0251021 (2021).

32.   Kim, K. *et al.* The roles of APOBEC-mediated RNA editing in SARS-CoV-2 mutations,

replication and fitness. *Sci Rep* **12**, 14972 (2022).

33.   Lei, C., Yang, J., Hu, J. & Sun, X. On the Calculation of TCID(50) for Quantitation of

Virus Infectivity. *Virol Sin* **36**, 141-144 (2021).

34.   Garrison, E.P.M.G. Haplotype-based variant detection from short-read sequencing. *arXiv*

(2012).

35.   N'Guessan A, T.A., Sanchez-Quete F, Goitom E, Reiling SJ, Galvez JH, Nguyen TL,

Nguyen HTL, Visentin F, Hachad M, Krylova K, Matthews S, Kraemer SA, Stretenowich

P, Bourgey M, Djambazian H, Chen S, Roy A, Brookes B, Lee S, Simon M, Maere T,

Vanrolleghem PA, Labelle M, Moreira S, Levade I, Bourque G, Ragoussis J, Dorner S,

Frigon D, Shapiro BJ. Detection of prevalent SARS-CoV-2 variant lineages in

wastewater and clinical sequences from cities in Québec, Canada. *medRxiv* (2022).

36.   Gerstung, M. *et al.* Reliable detection of subclonal single-nucleotide variants in tumour

cell populations. *Nat Commun* **3**, 811 (2012).

37.   Rao, R.S.P. *et al.* Evolutionary Dynamics of Indels in SARS-CoV-2 Spike Glycoprotein.

Evol Bioinform Online **17**, 11769343211064616 (2021).

38.   Ramachandran, V.K. *et al.* Comparison of variant callers for wastewater-based

epidemiology. *medRxiv* (2022).

39.   Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

arXiv (2013).

40.   McLaren, W. *et al.* The Ensembl Variant Effect Predictor. *Genome Biol* **17**, 122 (2016).