Nitrogen niche partitioning between tropical legumes and grasses strengthens under

elevated CO₂

Amber C. Churchill[1,2*], Haiyang Zhang[1,3], Gil Won Kim[1,4], Karen L. M. Catunda[1], Ian C.

Anderson[1], Forest Isbell[2], Ben Moore[1], Elise Pendall[1], Jonathan M. Plett[1], Jeff R. Powell[1],

Sally A. Power[1]

* Corresponding author: ac.churchill5@gmail.com; c: (1) 207-576-8591

[1]Hawkesbury Institute for the Environment, Western Sydney University, Hawkesbury

Campus, Ground Floor, Building R2, Locked Bag 1797, Penrith 2751, NSW Australia

[2]Department of Ecology, Evolution and Behavior, University of Minnesota, 140 Gortner

Laboratory, 1479 Gortner Ave., St. Paul, MN 55108, USA

[3]College of Life Sciences, Hebei University, Baoding, China

[4]Institute of Agriculture and Life Sciences, Gyeongsang National University, 501 Jinju

Daero, Jinju 52828, South Korea

Acknowledgements

The authors would like to acknowledge individuals whose efforts supported the data

collection of this work, including Kathryn Fuller, Alexandra Boyd, Shania Therese Didier

Serre, Samantha Weller, Benjamin Capel, Minh Doan. We thank Neha Mohan Babu for

comments on the final manuscript. Valuable input from PACE Advisory Board members -

Doug McNicholl (Meat and Livestock Australia), Cath Lescun (Dairy Australia), Catherine

Phelps (Dairy Australia), Tom Dickson (Heritage Seeds), Brendan Cullen (University

Melbourne) and Suzanne Boschman (Department of Primary Industries) - is also gratefully

acknowledged. This project was supported by funding from Meat and Livestock Australia's

Donor Company (P.PSH. 0793), Dairy Australia (C100002357) and Western Sydney

University. A version of this manuscript was included as a preprint on BioRxiv.

Conflict of Interest

The authors declare no conflicts of interest.

Author Contributions

Authors ACC, HZ, and GWK designed the experiment with input from SAP, EP, BM, JRP,

JMP, and KLMC. ACC, HZ, GWK, and KLMC performed experiments and conducted

glasshouse work. ACC conducted statistical analyses with input from HZ, JRP and FII; ACC

and HZ wrote the paper with input from all co-authors on draft iterations.

Data Availability

The data that support the findings of this study are openly available in Dryad at

https://doi.org/10.5061/dryad.nk98sf7ww.

Abstract

   1. Plant community biodiversity can be maintained, at least partially, by shifts in species interactions between facilitation and competition for resources as environmental conditions change. These interactions also drive ecosystem functioning, including productivity, and can promote over-yielding- an ecosystem service prioritized in working landscapes that occurs when there is either less competition, more facilitation, or both, between species in a community than within species. Importantly, shifts in species interactions that can result in over-yielding are unclear given rising CO_2 concentrations, especially in the context of tropical mixed-species grasslands.

   2. We examined the relative performance of two species pairs of tropical pasture grasses and legumes growing in monoculture and mixtures in a glasshouse experiment manipulating CO_2. We investigated how over-yielding can arise from nitrogen (N) niche partitioning and biotic facilitation using stable isotopes to differentiate soil N from biological N fixation (BNF) within N acquisition into aboveground biomass for these two-species mixtures.

   3. We found that N niche partitioning in species-level use of soil N vs. BNF drove species interactions in mixtures. Importantly partitioning, though not necessarily overyielding, was generally enhanced under elevated CO_2. However, this finding was mixture-dependent based on biomass of dominant species in mixtures and the strength of selection effects for the dominant mixture species.

   4. This study demonstrates that rising atmospheric CO_2 may alter niche partitioning between co-occurring species, with negative implications for the over-yielding benefits predicted for legume-grass mixtures in working landscapes with tropical species. Furthermore, these changes in inter-species interactions have consequences for shifts in grassland composition that are not yet considered in larger-scale projections for impacts of climate change and species distributions.

Graphical abstract (Image by H. Zhang): Among our tropical pasture species we found that grasses (dotted lines) grown in monoculture rely fully on soil nitrogen (N), while legumes (solid lines) grown in monoculture relied approximately equally on soil N and biological nitrogen fixation (BNF) to meet N requirements. When grown with tropical grasses, however, legumes shifted to rely more strongly on BNF, indicative of niche partitioning and decreased competition for soil nutrients with grasses. This separation of niche space was strengthened under elevated CO_2 conditions, ultimately reducing legume production.

Introduction

Interactions between plant species within communities and ecosystems vary over time and as environmental conditions change, thereby driving shifts between competition and facilitation for resources (Adler, Dalgleish, & Ellner, 2012; Arfin Khan, Grant, Beierkuhnlein, Kreyling, & Jentsch, 2014). Niche partitioning is a mechanism allowing the co-existence of numerous species that may overlap in fundamental niche space but differ in realized niche space to reduce impacts from competition (Jumpponen, Högberg, Huss-Danell, & Mulder, 2002; McKane et al., 2002; Zuppinger-Dingley et al., 2014). Additionally, changes in niche space associated with interspecific interactions are a common feature of net biodiversity effects whereby interactions among community members result in greater ecosystem functioning, relative to the sum of individual contributions (Barry et al., 2019; Loreau & Hector, 2001; Loreau et al., 2008); when applied to productivity this concept is often referred to as over-yielding (Vandermeer, 1981). Quantifying mechanisms underpinning interspecific interactions, such as niche partitioning, is fundamental to understanding relationships between biodiversity and ecosystem function.

Nitrogen (N) is a primary often-limiting nutrient that plays an important role in structuring community interactions and ecosystem function alongside phosphorus (P) and other mineral nutrients (Fay et al., 2015; Vitousek & Howarth, 1991). Plants that host N-fixing bacteria (i.e. N-fixers), such as legumes, are often considered to have an advantage under low soil N conditions as the mutualism can provide an alternative route for accessing N. This biological N fixation (BNF), therefore, can promote biodiversity and ecosystem function, via N niche partitioning or facilitation (Barry et al., 2019; Elias & Agrawal, 2021; Lee, Reich, & Tjoelker, 2003; Mulder, Jumpponen, Högberg, & Huss-Danell, 2002; Perring, Cullen, Johnson, & Hovenden, 2010).

In communities containing legumes, BNF is often assumed to result in N facilitation and an increase in the overall pool of ecosystem N available for neighbouring plants (Lee et al., 2003) that promote and subsequently increase production among comprising species (Barry et al., 2019; Fornara & Tilman, 2008; von Felton et al., 2009). There are two main pathways enabling this outcome: 1) direct acquisition of BNF resources from legumes by non-legumes (i.e. grasses) and 2) via niche partitioning where grasses compete less with legumes for shared soil nitrogen, given that legumes have access to a non-soil pool of N (Barry *et al.*, 2019). There are a number of biotic feedback-based mechanisms enabling the sharing of BNF

derived N to non-legume community members, including decomposition of legume leaf litter

(Makkonen et al., 2012; Wei, Reich, & Hobbie, 2019), or roots (Fustec, Lesuffleur, Mahieu,

& Cliquet, 2009; Mclaren & Turkington, 2010) that lead to higher soil N. Additionally, some

literature has highlighted the potential importance of root exudation of high N-containing

compounds that impact N (or carbon: C) availability in the rhizosphere (Henneron, Kardol,

Wardle, Cros, & Fontaine, 2020; Paynel, Murray, & Bernard Cliquet, 2001), or transfer

through the activity of soil fauna (Gilarte, Plett, Pendall, Carrillo, & Nielsen, 2020), thereby

promoting increased N availability for surrounding community members. Alternatively, niche

partitioning can support greater overall ecosystem N availability as legumes rely on BNF for

N resources, leaving the soil N pool available to neighbouring community members

(Ashworth, Toler, Allen, & Auge, 2018). These pathways offer numerous strategies that

ultimately promote increased community productivity in the presence of legumes

representing a potential for over-yielding.

Atmospheric CO_2 concentrations have increased by 50% since the industrial revolution

(IPCC, 2022). There are many ecosystem consequences associated with this change,

including an increase in plant productivity (de Graaff, van Groenigen, Six, Hungate, & van

Kessel, 2006; Gill et al., 2002), altered plant community composition and diversity (Reich,

2009; Zelikova et al., 2014), and modified plant interactions within a community that impact

ecosystem scale processes (Blumenthal et al., 2013; Gonçalves de Oliveira, Rios, Pereira, &

Souza, 2021; Maschler et al., 2022). However, some of these patterns may only occur when

water is limiting, nutrients are abundant (Reich, Hobbie, & Lee, 2014; Rogers, Ainsworth, &

Leakey, 2009; Soussana & Lüscher, 2007), or when plant diversity is high (Reich et al., 2001;

Shaw et al., 2002). Under well-watered conditions, eCO_2 generally benefits the C_3

photosynthetic pathway over the C_4 pathway (used by warm season or tropical grasses) due to

differences in carbon capture (Ainsworth & Long, 2005; Reich *et al.*, 2014, but see Reich *et*

al., 2018). Furthermore, eCO_2 is reported to increase belowground biomass, especially for

fine roots (Nie, Lu, Bell, Raut, & Pendall, 2013), which may increase access to soil nutrients

(de Graaff et al., 2006; Dieleman et al., 2012; Reich et al., 2001).

Many studies have found that eCO_2 fertilization effects are larger for N fixers than grasses

due to greater C resource availability to support the energetically demanding process of N-

fixation (Liang, Qi, Souza, & Luo, 2016). However, N-fixer responses in natural ecosystems

have been varied, as potential benefits from increased photosynthesis can become constrained

by other limiting nutrients such as phosphorus (Coventry, Hirth, Reeves, & Burnett, 1985; Hungate et al., 2004; Rogers et al., 2009; Soussana & Lüscher, 2007), or reduced due to temperature or drought conditions (Rogers et al., 2009). Even so, most studies find decreased foliar N concentrations for non-N-fixers under eCO_2 with subsequent increases C:N ratios in plant tissues (Ainsworth & Long, 2005; Augustine et al., 2018; Gojon, Cassan, Bach, Lejay, & Martin, 2022; Rogers et al., 2009). However, within mixed grass-legume grassland communities, leaf N and C:N ratios are less likely to be impacted by eCO_2 for the legume component of the sward (Rogers et al., 2009; Winkler & Herbst, 2004). These findings suggest that grassland responses to eCO_2 may depend on the presence of legumes (Jones & Donnelly, 2004). Importantly, the perceived 'benefit' of eCO_2 for tropical and subtropical ecosystems is less well documented, as most studies have focused on temperate species, with C_3 photosynthetic pathways (Naudts et al., 2013). The potential impacts of eCO_2 on nutrient dynamics in C_4 grasses, including those in mixed C_3-C_4 systems is of particular interest, due to the increasing abundance of warm season grasses in many areas of the world (McKeon et al., 2009). Given the diverse predicted outcomes for ecosystems as eCO_2 concentrations increase (Cannell & Thornley, 1998; de Graaff et al., 2006; Dijkstra et al., 2010), there is a need for improved understanding of the impacts of rising CO_2 in managed and natural grasslands. These include ongoing shifts in grassland composition, driven by species differences in response to climate change (McKeon et al., 2009; Munson & Long, 2017; Sherry et al., 2007; C. Wang, Hunt, Zhang, & Guo, 2013; Xie et al., 2022; Zelikova et al., 2014). Increased understanding of the processes that underlie plant-plant interactions between tropical grasses and legumes will therefore increase the accuracy of predictions for how grassland species composition and productivity will be impacted as atmospheric CO_2 concentrations continue to rise and human-induced shifts in ecosystem biodiversity are realized.

To address this area of research, we examined potential mechanisms by which eCO_2 might benefit warm season (C_4) grasses grown in mixed swards with tropical legumes. Specifically, we evaluated the potential for increased production and nutrient availability within two-species swards relative to individual species, driven by biologically fixed N (from rhizobia) supported by legumes and accessed by grasses. We hypothesized that 1) grasses grown with legumes benefit in terms of both nutrition and productivity, relative to those growing in monoculture, due to access to BNF from legume rhizobia; 2) tropical legumes grown under eCO_2 will increase production as a consequence of increased carbon-use-efficiency; 3)

increased C availability under eCO$_2$ will increase legume N fixation and both legume shoot N

and soil N content; and 4) increased available N due to BNF will promote overyielding and a

net biodiversity effect in two-species mixtures driven by a combination of complementarity

and selection effects that are enhanced by eCO$_2$.

Materials and Methods

Experimental design

This experiment was conducted in a glasshouse at Western Sydney University's Hawkesbury

Campus, Richmond, New South Wales (33°61'S,150°74'E), Australia. We used six separate

glasshouse chambers, with three randomly assigned to ambient conditions (410 - 430 ppm;

aCO$_2$) and three an elevated regime (630 - 650 ppm; eCO$_2$). The ambient treatment matches

contemporary atmospheric concentration, while the elevated regime was consistent with the

upper-end of predicted end-of-century increase in global atmospheric [CO$_2$] (Szulejko,

Kumar, Deep, & Kim, 2017). We maintained a consistent temperature regime across all

chambers at a daily average of 27°C (**Fig S1A**). This average included a temperature ramp to

mimic diurnal patterns (00-06 hours: 22°C, 06-09 hrs: 27°C, 09-18 hrs: 32°C, 18-21 hrs:

27°C, 21-00 hrs: 22°C) for spring to summer temperatures for areas currently cultivating

tropical pasture in northern NSW and southern Queensland (BOM, 2020). Additionally, all

chambers were controlled at 50-60% relative humidity (**Fig S1B**) with natural lighting

between 11 hours (August) and 14 hours (December).

The soil used in this experiment was collected adjacent to the Pastures and Climate Extremes

field experimental facility (Churchill et al., 2022). Soil was sieved to 5 mm before being air

dried and mixed with quartz sand (soil:sand 7:3 v/v). Each pot (3.7L, 150 mm diameter, 240

mm height) was filled with 3.9 kg of the soil-sand mixture before planting (Zhang et al.,

2021). The final soil mixture was ~ 84% sand, with an average pH of 5.6, plant available N at

55 mg kg^{-1}, plant available (bray) P at 39 mg kg^{-1} and soil organic carbon at 1.1%. All pots

were maintained under well-watered conditions using an automated irrigation system such

that soil water capacity was brought to 100% every two days (Zhang et al., 2021). In order to

promote legume nodulation, no nutrients were added to existing.

Two tropical grasses and two tropical legumes were selected, based on current widespread

use or consideration for future use by key pasture industries within Australia (**Table 1**). All

species are commonly grown in multi-species improved pastures in southern Queensland and

New South Wales, Australia, as well as among other sub-tropical grasslands world-wide

(Thomas & Sumberg, 1995; Bell *et al.*, 2012; Miegoue *et al.*, 2016; Melesse *et al.*, 2017;

DPI, 2018). Specific species-pairs were the legume *Macroptilium bracteatum* (Nees. &

Marti.) growing with the grass *Chloris gayana* (Kunth), and the legume *Desmodium intortum*

(Mill) Urb with grass *Panicum maximum* var. *trichoglume* (Robyns). All seeds were supplied

by Heritage Seeds and contained AgriCote advanced seed coating technology; legume seed

came pre-inoculated with appropriate rhizobia (Barenbrug Co., Heritage Seeds, New South

Wales, Australia). Seeds were planted directly into seedling trays (~5 seeds per cell) and

thinned to a single individual over time. After three weeks, two individuals for the

monoculture or one individual from each species for the mixture were then transplanted into

the pots. The experimental design therefore included the two CO_2 treatments (aCO_2 and

eCO_2), six types of plant combinations in pots ($grass_1$, $legume_1$, $mixed_1$, $grass_2$, $legume_2$,

$mixed_2$), and twelve replicates per plant combination per treatment (four replicates per

chamber).

Aboveground sampling

After transplanting the seedlings, all plants were grown for 11 weeks before a final harvest,

just as individuals were reaching the flowering phenophase. All aboveground standing plant

material (shoot; leaves and stems/tillers together) was cut at soil level and for mixed species

pots, separated by species, then weighed. The fresh aboveground biomass was frozen at -

18°C. Within one month all samples were then microwaved at medium power for 90 seconds

to halt enzymic degradation of plant tissue compounds, and oven dried at 70°C for 48 hours

then weighed (Landhäusser et al., 2018). All dried plant material was homogenized, ground

to pass through a 2-mm screen in a laboratory cyclone mill (Foss Cyclotec Mill, Denmark)

then ball-milled to a fine powder (Retsch® MM200; Hann, Germany) for nitrogen (N) and

isotopic (i.e. δ^{15}N) measurements, which were determined using an elemental analyzer

interfaced to a continuous flow isotope ratio mass spectrometer at the UC Davis Stable

Isotope Facility (Davis, CA, USA).

Belowground biomass

After aboveground material was removed, belowground biomass was determined. All pot

soils were carefully sieved using 2 mm sieves to retain root fragments and roots were oven

dried to constant mass. We assessed legume nodule number and weight for a subset of fresh

root material in each legume- or mixed species pot by randomly subsetting roots and nodules

from different locations within the root system. From this subset, we separated the nodules

from the roots and weighed the fresh biomass for each component separately. Fresh nodules

were incubated with acetylene for 1 hour and ethylene evolution tested as a proxy to estimate

the average instantaneous rate of nitrogenase activity of the bacteria within the nodules using

acetylene reduction assays (ARA; Plett *et al.*, 2016). After ARA measurement, root nodules

were oven dried and weighed as above. Final estimates of biological nitrogen fixation (BNF)

activity were calculated as the rate of ethylene production per gram of incubated nodule dry

biomass.

Soil nutrients

Nutrient availability was assessed using ion exchange resins for nitrate (NO_3^-), ammonia

(NH_4^+) and phosphate (PO_4^{-3}). We enclosed 5 g of mixed bed resin in a 3 cm x 5 cm nylon

bag (42 μm mesh) and placed one bag in each pot in the upper 10 cm. Resin bags were

collected during the biomass harvest, briefly rinsed to remove attached soil and root particles

before immediate storage at -70 °C prior to extraction with 0.5 M HCl solution. Extractable

NO_3^-, NH_4^+ and PO_4^{-3} concentrations in the resin bag were determined on a SEAL AQ2

Discrete Analyser (SEAL Analytical Inc., USA).

Quantifying contributions of biological nitrogen fixation

The contribution of BNF to plant total N was calculated as described in Chalk *et al.* (2016)

and Bell *et al.* (2017), where lower δ ^{15}N is indicative of a higher N-fixation amount. The

minimum observed values for δ^{15}N for both legume species in this experiment (*B*;

Macroptilium: -2.7 ‰; *Desmodium*: -2.0 ‰) were used to represent N derived solely from

BNF. Note that while this could overestimate the absolute estimate of BNF, the general

patterns among different treatments should remain unaffected (Chalk et al., 2016). Mean δ^{15}N

values for companion grass species in monoculture at each treatment level of CO_2 were used

as baselines for soil derived N (*Ref*; *Chloris*: 3.9 ‰, 3.7‰; *Panicum*: 4.9 ‰, 4.4 ‰; aCO_2,

eCO_2 respectively), and we then calculated the relative contribution of BNF vs. soil N to

plant aboveground N following Equation 1:

$$(Equation\ 1)\ BNF\ contribution\ to\ plant\ N = \frac{(Ref - Sample_{15_N})}{Ref - B}$$

Scale of inference	Scale at which factor of interest is applied	Number replicates at the appropriate scale

CO_2 treatment	Chamber	3 chambers for each CO_2 concentration
Legume-grass combinations	Monoculture and mixed pots	12 replicates per pot type; 4 in each chamber (legume, grass, mixed)
Species	Species	2 species included

Calculations and statistical analyses

Effects of CO_2 treatment on aboveground plant responses measured on pot level totals

(aboveground shoot biomass, total shoot N content, soil derived shoot N content,

belowground biomass, resin extractable soil nutrients, total shoot P content, and shoot N:P)

were analysed using linear mixed effects models. For these analyses, 'Species Pair', 'CO_2'

and 'Plant Type' treatments were included as fixed effects with all two-way and three-way

interactions; 'Chamber' was also included as a random effect in all models to account for

non-independence among replicates within each chamber due to potential between-chamber

environmental variation. Treatments of CO_2 included two levels (aCO_2 and eCO_2), 'Species

Pair' included '*Macroptilium & Chloris*' and '*Desmodium & Panicum*', while the 'Plant

Type' category included 'Grass', 'Legume', and 'Mixed' pots. Nodule data were also

analysed using linear mixed effects models, such that 'CO_2', 'Species Pair', and 'Plant Type'

were included as interactive fixed effects and 'Chamber' as a random effect. For these

analyses 'Plant Type' only included 'Legume' and 'Mixed' pots.

Over-yielding was determined as the increased productivity of a mixed pasture over the

discrete contributions of the constituent species grown in monoculture. We calculated over-

yielding for each species pair based on mixture total productivity (and total N content in

aboveground plant material) and the associated productivity of the monoculture grass and

legume species standardized by the number of individuals per pot, following Equation 2:

293
$$(Equation\ 2)\ Overyield\ effect\ of\ mixtures = \frac{Mixed}{\frac{Grass}{2} + \frac{Legume}{2}} - 1$$

Effects of eCO_2 on over-yielding were examined by calculating over-yield estimates based on

monoculture and mixed pots in each chamber (n = 6; 3 for each CO_2 level). We then ran a

linear mixed effects model with over-yield as the response, 'CO_2' level and 'Species Pair' as

the fixed effects and 'Chamber' as a random effect. Effect size of over-yielding was

determined by calculating mean treatment values and 95% confidence intervals (CI) across

all replicates (n = 12 for each plant type and CO_2 level) and over-yielding was deemed

significant if CIs didn't overlap with 0.

The net biodiversity effect, and its constituents- the complementarity effect and selection

effect- were calculated following Loreau & Hector (2001) based on averages for monoculture

and mixed pots of each species pair in each chamber and analysed statistically as described

for the over-yield metric above. Net biodiversity effect was therefore defined as the

difference between the observed yield and the expected yield in the absence of

complementarity or selection (Loreau & Hector, 2001). Complementarity effects are positive

when mixture yields are higher than expected based on the constituent monoculture species,

while selection effects are positive when species with greater than average yields dominate in

mixture.

All analyses were conducted in R version 4.1.1 (R Core Team, 2020) using the package lme4

(Bates, Maechler, Bolker, & Walker, 2015) with statistical contrasts based on Kenward-

Roger degrees of freedom calculated using the 'Anova' function from the 'car' package (Fox

& Weisberg, 2019). Pairwise comparisons to determine treatment differences in plant

responses for all models were conducted using the R package emmeans (Length, 2020) and

using the Tukey method for *P*-value adjustment.

Results

Shoot biomass and N content

Shoot biomass differed strongly between our species pairs, with greater productivity in pots

with the *Macroptilium* & *Chloris* monocultures and mixture, compared with *Desmodium* &

Panicum (**Fig 1**; **Table 2**). Elevated CO_2 (eCO_2) had limited effects on shoot biomass, with

increased productivity only observed for *Macroptilium* monoculture pots (**Fig 1a**). There

were no significant productivity differences among plant types (grass, legume, mixed) for the

Macroptilium & *Chloris* pairing, while *Desmodium* in monoculture produced less shoot

biomass than *Panicum* or the mixed plant type (**Fig 1b**). Despite little change in pot-level

biomass among monocultures and mixed pots, the contribution of grasses and legumes to

mixed pot total biomass was heavily weighted toward grass production in both species-pairs,

with *Chloris* contributing 65% and 61% of total pot biomass under aCO_2 and eCO_2 and

Panicum at 90% and 95%, respectively. Relative to growth by individual plants in

monoculture pots, this change represented a 38%/52% gain in biomass under aCO_2 and eCO_2

by *Chloris* and 20%/28% reduction in growth by *Macroptilium* (**Fig 1a**). This trend was

stronger for the *Desmodium & Panicum* species pair with 96%/87% gain for *Panicum* and

65%/86% loss for *Desmodium* between monoculture and mixed pots (**Fig 1b**).

Figure 1. Shoot biomass of grass and legume grown as monoculture or mixtures for two species pairs under ambient (aCO_2) and elevated CO_2 (eCO_2) comprising (a) *Macroptilium & Chloris* and (b) *Desmodium & Panicum*. Bars indicate mean values, plus standard error. Significant differences ($p < 0.05$) between CO_2 treatments are indicated by '*', and significant differences among plant types (grass, legume, mixed) are indicated by differing letter designations.

Figure 2. Shoot nitrogen (N) content (%N multiplied by shoot biomass) partitioned between grasses and legumes grown as monocultures or mixtures under ambient (aCO$_2$) and elevated CO$_2$ (eCO$_2$), for (a) *Macroptilium & Chloris* and (b) *Desmodium & Panicum*. Bars indicate mean values, plus standard error. Notations follow Fig. 1.

We found that legumes had a higher shoot %N than grasses, and this resulted in greater total

shoot N content for the legumes in both species pairs (**Fig 2**; **Table 2**). Additionally, the

Macroptilium & Chloris species pair generally had greater total shoot N content than

Desmodium & Panicum, driven by the differences in aboveground biomass rather than shoot

N concentrations. *Macroptilium* exhibited increased shoot N content under eCO$_2$ in

monoculture, however there was no change in N content in the mixed pots that included this

species (**Fig 2a**). In contrast, *Desmodium* N content was not significantly affected by eCO$_2$ in

monoculture, however pot total N content in the mixtures with *Panicum* was reduced under

eCO$_2$ (**Fig 2b**). Legume contributions to shoot N content in mixed pots also differed between

the two species pairs, with *Macroptilium* contributing 72%/67% under aCO$_2$ and eCO$_2$ and

Desmodium contributing 24%/18% of the total. Importantly, in both species pairs the shoot N

content per individual plant in grasses grown with legumes increased (*Chloris*: 51%/83%,

Panicum: 135%/84%; aCO_2/eCO_2, respectively), while the corresponding shoot N content per

individual in the mixed pot legumes declined compared with monoculture pots

(*Macroptilium*: -25%/-43%, *Desmodium*: -59%/-85%). For the *Desmodium* & *Panicum* pair

this change was large enough that the *Panicum* grown with *Desmodium* contained

comparable shoot N content to the *Desmodium* grown in monoculture.

N uptake source partitioning: soil N uptake

Despite variability in shoot N contents, the portion of shoot N that was derived from soil

nutrients (**Fig 3**) remained similar among plant types within each species pair (**Table 2**).

There were no effects of eCO_2 on soil-derived shoot N for either species pair, however the

fraction of soil-derived N in aboveground plant material in mixed pots was not equally

divided between grasses and legumes. Instead, grasses dominated soil nutrient uptake,

accounting for 74%/83% for aCO_2 and eCO_2 pots in the *Macroptilium* & *Chloris* species pair

and 94%/95% in *Desmodium* & *Panicum*. Raw values for $\delta^{15}N$ for each plant type and CO_2

treatment are included in **Table S1**. Resin extractable soil nutrients indicated that eCO_2 did

not impact availability (**Fig S2**) of NO_3^-, NH_4^+ or PO_4^{-3} ($p > 0.05$). We did, however, find

differences in the soil availability of NO_3^- and NH_4^+ among plant types within the

Macroptilium & *Chloris* species pair. *Chloris* monoculture pots had less NO_3^- than in mixed

pots (**Fig S2a**), and *Macroptilium* monoculture pots had less NH_4^+ than mixed pots (**Fig S2c**).

The ability of plants to take up soil nutrients is contingent on belowground traits, and we

found that there were key differences in the root biomass associated with eCO_2

(*Macroptilium* & *Chloris* **Fig S3a**; **Table S2**) and among plant types (legume/grass/mixed,

Desmodium & *Panicum*; **Fig S3b**). In general, there was increased root biomass in pots under

eCO_2, driven by significant increases in *Macroptilium* roots in monoculture and for mixed

roots in the *Macroptilium* & *Chloris* pots (**Fig S3a**). In the *Desmodium* & *Panicum* species

pair there was no impact of eCO_2, however *Panicum* had the greatest root biomass, followed

by the mixed pots and then *Desmodium* in monoculture (**Fig S3b**). Despite these changes in

belowground biomass, there were no shifts in the root mass fraction between species pairs,

among plant types or associated with eCO_2 (**Fig S3c** & **d**; **Table S2**).

N uptake source partitioning: biological nitrogen fixation

We found that a large fraction of the total shoot N was derived from BNF in legume
monoculture and mixed species pots, although this differed between legume species and
between plant types (**Fig 4**; **Table 3**). *Macroptilium* derived more shoot N via BNF than
Desmodium (*Macroptilium*: 77/82% of total shoot N in aCO_2 vs. eCO_2, *Desmodium*: 32/39%;
Table 3) in monoculture. Pot level shoot N relied more strongly on BNF in mixtures, with
Macroptilium & *Chloris* pots increasing reliance on BNF to 84/93% under aCO_2/eCO_2 and
Desmodium & *Panicum* shoot N increasing to 72/48%. Importantly, although both legumes
derived less shoot N from BNF by mass in mixture than monoculture (**Fig 4a** & **c**), BNF
contributed a greater percentage of the total shoot N in contrast to soil N (**Fig 4d**) for
Desmodium especially under aCO_2. Across both species pairs there were low levels of BNF-
derived N uptake by grasses (*Chloris*: 3/1%, *Panicum*: 6/1%). However, by mass, the BNF
derived N accounted for 29% of the shoot N in *Panicum* in mixtures under aCO_2, but only
2% under eCO_2 (**Fig 4b** & **d**) thereby providing evidence of direct facilitation under ambient
conditions within this species pair that was disrupted under eCO_2.
395

Fig 3. Shoot N content derived from soil for (a) *Macroptilium & Chloris* and (b) *Desmodium & Panicum* under ambient (aCO_2) and elevated (eCO_2) concentrations of CO_2. Values shown are pot-level means + 1 SE. Note no significant differences with CO_2 treatment or among plant types were observed.

Fig. 4. Shoot N content derived from biological N fixation (BNF; a & c) and the proportion of aboveground N content derived from BNF (b & d) for *Macroptilium* & *Chloris* (a & b) and *Desmodium* & *Panicum* (c & d). Values shown are means ± 1 SE. Colours in panels match Fig 1, with green indicating N in legume shoots, and orange indicating N in grass shoots. Notations from Fig. 1.

The ability of legumes to support rhizobia in nodules is the main constraint on the potential

role of BNF as a source for shoot N, and we found clear differences between species in terms

of root nodules (**Fig 5**; **Table 4**). In parallel with total shoot N content, *Macroptilium* had

greater nodule biomass (**Fig 5a**) and number of nodules (**Fig 5b**), compared to *Desmodium,*

in monoculture. Additionally, both legumes had a significant reduction in nodule biomass and

number of nodules when grown with grass, and there were no impacts of eCO_2 on these

measurements. Despite the difference in nodule biomass and number between legume

species, *Desmodium* had equivalent levels of instantaneous nodule activity to *Macroptilium*,

based on production of ethylene as a proxy for the rate of nitrogen fixation following harvest

(**Fig 5c**) and in monoculture there was no impact of eCO_2 for either species. However, for

Desmodium grown in mixture with *Panicum*, real-time nodule activity significantly increased

for individuals grown under eCO_2 (**Fig 5c**).

Fig 5. Effects of eCO_2 on metrics of nitrogen fixation capacity relating to root nodules for legumes grown in monoculture (Legume) and in mixtures with grasses (Mixed; *Macroptilium* and *Chloris*: Mac & Chl, *Desmodium* and *Panicum:* Des & Pan) including (a) the total biomass of nodules per pot, (b) the number of nodules per pot, and (c) root nodule activity as measured by the production of ethylene as a proxy for the rate of nitrogen fixation. Values shown are means ± 1 SE. Differences between plant types within

a species pair are indicated by letters (Mac & Chl uppercase, Des & Pan lowercase) and significant differences between CO_2 treatments are indicated by '*'.

Over-yielding and Complementarity vs. Selection Effects

We found a non-significant positive response of over-yielding for both species' pairs (**Fig.**

6a). In *Macroptilium & Chloris* mixture, over-yielding was mainly determined by the

complementarity effect but not the selection effect (**Fig. 6bc**). By contrast, for the

Desmodium & Panicum mixture, the over-yielding was driven more by selection compared

with the complementarity effect (**Fig. 6bc**).

We also found eCO_2 significantly reduced over-yielding (**Fig. 6a**). In the *Macroptilium &*

Chloris mixture, complementarity effects shifted from neutral under aCO_2 to significantly

positive under eCO_2. Meanwhile, selection effects shifted from neutral under aCO_2 to

significantly negative under eCO_2, meaning that the dominant species had less biomass than

expected in the mixture (**Fig. 6c**). In the *Desmodium & Panicum* mixture, however, the

complementarity effect shifted from significantly positive under aCO_2 to neutral under eCO_2,

meaning that niche partitioning was weakened. At the same time, the selection effect was

unaffected by CO_2 (**Fig. 6c**), although values were significantly positive under both CO_2

scenarios, meaning that the dominant species (*Panicum*) consistently had greater biomass in

mixtures than proportional production would suggest from monocultures, regardless of CO_2

treatment. There were no significant effects of eCO_2 among the two species pairs for either

complementarity (CO_2: $F_{1,4} = 0.3$, $p = 0.62$; Species pair: $F_{1,44} = 1.9$, $p = 0.18$; CO_2 x Species

pair: $F_{1,44} = 2.1$, $p = 0.15$) or selection effects (CO_2: $F_{1,4} = 1.0$, $p = 0.37$; Species pair: $F_{1,44} =$

94.9, $p < 0.01$; CO_2 x Species pair: $F_{1,44} = 2.6$, $p = 0.11$). Summing these components of the

net biodiversity effect supports the general outcomes of biomass over-yielding only in the

Desmodium & Panicum mixture, with reduced over-yielding under eCO_2 (**Fig. 6d**). There

were, however, no significant effects of CO_2 between the species pairs (CO_2: $F_{1,4} = 0.7$, $p =$

0.44; Species pair: $F_{1,44} = 3.0$, $p = 0.09$; CO_2 x Species pair: $F_{1,44} = 0.8$, $p = 0.37$).

Fig. 6. Over-yield effect size for (a) aboveground productivity for pots grown with mixed grass-legume plant types in comparison with standardized summed totals for grasses and legumes grown in monoculture. Standardized species-pair level calculations for examining (b) complementarity effect, (c) selection effect and (d) the net biodiversity effect. Positive values indicate an increase relative to monoculture pots (a) or a net positive effect in interaction type between species (b-d). Non-overlapping bars indicate a significant difference from 0. Points shown are means values with 95% CI for *Macroptilium & Chloris* (Mac & Chl) mixtures and *Desmodium & Panicum* (Des & Pan) mixtures.

Discussion

In mixed grasslands or pastures containing both grasses and legumes nutrient niche partitioning among community members can promote species coexistence and ecosystem function, particularly by influencing nutrient availability (Craven et al., 2018; da Silveira Pontes, Maire, Schellberg, & Louault, 2015; Fornara & Tilman, 2009). While direct-nutrient sharing is often predicted between community members (Pirhofer-Walzl et al., 2012), nutrient, especially N, niche partitioning is a more common finding in ecological and agricultural studies under field conditions (Nyfeler, Huguenin-Elie, Suter, Frossard, & Lüscher, 2011; Pelzer, Hombert, Jeuffroy, & Makowski, 2014; Zuppinger-Dingley et al., 2014). In these situations, community members access different pools of available resources, over time and/or space, enabling the persistence of many species by reducing competitive interactions (Adler et al., 2012; Zuppinger-Dingley et al., 2014). Our results found that (1) within tropical grass-legume mixtures, N niche partitioning drove plant-plant interactions such that legumes increased reliance on biological nitrogen fixation (BNF) relative to soil N when grown with grasses; (2) elevated CO_2 (eCO_2) had varied impact on individual tropical forage species, with benefits in growth and total N content for one legume, but reductions in pot-level total N content for mixed pots containing a second legume species- and no direct benefit for tropical grasses; (3) community productivity and N content from tropical legume-grass mixtures were lower under eCO_2 than aCO_2, suggesting that legume-grass pairings may benefit ecosystem function to a lesser extent in the face of rising CO_2; and (4) the species-specific responses associated with over-yielding and responses to eCO_2, were linked with differences in complementarity and selection effects operating between the two mixed communities.

Tropical grass-legume mixtures

Relationships between plant functional group diversity and ecosystem function typically predict and find that increased diversity supports greater ecosystem productivity (Hector et al., 1999; Küchenmeister, Küchenmeister, Wrage, Kayser, & Isselstein, 2012; Loreau & Hector, 2001; Mason, Palmer, Romera, Waugh, & Mudge, 2017; Tilman, Isbell, & Cowles, 2014). In testing this hypothesis, many studies have included grass-forb-legume or grass-shrub-forb combinations experimentally to examine potential over-yielding benefits (Finn et al., 2013), but fewer ecological studies have included C_4 grasses alongside legumes (Lee et al., 2003; Julianne M/ Lilley, Bolger, & Gifford, 2001; Rasmussen, Gylfadóttir, Loges, Eriksen, & Helgadóttir, 2013). This is despite early results suggesting that the high

competitive ability of some C_4 grasses may functionally contribute to over-yielding
(Lambers, Harpole, Tilman, Knops, & Reich, 2004) and the economic importance of such
mixtures in many agricultural grasslands around the world (Butler, Interrante, & Foster,
2013; Duchene, Vian, & Celette, 2017). Previous results have shown that, at a global scale
tropical grasses show lower productivity benefits from the addition of legumes, relative to
temperate mixed-species systems (Ashworth et al., 2018). Therefore, increased understanding
of the mechanisms that may promote over-yielding are of broad interest. Our study found
strong evidence for productivity gains by the grass component in tropical grass-legume
mixtures, accompanied by declines in the legume partner, which translated into minimal
changes at the pot or sward level. This result aligns with earlier tropical mixed pasture trials
that concluded that the relative abundance of grasses and legumes has an important impact on
the transfer of N and sward productivity benefits of mixtures, compared to monocultures of
the component species (Baba, Halim, Alimon, & Abubakar, 2011).
485
Increased productivity in mixed species grasslands, relative to monocultures, is primarily
attributed to increased access to resources, including light, water or nutrient availability in
ways that promote facilitation. This facilitation can include mechanisms of niche partitioning
such that different species access different resource pools in time or space to minimize
competition (da Silveira Pontes et al., 2015; Jumpponen et al., 2002; Thilakarathna, McElroy,
Chapagain, Papadopoulos, & Raizada, 2016). Biological or environmental conditions under
which different mechanisms promoting nutrient facilitation may dominate are still an area of
active research (Brooker et al., 2015), and our study addressed the potential for niche
partitioning vs. direct facilitation for N between co-occurring grasses and legumes (Ajayi,
Babayemi, & Taiwo, 2008; Baba et al., 2011; L. W. Bell et al., 2017; Miegoue et al., 2016).
Despite only limited gains in shoot N for mixed pots, relative to the grass monocultures, this
indicated niche partitioning as the main mechanism promoting plant N access between the
component species. Under these conditions, the proportion of shoot N derived from BNF
increased for legumes even as the total mass of shoot N declined in comparison with
monoculture pots. In agricultural-grassland settings mixtures of grasses and legumes are often
used to specifically improve forage quality and production in the absence of fertilizer
applications (Brooker, Karley, Newton, Pakeman, & Schöb, 2016; Li et al., 2015; Warwick et
al., 2016). This practice is predicted to play an important role in grassland management for
grazed systems as C_4 grasses become more dominant under changing climate (Butler et al.,
2013; McKeon et al., 2009; Still, Cotton, & Griffith, 2019).

There are a variety of mechanisms that enable N niche complementarity in mixed-species

grasslands, and it is worth noting that the duration of species associations may play a role in

the potential for direct-facilitation of BNF-derived N between for companion grasses in

mixed grasslands (Heichel & Henjum, 1991; Thilakarathna et al., 2016). Such a time lag may

be associated with the decomposition of legume leaf and root litter that can provide sources

for increased ecosystem N availability on longer time scales (Cannell & Thornley, 1998;

Kohmann, Sollenberger, Dubeux, Silveira, & Moreno, 2019). While there was likely root

turnover over the course of our glasshouse experiment, and any senesced leaves were retained

in pots, these mechanisms are likely more substantial in field settings especially where soil

disturbance may promote incorporation of plant materials directly into soils or where legume

residues have sufficient time to undergo decomposition (Brooker et al., 2015; Thilakarathna

et al., 2016). Furthermore, under field conditions, legumes often shift to BNF as plants

mature (Edmeades & Goh, 1984; Wery, Turc, & Salsac, 1986) and this can impact the

potential facilitation provided to companion species (L. W. Bell et al., 2017). In our study all

plants were harvested immediately prior to the first flowering phenophase, and consequently

observed patterns are from first-year perennial plants in a rapid growth phase when

individuals may rely more strongly on available soil N pools. Grasses in mixed pots in our

study did show some evidence for incorporation of BNF-derived N into aboveground plant

tissue (up to 29% of shoot N for *Panicum* grown with *Desmodium*), a contribution that could

increase as swards mature (Thilakarathna et al., 2016).

Consequences of elevated CO$_2$

Previous work has shown that eCO$_2$ can increase grassland and crop productivity (Shaw et

al., 2002; Terrer et al., 2021), or have no impact over differing time scales, and can result in

shifts in the abundance of component species (Bloor, Pichon, Falcimagne, Leadley, &

Soussana, 2010; Carroll, Caporn, Johnson, Morecroft, & Lee, 2003; Mueller et al., 2016).

Under well-watered conditions, C$_4$ grasses are not predicted to benefit directly from eCO$_2$

and our results follow those predictions (Ainsworth & Long, 2005; Soussana & Lüscher,

2007; Wang *et al.*, 2022; but see Reich *et al.*, 2018). On the other hand, we found some

limited effects of eCO$_2$ on legume production, with increased growth in *Macroptilium* likely

due to increases in shoot N and root biomass. In contrast, *Desmodium* did not respond to

eCO$_2$, potentially due to a greater reliance on soil N rather than BNF in monoculture, that

may have induced stoichiometric constraints in responding to eCO$_2$. This result is in line with

research showing that the strongest predictor for an individual's or species' positive response

to eCO_2 in legumes is the ability to form nodules (Cernusak et al., 2011; Parvin, Uddin,

Tausz-Posch, Armstrong, & Tausz, 2020). For example, *Macroptilium* responded positively

to eCO_2 and also had greater nodule production than *Desmodium*, although the number of

nodules was unaffected by eCO_2. Indeed, despite the positive effect of eCO_2 on *Desmodium*

nodule activity in mixtures, low nodule numbers and biomass overall resulted in a

proportional reduction in legume biomass within the mixed sward.

At the sward level, despite some gain in productivity for legume biomass and total shoot N in

monoculture (*Macroptilium* only), we didn't find an increase in productivity for mixed pots

under eCO_2. While this result is in contrast to C_3 grass-legume mixtures for temperate

pastures in Australia where eCO_2 has been shown to increase total shoot N yield (J. M.

Lilley, Bolger, Peoples, & Gifford, 2001), it is consistent with evidence for limited responses

to eCO_4 by C_4 grasses and legumes. One explanation may be related to shifts in the nutritional

quality of component species under eCO_2, as reductions in quality for aboveground biomass

have been reported across ecosystems (Augustine et al., 2018; Bhargava & Mitra, 2021). Our

species experienced limited shifts in quality based on shoot N%, including lower N in

Macroptilium when grown with *Chloris* and in *Panicum* when grown with *Desmodium*. Even

so, these patterns align with other studies measuring nutritional shifts in both tropical grasses

and legumes under low soil nutrients where legume shoot N% declined in mixture under

eCO_2 (Edwards, McCaffery, & Evans, 2006).

While the use of confined soil space to test mechanisms for the source of nutrient use and

acquisition between grasses and legumes is typical in glasshouse settings, there are some key

factors that are likely to impact the species interactions and dynamics in response to eCO_2

that are not captured here that are important for overarching messages in relation to nutrient

facilitation. Our study found that eCO_2 conditions differentially altered the shoot N%

between legumes in mixture but had no impact on legumes grown in monoculture. Under

field settings, the effects of herbivory on legumes that maintain a higher nutritive quality

relative to surrounding species may result in a comparative loss in abundance or persistence

within the community, ultimately reducing the potential over-yield or net biodiversity effect

(Rogers et al., 2009). Additionally, literature reviews on field studies have generally found an

increase in root length and biomass associated with eCO_2 in grasslands (Dieleman et al.,

2012; Nie et al., 2013), a pattern that was seen in pots containing *Macroptilium* in this study.

While our pots were not root-bound by the end of the experiment, other key differences

between field and glasshouse conditions may have limited further responses, for example Nie

et al. (2013) also found a shift in the depth distribution of roots that is not possible under non-

field settings. Such spatial re-distribution of roots with depth can play a major role in

complementarity effects between species within mixtures (Oram et al., 2018).

Over-yield implications under future CO_2 scenarios

Reliance on different sources of N among species within mixed plant communities can

ultimately reduce competition especially under low nutrient availability (Ball et al., 2021;

Elias & Agrawal, 2021). The increased N niche partitioning between grasses and legumes

grown in mixture found in this study support this conclusion. Furthermore, the species pair

that had a substantial shift in N source use between monoculture and mixtures was also

associated with a significant over-yielding benefit. This suggests that greater plasticity in

niche space may promote species coexistence and enhance ecosystem production.

Belowground, the idea of nutrient form (nitrate vs. ammonium) niche plasticity supporting

coexistence has been well-established for cold-climate grasslands (Ashton, Miller, Bowman,

& Suding, 2010) and aboveground light-use plasticity has been shown to translate into

increased production in experimental grasslands (Meilhac, Deschamps, Maire, Flajoulot, &

Litrico, 2020).

Importantly, however, we found that eCO_2 reduced over-yielding for both aboveground

biomass and total shoot N in our tropical grass-legume mixtures. While there were no

statistical effects of eCO_2 on complementarity and selection effects directly, we found key

differences in the relative importance of these effects in contributing to the over-yield/net

biodiversity effect, with positive complementarity and selection effects together resulting in

increased sward level over-yielding that was greatest under aCO_2.

Conclusions

Plant responses to eCO_2 are largely dependent on soil water and nutrient availability.

Therefore, shifts in plant-plant interactions including competition (evidenced by niche

partitioning) or facilitation among community members may drive diverse ecosystem

responses to changes in CO_2 concentrations. Among our tropical pasture species, we found

that legumes grown in monoculture relied approximately equally on available soil N pools

and BNF for their N requirements. When grown with tropical grasses, however, legumes

shifted to rely more strongly on BNF, indicative of stronger niche partitioning and less

successful competition for soil nutrients with grasses. This separation of niche space was

strengthened under elevated CO_2 conditions, ultimately reducing legume production and

minimizing over-yielding benefits. Disentangling contributions of niche partitioning and

abiotic facilitation, here quantified using stable isotopes, are key in understanding grass-

legume contributions to ecosystem function under future climate conditions.

References

Adler, P. B., Dalgleish, H. J., & Ellner, S. P. (2012). Forecasting plant community impacts of climate variability and change: when do competitive interactions matter? *Journal of Ecology*, *100*(2), 478–487. doi: 10.1111/j.1365-2745.2011.01930.x

Ainsworth, E. A., & Long, S. P. (2005). What have we learned from 15 years of free-air CO_2 enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO2. *New Phytologist*, *165*(2), 351–372. doi: 10.1111/j.1469-8137.2004.01224.x

Ajayi, F. T., Babayemi, O. J., & Taiwo, A. A. (2008). Effects of supplementation of *Panicum maximum* with four herbaceous forage legumes on performance, nutrient digestibility and nitrogen balance in West African dwarf goats. *Animal Science Journal*, *79*(6), 673–679. doi: 10.1111/j.1740-0929.2008.00579.x

Arfin Khan, M. A. S., Grant, K., Beierkuhnlein, C., Kreyling, J., & Jentsch, A. (2014). Climatic extremes lead to species-specific legume facilitation in an experimental temperate grassland. *Plant and Soil*, *379*(1–2), 161–175. doi: 10.1007/s11104-014-2050-8

Ashton, I. W., Miller, A. E., Bowman, W. D., & Suding, K. N. (2010). Niche complementarity due to plasticity in resource use: plant partitioning of chemical N forms. *Ecology*, *91*(11), 3252–3260. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/21141186

Ashworth, A. J., Toler, H. D., Allen, F. L., & Auge, R. M. (2018). Global meta-analysis reveals agro-grassland productivity varies based on species diversity over time. *PLoS ONE*, *13*(7), 1–20. doi: 10.1371/journal.pone.0200274

Augustine, D. J., Blumenthal, D. M., Springer, T. L., LeCain, D. R., Gunter, S. A., & Derner, J. D. (2018). Elevated CO_2 induces substantial and persistent declines in forage quality irrespective of warming in mixedgrass prairie. *Ecological Applications*, *28*(3), 721–735. doi: 10.1002/eap.1680

Baba, M., Halim, R. A., Alimon, A. R., & Abubakar, I. (2011). Grass-legume mixtures for enhanced forage production: Analysis of dry matter yield and competition indices. *African Journal of Agricultural Research*, *6*(23), 5242–5250. doi: 10.5897/AJAR11.615

Ball, K. R., Woodin, S. J., Power, S. A., Brien, C., Berger, B., Smith, P., & Pendall, E. (2021). Integrating ecological stoichiometry to understand nutrient limitation and potential for competition in mixed pasture assemblages. *Journal of Soil Science and Plant Nutrition*, *21*, 24989–2500.

Barry, K. E., Mommer, L., Ruijven, J. Van, Wirth, C., Wright, A. J., Bai, Y., … Scherer-lorenzen, M. (2019). The future of complementarity: disentangling causes from consequences. *Trends in Ecology & Evolution*, *34*(2), 167–180.

Bates, D., Maechler, M., Bolker, B. M., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, *67*(1), 1–48. Retrieved from http://arxiv.org/abs/1406.5823.

Bell, L. W., Lawrence, J., Johnson, B., & Peoples, M. B. (2017). New ley legumes increase nitrogen fixation and availability and grain crop yields in subtropical cropping systems. *Crop and Pasture Science*, *68*(1), 11–26. doi: 10.1071/CP16248

Bell, M., Harch, G., & Moody, P. (2012). Diversification from cropping into mixed crop-livestock systems–the sustainability risks posed by hay removal from pasture or forage blocks. *Proceedings of 16th Agronomy Conference 2012*, *18*(11), 75.

Bhargava, S., & Mitra, S. (2021). Elevated atmospheric CO_2 and the future of crop plants. *Plant Breeding*, *140*(1), 1–11. doi: 10.1111/pbr.12871

Bloor, J. M. G., Pichon, P., Falcimagne, R., Leadley, P., & Soussana, J.-F. (2010). Effects of warming, summer drought, and CO_2 enrichment on aboveground biomass production,

flowering phenology, and community structure in an upland grassland ecosystem. *Ecosystems*, *13*(6), 888–900. doi: 10.1007/s10021-010-9363-0

Blumenthal, D. M., Resco, V., Morgan, J. A., Williams, D. G., Lecain, D. R., Hardy, E. M., … Bladyka, E. (2013). Invasive forb benefits from water savings by native plants and carbon fertilization under elevated CO_2 and warming. *New Phytologist*, *200*(4), 1156–1165. doi: 10.1111/nph.12459

Brooker, R. W., Bennett, A. E., Cong, W. F., Daniell, T. J., George, T. S., Hallett, P. D., … White, P. J. (2015). Improving intercropping: A synthesis of research in agronomy, plant physiology and ecology. *New Phytologist*, *206*(1), 107–117. doi: 10.1111/nph.13132

Brooker, R. W., Karley, A. J., Newton, A. C., Pakeman, R. J., & Schöb, C. (2016). Facilitation and sustainable agriculture: A mechanistic approach to reconciling crop production and conservation. *Functional Ecology*, *30*(1), 98–107. doi: 10.1111/1365-2435.12496

Butler, T. J., Interrante, S. M., & Foster, J. L. (2013). Assessing the production and nutritive value of warm-season legumes in Oklahoma and Texas. *Forage & Grazinglands*, *11*(1), 1–6. doi: 10.2134/fg-2013-0026-rs

Cannell, M. G. R., & Thornley, J. H. M. (1998). N-poor ecosystems may respond more to elevated [CO_2] than N-rich ones in the long term. A model analysis of grassland. *Global Change Biology*, *4*(4), 431–442. doi: 10.1046/j.1365-2486.1998.00167.x

Carroll, J. A., Caporn, S. J. M., Johnson, D., Morecroft, M. D., & Lee, J. A. (2003). The interactions between plant growth, vegetation structure and soil processes in semi-natural acidic and calcareous grasslands receiving long-term inputs of simulated pollutant nitrogen deposition. *Environmental Pollution*, *121*, 363–376. doi: 10.1007/BF00141010

Cernusak, L. A., Winter, K., Martínez, C., Correa, E., Aranda, J., Garcia, M., … Turner, B. L. (2011). Responses of legume versus nonlegume tropical tree seedlings to elevated CO_2 concentration. *Plant Physiology*, *157*(1), 372–385. doi: 10.1104/pp.111.182436

Chalk, P. M., Inácio, C. T., Balieiro, F. C., & Rouws, J. R. C. (2016). Do techniques based on [15]N enrichment and [15]N natural abundance give consistent estimates of the symbiotic dependence of N_2-fixing plants? *Plant and Soil*, *399*(1–2), 415–426. doi: 10.1007/s11104-015-2689-9

Churchill, A. C., Zhang, H., Fuller, K. J., Amiji, B., Anderson, I. C., Barton, C. V. M., … Power, S. A. (2022). Pastures and climate extremes: impacts of cool season warming and drought on the productivity of key pasture species in a field experiment. *Frontiers in Plant Science*, *13*(March), 1–16. doi: 10.3389/fpls.2022.836968

Coventry, D. R., Hirth, J. R., Reeves, T. G., & Burnett, V. F. (1985). Growth and nitrogen fixation by subterranean clover in response to inoculation, molybdenum application and soil amendment with lime. *Soil Biology and Biochemistry*, *17*(6), 791–796. doi: 10.1016/0038-0717(85)90134-8

Craven, D., Eisenhauer, N., Pearse, W. D., Hautier, Y., Isbell, F. I., Roscher, C., … Manning, P. (2018). Multiple facets of biodiversity drive the diversity–stability relationship. *Nature Ecology and Evolution*, *2*(10), 1579–1587. doi: 10.1038/s41559-018-0647-7

da Silveira Pontes, L., Maire, V., Schellberg, J., & Louault, F. (2015). Grass strategies and grassland community responses to environmental drivers: a review. *Agronomy for Sustainable Development*, *35*(4), 1297–1318. doi: 10.1007/s13593-015-0314-1

de Graaff, M. A., van Groenigen, K. J., Six, J., Hungate, B., & van Kessel, C. (2006). Interactions between plant growth and soil nutrient cycling under elevated CO_2: A meta-analysis. *Global Change Biology*, *12*(11), 2077–2091. doi: 10.1111/j.1365-2486.2006.01240.x

Dieleman, W. I. J., Vicca, S., Dijkstra, F. A., Hagedorn, F., Hovenden, M. J., Larsen, K.

S., … Janssens, I. A. (2012). Simple additive effects are rare: a quantitative review of plant biomass and soil process responses to combined manipulations of CO_2 and temperature. *Global Change Biology*, *18*(9), 2681–2693. doi: 10.1111/j.1365-2486.2012.02745.x

Dijkstra, F. A., Blumenthal, D., Morgan, J. A., Pendall, E., Carrillo, Y., & Follett, R. F. (2010). Contrasting effects of elevated CO_2 and warming on nitrogen cycling in a semiarid grassland. *New Phytologist*, *187*(2), 426–437. doi: 10.1111/j.1469-8137.2010.03293.x

Duchene, O., Vian, J. F., & Celette, F. (2017). Intercropping with legume for agroecological cropping systems: Complementarity and facilitation processes and the importance of soil microorganisms. A review. *Agriculture, Ecosystems and Environment*, *240*, 148–161. doi: 10.1016/j.agee.2017.02.019

Edmeades, D. C., & Goh, K. M. (1984). Symbiotic nitrogen fixation in a sequence of pastures of increasing age measured by a ^{15}N dilution technique. *New Zealand Journal of Agricultural Research*, *21*(4), 623–628. doi: 10.1080/00288233.1978.10427458

Edwards, E. J., McCaffery, S., & Evans, J. R. (2006). Phosphorus availability and elevated CO_2 affect biological nitrogen fixation and nutrient fluxes in a clover-dominated sward. *New Phytologist*, *169*(1), 157–167. doi: 10.1111/j.1469-8137.2005.01568.x

Elias, J. D., & Agrawal, A. A. (2021). A private channel of nitrogen alleviates interspecific competition for an annual legume. *Ecology*, *102*(9), 1–10. doi: 10.1002/ecy.3449

Fay, P. A., Prober, S. M., Harpole, W. S., Knops, J. M. H., Bakker, J. D., Borer, E. T., … Yang, L. H. (2015). Grassland productivity limited by multiple nutrients. *Nature Plants*, *1*(July), 1–5. doi: 10.1038/nplants.2015.80

Finn, J. A., Kirwan, L., Connolly, J., Sebastià, M. T., Helgadottir, A., Baadshaug, O. H., … Lüscher, A. (2013). Ecosystem function enhanced by combining four functional types of plant species in intensively managed grassland mixtures: A 3-year continental-scale field experiment. *Journal of Applied Ecology*, *50*(2), 365–375. doi: 10.1111/1365-2664.12041

Fornara, D. A., & Tilman, D. G. (2008). Plant functional composition influences rates of soil carbon and nitrogen accumulation. *Journal of Ecology*, *96*(2), 314–322. doi: 10.1111/j.1365-2745.2007.01345.x

Fornara, D. A., & Tilman, D. G. (2009). Ecological mechanisms associated with the positive diversity-productivity relationship in an N-limited grassland. *Ecology*, *90*(2), 408–418. doi: 10.1890/08-0325.1

Fox, J., & Weisberg, S. (2019). *An R companion to applied regression* (No. Third Edition; Third). Thousand Oaks, CA: Sage.

Fustec, J., Lesuffleur, F., Mahieu, S., & Cliquet, J. B. (2009). Nitrogen rhizodeposition of legumes. *Sustainable Agriculture*, *2*, 869–881. doi: 10.1007/978-94-007-0394-0_38

Gilarte, P., Plett, J., Pendall, E., Carrillo, Y., & Nielsen, U. N. (2020). Direct and indirect trophic interactions of soil nematodes impact chickpea and oat nutrition. *Plant and Soil*, *457*(1–2), 255–268. doi: 10.1007/s11104-020-04735-6

Gill, R. A., Polley, H. W., Johnson, H. B., Anderson, L. J., Maherali, H., & Jackson, R. B. (2002). Nonlinear grassland responses to past and future atmospheric CO_2. *Nature*, *417*(6886), 279–282. doi: 10.1038/417279a

Gojon, A., Cassan, O., Bach, L., Lejay, L., & Martin, A. (2022). The decline of plant mineral nutrition under rising CO_2: physiological and molecular aspects of a bad deal. *Trends in Plant Science*, 1–14. doi: 10.1016/j.tplants.2022.09.002

Gonçalves de Oliveira, A. C., Rios, P. M., Pereira, E. G., & Souza, J. P. (2021). Growth and competition between a native leguminous forb and an alien grass from the Cerrado under elevated CO_2. *Austral Ecology*, *46*(5), 750–761. doi: 10.1111/aec.13006

Hector, A., Schmid, B., Beierkuhnlein, C., Caldeira, M. C., Diemer, M., Dimitrakopoulos, P.

G., … Lawton, J. H. (1999). Plant diversity and productivity experiments in European grasslands. *Science*, *286*(5442), 1123–1127. doi: 10.1126/science.286.5442.1123

Heichel, G. H., & Henjum, K. I. (1991). Dinitrogen fixation, nitrogen transfer, and productivity of forage legume-grass communities. *Crop Science*, *31*(1), 202–208. doi: 10.2135/cropsci1991.0011183x003100010045x

Henneron, L., Kardol, P., Wardle, D. A., Cros, C., & Fontaine, S. (2020). Rhizosphere control of soil nitrogen cycling: a key component of plant economic strategies. *New Phytologist*, *228*(4), 1269–1282. doi: 10.1111/nph.16760

Hungate, B. A., Stiling, P. D., Dijkstra, P., Johnson, D. W., Ketterer, M. E., Hymus, G. J., … Drake, B. G. (2004). CO_2 elicits long-term decline in nitrogen fixation. *Science*, *304*(5675), 1291. doi: 10.1126/science.1095549

IPCC. (2022). Climate Change 2022 Impacts, Adaptation and Vulnerability. In *Intergovernmental Panel on Climate Change*.

Jones, M. B., & Donnelly, A. (2004). Carbon sequestration in temperate grassland ecosytems and the inlfuence of management, climate and elevated CO_2. *New Phytologist*, *164*, 423–439.

Jumpponen, A., Högberg, P., Huss-Danell, K., & Mulder, C. P. H. (2002). Interspecific and spatial differences in nitrogen uptake in monocultures and two-species mixtures in north European grasslands. *Functional Ecology*, *16*(4), 454–461. doi: 10.1046/j.1365-2435.2002.00642.x

Kohmann, M. M., Sollenberger, L. E., Dubeux, J. C. B., Silveira, M. L., & Moreno, L. S. B. (2019). Legume proportion in grassland litter affects decomposition dynamics and nutrient mineralization. *Agronomy Journal*, *111*(3), 1079–1089. doi: 10.2134/agronj2018.09.0603

Küchenmeister, F., Küchenmeister, K., Wrage, N., Kayser, M., & Isselstein, J. (2012). Yield and yield stability in mixtures of productive grassland species: Does species number or functional group composition matter? *Grassland Science*, *58*(2), 94–100. doi: 10.1111/j.1744-697X.2012.00242.x

Lambers, J. H. R., Harpole, W. S., Tilman, D. G., Knops, J., & Reich, P. B. (2004). Mechanisms responsible for the positive diversity-productivity relationship in Minnesota grasslands. *Ecology Letters*, *7*(8), 661–668. doi: 10.1111/j.1461-0248.2004.00623.x

Landhäusser, S. M., Chow, P. S., Turin Dickman, L., Furze, M. E., Kuhlman, I., Schmid, S., … Adams, H. D. (2018). Standardized protocols and procedures can precisely and accurately quantify non-structural carbohydrates. *Tree Physiology*, *38*(12), 1764–1778. doi: 10.1093/treephys/tpy118

Lee, T. D., Reich, P. B., & Tjoelker, M. G. (2003). Legume presence increases photosynthesis and N concentrations of co-occurring non-fixers but does not modulate their responsiveness to carbon dioxide enrichment. *Oecologia*, *137*(1), 22–31. doi: 10.1007/s00442-003-1309-1

Length, R. (2020). *emmeans: estimated marginal means, aka least-squares means* (p. https://cran.r-project.org/package=emmeans). p. https://cran.r-project.org/package=emmeans. Retrieved from https://cran.r-project.org/package=emmeans

Li, Q., Song, Y., Li, G., Yu, P., Wang, P., & Zhou, D. (2015). Grass-legume mixtures impact soil N, species recruitment, and productivity in temperate steppe grassland. *Plant and Soil*, *394*(1–2), 271–285. doi: 10.1007/s11104-015-2525-2

Liang, J., Qi, X., Souza, L., & Luo, Y. (2016). Processes regulating progressive nitrogen limitation under elevated carbon dioxide: A meta-analysis. *Biogeosciences*, *13*(9), 2689–2699. doi: 10.5194/bg-13-2689-2016

Lilley, J. M., Bolger, T. P., Peoples, M. B., & Gifford, R. M. (2001). Nutritive value and the

nitrogen dynamics of *Trifolium subterraneum* and *Phalaris aquatica* under warmer, high
CO_2 conditions. *New Phytologist*, *150*(2), 385–395. doi: 10.1046/j.1469-
8137.2001.00101.x
Lilley, Julianne M/, Bolger, T. P., & Gifford, R. M. (2001). Productivity of *Trifolium*
subterraneum and *Phalaris aquatica* under warmer, high CO_2 conditions. *New*
Phytologist, *150*, 371–383.
Loreau, M., & Hector, A. (2001). Partitioning selection and complementarity in biodiversity
experiments. *Nature*, *412*(6842), 72–76. Retrieved from
http://www.biology.mcgill.ca/faculty/loreau/
Loreau, M., Naeem, S., Inchausti, P., Bengtsson, J., Grime, J. P., Hooper, D. U., … Wardle,
D. A. (2008). Biodiversity and ecosystem functioning: current knowledge and future
challenges. *Science*, *804*(2001), 804–809. doi: 10.1126/science.1064088
Makkonen, M., Berg, M. P., Handa, I. T., Hättenschwiler, S., van Ruijven, J., van Bodegom,
P. M., & Aerts, R. (2012). Highly consistent effects of plant litter identity and functional
traits on decomposition across a latitudinal gradient. *Ecology Letters*, *15*(9), 1033–1041.
doi: 10.1111/j.1461-0248.2012.01826.x
Maschler, J., Bialic-Murphy, L., Wan, J., Andresen, L. C., Zohner, C. M., Reich, P. B., …
Crowther, T. W. (2022). Links across ecological scales: Plant biomass responses to
elevated CO2. *Global Change Biology*, (March), 1–20. doi: 10.1111/gcb.16351
Mason, N. W. H., Palmer, D. J., Romera, A., Waugh, D., & Mudge, P. L. (2017). Combining
field experiments and predictive models to assess potential for increased plant diversity
to climate-proof intensive agriculture. *Ecology and Evolution*, *7*(13), 4907–4918. doi:
10.1002/ece3.3028
McKane, R. B., Johnson, L. C., Shaver, G. R., Nadelhoffer, K. J., Rastetter, E. B., Fry, B., …
Murray, G. (2002). Resource-based niches provide a basis for plant species diversity and
dominance in arctic tundra. *Nature*, *415*(6867), 68–71. doi: 10.1038/415068a
McKeon, G. M., Stone, G. S., Syktus, J. I., Carter, J. O., Flood, N. R., Ahrens, D. G., … Day,
K. A. (2009). Climate change impacts on northern Australian rangeland livestock
carrying capacity: A review of issues. *Rangeland Journal*, *31*(1), 1–29. doi:
10.1071/RJ08068
Mclaren, J. R., & Turkington, R. (2010). Plant functional group identity differentially affects
leaf and root decomposition. *Global Change Biology*, *16*(11), 3075–3084. doi:
10.1111/j.1365-2486.2009.02151.x
Meilhac, J., Deschamps, L., Maire, V., Flajoulot, S., & Litrico, I. (2020). Both selection and
plasticity drive niche differentiation in experimental grasslands. *Nature Plants*, *6*(1), 28–
33. doi: 10.1038/s41477-019-0569-7
Melesse, A., Steingass, H., Schollenberger, M., & Rodehutscord, M. (2017). Screening of
common tropical grass and legume forages in Ethiopia for their nutrient composition and
methane production profile in vitro. *Tropical Grasslands-Forrajes Tropicales*, *5*(3),
163–175. doi: 10.17138/TGFT(5)163-175
Meterology, B. of. (2020). Bureau of Meterology. Retrieved August 15, 2018, from
Government, Australian website: http://www.bom.gov.au/
Miegoue, E., Tendonkeng, F., Lemoufouet, J., Noumbissi, M. N. B., Mweugang, N. N.,
Zougou, T. G., … Tedonkeng, E. P. (2016). Pre-weaning growth of cavies feed on
Panicum maximum supplemented with diet containing *Arachis glabrata*, *Calliandra*
calothyrsus or *Desmodium intortum*. *International Journal of Biological and Chemical*
Sciences, *10*(1), 313. doi: 10.4314/ijbcs.v10i1.24
Mueller, K. E., Blumenthal, D. M., Pendall, E., Carrillo, Y., Dijkstra, F. A., Williams, D.
G., … Morgan, J. A. (2016). Impacts of warming and elevated CO_2 on a semi-arid
grassland are non-additive, shift with precipitation, and reverse over time. *Ecology*

Letters, *19*(8), 956–966. doi: 10.1111/ele.12634

Mulder, C. P. H., Jumpponen, A., Högberg, P., & Huss-Danell, K. (2002). How plant diversity and legumes affect nitrogen dynamics in experimental grassland communities. *Oecologia*, *133*(3), 412–421. doi: 10.1007/s00442-002-1043-0

Munson, S. M., & Long, A. L. (2017). Climate drives shifts in grass reproductive phenology across the western USA. *New Phytologist*, *213*(4), 1945–1955. doi: 10.1111/nph.14327

Naudts, K., Van den Berge, J., Farfan, E., Rose, P., AbdElgawad, H., Ceulemans, R., … Nijs, I. (2013). Future climate alleviates stress impact on grassland productivity through altered antioxidant capacity. *Environmental and Experimental Botany*, *99*(2014), 150–158. doi: 10.1016/j.envexpbot.2013.11.003

Nie, M., Lu, M., Bell, J., Raut, S., & Pendall, E. (2013). Altered root traits due to elevated CO_2: A meta-analysis. *Global Ecology and Biogeography*, *22*(10), 1095–1105. doi: 10.1111/geb.12062

Nyfeler, D., Huguenin-Elie, O., Suter, M., Frossard, E., & Lüscher, A. (2011). Grass-legume mixtures can yield more nitrogen than legume pure stands due to mutual stimulation of nitrogen uptake from symbiotic and non-symbiotic sources. *Agriculture, Ecosystems and Environment*, *140*(1–2), 155–163. doi: 10.1016/j.agee.2010.11.022

Oram, N. J., Ravenek, J. M., Barry, K. E., Weigelt, A., Chen, H., Gessler, A., … Mommer, L. (2018). Below-ground complementarity effects in a grassland biodiversity experiment are related to deep-rooting species. *Journal of Ecology*, *106*(1), 265–277. doi: 10.1111/1365-2745.12877

Parvin, S., Uddin, S., Tausz-Posch, S., Armstrong, R., & Tausz, M. (2020). Carbon sink strength of nodules but not other organs modulates photosynthesis of faba bean (Vicia faba) grown under elevated [CO_2] and different water supply. *New Phytologist*, *227*(1), 132–145. doi: 10.1111/nph.16520

Paynel, F., Murray, P. J., & Bernard Cliquet, J. (2001). Root exudates: A pathway for short-term N transfer from clover and ryegrass. *Plant and Soil*, *229*(2), 235–243. doi: 10.1023/A:1004877214831

Pelzer, E., Hombert, N., Jeuffroy, M. H., & Makowski, D. (2014). Meta-analysis of the effect of nitrogen fertilization on annual cereal–legume intercrop production. *Agronomy Journal*, *106*(5), 1775–1786. doi: 10.2134/agronj13.0590

Perring, M. P., Cullen, B. R., Johnson, I. R., & Hovenden, M. J. (2010). Modelled effects of rising CO_2 concentration and climate change on native perennial grass and sown grass-legume pastures. *Climate Research*, *42*(1), 65–78. doi: 10.3354/cr00863

Pirhofer-Walzl, K., Rasmussen, J., Høgh-Jensen, H., Eriksen, J., Søegaard, K., & Rasmussen, J. (2012). Nitrogen transfer from forage legumes to nine neighbouring plants in a multi-species grassland. *Plant and Soil*, *350*(1–2), 71–84. doi: 10.1007/s11104-011-0882-z

Plett, J. M., Plett, K. L., Bithell, S. L., Mitchell, C., Moore, K., Powell, J. R., & Anderson, I. C. (2016). Improved Phytophthora resistance in commercial chickpea (*Cicer arietinum*) varieties negatively impacts symbiotic gene signalling and symbiotic potential in some varieties. *Plant Cell and Environment*, *39*(8), 1858–1869. doi: 10.1111/pce.12757

R Core Team. (2020). *R: A language and environment for statistical computing* (p. http://www.r-project.org/). p. http://www.r-project.org/. Retrieved from http://www.r-project.org/

Rasmussen, J., Gylfadóttir, T., Loges, R., Eriksen, J., & Helgadóttir, Á. (2013). Spatial and temporal variation in N transfer in grass-white clover mixtures at three Northern European field sites. *Soil Biology and Biochemistry*, *57*, 654–662. doi: 10.1016/j.soilbio.2012.07.004

Reich, P. B. (2009). Elevated CO_2 reduces losses of plant diversity caused by nitrogen deposition. *Science*, *326*, 1399–1402.

Reich, P. B., Hobbie, S. E., & Lee, T. D. (2014). Plant growth enhancement by elevated CO_2
eliminated by joint water and nitrogen limitation. *Nature Geoscience*, *7*(12), 920–924.
doi: 10.1038/ngeo2284
Reich, P. B., Hobbie, S. E., Lee, T. D., & Pastore, M. A. (2018). Unexpected reversal of C3
versus C4 grass response to elevated CO_2 during a 20-year field experiment. *Science*,
361(6407), 317–320. doi: 10.1126/science.aau8982
Reich, P. B., Knops, J., Tilman, D. G., Craine, J. M., Ellsworth, D. S., Tjoelker, M. G., …
Bengston, W. (2001). Plant diversity enhances ecosystem responses to elevated CO_2 and
nitrogen deposition. *Nature*, *410*, 809–812.
Rogers, A., Ainsworth, E. A., & Leakey, A. D. B. (2009). Will elevated carbon dioxide
concentration amplify the benefits of nitrogen fixation in legumes? *Plant Physiology*,
151(3), 1009–1016. doi: 10.1104/pp.109.144113
Shaw, M. R., Zavaleta, E. S., Chiariello, N. R., Cleland, E. E., Mooney, H. A., & Field, C. B.
(2002). Grassland responses to global environmental changes suppressed by elevated
CO_2. *Science*, *298*(2002), 1987–1990. doi: 10.1126/science.1075312
Sherry, R. A., Zhou, X., Gu, S., Arnone, J. A., Schimel, D. S., Verburg, P. S., … Luo, Y.
(2007). Divergence of reproductive phenology under climate warming. *PNAS*, *104*(1),
198–202.
Soussana, J. F., & Lüscher, A. (2007). Temperate grasslands and global atmospheric change:
A review. *Grass and Forage Science*, *62*(2), 127–134. doi: 10.1111/j.1365-
2494.2007.00577.x
Still, C. J., Cotton, J. M., & Griffith, D. M. (2019). Assessing earth system model predictions
of C4 grass cover in North America: From the glacial era to the end of this century.
Global Ecology and Biogeography, *28*(2), 145–157. doi: 10.1111/geb.12830
Szulejko, J. E., Kumar, P., Deep, A., & Kim, K. H. (2017). Global warming projections to
2100 using simple CO_2 greenhouse gas modeling and comments on CO2 climate
sensitivity factor. *Atmospheric Pollution Research*, *8*(1), 136–140. doi:
10.1016/j.apr.2016.08.002
Terrer, C., Phillips, R., Hungate, B., Rosende, J., Pett-Ridge, J., Craig, M., … Jackson, R.
(2021). A trade-off between plant and soil carbon storage under elevated CO_2. *Nature*,
591(July 2020). doi: 10.1038/s41586-021-03306-8
Thilakarathna, M. S., McElroy, M. S., Chapagain, T., Papadopoulos, Y. A., & Raizada, M. N.
(2016). Belowground nitrogen transfer from legumes to non-legumes under managed
herbaceous cropping systems. A review. *Agronomy for Sustainable Development*, *36*(4).
doi: 10.1007/s13593-016-0396-4
Thomas, D., & Sumberg, J. E. (1995). A review of the evaluation and use of tropical forage
legumes in sub-Saharan Africa. *Agriculture, Ecosystems and Environment*, *54*(3), 151–
163. doi: 10.1016/0167-8809(95)00584-F
Tilman, D. G., Isbell, F. I., & Cowles, J. M. (2014). Biodiversity and ecosystem functioning.
Annual Review of Ecology, Evolution, and Systematics, *45*, 471–493. doi:
10.1146/annurev-ecolsys-120213-091917
Vandermeer, J. (1981). The interference production principle: an ecological theory for
agriculture. *BioScience*, *31*(5), 361–364. doi: 10.2307/1308400
Vitousek, P. M., & Howarth, R. W. (1991). Nitrogen limitation on land and in the sea: how
can it occur. *Biogeochemistry*, *13*(2), 87–115. doi: 10.1007/BF00002772
von Felton, S., Hector, A., Buchmann, N., Niklaus, P. A., Schmid, B., & Scherer-Lorenzen,
M. (2009). Belowground nitrogen partitioning in experimental grassland plant
communities of varying species richness. *Ecology*, *90*(3), 1389–1399.
Wales, D. of P. I. N. S. (2018). Pasture species and varieties. Retrieved August 5, 2018, from
Pastures and rangelands website: https://www.dpi.nsw.gov.au/agriculture/pastures-and-

rangelands/species-varieties

Wang, C., Hunt, E. R., Zhang, L., & Guo, H. (2013). Phenology-assisted classification of C3
and C4 grasses in the U.S. Great Plains and their climate dependency with MODIS time
series. *Remote Sensing of Environment*, *138*, 90–101. doi: 10.1016/j.rse.2013.07.025

Wang, Z., Wang, C., & Liu, S. (2022). Elevated CO_2 alleviates adverse effects of drought on
plant water relations and photosynthesis: A global meta-analysis. *Journal of Ecology*,
(August), 1–14. doi: 10.1111/1365-2745.13988

Warwick, K., Allen, F. L., Keyser, P. D., Ashworth, A. J., Bates, G. E., Tyler, D. D., …
Harper, C. A. (2016). Biomass and integrated forage/biomass yields of switchgrass as
affected by intercropped cool- and warm-season legumes. *Journal of Soil and Water*
Conservation, *71*(1), 21–28. doi: 10.2489/jswc.71.1.21

Wei, X., Reich, P. B., & Hobbie, S. E. (2019). Legumes regulate grassland soil N cycling and
its response to variation in species diversity and N supply but not CO2. *Global Change*
Biology, *25*(7), 2396–2409. doi: 10.1111/gcb.14636

Wery, J., Turc, O., & Salsac, L. (1986). Relationship between growth, nitrogen fixation and
assimilation in a legume (Medicago sativa L.). *Plant and Soil*, *9629*, 17–29.

Winkler, J. B., & Herbst, M. (2004). Do plants of a semi-natural grassland community benefit
from long-term CO_2 enrichment? *Basic and Applied Ecology*, *5*(2), 131–143. doi:
10.1078/1439-1791-00219

Xie, Q., Huete, A., Hall, C. C., Medlyn, B. E., Power, S. A., Davies, J. M., … Beggs, P. J.
(2022). Satellite-observed shifts in C3/C4 abundance in Australian grasslands are
associated with rainfall patterns. *Remote Sensing of Environment*, *273*(February),
112983. doi: 10.1016/j.rse.2022.112983

Zelikova, T. J., Blumenthal, D. M., Williams, D. G., Souza, L., LeCain, D. R., Morgan, J., &
Pendall, E. (2014). Long-term exposure to elevated CO_2 enhances plant community
stability by suppressing dominant plant species in a mixed-grass prairie. *Proceedings of*
the National Academy of Sciences of the United States of America, *111*(43), 15456–
15461. doi: 10.1073/pnas.1414659111

Zhang, H., Powell, J. R., Plett, J. M., Churchill, A. C., Power, S. A., Macdonald, C. A., …
Anderson, I. C. (2021). Climate warming negates arbuscular mycorrhizal fungal
reductions in soil phosphorus leaching with tall fescue but not lucerne. *Soil Biology and*
Biochemistry, *152*(November), 108075. doi: 10.1016/j.soilbio.2020.108075

Zuppinger-Dingley, D., Schmid, B., Petermann, J. S., Yadav, V., De Deyn, G. B., & Flynn,
D. F. B. (2014). Selection for niche differentiation in plant communities increases
biodiversity effects. *Nature*, *515*(7525), 108–111. doi: 10.1038/nature13869

1000

Table 1. Tropical species and associated mixtures for addressing legume-grass facilitation

Scientific Name*	Common Name	Variety	Growth form	Paired species
Chloris gayana	Rhodes grass	Katambora	Grass	Burgundy
Desmodium intortum	Greenleaf Desmodium	Greenleaf	Legume	Panic
Macroptilium bracteatum	Burgundy Bean	Presto	Legume	Rhodes
Panicum maximum var. trichoglume	Panic grass	Megamax 059	Grass	Desmodium

Species are referenced by genera in text

Table 2) Effects of CO_2 treatment (CO_2; ambient, elevated), species pair (SP; *Macroptilium*/*Chloris* and *Desmodium*/*Panicum*) and plant type (PT; grass, legume, mixed-species), for pot level total shoot biomass, total shoot N, total shoot N derived from soil (shoot N- soil), the percent of N derived from biologically fixed N (Shoot N- BNF %; legume and mixed plant types only) and total shoot N derived from BNF (Shoot N- BNF content; legume and mixed plant types only).

Response	Factors	F Statistic*	P Value	r^2-M	r^2-C
Shoot biomass^	CO_2	$5.23_{1,4}$	0.08	0.40	0.45
	Species pair	$49.9_{1,143}$	<0.01		
	Plant type	$12.1_{2,143}$	<0.01		
	CO_2 x SP	$1.9_{1,143}$	0.17		
	CO_2 x PT	$2.5_{2,143}$	0.08		
	SP x PT	$5.6_{2,143}$	<0.01		
	CO_2 x SP x PT	$1.0_{2,143}$	0.38		
Shoot N content^	CO_2	$0.3_{1,4}$	0.60	0.65	0.66
	Species pair	$102.7_{1,143}$	<0.01		
	Plant type	$79.4_{2,143}$	<0.01		
	CO_2 x SP	$1.9_{1,143}$	0.18		
	CO_2 x PT	$3.7_{2,143}$	0.03		
	SP x PT	$14.7_{2,143}$	<0.01		
	CO_2 x SP x PT	$0.4_{2,143}$	0.64		
Shoot N-soil	CO_2	$0.1_{1,4}$	0.79	0.14	0.15
	Species pair	$17.8_{1,143}$	<0.01		
	Plant type	$2.7_{2,143}$	0.07		
	CO_2 x SP	$0.4_{1,143}$	0.54		
	CO_2 x PT	$0.1_{1,143}$	0.92		
	SP x PT	$0.04_{2,143}$	0.95		
	CO_2 x SP x PT	$1.3_{2,143}$	0.27		

Subscripts indicate degrees of freedom, ^data were ln transformed to meet assumptions of normality. All models included growth chamber as a random effect.

Table 3) Effects of CO_2 treatment (CO_2; ambient, elevated), species pair (SP; *Macroptilium*/*Chloris* and *Desmodium*/*Panicum*) and plant type (PT; legume and mixed-species only), for pot level percent of N derived from biologically fixed N (Shoot N- BNF %; legume and mixed plant types only) and total shoot N derived from BNF (Shoot N-BNF content).

Response	Factors	F Statistic*	P Value	r^2-M	r^2-C
Shoot N- BNF %	CO_2	$0.03_{1,4}$	0.87	0.44	0.45
	Species pair	$59.0_{1,94}$	<0.01		
	Plant type	$13.8_{1,94}$	<0.01		
	CO_2 x SP	$2.5_{1,94}$	0.12		
	CO_2 x PT	$2.1_{1,94}$	0.15		
	SP x PT	$2.8_{1,94}$	0.10		
	CO_2 x SP x PT	$3.3_{1,94}$	0.07		
Shoot N- BNF content^	CO_2	$0.1_{1,4}$	0.80	0.59	0.63
	Species pair	$132.3_{1,94}$	<0.01		
	Plant type	$30.8_{1,94}$	<0.01		
	CO_2 x SP	$0.7_{1,94}$	0.39		
	CO_2 x PT	$3.5_{1,94}$	0.06		
	SP x PT	$1.0_{1,94}$	0.32		
	CO_2 x SP x PT	$0.2_{1,94}$	0.62		

Subscripts indicate degrees of freedom, ^data were ln transformed to meet assumptions of normality. All models included growth chamber as a random effect.

Table 4) Effects of CO_2 treatment (CO_2; ambient, elevated) for pot level total biomass of root nodules (Nodule biomass), number of root nodules present (Nodule number), and the activity of nodules at the time of harvest (Nodule activity) by species pair (SP; *Macroptilium*/*Chloris* and *Desmodium*/*Panicum*) and plant type (PT; legumes in monoculture vs mixed pots only)

Response	Factors	F Statistic*	P Value	r^2-M	r^2-C
Nodule biomass^	CO_2	$4.0_{1,4}$	0.12	0.68	0.69
	Species pair	$65.6_{1,68}$	<0.01		
	Plant type	$97.2_{1,68}$	<0.01		
	CO_2 x SP	$0.6_{1,68}$	0.43		
	CO_2 x PT	$0.1_{1,68}$	0.80		
	SP x PT	$4.6_{1,68}$	0.04		
	CO_2 x SP x PT	$0.3_{1,68}$	0.58		
Nodule number^	CO_2	$0.4_{1,4}$	0.55	0.55	0.58
	Species pair	$28.8_{1,68}$	<0.01		
	Plant type	$72.9_{1,68}$	<0.01		
	CO_2 x SP	$0.02_{1,68}$	0.88		
	CO_2 x PT	$0.1_{1,68}$	0.8		
	SP x PT	$4.0_{1,68}$	0.05		
	CO_2 x SP x PT	$0.4_{1,68}$	0.52		
Nodule activity^	CO_2	$4.7_{1,4}$	0.10	0.16	0.16
	Species pair	$0.7_{1,68}$	0.42		
	Plant type	$7.0_{1,68}$	0.01		
	CO_2 x SP	$0.6_{1,68}$	0.44		
	CO_2 x PT	$1.0_{1,68}$	0.33		

SP x PT	$0.01_{1,68}$	0.90
CO_2 x SP x PT	$1.9_{1,68}$	0.17