1

DksA homologs, ppGpp and RegAB regulate dissimilatory nitrogen metabolism in

Paracoccus denitrificans

Ashvini Ray and Stephen Spiro*

Department of Biological Sciences, University of Texas at Dallas, Richardson,

Texas 75080.

*For correspondence: stephen.spiro@utdallas.edu

Running title: Roles of *dksA*, *relA* and *regAB* in *P. denitrificans*

Keywords: DksA, guanosine tetraphosphate, *Paracoccus denitrificans*, redox homeostasis, RegAB

ABSTRACT

The *Paracoccus denitrificans* genome encodes three members of the DksA/TraR family of transcription factors, two of which appear to be *bona fide* DksA proteins (hence designated DksA1$_{Pd}$ and DksA2$_{Pd}$). There is a single *relA/spoT* gene in the genome encoding a predicted bifunctional ppGpp synthetase and hydrolase (designated RSH, for RelASpoT Homolog). Both DksA1$_{Pd}$ and DksA2$_{Pd}$ can rescue the amino acid auxotrophy of an *Escherichia coli dksA* mutant. Deletion of either *dksA1* or *relA/spoT* eliminates the upregulation of expression of the periplasmic nitrate reductase (NAP) that is seen when cultures are grown on a reduced substrate such as butyrate. Thus, we conclude that DksA1$_{Pd}$ and ppGpp activate *nap* transcription in response to growth on a reduced substrate and that these proteins therefore help to maintain redox homeostasis by activating a mechanism for the disposal of excess reductant. We also show that the redox-sensing RegAB two-component pair acts as a negative regulator of *nap* expression under anaerobic growth conditions. Under the same conditions, RegAB acts as a positive regulator of expression of the membrane-associated nitrate reductase Nar and thus mediates reciprocal regulation of the two nitrate reductases that have distinct physiological roles. The *dksA1* and *relA* genes are conditionally synthetically lethal; the double mutant has a null phenotype for growth on butyrate and other reduced substrates while growing normally on succinate and citrate. The flavohemoglobin gene *hmp* is modestly upregulated in *dksA2* and *relA* mutants, which are also defective for biofilm formation (possibly because of enhanced scavenging of nitric oxide).

IMPORTANCE

Paracoccus denitrificans is a metabolically versatile Gram-negative bacterium that is used as a model for studies of respiratory metabolism. The organism can utilize nitrate as an electron

acceptor for anaerobic respiration, reducing it to dinitrogen via nitrite, nitric oxide, and nitrous

oxide. This pathway (known as denitrification) is important as a route for loss of fixed nitrogen

from soil and as a source of the greenhouse gas nitrous oxide. Thus, it is important to understand

those environmental and genetic factors that govern flux through the denitrification pathway. Here

we identify novel genes, the protein products of which function as regulators of expression of

enzymes that reduce nitrate and nitric oxide.

INTRODUCTION

Paracoccus denitrificans is a gram-negative α-proteobacterium and a facultative anaerobe that can

respire using nitrogen oxyanions and oxides as terminal electron acceptors in a process known as

denitrification (1). In the denitrification pathway, the toxic intermediate nitric oxide (NO) is

generated from nitrite by the respiratory nitrite reductase and is subsequently reduced to nitrous

oxide (N_2O) by the respiratory NO reductase. *P. denitrificans* also has an *hmp* gene encoding a

flavohemoglobin (Hmp) that potentially contributes to the metabolism of NO (2, 3). NO acts in

conjunction with transcriptional regulators as an important signal molecule to regulate gene

expression during the transition from aerobic respiration to denitrification (4, 5). Pathogenic

strains of *Salmonella* and *E. coli* have several transcriptional regulators which control the

expression of genes encoding enzymes that scavenge NO. These regulators and their target genes

may help to protect the bacteria against the NO that is made by host phagocytes (6–8). One such

transcriptional regulator is DksA, which acts as a thiol based NO sensor in *Salmonella* and

regulates detoxification of NO through Hmp in *E. coli* (9–11).

DksA, first described in *E. coli* as DnaK suppressor A (12) is a founder member of

TraR/DksA family of transcriptional regulators (13, 14). $DksA_{Ec}$ (*E. coli* DksA) is 151 amino acids

long and consists of three major domains (Fig. S1): a coiled-coil (CC) domain with a conserved

DXXDXA motif at its tip, a globular domain with a conserved four-cysteine Zn-finger motif, and

a C-terminal α-helical domain (15). The conserved acidic residues located at the tip of the CC

domain in $DksA_{Ec}$ are critical for binding to RNA polymerase (RNAP) and for $DksA_{Ec}$ to function

as a transcriptional regulator (16, 17). The globular domain has been shown to interact with the

rim of the secondary channel of RNAP (18), and the conserved redox-sensitive cysteines act as a

sensor for NO in *Salmonella* (9). Some bacterial genomes encode multiple homologs of DksA,

which differ in the presence or absence of a complete DXXDXA motif or the presence of one, two,

or four conserved cysteines in the Zn-finger motif (19). For instance, *Pseudomonas aeruginosa*

has two functional homologs, DksA1 and DksA2 (20). DksA1 has the conserved four cysteine Zn-

finger motif whereas DksA2 does not (21, 22). In α-proteobacteria more closely related to *P.*

denitrificans, such as *Sinorhizobium meliloti* and *Rhodobacter sphaeroides*, the assigned *dksA*

gene encodes a protein with a complete DXXDXA motif and only one cysteine in the Zn-finger

motif (23, 24). Deletion or overexpression of *dksA* in various species has pleiotropic effects on

multiple physiological processes. These include amino acid biosynthesis (25), motility (26),

biofilm formation (26), plant symbiosis (23), DNA repair (27), quorum sensing and virulence (28)

and oxidative and nitrosative damage (9, 11).

DksA, along with the alarmone guanosine-5', 3'-tetraphosphate (ppGpp) is known to alter

global gene expression to adjust cellular metabolism in response to nutritional stress, a mechanism

known as the stringent response (when triggered by amino acid starvation) in *E. coli* and other

related γ-proteobacteria (29–31). ppGpp is found throughout the bacterial kingdom as a second

messenger and is associated with various stress response mechanisms and growth rate regulation

(29). The metabolism of ppGpp is controlled by two enzymes encoded by *relA* and *spoT* (32, 33).

RelA is a ribosome-associated ppGpp synthetase, whereas SpoT is a bifunctional enzyme responsible for both synthesis and hydrolysis of ppGpp. The presence of both *relA* and *spoT* is characteristic of γ-proteobacteria, while only one RelA/SpoT bifunctional homolog (RSH) possessing both conserved ppGpp synthase and hydrolase domains is typically present in α-proteobacteria (34, 35). While DksA and ppGpp coregulate many genes, some phenotypic and transcriptomic studies report independent effects of both DksA and ppGpp (36, 37). The independent effects of DksA and ppGpp might be attributed to their direct binding to RNAP; ppGpp is also a direct effector of a number of metabolic enzymes (38–40)

In *P. denitrificans* Pd1222, three gene products have been annotated as members of the TraR/DksA family: Pden_0547 (which we designate $DksA1_{Pd}$), Pden_0916 ($DksA2_{Pd}$), and Pden_1862, which are 45%, 26% and 30% identical to *E. coli* DksA, respectively. The *P. denitrificans* proteins are more similar to their homologs from related α-proteobacteria. $DksA1_{Pd}$, $DksA2_{Pd}$, and Pden_1862 are 90%, 64%, and 33% identical to $DksA_{Rsp}$, respectively, while they are 64%, 47%, and 36% identical to $DksA_{Sm}$, respectively. *P. denitrificans* also has a single *relA*/*spoT* gene: Pden_1400 (henceforward called *relA*) which encodes for a RelA/SpoT bifunctional homolog (RSH). The predicted RSH is 705 amino acids long and possesses a phosphohydrolase HD domain with a conserved HDXXED motif in the N-terminal region, a RelA-SpoT region, and conserved TGS and ACT domains in the C-terminal domain (Fig. S2). On the basis of sequence similarity, Pden_1400 is predicted to have both ppGpp synthetase and hydrolase activities. None of these *P. denitrificans* genes and proteins has been functionally characterized. A proteomic analysis addressing aerobic to anaerobic growth adaptation in *P. denitrificans* demonstrated a significant increase in expression of $DksA1_{Pd}$ (A1AZG5) under anaerobic

conditions. Another study using microarray analysis showed a 3.4-fold higher expression of *dksA2*

during anaerobic growth (41, 42).

In this work, we report distinct phenotypes associated with deletion of two of the *dksA*-like

genes and the *relA* homolog in *P. denitrificans* Pd1222. We show that deletion of *dksA1* causes an

aerobic growth defect on reduced carbon substrates such as butyrate and that *dksA1* and *relA* null

strains fail to up-regulate expression and activity of the periplasmic nitrate reductase NAP under

these conditions. Furthermore, we demonstrate that ppGpp is involved in the stringent response in

P. denitrificans and that a *dksA1 relA* double mutant is unable to grow on reduced carbon

substrates. In addition, we show that *dksA2* is involved in biofilm production, possibly indirectly

via negative regulation of *hmp* transcription.

RESULTS

***P. denitrificans dksA1* and *dksA2* complement Δ*dksA*$_{Ec}$ strains.**

Sequence alignment of the three DksA/TraR proteins revealed that two (Pden_0547 and

Pden_0916) have a complete DXXDXA motif which is required for interaction with RNA

polymerase (16, 17) while the third (Pden_1862) does not. Since we have also not found any

phenotype associated with deletion of Pden_1862 (data not shown) our attention has focused on

Pden_0547 and Pden_0916, which we have designated *dksA1* and *dksA2*, respectively. Unlike the

canonical DksA of *E. coli*, DksA1$_{Pd}$ has only one cysteine conserved in the zinc finger motif

whereas DksA2$_{Pd}$ has all four cysteines (Fig. S1).

E. coli MG1655 deleted for *dksA* was complemented with either *dksA1* or *dksA2* cloned in

pIND4 and expressed from their native promoter. *E. coli* Δ*dksA* is auxotrophic for multiple amino

acids and fails to grow on M9-glucose medium without amino acid supplementation (11).

Expression of either *dksA1* or *dksA2* rescued growth of Δ*dksA$_{EC}$* on M9-glucose plates suggesting

that either gene may complement the *E. coli dksA* mutation (Fig. 1).

Deletion of *dksA1$_{Pd}$* impairs nitrate stimulation of aerobic growth on butyrate

To characterize the *dksA*-like genes of *P. denitrificans*, we constructed and tested the growth of

deletion strains lacking *dksA1* and *dksA2*. The mutants grew as well as the parent strain on rich

(LB) medium under both aerobic and anaerobic conditions (Fig. S3). We also did not observe

amino acid auxotrophy when the mutants were grown in succinate or glucose minimal medium

(data not shown). In *Salmonella enterica* sv. *typhimurium*, DksA is involved in redox homeostasis

since it regulates the expression of enzymes from central metabolic pathways that generate reduced

pyridine nucleotides (43). *P. denitrificans* may experience redox imbalance during growth on

reduced carbon substrates due to the generation of excess reducing equivalents (3). Therefore, we

assayed aerobic growth in minimal medium supplemented with succinate (MM-S) or butyrate

(MM-B) as sole carbon and energy sources that are relatively oxidized and reduced, respectively.

Growth of the wild-type strain in MM-B was significantly enhanced by the addition of nitrate to

the medium. Reduction of nitrate by the periplasmic nitrate reductase NAP provides a mechanism

to dispose of excess reducing equivalents during growth on reduced carbon substrates (44). The

Δ*dksA1* mutant grew normally on MM-S but showed impaired growth on MM-B that was not

improved by addition of nitrate to the medium (Fig. 2A and B). These phenotypes suggest that

DksA1$_{Pd}$ may regulate aerobic nitrate reduction by NAP during growth on reduced carbon

substrates.

To determine if this Δ*dksA1* associated phenotype is ppGpp-dependent, we measured

growth of the Δ*relA* strain under similar conditions. The Δ*relA* mutant showed significantly

reduced growth in MM-B, however, unlike for Δ*dksA1*, growth was improved by addition of

nitrate to the medium (Fig. 3A and B), though the magnitude of nitrate stimulation was less than

was observed in the wild-type strain.

Δ*dksA1* and Δ*relA* mutants have reduced *napE* promoter activity and NAP enzyme activity

It has been shown previously that the periplasmic nitrate reductase, NAP, has increased expression

and activity in cells grown aerobically on reduced energy sources (45, 46). Our results suggest that

the absence of *dksA1* impairs nitrate reduction in *P. denitrificans* during aerobic growth. We

hypothesized that DksA1$_{Pd}$ regulates expression of NAP during aerobic growth on reduced energy

sources. To test this hypothesis, we constructed a *napE-lacZ* promoter transcriptional fusion and

measured *napE* promoter activity using β-galactosidase as the reporter. Consistent with previous

reports (5), we observed an increase in *napE* promoter activity in cells grown aerobically in MM-

B compared to those grown in MM-S (Fig. 4A). The increase in *napE* promoter activity caused by

growth on butyrate was eliminated in the Δ*dksA1* mutant. The increase in *napE* promoter activity

in MM-B versus MM-S also disappeared when the cultures were grown anaerobically (Fig. 4A).

In parallel with the *napE* promoter activity assays we also measured the activity of NAP in

periplasmic fractions using reduced methylviologen as the electron donor to the enzyme. As

previously reported (45), and consistent with our promoter activity data, NAP activity in the wild-

type was significantly higher during aerobic growth in MM-B compared to MM-S (Fig. 4B). The

stimulation of activity by growth on butyrate disappeared in the Δ*dksA1* mutants and in cells grown

anaerobically in either MM-S or MM-B (Fig. 4B). Our results suggest that DksA1$_{Pd}$ upregulates

(either directly or indirectly) the *napE* promoter during aerobic growth on reduced carbon

substrates such as butyrate.

The *relA* mutant was found to have the same phenotype as the *dksA1* mutant with respect

to *nap* promoter activity and NAP enzyme activity. That is, the butyrate-dependent stimulation

was also lost in the *relA* mutant. Since DksA and ppGpp frequently act in concert by binding

together to a single site on RNA polymerase (30), the likely explanation that DksA-dependent

regulation of *nap* requires ppGpp.

The RegAB system regulates anaerobic *napE* promoter activity and NAP enzyme activity

In *P. pantotrophus*, regulation of the *nap* operon in response to carbon substrate oxidation state

involves *cis*-acting regulatory sequences in the *napE* promoter region (48). Specifically, a

heptameric imperfect inverted repeat sequence (TGAGACAttTGTCGCA) was suggested to be the

binding site for a negatively acting redox-responsive regulator (48). The inverted repeat is a partial

match to the binding site (GCGGC-N$_5$-GTCGC) for the response regulator RegR (RegA) in

Bradyrhizobium japonicum (49). We found a similar sequence (5′-**GAGGC**GTTTT**GACGC**-3′)

in the promoter region of the *nap* operon in *P. denitrificans* Pd1222. Therefore, we examined *napE*

promoter activity in Δ*regA* and Δ*regB* strains to determine if *nap* is regulated by *dksA1* alongside

the RegAB system. In the Δ*regA* mutant, we observed ~2-fold increases in *napE* promoter activity

in cells grown aerobically on succinate (Fig. 5A). In cells grown aerobically on butyrate, *napE*

promoter activity was similar in both *regA* and *regB* mutants and the wild-type strain. Interestingly,

there were significant increases in promoter activities in *regA* and *regB* mutants grown

anaerobically on either succinate or butyrate (Fig. 5A). These data suggest that RegA is the

negatively acting regulator that Ellington et al. suggested (48) binds to the inverted repeat sequence

in the *napE* promoter, and that the RegAB system acts to down-regulate this promoter principally

under anaerobic growth conditions (but also to a lesser extent) during aerobic growth on succinate.

The less severe phenotype of the *regB* mutant under some conditions suggests the possibility that

RegA is phosphorylated by another histidine kinase in the absence of RegB. Assays of NAP

activity in periplasmic fractions prepared from anaerobically grown cells confirm that RegAB

functions as a negative regulator under these conditions (Fig. 5B).

The membrane-associated nitrate reductase, Nar, is upregulated in *P. denitrificans* in

oxygen-limiting conditions (1, 50). Known regulators of the *nar* promoter are NarR (50) and,

probably, FnrP (51). Assays of Nar activity in membrane fractions revealed a reduced activity in

both Δ*regA* and Δ*regB* mutants suggesting a positive regulation of the *nar* operon by RegAB under

these conditions (Fig. 5C). Addition of 20 μM sodium azide (a selective inhibitor of Nar) to the

reaction mixture inhibited nitrate reduction confirming that the reductase activity in the membrane

fraction was due to Nar (data not shown). Reduced Nar activity also coincided with reduced growth

yield for the Δ*regA* mutant when grown anaerobically (Fig. S4). Thus, the RegAB system mediates

reciprocal regulation of the two nitrate reductases under anaerobic conditions, being a negative

regulator of NAP expression and a positive regulator of Nar expression.

The *dksA1* and *relA* genes are synthetically lethal for growth on reduced substrates

The DksA protein interacts with ppGpp to regulate several functions and deletion of either *dksA*

or *relA* gives rise to overlapping pleiotropic phenotypes (36). The similar phenotypes of Δ*dksA1*

and *relA* mutants with respect to loss of butyrate stimulation of *nap* promoter activity suggests that

DksA1$_{Pd}$ and ppGpp act in concert to regulate expression of the periplasmic nitrate reductase. A

Δ*dksA1*Δ*relA* double mutant exhibited a complete growth defect on reduced carbon substrates such

as butyrate, caproate and choline (Fig. 3, Fig. S5), while growth on the relatively oxidized

substrates succinate and citrate was normal (Fig. S6). Expression of either *dksA1* or *relA* on a broad

host range plasmid from their native promoters restored growth of the Δ*dksA1*Δ*relA* mutant,

confirming that both mutations are required for the null growth phenotype (Fig. 3). Thus, *dksA1*

and *relA* exhibit synthetic lethality with respect to growth on reduced substrates. The likely

explanation is that two (or more) genes that are independently regulated by DksA1$_{Pd}$ and ppGpp

are together required for growth on reduced substrates.

RSH is essential for initiating the stringent response in *P. denitrificans*.

The *relA/spoT* gene product is a bifunctional ppGpp synthetase/hydrolase (RSH) that synthesizes

ppGpp in response to nutrient (for example, amino acid) starvation in many bacteria (29, 52). To

determine if the RSH of *P. denitrificans* is involved in the stringent response to amino acid

starvation, wild-type strains were treated with serine hydroxamate (SHX) during growth in MM-

S. SHX, a structural analog of serine is known to cause amino acid starvation in a concentration

dependent manner and has been used to induce the stringent response in *E. coli*, which is

characterized by accumulation of ppGpp (12, 31). When strain Pd1222 was treated with SHX, we

observed that, unlike *E. coli* (53), there was no inhibition of growth at concentrations below 3 mM

(data not shown). On treatment with 3 mM SHX the wild-type strain exhibited an unusually long

lag phase but recovered growth while the Δ*relA* strain was completely inhibited (Fig 6B). Addition

of casamino acids reversed the SHX-induced growth inhibition in the wild-type and Δ*relA* strains

(Fig. 6C). Constitutive expression of *relA*$_{Pd}$ in the deletion strain reversed the observed phenotypes

and the strains grow as well as the wild-type strain (Fig. 6B and C) suggesting that RSH is required

for initiating the stringent response following amino acid starvation.

DksA2$_{Pd}$ is a positive regulator of the *hmp* promoter

In *E. coli*, *dksA* regulates the expression and activity of Hmp, a flavohemoprotein. Hmp is a nitric

oxide dioxygenase or denitrosylase which oxidizes nitric oxide to nitrate, and is known to be the

dominant NO defense mechanism under aerobic conditions (11). *P. denitrificans* has an *hmp* gene,

Pden_1689 (2, 3) and we have shown that the Hmp is functional for NO scavenging (unpublished

data). To determine if regulation of *hmp* is mediated by DksA in this organism, we assayed *hmp*

promoter activity in *P. denitrificans* and the Δ*dksA* mutants. Our results indicate that *hmp* promoter

activity is modestly but significantly elevated in the Δ*dksA2* mutant but is unaffected by the

Δ*dksA1* deletion (Fig. 7A). The Δ*relA* mutant also shows higher *hmp* promoter activity similar to

the Δ*dksA2* strain suggesting that the regulation of *hmp* by DksA2 is ppGpp-dependent (Fig. 7A).

Both DksA2$_{Pd}$ and RSH regulate biofilm formation in *P. denitrificans*

Our previous work has shown that deletion of *hmp* increases biofilm formation in *P. denitrificans*

Pd1222 possibly due to accumulation of NO (2). Given the evidence that DksA2$_{Pd}$ regulates *hmp*

expression in *P. denitrificans*, we assayed attached growth to determine if DksA2$_{Pd}$ also affects

biofilm formation. We observed that Δ*dksA2* strains had greatly reduced levels of attached growth

(Fig. 7B and C). Addition of an *hmp* mutation to the *dksA2* mutant rescued biofilm formation to

wild-type levels (Fig. 7B and C) suggesting that the *dksA2* phenotype requires Hmp. Deletion of

nsrR, which encodes a transcriptional repressor of *hmp* (2, and unpublished data), led to

significantly reduced biofilm formation similar to Δ*dksA2* (Fig. 7B and C). We also observed

reduced biofilm in Δ*relA* strains suggesting that *dksA2* and *relA* could be jointly involved in

regulating biofilm formation in *P. denitrificans* via negative regulation of *hmp* (Fig. 7B and C).

DISCUSSION

Expression of the periplasmic nitrate reductase NAP is upregulated by aerobic growth on reduced

carbon and energy sources, the likely explanation being that nitrate reduction by NAP provides a

means to dispose of excess reducing equivalents (44–48). We have shown that DksA1$_{Pd}$ and RSH

are required for the butyrate-stimulation of *napE* promoter and NAP enzyme activity. RSH is the

only protein encoded in the *P. denitrificans* genome with predicted ppGpp synthetase activity and

that it is enzymatically functional is supported by the detection of ppGpp in slow-growing cultures

of this organism (54). It is likely that DksA1$_{Pd}$ and ppGpp directly activate the *napE* promoter,

although indirect effects cannot be excluded. In either case, the implication is that growth on

reduced substrates somehow triggers the synthesis of ppGpp by RSH. Data from continuous

cultures show that the *napE* promoter is regulated both by growth rate and by carbon substrate,

leading to the conclusion that "…*nap* regulation is affected by a physiological signal which

integrates aspects of growth rate and carbon substrate utilization" (55). Our work suggests that

ppGpp (together with DksA1$_{Pd}$) is the integrating signal, which is consistent with the well-known

role of ppGpp in growth rate control (29).

One difference between the phenotypes of the *dksA1* and *relA* mutants is that nitrate does

not improve aerobic growth of the *dksA1* mutant on butyrate (as it does in the case of the wild-

type strain) but does modestly improve growth of the *relA* mutant. This improvement of growth

cannot be attributed to butyrate stimulation of *nap* expression, since that is lost in the *relA* mutant.

It is also the case that the *relA* mutant grows less well on butyrate (in the absence of nitrate) than

the wild-type strain and the *dksA1* mutant, and that the double mutant does not grow at all on

reduced substrates. It is likely that there are DksA1$_{Pd}$ and ppGpp regulated genes that are involved

in redox homeostasis that are yet to be identified.

We showed that regulation of *nap* expression also involves the RegAB two-component

system, which is redox-sensing and involved in redox homeostasis (56). Our results show that

RegAB is not a major player in mediating up-regulation of the *nap* promoter by growth on butyrate,

but instead negatively regulates *nap* expression during anaerobic growth. The response regulator

RegA very likely binds to an inverted repeat in the *napE* promoter that was initially characterized

in the closely related organism *P. pantotrophus* and suggested to be the binding site for a

negatively acting regulator (48). Evidently, regulation of the *napE* promoter by carbon substrate

and by oxygen availability can be separated, with the former requiring DksA1$_{Pd}$ and ppGpp and

the latter RegAB. Interestingly, RegAB also contribute (either directly or indirectly) to the

upregulation of the membrane associated nitrate reductase under anaerobic growth conditions.

Thus, NAP and Nar are reciprocally regulated by oxygen availability to be the dominant nitrate

reductases under aerobic and anaerobic conditions, respectively.

A recent study has shown that RegAB acts both positively and negatively under hypoxic

conditions in *Burkholderia pseudomallei*, an α-proteobacterium quite closely related to *P.*

denitrificans (57). Specifically, RegAB are required for upregulation of Nar under hypoxic

conditions in this organism, though this may involve both direct regulation by RegA and indirect

regulation, via FNR (57). Indirect effects of RegAB via NarR and/or FnrP are also possible in *P.*

denitrificans; we are able to discern a potential RegA binding site in the *narR-narK* intergenic

region that controls expression of the *narGHJI* genes encoding Nar (50).

In *E. coli*, DksA and ppGpp contribute to regulation of the *hmp* gene encoding the

flavohemoglobin, Hmp, which is responsible for detoxifying nitric oxide (11, 58, 59).   In *P.*

denitrificans, we find that deletion of *dksA2* or *relA* leads to modestly increased *hmp* promoter

activity and dramatically reduced ability to form biofilm. Since the biofilm-defective phenotype

requires Hmp, we suggest that reduced expression of *hmp* in the mutants causes a reduced

accumulation of nitric oxide, which stimulates biofilm formation in this organism (2). Reduced

growth as biofilm in *dksA* mutants has been reported for other organisms, though the regulatory

mechanisms are likely different (26, 60).

Our study has identified two functional DksA proteins in *P. denitrificans* that have distinct

roles in growth on reduced carbon substrates, aerobic and anaerobic nitrate respiration, expression

of flavohemoglobin, and biofilm formation. In other α-proteobacteria which have at least two

DksA homologs, physiological roles have so far been ascribed to only one of them (23, 24). Further

functional analysis of the *P. denitrificans* DksA-like proteins and of the role of ppGpp will advance

our broader understanding of DksA functions in α-proteobacteria.

MATERIALS AND METHODS

Bacterial strains and growth conditions. The bacterial strains and plasmids used are listed in

Table S1. All *Paracoccus denitrificans* strains were derived from the wild-type Pd1222 (61).

Escherichia coli S.17-1 was used as the donor and *E. coli* DH5α (pRK2013) as the helper strain

for the introduction of plasmids into *P. denitrificans* by conjugation (62, 63). *E. coli* Novablue

competent cells (Novagen) were used as the host for all genetic manipulations. The rich medium

used for routine growth was LB (Miller) Broth (10 g/l tryptone, 5 g/l yeast extract, 10 g/l NaCl),

and the defined minimal medium (MM) for *P. denitrificans* was as previously described (64),

solidified with 1.5 % agar (w/v) as needed and supplemented with carbon and energy sources and

with 20 mM nitrate as indicated. For growth assays, *P. denitrificans* strains were first streaked on

LB agar plates, single colonies were inoculated into LB broth and grown overnight in 15 ml tubes

at 250 rpm. These cells were washed with MM, inoculated into succinate-supplemented MM, and

grown overnight in 15 ml tubes at 250 rpm. Cells were pelleted, washed, and finally inoculated

into MM supplemented with different carbon sources with or without nitrate. For aerobic growth

assays, cultures were grown in triplicate wells in a BioTek plate reader with horizontal orbital

shaking at 280 rpm and a volume of 0.2 ml (<50% of total volume) to ensure proper aeration while

preventing cross contamination. The optical density was measured at 600 nm and adjusted for path

length variation. Serine hydroxamate was added directly from a freshly made 100 mM stock to the

growth medium to a final concentration of 3 mM at the time of inoculation, where indicated.

Anaerobic cultures for β-galactosidase assays were grown in filled glass cuvettes sealed

with a cap and septum, with medium supplemented with 20 mM nitrate. The cuvettes were gassed

with N_2 and incubated statically. Aerobic cultures were grown in 250 ml flasks shaking at 250 rpm

in MM supplemented with the indicated carbon substrate and with or without 20 mM nitrate. For

E. coli complementation assays, solid M9 medium (65) with 10 mM glucose was used, with or

without 0.2 % (w/v) casamino acids.

Plasmids were maintained by addition of antibiotics at the following concentrations:

ampicillin 100 µg/ml, kanamycin 25 µg/ml, rifampicin 100 µg/ml, tetracycline 2 µg/ml for *P.*

denitrificans and 15 µg/ml for *E. coli*. All *P. denitrificans* strains were grown at 30 ⁰C and *E. coli*

at 37 ⁰C.

Genetic manipulations. All primers used are listed in Table S2. Unmarked deletions were made

by allelic replacement using pK18*mobsacB* as described by Sullivan et al. with minor

modifications (66, 67). Briefly, in-frame marker-less deletions were constructed by PCR

amplification of ~500-1000 bp fragments from either side of the gene of interest. The two

amplicons were ligated together either by PCR using overlapping regions added to the primers or

by digestion and ligation using a restriction site incorporated in the primers (68). The ligated

fragment was first cloned into pSTBlue-1 (Novagen), then sub-cloned into pK18*mobsacB*,

transformed into *E. coli* S.17-1 and mobilized into *P. denitrificans* by conjugation. Triparental

mating was performed by mixing *P. denitrificans*, *E. coli* S.17-1 harboring recombinant

pK18*mobsacB* or pMP220 or pIND4 derivatives, and *E. coli* DH5α (pRK2013) cells grown to log

phase, in a 2:1:1 ratio and by incubation on LB agar plate for 2-3 days at 30 ⁰C. Single-crossover

recombination events were selected on rifampicin and kanamycin plates, and then recombinants

with double crossovers were isolated by growth on 6 % (w/v) sucrose plates and kanamycin

sensitivity as described by Schafer et al. (66). All mutations were designed to be in-frame deletions

to prevent polarity. Promoter-*lacZ* reporter fusions were constructed by amplifying ~300-400 bp

upstream of the start codon of genes of interest and first cloning into pSTBlue-1. The sequence

was verified and the fragment cloned into the low copy number vector pMP220 between *Eco*RI

and *Pst*I restriction sites (69). Plasmids were introduced into *P. denitrificans* by conjugation from

E. coli S.17-1. *P. denitrificans* exconjugants were selected by rifampicin and tetracycline

resistance. For complementation tests in *P. denitrificans*, the gene of interest with its native

promoter was amplified and cloned directly into pIND4 (70) and the sequence verified. The

cloning was done using the NEBuilder HiFi DNA assembly cloning kit (NEB). Plasmids were

transformed into *E. coli* S.17-1 and mobilized into *P. denitrificans* strains by conjugation as

described above.

β-galactosidase assay: Liquid cultures were grown aerobically and anaerobically in minimal

medium with the indicated carbon sources and with or without 20 mM nitrate. Cells were harvested

at the mid-log phase of growth and the assay was performed as described by Miller (71). B-

galactosidase activity was measured using 0.1 ml of cells made permeable by treatment with

sodium dodecyl sulfate and chloroform. In the case of the *hmp-lacZ* fusion, strains were grown to

mid-log phase in the same medium and growth conditions as used for the biofilm assays.

Biofilm assays: Attached growth was assayed in 60 mm x 15 mm polystyrene petri dishes (2). A

70 µl aliquot of overnight culture was inoculated into 7 ml of LB broth supplemented with 10 mM

calcium chloride. The petri dishes were incubated statically at 30 ºC for 72 hours. The plates were

then washed thrice under gently running tap water and stained with crystal violet (0.1 % in water)

for 20 minutes, washed three times with water, and air dried. For quantitation, the crystal violet

stain was extracted into 3 ml 95 % ethanol, diluted as necessary, and absorbance measured at 595

nm (2, 72).

Cell fractionation: *P. denitrificans* strains were grown aerobically and anaerobically in MM with

either succinate (10 mM) or butyrate (10 mM) as the sole carbon source; nitrate (20 mM) was

added as indicated. Aerobic cultures (50 ml) were grown at 30 ⁰C in 250 ml conical flasks shaken

at 250 rpm. Anaerobic cultures were grown in 100 ml bottles filled to the neck with medium and

incubated at 30 ⁰C without shaking. Periplasmic fractions for the nitrate reductase assay were

prepared as described in Quan et al (73) . In short, mid-exponential phase cells were harvested by

centrifugation at 3,000 g for 20 minutes at 4 ⁰C. Cells were then gently resuspended in TSE buffer

(200 mM Tris–HCl, pH 8.0, 500 mM sucrose, 1 mM EDTA without any protease inhibitor) and

incubated on ice for 30 minutes, and centrifuged at 16,000 g for 30 minutes at 4 ⁰C. The supernatant

was transferred to a fresh tube and used as the periplasmic fraction. Membrane fractions were

prepared according to Dunn et al. with some modifications (74). Briefly, cultures were harvested

in the mid-exponential phase of growth by centrifugation at 8,000 g for 5 minutes at 4 ⁰C and

washed twice in 10 mM potassium phosphate buffer (pH 7.5) in 50 ml Falcon tubes. Cells were

resuspended in 2.5 ml of ice cold 10 mM phosphate buffer, 10 mM $MgSO_4$, 0.2 M NaCl. Samples

were sonicated on ice using a Fisher Scientific Sonic Dismembrator model 500 outfitted with a 3

mm diameter tip at 20 % amplitude (10 W) for five 20 second bursts at 1 minute intervals. To

clarify the lysate, samples were centrifuged at 16,000 g for 20 minutes at 4 ⁰C. The clarified total

lysate was further centrifuged at 100,000 g for 2 hours at 4 ⁰C. The pellet was suspended in 0.2 ml

of phosphate buffer and retained as the membrane fraction for the enzyme assay. Fractions were

kept on ice and used within hours of preparation.

Enzyme activity: Nitrate reductase activity was determined spectrophotometrically using reduced

methylviologen (MV^+) as the electron donor and 10 mM sodium nitrate as the electron acceptor

(45). The extinction coefficient of reduced methylviologen used was 8.25 mM^{-1} cm^{-1} at 600 nm

(75). The protein content of the periplasmic and membrane fractions was measured by the protein

estimation method of Lowry et al., using bovine serum albumin as the standard (76).

ACKNOWLEDGMENTS

417   We are grateful to Victor Luque-Almagro, Andrew Gates and David Richardson for generously

418   providing the *regA*::*kan* mutant and to Aashna Pathi for her contribution to strain construction.

REFERENCES

1.   Baker Simon C., Ferguson Stuart J., Ludwig Bernd, Page M. Dudley, Richter Oliver-Matthias H., van
Spanning Rob J. M. 1998. Molecular Genetics of the Genus *Paracoccus*: Metabolically Versatile
Bacteria with Bioenergetic Flexibility. Microbiology and Molecular Biology Reviews 62:1046–1078.

2.   Kumar S, Spiro S. 2017. Environmental and Genetic Determinants of Biofilm Formation in
Paracoccus denitrificans. mSphere 2.

3.   Hartop KR, Sullivan MJ, Giannopoulos G, Gates AJ, Bond PL, Yuan Z, Clarke TA, Rowley G,
Richardson DJ. 2017. The metabolic impact of extracellular nitrite on aerobic metabolism of
Paracoccus denitrificans. Water Res 113:207–214.

4.   Hassan J, Bergaust LL, Wheat ID, Bakken LR. 2014. Low probability of initiating *nirS* transcription
explains observed gas kinetics and growth of bacteria switching from aerobic respiration to
denitrification. PLoS Comput Biol 10:e1003933.

5.   Hassan J, Bergaust LL, Molstad L, de Vries S, Bakken LR. 2016. Homeostatic control of nitric oxide
(NO) at nanomolar concentrations in denitrifying bacteria - modelling and experimental
determination of NO reductase kinetics in vivo in *Paracoccus denitrificans*. Environ Microbiol
18:2964–2978.

6.   Fang FC. 1997. Perspectives series: host/pathogen interactions. Mechanisms of nitric oxide-related
antimicrobial activity. J Clin Invest 99:2818–2825.

7.   Henard C, Vazquez-Torres A. 2011. Nitric Oxide and *Salmonella* Pathogenesis. Frontiers in
Microbiology 2.

8.   Hyduke DR, Jarboe LR, Tran LM, Chou KJY, Liao JC. 2007. Integrated network analysis identifies
nitric oxide response networks and dihydroxyacid dehydratase as a crucial target in *Escherichia*
coli. Proceedings of the National Academy of Sciences 104:8484–8489.

9.   Henard CA, Tapscott T, Crawford MA, Husain M, Doulias P-T, Porwollik S, Liu L, McClelland M,
Ischiropoulos H, Vázquez-Torres A. 2014. The 4-cysteine zinc-finger motif of the RNA polymerase
regulator DksA serves as a thiol switch for sensing oxidative and nitrosative stress. Mol Microbiol
91:790–804.

10. Crawford MA, Henard CA, Tapscott T, Porwollik S, McClelland M, Vázquez-Torres A. 2016. DksA-
Dependent Transcriptional Regulation in *Salmonella* Experiencing Nitrosative Stress. Frontiers in
Microbiology 7.

11. Chou WK, Brynildsen MP. 2019. Loss of DksA leads to multi-faceted impairment of nitric oxide
detoxification by *Escherichia coli*. Free Radical Biology and Medicine 130:288–296.

12. Kang PJ, Craig EA. 1990. Identification and characterization of a new *Escherichia coli* gene that is a
dosage-dependent suppressor of a dnaK deletion mutation. Journal of Bacteriology.

13. Paul BJ, Barker MM, Ross W, Schneider DA, Webb C, Foster JW, Gourse RL. 2004. DksA: A Critical
Component of the Transcription Initiation Machinery that Potentiates the Regulation of rRNA
Promoters by ppGpp and the Initiating NTP. Cell 118:311–322.

14. Rutherford ST, Lemke JJ, Vrentas CE, Gaal T, Ross W, Gourse RL. 2007. Effects of DksA, GreA, and
GreB on transcription initiation: insights into the mechanisms of factors that bind in the secondary
channel of RNA polymerase. J Mol Biol 366:1243–1257.

15. Vassylyeva MN, Perederina AA, Svetlov V, Yokoyama S, Artsimovitch I, Vassylyev DG. 2004.
Cloning, expression, purification, crystallization and initial crystallographic analysis of transcription
factor DksA from *Escherichia coli*. Acta Crystallogr D Biol Crystallogr 60:1611–1613.

16. Perederina A, Svetlov V, Vassylyeva MN, Tahirov TH, Yokoyama S, Artsimovitch I, Vassylyev DG.
2004. Regulation through the Secondary Channel—Structural Framework for ppGpp-DksA
Synergism during Transcription. Cell 118:297–309.

17. Lennon CW, Ross W, Martin-Tumasz S, Toulokhonov I, Vrentas CE, Rutherford ST, Lee J-H, Butcher
SE, Gourse RL. 2012. Direct interactions between the coiled-coil tip of DksA and the trigger loop of
RNA polymerase mediate transcriptional regulation. Genes Dev 26:2634–2646.

18. Shin Y, Qayyum MZ, Pupov D, Esyunina D, Kulbachinskiy A, Murakami KS. 2021. Structural basis of
ribosomal RNA transcription regulation. 1. Nat Commun 12:528.

19. Furman R, Tsodikov OV, Wolf YI, Artsimovitch I. 2013. An Insertion in the Catalytic Trigger Loop
Gates the Secondary Channel of RNA Polymerase. Journal of Molecular Biology 425:82–93.

20. Min KB, Yoon SS. 2020. Transcriptome analysis reveals that the RNA polymerase-binding protein
DksA1 has pleiotropic functions in *Pseudomonas aeruginosa*. J Biol Chem 295:3851–3864.

21. Blaby-Haas CE, Furman R, Rodionov DA, Artsimovitch I, de Crécy-Lagard V. 2011. Role of a Zn-
independent DksA in Zn homeostasis and stringent response. Mol Microbiol 79:700–715.

22. Furman R, Biswas T, Danhart EM, Foster MP, Tsodikov OV, Artsimovitch I. 2013. DksA2, a zinc-
independent structural analog of the transcription factor DksA. FEBS Lett 587:614–619.

23. Wippel K, Long SR. 2016. Contributions of *Sinorhizobium meliloti* Transcriptional Regulator DksA to
Bacterial Growth and Efficient Symbiosis with Medicago sativa. Journal of Bacteriology.

24. Lennon CW, Lemmer KC, Irons JL, Sellman MI, Donohue TJ, Gourse RL, Ross W. 2014. A
Rhodobacter sphaeroides Protein Mechanistically Similar to *Escherichia coli* DksA Regulates
Photosynthetic Growth. mBio.

25. Paul BJ, Berkmen MB, Gourse RL. 2005. DksA potentiates direct activation of amino acid promoters
by ppGpp. Proceedings of the National Academy of Sciences.

26. Azriel S, Goren A, Rahav G, Gal-Mor O. 2016. The Stringent Response Regulator DksA Is Required
for *Salmonella enterica* Serovar *Typhimurium* Growth in Minimal Medium, Motility, Biofilm
Formation, and Intestinal Colonization. Infection and Immunity.

27. Trautinger BW, Jaktaji RP, Rusakova E, Lloyd RG. 2005. RNA polymerase modulators and DNA
repair activities resolve conflicts between DNA replication and transcription. Mol Cell 19:247–258.

28. Jude F, Köhler T, Branny P, Perron K, Mayer MP, Comte R, van Delden C. 2003. Posttranscriptional
Control of Quorum-Sensing-Dependent Virulence Genes by DksA in *Pseudomonas aeruginosa*.
Journal of Bacteriology.

29. Hauryliuk V, Atkinson GC, Murakami KS, Tenson T, Gerdes K. 2015. Recent functional insights into
the role of (p)ppGpp in bacterial physiology. 5. Nat Rev Microbiol 13:298–309.

30. Gourse RL, Chen AY, Gopalkrishnan S, Sanchez-Vazquez P, Myers A, Ross W. 2018. Transcriptional
Responses to ppGpp and DksA. Annu Rev Microbiol 72:163–184.

31. Traxler MF, Summers SM, Nguyen H-T, Zacharia VM, Smith JT, Conway T. 2008. The global, ppGpp-
mediated stringent response to amino acid starvation in *Escherichia coli*. Mol Microbiol 68:1128–
1148.

32. Sarubbi E, Rudd KE, Xiao H, Ikehara K, Kalman M, Cashel M. 1989. Characterization of the *spoT*
gene of Escherichia coli. J Biol Chem 264:15074–15082.

33. Murray DK, Bremer H. 1996. Control ofspoT-dependent ppGpp Synthesis and Degradation in
Escherichia coli. Journal of Molecular Biology 259:41–57.

34. Mittenhuber G. 2001. Comparative genomics and evolution of genes encoding bacterial (p)ppGpp
synthetases/hydrolases (the Rel, RelA and SpoT proteins). J Mol Microbiol Biotechnol 3:585–600.

35. Richardson* DJ, Berks BC, Russell DA, Spiro S, Taylor CJ. 2001. Functional, biochemical and genetic
diversity of prokaryotic nitrate reductases. Cellular and Molecular Life Sciences CMLS 58:165–178.

36. Magnusson LU, Gummesson B, Joksimović P, Farewell A, Nyström T. 2007. Identical, Independent,
and Opposing Roles of ppGpp and DksA in *Escherichia coli*. J Bacteriol 189:5193–5202.

37. Åberg A, Fernández-Vázquez J, Cabrer-Panes JD, Sánchez A, Balsalobre C. 2009. Similar and
Divergent Effects of ppGpp and DksA Deficiencies on Transcription in *Escherichia coli*. Journal of
Bacteriology 191:3226–3236.

38. Costanzo A, Nicoloff H, Barchinger SE, Banta AB, Gourse RL, Ades SE. 2008. ppGpp and DksA likely
regulate the activity of the extracytoplasmic stress factor sigmaE in *Escherichia coli* by both direct
and indirect mechanisms. Mol Microbiol 67:619–632.

39. Szalewska-Palasz A, Johansson LUM, Bernardo LMD, Skärfstad E, Stec E, Brännström K, Shingler V.
2007. Properties of RNA polymerase bypass mutants: implications for the role of ppGpp and its co-
factor DksA in controlling transcription dependent on sigma54. J Biol Chem 282:18046–18056.

40. Wang B, Dai P, Ding D, Del Rosario A, Grant RA, Pentelute BL, Laub MT. 2019. Affinity-based
capture and identification of protein effectors of the growth regulator ppGpp. Nat Chem Biol
15:141–150.

41. Giannopoulos G, Sullivan MJ, Hartop KR, Rowley G, Gates AJ, Watmough NJ, Richardson DJ. 2017.
Tuning the modular *Paracoccus denitrificans* respirome to adapt from aerobic respiration to
anaerobic denitrification. Environ Microbiol 19:4953–4964.

42. Olaya-Abril A, Hidalgo-Carrillo J, Luque-Almagro VM, Fuentes-Almagro C, Urbano FJ, Moreno-
Vivián C, Richardson DJ, Roldán MD. 2018. Exploring the Denitrification Proteome of *Paracoccus
denitrificans* PD1222. Frontiers in Microbiology 9.

43. Henard CA, Bourret TJ, Song M, Vázquez-Torres A. 2010. Control of Redox Balance by the Stringent
Response Regulatory Protein Promotes Antioxidant Defenses of Salmonella*. Journal of Biological
Chemistry 285:36785–36793.

44. Sears HJ, Spiro S, Richardson DJ. 1997. Effect of carbon substrate and aeration on nitrate reduction
and expression of the periplasmic and membrane-bound nitrate reductases in carbon-limited
continuous cultures of *Paracoccus denitrificans* Pd1222. Microbiology (Reading) 143:3767–3774.

45. Sears HJ, Ferguson SJ, Richardson DJ, Spiro S. 1993. The identification of a periplasmic nitrate
reductase in *Paracoccus denitrificans*. FEMS Microbiology Letters.

46. Sears HJ, Sawers G, Berks BC, Ferguson SJ, Richardson DJ. 2000. Control of periplasmic nitrate
reductase gene expression (*napEDABC*) from *Paracoccus pantotrophus* in response to oxygen and
carbon substrates. Microbiology,. Microbiology Society,.

47. Ellington MJK, Bhakoo KK, Sawers G, Richardson DJ, Ferguson SJ. 2002. Hierarchy of Carbon Source
Selection in *Paracoccus pantotrophus*: Strict Correlation between Reduction State of the Carbon
Substrate and Aerobic Expression of the *nap* Operon. Journal of Bacteriology.

48. Ellington MJK, Fosdike WLJ, Sawers RG, Richardson DJ, Ferguson SJ. 2006. Regulation of the nap
operon encoding the periplasmic nitrate reductase of *Paracoccus pantotrophus*: delineation of
DNA sequences required for redox control. Archives of Microbiology 184:298–304.

49. Emmerich R, Strehler P, Hennecke H, Fischer H-M. 2000. An imperfect inverted repeat is critical for
DNA binding of the response regulator RegR of *Bradyrhizobium japonicum*. Nucleic Acids Res
28:4166–4171.

50. Wood NJ, Alizadeh T, Bennett S, Pearce J, Ferguson SJ, Richardson DJ, Moir JWB. 2001. Maximal
Expression of Membrane-Bound Nitrate Reductase in *Paracoccus* Is Induced by Nitrate via a Third
FNR-Like Regulator Named NarR. J Bacteriol 183:3606–3613.

51. Van Spanning RJ, De Boer AP, Reijnders WN, Westerhoff HV, Stouthamer AH, Van Der Oost J. 1997.
FnrP and NNR of *Paracoccus denitrificans* are both members of the FNR family of transcriptional
activators but have distinct roles in respiratory adaptation in response to oxygen limitation. Mol
Microbiol 23:893–907.

52. Lemke JJ, Sanchez-Vazquez P, Burgos HL, Hedberg G, Ross W, Gourse RL. 2011. Direct regulation of
Escherichia coli ribosomal protein promoters by the transcription factors ppGpp and DksA.
Proceedings of the National Academy of Sciences.

53. Tosa T, Pizer LI. 1971. Biochemical bases for the antimetabolite action of L-serine hydroxamate. J
Bacteriol 106:972–982.

54. van Verseveld HW, Chesbro WR, Braster M, Stouthamer AH. 1984. Eubacteria have 3 growth
modes keyed to nutrient flow. Arch Microbiol 137:176–184.

55. Ellington MJK, Sawers G, Sears HJ, Spiro S, Richardson DJ, Ferguson SJ. 2003. Characterization of
the expression and activity of the periplasmic nitrate reductase of *Paracoccus pantotrophus* in
chemostat cultures. Microbiology 149:1533–40.

56. Elsen Sylvie, Swem Lee R., Swem Danielle L., Bauer Carl E. 2004. RegB/RegA, a Highly Conserved
Redox-Responding Global Two-Component Regulatory System. Microbiology and Molecular
Biology Reviews 68:263–279.

57. Phenn J, Pané-Farré J, Meukow N, Klein A, Troitzsch A, Tan P, Fuchs S, Wagner GE, Lichtenegger S,
Steinmetz I, Kohler C. 2021. RegAB Homolog of *Burkholderia pseudomallei* is the Master Regulator
of Redox Control and involved in Virulence. PLOS Pathogens 17:e1009604.

58. Membrillo-Hernandez J, Cook GM, Poole RK. 1997. Roles of RpoS (sigmaS), IHF and ppGpp in the
expression of the *hmp* gene encoding the flavohemoglobin (Hmp) of Escherichia *coli* K-12. Mol Gen
Genet 254:599–603.

59. Wan X, Brynildsen MP. 2021. Robustness of nitric oxide detoxification to nitrogen starvation in
Escherichia coli requires RelA. Free Radic Biol Med 176:286–297.

60. Pal RR, Bag S, Dasgupta S, Das B, Bhadra RK. 2012. Functional Characterization of the Stringent
Response Regulatory Gene *dksA* of *Vibrio cholerae* and Its Role in Modulation of Virulence
Phenotypes. Journal of Bacteriology 194:5638–5648.

61. de Vries GE, Harms N, Hoogendijk J, Stouthamer AH. 1989. Isolation and characterization of
Paracoccus denitrificans mutants with increased conjugation frequencies and pleiotropic loss of a
(nGATCn) DNA-modifying property. Arch Microbiol 152:52–57.

62. Simon R, Priefer U, Pühler A. 1983. A Broad Host Range Mobilization System for In Vivo Genetic
Engineering: Transposon Mutagenesis in Gram Negative Bacteria. Bio/Technology 1:784–791.

63. Figurski DH, Helinski DR. 1979. Replication of an origin-containing derivative of plasmid RK2
dependent on a plasmid function provided in trans. Proc Natl Acad Sci U S A 76:1648–1652.

64. Hahnke S, Moosmann P, Erb T, Strous M. 2014. An improved medium for the anaerobic growth of
Paracoccus denitrificans Pd1222. Frontiers in Microbiology 5.

65. Miller JH. 1992. A short course in bacterial genetics. Cold Spring Harbor Laboratory Press, Cold
Spring Harbor, N.Y.

66. Schäfer A, Tauch A, Jäger W, Kalinowski J, Thierbach G, Pühler A. 1994. Small mobilizable multi-
purpose cloning vectors derived from the *Escherichia coli* plasmids pK18 and pK19: selection of
defined deletions in the chromosome of *Corynebacterium glutamicum*. Gene 145:69–73.

67. Sullivan MJ, Gates AJ, Appia-Ayme C, Rowley G, Richardson DJ. 2013. Copper control of bacterial
nitrous oxide emission and its impact on vitamin B12-dependent metabolism. Proceedings of the
National Academy of Sciences 110:19926–19931.

68. Wurch T, Lestienne F, Pauwels PJ. 1998. A modified overlap extension PCR method to create
chimeric genes in the absence of restriction enzymes. Biotechnology Techniques 12:653–657.

69. Spaink HP, Okker RJH, Wijffelman CA, Pees E, Lugtenberg BJJ. 1987. Promoters in the nodulation
region of the *Rhizobium leguminosarum* Sym plasmid pRL1JI. Plant Molecular Biology 9:27–39.

70. Ind Alice C., Porter Steven L., Brown Mostyn T., Byles Elaine D., de Beyer Jennifer A., Godfrey Scott
A., Armitage Judith P. 2009. Inducible-Expression Plasmid for *Rhodobacter sphaeroides* and
Paracoccus denitrificans. Applied and Environmental Microbiology 75:6613–6615.

71. 1993. Jeffrey H. Miller, A Short Course in Bacterial Genetics – A Laboratory Manual and Handbook
for *Escherichia coli* and Related Bacteria. Cold Spring Harbor 1992. Cold Spring Harbor Laboratory
Press. ISBN: 0–87969-349–5. Journal of Basic Microbiology.

72. Pratt LA, Kolter R. 1998. Genetic analysis of *Escherichia coli* biofilm formation: roles of flagella,
motility, chemotaxis and type I pili. Molecular Microbiology 30:285–293.

73. Quan S, Hiniker A, Collet J-F, Bardwell JCA. 2013. Isolation of Bacteria Envelope Proteins, p. 359–
366. *In* Delcour, AH (ed.), Bacterial Cell Surfaces: Methods and Protocols. Humana Press, Totowa,
NJ.

74. Dunn AK, Karr EA, Wang Y, Batton AR, Ruby EG, Stabb EV. 2010. The alternative oxidase (AOX)
gene in *Vibrio fischeri* is controlled by NsrR and upregulated in response to nitric oxide. Molecular
Microbiology 77:44–55.

75.   Yu L, Wolin MJ. 1969. Hydrogenase Measurement with Photochemically Reduced Methyl
Viologen1. J Bacteriol 98:51–55.

76.   Oh L, Nj R, Al F, Rj R. 1951. Protein measurement with the Folin phenol reagent. The Journal of
biological chemistry 193.

FIGURE LEGENDS

FIGURE 1.

P. denitrificans dksA1$_{Pd}$ and *dksA2$_{Pd}$* complement an *E. coli* $\Delta dksA$ mutant ($\Delta dksA_{EC}$). Strains

were streaked on M9 minimal (M9) agar plates containing 10 mM glucose (G) with or without

0.2% (w/v) casamino acids (CAA). Plates were incubated overnight at 37 ºC. 1 - *E. coli*

MG1655; 2 - $\Delta dksA_{EC}$ complemented with *dksA1$_{Pd}$*; 3 - $\Delta dksA_{EC}$ complemented with *dksA2$_{Pd}$*; 4 -

$\Delta dksA_{EC}$ carrying empty pIND4.

FIGURE 2.

Deletion of *dksA1* impairs nitrate utilization during aerobic growth. Aerobic growth of the

indicated strains in: (A) minimal medium containing succinate or (B) minimal medium

containing butyrate, with (+ NO_3^-) or without (- NO_3^-) 20 mM nitrate. Optical density was

measured (at 600 nm) at the indicated time points (the means from three independent

experiments).

FIGURE 3.

Loss of both *dksA1* and *relA* leads to failure to grow on reduced carbon substrates. Aerobic growth

of indicated strains in liquid minimal medium with butyrate (A) without (- NO_3^-) or (B) with 20

mM nitrate (+ NO$_3^-$). Optical density was measured (at 600 nm) at the indicated time points (the

means from three independent experiments).

FIGURE 4.

The Δ*dksA1* mutant has reduced *nap* promoter and NAP enzyme activity. (A) β-galactosidase

activity in *P. denitrificans* wild-type, Δ*dksA1* and Δ*relA* carrying plasmid p*nap-lacZ*. (B) NAP

enzyme activity in the periplasmic fractions of *P. denitrificans* wild-type, Δ*dksA1* and Δ*relA* cells.

Cultures were grown in minimal medium containing 10 mM succinate (S) or butyrate (B) as sole

carbon and energy source with or without 20 mM nitrate (N). Aerobic (+O$_2$) cultures were grown

in shake flasks at 250 rpm. Anaerobic (-O$_2$) cultures were grown statically in filled closed cap

bottles. The bars represent the mean of at least three separate experiments and the error bars

represent the standard deviation of the mean. Asterisks on the bars indicate a significant difference

compared to the wild-type within the group according to Student's t-test (*$P \leq 0.05$, **$P \leq 0.01$,

***$P \leq 0.001$).

FIGURE 5.

The RegAB system regulates anaerobic *nap* promoter and NAP activity. (A) β-galactosidase

activity in *P. denitrificans* wild-type, Δ*regA* and Δ*regB* carrying plasmid p*nap-lacZ*. (B) NAP

enzyme activity in the periplasmic fractions of anaerobically grown *P. denitrificans* wild-type,

Δ*regA* and Δ*regB* cells. (C) NAR enzyme activity in the membrane fractions of *P. denitrificans*

wild-type, Δ*regA* and Δ*regB* cells. Cultures were grown in minimal medium containing 10 mM

succinate (S) or butyrate (B) as sole carbon source with or without 20 mM nitrate (N). Aerobic

(+O$_2$) cultures were grown in shake flasks at 250 rpm. Anaerobic (-O$_2$) cultures were grown in

filled closed cap bottles statically. The bars represent the mean of at least three separate

experiments and the error bars represent the standard deviation of the mean. Asterisks on the bars

indicate a significant difference compared to the wild-type within the group according to Student's

t-test (*P $\leq$ 0.05, **P $\leq$ 0.01, ***P $\leq$ 0.001).

FIGURE 6.

The *relA* gene is required for the response to amino acid starvation in *P. denitrificans.* Aerobic

growth of indicated strains in minimal medium containing: (A) succinate only, (B) succinate and

3 mM serine hydroxamate, and (C) succinate, 3 mM serine hydroxamate and 0.2% (w/v) casamino

acids. Optical density was measured (at 600 nm) at the indicated time points (the means from three

independent experiments).

FIGURE 7.

DksA2$_{Pd}$ and RSH regulate *hmp* promoter activity and biofilm formation in *P. denitrificans*. (A)

β-galactosidase activities in *P. denitrificans* wild-type, Δ*dksA2* and Δ*relA* carrying plasmid p*hmp-*

lacZ. (B) Assays of biofilm formation by the wild-type, deletion, and complemented strains.

Dishes were washed, stained with crystal violet, and photographed from above. (C) The crystal

violet was extracted with 95% ethanol, and the absorbance measured at 595 nm. Means and

standard deviations were calculated from results of three separate assays. Cultures were grown in

6-cm-diameter petri dishes in liquid L broth (Miller), supplemented with 10 mM CaCl$_2$ and

incubated without shaking at 30 ºC. Cells were harvested at mid-log phase for the β-galactosidase

assay or left undisturbed for 72 hours for the biofilm assay. Asterisks on the bars indicate a

significant difference compared to the wild-type according to Student's t-test (**$P \leq 0.01$, ***P

≤ 0.001).

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7