1

4 DksA, ppGpp and RegAB regulate nitrate respiration in *Paracoccus denitrificans*

6 Ashvini Ray and Stephen Spiro*

8 Department of Biological Sciences, University of Texas at Dallas, Richardson,

9 Texas 75080.

11 *For correspondence: stephen.spiro@utdallas.edu

Running title: Roles of *dksA*, *relA/spoT* and *regAB* in *P. denitrificans*

Keywords: DksA, guanosine tetraphosphate, nitrate respiration, *Paracoccus*

denitrificans, redox homeostasis, RegAB

ABSTRACT

The periplasmic (NAP) and membrane-associated (Nar) nitrate reductases of *Paracoccus denitrificans* are

responsible for nitrate reduction under aerobic and anaerobic conditions, respectively. Expression of

NAP is elevated in cells grown on a relatively reduced carbon and energy source (such as butyrate); it is

believed that NAP contributes to redox homeostasis by coupling nitrate reduction to the disposal of

excess reducing equivalents. Here we show that deletion of either *dksA1* (encoding one of two *dksA*

homologs in the *P. denitrificans* genome) or *relA/spoT* (encoding a bifunctional ppGpp synthetase and

hydrolase) eliminates the butyrate-dependent increase in *nap* promoter and NAP enzyme activity. We

conclude that ppGpp likely signals growth on a reduced substrate and, together with DksA1, mediates

increased expression of the genes encoding NAP. Support for this model comes from the observation that

nap promoter activity is increased in cultures exposed to a protein synthesis inhibitor that is known to

trigger ppGpp synthesis in other organisms. We also show that, under anaerobic growth conditions, the

redox-sensing RegAB two-component pair acts as a negative regulator of NAP expression and as a

positive regulator of expression of the membrane-associated nitrate reductase Nar. The *dksA1* and

relA/spoT genes are conditionally synthetically lethal; the double mutant has a null phenotype for growth

on butyrate and other reduced substrates while growing normally on succinate and citrate. We also show

that the second *dksA* homolog (*dksA2*) and *relA/spoT* have roles in regulation of expression of the

flavohemoglobin Hmp and in biofilm formation.

IMPORTANCE

Paracoccus denitrificans is a metabolically versatile Gram-negative bacterium that is used as a

model for studies of respiratory metabolism. The organism can utilize nitrate as an electron

acceptor for anaerobic respiration, reducing it to dinitrogen via nitrite, nitric oxide, and nitrous

oxide. This pathway (known as denitrification) is important as a route for loss of fixed nitrogen

from soil and as a source of the greenhouse gas nitrous oxide. Thus, it is important to understand

those environmental and genetic factors that govern flux through the denitrification pathway. Here

we identify four proteins and a small molecule (ppGpp) which function as previously unknown

regulators of expression of enzymes that reduce nitrate and oxidize nitric oxide.

INTRODUCTION

Paracoccus denitrificans is a Gram-negative α-proteobacterium and a facultative anaerobe that

can respire using nitrogen oxyanions and oxides as terminal electron acceptors in an anaerobic

process known as denitrification (1). In the first step of denitrification, nitrate is reduced to nitrite

by the membrane-associated nitrate reductase, Nar (2). *P. denitrificans* also expresses a

periplasmic nitrate reductase, NAP, which makes a smaller contribution to the generation of a

proton motive force than does Nar (2). Maximal expression of NAP occurs during aerobic growth

on relatively reduced substrates, and it is thought that NAP facilitates redox balancing by allowing

for the disposal of excess reducing equivalents (3–5).

In the denitrification pathway, the toxic intermediate nitric oxide (NO) is generated from

nitrite by the respiratory nitrite reductase and is subsequently reduced to nitrous oxide (N_2O) by

the respiratory NO reductase. *P. denitrificans* also has an *hmp* gene encoding a flavohemoglobin

(Hmp) that potentially contributes to the metabolism of NO (6, 7). NO acts in conjunction with

transcriptional regulators as an important signal molecule to regulate gene expression during the

transition from aerobic respiration to denitrification (8, 9). Pathogenic strains of *Salmonella* and

E. coli have several transcriptional regulators which control the expression of genes encoding

enzymes that scavenge NO. These regulators and their target genes may help to protect the bacteria

against the NO that is made by host phagocytes (10–12). One such transcriptional regulator is

DksA, which acts as a thiol based NO sensor in *Salmonella* and regulates detoxification of NO

through Hmp in *E. coli* (13–15).

DksA, first described in *E. coli* as DnaK suppressor A (16) is a founder member of

TraR/DksA family of transcriptional regulators (17, 18). DksA$_{Ec}$ (*E. coli* DksA) is 151 amino acids

long and consists of three major domains (Fig. S1): a coiled-coil (CC) domain with a conserved

DXXDXA motif at its tip, a globular domain with a conserved four-cysteine Zn-finger motif, and

a C-terminal α-helical domain (19). The conserved acidic residues located at the tip of the CC

domain in DksA$_{Ec}$ are critical for binding to RNA polymerase (RNAP) and for DksA$_{Ec}$ to function

as a transcriptional regulator (20, 21). The globular domain has been shown to interact with the

rim of the secondary channel of RNAP (22), and the conserved redox-sensitive cysteines act as a

sensor for NO in *Salmonella* (13). Some bacterial genomes encode multiple homologs of DksA,

which differ in the presence or absence of a complete DXXDXA motif or the presence of one, two,

or four conserved cysteines in the Zn-finger motif (23). For instance, *Pseudomonas aeruginosa*

has two functional homologs, DksA1 and DksA2 (24). DksA1 has the conserved four cysteine Zn-

finger motif whereas DksA2 does not (25, 26). In α-proteobacteria more closely related to *P.*

denitrificans, such as *Sinorhizobium meliloti* and *Rhodobacter sphaeroides*, the assigned *dksA*

gene encodes a protein with a complete DXXDXA motif and only one cysteine in the Zn-finger

motif (27, 28). Deletion or overexpression of *dksA* in various species has pleiotropic effects on

multiple physiological processes. These include amino acid biosynthesis (29), motility (30),

biofilm formation (30), plant symbiosis (27), DNA repair (31), quorum sensing and virulence (32)

and oxidative and nitrosative damage (13, 15).

DksA, along with the alarmone guanosine-5', 3'-tetraphosphate (ppGpp) is known to alter

global gene expression to adjust cellular metabolism in response to nutritional stress, a mechanism

known as the stringent response (when triggered by amino acid starvation) in *E. coli* and other related γ-proteobacteria (33–35). ppGpp is found throughout the bacterial kingdom as a second messenger and is associated with various stress response mechanisms and with growth rate regulation (33). The metabolism of ppGpp is controlled by two enzymes encoded by *relA* and *spoT* (36, 37). RelA is a ribosome-associated ppGpp synthetase, whereas SpoT is a bifunctional enzyme responsible for both synthesis and hydrolysis of ppGpp. The presence of both *relA* and *spoT* is characteristic of γ-proteobacteria, while only one RelA/SpoT bifunctional homolog (RSH) possessing both conserved ppGpp synthetase and hydrolase domains is typically present in α-proteobacteria (2, 38). While DksA and ppGpp coregulate many genes, some phenotypic and transcriptomic studies report independent effects of both DksA and ppGpp (39, 40). The independent effects of DksA and ppGpp might be attributed to their direct binding to RNAP; ppGpp is also a direct effector of a number of metabolic enzymes (41–43)

In *P. denitrificans* Pd1222, three gene products have been annotated as members of the TraR/DksA family: Pden_0547 (which we designate DksA1$_{Pd}$), Pden_0916 (DksA2$_{Pd}$), and Pden_1862, which are 45%, 26% and 30% identical to *E. coli* DksA, respectively. The *P. denitrificans* proteins are more similar to their homologs from related α-proteobacteria. DksA1$_{Pd}$, DksA2$_{Pd}$, and Pden_1862 are 90%, 64%, and 33% identical to DksA$_{Rsp}$ (*R. sphaeroides* DksA), respectively, while they are 64%, 47%, and 36% identical to DksA$_{Sm}$ (*S. meliloti* DksA), respectively. *P. denitrificans* also has a single *relA/spoT* gene: Pden_1400 (henceforward called *rel*) which encodes for a RelA/SpoT bifunctional homolog (RSH). The predicted RSH is 705 amino acids long and possesses a phosphohydrolase HD domain with a conserved HDXXED motif in the N-terminal region, a RelA-SpoT region, and conserved TGS and ACT domains in the C-terminal domain (Fig. S2). On the basis of sequence similarity, Pden_1400 is predicted to have

both ppGpp synthetase and hydrolase activities. None of these *P. denitrificans* genes and proteins has been functionally characterized. A proteomic analysis addressing aerobic to anaerobic growth adaptation in *P. denitrificans* demonstrated a significant increase in expression of DksA1$_{Pd}$ (A1AZG5) under anaerobic conditions. Another study using microarray analysis showed a 3.4-fold higher expression of *dksA2* during anaerobic growth (44, 45). Our attention was drawn to the DksA homologs of *P. denitrificans* because they are differentially expressed under denitrifying conditions, and by the report that DksA regulates *hmp* expression in *E. coli* (15). Our goal for the work described here was therefore to use a reverse-genetics approach in an effort to identify physiological roles for the two DksA proteins and for RSH.

In this work, we report distinct phenotypes associated with deletion of two of the *dksA*-like genes and the *relA/spoT* homolog in *P. denitrificans* Pd1222. We show that deletion of *dksA1* causes an aerobic growth defect on reduced carbon substrates such as butyrate and that *dksA1* and *rel* null strains fail to up-regulate expression and activity of the periplasmic nitrate reductase NAP under these conditions. Furthermore, we demonstrate that a *dksA1 rel* double mutant is unable to grow on reduced carbon substrates. In addition, we show that *dksA2* is involved in biofilm production, possibly indirectly via negative regulation of *hmp* transcription.

RESULTS

***P. denitrificans* dksA1 and dksA2 complement Δ*dksA*$_{Ec}$ strains.**

Sequence alignment of the three DksA/TraR proteins revealed that two (Pden_0547 and Pden_0916) have a complete DXXDXA motif which is required for interaction with RNA polymerase (20, 21) while the third (Pden_1862) does not. For this reason, our attention has focused on Pden_0547 and Pden_0916, which we have designated *dksA1* and *dksA2*, respectively.

139   Unlike the canonical DksA of *E. coli*, DksA1$_{Pd}$ has only one cysteine conserved in the zinc finger

140   motif whereas DksA2$_{Pd}$ has all four cysteines (Fig. S1).

141   *E. coli* MG1655 deleted for *dksA* was complemented with either *dksA1* or *dksA2* cloned in

pIND4 and expressed from their native promoter. *E. coli* Δ*dksA* is auxotrophic for multiple amino

acids and fails to grow on M9-glucose medium without amino acid supplementation (15).

Expression of either *dksA1* or *dksA2* rescued growth of Δ*dksA$_{EC}$* on M9-glucose plates suggesting

that either gene may complement the *E. coli dksA* mutation (Fig. 1).

Deletion of *dksA1$_{Pd}$* impairs nitrate stimulation of aerobic growth on butyrate

To characterize the *dksA*-like genes of *P. denitrificans*, we constructed and tested the growth of

deletion strains lacking *dksA1* or *dksA2*. Each mutant grew as well as the parent strain on rich (LB)

medium under both aerobic and anaerobic conditions (Fig. S3). We also did not observe amino

acid auxotrophy when the mutants were grown in succinate or glucose minimal medium (data not

shown). In *Salmonella enterica* sv. *typhimurium*, DksA is involved in redox homeostasis since it

regulates the expression of enzymes from central metabolic pathways that generate reduced

pyridine nucleotides (46). *P. denitrificans* may experience redox imbalance during growth on

reduced carbon substrates due to the generation of excess reducing equivalents (3). Therefore, we

assayed aerobic growth in minimal medium supplemented with succinate (MM-S) or butyrate

(MM-B) as sole carbon and energy sources that are relatively oxidized and reduced, respectively.

Growth of the wild-type strain in MM-B was significantly enhanced by the addition of nitrate to

the medium. Reduction of nitrate by the periplasmic nitrate reductase NAP provides a mechanism

to dispose of excess reducing equivalents during growth on reduced carbon substrates (4). The

Δ*dksA1* mutant grew normally on MM-S but showed slightly impaired growth on MM-B relative

to the wild type and growth on MM-B was not improved by addition of nitrate to the medium

(Fig. 2A and B). Expression of *dksA1* from a recombinant plasmid in the *dksA1* mutant restored a

wild type phenotype for growth on MM-B both in the presence and absence of nitrate (Fig. S4).

These phenotypes suggest that DksA1$_{Pd}$ may regulate aerobic nitrate reduction by NAP during

growth on reduced carbon substrates.

To determine if the Δ*dksA1* associated phenotype is ppGpp-dependent, we measured

growth of the Δ*rel* strain under similar conditions. The Δ*rel* mutant showed significantly reduced

growth in MM-B compared to wild type, however, unlike for Δ*dksA1*, growth was improved by

addition of nitrate to the medium (Fig. 3 and S5), though the magnitude of nitrate stimulation was

less than was observed in the wild-type strain. Complementation of the Δ*rel* mutant with the *rel*

gene cloned in a plasmid restored the wild type growth phenotypes on MM-B (Fig. S6).

Δ*dksA1* and Δ*rel* mutants have reduced *napE* promoter activity and NAP enzyme activity

It has been shown previously that the periplasmic nitrate reductase, NAP, has increased expression

and activity in cells grown aerobically on reduced energy sources (3, 5). Our results suggest that

the absence of *dksA1* impairs nitrate reduction in *P. denitrificans* during aerobic growth. We

hypothesized that DksA1$_{Pd}$ regulates expression of NAP during aerobic growth on reduced energy

sources. To test this hypothesis, we constructed a *napE-lacZ* promoter transcriptional fusion and

measured *napE* promoter activity using β-galactosidase as the reporter. Consistent with previous

reports (5), we observed a ~3-fold increase in *napE* promoter activity in cells grown aerobically in

MM-B compared to those grown in MM-S (Fig. 4A). The increase in *napE* promoter activity

caused by growth on butyrate was eliminated in the Δ*dksA1* mutant. The increase in *napE* promoter

activity in MM-B versus MM-S also disappeared when the cultures were grown anaerobically

(Fig. 4A). To confirm that these observations were due to loss of *dksA1,* we measured *napE*

promotor activity in the complemented Δ*dksA1* strains. Complementation with the *dksA1* gene

restored the butyrate stimulation of *napE* promotor activity (Fig. S7)

In parallel with the *napE* promoter activity assays we also measured the activity of NAP in

periplasmic fractions using reduced methylviologen as the electron donor to the enzyme. As

previously reported (3), and consistent with our promoter activity data, NAP activity in the wild-

type was significantly (~4-fold) higher during aerobic growth in MM-B compared to MM-S (Fig.

4B). The stimulation of activity by growth on butyrate disappeared in the Δ*dksA1* mutants and in

cells grown anaerobically in either MM-S or MM-B (Fig. 4B). Our results suggest that DksA1$_{Pd}$

upregulates (either directly or indirectly) the *napE* promoter during aerobic growth on reduced

carbon substrates such as butyrate.

The *rel* mutant was found to have the same phenotype as the *dksA1* mutant with respect to

nap promoter activity and NAP enzyme activity. That is, the butyrate-dependent stimulation was

also lost in the *rel* mutant (Fig. 4 and S7). Since DksA and ppGpp frequently act in concert by

binding together to a single site on RNA polymerase (34), the likely explanation that DksA-

dependent regulation of *nap* requires ppGpp.

The RegAB system regulates anaerobic *napE* promoter activity and NAP enzyme activity

In *P. pantotrophus*, regulation of the *nap* operon in response to carbon substrate oxidation state

involves *cis*-acting regulatory sequences in the *napE* promoter region (48). Specifically, a

heptameric imperfect inverted repeat sequence (TG<u>A</u>G<u>A</u>C<u>A</u>tttT<u>GTCGC</u>A) was suggested to be

the binding site for a negatively acting redox-responsive regulator (48). The inverted repeat is a

partial match to the binding site (GCGGC-N$_5$-GTCGC) for the response regulator RegR (RegA)

from *Bradyrhizobium japonicum* (49). We found a similar sequence (5′-<u>GA</u>GG<u>C</u>gtttt<u>GA</u>C<u>GC</u>-3′)

in the promoter region of the *nap* operon in *P. denitrificans* Pd1222 (matches to the

experimentally-determined sequence from *B. japonicum* are underlined). Therefore, we examined

napE promoter activity in Δ*regA* and Δ*regB* strains to determine if *nap* is regulated by RegAB

alongside *dksA1*. In the Δ*regA* mutant, we observed ~2-fold increases in *napE* promoter activity

in cells grown aerobically on succinate (Fig. 5A). In cells grown aerobically on butyrate, *napE*

promoter activity was similar in both *regA* and *regB* mutants and the wild-type strain. Interestingly,

there were significant increases in promoter activities in *regA* and *regB* mutants grown

anaerobically on either succinate or butyrate, compared to the wild type under the same conditions

(Fig. 5A). Expression of *regA* and *regB* in the Δ*regA* and Δ*regB* strains respectively reduced *napE*

promoter activity to levels below those seen in the wild type strain, confirming that RegAB acts

negatively on the *napE* promoter (Fig. S8).

Our data suggest that RegA is the negatively acting regulator that Ellington et al. suggested

(48) binds to the inverted repeat sequence in the *napE* promoter, and that the RegAB system acts

to down-regulate this promoter principally under anaerobic growth conditions (but also to a lesser

extent during aerobic growth on succinate). The less severe phenotype of the *regB* mutant under

some conditions suggests the possibility that RegA is phosphorylated by another histidine kinase

(or by a low molecular weight phospho-donor) in the absence of RegB. Assays of NAP activity in

periplasmic fractions prepared from anaerobically grown cells confirm that RegAB functions as a

negative regulator under these conditions (Fig. 5B).

The membrane-associated nitrate reductase, Nar, is upregulated in *P. denitrificans* in

oxygen-limiting conditions (1, 50). Known regulators of the *nar* promoter are NarR (50) and,

probably, FnrP (51). Assays of Nar activity in membrane fractions revealed a reduced activity in

both Δ*regA* and Δ*regB* mutants suggesting positive regulation of the *nar* operon by RegAB under

these conditions (Fig. 5C). Addition of 20 μM sodium azide (a selective inhibitor of Nar) to the

reaction mixture inhibited nitrate reduction confirming that the reductase activity in the membrane

fraction was due to Nar (data not shown). Reduced Nar activity also coincided with reduced growth

yield for the $\Delta regA$ mutant when grown anaerobically (Fig. S9). Thus, the RegAB system mediates

reciprocal regulation of the two nitrate reductases under anaerobic conditions, being a negative

regulator of NAP expression and a positive regulator of Nar expression.

The *dksA1* and *rel* genes are synthetically lethal for growth on reduced substrates

The DksA protein interacts with ppGpp to regulate several functions and deletion of either *dksA*

or *rel* gives rise to overlapping pleiotropic phenotypes (39). The similar phenotypes of $\Delta dksA1$ and

rel mutants with respect to loss of butyrate stimulation of *nap* promoter activity suggests that

DksA1$_{Pd}$ and ppGpp act in concert to regulate expression of the periplasmic nitrate reductase. A

$\Delta dksA1\Delta rel$ double mutant exhibited a complete growth defect on reduced carbon substrates such

as butyrate, caproate and choline (Fig. 3, Fig. S10), while growth on the relatively oxidized

substrates succinate and citrate was normal (Fig. S11). Expression of either *dksA1* or *rel* on a broad

host range plasmid from their native promoters restored growth of the $\Delta dksA1\Delta rel$ mutant,

confirming that both mutations are required for the null growth phenotype (Fig. S12). Thus, *dksA1*

and *rel* exhibit synthetic lethality with respect to growth on reduced substrates. The likely

explanation is that two (or more) genes that are independently regulated by DksA1$_{Pd}$ and ppGpp

are together required for growth on reduced substrates.

Partial activation of the *napE* promoter by a protein synthesis inhibitor

In *E. coli*, ppGpp synthesis by RelA can be stimulated by manipulations that reduce the extent of

aminoacyl tRNA charging, including amino acid limitation or inhibition of an amino acyl tRNA

synthetase (52). For example, serine hydroxamate (SHX) is a seryl-tRNA synthetase inhibitor that

triggers ppGpp synthesis (52–54). SHX (1.28 mM) inhibits *E. coli* growth, and inhibition can be

reversed by exogenous serine, presumably because serine overcomes competitive inhibition

mediated by SHX (53). Addition of 3 mM SHX to MM-S caused a prolonged growth inhibition

of wild type *P. denitrificans* that was reversed by the addition of casamino acids to the medium

(Fig. 6). The physiological basis for the eventual recovery of the culture is not known but it may

be ppGpp-dependent since the *rel* mutant did not recover from the SHX treatment (Fig. 6). The

simplest interpretation of these observations is that SHX treatment triggers ppGpp synthesis by

RSH and that ppGpp is required for the recovery from growth arrest. A prediction that follows is

that treatment of a culture with a sub-inhibitory concentration of SHX will stimulate a promoter

activated by ppGpp. Therefore, we measured activity of the *napE-lacZ* fusion in cultures grown

aerobically in MM-S and treated with 2 mM SHX while in the logarithmic phase of growth. Ninety

minutes after SHX addition, promoter activity in the treated cultures was ~2-fold higher than in

the untreated cultures (1101 ± 96 versus 609 ± 60 units of β-galactosidase activity). In the *rel*

mutant, we saw no induction of the *napE* promoter in response to 2 mM SHX. Partial induction of

the *napE* promoter by a treatment known to trigger ppGpp synthesis in other organisms is

consistent with the idea that the promoter is regulated by ppGpp and DksA1.

DksA2$_{Pd}$ is a positive regulator of the *hmp* promoter

In *E. coli*, *dksA* regulates the expression and activity of Hmp, a flavohemoprotein. Hmp is a nitric

oxide dioxygenase or denitrosylase which oxidizes nitric oxide to nitrate, and is known to be the

dominant NO defense mechanism under aerobic conditions (15). *P. denitrificans* has an *hmp* gene,

Pden_1689 (6, 7) and we have shown that the Hmp is functional for NO scavenging (unpublished

data). To determine if regulation of *hmp* is mediated by DksA in this organism, we assayed *hmp*

promoter activity in *P. denitrificans* and the Δ*dksA* mutants. Our results indicate that *hmp* promoter

activity is modestly but significantly elevated in the Δ*dksA2* mutant but is unaffected by the

Δ*dksA1* deletion (Fig. 7A). The Δ*rel* mutant also shows higher *hmp* promoter activity similar to

the Δ*dksA2* strain suggesting that the regulation of *hmp* by DksA2 is ppGpp-dependent (Fig. 7A).

Both DksA2_{Pd} and RSH regulate biofilm formation in *P. denitrificans*

Our previous work has shown that deletion of *hmp* increases biofilm formation in *P. denitrificans*

Pd1222 possibly due to accumulation of NO (6). Given the evidence that DksA2_{Pd} regulates *hmp*

expression in *P. denitrificans*, we assayed attached growth to determine if DksA2_{Pd} also affects

biofilm formation. We observed that Δ*dksA2* strains had greatly reduced levels of attached growth

(Fig. 7B and C). Addition of an *hmp* mutation to the *dksA2* mutant rescued biofilm formation to

wild-type levels (Fig. 7B and C) suggesting that the *dksA2* phenotype requires Hmp. Deletion of

nsrR, which encodes a transcriptional repressor of *hmp* (6), led to significantly reduced biofilm

formation similar to Δ*dksA2* (Fig. 7B and C). We also observed reduced biofilm in a Δ*rel* strain

(and in a Δ*dksA2* Δ*rel* double mutant, not shown) suggesting that *dksA2* and *rel* could be jointly

involved in regulating biofilm formation in *P. denitrificans* via negative regulation of *hmp* (Fig.

7B and C).

DISCUSSION

Expression of the periplasmic nitrate reductase NAP is upregulated by aerobic growth on reduced

carbon and energy sources, the likely explanation being that nitrate reduction by NAP provides a

means to dispose of excess reducing equivalents (3–5, 47, 48). We have shown that DksA1_{Pd} and

RSH are required for the butyrate-stimulation of *napE* promoter and NAP enzyme activity. RSH

is the only protein encoded in the *P. denitrificans* genome with predicted ppGpp synthetase activity

and that it is enzymatically functional is supported by the detection of ppGpp in slow-growing

cultures of this organism (55). It is likely that DksA1_{Pd} and ppGpp directly activate the *napE*

promoter, although indirect effects cannot be excluded. In either case, the implication is that

growth on reduced substrates somehow triggers the synthesis of ppGpp by RSH. Data from

continuous cultures show that the *napE* promoter is regulated both by growth rate and by carbon

substrate, leading to the conclusion that "…*nap* regulation is affected by a physiological signal

which integrates aspects of growth rate and carbon substrate utilization" (56). Our work suggests

that ppGpp (together with DksA1$_{Pd}$) is the integrating signal, which is consistent with the well-

known role of ppGpp in growth rate control (33).

One difference between the phenotypes of the *dksA1* and *rel* mutants is that nitrate does

not improve aerobic growth of the *dksA1* mutant on butyrate (as it does in the case of the wild-

type strain) but does modestly improve growth of the *rel* mutant. This improvement of growth

cannot be attributed to butyrate stimulation of *nap* expression, since that is lost in the *rel* mutant.

It is also the case that the *rel* mutant grows less well on butyrate (in the absence of nitrate) than

the wild-type strain and the *dksA1* mutant, and that the double mutant does not grow at all on

reduced substrates. It is likely that there are DksA1$_{Pd}$ and ppGpp regulated genes that are involved

in redox homeostasis that are yet to be identified.

We showed that regulation of *nap* expression also involves the RegAB two-component

system, which is redox-sensing and involved in redox homeostasis (57). Our results show that

RegAB is not a major player in mediating up-regulation of the *nap* promoter by growth on butyrate,

but instead negatively regulates *nap* expression during anaerobic growth. The response regulator

RegA very likely binds to an inverted repeat in the *napE* promoter that was initially characterized

in the closely related organism *P. pantotrophus* and suggested to be the binding site for a

negatively acting regulator (48). Evidently, regulation of the *napE* promoter by carbon substrate

and by oxygen availability can be separated, with the former requiring DksA1$_{Pd}$ and ppGpp and

the latter RegAB. Interestingly, RegAB also contribute (either directly or indirectly) to the

upregulation of the membrane associated nitrate reductase under anaerobic growth conditions.

Thus, NAP and Nar are reciprocally regulated by oxygen availability to be the dominant nitrate

reductases under aerobic and anaerobic conditions, respectively.

A recent study has shown that RegAB acts both positively and negatively under hypoxic

conditions in *Burkholderia pseudomallei*, an α-proteobacterium quite closely related to *P.*

denitrificans (58). Specifically, RegAB are required for upregulation of Nar under hypoxic

conditions in this organism, though this may involve both direct regulation by RegA and indirect

regulation, via FNR (58). Indirect effects of RegAB via NarR and/or FnrP are also possible in *P.*

denitrificans; we are able to discern a potential RegA binding site in the *narR-narK* intergenic

region that controls expression of the *narGHJI* genes encoding Nar (50).

In *E. coli*, DksA and ppGpp contribute to regulation of the *hmp* gene encoding the

flavohemoglobin, Hmp, which is responsible for detoxifying nitric oxide (15, 59, 60).   In *P.*

denitrificans, we find that deletion of *dksA2* or *rel* leads to modestly increased *hmp* promoter

activity and dramatically reduced ability to form biofilm. Since the biofilm-defective phenotype

requires Hmp, we suggest that reduced expression of *hmp* in the mutants causes a reduced

accumulation of nitric oxide, which stimulates biofilm formation in this organism (6). Reduced

growth as biofilm in *dksA* mutants has been reported for other organisms, though the regulatory

mechanisms are likely different (30, 61).

Our study has identified two functional DksA proteins in *P. denitrificans* that have distinct

roles in growth on reduced carbon substrates, aerobic and anaerobic nitrate respiration, expression

of flavohemoglobin, and biofilm formation. In other α-proteobacteria which have at least two

DksA homologs, physiological roles have so far been ascribed to only one of them (27, 28). Further

functional analysis of the *P. denitrificans* DksA-like proteins and of the role of ppGpp will advance

our broader understanding of DksA functions in α-proteobacteria.

MATERIALS AND METHODS

Bacterial strains and growth conditions. The bacterial strains and plasmids used are listed in

Table S1. All *Paracoccus denitrificans* strains were derived from the wild-type Pd1222 (62).

Escherichia coli S.17-1 was used as the donor and *E. coli* DH5α (pRK2013) as the helper strain

for the introduction of plasmids into *P. denitrificans* by conjugation (63, 64). *E. coli* Novablue

competent cells (Novagen) were used as the host for all genetic manipulations. The rich medium

used for routine growth was LB (Miller) Broth (10 g/l tryptone, 5 g/l yeast extract, 10 g/l NaCl),

and the defined minimal medium (MM) for *P. denitrificans* was as previously described (65),

solidified with 1.5 % agar (w/v) as needed and supplemented with carbon and energy sources and

with 20 mM nitrate as indicated. For growth assays, *P. denitrificans* strains were first streaked on

LB agar plates, single colonies were inoculated into LB broth and grown overnight in 15 ml tubes

at 250 rpm. These cells were washed with MM, inoculated into succinate-supplemented MM, and

grown overnight in 15 ml tubes at 250 rpm. Cells were pelleted, washed, and finally inoculated

into MM supplemented with different carbon sources with or without nitrate. For aerobic growth

assays, cultures were grown in triplicate wells in a BioTek plate reader with horizontal orbital

shaking at 280 rpm and a volume of 0.2 ml (<50% of total volume) to ensure proper aeration while

preventing cross contamination. The optical density was measured at 600 nm and adjusted for path

length variation. Serine hydroxamate was added directly from a freshly made 100 mM stock to the

growth medium to a final concentration of 2-3 mM either at the time of inoculation or to a growing

culture, as indicated.

Anaerobic cultures for β-galactosidase assays were grown in filled glass cuvettes sealed with a cap and septum, with medium supplemented with 20 mM nitrate. The cuvettes were gassed with N_2 and incubated statically. Aerobic cultures were grown in 250 ml flasks shaking at 250 rpm in MM supplemented with the indicated carbon substrate and with or without 20 mM nitrate. For *E. coli* complementation assays, solid M9 medium (66) with 10 mM glucose was used, with or without 0.2 % (w/v) casamino acids.

Plasmids were maintained by addition of antibiotics at the following concentrations: ampicillin 100 µg/ml, kanamycin 25 µg/ml, rifampicin 100 µg/ml, tetracycline 2 µg/ml for *P. denitrificans* and 15 µg/ml for *E. coli*. All *P. denitrificans* strains were grown at 30 ºC and *E. coli* at 37 ºC.

Genetic manipulations. All primers used are listed in Table S2. Unmarked deletions were made by allelic replacement using pK18*mobsacB* as described by Sullivan et al. with minor modifications (67, 68). Briefly, in-frame marker-less deletions were constructed by PCR amplification of ~500-1000 bp fragments from either side of the gene of interest. The two amplicons were ligated together either by PCR using overlapping regions added to the primers or by digestion and ligation using a restriction site incorporated in the primers (69). The ligated fragment was first cloned into pSTBlue-1 (Novagen), then sub-cloned into pK18*mobsacB*, transformed into *E. coli* S.17-1 and mobilized into *P. denitrificans* by conjugation. Triparental mating was performed by mixing *P. denitrificans*, *E. coli* S.17-1 harboring recombinant pK18*mobsacB* or pMP220 or pIND4 derivatives, and *E. coli* DH5α (pRK2013) cells grown to log phase, in a 2:1:1 ratio and by incubation on LB agar plate for 2-3 days at 30 ºC. Single-crossover recombination events were selected on rifampicin and kanamycin plates, and then recombinants with double crossovers were isolated by growth on 6 % (w/v) sucrose plates and kanamycin

sensitivity as described by Schafer et al. (67). All mutations were designed to be in-frame deletions

to prevent polarity. Promoter-*lacZ* reporter fusions were constructed by amplifying ~300-400 bp

upstream of the start codon of genes of interest and first cloning into pSTBlue-1. The sequence

was verified and the fragment cloned into the low copy number vector pMP220 between *Eco*RI

and *Pst*I restriction sites (70). Plasmids were introduced into *P. denitrificans* by conjugation from

E. coli S.17-1. *P. denitrificans* exconjugants were selected by rifampicin and tetracycline

resistance. For complementation tests in *P. denitrificans*, the gene of interest with its native

promoter was amplified and cloned directly into pIND4 (71) and the sequence verified. The

cloning was done using the NEBuilder HiFi DNA assembly cloning kit (NEB). Plasmids were

transformed into *E. coli* S.17-1 and mobilized into *P. denitrificans* strains by conjugation as

described above.

β-galactosidase assay: Liquid cultures were grown aerobically and anaerobically in minimal

medium with the indicated carbon sources and with or without 20 mM nitrate. Cells were harvested

at the mid-log phase of growth and the assay was performed as described by Miller (66). β-

galactosidase activity was measured using 0.1 ml of cells made permeable by treatment with

sodium dodecyl sulfate and chloroform. In the case of the *hmp-lacZ* fusion, strains were grown to

mid-log phase in the same medium and growth conditions as used for the biofilm assays.

Biofilm assays: Attached growth was assayed in 60 mm x 15 mm polystyrene petri dishes (6). A

70 µl aliquot of overnight culture was inoculated into 7 ml of LB broth supplemented with 10 mM

calcium chloride. The Petri dishes were incubated statically at 30 ºC for 72 hours. The plates were

then washed thrice under gently running tap water and stained with crystal violet (0.1 % in water)

for 20 minutes, washed three times with water, and air dried. For quantitation, the crystal violet

stain was extracted into 3 ml 95 % ethanol, diluted as necessary, and absorbance measured at 595

nm (6, 72).

Cell fractionation: *P. denitrificans* strains were grown aerobically and anaerobically in MM with

either succinate (10 mM) or butyrate (10 mM) as the sole carbon source; nitrate (20 mM) was

added as indicated. Aerobic cultures (50 ml) were grown at 30 ºC in 250 ml conical flasks shaken

at 250 rpm. Anaerobic cultures were grown in 100 ml bottles filled to the neck with medium and

incubated at 30 ºC without shaking. Periplasmic fractions for the nitrate reductase assay were

prepared as described in Quan et al (73) . In short, mid-exponential phase cells were harvested by

centrifugation at 3,000 g for 20 minutes at 4 ºC. Cells were then gently resuspended in TSE buffer

(200 mM Tris–HCl, pH 8.0, 500 mM sucrose, 1 mM EDTA without any protease inhibitor) and

incubated on ice for 30 minutes, and centrifuged at 16,000 g for 30 minutes at 4 ºC. The supernatant

was transferred to a fresh tube and used as the periplasmic fraction. Membrane fractions were

prepared according to Dunn et al. with some modifications (74). Briefly, cultures were harvested

in the mid-exponential phase of growth by centrifugation at 8,000 g for 5 minutes at 4 ºC and

washed twice in 10 mM potassium phosphate buffer (pH 7.5) in 50 ml Falcon tubes. Cells were

resuspended in 2.5 ml of ice cold 10 mM phosphate buffer, 10 mM $MgSO_4$, 0.2 M NaCl. Samples

were sonicated on ice using a Fisher Scientific Sonic Dismembrator model 500 outfitted with a 3

mm diameter tip at 20 % amplitude (10 W) for five 20 second bursts at 1 minute intervals. To

clarify the lysate, samples were centrifuged at 16,000 g for 20 minutes at 4 ºC. The clarified total

lysate was further centrifuged at 100,000 g for 2 hours at 4 ºC. The pellet was suspended in 0.2 ml

of phosphate buffer and retained as the membrane fraction for the enzyme assay. Fractions were

kept on ice and used within hours of preparation.

Enzyme activity: Nitrate reductase activity was determined spectrophotometrically using reduced

methylviologen (MV^+) as the electron donor and 10 mM sodium nitrate as the electron acceptor

(3). The extinction coefficient of reduced methylviologen used was 8.25 $mM^{-1} cm^{-1}$ at 600 nm (75).

The protein content of the periplasmic and membrane fractions was measured by the protein

estimation method of Lowry et al., using bovine serum albumin as the standard (76).

ACKNOWLEDGMENTS

We are grateful to Victor Luque-Almagro, Andrew Gates and David Richardson for generously

providing the *regA*::*kan* mutant and to Aashna Pathi for her contribution to strain construction.

1.   Baker SC, Ferguson SJ, Ludwig B, Page MD, Richter OH, van Spanning RJM. 1998.

Molecular genetics of the Genus *Paracoccus*: Metabolically versatile bacteria with

bioenergetic flexibility. *Microbiol Mol Biol Rev* 62: 1046–1078.

2.   Richardson DJ, Berks BC, Russell DA, Spiro S, Taylor CJ. 2001. Functional, biochemical

and genetic diversity of prokaryotic nitrate reductases. *Cell Molecular Life Sci* 58: 165–178.

3.   Sears HJ, Ferguson SJ, Richardson DJ, Spiro S. 1993. The identification of a periplasmic

nitrate reductase in *Paracoccus denitrificans*. *FEMS Microbiol Lett* . 113: 107-112.

4.   Sears HJ, Spiro S, Richardson DJ. 1997. Effect of carbon substrate and aeration on nitrate

reduction and expression of the periplasmic and membrane-bound nitrate reductases in

carbon-limited continuous cultures of *Paracoccus denitrificans* Pd1222. *Microbiology* 143:

3767–3774.

5. Sears HJ, Sawers G, Berks BC, Ferguson SJ, Richardson DJ. 2000. Control of periplasmic

nitrate reductase gene expression (*napEDABC*) from *Paracoccus pantotrophus* in response

to oxygen and carbon substrates. *Microbiology* 146: 2977-2985.

6. Kumar S, Spiro S. 2017. Environmental and genetic determinants of biofilm formation in

Paracoccus denitrificans. mSphere 2: e00350-17.

7. Hartop KR, Sullivan MJ, Giannopoulos G, Gates AJ, Bond PL, Yuan Z, Clarke TA,

Rowley G, Richardson DJ. 2017. The metabolic impact of extracellular nitrite on aerobic

metabolism of *Paracoccus denitrificans*. *Water Res* 113: 207–214.

8. Hassan J, Bergaust LL, Wheat ID, Bakken LR. 2014. Low probability of initiating *nirS*

transcription explains observed gas kinetics and growth of bacteria switching from aerobic

respiration to denitrification. PLoS Comput Biol 10: e1003933.

9. Hassan J, Bergaust LL, Molstad L, de Vries S, Bakken LR. 2016. Homeostatic control of

nitric oxide (NO) at nanomolar concentrations in denitrifying bacteria - modelling and

experimental determination of NO reductase kinetics in vivo in *Paracoccus denitrificans*.

Environ Microbiol 18: 2964–2978.

10. Fang FC. 1997. Perspectives series: host/pathogen interactions. Mechanisms of nitric oxide-

related antimicrobial activity. J Clin Invest 99: 2818–2825.

11. Henard C, Vazquez-Torres A. 2011. Nitric oxide and *Salmonella* pathogenesis. *Front*

Microbiol 2: 84.

12. Hyduke DR, Jarboe LR, Tran LM, Chou KJY, Liao JC. 2007. Integrated network analysis

identifies nitric oxide response networks and dihydroxyacid dehydratase as a crucial target

in *Escherichia coli*. Proc Natl Acad Sci USA 104: 8484–8489.

13. Henard CA, Tapscott T, Crawford MA, Husain M, Doulias PT, Porwollik S, Liu L,

McClelland M, Ischiropoulos H, Vázquez-Torres A. 2014. The 4-cysteine zinc-finger motif

of the RNA polymerase regulator DksA serves as a thiol switch for sensing oxidative and

nitrosative stress. Mol Microbiol 91: 790–804.

14. Crawford MA, Henard CA, Tapscott T, Porwollik S, McClelland M, Vázquez-Torres A.

2016. DksA-dependent transcriptional regulation in *Salmonella* experiencing nitrosative

stress. *Front Microbiol* 7: 444.

15. Chou WK, Brynildsen MP. 2019. Loss of DksA leads to multi-faceted impairment of nitric

oxide detoxification by *Escherichia coli*. *Free Rad Biol Med* 130: 288–296.

16. Kang PJ, Craig EA. 1990. Identification and characterization of a new *Escherichia coli*

gene that is a dosage-dependent suppressor of a *dnaK* deletion mutation. *J Bacteriol* 172:

2055-2064.

17. Paul BJ, Barker MM, Ross W, Schneider DA, Webb C, Foster JW, Gourse RL. 2004.

DksA: A critical component of the transcription initiation machinery that potentiates the

regulation of rRNA promoters by ppGpp and the initiating NTP. *Cell* 118: 311–322.

18. Rutherford ST, Lemke JJ, Vrentas CE, Gaal T, Ross W, Gourse RL. 2007. Effects of DksA,

GreA, and GreB on transcription initiation: insights into the mechanisms of factors that

bind in the secondary channel of RNA polymerase. *J Mol Biol* 366: 1243–1257.

19. Vassylyeva MN, Perederina AA, Svetlov V, Yokoyama S, Artsimovitch I, Vassylyev DG.

2004. Cloning, expression, purification, crystallization and initial crystallographic analysis

of transcription factor DksA from *Escherichia coli. Acta Crystallogr D Biol Crystallog* 60:

1611–1613.

20. Perederina A, Svetlov V, Vassylyeva MN, Tahirov TH, Yokoyama S, Artsimovitch I,

Vassylyev DG. 2004. Regulation through the secondary channel - structural framework for

ppGpp-DksA synergism during transcription. *Cell* 118: 297–309.

21. Lennon CW, Ross W, Martin-Tumasz S, Toulokhonov I, Vrentas CE, Rutherford ST, Lee

J-H, Butcher SE, Gourse RL. 2012. Direct interactions between the coiled-coil tip of DksA

and the trigger loop of RNA polymerase mediate transcriptional regulation. *Genes Dev* 26:

2634–2646.

22. Shin Y, Qayyum MZ, Pupov D, Esyunina D, Kulbachinskiy A, Murakami KS. 2021.

Structural basis of ribosomal RNA transcription regulation. *Nat Commun* 12: 528.

23. Furman R, Tsodikov OV, Wolf YI, Artsimovitch I. 2013. An insertion in the catalytic

trigger loop gates the secondary channel of RNA polymerase. *J Mol Biol* 425: 82–93.

24. Min KB, Yoon SS. 2020. Transcriptome analysis reveals that the RNA polymerase-binding

protein DksA1 has pleiotropic functions in *Pseudomonas aeruginosa*. *J Biol Chem* 295:

3851–3864.

25. Blaby-Haas CE, Furman R, Rodionov DA, Artsimovitch I, de Crécy-Lagard V. 2011. Role

of a Zn-independent DksA in Zn homeostasis and stringent response. *Mol Microbiol* 79:

700–715.

26.   Furman R, Biswas T, Danhart EM, Foster MP, Tsodikov OV, Artsimovitch I. 2013. DksA2,

518   a zinc-independent structural analog of the transcription factor DksA. *FEBS Lett* 587: 614–

519   619.

520   27.   Wippel K, Long SR. 2016. Contributions of *Sinorhizobium meliloti* transcriptional regulator

521   DksA to bacterial growth and efficient symbiosis with *Medicago sativa*. *J Bacteriol* 198:

522   1374-1383.

523   28.   Lennon CW, Lemmer KC, Irons JL, Sellman MI, Donohue TJ, Gourse RL, Ross W. 2014.

524   A *Rhodobacter sphaeroides* protein mechanistically similar to *Escherichia coli* DksA

525   regulates photosynthetic growth. *mBio* 5: e01105-14.

526   29.   Paul BJ, Berkmen MB, Gourse RL. 2005. DksA potentiates direct activation of amino acid

527   promoters by ppGpp. *Proc Natl Acad Sci USA* 102: 7823-7828.

528   30.   Azriel S, Goren A, Rahav G, Gal-Mor O. 2016. The stringent response regulator DksA is

529   required for *Salmonella enterica* serovar *Typhimurium* growth in minimal medium,

530   motility, biofilm formation, and intestinal colonization. *Inf Immun* 84: 375-384.

531   31.   Trautinger BW, Jaktaji RP, Rusakova E, Lloyd RG. 2005. RNA polymerase modulators

532   and DNA repair activities resolve conflicts between DNA replication and transcription. *Mol*

533   *Cell* 19: 247–258.

534   32.   Jude F, Köhler T, Branny P, Perron K, Mayer MP, Comte R, van Delden C. 2003.

535   Posttranscriptional control of quorum-sensing-dependent virulence genes by DksA in

536   *Pseudomonas aeruginosa*. *J Bacteriol* 185: 3558-3566.

33.   Hauryliuk V, Atkinson GC, Murakami KS, Tenson T, Gerdes K. 2015. Recent functional

insights into the role of (p)ppGpp in bacterial physiology. *Nat Rev Microbiol* 13: 298–309.

34.   Gourse RL, Chen AY, Gopalkrishnan S, Sanchez-Vazquez P, Myers A, Ross W. 2018.

Transcriptional responses to ppGpp and DksA. *Ann Rev Microbiol* 72: 163–184.

35.   Traxler MF, Summers SM, Nguyen H-T, Zacharia VM, Smith JT, Conway T. 2008. The

global, ppGpp-mediated stringent response to amino acid starvation in *Escherichia coli*.

Mol Microbiol 68: 1128–1148.

36.   Sarubbi E, Rudd KE, Xiao H, Ikehara K, Kalman M, Cashel M. 1989. Characterization of

the *spoT* gene of *Escherichia coli. J Biol Chem* 264: 15074–15082.

37.   Murray DK, Bremer H. 1996. Control of *spoT*-dependent ppGpp Synthesis and degradation

in *Escherichia coli. J Mol Biol* 259: 41–57.

38.   Mittenhuber G. 2001. Comparative genomics and evolution of genes encoding bacterial

(p)ppGpp synthetases/hydrolases (the Rel, RelA and SpoT proteins). J Mol Microbiol

Biotechnol 3: 585–600.

39.   Magnusson LU, Gummesson B, Joksimović P, Farewell A, Nyström T. 2007. Identical,

independent, and opposing roles of ppGpp and DksA in *Escherichia coli*. *J Bacteriol* 189:

5193–5202.

40.   Åberg A, Fernández-Vázquez J, Cabrer-Panes JD, Sánchez A, Balsalobre C. 2009. Similar

and divergent effects of ppGpp and DksA deficiencies on transcription in *Escherichia coli.*

J Bacteriol 191:3226–3236.

41. Costanzo A, Nicoloff H, Barchinger SE, Banta AB, Gourse RL, Ades SE. 2008. ppGpp and

DksA likely regulate the activity of the extracytoplasmic stress factor sigmaE in

Escherichia coli by both direct and indirect mechanisms. *Mol Microbiol* 67: 619–632.

42. Szalewska-Palasz A, Johansson LUM, Bernardo LMD, Skärfstad E, Stec E, Brännström K,

Shingler V. 2007. Properties of RNA polymerase bypass mutants: implications for the role

of ppGpp and its co-factor DksA in controlling transcription dependent on sigma54. *J Biol*

Chem 282: 18046–18056.

43. Wang B, Dai P, Ding D, Del Rosario A, Grant RA, Pentelute BL, Laub MT. 2019. Affinity-

based capture and identification of protein effectors of the growth regulator ppGpp. *Nat*

Chem Biol 15: 141–150.

44. Giannopoulos G, Sullivan MJ, Hartop KR, Rowley G, Gates AJ, Watmough NJ,

Richardson DJ. 2017. Tuning the modular *Paracoccus denitrificans* respirome to adapt

from aerobic respiration to anaerobic denitrification. *Environ Microbiol* 19: 4953–4964.

45. Olaya-Abril A, Hidalgo-Carrillo J, Luque-Almagro VM, Fuentes-Almagro C, Urbano FJ,

Moreno-Vivián C, Richardson DJ, Roldán MD. 2018. Exploring the denitrification

proteome of *Paracoccus denitrificans* PD1222. *Front Microbiol* 9: 1137.

46. Henard CA, Bourret TJ, Song M, Vázquez-Torres A. 2010. Control of redox balance by the

stringent response regulatory protein promotes antioxidant defenses of *Salmonella*. *J Biol*

Chem 285: 36785–36793.

47. Ellington MJK, Bhakoo KK, Sawers G, Richardson DJ, Ferguson SJ. 2002. Hierarchy of

carbon source selection in *Paracoccus pantotrophus*: Strict correlation between reduction

state of the carbon substrate and aerobic expression of the *nap* operon. *J Bacteriol* 184:

4767-4774.

48.   Ellington MJK, Fosdike WLJ, Sawers RG, Richardson DJ, Ferguson SJ. 2006. Regulation

of the *nap* operon encoding the periplasmic nitrate reductase of *Paracoccus pantotrophus*:

delineation of DNA sequences required for redox control. *Arch Microbiol* 184: 298–304.

49.   Emmerich R, Strehler P, Hennecke H, Fischer H-M. 2000. An imperfect inverted repeat is

critical for DNA binding of the response regulator RegR of *Bradyrhizobium japonicum*.

Nucl Acids Res 28: 4166–4171.

50.   Wood NJ, Alizadeh T, Bennett S, Pearce J, Ferguson SJ, Richardson DJ, Moir JWB. 2001.

Maximal expression of membrane-bound nitrate reductase in *Paracoccus* is induced by

nitrate via a third FNR-like regulator named NarR. *J Bacteriol* 183:3606–3613.

51.   Van Spanning RJ, De Boer AP, Reijnders WN, Westerhoff HV, Stouthamer AH, Van Der

Oost J. 1997. FnrP and NNR of *Paracoccus denitrificans* are both members of the FNR

family of transcriptional activators but have distinct roles in respiratory adaptation in

response to oxygen limitation. *Mol Microbiol* 23: 893–907.

52.   Dworkin J. 2023. Understanding the stringent response: Experimental context matters.

2023. *mBio* e03404-22.

53.   Tosa T, Pizer LI. 1971. Biochemical bases for the antimetabolite action of L-serine

hydroxamate. *J Bacteriol* 106: 972–982.

54. Ross W, Sanchez-Vazquez P, Chen AY, Lee J-H, Burgos HL, Gourse RL. 2016. ppGpp

binding to a site at the RNAP-DksA interface accounts for its dramatic effects on

transcription initiation during the stringent response. *Mol Cell* 62: 811–823.

55. van Verseveld HW, Chesbro WR, Braster M, Stouthamer AH. 1984. Eubacteria have 3

growth modes keyed to nutrient flow. *Arch Microbiol* 137: 176–184.

56. Ellington MJK, Sawers G, Sears HJ, Spiro S, Richardson DJ, Ferguson SJ. 2003.

Characterization of the expression and activity of the periplasmic nitrate reductase of

Paracoccus pantotrophus in chemostat cultures. *Microbiology* 149: 1533–40.

57. Elsen S, Swem LR, Swem DL, Bauer CE. 2004. RegB/RegA, a highly conserved redox-

responding global two-component regulatory system. *Microbiol Mol Biol Rev* 68: 263–279.

58. Phenn J, Pané-Farré J, Meukow N, Klein A, Troitzsch A, Tan P, Fuchs S, Wagner GE,

Lichtenegger S, Steinmetz I, Kohler C. 2021. RegAB homolog of *Burkholderia*

pseudomallei is the master regulator of redox control and involved in virulence. *PLoS*

Pathogens 17:e1009604.

59. Membrillo-Hernandez J, Cook GM, Poole RK. 1997. Roles of RpoS (sigmaS), IHF and

ppGpp in the expression of the *hmp* gene encoding the flavohemoglobin (Hmp) of

Escherichia coli K-12. *Mol Gen Genet* 254: 599–603.

60. Wan X, Brynildsen MP. 2021. Robustness of nitric oxide detoxification to nitrogen

starvation in *Escherichia coli* requires RelA. *Free Radic Biol Med* 176: 286–297.

61.   Pal RR, Bag S, Dasgupta S, Das B, Bhadra RK. 2012. Functional characterization of the

617   stringent response regulatory gene *dksA* of *Vibrio cholerae* and its role in modulation of

618   virulence phenotypes. *J Bacteriol* 194: 5638–5648.

619   62.   de Vries GE, Harms N, Hoogendijk J, Stouthamer AH. 1989. Isolation and characterization

620   of *Paracoccus denitrificans* mutants with increased conjugation frequencies and pleiotropic

621   loss of a (nGATCn) DNA-modifying property. *Arch Microbiol* 152: 52–57.

622   63.   Simon R, Priefer U, Pühler A. 1983. A broad host range mobilization system for in vivo

623   genetic engineering: Transposon mutagenesis in Gram negative bacteria. *Bio/Technology* 1:

624   784–791.

625   64.   Figurski DH, Helinski DR. 1979. Replication of an origin-containing derivative of plasmid

626   RK2 dependent on a plasmid function provided in trans. *Proc Natl Acad Sci USA* 76: 1648–

627   1652.

628   65.   Hahnke S, Moosmann P, Erb T, Strous M. 2014. An improved medium for the anaerobic

629   growth of *Paracoccus denitrificans* Pd1222. *Front Microbiol* 5: 18.

630   66.   Miller JH. 1992. A Short Course in Bacterial Genetics. Cold Spring Harbor Laboratory

631   Press, Cold Spring Harbor, N.Y.

632   67.   Schäfer A, Tauch A, Jäger W, Kalinowski J, Thierbach G, Pühler A. 1994. Small

633   mobilizable multi-purpose cloning vectors derived from the *Escherichia coli* plasmids

634   pK18 and pK19: selection of defined deletions in the chromosome of *Corynebacterium*

635   *glutamicum*. *Gene* 145: 69–73.

68.   Sullivan MJ, Gates AJ, Appia-Ayme C, Rowley G, Richardson DJ. 2013. Copper control of

bacterial nitrous oxide emission and its impact on vitamin B12-dependent metabolism. *Proc*

Natl Acad Sci USA 110: 19926–19931.

69.   Wurch T, Lestienne F, Pauwels PJ. 1998. A modified overlap extension PCR method to

create chimeric genes in the absence of restriction enzymes. *Biotechnology Techniques* 12:

653–657.

70.   Spaink HP, Okker RJH, Wijffelman CA, Pees E, Lugtenberg BJJ. 1987. Promoters in the

nodulation region of the *Rhizobium leguminosarum* Sym plasmid pRL1JI. *Plant Mol Biol* 9:

27–39.

71.   Ind AC, Porter SL, Brown MT, Byles ED, de Beyer JA, Godfrey SA, Armitage JP. 2009.

Inducible-expression plasmid for *Rhodobacter sphaeroides* and *Paracoccus denitrificans.*

Appl Environ Microbiol 75: 6613–6615.

72.   Pratt LA, Kolter R. 1998. Genetic analysis of *Escherichia coli* biofilm formation: roles of flagella,

motility, chemotaxis and type I pili. *Mol Microbiol* 30: 285–293.

73.   Quan S, Hiniker A, Collet J-F, Bardwell JCA. 2013. Isolation of bacteria envelope proteins. pp.

359–366 in Delcour, AH (ed.), Bacterial Cell Surfaces: Methods and Protocols. Humana Press,

Totowa, NJ.

74.   Dunn AK, Karr EA, Wang Y, Batton AR, Ruby EG, Stabb EV. 2010. The alternative oxidase

(AOX) gene in *Vibrio fischeri* is controlled by NsrR and upregulated in response to nitric oxide.

Mol Microbiol 77: 44–55.

75.   Yu L, Wolin MJ. 1969. Hydrogenase measurement with photochemically reduced methyl

viologen1. *J Bacteriol* 98: 51–55.

76.   Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. 1951. Protein measurement with the Folin phenol

reagent. *J Biol Chem* 193: 265-275.

FIGURE LEGENDS

FIGURE 1.

P. denitrificans dksA1$_{Pd}$ and *dksA2$_{Pd}$* complement an *E. coli* Δ*dksA* mutant (Δ*dksA$_{EC}$*). Strains

were streaked on M9 minimal (M9) agar plates containing 10 mM glucose (G) with or without

0.2% (w/v) casamino acids (CAA). Plates were incubated overnight at 37 ºC. 1 - *E. coli*

MG1655; 2 - Δ*dksA$_{EC}$* complemented with *dksA1$_{Pd}$*; 3 - Δ*dksA$_{EC}$* complemented with *dksA2$_{Pd}$*; 4 -

Δ*dksA$_{EC}$* carrying empty pIND4.

FIGURE 2.

Deletion of *dksA1* impairs nitrate utilization during aerobic growth. Aerobic growth of the

indicated strains in: (A) minimal medium containing succinate or (B) minimal medium

containing butyrate, with (+ NO$_3^-$) or without (- NO$_3^-$) 20 mM nitrate. Optical density was

measured (at 600 nm) at the indicated time points (the means from three independent

experiments). The estimated doubling times for growth assays are reported in Table S3.

FIGURE 3.

Loss of both *dksA1* and *rel* leads to failure to grow on reduced carbon substrates. Aerobic growth of indicated strains in liquid minimal medium with butyrate (A) without (- NO_3^-) or (B) with 20 mM nitrate (+ NO_3^-). Optical density was measured (at 600 nm) at the indicated time points (the means from three independent experiments). The estimated doubling times for growth assays are reported in Table S3.

FIGURE 4.

The Δ*dksA1* mutant has reduced *nap* promoter and NAP enzyme activity. (A) β-galactosidase activity in *P. denitrificans* wild-type, Δ*dksA1* and Δ*rel* carrying plasmid p*nap-lacZ*. (B) NAP enzyme activity in the periplasmic fractions of *P. denitrificans* wild-type, Δ*dksA1* and Δ*rel* cells. Cultures were grown in minimal medium containing 10 mM succinate (S) or butyrate (B) as sole carbon and energy source with or without 20 mM nitrate (N). Aerobic (+O_2) cultures were grown in shake flasks at 250 rpm. Anaerobic (-O_2) cultures were grown statically in filled closed cap bottles. The bars represent the mean of at least three separate experiments and the error bars represent the standard deviation of the mean. Asterisks on the bars indicate a significant difference compared to the wild-type within the group according to Student's t-test (*$P \leq 0.05$, **$P \leq 0.01$, ***$P \leq 0.001$).

FIGURE 5.

The RegAB system regulates anaerobic *nap* promoter and NAP activity. (A) β-galactosidase

activity in *P. denitrificans* wild-type, Δ*regA* and Δ*regB* carrying plasmid p*nap-lacZ*. (B) NAP

enzyme activity in the periplasmic fractions of anaerobically grown *P. denitrificans* wild-type,

Δ*regA* and Δ*regB* cells. (C) NAR enzyme activity in the membrane fractions of *P. denitrificans*

wild-type, Δ*regA* and Δ*regB* cells. Cultures were grown in minimal medium containing 10 mM

succinate (S) or butyrate (B) as sole carbon source with or without 20 mM nitrate (N). Aerobic

($+O_2$) cultures were grown in shake flasks at 250 rpm. Anaerobic ($-O_2$) cultures were grown in

filled closed cap bottles statically. The bars represent the mean of at least three separate

experiments and the error bars represent the standard deviation of the mean. Asterisks on the bars

indicate a significant difference compared to the wild-type within the group according to Student's

t-test (*$P \leq 0.05$, **$P \leq 0.01$, ***$P \leq 0.001$).

FIGURE 6.

The *rel* gene is required for recovery from serine hydroxamate treatment. Aerobic growth of

indicated strains in minimal medium containing: (A) succinate only, (B) succinate and 3 mM

serine hydroxamate, and (C) succinate, 3 mM serine hydroxamate and 0.2% (w/v) casamino

acids. Optical density was measured (at 600 nm) at the indicated time points (the means from

three independent experiments). The estimated doubling times for growth assays are reported in

Table S4.

FIGURE 7.

DksA2$_{Pd}$ and RSH regulate *hmp* promoter activity and biofilm formation in *P. denitrificans*. (A) β-galactosidase activities in *P. denitrificans* wild-type, Δ*dksA2* and Δ*rel* carrying plasmid p*hmp-lacZ*. (B) Assays of biofilm formation by the wild-type, deletion, and complemented strains. Dishes were washed, stained with crystal violet, and photographed from above. (C) The crystal violet was extracted with 95% ethanol, and the absorbance measured at 595 nm. Means and standard deviations were calculated from results of three separate assays. Cultures were grown in 6-cm-diameter petri dishes in liquid L broth (Miller), supplemented with 10 mM CaCl$_2$ and incubated without shaking at 30 ºC. Cells were harvested at mid-log phase for the β-galactosidase assay or left undisturbed for 72 hours for the biofilm assay. Asterisks on the bars indicate a significant difference compared to the wild-type according to Student's t-test (**P ≤ 0.01, ***P ≤ 0.001).

Fig. 1

Figure 1. *P. denitrificans dksA1*$_{Pd}$ and *dksA2*$_{Pd}$ complement an *E. coli* Δ*dksA* mutant (Δ*dksA*$_{EC}$). Strains were streaked on M9 minimal (M9) agar plates containing 10 mM glucose (G) with or without 0.2% (w/v) casamino acids (CAA). Plates were incubated overnight at 37 ºC. 1 - *E. coli* MG1655; 2 - Δ*dksA*$_{EC}$ complemented with *dksA1*$_{Pd}$; 3 - Δ*dksA*$_{EC}$ complemented with *dksA2*$_{Pd}$; 4 - Δ*dksA*$_{EC}$ carrying empty pIND4.

Fig. 2

Figure 2. Deletion of *dksA1* impairs nitrate utilization during aerobic growth. Aerobic growth of indicated strains in, **(A)** minimal medium containing succinate or **(B)** minimal medium containing butyrate, with (+ NO_3^-) or without (- NO_3^-) 20 mM nitrate. Optical density was measured (at 600 nm) at the indicated time points (the means from three independent experiments).

Fig. 3

Figure 3. Loss of both *dksA1* and *relA* leads to failure to grow on reduced carbon substrates. Aerobic growth of indicated strains in liquid minimal medium with butyrate **(A)** without (- NO_3^-) or **(B)** with 20 mM nitrate (+ NO_3^-). Optical density was measured (at 600 nm) at the indicated time points (the means from three independent experiments).

Fig. 4

Figure 4. The ΔdksA1 mutant has reduced *nap* promoter and NAP enzyme activity. (A) β-galactosidase activity in *P. denitrificans* wild-type, Δ*dksA1* and Δ*relA* carrying plasmid p*nap-lacZ*. **(B)** NAP enzyme activity in the periplasmic fraction of *P. denitrificans* wild-type, Δ*dksA1* and Δ*relA* cells. Cultures were grown in minimal medium containing 10 mM succinate (S) or butyrate (B) as sole carbon and energy source with or without 20 mM nitrate (N). Aerobic ($+O_2$) cultures were grown in shake flasks at 250 rpm. Anaerobic ($-O_2$) cultures were grown statically in filled closed cap bottles. The bars represent the mean of at least three separate experiments and the error bars represent the standard deviation of the mean. Asterisks on the bars indicate a significant difference compared to the wild-type within the group according to Student's t-test (*$P \leq 0.05$, **$P \leq 0.01$, ***$P \leq 0.001$).

Main Figures

Figure 5. The RegAB system regulates anaerobic *nap* promoter and NAP activity. (A) β-galactosidase activity in *P. denitrificans* wild-type, Δ*regA* and Δ*regB* carrying plasmid p*nap-lacZ*. **(B)** NAP enzyme activity in the periplasmic fractions of anaerobically grown *P. denitrificans* wild-type, Δ*regA* and Δ*regB* cells. **(C)** NAR enzyme activity in the membrane fractions of *P. denitrificans* wild-type, Δ*regA* and Δ*regB* cells. Cultures were grown in minimal medium containing 10 mM succinate (S) or butyrate (B) as sole carbon source with or without 20 mM nitrate (N). Aerobic (+O_2) cultures were grown in shake flasks at 250 rpm. Anaerobic (-O_2) cultures were grown in filled closed cap bottles statically. The bars represent the mean of at least three separate experiments and the error bars represent the standard deviation of the mean. Asterisks on the bars indicate a significant difference compared to the wild-type within the group according to Student's t-test (*P ≤ 0.05, **P ≤ 0.01, ***P ≤ 0.001).

Fig. 6

Figure 6. The *relA* gene is required for the response to amino acid starvation in *P. denitrificans*. Aerobic growth of indicated strains in minimal medium containing: **(A)** succinate only, **(B)** succinate and 3 mM serine hydroxamate, and **(C)** succinate, 3 mM serine hydroxamate and 0.2% (w/v) casamino acids. Optical density was measured (at 600 nm) at the indicated time points (the means from three independent experiments).

Fig. 7

Figure 7. DksA2$_{Pd}$ and RSH regulate *hmp* promoter activity and biofilm formation in *P. denitrificans*. (A) β-galactosidase activities in *P. denitrificans* wild-type, Δ*dksA2* and Δ*relA* carrying plasmid p*hmp-lacZ*. **(B)** Assays of biofilm formation by the wild-type, deletion, and complemented strains. Dishes were washed, stained with crystal violet, and photographed from above. **(C)** The crystal violet was extracted with 95% ethanol, and the absorbance measured at 595 nm. Means and standard deviations were calculated from results of three separate assays. Cultures were grown in 6-cm-diameter petri dishes in liquid L broth (Miller), supplemented with 10 mM CaCl$_2$ and incubated without shaking at 30 ºC. Cells were harvested at mid-log phase for the β-galactosidase assay or left undisturbed for 72 hours for the biofilm assay. Asterisks on the bars indicate a significant difference compared to the wild-type according to Student's t-test (**$P \leq 0.01$, ***$P \leq 0.001$).