**Genomic screening of 16 UK native bat species through conservationist networks uncovers coronaviruses with zoonotic potential**

Cedric C.S. Tan[1,2,*], Jahcub Trew[3,*], Thomas P. Peacock[4,*], Kai Yi Mok[4], Charlie Hart[3], Kelvin Lau[5], Dongchun Ni[6], C. David L. Orme[3], Emma Ransome[3], William D. Pearse[3], Christopher M. Coleman[7], Dalan Bailey[8], Nazia Thakur[8,9], Jessica L. Quantrill[4], Ksenia Sukhova[4], Damien Richard[1], Laura Kahane[3], Guy Woodward[3], Thomas Bell[3], Lisa Worledge[10], Joe Nunez-Mino[10], Wendy Barclay[4], Lucy van Dorp[1], Francois Balloux[1], Vincent Savolainen[3]

[1]UCL Genetics Institute, University College London, Gower St, London WC1E 6BT, UK.

[2]The Francis Crick Institute, 1 Midland Rd, London NW1 1AT, UK.

[3]Georgina Mace Centre for the Living Planet, Department of Life Sciences, Imperial College London, Silwood Park Campus, Ascot SL5 7PY, UK.

[4]Department of Infectious Disease, Imperial College London, St Marys Medical School, Paddington, London W2 1PG, UK.
[5]Protein Production and Structure Core Facility (PTPSP), School of Life Sciences, École Polytechnique Fédérale de Lausanne, Rte Cantonale, 1015 Lausanne, Switzerland.
[6]Laboratory of Biological Electron Microscopy (LBEM), School of Basic Science, École Polytechnique Fédérale de Lausanne, Rte Cantonale, 1015 Lausanne, Switzerland.
[7]Queen's Medical Centre, University of Nottingham, Derby Rd, Lenton, Nottingham NG7 2UH, UK.
[8]The Pirbright Institute, Surrey GU24 0NF, UK.
[9]Nuffield Department of Medicine, University of Oxford, Oxford OX3 7BN, UK.
[10]The Bat Conservation Trust, Studio 15 Cloisters House, Cloisters Business Centre, 8 Battersea Park Road, London SW8 4BG, UK.

* These authors contributed equally

Correspondence: V.S. (v.savolainen@imperial.ac.uk)

**Abstract (max 150 words)**

There has been limited characterisation of bat-borne coronaviruses in Europe. Here, we
screened for coronaviruses in 48 faecal samples from 16 of the 17 bat species breeding
in the UK, collected through a bat rehabilitation and conservationist network. We
recovered nine (two novel) complete genomes across six bat species: four
alphacoronaviruses, a MERS-related betacoronavirus, and four closely related
sarbecoviruses. We demonstrate that at least one of these sarbecoviruses can bind and
use the human ACE2 receptor for infecting human cells, albeit suboptimally.
Additionally, the spike proteins of these sarbecoviruses possess an R-A-K-Q motif,
which lies only one nucleotide mutation away from a furin cleavage site (FCS) that
enhances infectivity in other coronaviruses, including SARS-CoV-2. However, mutating
this motif to an FCS does not enable spike cleavage. Overall, while UK sarbecoviruses
would require further molecular adaptations to infect humans, their zoonotic risk is
unknown and warrants closer surveillance.

## Introduction

The majority of emerging infectious diseases in humans are zoonotic - arising from animal-to-human transmission of a pathogen[1] - and more than 70% originate in wildlife[2]. *Coronaviridae* is a diverse family of viruses that can infect a broad range of animals and are prone to zoonotic spillovers. There are seven major coronaviruses known to infect humans: SARS-CoV-2 is the agent of the COVID-19 pandemic whose direct ancestor has not yet been identified but its closest relatives have been isolated from horseshoe bats. SARS-CoV-1 caused a major international outbreak in 2002-2004 with around 8,000 recorded cases and at least 774 deaths[3]. MERS-CoV fuels recurrent disease outbreaks in humans through repeated host jumps from its reservoir in camels[4]. Four coronaviruses (HCoV-229, -NL63, -OC43 and -HKU1) circulate endemically in humans and their ancestral reservoirs are believed to be species of bats and rodents, with host jumps into humans likely facilitated by other mammals as bridging hosts[5–7]. Additionally, multiple host jumps from animals into humans leading to isolated or small clusters of cases have been documented for other coronavirus species[5]. Given the current health burden exerted by coronaviruses and the risk they pose as possible agents of future epidemics and pandemics, surveillance of animal-borne coronaviruses should be a public health priority. Indeed, the discovery and characterisation of the diversity of coronaviruses harboured by mammals across the world is the first step for designing pre-emptive measures to minimise human or animal exposure. Here, we focus on bats since some human coronaviruses have their ancestral origins in these species.

Several studies over the last decade have screened bats across Asia, Africa, the Middle East and Europe for coronaviruses, finding anywhere from 1.5-23% coronavirus prevalence in animals tested[8–19]. A selection of studies representing the diversity of previous screening efforts is listed in Supplementary Data 1. These prevalence estimates were primarily obtained via a reverse transcription real-time PCR (RT-PCR) using degenerate primers designed to target most coronaviruses species (i.e., pan-coronavirus primers; Supplementary Data 1). Given the vast diversity of coronaviruses, including those yet to be discovered, it is difficult to design primers that can amplify and capture the full diversity of coronaviruses. Our own comparative analysis of published

primer sets show that existing RT-PCR assays[20–24] underestimate coronavirus

prevalence (Supplementary Information; Supplementary Fig. 1). Difficulties with primer

design is exacerbated by low RNA concentrations in field samples and RNA

degradation, so the large variability in prevalence estimates in these studies may be

due to the sensitivity of the primer set used rather than the epidemiology of bat-

coronaviruses. While sample RNA quality remains mainly dependent on sample

collection and laboratory practices, because untargeted RNA sequencing does not

require *a priori* knowledge of sequence information, it provides a more accurate

estimate of viral diversity and prevalence. Hence, we chose this approach over RT-PCR

specifically to screen UK bats for coronaviruses.

For a zoonotic event to occur, a virus must be able to transmit efficiently between

animals and humans and be able to infect and replicate in human cells through

interactions with host cellular machinery. Additionally, zoonotic pathogens of most

concern are those that can transmit efficiently between humans. As such, the true

zoonotic potential (i.e., the likelihood of infecting humans in the future) of a virus can

only be determined by assessing all these prerequisite abilities. Therefore, excluding

unethical experimental infection or transmission studies involving humans, assessments

of zoonotic potential should be done via alternative approaches. These include *in silico*

analyses that determine the degree of sequence and structural homology to other

known and closely-related human-infecting viruses[11,25]. Even more compelling evidence

for zoonotic potential can be obtained *in vitro*, and one of the most direct assessments

is to isolate and test the infectivity of novel viruses in human cells[25,26]. However, this

would increase the risk of exposure to these potentially infectious agents, necessitating

stringent biosafety precautions. Additionally, isolation of novel viruses via cell culture,

without prior knowledge of their cell tropism and receptor usage, can be challenging.

A low risk and effective alternative is to measure the binding efficiency of viral entry

proteins to host receptors[10], or to assess efficiency of viral entry into human cell lines

via a pseudovirus assay[25], which expresses only the viral entry protein in a non-

infectious reporter system. While observed binding and cellular entry in these low-risk

assays do not show that a virus can replicate effectively in human cells, infect humans,
or transmit efficiently between humans, they provide an indication of which human cell
receptors can be exploited by novel viruses, which are one key determinant of viral
infection. However, despite the importance of functional validation, many studies to date
fall short of providing *in vitro* or even *in silico* assessments of zoonotic risk
(Supplementary Data 1).

There are 17 bat species that breed in the United Kingdom (UK), most of which often
roost in domestic buildings, churches, barns and other man-made structures. This
frequent habitat overlap with humans places bats in close proximity to domesticated and
farmed animals, which can serve as potential bridging hosts for transmitting bat-borne
viruses to humans[27]. However, multiple factors have to align for the successful
emergence of a zoonotic pathogen in humans, including the frequency of exposure, the
ability of the pathogen to infect humans and its capacity for onward human-to-human
transmission[28]. The relative risks of these various factors for zoonotic spillover remains
largely unknown and may vary depending on pathogen and geographical context. All
UK bat species are protected by law across the UK with licences required for work
related to bats. So, although direct contact is rare among the general public, it is far
more common for the small proportion of the population comprising bat scientists,
ecologists, conservationists and bat rehabilitators that undertake regular research,
monitoring, surveillance, and bat rehabilitation work.

Only two studies screening UK bats for coronaviruses have been conducted to date[11,19].
The first, published over a decade ago, screened seven bat species and detected
alphacoronaviruses in Daubenton's bat and Natterer's bat (*Myotis daubentonii and M.*
*nattereri, respectively*)[19]. The other, from 2021, screened faecal samples from lesser
horseshoe bats (*Rhinolophus hipposideros*) and recovered the whole genome
sequence of a single sarbecovirus, RhGB01 (MW719567)[11]. However, neither study
provided direct *in vitro* assessments of zoonotic risk. Accordingly, the viral diversity and
zoonotic potential of UK bat viruses remains largely unknown. This is equally true of
most other UK mammals. However, given that the evolutionary origins of many

coronaviruses of human health concern can be traced back to bats, assessing their
zoonotic potential in UK bats is a top priority, before moving on to other animal groups.
138
To address this knowledge gap, we used an existing UK network of bat rehabilitators
and conservationists to collect faecal samples from UK bats. Faeces from all but one
bat species breeding in the UK (the grey long-eared bat, *Plecotus austriacus*, the rarest
species in the UK) were collected and subsequently screened using deep RNA
sequencing to characterise the genomic diversity of bat-borne coronaviruses in the UK.
To assess their zoonotic potential, we then tested the ability of a subset of these
coronaviruses to bind human-cell receptors *in vitro*, which is a key requisite for human
infection.

**Results**

*Untargeted RNA sequencing recovers nine complete coronavirus genomes including two new species*

We performed deep RNA sequencing on 48 faecal samples from 16 of the 17 UK breeding bat species, with wide geographic coverage and spanning two years of sampling (Supplementary Fig. 2). Through taxonomic assignment of sequencing reads using Kraken2[29], we detected the presence of at least 30 viral families, 53% of which primarily infect non-mammalian hosts such as plants, insects and bacteria (Supplementary Fig. 3a). Additionally, the total relative abundance of viral species that infect non-mammalian hosts was significantly higher than that for mammalian viruses (two-sided Mann-Whitney U test, $U$=1393, $p$=0.004; Supplementary Fig. 3b). These findings indicate that the faecal virome in UK bats largely comprises viruses that do not necessarily infect them, nor other mammals, including humans.

We next focused on coronaviruses due to their relevance to human health and recovered nine complete genomes (96-100% completeness; assessed by CheckV[30]) and five partial contigs (<3%) across six UK bat species (*M. daubentonii*, *Pipistrellus pipistrellus*, *P. pygmaeus*, *P. auritus*, *R. ferrumequinum*, and *R. hipposideros*), detecting coronaviruses amongst 29% of the samples. The nine complete genomes were assessed by CheckV to be of high-quality[30] and read alignments to these genomes indicated an even coverage of reads with a median coverage of 548-7958 reads per position (Supplementary Fig. 4; Table 1).

A global phylogenetic tree based on alignment-free genetic distances[31] revealed the genus and subgenus membership of these new coronaviruses (Fig. 1a; see Methods). We then followed with local maximum-likelihood phylogenetic analyses to determine their precise placement within each subgenus (Fig. 1b-d). These phylogenetic analyses reveal that the nine novel genomes we recovered comprise four alphacoronaviruses from the *Pedacovirus* subgenus, five betacoronaviruses including one merbecovirus, and four sarbecoviruses (Fig. 1). Three of the coronaviruses recovered from *M.*

*daubentonii* (which we call MdGB01-03) form a well-supported clade with other
pedacoviruses isolated from the same bat species in Denmark (Fig. 1b). One
coronavirus sequenced from *P. pipistrellus* (PpiGB01) falls as a sister lineage to the
above clade. Another coronavirus from *P. auritus* (PaGB01) is related to MERS-CoV-
like merbecoviruses isolated from *Hypsugo*, *Pipistrellus*, and *Vespertilio* spp. from
Western Europe and China (Fig. 1c). Four coronaviruses isolated from *R.*
*ferrumequinum and R. hipposideros* (RfGB01-02 and RhGB07-08, respectively) are
closely related to the previously described UK bat sarbecovirus, RhGB01[32] (Fig. 1d).
186
Of the nine coronaviral genomes recovered here, two represent new species. Indeed,
pedacovirus PpiGB01 from *P. pipistrellus* was relatively divergent from its closest
match, a pedacovirus previously isolated from *M. daubentonii* (less than 81% nucleotide
sequence identity; Table 1). Similarly, merbecovirus PaGB01 shares less than 82%
sequence identity to its closest match, a merbecovirus isolated from *P. kuhlii* in Italy
(Table 1). Overall, our coronavirus screening efforts have extended our knowledge of
the existing diversity of coronaviruses. Looking at their genomic structures, we identified
one new gene in each of these new species (Supplementary Information;
Supplementary Fig. 5).
196
Viruses that are able to infect a broad range of hosts have been associated with a
higher risk of emerging as infectious diseases among humans[33,34]. Here, the four
sarbecovirus genomes, representing one viral species, were recovered from two distinct
horseshoe bat species, *R. ferrumequinum* and *R. hipposideros*. RhGB07, RhGB08,
RfGB01, RfGB02 share 97-100% identity with RhGB01 previously described in *R.*
*hipposideros*[11]. To better understand how these viruses might be shared among the two
hosts, we looked at the habitat distribution of each horseshoe bat species. The two
horseshoe bat species share a large proportion of their habitats, with 33% of their
occurrence records reported at the same geographical coordinates. Furthermore,
species distribution modelling predicted that 45% of the total land area occupied by the
two species is shared (Supplementary Fig. 6a). Since the two *Rhinolophus* species can
share roosts[35], these results indicate a potentially high frequency of direct contact,

which may facilitate viral sharing and thus account for the isolation of RhGB01-like
sarbecoviruses that are closely related from these two species.

To extend this analysis, we examined both observed and predicted distributions of all 17
UK breeding bat species to identify potential viral sharing hotspots for future
surveillance work. By analysing 42,953 occurrence records, we identified three regions
near Bristol, Birmingham and Brighton with particularly high species diversity (the most
diverse regions having 16 species in a single 5x5 km grid; Supplementary Fig. 6b).
Additionally, we identified regions within the UK, especially in Wales and the south
coast of England where the habitats of the greatest number of different bat species are
predicted to coincide (Supplementary Fig. 6c). Alongside an understanding of the
ecology of native species, including co-roosting and foraging behaviours, such
information is a useful resource for future surveillance studies, and for prioritising focal
areas of zoonotic risk.

*Sarbecoviruses recovered from UK bats can bind the human ACE2 receptor for cellular*
*entry*

We tested whether representatives of the newly identified UK coronaviruses (the
sarbecoviruses RhGB07 and RfGB02, the merbecovirus PaGB01, and the outlier
pedacovirus PpiGB01) could use human cellular receptors for viral entry as a proxy for
assessing their zoonotic potential. We successfully incorporated the spike proteins of
these UK coronaviruses into lentivirus-based pseudoviruses (see Methods,
Supplementary Figure 7c). We then tested the ability of these spike-expressing
pseudoviruses to infect human cells expressing the human receptors, angiotensin-
converting enzyme 2 (hACE2), dipeptidyl peptidase-4 (hDPP4) and aminopeptidase N
(hAPN), which are the primary receptors exploited by SARS-CoV-2, MERS-CoV and
HCoV-229E, respectively.

None of the spike pseudoviruses of the UK coronaviruses could enter cells using any of
the receptors except RhGB07, which showed significantly higher entry into cells

overexpressing hACE2 compared to those not expressing hACE2 (Fig. 2a; $p<0.0001$).

As expected, SARS-CoV-2, MERS-CoV and HCoV-229E showed significantly higher

entry into cells overexpressing hACE2, hDPP4 and hAPN, respectively (Fig. 2a;

$p<0.0001$). Further, VSV-G expressing pseudovirus, which enters cells using ubiquitous

protein receptors, showed comparably high entry across all groups (Fig. 2a).

Additionally, using biolayer-inteferometry (BLI), we confirmed that the RhGB07 spike is

able to bind hACE2 with a dissociation constant, $K_d=253nM$ (Fig. 2b). However, the

binding affinity of RhGB07 spike to hACE2 is approximately 17-fold lower than that for

the SARS-CoV-2 spike ($K_d=15nM$) (Fig. 2b).

Given the lower binding affinity of RhGB07 spike compared to SARS-CoV-2, we then

investigated if, like SARS-CoV-2, RhGB07 spike-expressing pseudoviruses can infect

human cells expressing lower (HEK293T-hACE2 – HEK293Ts stably transduced with

hACE2; Supplementary Fig. 7a) or physiological levels of hACE2 (Calu-3 lung, and

Caco-2 colorectal cell lines). Alongside this, we tested the entry of RfGB02, PaGB01

and PpiGB01 spike pseudoviruses in the same cell lines in case they use a human

receptor not otherwise tested as in Fig. 2a. As positive controls, we included the spike

proteins from other coronaviruses, BANAL-20-52/SARS-CoV-2 (wild-type Wuhan-Hu-1

with D614G), MERS-CoV, and HCoV-229E, which can efficiently enter these cell lines

using hACE2[36,37], hDPP4 and hAPN, respectively. We also included the negative

control, RaTG13, which can bind hACE2 but cannot enter cells expressing lower or

physiological levels of hACE2[37]. As expected, RaTG13 could not enter any of these cell

lines, while all positive controls showed significantly higher entry into these cell lines

than 'bald' pseudoviruses not expressing any spike protein ($p<0.01$; Fig. 2c). In contrast,

none of the UK spike pseudoviruses tested, including RhGB07, displayed significant

entry into any of these human cell lines ($p>0.05$; Fig. 2c).

Separately, we asked if other host proteins are necessary for efficient cellular entry of

the UK coronaviruses. In particular, the transmembrane serine protease 2 (TMPRSS2)

protease has been shown to greatly enhance the entry efficiency of MERS-CoV[38] and

HCoV-229E[39] spike pseudoviruses into human cells. However, PaGB01 and PpiGB01,

which fall in the same subgenus as MERS-CoV and HCoV-229E, respectively, could not

enter TMPRSS2-overexpressing cells (Fig. 2d).

RhGB07 can bind and use overexpressed hACE2 for cellular entry but RfGB02 cannot,

despite the high 98% sequence identity of their spike proteins. This begs the questions

as to how RhGB07 might have acquired the ability to use hACE2, and whether this

might be associated with the usage of bat ACE2 orthologues. To investigate this, we

tested the entry of RhGB07 and RfGB02 spike pseudoviruses into human cells

expressing the ACE2 orthologues from four bat species (*R. ferrumequinum*, *R. pusillus*,

*Myotis lucifugus*, and *Rousettus leschenaultia*). These bat ACE2 proteins all expressed

robustly, albeit to slightly differing efficiencies (Supplementary Fig. 7b), as seen

previously[40]. Unfortunately, throughout the course of this study, there was no publicly

available ACE2 sequence for *R. hipposideros* (from which RhGB07 was recovered),

and no ACE2 transcripts could be identified directly from our metatranscriptomic

libraries.

Nevertheless, we detected significant cell entry but only through *M. lucifugus* ACE2

($p<0.0001$; Fig. 2e), and neither RhGB07 nor RfGB02 could use *R. ferrumequinum*

ACE2 receptors, despite RfGB02 being sampled from this species. In contrast, SARS-

CoV-2, BANAL-20-52 and RaTG13 were all able to efficiently use *R. ferrumequinum*

ACE2 ($p<0.0001$; Fig. 2e), indicating that this ACE2 construct could allow sarbecovirus

entry. Surprisingly, BLI measurements indicate detectable binding of RhGB07 spike to

both *R. ferrumequinum* and *M. lucifugus* ACE2 (Fig. 2f), which means that RhGB07 can

bind *R. ferrumequinum* ACE2 but not efficiently enter cells expressing this receptor.

This highlights that binding of host ACE2 alone may not be sufficient for efficient viral

entry, and that other host cell-virus interactions (e.g., presence of suitable co-receptors)

may be required. Taken together, our results suggest that RhGB01-like viruses may not

be using ACE2 to infect their native *Rhinolophus* hosts.

*Structural and sequence features of RhGB07 spike explain detectable but inefficient usage of hACE2*

To gain deeper insight into the results of the above assays, we used the AlphaFold artificial intelligence program[41] to predict the 3D structure of the receptor-binding domain (RBD) of the RhGB07 spike protein. We then compared it to the resolved RBD structures of SARS-CoV-2[42], BANAL-236 (a close relative of BANAL-20-52)[8], and RaTG13[37] bound to hACE2. Superposition of the RBD structures showed high structural conservation across all four sarbecoviruses (Fig. 3a). Additionally, the 3D structure of the RhGB07 RBD near the RBD-hACE2 binding interface was highly similar to that of SARS-CoV-2 (Fig. 3b), which was confirmed by comparing the area of contact surface ($894 \text{ Å}^2$ and $850 \text{ Å}^2$, respectively; Supplementary Fig. 8).

These findings account for the ability of the RhGB07 spike protein to bind hACE2 (Fig. 2a, b). To understand why RhGB07 spike pseudoviruses could not enter cells expressing the ACE2 receptor at physiological levels (Fig. 2c), we compared the level of conservation at key RBD residues of SARS-CoV-2[42] and SARS-CoV[43] in contact with hACE2. This included sarbecoviruses isolated from Asia, Europe, and Africa, which bind hACE2 with various affinities[44–46]. All these sarbecoviruses showed conservation at more than half of SARS-CoV-2 (Fig. 3c) or SARS-CoV (Fig. 3d) contact residues, with high levels of conservation at certain positions like Y453, N487, Y489, G502 and Y505 (SARS-CoV-2 numbering; Fig. 3c). Previous deep mutational scanning experiments showed that all these positions, except Y453, cannot be mutated without considerable loss of hACE2 binding affinity[47], indicating that contact residues are important determinants of hACE2 binding. Notably, the novel RhGB01-like sarbecoviruses share only 9/17 SARS-CoV-2 (Fig. 3c) and 7/14 SARS-CoV contact residues (Fig. 3d). This is slightly below RaTG13, which shares 11/17 SARS-CoV-2 and 8/14 SARS-CoV contact residues, respectively. In contrast, BANAL-20-52 shares 15/17 contact residues with SARS-CoV-2.

These results indicate poorer conservation of these key contact residues in UK
sarbecoviruses, which would explain the relatively lower hACE2 usage efficiency, and
hence the ability to infect human cells, of RhGB07 and RaTG13 compared to BANAL-
20-52. Similarly, the poorer conservation of SARS-CoV-2 contact residues in BtKY72
than Khosta-2 (Fig. 3c) may explain the lower binding affinity to hACE2 of the former[45].
Notably, RhGB07 - but not RfGB02 - could enter cells using hACE2 (Fig. 2a), despite
their spike proteins sharing the same variants at all SARS-CoV-2 key contact residues
(Fig. 3c). The RfGB02 spike has 26 amino acid mutations relative to RhGB07, and only
four of these were within the RBD (K337N, H432L, T470A, P487Q; Supplementary Data
2). As such, either the residues at these positions are, in addition to the SARS-CoV-2
contact residues, important mediators of hACE2-binding, or the remaining 22 non-RBD
mutations have caused structural changes that reduce the binding affinity to hACE2, or
both. Further experiments delineating these mutational effects could help to shed light
on the molecular determinants of sarbecoviral entry into human cells.

Remarkably, the RhGB01-like sarbecoviruses all possess a R-A-K-Q sequence at the
S1/S2 cleavage site (spike residues 669-672; Supplementary Fig. 9a), which is one
nucleotide away (Gln/C**A**A to Arg/C**G**A) from the canonical R-X-K/R-R motif, a potential
furin cleavage site (FCS) that allows cleavage by host furin-like proteases, enhancing
the ability of many coronaviruses, including SARS-CoV-2, to infect human cells[48,49].
This R-A-K-Q motif is also found in Khosta-2[50], a sarbecovirus recovered from *R.*
*hipposideros* in Khosta, Russia, which is at the south-eastern extremes of Europe, but
not in BtKY72 from *Rhinolophus* sp. in Kenya[51] or other sarbecoviruses isolated from
Asia. However, Western blot analyses indicated that even when we mutated R-A-K-Q to
R-A-K-R (i.e., a Q672R mutation), the RhGB07 spike is not cleaved by any human host
protease (Supplementary Fig. 9b). Previous studies have shown that the FCS on
SARS-CoV-2 (681-RRAR-684) lies on an extended flexible loop that protrudes out of
the spike structure, which allows access by host furin[52,53]. Also, it has been shown that
deletions that shortened this extended loop, while leaving the multibasic site intact.
prevented efficient cleavage of SARS-CoV-2 spike, which was likely due to reduced
accessibility of the FCS[49]. This loop is seven residues shorter in RhGB01-like viruses

(Supplementary Fig. 9b), which may explain why no cleavage was observed for the

RhGB07 R-A-K-R pseudovirus mutant.

Finally, a recombination analysis of the RhGB01-like and other representative

sarbecoviruses indicates a high prevalence of recombination (Supplementary

Information; Supplementary Fig. 10), which may accelerate adaptation for infecting

novel hosts. Given these findings, the current zoonotic risk of sarbecoviruses in UK

bats, while small, cannot be ignored and warrants more extensive surveillance of bats at

the national scale.

**Discussion**

The emergence of the COVID-19 pandemic in 2019 is a sobering reminder of the massive impact of zoonotic viruses on global health and economy. Despite this, genomic surveillance in wildlife remains limited. In this study, we used an existing network of bat rehabilitators and conservationists to obtain geographically and temporally diverse samples from almost all bat species in the UK. We argue that this can be a sustainable and effective surveillance model to identify and characterise novel animal-borne viruses that may or may not yet be able to infect humans but might evolve the ability to do so in the future.

We provided evidence that at least one sarbecovirus isolated from UK horseshoe bats can bind hACE2 *in vitro* and discuss these patterns relative to our *in silico* analyses. Crook *et al.*[11] performed a contact residue analysis, similar to the one we report in Fig. 3c and Fig. 3d, on RhGB01 and suggested that moderate similarity in its key contact residues indicates that it is unlikely to bind hACE2. However, our *in vitro* (Fig. 2) and *in silico* (Fig. 3) results highlight that despite having only moderate conservation of key contact residues, RhGB07 can bind and use hACE2. Additionally, the spike of RhGB07, but not RfGB02, can use hACE2 for cellular entry, despite identical conservation levels to SARS-CoV-2 at key contact residues. These findings indicate that assessing the conservation of key contact residues (Fig. 3c, d) may have limited predictive power for whether a spike protein can bind hACE2, possibly due to multiple structural configurations allowing hACE2 binding. This is further evidenced by the different contact residues for SARS-CoV and SARS-CoV-2.

Our findings indicate that the RhGB01-like viruses likely require further adaptations, particularly in their spike proteins, before they can make a zoonotic jump. Notably, single mutations of some of the SARS-CoV-2 contact residues in sarbecoviral spike proteins have been shown to enable binding of ACE2 from novel host species and improve binding affinity by greater than fivefold[45]. Additionally, a single T403R mutation in the RaTG13 spike has been shown to allow the virus to infect human cells[54]. Given this, we speculate that the genetic barrier precluding effective hACE2 usage for cellular

entry into human cells may be small. This may also be the case for the other RhGB01-
like sarbecoviruses sampled previously[11]. Strikingly, of the two RhGB01-like viruses we
investigated here, one was capable of infecting hACE2-overexpressing cells and the
other not, despite 98% spike sequence similarity and identical SARS-CoV-2 residues.
This further indicates that minor adaptations in the spike protein may significantly affect
binding affinity with host receptors, and hence the zoonotic potential of different viral
lineages.

We also identified a R-A-K-Q sequence in all European sarbecoviruses that resembles
an FCS precursor, but which is absent in all Asian sarbecoviruses considered
(Supplementary Fig. 9a). This supports previous observations that FCSs naturally occur
in coronaviruses and have emerged independently at least six times amongst
betacoronaviruses[55]. However, even after mutating R-A-K-Q to R-A-K-R, we could not
detect any cleavage during expression of RhGB07 spike by human proteases
(Supplementary Fig. 9b). These findings indicate that, in addition to acquiring a
functional FCS via substitution, European sarbecoviruses would likely have to acquire
an extended loop structure (like in SARS-CoV-2) via insertion for efficient spike
cleavage.

We found a high prevalence of genetic recombination amongst sarbecoviruses,
particularly in the spike gene (Supplementary Fig. 10), which may facilitate viral
adaptations to overcome the genetic barrier for a zoonotic jump. This observation is
corroborated by other studies that have also suggested an enrichment of recombination
signals in or surrounding the sarbecovirus spike gene[56,57]. Co-infections and
subsequent recombination of RhGB01-like viruses with other coronaviruses that already
effectively use hACE2 may therefore facilitate zoonotic transmission. As such, the
possibility of a future host-jump into humans cannot be ruled out, even if the risk is small.
This reiterates the need for individuals that are in frequent contact with bats, such as bat
rehabilitators, to adhere to current biosafety practices to reduce their exposure to bat
coronaviruses and likewise to reduce the risk of the exposure of bats to human-borne
coronaviruses[58], such as SARS-CoV-2 or the endemic HCoVs. Fortunately, in the UK,

the risk of zoonotic exposure is minimised for most people through a lack of direct

contact (roosting spaces are often well away from human inhabitants) along with the

provision of science-based information to roost owners by organisations such as the Bat

Conservation Trust (https://www.bats.org.uk).

Our *in vitro* assays indicate that RhGB01-like sarbecoviruses, including RfGB02 that

was directly sampled from this species, do not use *R. ferrumequinum* ACE2 as their

primary receptor, which is in line with other studies of bat coronaviruses[45,59].

Importantly, this raises the question as to what evolutionary mechanisms drive the

acquisition of the ability to use hACE2 in bat sarbecoviruses. Given previous

associations between pathogen host breadth and their capacity to emerge as zoonotic

diseases[33,34], we speculate that multi-host viruses tend to have 'generalist' cell entry

receptors that possess a low genetic barrier to the evolution of zoonotic transmission.

More extensive surveillance of the viral sharing dynamics in mammalian hosts, including

bats, may provide key insights into the molecular and ecological determinants of

zoonotic events. Such studies can leverage both species occurrence data and niche

modelling to prioritise regions where a high number of species are likely to be found

combined with an understanding of species ecology for quantification of risk.

The initial spread of SARS-CoV-2 in China, its likely evolutionary origin in *Rhinolophus*

bats[60], and the subsequent identification of other bat-borne sarbecoviruses in Southeast

Asia[8,10], has focused attentions on the zoonotic risk of coronaviruses in those

geographical regions. However, our findings highlight the zoonotic risk of

sarbecoviruses may extend beyond Asia, stressing the importance of more extensive

surveillance globally.

Finally, whilst it is imperative to quantify the risk of zoonotic events from bats and design

approaches to mitigate this risk more effectively, bats fulfil important roles in

ecosystems globally, including services such as arthropod pest suppression, pollination

and seed dispersal[61]. Some bat species have rapidly declining populations – for

example, one third of the most threatened mammalian species in the UK are bats[62,63].

Recent studies have shown that human-associated stressors such as habitat loss and

changes in land-use can be important drivers of zoonotic spillover from wildlife[64,65], and

that bat culls are ineffective in minimising cross-species transmission[66]. As such, it is

vitally important that an integrated ecological conservation approach is taken that

includes maintaining legal protection, rather than destruction of wildlife and its habitat, in

future efforts to mitigate zoonotic risk.

## Methods

*Sample collection*

Sampling kits were sent out to various bat rehabilitators in the UK, as described previously[58], for the collection of faeces from bats. These faecal samples (0.02-1g) were immediately stored in 5 ml of RNAlater solution to prevent degradation of RNA. The geographical locations and collection dates for all samples are provided in Supplementary Data 3. Registered bat rehabilitators have received relevant training to aide in this role, including bat health assessment and identification, public engagement, risk assessment, and legal requirements, approved by experienced trainers. Unlike other activities which involve handling bats, a license is not required for care and rehabilitation purposes in the UK, except where bats are to be kept in captivity for six months or more. Therefore, for this study, faecal samples were collected by bat rehabilitators without the need for legal or ethical approval, in particular because all sampling was non-invasive and did not require handling of bats.

*Murine hepatitis virus (MHV) spike-in control culture*

MHV (GenBank AY700211.1) was propagated in an NCTC 1469 clone derivative (ATCC CCL-9.1) cell line in high glucose DMEM and 10% horse serum. Both the MHV and NCTC cell line were acquired from the American Type culture Collection (ATCC, Manassas, Virginia, USA). Cell culture supernatant was isolated for later RNA extraction.

*RNA extraction*

RNA was extracted from faecal samples using the QIAamp Viral RNA Mini Kit (Qiagen) following the protocol for extracting RNA from stool samples. We used up to 0.5 g of faeces, which was vortexed in 2ml of 0.9% NaCl solution, at 6000rpm for 2 minutes. The supernatant was filtered using a 0.2μm syringe filter, 280 μl of which was used for RNA extraction. For the MHV spike-in control, we used 140 μl of culture supernatant for RNA extraction. Total RNA was eluted in 80 μl of AVE buffer and stored at -80$^{o}$C. RNA was quantified using Qubit 2.0 fluorometer (Invitrogen). All faecal extractions were

499    spiked with 20 µl of MHV RNA prior to library preparation to act as a sequencing quality
500    control.

*Coronavirus database*

To create a database representing the extant global genomic diversity of coronaviruses,
we downloaded all complete *Coronaviridae* (taxid:11118) genomes from NCBI Virus,
excluding provirus sequences (accessed 4th July 2022). Additionally, we downloaded all
non-human-associated and non-SARS-CoV-2 betacoronaviruses from GISAID[67] (*n*=29).
To minimise the overrepresentation of certain viral species, we randomly retained 50
genomes for each of the following species: porcine epidemic diarrhoea virus, avian
infectious bronchitis virus, MERS-CoV, SARS-CoV and SARS-CoV-2 sequences. This
yielded a final dataset comprising 2118 genomes.

*Metagenomic sequencing and assembly*

All samples were prepared for sequencing using the NEBNext® Ultra™ Directional RNA
Library Prep Kit, with a QIAseq FastSelect rRNA depletion step. Prior to library
preparation, we also spiked in MHV RNA (GenBank AY700211.1) as a positive control.
Sequencing was carried out using Illumina NovaSeq, paired end 150 bp. Quality control
of reads was performed using bbduk.sh v39.01 from the BBTools Suite
(https://sourceforge.net/projects/bbmap/). Briefly, we trimmed adapter sequences and
read ends below Q10, and discarded trimmed reads with average quality below Q10.
Reads that mapped to the positive control using Bowtie2 v2.4.5[68] were removed prior to
all downstream analyses. *De novo* metagenomic assembly was performed on quality-
controlled or raw reads for each sample using coronaSPAdes v3.15.4[69]. Assembled
scaffolds were then queried using BLASTn against all 2118 genomes in our coronavirus
database to determine their most related reference. Scaffolds that could be aligned
using BLASTn to coronaviruses in our database and that were already longer than 28kb
were considered as complete genomes.

In some cases, *de novo* assembly yielded multiple scaffolds that were shorter than 28kb

but shared the same closest reference. We 'stitched' these scaffolds together using the

BLASTn alignment coordinates to the closest coronavirus reference and replaced any

gaps with Ns. *De novo* assembly using adapter-trimmed reads, without quality trimming

or filtering, produced better results, producing longer and more complete scaffolds,

yielding six >28kb scaffolds (MdGB01, MdGB02, MdGB03, PpiGB01, RfGB01,

RfGB02), compared to quality-controlled read assembly which yielded only two

(RfGB01, PpiGB01). Further, the two >28kb scaffolds, RfGB01 and PpiGB01, generated

using either adapter-trimmed or quality-controlled assemblies were identical, suggesting

that *de novo* assembly using adapter-trimmed reads were reliable. We hence chose the

assemblies generated using adapter-trimmed reads for our downstream analyses. We

named the novel complete genomes following the naming convention for the

sarbecovirus previously described in a UK bat, RhGB01 – species: 'Rh' (*R.*

*hipposideros*), region the coronavirus was found in: 'GB' (Great Britain) and the

frequency of description: '01' (the first described in that species and country).

*Genome annotation and characterisation of novel genes*

Assembled genomes were annotated using Prokka v1.14.6[70], and annotated genes

were inspected to identify and correct erroneous frameshifts that were present in the

raw assemblies to produce the final genomes. For the four novel sarbecoviruses

(RhGB07, RhGB08, RfGb01, RfGB02) and three of the pedacoviruses (MdGB01,

MdGB02, MdG03), we also performed genome alignments to their closest known

relative shown in Table 1 to check if erroneous indels were present. The gene

annotations were also analysed to determine if these genomes carry any novel genes.

We used PSI-BLAST on the online webserver (https://blast.ncbi.nlm.nih.gov/), an

iterative search program that is more sensitive than the conventional protein BLAST[71],

to identify distant homologues of annotated genes. We additionally used

InterProScan[72,73] to make functional predictions for potentially novel proteins.

*Taxonomic classification of sequencing reads*

Taxonomic classification of reads was done using Kraken2 v2.1.2[29] with the '--paired' flag and using the Viral database maintained by Ben Langmead (7 June 2022 release; https://genome-idx.s3.amazonaws.com/kraken/k2_viral_20220607.tar.gz). This database comprises all genomes available on NCBI RefSeq as of June 2022. We then extracted reads assigned to each viral family (Supplementary Fig. 3a) or viral species (Supplementary Fig. 3b). To minimise the effects of potential read misclassifications, we applied abundance thresholds as described previously[74]. Briefly, we considered a taxon to be present if greater than 10 read pairs were assigned and if its relative abundance was greater than 0.005.

*Species niche modelling*

Bat occurrence records data were gathered from the online databases NBN Atlas (https://nbnatlas.org/) and GBIF (www.gbif.org). Records from year 2000-present were included, removing replicate records and those with high coordinate uncertainty. The number of occurrence points used for modelling ranged from 32 (*Myotis alcathoe*) to 16,403 (*Pipistrellus pipistrellus*). An initial 17 environmental variables were identified *a priori* to be important for predicting bat distributions. Nine were climatic variables averaged across 1980-2010 as described by Karger et al.[75], and were reduced to five variables using Variance Inflation Factor (VIF), retaining only those with a VIF < 0.5. These were mean annual air temperature, mean diurnal air temperature range, mean daily mean air temperature of the wettest quarter, precipitation seasonality and mean monthly precipitation amount of the warmest quarter. Four variables were derived from the UKCEH Land Cover Map 2019[76]. After merging similar land use classes, distance to woodland, distance to grassland, distance to arable and horticulture, and distance to urban were measured using Euclidean distance tools in ArcMap version 10.8. Two further distance variables were derived from Ordnance Survey polygons (2019, 2021): distance to the nearest road[77] and distance to the nearest river[78]. Elevation and slope were included to describe the topography of Great Britain, and were taken from the

LiDAR Composite Digital Terrain Model data at 10m resolution[79]. All spatial data were

subsequently reduced to 1000m resolution and projected to British National Grid.

An ensemble of five supervised binary classifiers was trained to predict the suitability of

a land area for each of the 17 UK bat species using the R package sdm[80]: random

forest (RF), maximum entropy (MaxEnt), multivariate adaptive regression splines

(MARS), boosted regression trees (BRT), and support vector machines (SVM).

Classifiers were trained to predict the relative probabilities of species occurrence based

on the 13 ecological variables described above, using the occurrence data for each

species and an equal number of randomly generated pseudo-absence data points

across the study area. Training and evaluation were performed using a 5-fold cross-

validation protocol, where a random subset comprising 80% of the dataset is used for

training and the remaining 20% use for the final evaluation. A final ensemble of all five

classifiers that were trained was used to generate the species distribution maps, with

the contribution of each individual classifier weighted based on its area under the

receiver operating characteristic curve (AUROC) score obtained during training. The

resultant species distribution maps indicate habitat suitability as a probability score for

each 1 km square grid on the study area, which ranges from 0 (unsuitable habitat) to 1

(suitable habitat). All models across all species performed well, with a median AUROC,

sensitivity and specificity of 0.827, 0.854, and 0.78, respectively. The individual species

distribution maps and model performance metrics are provided in Supplementary Fig.

11 and serve as a useful resource for future studies that seek to understand the

geographical range of UK bat species.

*Phylogenetic analyses*

To place the novel sequences within the global diversity of coronaviruses sequenced to

date, we computed alignment-free pairwise Mash distances using Mash v2.3[31] with a *k*-

mer length of 12, and reconstructed neighbour-joining trees[81] using the nj function from

the Ape v5.6.2 package in R (Fig. 1a). This alignment-free phylogenetic reconstruction

approach circumvents the challenge of aligning highly diverse sequences at the family

level, where high frequency of viral recombination may obscure true evolutionary

histories[82] and prevent dataset wide alignments. In accordance with previous work[83], we

rooted the neighbour-joining tree to a monophyletic *Deltacoronavirus* clade comprising

all 10 representative *Deltacoronavirus* genomes downloaded from NCBI RefSeq.

From this global phylogeny, we retrieved the pedacovirus ($n$=106), merbecovirus

($n$=113) and sarbecovirus genomes ($n$=534) most proximal to the novel assembled

genomes. We then aligned genomes from these subgenera separately using the Augur

v14.0.0[84] wrapper for MAFFT v7.490[85]. Genome positions where more than 20% of

sequences were assigned gaps were removed from the alignment. We subsequently

reconstructed finer-scale maximum-likelihood trees with IQTree v2.1.4-beta under a

GTR+G model, using ultrafast bootstrapping (UFBoot)[86] and approximate likelihood-

ratio tests (SH-aLRT)[87] with 1000 replicates. All phylogenetic trees were visualised

either using FigTree v1.4.4 or ggtree v3.2.1[88].

*Recombination analysis*

We selected 218 sarbecovirus genomes from the local sarbecovirus tree ($n$=534) by

retaining only one representative each for SARS-CoV (NC_004718) and SARS-CoV-2

(MW206198). We subsequently aligned these genomes via the same approach

described above but masked all positions with >20% of gaps by replacing the positions

with Ns, and removed gaps in the alignment relative to the genome used to root the

local sarbecovirus tree, NC_025217. This masked alignment was then analysed using

RDP v4.101[89]. Gene annotations for NC_025217 were obtained from GenBank and

used to annotate predicted recombinant positions.

*Spike protein homology and conservation of contact residues*

We extracted the Prokka-annotated spike protein sequences from our novel genomes

for further analysis. We peformed multiple sequence alignments of spike proteins from

our novel genomes and other sarbecoviruses that have been shown to bind human

ACE2[25,44,45] (BANAL-236, MZ937003.2; SARS-CoV-2, NC_045512.2; SARS-CoV-1,

NC_004718.3; Rs4084, KY417144.1; RsSHC014, KC881005.1; WIV1, KF367457.1;

Rs7327, KY417151.1; Rs4231, KY417146.1; LYRa11, KF569996.1; Pangolin GD-1,
EPI_ISL_410721; Pangolin GX-P2V, EPI_ISL_410542; RhGB01, MW719567.1; Khosta-
2, MZ190138.1; BtKY72, KY352407.1) using Mafft v7.490[85]. Subsequently, pairwise
amino acid similarity scores, visualisation of the alignments, and annotatation were
performed using the spike alignments using UGENE v42.0[90]. The accessions of all
genome records used in these analyses are provided in Supplementary Data 4.

*Pseudovirus assays*
To further test the capability of the coronaviruses we identified to infect human cells, we
synthesised human codon-optimised, Δ19-truncated (or equivalent) spike constructs in
pcDNA.3.1. The merbecovirus PaGB01 and pedacovirus PpiGB01 were additionally
synthesised with GSG-linker Myc tags for detection of spike incorporation into
pseudoparticles. Gene synthesis and codon optimisation was performed by GeneArt
(Thermo Fisher). Plasmids for human (*Homo sapiens*; BAB40370.1), least horseshoe
bat (*Rhinolophus pusillus*; ADN93477.1), Leschenault's rousette fruit bat (*Rousettus
leschenaultia*; BAF50705.1), and little brown bat (*Myotis lucifugus;* XP_023609438.1) in
pDisplay were used as previously described[40]. Additionally, Greater horseshoe bat
(*Rhinolophus ferrumequinum;* BAH02663.1) ACE2 was synthesised and cloned into
pDISPLAY for this study.

We maintained human embryonic kidney cells (HEK 293T; ATCC CRL-11268) and
human Hepatocyte carcinoma clone 5 (Huh7.5; C. Rice, Rockefeller University, New
York, NY) in complete media (DMEM, 10% FBS, 1% non-essential amino acids (NEAA)
and 1% penicillin-streptomycin (P/S)). Human lung cancer cells (Calu-3; ATCC HTB-55)
and Human epithelial colorectal adenocarcinoma cells (Caco-2; ATCC HTB-37) were
maintained in DMEM, 20% FBS, 1% NEAA and 1% P/S. All cells were kept at 5% $CO_2$,
37°C. 293T-hACE2 and Huh7.5-TMPRSS2 cells were generated by transducing HEK
293T or Huh7.5 cells with an ACE2 or TMPRSS2-expressing lentiviral vector, MT126[91]
and selecting with 2 µg ml$^{-1}$ puromycin or 4 mg ml$^{-1}$ G418; after selection, cells were
subsequently maintained with 1 µg ml$^{-1}$ puromycin or 2 mg ml$^{-1}$ G418, respectively.

Lentiviral based pseudotyped viruses were generated as previously described[49]. Briefly,
100 mm dishes of 293T cells were transfected using lipofectamine 3000 (Invitrogen)
with a mixture of 1 µg of the HIV packaging plasmid pCAGGs-GAG-POL, 1.5 µg of the
luciferase reporter construct (pCSFLW), and 1 µg of the plasmid encoding the spike or
glycoprotein of interest in pcDNA3.1. After 24 h supernatant was discarded and
replaced. PV-containing supernatants were collected at 48 and 72 h post-transfection,
passed through a 0.45 µM filter, and aliquoted and frozen at -80°C.

Pseudovirus entry assays were performed as previously described[49]. Briefly, 100 mm
dishes of 293T cells were transfected using lipofectamine 3000 (Invitrogen) with 2 µg of
the ACE2 encoding plasmid or empty vector. After 24 h, cells were resuspended by
scraping and plated into 96 well plates. Cells were overlaid with pseudovirus for 48 h
before lysis with reporter lysis buffer (Promega). Caco-2, Calu-3, and 293T-hACE2 cells
were seeded into 96 well plates. Cells were overlaid with pseudovirus for 48 h before
lysis with cell culture lysis buffer (Promega). We determined luciferase luminescence on
a FLUOstar Omega plate reader (BMF Labtech) using the Luciferase Assay System
(Promega). For all pseudovirus experiments, the amount of pseudovirus added was
standardised by quantifying p24 protein by western blot in a matched concentrated
pseudovirus stock.

We assessed expression of transfected receptors using Western blot assays. Cell
suspensions were pelleted by centrifugation at 200 x g for 7 min at 4°C, then
supernatant was removed. Cells were resuspended in 150 µl of cold
radioimmunoprecipitation assay (RIPA) buffer (Thermo Fisher) and incubated on ice for
30 min. Then, they were spun down at 3750 RPM for 30 min at 4°C. The protein-
containing supernatants were transferred to sterile Eppendorfs and frozen down at -
20°C. Before running a gel, 50 µl of 2-Mercaptoethanol (BME; Sigma) diluted 1:10 in 4X
Laemmli Sample Buffer (Bio-Rad, USA) was added to lysates and incubated at 80°C for
10 min.

To analyse incorporation of spike into the different sarbecovirus pseudoviruses, we
concentrated pseudovirus by ultracentrifugation at 100,000 x g for 2 h over a 20%
sucrose cushion.

In all experiments, we confirmed the successful expression of host receptors and spike
pseudoviruses using Western blot analyses (Supplementary Fig. 7c and d,
respectively). For Western blotting, membranes were probed with mouse anti-tubulin
(diluted 1/5,000; abcam; ab7291), mouse anti-p24 (diluted 1/2,000; abcam; ab9071),
rabbit anti-SARS spike protein (diluted 1/2,000; NOVUS; NB100-56578), rabbit anti-HA
tag (diluted 1/2000; abcam; ab9110), rabbit anti-ACE2 antibody (diluted 1/500; abcam;
ab15348), or rabbit anti-Myc tag (diluted 1/2000; abcam; ab9106). Near infra-red
secondary antibodies, IRDye[®] 680RD Goat anti-mouse (diluted 1/10,000; abcam;
ab216776), IRDye[®] 680RD Goat anti-rabbit (diluted 1/10,000; abcam; ab216777), were
subsequently used. Western blots were visualized using an Odyssey DLx Imaging
System (LI-COR Biosciences). The ACE2 constructs used in this work are from distinct
bat species and exhibited differing levels of expression or stability, as observed
previously[40]. Expression levels of ACE2 do not correlate with efficiency of cell entry. All
raw, uncropped western blot images are provided in Supplementary Fig. 12.

*Alphafold2 (ColabFold) structural analysis*

The protein structure model of the RhGB07 RBD was predicted using Alphafold2 as
implemented in ColabFold[92]. Default settings were used. The top ranked model was
used for all analyses. Structural representations and calculations were done within
ChimeraX[93,94]. RMSD values for structural superpositions were calculated using the
matchmaker command. Reported values represent the RMSD of all $C_\alpha$ atoms. Buried
surface area calculations were performed using the measure buriedarea command**.**

*Biolayer Interferometry (BLI)*

The RhGB07 spike trimer was designed to mimic the native trimeric conformation of the
protein. It consists of a gene synthesized by Genscript of CHO codon-optimized

sequence of RhGB07, residues 1-1191, preceded by a u-phosphatase signal peptide[95],
residues 969 and 970 mutated to proline (2P) to stabilize the prefusion state of the spike
trimer, a putative basic site that may be the site of proteolysis (RAKQ, residues 669-
672, was mutated to GASQ), a C-terminal T4 foldon fusion domain to stabilize the trimer
complex, followed by C-terminal 8x His and 2x Strep tags for affinity purification. This
gene was cloned with the pcDNA3.1(+) vector. The trimeric RhGB07 spike protein was
expressed as previously reported as for the SARS-CoV-2 spike transiently expressed in
suspension-adapted ExpiCHO cells (Thermo Fisher) in ProCHO5 medium (Lonza) at 5
$\times 10^6$ cells/mL using PEI MAX (Polysciences) for DNA delivery[96]. At 1 h post-
transfection, dimethyl sulfoxide (DMSO; AppliChem) was added to 2% (v/v). Following a
7-day incubation with agitation at 31 °C and 4.5% $CO_2$, the cell culture medium was
harvested and clarified using a 0.22 µm filter. The conditioned medium was loaded onto
Streptactin XT columns (IBA) washed with PBS and eluted with 50 mM biotin in 150 mM
NaCl, 100 mM HEPES 7.5. Eluted protein was then dialyzed overnight into PBS. The
purity of spike trimers was determined to be >99% pure by SDS-PAGE analysis.
753
Human (residues 19-615), little brown bat (19-629) and greater horseshoe bat (19-615)
ACE2 genes were synthesized by Genscript and cloned in after the human pregnancy
specific glycoprotein 1 signal peptide and is followed by a 3C protease cleavage site, a
mouse IgG2a Fc fragment and a 10x His tag (only for the hACE2 construct). Protein
production was produced exactly as for the RhGB07 spike. The filtered conditioned
media was then subjected to Protein A purification. Eluted protein was dialyzed into
PBS.
761
Experiments were performed on a Gator BLI system. Running buffer was 1X PBS.
Dimeric mFc-hACE2 and bat ACE2 were diluted to 10 µg/mL and captured with MFc
tips (GatorBio). Loaded tips were dipped into 2-fold serial dilution series (highest
concentration 3000 nM) of the RhGB07 spike protein. Curves were processed using the
Gator software with a 1:1 fit after background subtraction. Plots were generated in
Prism v9.
768

769 *Data analysis and visualisation*

770 All data analyses were performed using R v4.1.0 or Python v3.9.12. Visualizations were
771 performed using ggplot v3.3.5[97]. Genomic and protein sequence data were analysed
772 using Biostrings v2.62.0[98]. Supplementary Fig. 5 was made using Adobe Illustrator
773 v27.1.1 and Geneious v11.1.5 (https://www.geneious.com).
774

**Data availability**

All novel genomes are available in NCBI GenBank under the accessions OQ401247-
OQ401251, and OQ401253-OQ401255 (BioProject accession PRJNA929706). The raw
sequencing reads generated and analysed in this study have also been uploaded to the
SRA under the accessions SRX19257406- SRX19257414. All GenBank and GISAID
accessions for the sequences included in the coronavirus database are provided in
Supplementary Data 4. Other sequences used are as follows: MHV, AY700211.1;
BANAL-236, MZ937003.2; SARS-CoV-2, NC_045512.2; SARS-CoV-1, NC_004718.3;
Rs4084, KY417144.1; RsSHC014, KC881005.1; WIV1, KF367457.1; Rs7327,
KY417151.1; Rs4231, KY417146.1; LYRa11, KF569996.1; Pangolin GD-1,
EPI_ISL_410721; Pangolin GX-P2V, EPI_ISL_410542; RhGB01, MW719567.1; Khosta-
2, MZ190138.1; BtKY72, KY352407.1. The NBN Atlas datasets used for species niche
modelling are listed in Supplementary Data 5.
788

789 **Code availability**

790 All custom code used to perform the analyses reported here are hosted on GitHub
791 (https://github.com/cednotsed/bat-CoVs.git)[99].
792

**References**

1.  Taylor, L. H., Latham, S. M. & Woolhouse, M. E. J. Risk factors for human disease

emergence. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **356**, 983–989 (2001).

2.  Jones, K. E. *et al.* Global trends in emerging infectious diseases. *Nature* **451**, 990–

993 (2008).

3.  Peiris, J. S., Guan, Y. & Yuen, K. Severe acute respiratory syndrome. *Nat. Med.* **10**,

S88–S97 (2004).

4.  Memish, Z. A. *et al.* Human infection with MERS coronavirus after exposure to

infected camels, Saudi Arabia, 2013. *Emerg. Infect. Dis.* **20**, 1012–1015 (2014).

5.  Balloux, F. *et al.* The past, current and future epidemiological dynamic of SARS-

CoV-2. *Oxf. Open Immunol.* **3**, iqac003 (2022).

6.  Ye, Z.-W. *et al.* Zoonotic origins of human coronaviruses. *Int. J. Biol. Sci.* **16**, 1686

(2020).

7.  Corman, V. M., Muth, D., Niemeyer, D. & Drosten, C. Hosts and sources of endemic

human coronaviruses. *Adv. Virus Res.* **100**, 163–188 (2018).

8.  Temmam, S. *et al.* Coronaviruses with a SARS-CoV-2-like receptor-binding domain

allowing ACE2-mediated entry into human cells isolated from bats of Indochinese

peninsula. (2021).

9.  Kumakamba, C. *et al.* Coronavirus surveillance in wildlife from two Congo basin

countries detects RNA of multiple species circulating in bats and rodents. *PloS One*

**16**, e0236971 (2021).

10. Wacharapluesadee, S. *et al.* Evidence for SARS-CoV-2 related coronaviruses

circulating in bats and pangolins in Southeast Asia. *Nat. Commun.* **12**, 972 (2021).

11. Crook, J. M. *et al.* Metagenomic identification of a new sarbecovirus from horseshoe

bats in Europe. *Sci. Rep.* **11**, 1–9 (2021).

12. Yadav, P. D. *et al.* Detection of coronaviruses in Pteropus & Rousettus species of

bats from different States of India. *Indian J. Med. Res.* **151**, 226 (2020).

13. Valitutto, M. T. *et al.* Detection of novel coronaviruses in bats in Myanmar. *PloS One*

**15**, e0230802 (2020).

14. Rizzo, F. *et al.* Coronavirus and paramyxovirus in bats from Northwest Italy. *BMC*

*Vet. Res.* **13**, 1–11 (2017).

15. Lin, X.-D. *et al.* Extensive diversity of coronaviruses in bats from China. *Virology*

**507**, 1–10 (2017).

16. Waruhiu, C. *et al.* Molecular detection of viruses in Kenyan bats and discovery of

novel astroviruses, caliciviruses and rotaviruses. *Virol. Sin.* **32**, 101–114 (2017).

17. Shehata, M. M. *et al.* Surveillance for coronaviruses in bats, Lebanon and Egypt,

2013–2015. *Emerg. Infect. Dis.* **22**, 148 (2016).

18. Kemenesi, G. *et al.* Molecular survey of RNA viruses in Hungarian bats: discovering

novel astroviruses, coronaviruses, and caliciviruses. *Vector-Borne Zoonotic Dis.* **14**,

846–855 (2014).

19. August, T. A., Mathews, F. & Nunn, M. A. Alphacoronavirus detected in bats in the

United Kingdom. *Vector-Borne Zoonotic Dis.* **12**, 530–533 (2012).

20. De Souza Luna, L. K. *et al.* Generic Detection of Coronaviruses and Differentiation

at the Prototype Strain Level by Reverse Transcription-PCR and Nonfluorescent

Low-Density Microarray. *J. Clin. Microbiol.* **45**, 1049 (2007).

21. Xiu, L. *et al.* A RT-PCR assay for the detection of coronaviruses from four genera. *J.*

*Clin. Virol.* **128**, 104391 (2020).

22. Watanabe, S. *et al.* Bat coronaviruses and experimental infection of bats, the

Philippines. *Emerg. Infect. Dis.* **16**, 1217 (2010).

23. Vijgen, L., Moës, E., Keyaerts, E., Li, S. & Ranst, M. V. A pancoronavirus RT-PCR

assay for detection of all known coronaviruses. in *SARS-and Other Coronaviruses*

3–12 (Springer, 2008).

24. Holbrook, M. G. *et al.* Updated and validated pan-coronavirus PCR assay to detect

all coronavirus genera. *Viruses* **13**, 599 (2021).

25. Temmam, S. *et al.* Bat coronaviruses related to SARS-CoV-2 and infectious for

human cells. *Nature* **604**, 330–336 (2022).

26. Chen, J. *et al.* A bat MERS-like coronavirus circulates in pangolins and utilizes

human DPP4 and host proteases for cell entry. *Cell* **186**, 850–863 (2023).

27. Hassell, J. M., Begon, M., Ward, M. J. & Fèvre, E. M. Urbanization and disease

emergence: dynamics at the wildlife–livestock–human interface. *Trends Ecol. Evol.*

**32**, 55–67 (2017).

28. Plowright, R. K. *et al.* Pathways to zoonotic spillover. *Nat. Rev. Microbiol.* **15**, 502–

510 (2017).

29. Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2.

*Genome Biol.* **20**, 257 (2019).

30. Nayfach, S. *et al.* CheckV assesses the quality and completeness of metagenome-

assembled viral genomes. *Nat. Biotechnol.* **39**, 578–585 (2021).

31. Ondov, B. D. *et al.* Mash: fast genome and metagenome distance estimation using

MinHash. *Genome Biol.* **17**, 132 (2016).

32. Crook, J. M. *et al.* Metagenomic identification of a new sarbecovirus from horseshoe

bats in Europe. *Sci. Rep.* **11**, 1–9 (2021).

33. Kreuder Johnson, C. *et al.* Spillover and pandemic properties of zoonotic viruses

with high host plasticity. *Sci. Rep.* **5**, 14830 (2015).

34. Woolhouse, M. E. J. & Gowtage-Sequeria, S. Host range and emerging and

reemerging pathogens. *Emerg. Infect. Dis.* **11**, 1842 (2005).

35. Winter, R., Mantilla-Contreras, J. & Schmidt, S. Usage of buildings in the life cycle of

two endangered Rhinolophus species in the Mediterranean region: implications for

roost protection. *Eur. J. Wildl. Res.* **66**, 1–13 (2020).

36. Temmam, S. *et al.* Coronaviruses with a SARS-CoV-2-like receptor-binding domain

allowing ACE2-mediated entry into human cells isolated from bats of Indochinese

peninsula. (2021).

37. Liu, K. *et al.* Binding and molecular basis of the bat coronavirus RaTG13 virus to

ACE2 in humans and other species. *Cell* **184**, 3438–3451 (2021).

38. Park, J.-E. *et al.* Proteolytic processing of Middle East respiratory syndrome

coronavirus spikes expands virus tropism. *Proc. Natl. Acad. Sci.* **113**, 12262–12267

(2016).

39. Bertram, S. *et al.* TMPRSS2 activates the human coronavirus 229E for cathepsin-

independent host cell entry and is expressed in viral target cells in the respiratory

epithelium. *J. Virol.* **87**, 6150–6160 (2013).

40. Conceicao, C. *et al.* The SARS-CoV-2 Spike protein has a broad tropism for

mammalian ACE2 proteins. (2020) doi:10.1371/journal.pbio.3001016.

41. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature*

**596**, 583–589 (2021).

42. Lan, J. *et al.* Structure of the SARS-CoV-2 spike receptor-binding domain bound to

the ACE2 receptor. *Nature* **581**, 215–220 (2020).

43. Li, F., Li, W., Farzan, M. & Harrison, S. C. Structure of SARS coronavirus spike

receptor-binding domain complexed with receptor. *Science* **309**, 1864–1868 (2005).

44. Seifert, S. N. *et al.* An ACE2-dependent Sarbecovirus in Russian bats is resistant to

SARS-CoV-2 vaccines. *PLoS Pathog.* **18**, e1010828 (2022).

45. Starr, T. N. *et al.* ACE2 binding is an ancestral and evolvable trait of sarbecoviruses.

*Nature* **603**, 913–918 (2022).

46. Roelle, S. M., Shukla, N., Pham, A. T., Bruchez, A. M. & Matreyek, K. A. Expanded

ACE2 dependencies of diverse SARS-like coronavirus receptor binding domains.

*PLoS Biol.* **20**, e3001738 (2022).

47. Starr, T. N. *et al.* Deep Mutational Scanning of SARS-CoV-2 Receptor Binding

Domain Reveals Constraints on Folding and ACE2 Binding. *Cell* **182**, 1295-

1310.e20 (2020).

48. Sander, A.-L. *et al.* Genomic determinants of Furin cleavage in diverse European

SARS-related bat coronaviruses. *Commun. Biol.* **5**, 1–8 (2022).

49. Peacock, T. P. *et al.* The furin cleavage site in the SARS-CoV-2 spike protein is

required for transmission in ferrets. *Nat. Microbiol. 2021 67* **6**, 899–909 (2021).

50. Alkhovsky, S. *et al.* SARS-like coronaviruses in horseshoe bats (Rhinolophus spp.)

in Russia, 2020. *Viruses* **14**, 113 (2022).

51. Tao, Y. *et al.* Surveillance of bat coronaviruses in Kenya identifies relatives of

human coronaviruses NL63 and 229E and their recombination history. *J. Virol.* **91**,

(2017).

52. Lemmin, T., Kalbermatter, D., Harder, D., Plattet, P. & Fotiadis, D. Structures and

dynamics of the novel S1/S2 protease cleavage site loop of the SARS-CoV-2 spike

glycoprotein. *J. Struct. Biol. X* **4**, 100038 (2020).

53. Wrapp, D. *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion

conformation. *Science* **367**, 1260–1263 (2020).

54. Zech, F. *et al.* Spike mutation T403R allows bat coronavirus RaTG13 to use human

ACE2. *bioRxiv* 2021.05.31.446386 (2021) doi:10.1101/2021.05.31.446386.

55. Wu, Y. & Zhao, S. Furin cleavage sites naturally occur in coronaviruses. *Stem Cell*

*Res.* **50**, 102115 (2021).

56. Boni, M. F. *et al.* Evolutionary origins of the SARS-CoV-2 sarbecovirus lineage

responsible for the COVID-19 pandemic. *Nat. Microbiol.* **5**, 1408–1417 (2020).

57. Bobay, L.-M., O'Donnell, A. C. & Ochman, H. Recombination events are

concentrated in the spike protein region of Betacoronaviruses. *PLoS Genet.* **16**,

e1009272 (2020).

58. Jones, S. *et al.* Testing bats in rehabilitation for SARS-CoV-2 before release into the

wild. *Conserv. Sci. Pract.* e12707 (2022) doi:10.1111/CSP2.12707.

59. Menachery, V. D. *et al.* Trypsin treatment unlocks barrier for zoonotic bat

coronavirus infection. *J. Virol.* **94**, e01774-19 (2020).

60. Boni, M. F. *et al.* Evolutionary origins of the SARS-CoV-2 sarbecovirus lineage

responsible for the COVID-19 pandemic. *Nat. Microbiol.* **5**, 1408–1417 (2020).

61. Kunz, T. H., Braun de Torrez, E., Bauer, D., Lobova, T. & Fleming, T. H. Ecosystem

services provided by bats. *Ann. N. Y. Acad. Sci.* **1223**, 1–38 (2011).

62. Bat Conservation Trust. *The National Bat Monitoring Programme Annual Report*

*2021.* (2021).

63. Mathews, F. & Harrower, C. IUCN–compliant Red List for Britain's Terrestrial

Mammals. Assessment by the Mammal Society under contract to Natural England,

Natural Resources Wales and Scottish Natural Heritage. (2020).

64. Gibb, R. *et al.* Zoonotic host diversity increases in human-dominated ecosystems.

*Nature* **584**, 398–402 (2020).

65. Eby, P. *et al.* Pathogen spillover driven by rapid changes in bat ecology. *Nature* 1–3

(2022).

66. Viana, M. *et al.* Effects of culling vampire bats on the spatial spread and spillover of

rabies virus. *Sci. Adv.* **9**, eadd7437 (2023).

67. Khare, S. *et al.* GISAID's Role in Pandemic Response. *China CDC Wkly.* **3**, 1049–

1051 (2021).

68. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat.*

*Methods* **9**, 357–359 (2012).

69. Meleshko, D., Hajirasouliha, I. & Korobeynikov, A. coronaSPAdes: from biosynthetic

gene clusters to RNA viral assemblies. *Bioinformatics* **38**, 1–8 (2021).

70. Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**,

2068–2069 (2014).

71. Altschul, S. F. *et al.* Gapped BLAST and PSI-BLAST: a new generation of protein

database search programs. *Nucleic Acids Res.* **25**, 3389–3402 (1997).

72. Blum, M. *et al.* The InterPro protein families and domains database: 20 years on.

*Nucleic Acids Res.* **49**, D344–D354 (2021).

73. Zdobnov, E. M. & Apweiler, R. InterProScan–an integration platform for the

signature-recognition methods in InterPro. *Bioinformatics* **17**, 847–848 (2001).

74. Tan, C. C. S. *et al.* No evidence for a common blood microbiome based on a

population study of 9,770 healthy humans. *bioRxiv* 2022.07.29.502098 (2022)

doi:10.1101/2022.07.29.502098.

75. Karger, D. N. *et al.* Climatologies at high resolution for the earth's land surface

areas. *Sci. Data* **4**, 1–20 (2017).

76. Morton, R. D., Marston, C. G., O'Neil, A. W. & Rowland, C. S. Land Cover Map 2019

(land parcels, GB). (2020).

77. Ordnance Survey. OS Open Roads. ESRI® Shapefile. (2021).

78. Ordnance Survey. OS Open Rivers. ESRI® Shapefile. (2019).

79. Defra Data Services Platform. LIDAR Composite DTM 2019 – 10m. (2019).

80. Naimi, B. & Araújo, M. B. sdm: a reproducible and extensible R platform for species

distribution modelling. *Ecography* **39**, 368–375 (2016).

81. Saitou, N. & Nei, M. The neighbor-joining method: a new method for reconstructing

phylogenetic trees. *Mol. Biol. Evol.* **4**, 406–425 (1987).

82. Zielezinski, A., Vinga, S., Almeida, J. & Karlowski, W. M. Alignment-free sequence

comparison: benefits, applications, and tools. *Genome Biol.* **18**, 186 (2017).

83. Tan, C. C. S. *et al.* Pre-existing T cell-mediated cross-reactivity to SARS-CoV-2

cannot solely be explained by prior exposure to endemic human coronaviruses.

*Infect. Genet. Evol.* **95**, 105075 (2021).

84. Huddleston, J. *et al.* Augur: a bioinformatics toolkit for phylogenetic analyses of

human pathogens. *J. Open Source Softw.* **6**, 2906 (2021).

85. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version

7: Improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

86. Minh, B. Q., Nguyen, M. A. T. & von Haeseler, A. Ultrafast Approximation for

Phylogenetic Bootstrap. *Mol. Biol. Evol.* **30**, 1188–1195 (2013).

87. Guindon, S. *et al.* New Algorithms and Methods to Estimate Maximum-Likelihood

Phylogenies: Assessing the Performance of PhyML 3.0. *Syst. Biol.* **59**, 307–321

(2010).

88. Yu, G., Smith, D. K., Zhu, H., Guan, Y. & Lam, T. T. ggtree: an R package for

visualization and annotation of phylogenetic trees with their covariates and other

associated data. *Methods Ecol. Evol.* **8**, 28–36 (2017).

89. Martin, D. P., Murrell, B., Golden, M., Khoosal, A. & Muhire, B. RDP4: Detection and

analysis of recombination patterns in virus genomes. *Virus Evol.* **1**, (2015).

90. Okonechnikov, K. *et al.* Unipro UGENE: a unified bioinformatics toolkit.

*Bioinformatics* **28**, 1166–1167 (2012).

91. Rebendenne, A. *et al.* SARS-CoV-2 triggers an MDA-5-dependent interferon

response which is unable to control replication in lung epithelial cells. *J. Virol.* **95**,

e02415-20 (2021).

92. Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat. Methods*
**19**, 679–682 (2022).

93. Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers,
educators, and developers. *Protein Sci.* **30**, 70–82 (2021).

94. Goddard, T. D. *et al.* UCSF ChimeraX: Meeting modern challenges in visualization
and analysis. *Protein Sci.* **27**, 14–25 (2018).

95. Wrobel, A. G. *et al.* SARS-CoV-2 and bat RaTG13 spike glycoprotein structures
inform on virus evolution and furin-cleavage effects. *Nat. Struct. Mol. Biol.* **27**, 763–
767 (2020).

96. Fenwick, C. *et al.* Patient-derived monoclonal antibody neutralizes SARS-CoV-2
Omicron variants and confers full protection in monkeys. *Nat. Microbiol.* **7**, 1376–
1389 (2022).

97. Wickham, H. ggplot2. *Wiley Interdiscip. Rev. Comput. Stat.* **3**, 180–185 (2011).

98. Pagès, H., Aboyoun, P., Gentleman, R. & DebRoy, S. Biostrings: Efficient
manipulation of biological strings. *R Package Version* **2**, 10–18129 (2019).

99. Tan, Cedric. Source code: bat-CoVs. *GitHub* doi:10.5281/zenodo.7852230.

1010

**Author contributions statement**

VS and FB contributed equally and co-supervised the research. VS, CC, ER, GW and
TB wrote the grant application that supported this research. Primary analysis on the
project was carried out by CCST, TPP, KYM, KL and CH with contributions from FB,
LvD, WDP, DO, and WB. CCST wrote the initial draft of manuscript, with subsequent
rounds of editing from VS, FB and LvD. All authors provided intellectual contributions to
the manuscript.

## Competing interests statement

Authors declare that to current knowledge, there are no legal, financial or personal competing interests.

## Acknowledgments

We would like to acknowledge the following bat rehabilitators and conservationists for the collection of bat samples that were crucial for this study: Daniel Whitby, Dorset Bat Group, Claire Andrews, Caitlin Woodfield, Emma Turnbull, Elaine Charlson, Rose-Ann Movsovic, Margaret Grimsey, Hazel Ryan, Gareth Harris, Danielle Linton, Sam Smith, Ross Baker, Lynn Whitfield, Colleen Hope, Josh Sowden, Emily Dickens, Tricia Scott, Jonah Tosney, Eilish Rothney, Dale Irvine, Joe Salkeld, Alice Samuel, Amanda Millar, Sheila Wright, Stewart Rowden, Joy Hall, Jessica Dangerfield, Catherine Wood, Rachael Tarlinton, Ternenge Apaa, and Fiona Mathews. We thank Rachael Tarlinton, Ternenge Apaa, and Fiona Mathews for comments on the manuscript, Scott Jones, Faye Hobbs and Danielle Harris for assisting with sample processing, and Xavier Didelot for advice on genetic recombination analyses. We also thank all who had previously deposited on NCBI GenBank and GISAID the genomes used here (Supplementary Data 4 and 6, respectively), as well as the NERC Omics facility for sequencing. VS, CC, ER, GW and TB were funded by NERC (UKRI Covid Urgency grant). FB, LvD and DR are funded by the European Commission (Horizon 2021-2024, END-VOC Project). Views and opinions expressed are however those of the authors only and do not necessarily reflect those of the European Union or the European Health and Digital Executive Agency. For the purpose of open access, the corresponding author has applied a 'Creative Commons Attribution' (CC BY) licence to any Author Accepted Manuscript version arising.

 **Tables**

 **Table 1: Summary statistics for novel coronavirus genomes assembled in this**

 **study**

| Host species | Common name | Genome name | Subgenus | Length (bp) | Closest hit | | BLASTn identity (%) | Prop. of query aligned (%) | CheckV completeness | Median coverage |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | Accession | Name | | | | |
| *Rhinolophus ferrumequinum* | Greater horseshoe bat | RfGB01 | *Sarbecovirus* | 29308 | MW719567 | RhGB01 | 98.1 | 99.7 | 97 | 689 |
| | | RfGB02 | *Sarbecovirus* | 29375 | MW719567 | RhGB01 | 98.1 | 99.4 | 97 | 7178 |
| *Rhinolophus hipposideros* | Lesser horseshoe bat | RhGB07 | *Sarbecovirus* | 29224 | MW719567 | RhGB01 | 97.9 | 100 | 97 | 3809 |
| | | RhGB08 | *Sarbecovirus* | 29217 | MW719567 | RhGB01 | 98 | 100 | 96 | 548 |
| *Plecotus auritus* | Brown long-eared bat | PaGB01 | *Merbecovirus* | 30018 | MG596803 | *P. kuhlii* MERS-related CoV | 81.5 | 99.7 | 99 | 6237.5 |
| *Pipistrellus pipistrellus* | Common pipistrelle | PpiGB01 | *Pedacovirus* | 28247 | MN535731 | *M. daubentonii* pedacovirus | 80.8 | 82.9 | 100 | 7438 |
| *Myotis daubentonii* | Daubenton's bat | MdGB01 | *Pedacovirus* | 28224 | MN535731 | *M. daubentonii* pedacovirus | 95.5 | 99.6 | 100 | 7938 |
| | | MdGB02 | *Pedacovirus* | 28010 | MN535733 | *M. daubentonii* pedacovirus | 95.4 | 99.8 | 100 | 7874 |
| | | MdGB03 | *Pedacovirus* | 28227 | MN535733 | *M. daubentonii* pedacovirus | 95.4 | 99.8 | 100 | 7958 |

**Figure legends**

1049

**Fig. 1. Phylogenetic placement of novel coronaviruses.** (a) Alignment-free phylogeny of the global
diversity of coronavirus genomes ($n$=2118) and our nine novel genomes. Host genus (inner ring) and their
broader host groups (outer ring) are annotated. Local maximum likelihood trees of (b) pedacoviruses
($n$=106), (c) merbecoviruses ($n$=113) and (d) sarbecoviruses ($n$=534).

**Fig. 2. RhGB07 can bind and use human ACE2 for cell entry *in vitro*.** (a) Entry of different spike pseudoviruses expressing viral glycoproteins into HEK293T cells transfected with (a) human receptors known to allow entry of human coronaviruses or (e) ACE2 homologues from different species. For (a) and (e), the raw entry values for each pseudoviruses were normalised by their entry into cells transfected with a vector containing no receptor sequence (i.e., 'empty'). The raw entry values of representative repeats (*n*=3 independent experiments) are also shown for direct comparisons of absolute entry. Representative bio-layer interferometry binding curves showing the association and dissociation of SARS-CoV-2 and RhGB07 spike proteins with (b) hACE2 (*n*=2 and 3 independent experiments, at three and seven protein concentrations, respectively), or (f) with *R. ferrumequinum* or *M. lucifugus* ACE2 (*n*=1 independent experiment, at seven protein concentrations). (c) Entry of pseudoviruses into different 'normal' human cell lines that stably express lower or physiological levels of hACE2. All entry measurements are normalised to those for the 'bald' pseudovirus not expressing any spike protein. (d) Entry of pseudoviruses into Huh7.5 cells transduced with a human TMPRSS2 vector, normalised to 'bald'. Data from panels (a), (c), (d) and (e) are compiled from *n*=3-8 independent experiments and plotted as mean + s.d.. Statistical

significance was determined by (a, e) two-way ANOVA or (c, d) one-way ANOVA on log-transformed data (after determining log normality by the Shapiro–Wilk test and QQ plot) with multiple comparisons against 'empty' vector or 'bald' pseudovirus, respectively. *$0.05 \geq p > 0.01$; **$0.01 \geq p > 0.001$; ***$0.001 \geq p > 0.0001$; ****$p \leq 0.0001$. Exact p values annotated for each graph are as follows (from left to right), (a) <0.0001, <0.0001, <0.0001, <0.0001; (c, Calu-3) <0.0001, <0.0001, 0.001, <0.0001, <0.0001; (c, Caco-2) <0.0001, <0.0001, 0.022, <0.0001, <0.0001; (c, HEK 293T-ACE2) <0.0001, <0.0001, <0.0001; (d) <0.0001, <0.0001, <0.0001. (e, SARS-CoV-2) <0.0001, 0.0003, 0.12, <0.0001, <0.0001 (e, BANAL-20-52) <0.0001, <0.0001, <0.0001, <0.0001, <0.0001; (e RatG13) <0.0001, <0.0001, <0.0001, <0.0001 (e, RhGB07) <0.0001, <0.0001; (e, RfGB02) <0.0001.

**Fig. 3. Structural and sequence features of RhGB01-like sarbecoviruses.** (a) The solved RBD structures of SARS-CoV-2, RaTG13, BANAL-236 (close relative of BANAL-20-52[8]) and the AlphaFold-

predicted structure of RhGB07 were superposed. (b) The 3D surfaces of the RBD-hACE2 binding

interface for SARS-CoV-2 and RhGB07. Alignment of sarbecovirus spike proteins showing the

conservation of key contact residues involved interactions between (c) SARS-CoV-1 spike and (d) SARS-

CoV-2 spike with hACE2. The sequences shown in the alignments are from Asian, European and African

sarbecoviruses that have been shown to bind hACE2[44–46]. These sequences were ordered based on their

genetic relatedness, as inferred from a consensus maximum likelihood phylogenetic tree reconstructed

from their whole genomes (bottom left).