Deprivation of arginine and lysine interferes with growth of patient-derived tumor xenografts

Wolkersdorfer A.M.[1], Bergmann B.[2], Adelmann J.[3], Ebbinghaus, M.[4], Günther, E.[5], Gutmann M.[1], Hahn L.[1], Hurwitz R.[6], Lühmann T.[1]., Schueler J.[4], Schmidt L.[1], Teifel M.[5], Meinel L.[1,7,*], Rudel T.[2,7,*]

[1] Institute of Pharmacy and Food Chemistry, University of Würzburg, Würzburg, Germany

[2] Chair of Microbiology, Biocentre, University of Würzburg, Würzburg, Germany

[3] Institute of Organic Chemistry, University of Würzburg, Würzburg, Germany

[4] Charles River Laboratories Germany GmbH, Am Flughafen 12–14, Freiburg, Germany

[5] Aeterna Zentaris GmbH / Weismuellerstr. 50, Frankfurt am Main, Germany

[6] Max Planck Institute for Infection Biology, Berlin, Germany

[7] Helmholtz-Institute for RNA-based Infection Research (HIRI), Würzburg, Germany

Running title: Tumor therapy by amino acid depletion

Keywords: L-amino acid oxidase, PEGylation, arginine-auxothrophy, arginine lysine starvation, tumor therapy, patient-derived tumor

Funding Source:

Aeterna Zentaris GmbH, Frankfurt am Main, Germany (in part)

Conflicts of Interest:

The authors declare no potential conflicts of interest.

*Corresponding authors:

Prof. Dr. Thomas Rudel, Chair of Microbiology, Biocentre, University of Wuerzburg, Am Hubland, DE-97074 Wuerzburg, Germany, E-Mail: thomas.rudel@uni-wuerzburg.de

Prof. Dr. Lorenz Meinel, Institute of Pharmacy and Food Chemistry, University of Würzburg, Würzburg, Germany, E-Mail: lorenz.meinel@uni-wuerzburg.de

Abstract

In recent years, a novel treatment method for cancer emerged, which is based on the starvation of tumors of amino acids like arginine. The deprivation of arginine in serum is based on enzymatic degradation and can be realized by arginine deaminases like the L-amino acid oxidase found in the ink toxin of the sea hare *Aplysia punctata*. Previously isolated from the ink, the L-amino acids oxidase was described to oxidate the essential amino acid L-lysine and L-arginine to their corresponding deaminated alpha-keto acids. Here, we present the recombinant production and functionalization of amino acid oxidase *Aplysia Punctata* ink toxin (APIT). PEGylated APIT (APIT-PEG) increased the blood circulation time. APIT-PEG treatment of patient-derived xenografted mice shows a significant dose dependent reduction of tumor growth over time mediated by amino acid starvation of the tumor. Treatment of mice with APIT-PEG which lead to deprivation of arginine was well tolerated.

Significance: Inhibiting the growth of human head and neck cancer in a mouse model by the depletion of amino acids L-lysine and L-arginine provides proof of principle for tumor therapy based on APIT-PEG.

Introduction

Increasing resistance of tumors towards chemotherapies can be observed with standard of care DNA-damaging agents, antimetabolites, mitotic inhibitors, nucleotide analogous or inhibitors of topoisomerase.[1] The cytotoxic effect of the drugs is primarily by the induction of apoptosis. In treated tumor cells mutations of the general apoptosis signaling pathways can be observed, which leads to broad resistance to drug treatment due to the lag of activation of the apoptotic pathway.[2] This constitutes the need for alternative anti-tumor therapy approaches to tackle these limitations of current anti-tumor treatments.

An increasingly recognized strategy is the enzymatic depletion of amino acids in serum to target metabolic deficient malignances. A prime example of amino acid depletion cancer therapy is the treatment of leukemias with asparaginase. Normal cells convert aspartate to asparagine; however, this pathway is insufficient in leukemic cells. Treatment with asparaginase cures more than 90% of pediatric acute lymphoblastic leukemia patients.[3, 4]

Another highly discussed therapeutic approach is the depletion of arginine in arginine auxotroph tumors. Auxotrophy to the semi-essential amino acid arginine occurs through epigenetic silencing of *arginosuccinate synthetase 1* (ASS1) or *arginine lyase (AL)* genes, preventing the synthesis of arginine through citrulline in the urea cycle.[5-8] In many cancer types arginine-auxotrophy reaches an average of 60%-100%.[9] ASS1-deficient cells have a generally poor survival rate and an increased requirement for aspartate used for pyrimidine synthesis to support the proliferation of cells .[10-17]. This property can be therapeutically exploited by treating these tumor cells with **a**rginine **dei**minases (ADI), which deaminate arginine to citrulline and thereby deplete arginine. In combination with arginine auxotrophy, cells are then starved from arginine and undergo apoptotic cell death.[17, 18] Therefore, a novel strategy for targeting auxotrophic cancers by starvation of the tumor of arginine emerged.[19]

A promising example for this treatment method is ADI-PEG20. First clinical trials with a mycoplasma-derived PEGylated ADI demonstrated the sensitivity of tumors towards arginine deprivation.[20, 21] Based on this finding the FDA and the EMA granted orphan drug designation for ADI-PEG20 for the treatment of hepatocellular carcinoma and melanoma. Phase I clinical trials combined arginine deprivation with conventional cytostatically chemotherapy (cisplatin, pemetrexed-cisplatin, Nab-Paclitaxel with Gemcitabine, FOXLFOX6).[22-25] In the studies patients showed an increase of immunogenic reaction against ADI-PEG20 after few weeks, which decreased the effectiveness of the combination therapy.

To overcome these limitations, the necessity of alternative products for ADI-PEG20 emerges

including an arginine and lysine specific **L-a**mino **a**cid **o**xidases (LAAO) in the defense ink of

Aplysia punctata (*Aplysia punctata* **i**nk **t**oxin = APIT). APIT was first isolated from crude ink,

which deaminates L-lysine and L-arginine to their corresponding α-keto acids.(26) APITs

tumor specific cytotoxicity was demonstrated on apoptosis-resistant tumor cell.(27) However,

in preliminary *in vivo* studies, intravenous injection of APIT into mice had an unfavorable renal

clearance shortly after application.(28) To increase serum half-life, polymer modifications such

as the coupling of **p**oly**e**thylene **g**lycol (PEG) to the protein are well-known strategies.

Unspecific PEGylation of APIT, in which the lysine residues of the enzyme are targeted by N-

hydroxysuccinimidyl chemistry, was performed, leading to increased serum half-life and

exhibited comparable anticancer activity as ADI-PEG20.(28)

In this study, the effect of arginine and lysine deprivation mediated by APIT-PEG treatment

was tested in the patient-derived HNXF 536 head and neck xenograft model (PDX). The results

show that APIT-PEG efficiently and continuously depleted L-arginine and L-lysine levels in

treated animals and significantly inhibited tumor growth in this model.

Material and Methods

Recombinant expression of APIT

Competent *E. coli* HMS174 were transformed with the pET26b-based plasmid with inserted

APIT encoding gene between *Nde*I and *Eco*RI restriction sites. Protein sequence of APIT is

supplied in the SI. Expression was controlled by T7 promoter and induced by 1 mM IPTG

(Sigma Aldrich, Steinheim, Germany) at an OD_{600} of 0.8-1.0. After 4 h cells were harvested by

centrifugation at 5000 g, 4°C. Expression resulted in inclusion bodies. Pelleted cells were

resuspended in 50 mM Tris pH 8.0. Inclusion bodies were isolated in several sonification steps

using a Bandelin Sonoplus ultrasonic device (Berlin, Germany).

Refolding of APIT

0.5 g of purified inclusion bodies were solubilized in 100 mL 6 M Urea, 50 mM Tris, 200 mM

NaCl, 100 mM DTT at pH 8.0 overnight at 4°C. Solubilized inclusion bodies were centrifuged

at 5000 g, 4°C for 30 min and filtrated using a 0.2 μm PS-filter. The protein was diluted in

refolding buffer 1 M Tris, 1 M NaCl, 1 mM EDTA, 25.6 μM FAD, 20 μM Cysteine, 0.5%

PEG6000 pH 8.6 and stirred at 4°C for 7 days.

Purification of APIT

The subsequent purification of refolded APIT was performed in two steps using an anion exchange chromatography (HiTrap™ Q FF) and a size exclusion chromatography (HiLoad™ 16/600 Superdex™ 200 pg) from Cytiva (Uppsala, Sweden).

Elution was performed with buffer A (50 mM Tris, pH 8.0) and buffer B (50 mM Tris, 1 M NaCl, pH 8.0), starting with a washing step of 25% B in 10 CV, followed by an elution gradient of 25-60 in 10 CV. APIT was then loaded onto a HiLoad™ 16/600 Superdex 200 pg. Column was then run using PBS pH 7.4 at flowrate of 1 mL/min.

PEGylation and purification of APIT-PEG5kDa

APIT was dialyzed in PBS pH 7.4 overnight at 4°C and concentration was adjusted to 1.1 mg/mL. 20 eq of PEG-NHS 5kDa (Celares GmbH, Berlin, Germany) was dissolved at a concentration of 408 mg/mL in PBS pH 7.4 and 1:10 (v:v) added to APIT. The reaction was performed overnight at 4°C continuously shaking. Reaction was purified using an anion exchange chromatography (HiTrap™ Q HP, Cytiva, Uppsala, Sweden) with buffer A (50 mM Tris, pH 8.0) and buffer B (50 mM Tris, 1 M NaCl, pH 8.0). The reaction was diluted 1:3 in buffer A, a maximum of 10 mg of protein was loaded onto 1 mL of resin at a flowrate of 0.5 mL/min. Afterwards a washing step at 0% B for at least 10 CV and elution gradient from 0-100% B was performed. Elution peak was pooled and dialyzed against PBS pH 7.4, concentration was adjusted to 2 mg/mL PEGylated APIT was filtered using a 0.2 μm PS filter and stored light protected.

Enzyme activity

The enzyme activity in the presence of the substate L-lysine was determined by measuring the production of hydrogen peroxide. H_2O_2 is reduced stoichiometrically by the chromogenic ABTS in a coupled HRP-catalyzed reaction as previously described by Butzke et al. (27). The enzyme activity is expressed in units [U], 1 U is equivalent to the production of 1 μmol H_2O_2 per minute.

HPLC Analysis

Purity of produced monomeric APIT and PEGylated APIT was assessed by HPLC analysis using an Agilent Technologies 1100 series system (Waldbronn, Germany). A ZORBAX

109 300SB-CN, 300 A, 5 µm, 4.6 x 250 nm was used. For elution was performed with Water in

110 HPLC grade + 0.1% TFA and increasing concentrations of Acetonitrile HPLC grade + 0.1 TFA.

Degree of PEGylation

The degree of PEGylation was assessed by determination of the mass increase measured by

MALDI mass spectra with an ultrafleXtreme mass spectrometer (Bruker Daltonics, Bremen,

Germany) equipped with a 355 nm smartbeam-II™ laser. 20-50 µg of protein were desalted

using Sep-Pak Vac C18 cartridges. The recommended manufacturers instruction was followed.

Bound protein was eluted using 80% acetonitrile in ddH$_2$O+ 0.1% TFA and subsequently

lyophilized at -100°C, 0.001 mbar. For external calibration Bruker Protein II standard (Bruker

Daltonics #8207234) was mixed with sinapinic acid solution (saturated in a 1:2 mixture (v/v)

of acetonitrile and 0.1% TFA). Drops were measured on a stainless-steel target (MTP 384 target

plate ground steel, Bruker Daltonics #8280784). The proteins were analyzed after double layer

preparation. First a thin matrix layer of SA solution (saturated in ethanol) was applied onto the

stainless-steel target, second the sample solution (4 mg/mL in a 1:1 mixture of acetonitrile and

ddH2O) was mixed with sinapinic acid solution (saturated in a 1:2 mixture (v/v) of acetonitrile

and 0.1% TFA) and dropped onto the first layer. Several thousand shots were accumulated for

the spectrum.

Pharmacodynamics of APIT-PEG on plasma L-arginine and L-lysine levels

127 Depletion of arginine and lysine in blood serum was assessed after a single intravenous

128 administration of 250 U/kg of APIT, 1000 U/kg, 250 U/kg APIT-PEG or 250 mL/kg PBS as

129 control vehicle into 5 tumor-free mice each group. Blood was collected 1 h prior to injection

130 and 0.25 h, 1 h, 3 h, 6 h, 24 h and 48 h after injection from each mouse. Samples of each

131 timepoint were pooled.

Dose-finding experiments and efficacy experiments of APIT-PEG in HNXF 536 PDX mice model

The maximum tolerated dose (MTD) of APIT-PEG was evaluated at three dose levels of 15

U/kg, 30 U/kg and 50 U/kg in 4 tumor-free mice per group. APIT-PEG was intravenously

injected every other day for one week and the tolerability was assessed by observing clinical

symptoms and body weight development.

Tumor fragments of HNXF 536 of a serial passage in nude mice were placed in PBS containing

10% penicillin/streptomycin and cut into fragments of 3-4 mm diameter. Prior to tumor

implantation, NMRI nu/nu (Crl:NMRI-*Foxn1^{nu}*) mice were anesthetized by inhalation of isoflurane. Animals were monitored until the tumor implants reached an appropriate size of 50-250 mm^3.

Mice bearing the HNXF 536 were randomly divided into control and treatment groups. The day of randomization was designated as Day 0 of the experiment. Mice were treated with 1) control vehicle 5 ml/kg/day every other day 2) APIT-PEG 15 U/kg/day every other day 3) APIT-PEG 40 U/kg/day every other day 4) doxorubicin-HCl 6 mg/kg every 7 days (See Tab. SI 1). Injection was performed via the lateral tail vein. Intraperitoneal injections were carried out if an intravenous injection was not possible due to an inflammation at the injection site. To control arginine and lysine depletion during the therapy, blood was collected 6 h, 24 h and 48 h after first treatment.

Analysis of amino acids in serum

For analysis of arginine and lysine levels in serum, blood was collected for by retrobulbar sinus puncture under isoflurane anesthesia. Plasma was stored in lithium heparin vials on ice. Vials were centrifuged at 2,000 x g for 5 min at 4°C. Separated plasma was transferred to a cooled tube on ice. 60 µL sample was diluted with 15 µL precipitation buffer and stored at 5°C for 60 min to precipitate proteins present. After centrifugation at 2 000 x g for 5 min at 4°C, the supernatants were diluted 1:1 with sample dilution buffer containing 200 nmol/ml Norleucin, which was used as internal standard. For quantitative and qualitative analysis of the amino acids serin, alanine, lysine and arginine at 570 nm the ARACUS Amino Acid Analyzer was used (MembraPure, Hennigsdorf, Germany).

H&E staining of dissected tumors

Tumors were dissected at day 42 after randomization. Tumor blocks were cut into 5 µm sections using a Leica RM2135 microtome (Leica Biosystems, Wetzlar, Germany). Slides were dried overnight at 40°C. Prior to staining slides were heated at 60°C for 30 min. Paraffin was removed by using xylene and rehydrated in ethanol solutions and washed with distilled water. Sections were then stained with hematoxylin for 8 min and washed in with water for 15 min. Prior to the eosin staining for 2 min, the slide was dipped in 96% ethanol. The slide was washed with water and dehydrated in 100% ethanol and xylene and mounted using the Merck Neo-Mount solution.

Statistics

170 Relative tumor growth over time was compared to Day 0. The relative body weight changes

171 and relative tumor growth are depicted as the group median and 5-95% confidence interval.

172 Statistical analysis was performed on the relative tumor growth comparing the treatments

173 separately to the control using the unpaired, nonparametric Mann-Whitney-U test. A p value

174 less than 0.05 was considered significant (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

Results

Expression and purification of monomeric APIT

We have previously demonstrated that the arginine and lysine deaminase APIT, an enzyme

found in the *Aplysia punctata* ink toxin, induces cell death in cancer cells, but not in normal

cells. To develop APIT for potential future pharmaceutical application, the serum half-life of

APIT had to be increased through PEGylation. We therefore developed the recombinant

expression and subsequent purification of APIT, which resulted in approximately 8 mg/L

monomeric APIT after the last purification step via size-exclusion chromatography. The most

crucial part of the purification was the purity of the inclusion bodies prior to refolding, which

was optimized by a serial of sonification steps. Refolded APIT was purified via anion exchange

chromatography, which still contained high molecular APIT (Fig. 1A). To obtain pure

monomeric APIT, a reductive step using tris(2-carboxyethyl) phosphine (TCEP) was

introduced prior to size exclusion chromatography (SEC) (Fig. 1B). Over time, monomeric

APIT tended to form dimers, which did not interfere with the enzymatic activity. To avoid

dimerization, monomeric APIT was stored at -80°C with the addition of small amounts of

TCEP. Monomeric APIT had a mass of approximately 59592.4 g/mol and a purity of ≥98%

measured by HPLC (Fig. 1 C), with a small amount of dimeric APIT with a mass of

121457.8 g/mol (Fig. 1D) and a batch dependent enzyme activity of 80-100 U/mg

Figure 1: Purification and characterization of recombinant APIT A) Chromatogram of anion
exchange after refolding of APIT B) Size-exclusion chromatogram of pooled and reduced APIT
C) HPLC analysis of monomeric recombinant APIT after purification on a normal phase
column D) MALDI-Mass of purified APIT showing small amounts of remaining dimeric APIT.

Production of APIT-PEG for *in vivo* application

To enhance the serum half-life, we PEGylated purified APIT at lysine residues and purified the
conjugates via anion-exchange chromatography (Fig. 2A). After conjugation, a protein
recovery of 60% of pure APIT-PEG was achieved. Compared to APIT the enzyme activity of
APIT-PEG was reduced by 20%. Purity of APIT-PEG was ≥95% measured by HPLC and SDS-
PAGE (Fig. 2B-C). APIT-PEG showed an average increase of mass by 55 507 g/mol indicating
a degree of PEGylation of approximately 8-10 mol PEG/ mol measured by MALDI-MS (Fig.
2D). APIT-PEG was formulated in PBS pH 7.2 and stored at 4°C. Prior to release for animal
experiments, the endotoxin level was determined and shown to have a concentration of 9
EU/mg, 0.14 EU/U. Masking of endotoxin by formulated APIT-PEG was controlled by the
spike-test. Spike recovery was 103-110%, concluding formulated APIT-PEG is not interacting
with endotoxins and therefore hiding endotoxins.

Figure 2: Purification of PEGylated APIT A) Chromatogram of Sepharose Q HP column B) 5-
15% Tris-SDS-PAGE of non-reducing APIT (1) and APIT-PEG 2 µg (2) 4 µg (3) stained with
Coomassie blue G250 C) CN-HPLC profile of PEGylated APIT after purification D) MALDI-
mass spectrum of PEGylated APIT shows an increase of mass by 55507 g/mol, indicating a
degree of PEGylation of 8-10 PEG per APIT.

PEGylated APIT depletes serum arginine and lysine for at least 48 h

Depletion of arginine and lysine in mice blood was tested after a single injection of A) 250 U/kg APIT B) 250 U/kg APIT-PEG or C) 25 mL/kg PBS as control vehicle for dilution effects. Serum of mice treated with APIT showed a beginning recovery of arginine (Fig. 3 A) and lysine levels after 6 h (Fig. 3 B). A single injection of 1000 U/kg APIT showed the depletion of arginine and lysin for up to 6 h. Nevertheless, after 24 h a complete recovery of amino acids in serum were observable (Fig. SI 1). Only the injection of APIT-PEG showed a complete depletion of serum arginine and lysine during the 48 h period (Fig. 3).

Figure 3: Pharmacodynamics of a single administration of 250 U/kg APIT or 250 U/kg APIT-PEG on A) arginine concentration in Serum and B) lysine concentration in serum showing beginning recovery of arginine and lysine concentration after 6 h, whereas APIT-PEG showed a complete depletion of serum arginine and lysine at least 48 h.

Tolerability experiments in tumor-free nu/nu mice

Dose finding tolerability tests were performed to define the suitable dose that does not affect the animal's health during the continuous treatment with APIT-PEG. Therefore, the relative change in body weight was taken as an indicator of tolerability and health since weight loss or gain should not exceed 10% in seven days during chemotherapy. Mice were treated with 15 U/kg, 30 U/kg or 50 U/kg of APIT-PEG every other day. The median relative body weight change of mice administered with 50 U/kg APIT-PEG showed a body weight loss of 7.3% after one week, which was declared as the upper limit (Fig. 4). The results of these experiments demonstrated that doses up to 50 U/kg were well tolerated.

240 Figure 4: The median and 5-95% confidence interval (CI) of relative body weight change of
241 mice treated with APIT-PEG every other day for one week shows a stronger impact of higher
242 doses of APIT-PEG on the well-being of mice.

Efficacy experiment of APIT-PEG in the HNXF 536 PDX mouse model

To evaluate the anti-tumor effect of arginine and lysine deprivation, we tested APIT-PEG in
mice xenografted with HNXF 536, a patient-derived head and neck cancer (SI Fig. 2). Based
on the tolerability experiment APIT-PEG was applied intravenously at 15 U/kg and 40 U/kg
every other day in this efficacy experiment in mice. As standard-of-care drug doxorubicin was
administered once a week at a concentration of 6 mg/kg and the control group received PBS
every other day (for details see Tab. SI 1). At day 21 the treatment was terminated due to the
high tumor load in the control group and lower dose group of 15 U/kg APIT-PEG. Tumor
growth was monitored until day 28 and last observations were carried forward. Mice were
weighed until day 42, relative body weight of doxorubicin showed a loss of weight during the
treatment, the well-known impact of standard chemotherapeutic treatment approaches on health
(Fig. 5A). Body weight of mice was slowly regained after the end of the treatment. APIT-PEG
treatment did not result in severe changes of body weight and did not affect the mice's health,
when compared to the control group. After the end of the treatment all mice gained weight
comparable to the control group. The tumor volume was determined from all mice as a measure
for tumor growth (Fig. 5B). Treatment with doxorubicin resulted in significant loss of tumor
volume starting at day 10. Treatment with 15 U/kg APIT-PEG did not show significant
difference in relative tumor growth compared to the control group (Fig. 5B-D). Serum analysis

of amino acids showed an increase of serum arginine after 48 h, probably leading to the lag of effectiveness due to incomplete arginine and lysine depletion (SI Fig. 3, 4). In contrast, treatment with 40 U/kg APIT-PEG successfully depleted arginine and lysine in blood serum during the treatment and showed a significant reduced tumor growth on the last day of injection ($p= 0.0160$) and one week after the end of treatment ($p= 0.0075$) compared to the non-treated control group (Fig. 5C-D).

Figure 5: Efficacy experiment of treatment of patient derived HNXF 536 xenografted mice with APIT-PEG or Doxorubicin the end of treatments is indicated with dashes lines A) Median and 5-95% CI of relative body weight changes for monitoring the mice's health B) Median and 5-95% CI of relative tumor growth during and one week after the treatment C) Boxplot representation of median and 5-95% confidence interval of relative tumor growth on day 21 (last day of treatment) D) and one week after the end of treatment on day 28. (Whiskers indicate 5-95% percentile values, the line in the middle of the box represents the median; Significance was tested with the Mann-Whitney U test, *$p < 0.05$, **$p < 0.01$, *** $p < 0.001$).

Discussion and Outlook

Targeting amino acid metabolism is an attractive anticancer strategy since it may provide alternatives for treatment of therapy resistant tumors (for review see (19)). Killing of therapy resistant tumor cell lines has also been demonstrated for APIT, which kills cancer cells independent of the induction of apoptosis (27), the mechanism induced by many of the genotoxic agents used for cancer therapy.(1) Resistances of tumors to these canonical therapies occur frequently by mutations in the general apoptosis signaling pathways.(2) This explains why anti-tumor therapies provoke the emergence of broad resistance to drugs with the same mode of action, the induction of apoptotic cell death. In addition, investigations in amino acid depletion cancer therapies suggest that side effects like those elicited by current treatments with genotoxic agents are less severe or even not observed, since amino acid auxotrophies are often highly tumor specific.(3) A prime example of amino acid depletion cancer therapy is the treatment of leukemias asparaginase curing more than 90% of pediatric acute lymphoblastic leukemia patients.(4)

Since APIT depletes the essential amino acid lysine and the semi-essential amino acid arginine (26), both activities interfere with tumor cell amino acid metabolism. Besides increased need of proliferating cells for lysine, the enzymes involved in *de novo* synthesis of arginine, argininosuccinate synthetase (ASS1) and argininosuccinate lyase (ASL) are frequently deregulated in cancer cells (5, 6), leading to arginine auxotrophies in many different tumor cells (for review see (9)) This metabolic Achilles heel of tumor cells can be used for therapeutic approaches based on the activities of APIT. However, the blood circulation time of active purified APIT is insufficient to achieve the long-lasting depletion of L-lysine and L-arginine that is required to starve tumor cells.

Here we developed the PEGylation of purified APIT and thereby achieved a dramatic increase in the blood circulation time of the enzyme. A single dose of PEG-APIT depleted L-lysine and L-arginine for more than 50 hours whereas the amino acid oxidase activity of non-PEGylated APIT was cleared in less than 10 hours. For the current experiments, the purification yield was 8 mg monomeric APIT per liter culture with a batch-dependent enzyme activity between 80-100 U/mg. Further optimization of expression and large-scale production using fermenters will enable the production of PEG-APIT at a scale required for clinical testing. So far, size-exclusion chromatography step is necessary to yield pure monomeric APIT. Further optimization to exclude higher molecular APIT might be achieved by selection of suitable hydrophobic

interaction chromatography media to improve the purification procedure also for large scale approaches. Longer incubation times of APIT without reducing agents showed the formation of dimers, without changes in enzyme activity. Monomeric APIT can be stored using reducing agents like TCEP. Nevertheless, several other L-amino acids oxidases are reported to be dimeric, the dimerization of APIT over time should further be evaluated.

APIT was PEGylated on lysine residues using monofunctionalized NHS-activated polyethylene glycol. The high PEG concentration during the PEGylation resulted in minor precipitation of protein, explaining the protein recovery of 60% APIT-PEG. PEGylated APIT was successfully isolated by anion-exchange chromatography. The degree of PEGylation was further determined by MALDI-mass spectrometry and was between 8-10 mol PEG per mol APIT. The enzyme activity of APIT-PEG was about 20% reduced compared to APIT, indicating that PEGylation causes a minor loss of enzyme activity. Furthermore, a critical step for the application in animals is the endotoxin amount. Reduction of endotoxin might be achieved by endotoxin reduced workflow via anion exchange chromatography and could be integrated in the existing purification steps in larger scale productions.

For intravenous injection into mice a formulation of APIT-PEG in endotoxin-free PBS was used. In this formulation no endotoxin masking was observed. Mice were administered different concentration of APIT-PEG according to the previously tested tolerated doses. APIT-PEG treatment had no negative effect on the well-being of mice, whereas doxorubicin treatment resulted in weight loss. APIT-PEG showed dose-dependent, significant decrease of the head and neck cancer HNXF 536 in patient derived xenografted mice compared to the control group. Interestingly, treatment with the lower dose of 15 U/kg APIT-PEG caused a non-significant reduction of the tumor volume and this correlated with a low but measurable level of L-arginine and L-lysine in the plasma of these mice. Due to the very good tolerability of the treatment, the dose and thus the efficacy could be optimized in further experiments. Due to its broader action by depleting of two amino acids instead of only one, APIT-PEG could be an interesting alternative for ADI-PEG20, which was granted the orphan drug status and is currently used for the treatment of malignant pleural mesothelioma.

Acknowledgements

We thank Prof. Dr. Thomas Dandekar (Chair of Bioinformatics) for the help with the statistical analysis, and Prof. Dr. Andreas Rosenwald (Institute for Pathology) for the help with the analysis of histological sections.

References

1.   Herr I, Debatin KM. Cellular stress response and apoptosis in cancer therapy. Blood. 2001;98(9):2603-14.

2.   Igney FH, Krammer PH. Death and anti-death: tumour resistance to apoptosis. Nat Rev Cancer. 2002;2(4):277-88.

3.   Egler RA, Ahuja SP, Matloub Y. L-asparaginase in the treatment of patients with acute lymphoblastic leukemia. J Pharmacol Pharmacother. 2016;7(2):62-71.

4.   Maese L, Rau RE. Current Use of Asparaginase in Acute Lymphoblastic Leukemia/Lymphoblastic Lymphoma. Front Pediatr. 2022;10:902117.

5.   Cheng CT, Qi Y, Wang YC, Chi KK, Chung Y, Ouyang C, et al. Arginine starvation kills tumor cells through aspartate exhaustion and mitochondrial dysfunction. Commun Biol. 2018;1:178.

6.   Cancer Genome Atlas Research N, Weinstein JN, Collisson EA, Mills GB, Shaw KR, Ozenberger BA, et al. The Cancer Genome Atlas Pan-Cancer analysis project. Nat Genet. 2013;45(10):1113-20.

7.   Qiu F, Huang J, Sui M. Targeting arginine metabolism pathway to treat arginine-dependent cancers. Cancer Lett. 2015;364(1):1-7.

8.   Morris SM, Jr. Regulation of enzymes of the urea cycle and arginine metabolism. Annu Rev Nutr. 2002;22:87-105.

9.   Riess C, Shokraie F, Classen CF, Kreikemeyer B, Fiedler T, Junghanss C, et al. Arginine-Depleting Enzymes - An Increasingly Recognized Treatment Strategy for Therapy-Refractory Malignancies. Cell Physiol Biochem. 2018;51(2):854-70.

10.   Huang HY, Wu WR, Wang YH, Wang JW, Fang FM, Tsai JW, et al. ASS1 as a novel tumor suppressor gene in myxofibrosarcomas: aberrant loss via epigenetic DNA methylation confers aggressive phenotypes, negative prognostic impact, and therapeutic relevance. Clin Cancer Res. 2013;19(11):2861-72.

11.   Kobayashi E, Masuda M, Nakayama R, Ichikawa H, Satow R, Shitashige M, et al. Reduced argininosuccinate synthetase is a predictive biomarker for the development of pulmonary metastasis in patients with osteosarcoma. Mol Cancer Ther. 2010;9(3):535-44.

12.   Kremer JC, Van Tine BA. Therapeutic arginine starvation in ASS1-deficient cancers inhibits the Warburg effect. Mol Cell Oncol. 2017;4(3):e1295131.

13.   Liu Q, Stewart J, Wang H, Rashid A, Zhao J, Katz MH, et al. Reduced expression of argininosuccinate synthetase 1 has a negative prognostic impact in patients with pancreatic ductal adenocarcinoma. PLoS One. 2017;12(2):e0171985.

14.   Nicholson LJ, Smith PR, Hiller L, Szlosarek PW, Kimberley C, Sehouli J, et al. Epigenetic silencing of argininosuccinate synthetase confers resistance to platinum-induced cell death but collateral sensitivity to arginine auxotrophy in ovarian cancer. Int J Cancer. 2009;125(6):1454-63.

15.   Qiu F, Chen YR, Liu X, Chu CY, Shen LJ, Xu J, et al. Arginine starvation impairs mitochondrial respiratory function in ASS1-deficient breast cancer cells. Sci Signal. 2014;7(319):ra31.

16.   Rabinovich S, Adler L, Yizhak K, Sarver A, Silberman A, Agron S, et al. Diversion of aspartate in ASS1-deficient tumours fosters de novo pyrimidine synthesis. Nature. 2015;527(7578):379-83.

17.   Allen MD, Luong P, Hudson C, Leyton J, Delage B, Ghazaly E, et al. Prognostic and therapeutic impact of argininosuccinate synthetase 1 control in bladder cancer as monitored longitudinally by PET imaging. Cancer Res. 2014;74(3):896-907.

18.   Kim RH, Coates JM, Bowles TL, McNerney GP, Sutcliffe J, Jung JU, et al. Arginine deiminase as a novel therapy for prostate cancer induces autophagy and caspase-independent apoptosis. Cancer Res. 2009;69(2):700-8.

19.   Butler M, van der Meer LT, van Leeuwen FN. Amino Acid Depletion Therapies:
Starving Cancer Cells to Death. Trends Endocrinol Metab. 2021;32(6):367-81.
20.   Dillon BJ, Holtsberg FW, Ensor CM, Bomalaski JS, Clark MA. Biochemical
characterization of the arginine degrading enzymes arginase and arginine deiminase and their
effect on nitric oxide production. Med Sci Monit. 2002;8(7):BR248-53.
21.   Holtsberg FW, Ensor CM, Steiner MR, Bomalaski JS, Clark MA. Poly(ethylene
glycol) (PEG) conjugated arginine deiminase: effects of PEG formulations on its
pharmacological properties. J Control Release. 2002;80(1-3):259-71.
22.   Hall PE, Lewis R, Syed N, Shaffer R, Evanson J, Ellis S, et al. A Phase I Study of
Pegylated Arginine Deiminase (Pegargiminase), Cisplatin, and Pemetrexed in
Argininosuccinate Synthetase 1-Deficient Recurrent High-grade Glioma. Clin Cancer Res.
2019;25(9):2708-16.
23.   Harding JJ, Do RK, Dika IE, Hollywood E, Uhlitskykh K, Valentino E, et al. A phase
1 study of ADI-PEG 20 and modified FOLFOX6 in patients with advanced hepatocellular
carcinoma and other gastrointestinal malignancies. Cancer Chemother Pharmacol.
2018;82(3):429-40.
24.   Lowery MA, Yu KH, Kelsen DP, Harding JJ, Bomalaski JS, Glassman DC, et al. A
phase 1/1B trial of ADI-PEG 20 plus nab-paclitaxel and gemcitabine in patients with
advanced pancreatic adenocarcinoma. Cancer. 2017;123(23):4556-65.
25.   Yao S, Janku F, Subbiah V, Stewart J, Patel SP, Kaseb A, et al. Phase 1 trial of ADI-
PEG20 plus cisplatin in patients with pretreated metastatic melanoma or other advanced solid
malignancies. Br J Cancer. 2021;124(9):1533-9.
26.   Butzke D, Hurwitz R, Thiede B, Goedert S, Rudel T. Cloning and biochemical
characterization of APIT, a new l-amino acid oxidase from Aplysia punctata. Toxicon.
2005;46(5):479-89.
27.   Butzke D, Machuy N, Thiede B, Hurwitz R, Goedert S, Rudel T. Hydrogen peroxide
produced by Aplysia ink toxin kills tumor cells independent of apoptosis via peroxiredoxin I
sensitive pathways. Cell Death Differ. 2004;11(6):608-17.
28.   Butzke D. GS, Dittrich M., Rudel T., Meyer T.   , inventor; Max-Planck-Gesellschaft
zur Foerderung der Wissenschaften E.V., assignee. L-Amino acid oxidase with cytotoxic
activity from Aplysia Punctata 2008.

424