Metagenomic analysis of ecological niche overlap and community collapse in microbiome dynamics

Hiroaki Fujita[1*], Masayuki Ushio[1,2], Kenta Suzuki[3], Masato S. Abe[4], Masato Yamamichi[5,6], Yusuke Okazaki[7], Alberto Canarini[1], Ibuki Hayashi[1], Keitaro Fukushima[8], Shinji Fukuda[9-12], E. Toby Kiers[13], and Hirokazu Toju[1*]

[1]Center for Ecological Research, Kyoto University, Otsu, Shiga 520-2133, Japan

[2]Department of Ocean Science (OCES), The Hong Kong University of Science and Technology (HKUST), Clear Water Bay, Kowloon, Hong Kong SAR, China

[3]Integrated Bioresource Information Division, BioResource Research Center, RIKEN, Tsukuba, Ibaraki 305-0074, Japan

[4]Faculty of Culture and Information Science, Doshisha University, Kyotanabe, Kyoto 610-0321, Japan

[5]School of Biological Sciences, The University of Queensland, St. Lucia, Brisbane, QLD 4072, Australia

[6]Department of International Health and Medical Anthropology, Institute of Tropical Medicine, Nagasaki University, Nagasaki 852-8523, Japan

[7]Institute for Chemical Research, Kyoto University, Gokasho, Uji, Kyoto 611-0011, Japan

[8]Faculty of Food and Agricultural Sciences, Fukushima University, Kanayagawa 1, Fukushima, Fukushima 960-1296, Japan.

[9]Institute for Advanced Biosciences, Keio University, Tsuruoka, Yamagata 997-0052, Japan.

[10]Gut Environmental Design Group, Kanagawa Institute of Industrial Science and Technology, Kawasaki, Kanagawa 210-0821, Japan.

[11]Transborder Medical Research Center, University of Tsukuba, Tsukuba, Ibaraki 305-8575,

Japan.

[12]Laboratory for Regenerative Microbiology, Juntendo University Graduate School of

Medicine, Tokyo 113-8421, Japan.

[13]Department of Ecological Science, Vrije Universiteit Amsterdam, Amsterdam, the

Netherlands

Correspondence and requests for materials should be addressed to H.F. (email:

fujita.h@ecology.kyoto-u.ac.jp) or H.T. (email: toju.hirokazu.4c@kyoto-u.ac.jp).

Abstract

Species utilizing the same resources ultimately do not coexist for long periods of time. Such competitive exclusion mechanisms potentially underly dynamics of microbiomes, causing breakdowns of communities constituted by species with similar genetic backgrounds of resource use. Nonetheless, it remains a major challenge to integrate genomics and ecology for understanding deterministic processes of species coexistence in species-rich communities. We here show that community-scale analyses of functional gene redundancy provide statistical platforms for interpreting and predicting collapse of bacterial communities. Through 110-day time-series of experimental microbiome dynamics, we analyzed the metagenome-assembled genomes of coexisting bacterial species. We then reconstructed ecological niche space based on the multivariate analysis of the genome compositions in order to evaluate potential shifts in the level of niche overlap between species through time. Specifically, we hypothesized that community-scale pressure of competitive exclusion could be evaluated by quantifying overlap of genetically determined resource-use profiles (metabolic pathway profiles) among coexisting species. We found that the degree of community compositional changes observed in the experimental microbiome was explained by the magnitude of metabolic pathway (gene repertoire) overlaps among bacterial species. The metagenome-based analysis of genetic potential for competitive exclusion will help us forecast major events in microbiome dynamics such as sudden community collapse (i.e., dysbiosis).

Classic niche theory predicts that coexistence of species requires interspecific difference in resource ranges[1–6]. Although some specific mechanisms can guarantee coexistence even in the presence of niche overlap (e.g., spatial structure of habitats and temporal variability in resource availability), similarity/dissimilarity in basic resource dependency among species is the key factor determining the occurrence of competitive exclusion[7–9]. Therefore, evaluating the overlap of "fundamental niches", which are defined by species' fundamental resource requirements and resource-use capabilities[10,11], is an essential step for understanding and predicting community-level dynamics.

Insights into fundamental niches are encrypted in species' genomes[12–14]. In other words, as species' traits are encoded in their DNA, reconstructed genomes provide the ultimate basis for evaluating target species' fundamental niches[15,16]. Thus, potential strength of competitive

interactions within ecological guilds or communities could be evaluated based on the

distribution of species' gene repertoires within ecological niche space reconstructed with

metagenomic data[12,15,16] ("metagenomic niche space"). Although overlap of niches does not

always cause competitive exclusion[7–9], higher levels of gene repertoire overlap within a

community may impose greater impacts on population dynamics of constituent species.

It is essential to examine whether such competition-driven population-level

phenomena underly drastic ecological events observed at the community level. Microbial

communities sometimes show sudden and substantial changes in species and/or taxonomic

compositions[17–19]. Human gut microbiomes, for example, have been reported to show drastic

shifts from species-rich states to "imbalanced" states with low α-diversity and

overrepresentation of pathogenic species[20–23] (e.g., *Clostridium difficile*). Elucidating the

ecological mechanisms by which such drastic community-level events are caused provide

fundamental insights into microbiome dynamics[23–25]. In this respect, an important challenge is

to test the hypothesis that higher levels of gene-repertoire overlap are observable prior to

drastic community compositional changes than after such changes. However, this hypothesis,

to our knowledge, has not yet been tested presumably due to the paucity of time-series

observations of microbiomes with substantial compositional changes. Even if such

microbiome time-series data are available, analyses of potential niche (gene repertoire)

overlap require another line of information, specifically, data of respective species' genomes

at multiple time points. Therefore, developing research systems that overcome these current

constrains is expected to deepen our understanding of microbiome ecological processes.

We here show how degree of gene-repertoire overlap changes through dynamics of

species-rich microbial communities. By targeting an experimental microbial system showing

rapid and substantial changes in taxonomic compositions[19], we reconstruct niche space

depicting species' gene repertoires. Based on a whole-genome shotgun metagenomic analysis

at 13 time points within the 110-day time-series of the microbiome experiment, we reveal

temporal shifts in the magnitude of gene repertoire overlap among microbial species. We then

examine whether a high level of fundamental-niche overlap is observed prior to drastic

changes in community structure. Overall, we explore how signs of drastic shifts in community

structure are detected by reconstructing community-scale degree of fundamental niche

overlap with the aid of genomic information.

Results

Target microbiome. We focused on the experimental microbiome showing drastic shifts in taxonomic compositions[19]. In a previous study[19], a 110-day monitoring of microbiomes was performed with six experimental settings. To set up experimental microbiomes with high diversity of bacterial species/taxa, natural microbial communities derived from soil or pond-water ecosystems, rather than "synthetic" communities with pre-defined diversity, were used as source inocula. Specifically, microbiomes were set up with combinations of two source inoculum types (soil- or pond-water-derived inoculum microbiomes) and three medium types (oatmeal, oatmeal-peptone, or peptone broth medium) with eight replications ($2 \times 3 \times 8 = 48$ microbiomes; see Methods for details). From each of the 48 microbiomes, a fraction of each replicate community was sampled every 24 hours. The collected samples were subjected to the amplicon sequencing of the 16S rRNA region and the temporal changes in community compositions were monitored throughout the time-series[19]. By calculating the magnitude of time-series changes in community compositions[19] ("abruptness" index; Fig. 1a; Supplementary Fig. 1), we focused on a water-inoculum/oatmeal-medium replicate community showing the most abrupt (rapid and substantial) changes in community compositions among the 48 microbiomes examined (Fig. 1a; Supplementary Fig. 1).

Functional dynamics of microbiomes. By targeting the replicate community mentioned above, we performed whole-genome shotgun sequencing at 13 time points across the time-series. In total, 32 high-quality (> 80 % completeness and < 5 % contamination) metagenome-assembled genomes (MAGs) belonging to 20 genera (16 families; 12 orders) were detected (Figs. 1b-c and 2; Supplementary Fig. 2; Supplementary Table 1). As indicated in the amplicon sequencing analysis[19] (Fig. 1a), drastic shifts from taxon-rich community states to oligopolistic states was observed around Day 20 in the shotgun sequencing analysis (Fig. 1b).

After the drastic community compositional change, the system reached a quasi-stable state represented by the dominance of a *Hydrotalea* (Chitinophagaceae) bacterium (Fig. 1b). The MAG of the *Hydrotalea* was characterized by relatively low GC content (38 %) and relatively small genome size within the community (ca. 3.1 Mb; Fig. 2a). In contrast, the two bacterial MAGs consistently coexisted with the dominant *Hydrotalea* through the time-series (i.e., *Terracidiphilus* and *Mangrovibacter*) had larger genome size (4.2 and 5.4 Mb, respectively; Fig. 1c), characterized by various genes absent from the *Hydrotalea* genome (Fig. 2; Supplementary Fig. 3). Specifically, the *Terracidiphilus* MAG showed metabolic

pathways/processes for degrading plant-derived biopolymers (e.g., cellulose; Fig. 2),

potentially surviving as a primary user of polymer compounds within the plant-derived

(oatmeal) medium. Meanwhile, the *Mangrovibacter* MAG had pathways/processes related to

starch degradation (e.g., amylase) and vitamin-B_{12} transportation, which were absent from the

genomes of *Hydrotalea, Terracidiphilus*, and the other MAG (*Rhizomicrobium*) detected on

Day 40-60 (Fig. 2).

Multivariate analysis of gene repertoires. The whole-genome shotgun metagenomic data

were used to evaluate how the level of gene repertoire overlap among microbes shifted

through time. We anticipated that microbial species with similar resource-use abilities or

restrictions have similar genomic structure. Therefore, it is expected that species competing

for the same resource tend to form clusters within the space defined based on the principal

coordinate analysis (PCoA) of dissimilarity in gene repertoires. For each pair of the 32

MAGs, dissimilarity (Jaccard β-diversity) of gene repertoires was calculated based on the

matrix representing the presence/absence of the 6,999 genes annotated with the program

Prokka[26]. A PCoA was then performed using the β-diversity information (Fig. 3a). At each of

the 13 time point, detected MAGs were plotted on the PCoA space. Since we did not have *a*

priori knowledge of specific metabolic pathways keys to the microbe-to-microbe competition

within the experimental microbiome, the entire datasets were used in this multivariate

analysis. Given general characteristics of multivariate analysis based on β-diversity metrics,

the multivariate reconstruction of ecological niche space depends greatly on the genes whose

presence/absence profiles vary among species, while housekeeping genes possessed by most

species are expected to contribute little to the multivariate analysis.

We then found that alphaproteobacterial and gammaproteobacterial MAGs respectively

constituted some clusters within the niche space reconstructed based on the multivariate

analysis early in the microbiome dynamics (Day 1-20; Fig. 3b). This state with high niche

overlap and potential within-guild competition for resources then collapsed into a simpler

community state represented by *Hydrotalea*, *Mangrovibacter*, *Terracidiphilus*, *and*

Rhizomicrobium as detailed above (Fig. 3b). The space once occupied by many

alphaproteobacterial and gammaproteobacterial MAGs remained unoccupied or sparsely

occupied after the community compositional collapse. Even when the number of MAGs

detectable with our shotgun-metagenomic sequencing increased again late in the time-series,

aggregations of microbes with similar genomic compositions remained unobserved (Fig. 3b).

Metagenomic niche overlap. We next quantitatively evaluated dynamics in the magnitude of community-scale niche overlap within the multivariate space (Fig. 3). In our analysis, the niche overlap index was defined as:

$$1 - \frac{\sum_{i \in D, j \in D} \beta_{ij}}{N_D},$$

where D is the set of MAGs detected on a focal day (time point), β_{ij} is the Jaccard metric of dissimilarity in gene compositions, and N_D is the number of MAGs detected on the day. By definition, this niche overlap value varies from 0 (completely different repertoires of genes in all pairs of MAGs) and 1 (completely identical gene repertoires in all pairs of MAGs), allowing us to evaluate niche overlap levels of target communities within the standardized ranges. The results indicated that the level of niche overlap was the highest on Day 1 and that it gradually decreased until Day 20 (Fig. 4a-b). A slight increase in the niche overlap index was observed on Day 24, but it dropped sharply by Day 30 (Fig. 4a-b). Although the niche overlap score remained low between Day 40 and 60, it increased again on Day 70 (Fig. 4b).

We then found that the estimated niche overlap level significantly explained the magnitude of the observed community compositional changes ($t = 5.525$, df = 10, $P = 0.00025$; Fig. 5). In other words, higher levels of gene-repertoire overlap within a community were followed by larger shifts in community compositions at subsequent time points.

Discussion

We here showed that among-species overlap of gene repertoires are observable prior to drastic changes in community structure. Early in the experimental microbiome dynamics, alphaproteobacterial and gammaproteobacterial species were present, resulting in relatively high niche-overlap scores at the community level (Figs. 3 and 4). The quasi-equilibrium state then collapsed into another quasi-equilibrium represented by a small number of bacteria varying in genome size and metabolic capabilities. Throughout the time-series, higher niche overlap levels entailed greater changes in microbial community compositions (Fig. 5). These findings lead to the working hypothesis that collapse of microbiome structure is predicted by the level of potential niche overlap within multivariate metagenomic space. In light of the "limiting similarity" rule of ecological niches[27], pairs of microbial species that exceed a

critical limit of genome compositional similarity are expected to compete for the same

resources, eventually driving competitive exclusion processes. Thus, as examined in this

study, similarity/dissimilarity in genetically determined resource-use properties (i.e.,

fundamental niches) sets baselines for consequences of interspecific interactions.

The results also indicated that niche overlap level does not necessarily show monotonic

decrease through microbial community processes. Although gene-repertoire overlap level and

detectable species richness sharply declined early in the microbiome dynamics, both variables

gradually increased again around Day 80 (Figs. 1a and 4b). In the resurgence process,

however, the dense clusters of alphaproteobacterial or gammaproteobacterial species detected

until Day 20 did not appear again within the niche space (Fig. 3b). These observations suggest

that once collapsed, microbial communities may not return to previous states with highest

levels of niche overlap, but refilling of poorly-used niches can occur under the constraint of

limiting similarity within niche space.

The simple framework for evaluating overlap of fundamental niches is applicable to

diverse types of microbiomes. Given that our β-diversity-based index is standardized within

the range from 0 to 1, next crucial step is to examine how threshold niche overlap values for

anticipating microbial community collapse vary among different types of ecosystems. Such

threshold values can vary among ecosystems depending on their basic levels of sustainable

functional redundancy. In our laboratory microbiome, for example, the lack of environmental

fluctuations (e.g., temperature fluctuations) and spatial structure (e.g., refuges for inferior

species) might severely limited coexistence of functionally similar species (species with

similar metabolic capabilities). In contrast, in human gut microbiomes, spatially

complexity[28,29] and fluctuating environmental conditions[21] may reduce the risk of competitive

exclusion, allowing higher levels of niche overlap within communities. Thus, extension of

time-series metagenomic analyses to diverse types of ecosystems[30–33] will enhance our

knowledge of relationship among ecosystem properties, functional redundancy, and

community stability.

In this study, we included whole metagenomic datasets of the examined microbes due to

the lack of *a priori* insights into the metabolic pathways/processes playing essential roles in

interspecific competition for resources. In this respect, our analysis is a preliminary

conceptual step for evaluating potential overlap of fundamental niches at the community

level. In future studies, analyses excluding housekeeping genes[34,35] or those focusing on

specific functional groups of genes (e.g., carbohydrate degrading genes[36]) may provide more

reliable inference of niche overlap. Because such selection of genes can critically influence

threshold niche-overlap values for anticipating abrupt community compositional changes,

setting a commonly applicable criterion of choosing target gene sets will help us perform

comparative analyses across a wide range of microbial communities.

While genomic information provides an ultimate platform for inferring fundamental

niches[12–14], overlap of gene repertoires may not always result in competitive exclusion of

species within communities. Even in a pair of species with similar gene repertoires,

differentiation in gene expression patterns may occur to avoid overlap of resource-use

patterns between species, allowing coexistence of the two species in an environment. Such

differentiation of "realized niches[10]" through phenotypic plasticity is potentially evaluated by

transcriptomic or metabolomic analyses[37,38]. Consequently, integration of (meta)transcriptome

and (meta)metabolome analyses[39–41] with metagenome-based analyses will reorganize our

understanding of deterministic processes in microbiome dynamics.

Methods

Time-series data of experimental microbiomes. We used the experimental system of the microbiome time-series monitoring described in a previous study[19]. In the experiment, microbiomes differing in the magnitude of community compositional shifts were constructed across the six treatments defined by the combinations of two inoculum sources and three types of media. One of the source microbiomes derived from the soil collected from the A layer (0-10 cm in depth) in the research forest of Center for Ecological Research, Kyoto University, Kyoto, Japan (34.972 ºN; 135.958 ºE). The other source inoculum was prepared by collecting water from a pond ("Shoubuike") near Center for Ecological Research (34.974 ºN, 135.966 ºE). Each of the source inocula was introduced into oatmeal (Medium-A), oatmeal-peptone (Medium-B), or peptone (Medium-C) broth media with eight replicates. Thus, in total, 48 experimental microcosms (two source microbiomes × three media × eight replicates) were constructed in a deep-well plate (1000-µL-scale culture in each well). The plate was kept shaken at 1,000 rpm at 23 ºC. After five-day pre-incubation, 200 µL out of the 1,000-µL culture medium was sampled from each well every 24 hours for 110 days. In each sampling event, 200 µL of fresh medium was added to each well so that the total culture volume was kept constant. In total, 5,280 samples (48 communities/day × 110 days) were collected through the time-series experiment. After DNA extraction, the samples were subjected to the amplicon sequencing analysis of the 16S rRNA region[19]. To quantify the speed and magnitude of community shifts through time, the "abruptness" index was calculated through the time-series of each replicate microcosm in each experimental treatment[19]. Specifically, an estimate of the abruptness index for time point t was obtained as the Bray-Curtis β-diversity between average community compositions from time points $t - 4$ to t and those from $t + 1$ to $t + 5$ (i.e., dissimilarity between 5-day time-windows). The Bray-Curtis β-diversity[42] was calculated as $\frac{\sum_{i=1}^{n} |X_{ij} - X_{ik}|}{\sum_{i=1}^{n} (X_{ij} + X_{ik})}$, where X_{ij} and X_{ik} denoted relative abundance of microbial amplicon sequence variant (ASV) i in the compared time windows (j, from $t - 4$ to t; k, from $t + 1$ to $t + 5$). An abruptness score larger than 0.5 indicates that turnover of more than 50 % of community compositions occurred between the time-windows[19].

Whole-genome shotgun metagenomics. Focusing on a replicate microcosm in which the most rapid and substantial turnover of community compositions was observed (replicate no. 5

of Water/Medium-A treatment; Supplementary Fig. 1), whole-genome shotgun metagenomics

was conducted by targeting 13 samples (Day 1, 10, 20, 24, 30, 40, 50, 60, 70, 80, 90, 100,

110). Each DNA sample was processed with Nextera XT DNA Library Preparation Kit

(Illumina) and sequenced with the DNBSEQ-G400 (BGI; 200-bp paired-end sequencing).

From the output data, sequencing adaptors were removed using Cutadapt[43] 2.5 and quality

filtering was performed with Fastp[44] 0.21.0: ca. 10 Gb/sample was subjected to the analysis

[in total, 159.96 Gb (1000.301 M reads)].

The sequences of each sample were assembled with metaSPAdes[45] 3.15.2. Binning was

then performed with MetaWRAP[46] 1.3.2, followed by quality assessing with CheckM[47] 1.1.3.

Only the MAGs with > 80 % completeness and < 5 % contamination were used in the

downstream analyses. The identity between MAGs were calculated using FastANI[48] 1.33 and

MAGs with ≥ 98 % identity were dereplicated through the time-series (Supplementary Table

1). Read-coverage was then calculated with CoverM[49] 0.6.0, followed by taxonomic

annotation was performed using GTDB-Tk[50,51] 1.6. Gene annotation was performed with

Prokka[26] 1.14.6, yielding 6,999 annotated genes (Supplementary Data 1). To conduct

additional functional annotation of genes, the orthology numbers of Kyoto Encyclopedia of

Genomes (KEGG) were retrieved using GhostKOALA[52] 2.2. For respective microbial MAGs

(bins), completeness of metabolic pathways was estimated with KEGG decoder[53] 1.3. Based

on the matrix representing KEGG metabolic pathway/process profiles of respective MAGs

(Supplementary Data 2), a heatmap showing pathway/process completeness was drawn

(Supplementary Fig. 3).

Background environmental conditions. For the 13 samples subjected to the shotgun

metagenomic analysis, concentrations of ammonium (NH_4^+) and nitrate (NO_3^-) were

measured to obtain supplementary information of background environmental conditions.

Colorimetric methods with a modified indophenol reaction[54,55] and the VCl3/Griess assay

were applied for the measurements of NH_4^+ and NO_3^-, respectively. Samples were run in

triplicates via a standard addition method to account for individual matrix effects[56].

Multivariate analysis of the metagenomic space. Based on the whole matrix representing

the profiles of the 6,999 genes (Supplementary Data 1), the Jaccard metric of β-diversity was

calculated for each pair of the 32 microbial MAGs (β_{ij}, where i and j represent MAGs). The

β-diversity estimates were then used to perform a principal coordinate analysis. Using the

obtained principal coordinate scores, all the microbial MAGs detected through the time-series

were plotted on a multivariate space consisting of the first three PCoA axes (PCoA 1, PCoA 2,

and PCoA 3). At each time point, the MAGs detected with the shotgun metagenomic

sequencing (defined as the MAGs whose relative abundance is greater than 0.1 %) was

plotted on the three-dimensional space defined with the PCoA axes.

Evaluation of niche overlap level. The community-scale magnitude of potential niche

overlap among species was evaluated based on the whole-genome shogun sequencing dataset.

Specifically, the niche overlap index was defined as:

317
$$1 - \frac{\sum_{i \in D, j \in D} \beta_{ij}}{N_D},$$

where D is the set of MAGs detected on a focal day (relative abundance > 0.1 %), β_{ij} is the

Jaccard metric of dissimilarity in gene compositions (defined in the previous section), and N_D

is the number of MAGs detected on the day.

The scores of the niche overlap index were shown on a two-dimensional space

representing metabolic pathway/process compositions of the whole community at respective

time points. At each time point, the gene repertoires of the detected MAGs (the MAGs whose

relative abundance is greater than 0.1 %) were summed, yielding a matrix representing 13

time points × 6,999 genes. The matrix was used to calculate dissimilarity (Jaccard β-diversity)

in microbiome-scale gene repertoires among time points to perform a PCoA analysis.

To test whether a high level of fundamental-niche overlap is observed prior to drastic

changes in microbial community structure, we examined relationship between the above niche

overlap index and time-series shifts in community structure (Bray-Curtis β-diversity between

present and next time points through the time-series of the shotgun metagenomic data).

Data availability

333 The 16S rRNA sequencing data reported in a previous study[19] are available from the DNA

334 Data Bank of Japan (DDBJ) with the accession number DRA013352, DRA013353,

335 DRA013354, DRA013355, DRA013356, DRA013368 and DRA013379. The whole-genome

shotgun metagenomics data are available with the DDBJ accession number DRA013382. The microbial community data are deposited at our GitHub repository (https://github.com/hiroakif93/MTS_nicheSpace) [to be publicly available after acceptance of the paper]. The matrices of the shotgun metagenomic data are available as Supplementary Data 1 and 2.

Code availability

All the scripts used to analyze the data are available at the GitHub repository (https://github.com/hiroakif93/MTS_nicheSpace) [to be publicly available after acceptance of the paper].

Acknowledgements. Computation time was provided by the SuperComputer System, Institute for Chemical Research, Kyoto University. This work was financially supported by JST PRESTO (JPMJPR16Q6), Human Frontier Science Program (RGP0029/2019), JSPS Grant-in-Aid for Scientific Research (20K20586), NEDO Moonshot Research and Development Program (JPNP18016), and JST FOREST (JPMJFR2048) to H.T., JSPS Grant-in-Aid for Scientific Research (20K06820 and 20H03010) to K.S., and JSPS Fellowship to H.F. and A.C..

Author Contributions. H.T. designed the work with H.F.. H.F. and A.C. performed experiments. H.F. analyzed the data with Y.O., and H.T.. H.F. and H.T. wrote the paper with all the authors.

Competing Interests. The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at [URL to be supplied by the publisher]

364 **Correspondence and requests for materials** should be addressed to fujita.h@ecology.kyoto-
365 u.ac.jp or toju.hirokazu.4c@kyoto-u.ac.jp.

References

1. Gause, G. F. *The Struggle for Coexistence*. (Williams & Wilkins, 1934).

2. Hardin, G. The competitive exclusion principle. *Science (1979)* **131**, 1292–1297
(1960).

3. Volterra, V. Variations and fluctuations of the number of individuals in animal species
living together. *ICES Journal of Marine Science* **3**, 3–51 (1928).

4. Zaret, T. M. & Rand, A. S. Competition in tropical stream fishes: support for the
competitive exclusion principle. *Ecology* **52**, 336–342 (1971).

5. Mayfield, M. M. & Levine, J. M. Opposing effects of competitive exclusion on the
phylogenetic structure of communities. *Ecol Lett* **13**, 1085–1093 (2010).

6. Grime, J. P. Competitive exclusion in herbaceous vegetation. *Nature* **242**, 344–347
(1973).

7. Letten, A. D., Ke, P. J. & Fukami, T. Linking modern coexistence theory and
contemporary niche theory. *Ecol Monogr* **87**, 161–177 (2017).

8. Chesson, P. Updates on mechanisms of maintenance of species diversity. *Journal of*
Ecology **106**, 1773–1794 (2018).

9. Chesson, P. Mechanisms of maintenance of species diversity. *Annu Rev Ecol Syst* **31**,
343–366 (2000).

10. Chase, J. M. & Leibold, M. A. *Ecological niches: linking classical and contemporary*
approaches. *Biodiversity and Conservation* (University of Chicago Press, 2004).
doi:10.1023/b:bioc.0000029366.24837.fc.

11. Hutchinson, G. E. Concluding Remarks. *Cold Spring Harb Symp Quant Biol* **22**, 415–
427 (1957).

12. Régimbeau, A. *et al.* Contribution of genome-scale metabolic modelling to niche

theory. *Ecol Lett* **25**, 1352–1364 (2022).

13. Smith, S. R. *et al.* Evolution and regulation of nitrogen flux through

compartmentalized metabolic networks in a marine diatom. *Nat Commun* **10**, 4552

(2019).

14. Palomo, A. *et al.* Comparative genomics sheds light on niche differentiation and the

evolutionary history of comammox *Nitrospira*. *ISME Journal* **12**, 1779–1793 (2018).

15. Alneberg, J. *et al.* Ecosystem-wide metagenomic binning enables prediction of

ecological niches from genomes. *Commun Biol* **3**, 119 (2020).

16. Fahimipour, A. K. & Gross, T. Mapping the bacterial metabolic niche space. *Nat*

Commun **11**, 4887 (2020).

17. Carding, S., Verbeke, K., Vipond, D. T., Corfe, B. M. & Owen, L. J. Dysbiosis of the

gut microbiota in disease. *Microb Ecol Health Dis* **26**, 26191 (2015).

18. Ravel, J. *et al.* Daily temporal dynamics of vaginal microbiota before, during and after

episodes of bacterial vaginosis. *Microbiome* **1**, 29 (2013).

19. Fujita, H. *et al.* Alternative stable states, nonlinear behavior, and predictability of

microbiome dynamics. *bioRxiv* https://doi.org/10.1101/2022.08.23.505041 (2022).

20. Lahti, L., Salojärvi, J., Salonen, A., Scheffer, M. & de Vos, W. M. Tipping elements in

the human intestinal ecosystem. *Nat Commun* **5**, 4344 (2014).

21. David, L. A. *et al.* Diet rapidly and reproducibly alters the human gut microbiome.

Nature **505**, 559–563 (2014).

22. Kho, Z. Y. & Lal, S. K. The human gut microbiome - A potential controller of wellness

and disease. *Front Microbiol* **9**, 1835 (2018).

23. Kriss, M., Hazleton, K. Z., Nusbacher, N. M., Martin, C. G. & Lozupone, C. A. Low

diversity gut microbiota dysbiosis: drivers, functional implications and recovery. *Curr*

Opin Microbiol **44**, 34–40 (2018).

24. Costello, E. K., Stagaman, K., Dethlefsen, L., Bohannan, B. J. M. & Relman, D. A. The

application of ecological theory toward an understanding of the human microbiome.

Science (1979) **336**, 1255–1262 (2012).

25. Huttenhower, C. *et al.* Structure, function and diversity of the healthy human
microbiome. *Nature* **486**, 207–214 (2012).

26. Seemann, T. Prokka: Rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068–
2069 (2014).

27. Macarthur, R. & Levins, R. The limiting similarity, convergence, and divergence of
coexisting species. *American Naturalist* **101**, 377–385 (1967).

28. Tropini, C., Earle, K. A., Huang, K. C. & Sonnenburg, J. L. The Gut Microbiome:
Connecting Spatial Organization to Function. *Cell Host Microbe* **21**, 433–442 (2017).

29. Earle, K. A. *et al.* Quantitative Imaging of Gut Microbiota Spatial Organization. *Cell
Host Microbe* **18**, 478–488 (2015).

30. Jansson, J. K. & Hofmockel, K. S. Soil microbiomes and climate change. *Nat Rev
Microbiol* **18**, 35–46 (2020).

31. Fierer, N. Embracing the unknown: Disentangling the complexities of the soil
microbiome. *Nat Rev Microbiol* **15**, 579–590 (2017).

32. Trivedi, P., Leach, J. E., Tringe, S. G., Sa, T. & Singh, B. K. Plant–microbiome
interactions: from community assembly to plant health. *Nat Rev Microbiol* **18**, 607–621
(2020).

33. Venter, J. C. *et al.* Environmental Genome Shotgun Sequencing of the Sargasso Sea.
Science (1979) **304**, 66–74 (2004).

34. Gibson, D. G. *et al.* Creation of a bacterial cell controlled by a chemically synthesized
genome. *Science* **329**, 52–6 (2010).

35. Maiden, M. C. J. *et al.* MLST revisited: The gene-by-gene approach to bacterial
genomics. *Nat Rev Microbiol* **11**, 728–736 (2013).

36. Flint, H. J., Scott, K. P., Duncan, S. H., Louis, P. & Forano, E. Microbial degradation of
complex carbohydrates in the gut. *Gut Microbes* **3**, 289–306 (2012).

37. Nowinski, B. & Moran, M. A. Niche dimensions of a marine bacterium are identified
using invasion studies in coastal seawater. *Nat Microbiol* **6**, 524–532 (2021).

38. Pereira, F. C. & Berry, D. Microbial nutrient niches in the gut. *Environ Microbiol* **19**,

447   1366–1378 (2017).

39.   Heintz-Buschart, A. & Wilmes, P. Human gut microbiome: function matters. *Trends Microbiol* **26**, 563–574 (2018).

40.   Schirmer, M. *et al.* Dynamics of metatranscription in the inflammatory bowel disease gut microbiome. *Nat Microbiol* **3**, 337–346 (2018).

41.   Turner, T. R. *et al.* Comparative metatranscriptomics reveals kingdom level changes in the rhizosphere microbiome of plants. *ISME Journal* **7**, 2248–2258 (2013).

42.   Legendre, P. & de Cáceres, M. Beta diversity as the variance of community data: Dissimilarity coefficients and partitioning. *Ecol Lett* **16**, 951–963 (2013).

43.   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J* **17**, https://doi.org/10.14806/ej.17.1.200 (2011).

44.   Chen, S., Zhou, Y., Chen, Y. & Gu, J. Fastp: An ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**, i884–i890 (2018).

45.   Bankevich, A. *et al.* SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing. *Journal of Computational Biology* **19**, 455–477 (2012).

46.   Uritskiy, G. v., DiRuggiero, J. & Taylor, J. MetaWRAP—a flexible pipeline for genome-resolved metagenomic data analysis. *Microbiome* **6**, 158 (2018).

47.   Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM: Assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* **25**, 1043–1055 (2015).

48.   Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun* **9**, 5114 (2018).

49.   Woodcroft B. CoverM: program available at https://github.com/wwood/CoverM. Preprint at (2021).

50.   Chaumeil, P. A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: A toolkit to classify genomes with the genome taxonomy database. *Bioinformatics* **36**, 1925–1927 (2020).

51.   Parks, D. H. *et al.* GTDB: an ongoing census of bacterial and archaeal diversity
through a phylogenetically consistent, rank normalized and complete genome-based
taxonomy. *Nucleic Acids Res* **50**, D785–D794 (2022).

52.   Kanehisa, M., Sato, Y. & Morishima, K. BlastKOALA and GhostKOALA: KEGG
Tools for Functional Characterization of Genome and Metagenome Sequences. *J Mol*
Biol **428**, 726–731 (2016).

53.   Graham, E. D., Heidelberg, J. F. & Tully, B. J. Potential for primary productivity in a
globally-distributed bacterial phototroph. *ISME Journal* **12**, 1861–1866 (2018).

54.   Kandeler, E. & Gerber, H. Short-term assay of soil urease activity using colorimetric
determination of ammonium. *Biol Fertil Soils* **6**, 68–72 (1988).

55.   Hood-Nowotny, R., Umana, N. H.-N., Inselbacher, E., Oswald- Lachouani, P. &
Wanek, W. Alternative methods for measuring inorganic, organic, and total dissolved
nitrogen in soil. *Soil Science Society of America Journal* **74**, 1018–1027 (2010).

56.   Taylor, B. W. *et al.* Improving the fluorometric ammonium method: Matrix effects,
background fluorescence, and standard additions. *J North Am Benthol Soc* **26**, 167–177
(2007).

Fig. 1 | Community and ecosystem dynamics. a, Time-series data of community structure.

For the replicate microcosm that showed the most abrupt community compositional changes

through the 110-day microbiome experiment[19] (Supplementary Fig. 1), family-level

taxonomic compositions inferred with 16S rRNA sequencing are shown. The blue line

represents the speed and magnitude of community compositional changes around each time

point ("abruptness" index[19]; see Methods). The red line indicates α-diversity (Shannon's H')

of microbial ASVs[19]. Note that a value larger than 0.5 represents turnover of more than 50 %

of microbial ASV compositions. See Supplementary Figure 1 for color profiles of bacterial

families. Reproduced from the data of a previous study[19]. **b**, Taxonomic compositions inferred

with whole-genome shotgun sequencing. At each of the 13 time points through the time-series

of the target microcosm, the relative abundance of each MAG was estimated based on the

normalized read coverage value (reads per kilobase of genome per million reads mapped). **c**,

Genome size and GC nucleotide content of the MAGs detected in the target microcosm. See

panel **b** for colors and symbols.

Fig. 2 | Metabolic pathway/process profiles of the MAGs. KEGG metabolic pathways/profiles of the reconstructed bacterial genomes (MAGs) are shown. The detection (relative abundance > 0.1 %) of each microbial MAG on each day within the whole-genome shotgun data is indicated in the panel below. Only the microbial MAGs with > 80 % completeness and < 5 % contamination were included (Supplementary Table 1). The five MAGs that co-occurred from Day 40 to 60 and metabolic pathways/processes mentioned in the main text are highlighted. Only the metabolic pathways/processes with highly heterogeneous patterns across microbial MAGs are shown. See Supplementary Figure 3 for

517   detailed profiles of the metabolic pathways/processes.

Fig. 3 | Metagenomic niche space. **a**, Distributions of MAGs within metagenomic niche space. Based on dissimilarity in gene repertoires, microbial MAGs that appeared in the time-series of the target microcosm were plotted on the three-dimensional space defined by the principal coordinate analysis (PCoA) of 6,999 genes. **b**, Changes in the distributions of microbial MAGs within niche space. At each time point, detected MAGs (relative abundance > 0.1 %) were plotted on the space defined in the multivariate analysis in the in the panel **a**.

Fig. 4 | Dynamics of niche-overlap level. a, Community-level profiles of metabolic

pathways/processes and niche overlap index. The niche overlap index was defined as $1 - \bar{\beta}$,

where $\bar{\beta}$ was mean Jaccard dissimilarity (β-diversity) of gene compositions between pairs of

the microbial MAGs detected at a target time point. The scores of the niche overlap index

were shown on a PCoA surface representing community-level compositions of genes. On the

PCoA surface, time points are distributed based on the sum of the gene repertoires of the

detected MAGs. **b**, Dynamics of niche-overlap levels. Niche overlap scores are shown across

the time-series.

Fig. 5 | Niche overlap level and community compositional shifts. The magnitude of

community compositional changes observed in the microbiome was regressed on niche

overlap index obtained based on the whole-genome shotgun analysis. Niche overlap index at

each time point and time-series shifts in community structure (Bray-Curtis β-diversity

between present and next time points through the time-series of the shotgun metagenomic

data) are shown along horizontal and vertical axes, respectively. The regression line is shown

with 95 % confidence intervals.

Supplementary Figure Captions

Supplementary Fig. 1 | Dynamics of family-level community structure. The dynamics of microbial family-level compositions were visualized based on the 16S rRNA sequencing data of the previous study[19]. The replicate microcosm (replicate no. 5 in Water/Medium-A treatment), which is subjected to the whole-genome shotgun sequencing analysis, is highlighted. The blue line represents the speed and magnitude of community compositional changes around each time point ("abruptness" index[19]; see Methods). The red line indicates α-diversity (Shannon's H') of microbial ASVs[19]. Reproduced from the data of a previous study[19].

Supplementary Fig. 2 | Overview of the whole-genome shotgun sequencing data. a, Comparison of relative abundance of bacterial taxa (families) between 16S rRNA amplicon sequencing[19] (reproduced from the previous study[19]; top) and whole-genome shotgun sequencing (this study; bottom). **b**, Correlation between the family-level relative abundance of 16S rRNA and whole-genome shotgun sequencing data (Spearman's correlation; $\rho = 0.667$, df = 794, $P < 0.05$). Each point represents each family at each time point. **c**, Background chemical properties. Changes in NO_3^- and NH_4^+ concentrations in the ecosystem are shown for the time points with whole-genome shotgun metagenomic data.

Supplementary Fig. 3 | Detailed information of the metabolic pathway/process profiles of the MAGs. The KEGG metabolic pathways/processes of the reconstructed bacterial genomes (MAGs) are shown. The detection (relative abundance > 0.1 %) of each microbial MAG on each day within the whole-genome shotgun data is indicated in the panel below. Only the microbial MAGs with > 80 % completeness and < 5 % contamination were included (Supplementary Table 1). The five MAGs that co-occurred from Day 40 to 60 and metabolic pathways/profiles mentioned in the main text are highlighted. The detailed definition of the KEGG metabolic pathways/processes is available at https://github.com/bjtully/BioData/blob/master/KEGGDecoder/KOALA_definitions.txt.