Dating in the Dark: Elevated Substitution Rates in Cave Cockroaches

(Blattodea: Nocticolidae) Have Negative Impacts on Molecular Date

Estimates

Toby G. L. Kovacs[1], James Walker[2], Simon Hellemans[3], Thomas Bourguignon[3,4],

Nikolai J. Tatarnic[5,6], Jane M. McRae[7], Simon Y. W. Ho[1], and Nathan Lo[1]

[1]School of Life and Environmental Sciences, University of Sydney, Sydney NSW 2006,

Australia

[2]Department of Agriculture, Fisheries and Forestry, Canberra ACT 2601, Australia

[3]Okinawa Institute of Science & Technology Graduate University, 1919-1 Tancha, Onna-son,

Okinawa 904-0495, Japan

[4]Faculty of Tropical AgriScience, Czech University of Life Sciences, Kamýcka 129, 16521

Prague, Czech Republic

[5]Collections & Research, Western Australian Museum, 49 Kew Street, Welshpool WA 6106,

Australia

[6]Centre for Evolutionary Biology, The University of Western Australia, Perth WA 6009,

Australia

[7]Bennelongia Environmental Consultants, 5 Bishop Street, Jolimont WA 6014, Australia

Abstract.—Rates of nucleotide substitution vary substantially across the Tree of Life, with potentially confounding effects on phylogenetic and evolutionary analyses. A large acceleration in mitochondrial substitution rate occurs in the cockroach family Nocticolidae, which predominantly inhabit subterranean environments. To evaluate the impacts of this among-lineage rate heterogeneity on estimates of phylogenetic relationships and evolutionary timescales, we analysed nuclear ultraconserved elements (UCEs) and mitochondrial genomes from nocticolids and other cockroaches. Substitution rates were substantially elevated in nocticolid lineages compared with other cockroaches, especially in mitochondrial protein-coding genes. This disparity in evolutionary rates is likely to have led to different evolutionary relationships being supported by phylogenetic analyses of mitochondrial genomes and UCE loci. Furthermore, Bayesian dating analyses using relaxed-clock models inferred much deeper divergence times compared with a flexible local clock. Our phylogenetic analysis of UCEs, which is the first genome-scale study to include all ten major cockroach families, unites Corydiidae and Nocticolidae and places Anaplectidae as the sister lineage to the rest of Blattoidea. We uncover an extraordinary level of genetic divergence in Nocticolidae, including two highly distinct clades that separated ~115 million years ago despite both containing representatives of the genus *Nocticola*. The results of our study highlight the potential impacts of high among-lineage rate variation on estimates of phylogenetic relationships and evolutionary timescales.

Keywords: Nocticolidae, ultraconserved elements, molecular dating, divergence times, among-lineage rate heterogeneity, clock models, flexible local clock

Rates of molecular evolution vary by several orders of magnitude across the Tree of Life.

This among-lineage rate variation has been linked to differences in life-history traits such as

body size, population size, longevity, metabolic rate, and generation time (Bromham 2009).

Remarkably, large disparities in evolutionary rates can even be found between closely related

lineages, such as in burrowing crayfish (Gan et al. 2018), crabronid wasps (Kaltenpoth et al.

2012), and parasitic plants (Lemaire et al. 2011; Bromham et al. 2013).

When molecular evolutionary rates show extreme or complex variation across

lineages, they can pose difficulties for phylogenetic inference and molecular dating

(Kolaczkowski and Thornton 2004, 2009; Dornburg et al. 2012; Crisp et al. 2014; Susko

2015; Roch et al. 2019). A range of clock models have been developed in order to account for

among-lineage rate variation in Bayesian phylogenetic dating analyses. These include local

clocks, in which different rates are allowed in groups of neighbouring branches, and relaxed

clocks, in which a different rate is allowed along each branch (Ho and Duchêne 2014). The

placement of local clocks can either be selected a priori or jointly estimated with the

phylogeny, which is possible using the random local clock (Drummond and Suchard 2010)

and the shrinkage-based random local clock (Fisher et al. 2021). The most widely used

relaxed clocks either treat the branch rates as autocorrelated (Thorne et al. 1998; Sanderson

2002; Lepage et al. 2006), where they are assumed to be related between adjacent branches,

or uncorrelated, where they are drawn independently from a chosen parametric distribution

(Drummond et al. 2006; Rannala and Yang 2007). These relaxed clocks typically rely on the

branch rates fitting a simple unimodal distribution and might provide a poor fit under

conditions of complex rate variation.

Although rate variation is ubiquituous across the tree of life (Ho 2020), few studies

have focused on the behaviour and performance of clock models when challenged with

extreme among-lineage rate variation. Studies of simulations have demonstrated that

misspecification of relaxed-clock models can lead to imprecise estimates of evolutionary

rates and timescales (Dornburg et al. 2012; Duchêne et al. 2014; Fourment and Darling 2018).

There has also been varying support for uncorrelated and autocorrelated relaxed clocks, with

substantial impacts on the inference of evolutionary histories (Lepage et al. 2007; Linder et al.

2011; Ho et al. 2015; dos Reis et al. 2018). When applied to phylogenies with among-lineage

rate variation, uncorrelated relaxed clocks tend to infer a distribution of substitution rates

centred on the average rate across the tree, leading to poor estimation of divergence times

(Dornburg et al. 2012; Crisp et al. 2014). Data sets containing such rate variation produce

more intuitive date estimates when analysed using a random local clock (Dornburg et al.

2012; Crisp et al. 2014), although this might depend on the distribution and extent of rate

variation across the tree (Gan et al. 2018). Despite its potential utility, the random local clock

can be difficult to implement and so other clock models are usually preferred in practice

(Drummond and Suchard 2010; Dornburg et al. 2012).

The flexible local clock combines some of the features of local clocks and relaxed

clocks by allowing multiple, independent uncorrelated relaxed clocks for different clades

across the tree (Fourment and Darling 2018). This model can capture large-scale rate

disparities among clades while also accounting for local variation in rates. Although the

flexible local clock provides an interesting framework that allows distinct evolutionary rates

between clades, it has not been widely employed. This is partly due to the challenge of

selecting local clocks a priori, but possibly also because its implementation is non-trivial. To

date, there has been no study that has compared the behaviour of a diverse range of clock

models, including local clocks and relaxed clocks, when challenged by extreme rate variation

among lineages.

One group of organisms that appears to have experienced a dramatic shift in

mitochondrial substitution rates is the cockroach family Nocticolidae (Lo et al. 2007;

Legendre et al. 2008, 2015; Djernæs et al. 2015; Wang et al. 2017; Bourguignon et al. 2018;

Li and Huang 2020). The causes and timing of this rate elevation, and whether such a pattern

also occurs in the nuclear genome of nocticolids, are unknown. Dominated by cavernicolous

taxa exhibiting troglomorphic (cave adapted) characters, nocticolids are small (<10 mm body

length), pale, and delicate, and often have absent or reduced eyes and wings (Fig. 1). There

are 38 described species of Nocticolidae across nine genera, found in Africa, Asia, and

Oceania (Supplementary Table S1). The family is in desperate need of taxonomic revision,

with 26 species placed in a poorly defined genus (*Nocticola*) accompanied by six monotypic

genera, a genus (*Alluaudellina*) that is likely to be a synonym of *Nocticola* (Chopard 1932;

Roth 1988), and a genus (*Spelaeoblatta*) containing four species only from Thailand and

Myanmar (Roth and McGavin 1994; Vidlička et al. 2003). There are even doubts that four of

the genera (including *Spelaeoblatta*) should be included in the family at all, based on

morphology (Li and Huang 2020). The majority (24) of species are subterranean, including

three from non-cave subterranean environments (Trotter et al. 2017) and two that are known

to move in and out of caves. Ten species are epigean and three are termitophilous (living

within termite nests). Epigean taxa typically have fully developed wings, compound eyes,

and dark pigmentation, while subterranean taxa rarely possess these features. Owing to their

small size, great speed, and inaccessible habitats, nocticolids remain the most poorly studied

of the ten recognized cockroach families (Karny 1924; Fernando 1957; Asahina 1974; Roth

1988).

The phylogenetic position of Nocticolidae remains uncertain because of limited taxon

sampling in the family. Several phylogenetic studies have found a close relationship between

Nocticolidae and the family Corydiidae, though these analyses have typically included only a

small number of nocticolids (Inward et al. 2007; Legendre et al. 2008; Djernæs et al. 2012;

Legendre et al. 2015; Djernæs et al. 2015; Wang et al. 2017; Bourguignon et al. 2018;

Evangelista et al. 2019; Li and Huang 2020; Djernæs and Murienne 2022). There has so far

been no phylogenomic analysis of all ten cockroach families, nor has there been a

comprehensive phylogenetic study of nocticolids themselves. Furthermore, previous genome-

scale dating analyses of Blattodea have lacked sufficient sampling of Nocticolidae to infer

divergence times within the family. A single specimen was included in an analysis of

transcriptomes that inferred minimal among-lineage rate variation and a Nocticolidae–

Corydiidae divergence in the Jurassic (Evangelista et al. 2019). Contrastingly, a single

specimen was included in an analysis of mitochondrial genomes that inferred a long

nocticolid branch and an older Nocticolidae–Corydiidae divergence in the Triassic, possibly

because the higher substitution rate pushed back the date estimate for this node (Bourguignon

et al. 2018).

Studies based on a handful of mitochondrial and nuclear molecular markers have

included up to six representatives of Nocticolidae, again highlighting increased substitution

rates across the family and deep divergences (100–200 Ma) among representatives of the

genus *Nocticola* (Djernæs et al. 2015; Legendre et al. 2015; Wang et al. 2017; Li and Huang

2020). Li and Huang (2020) used an autocorrelated relaxed clock to infer large increases in

substitution rates within Nocticolidae and the sister lineage Latindiinae, but their divergence

times remain uncertain due to sparse taxon sampling and imprecise calibrations. A

population-level study of three Australian species of *Nocticola* found very large

mitochondrial COI distances (17.3–25.8 %) between species, indicative of high substitution

rates, deep divergences, or both (Trotter et al. 2017). There has not yet been any

phylogenomic analysis of a diverse range of nocticolids, so uncertainties about their

taxonomy and evolutionary history have persisted.

Here we evaluate the impact of extreme among-lineage rate variation on the inference

of phylogenetic relationships and evolutionary timescales in Nocticolidae and Blattodea. We

present a detailed phylogenetic analysis of all ten recognized families of cockroaches, using

the largest data set so far assembled for Nocticolidae. We design the first baits for

ultraconserved elements (UCEs) from cockroaches, and assemble a data set comprising UCE

loci and mitochondrial genomes. Our study resolves the major branching order of all ten

cockroach families, identifies two highly divergent and rapidly evolving clades within

Nocticolidae, and shows that widely used relaxed-clock models have difficulty in accounting

for lineage-specific accelerations in evolutionary rates.

MATERIALS AND METHODS

We generated three genetic sequence data sets for our phylogenetic analyses of nocticolids

and other cockroaches. We began by using a barcoding marker, *16S*, to identify specimens

for genetic sequencing, given that the majority of samples either represented undescribed

species or could not be identified based on morphology. From the observed diversity in the

family, we chose 22 samples for short-read metagenomic 'genome skim' sequencing. Using

these libraries, we assembled our first data set, comprising sequences from mitochondrial

protein-coding genes (mtPCG), and our second data set, comprising sequences from nuclear

ultraconserved elements (UCE). We combined sequences of UCE loci, mitochondrial

protein-coding genes, and mitochondrial *16S* to form a third data set with maximum taxon

coverage (UCE-mtPCG-*16S*).

Taxon Sampling and Sequencing

We sampled 131 specimens of Nocticolidae from museums and private collections. The

majority of samples were from Australia (including Christmas Island), but we also included

taxa from New Guinea (Indonesia) and Singapore. For non-type samples and taxa with

multiple specimens, genomic DNA was extracted from whole organisms. For type specimens

or individuals from unique locations for which no known species have been described, a

single hind leg was used. We used PCR to amplify a 440 bp fragment of the mitochondrial

gene encoding 16S rRNA (*16S*). These 115 newly generated sequences were combined with

six nocticolid sequences available on GenBank, as well as outgroup sequence data from 37

non-nocticolid cockroaches and termites and five other insects (Supplementary Tables S2–

S4). We aligned these sequences using MAFFT 7.475 (Katoh and Toh 2008) and manually

removed poorly aligned regions. The final *16S* alignment comprised 438 bp from 163 taxa.

Using this alignment, we estimated phylogenetic relationships using maximum likelihood in

IQ-TREE v2 (Bui et al. 2020) to identify representative taxa for further sequencing.

For our first data set, we selected 23 representative lineages (Supplementary Table

S5) for short-read metagenomic 'genome skim' sequencing by BGI (Shenzhen, China). Using

the read libraries produced, we assembled mitochondrial genomes by mapping reads to an

available nocticolid mitochondrial genome (Bourguignon et al. 2018) in Geneious 2020.0.5

(www.geneious.com). Samples with incomplete mapping were assembled *de novo* using

NOVOplasty (Dierckxsens et al. 2017) and the resulting assembly was then used as a

reference for read mapping. We annotated mitochondrial genes using MITOS2 (Donath et al.

2019). Mitochondrial genomes from nocticolids were combined with those from

representatives of the diversity in each of the nine other cockroach families as well as

outgroup taxa from related insect lineages (Supplementary Table S6; Bourguignon et al.

2018). Nucleotide sequences of each of the 13 mitochondrial protein-coding genes were

aligned at the amino acid level using MUSCLE through the TranslatorX web server (Abascal

et al. 2010). Using PhyloMAd (Duchêne et al. 2018, 2022), we found evidence of saturation

at first and third codon sites and removed these from the sequences. The resulting data set

comprised the second codon sites (3793 bp) of the 13 mitochondrial protein-coding genes

(mtPCG) from 23 nocticolid taxa and 87 outgroup taxa.

To construct our second data set, we designed UCE baits using PHYLUCE 1.6.6

(Faircloth 2016). Baits were designed from whole genomes of five cockroaches and one

termite, using the same parameters as Hellemans et al. (2022) for the termite-specific set. Our

sampling for this approach included our nocticolid taxa, at least one representative from each

of the other nine cockroach families, as well as one mantid and five phasmids as outgroup

taxa. Reads were assembled using metaSPAdes 3.13 (Nurk et al. 2017), and UCEs were

extracted and aligned using the PHYLUCE suite. The concatenated UCE data set comprised

1676 loci (total length 383,325 bp) from 61 samples (Supplementary Table S7). To maximize

the taxon coverage in Nocticolidae and the outgroup, we constructed a third data set by

combining the UCE data with mtPCG and *16S* sequences (UCE-mtPCG-*16S*). Further details

of taxon selection, DNA extraction, amplification, library preparation, museum vouchers,

sample locations, and accession numbers are given in the Supplementary Material.

Phylogenetic Analyses

We inferred phylogenetic relationships using the mtPCG, UCE, and combined UCE-mtPCG-

16S data sets using maximum likelihood in IQ-TREE. For our data sets containing UCEs

(UCE and UCE+mtPCG+*16S*), we selected the partitioning scheme in ModelFinder using a

greedy algorithm to merge individual UCEs and mitochondrial genes (-m TESTMERGE)

(Lanfear et al. 2012; Chernomor et al. 2016). We allowed branch lengths to be proportionate

among subsets of the data (-p), which has been found to result in the highest statistical

support in many cases (Duchêne et al. 2020). The ModelFinder option in IQ-TREE was used

to select the best-fitting substitution models according to the Bayesian information criterion

(Kalyaanamoorthy et al. 2017). We also performed Bayesian phylogenetic analyses using

ExaBayes v1.5 (Aberer et al. 2014) for the mtPCG and UCE data sets, checking convergence

over four independent runs. In these analyses, we managed computational load by running

the analysis with the GTR+G substitution model on an unpartitioned data set.

To account for gene-tree incongruence among loci, we also analysed the UCE data set

using the summary-coalescent method in ASTRAL-III (Zhang et al. 2018). The gene tree for

each UCE locus was estimated using IQ-TREE with substitution models selected using

ModelFinder. In order to compare substitution rates across Blattodea, we calculated the mean

root-to-tip distance for each cockroach family based on the maximum-likelihood trees

inferred from the mtPCG and UCE data sets. We assessed the adequacy of substitution

models using PhyloMAd (Duchêne et al. 2018).

We inferred the evolutionary timescale of cockroaches using Bayesian phylogenetic

analyses of the mtPCG data set using BEAST v2.5 (Bouckaert et al. 2019) and the

approximate likelihood method in MCMCTree (Yang 2007; dos Reis and Yang 2011). We

also analysed our UCE data set in MCMCTree. In BEAST, we ran analyses using five

different clock models: strict clock, uncorrelated exponential relaxed clock, uncorrelated

lognormal relaxed clock, random local clock, and flexible local clock. We used the flexible

local clock to model substitution rates across the tree under two independent lognormal

distributions, one within Nocticolidae (including its stem branch) and one for the rest of the

tree. The five clock models were compared using marginal likelihoods estimated by nested

sampling (Maturana Russel et al. 2019). We used a set of fossil calibrations selected by

Evangelista et al. (2019), with the addition of *Crenocticola* (Li and Huang 2020), to inform

minimum bounds of uniform distributions (Supplementary Table S8). Both the mtPCG and

UCE data sets were analysed in MCMCtree using the two available relaxed clock models: the

independent (uncorrelated) lognormal relaxed clock and autocorrelated relaxed clock. Our

UCE data set had fewer outgroup taxa, reducing the number of fossil calibrations that could

be applied; we added secondary calibrations for the root of the tree and crown Dictyoptera,

based on estimates from a recent phylogenomic study (Evangelista et al. 2019).

In order to test the influence of including Nocticolidae on estimates of topology and

divergence times, we repeated all of our phylogenetic analyses (IQTREE, ExaBayes, BEAST,

and MCMCtree) using the mtPCG and UCE data sets after removing the nocticolid taxa. For

our Bayesian dating analyses, we compared all of the clock models except for the flexible

local clock (for which a separate relaxed clock had previously been assigned to Nocticolidae).

A detailed description of all methods can be found in the Supplementary Material.

RESULTS

Phylogenetic Relationships

Phylogenetic relationships among cockroach families could not be confidently resolved using

the mtPCG data set alone (Supplementary Fig. S2−S4). In contrast, our analyses of the UCE

data set produced well-resolved phylogenetic trees with strong support for relationships

among cockroach families (Fig. 1, Supplementary Fig. S5−S7). All of our phylogenetic

analyses strongly supported the monophyly of Nocticolidae (Fig. 1, Supplementary Fig. S1-

8). However, the inferred relationships between Nocticolidae and the other cockroach

families differed between our mtPCG and UCE analyses. In our mtPCG analyses, the

relationship between Nocticolidae and the other cockroach families varied between

maximum-likelihood and Bayesian analyses and even between replicate Bayesian analyses,

with Anaplectidae usually being placed as the sister group. In our analyses of UCE data,

Corydiidae was consistently and confidently placed as the sister group to Nocticolidae (Fig.

1; UFb = 100, SH-aLRT = 100, ASTRAL-PP = 0.99, ExaBayes-PP = 1) and

FIGURE 1. Evolutionary timescale of Blattodea, with Mantodea and Phasmatodea as
outgroup taxa. Divergence times were estimated based on UCE loci, using a Bayesian
phylogenomic approach with an uncorrelated relaxed clock in MCMCTree. The fixed tree
topology used in this analysis was inferred using maximum-likelihood and Bayesian analyses
of UCE loci. Node support from these analyses is represented by squares: outline colours
indicate values of the SH-like approximate likelihood-ratio test (SH-alrt), fill colours indicate
ultrafast bootstrap (UFb) values. Nodes with SH-alrt and UFb > 99 are not labelled. Posterior
probabilities were 1.0 for all nodes, with the exception of the node labelled with a star, which
had a different topology. Insert shows the diverse morphology present in Nocticolidae: a)
female *Nocticola australiensis* (Royal Arch Cave) with highly reduced eyes (photograph by
Braxton Jones); b) blind female *Nocticola baumi* with elongated appendages (photograph by
Martin Bláha); and c) male *Nocticola pheromosa* with fully developed eyes and wings
(photograph by Maimon Hussin).

Anaplectidae was inferred to be the sister lineage to the rest of Blattoidea, with high support

(UFb = 100, SH-aLRT = 100, ASTRAL-PP = 0.99, ExaBayes-PP = 1). After removing

nocticolid taxa, we found that the phylogenetic relationships among cockroach families could

still not be confidently resolved using the mtPCG data set (Supplementary Fig, S13-15).

However, the topologies inferred using the UCE data set were congruent to those from our

analyses that included Nocticolidae (Supplementary Fig, S16-17). Our assessment of

substitution model adequacy found that the full mtPCG data set had a high risk of biased

inferences according to the consistency index. This statistic reported a low risk of biased

inferences after removing nocticolid taxa. Unfortunately our UCE data set contained too

much missing data to assess substitution model adequacy.

286   Within Nocticolidae, our analyses found uniformly strong support for the mutual

monophyly of two divergent clades. We refer to these clades as Nocticolidae A and B. The

inferred relationships within each of these two clades varied slightly between mtPCG and

UCE analyses. Here we focus on the results of our analyses of the data sets that included the

UCE loci, which yielded the highest phylogenetic support. Within Nocticolidae, consistent

relationships were found between our analyses of the UCE data set (Fig. 1) and the

UCE+mtPCG+*16S* data set (Fig. 2). A nocticolid sequence obtained from GenBank

("*Nocticola* sp. Unknown"; Legendre et al. 2015), for which collection location details are

not available, was found to be the sister lineage to all other nocticolids examined.

295   Nocticolidae A contains approximately half of the Australian mainland taxa, nested

within three lineages containing taxa from Vietnam, Singapore and Malaysia, and Christmas

Island, respectively (Fig. 2). Among the Australian taxa in Nocticolidae A, Queensland taxa

formed the sister clade to a group comprising taxa from Western Australia and the Northern

Territory. The second clade, Nocticolidae B, comprised a sister grouping between the

remaining Australian taxa and one taxon from New Guinea (*N. baumi*). The Queensland taxa,

FIGURE 2. Phylogeny of Nocticolidae. Maximum-likelihood tree estimated using

concatenated UCE loci, second codon sites of 13 mitochondrial protein-coding genes, and

mitochondrial *16S*. There are two major clades in Nocticolidae, A (numbers in black) and B

(numbers in grey). Node support is represented by squares: outline colours indicate values of

the SH-like approximate likelihood-ratio test (SH-alrt), fill colours indicate ultrafast bootstrap

(UFb) values. Nodes with SH-alrt and UFb = 100 are not labelled. Colours represent the

geographical location of taxa from the Australian mainland (blue), South-East Asia (green),

Christmas Island (orange), and New Guinea (purple).

N. babindaensis and *N*. sp. (Mt Haig), again formed the sister clade to taxa from Western

Australia and the Northern Territory.

Evolutionary Timescales and Substitution Rates

Our phylogenetic analyses suggest that an acceleration in mitochondrial substitution rates

occurred early in the evolution of Nocticolidae. In our maximum-likelihood analyses of

the mtPCG and UCE data sets, root-to-tip distances for nocticolid clades were considerably

greater than those of other cockroach families (Fig. 3). For the mtPCG data set, Nocticolidae

A and B had root-to-tip distances that were 3–5 times those of representatives of other

cockroach families. In comparison, the UCE sequences showed root-to-tip distances for

nocticolids that were ~1.5 times those of other cockroach families.

In the Bayesian phylogenetic analysis of the mtPCG data set using BEAST, the

flexible local clock was decisively preferred over other clock models (minimum log Bayes

factor = 30.8, SD = 6.0, Supplementary Table S9). The flexible local clock inferred a

substantially higher rate in Nocticolidae (posterior mean 3.7×10^{-3} substitutions/site/Myr; 95%

CI $2.6–6.3 \times 10^{-3}$) than across the rest of the tree (posterior mean 8.0×10^{-4}

substitutions/site/Myr; 95% CI $6.4–9.4 \times 10^{-4}$) (Fig. 4). Our analyses using the uncorrelated

exponential and lognormal relaxed-clock models inferred very high rates along the stem

branches leading to Nocticolidae and to Nocticolidae A and B. Within each of these clades,

however, the inferred rates were similar to the background rate across the tree (Fig. 4).

Our Bayesian analyses inferred deeper divergence times when using mtPCG data than

UCE data. Among our analyses of the mtPCG data, the flexible local clock produced the

youngest estimates of divergence times, particularly for crown Nocticolidae (Fig. 4). These

estimates were the most similar to those inferred using the UCE data set. Our analyses using

the random local clock estimated the second most similar divergence times to those inferred

FIGURE 3. Accelerated mitochondrial substitution rates in Nocticolidae compared

with other families in Blattodea (cockroaches). (a) Maximum-likelihood tree of Blattodea

inferred using the second codon sites of the 13 mitochondrial protein-coding genes. (b)

Phylogenetic root-to-tip distances for the concatenated mitochondrial protein-coding genes

and the concatenated UCE loci. Root-to-tip distances were calculated for each of the major

cockroach families, with Nocticolidae split into its two major clades, A and B. Root-to-tip

distances were normalized by dividing the mean for each family by the mean root-to-tip

distance across all taxa.

using the UCE data set, and also identified a clear distinction between substitution rates in

Nocticolidae and across the rest of the tree. The strict clock estimated the oldest dates for

nodes within Nocticolidae, presumably to accommodate an excess of substitutions in

nocticolid lineages.

In the molecular dating analysis of the mtPCG data set using MCMCtree, the

autocorrelated relaxed clock gave very similar estimates of substitution rates and divergence

times to those inferred using the flexible local clock in BEAST (Fig. 4). The independent

lognormal relaxed clock in MCMCtree gave similar estimates of substitution rates and

divergence times to those inferred using the uncorrelated relaxed-clock models in BEAST

(Fig. 4). For our UCE data set, the independent lognormal and autocorrelated relaxed clocks

yielded different estimates of substitution rates and divergence times. The independent

lognormal relaxed clock estimated slightly higher substitution rates in Nocticolidae compared

with the rest of the tree, aligning with the results from our investigation of root-to-tip

distances (Supplementary Fig. S10). The autocorrelated relaxed clock estimated that the

highest substitution rates occurred along the branches leading to the phasmid and mantid

outgroups, the early branches among the major cockroach families including the stem of

Noctioclidae, and the stem branches leading to the two nocticolid clades (Supplementary Fig.

S10). Estimated substitution rates within each of the families were generally lower than those

along the respective stem branches, with branches within each of the major Nocticolidae

clades having the lowest rates in the tree (Supplementary Fig. S10). Consequently, for the

UCE data set the autocorrelated relaxed clock yielded older estimates of divergence times

within Nocticolidae than did the independent lognormal relaxed clock (Fig. 4).

Our dating analyses after removing nocticolid taxa support our contention that high

rates in Nocticolidae are driving older estimates of divergence times using some clock

models. For the mtPCG data set, the dates of crown Blattodea and Dictyoptera estimated

FIGURE 4. Estimates of divergence times and substitution rates using a range of molecular-clock models. (a) Dates for crown Nocticolidae, crown Nocticolidae A, and crown Nocticolidae B estimated from mitochondrial protein-coding genes using a strict clock (strict), uncorrelated lognormal relaxed clock (UCLN), uncorrelated exponential relaxed clock (UCEXP), random local clock (RLC), flexible local clock (FLC), autocorrelated relaxed clock (AC), and independent lognormal relaxed clock (ILN). Date estimates for the same nodes using our UCE data set are also presented for comparison using an autocorrelated relaxed clock (UCE-AC) and independent lognormal relaxed clock (UCE-ILN). Circles indicate posterior means and bars indicate 95% credibility intervals. b) Estimates of mitochondrial substitution rates in Nocticolidae compared with other families in Blattodea (cockroaches). Violin plots represent the mean substitution rate (substitutions per site per Myr) for all terminal and internal branches within Nocticolidae (including stem) and outside Nocticolidae (non-Nocticolidae). Rates were estimated using concatenated mitochondrial protein-coding genes.

using the lognormal relaxed clock, exponential relaxed clock, and strict clock were reduced

by the removal of nocticolid taxa, and similar to those inferred using the flexible local clock

when including Nocticolidae (Supplementary Figures S15,20).

We consider our estimate of evolutionary timescales based on UCE data and the

independent lognormal relaxed clock to be the most reliable for various reasons. First, the

inferences are based on a large genome-scale data set with representatives from all 10

cockroach families. Second, the UCE data have low among-lineage rate variation, especially

compared with the mtPCG data set. Third, the independent lognormal relaxed clock appears

to be the most appropriate because the distribution of branch rates estimated by the

autocorrelated relaxed clock is inconsistent with the root-to-tip distances; branches within

Nocticolidae were inferred to have a lower substitution rate than the rest of the tree, despite

its taxa having greater root-to-tip distances. We believe that this led to overestimation of

divergence times in Nocticolidae using this model.

Based on our analysis of UCE data with the independent lognormal relaxed clock, the

most recent common ancestor of cockroaches and mantids is dated at 272 Ma (95% CI 241–

292 Ma), while that of cockroaches is dated at 209 Ma (95% CI 186–233 Ma). The age of

crown Nocticolidae was inferred to be 115 Ma (95% CI 93–138 Ma), with the family

diverging from Corydiidae at 172 Ma (95% CI 152–193 Ma). The crown age of Nocticolidae

A was estimated at 39 Ma (95% CI 32–48 Ma), with the Australian lineages diverging from

each other at 21 Ma (95% CI 17–26 Ma). Nocticolidae B was inferred to have a crown age of

42 Ma (95% CI 34–51 Ma), with the Australian lineages diverging from each other at 32 Ma

(95% CI 26–39 Ma).

DISCUSSION

Our analyses have highlighted a substantial elevation of evolutionary rates in the

mitochondrial genomes of nocticolids, particularly along the basal branches of the clade, and

that a similar but smaller increase in rates has occurred in the nuclear genome. These results

are partly consistent with previous findings of long nocticolid branches in mitochondrial trees

(Lo et al. 2007; Legendre et al. 2008, 2015; Djernæs et al. 2015; Wang et al. 2017;

Bourguignon et al. 2018; Li and Huang 2020) but not in trees inferred from nuclear protein-

coding genes from transcriptomic data (Evangelista et al. 2019). We have also found that

some of the widely used relaxed-clock models are unable to accommodate the large

disparities in mitochondrial substitution rates across branches. This rate heterogeneity is

likely to be at least partly responsible for the differences in evolutionary relationships and

divergence times between the phylogenetic trees inferred from mitochondrial genomes and

nuclear ultraconserved elements.

Molecular Dating When Rates Vary

Mitochondrial substitution rates differed by a factor of more than 3 between nocticolids and

other cockroaches, representing a marked elevation of rates in a single family compared with

its relatives. This disparity in rates was apparent in the root-to-tip distances of the phylograms

and was identified in the branch rates inferred under the flexible local clock and random local

clock in BEAST, as well as the autocorrelated relaxed clock in MCMCtree. The ability of

these models to infer such highly distinct rates across the tree aligns with previous

applications of these models, and are expected to provide a better fit under conditons of

complex rate variation among lineages (Fourment and Darling 2018; Drummond and Suchard

2010). Among the clock models compared in BEAST, the flexible local clock had the highest

marginal likelihood and yielded mitochondrial estimates of divergence times that were most

similar to those inferred from UCE data. Although the random local clock was also able to

identify an elevation of mitochondrial rates in nocticolids, it had a much lower marginal

likelihood that indicated poorer fit. This model produced a date estimate for crown

Nocticolidae that was similar to that inferred using the flexible local clock, but appeared to

have difficulty in resolving the location of an early shift in rates within the family.

The uncorrelated relaxed clocks in BEAST and MCMCtree were unable to

reconstruct the disparity in substitution rates between nocticolids and other cockroaches.

Notably, the three relaxed-clock models inferred dates for nocticolids that were broadly

similar to those obtained under a strict clock, but widely divergent from those inferred from

the UCE data. Furthermore, the inferred topologies varied across BEAST analyses,

illustrating that the impacts of clock-model misspecification extend beyond estimates of

substitution rates and divergence times. These results call into question the reliability of the

divergence times inferred using these widely used relaxed-clock models under conditions of

extreme rate variation. Our analysis of UCEs with an independent lognormal relaxed clock

produced younger estimates of divergence times within Nocticolidae compared with our

other analyses and those inferred previously (Djernæs et al. 2015; Wang et al. 2017; Li and

Huang 2020), which were likely to have been pushed back to accommodate the greater

numbers of substitutions on nocticolid branches. However, we also used slightly different

calibration methods between our analyses of UCE data and mtPCG data.

Although we focus on Nocticolidae as a case study, our results have broad

implications for estimating evolutionary timescales for groups with high among-lineage rate

variation. We find that the flexible local clock, random local clock, and autocorrelated clock

models produce the most concordant estimates of substitution rates and divergence times

when analysing data containing high among-lineage rate variation. The unifying feature of

the local clock models is their ability to allow a distinct rate, or distribution of rates, in

Nocticolidae. The reasons for the performance of the autocorrelated clock model are less

clear, but it could be that the model does not substantially penalize a single jump in rates at

the base of Nocticolidae. Nevertheless, the different estimates of rate variation within

Nocticolidae highlight the problems associated with each model. For example, the two

replicate analyses using a random local clock had identified a separate local clock for each

clade in Nocticolidae, but had grouped the rate of stem Nocticolidae with different clades of

Nocticolidae (A or B). This led to age estimates for crown Nocticolidae that differed between

the two analyses. Although we attempted to account for this uncertainty by combining the

posterior samples, poor Markov chain convergence appears to be a common issue for this

model when applied to large data sets (Dornburg et al. 2012, Drummond and Suchard 2010).

Applying an independent unimodal parametric distribution of branch rates to

Nocticolidae through the flexible local clock appeared to model rate variation between

Nocticolidae and the rest of the tree effectively. However, this restricted separation of rates

between the two clades within Nocticolidae, instead accounting for rate variation in the

family by estimating higher substitution rates on a few branches. This resulted in the highest

substitution rate on the stem branch of Nocticolidae B while comparable rates were present

within Nocticolidae A and B and on the stem branch of Nocticolidae A, likely leading to an

overestimation of divergence times in Nocticolidae B.

The random local clock is the only model that maintained a high substitution rate

throughout Nocticolidae B, which appears to be more biologically plausible, leading to the

youngest age estimate for crown Nocticolidae B. Even our analyses of UCEs, which had only

slightly greater root-to-tip distances in Nocticolidae compared with the rest of the tree,

inferred a slightly increased rate in stem Nocticolidae and stem Nocticolidae B compared

with the rest of the group, potentially leading to slight overestimation of divergence dates in

Nocticolidae B. Perplexingly, analysis of UCE data using the autocorrelated clock model

estimated substitution rates in Nocticolidae to be lower than across the rest of the tree,

highlighting that this model can misbehave when there is minimal among-lineage rate

variation. Our results suggest that molecular dating analyses should be carried out using data

sets with minimal rate variation among lineages, where posible. This could include data sets

such as the UCEs analysed here or, when dealing with whole-genome data, this could involve

filtering loci for clocklikeness (Jarvis et al. 2014; Smith et al. 2018). Selecting loci with

lower among-lineage rate variation has additional benefits for phylogenetic resolution

(Vankan et al. 2021).

Phylogenetic Inference When Rates Vary

Although not the focus of this study, we briefly address the potential effects of extreme

among-lineage rate variation on our phylogenetic inference. Our analysis of UCE data found

strong support for Corydiidae as the sister group to Nocticolidae, confirming the results of

previous analyses. The relationships among the major cockroach families are also congruent

to those inferred using transcriptomes (Evangelista et al. 2019), with the addition of the

family Anaplectidae as the sister group to the rest of Blattoidea. This placement conflicts

with the results of previous studies that supported a close relationship between Anaplectidae

and Lamproblattidae based on mitochondrial genomes (Bourguignon et al. 2018), or that

placed Lamproblattidae as the sister group to the rest of Blattoidea with Anaplectidae as the

sister group to Cryptocercidae+Isoptera and Tryonicidae based on mitochondrial and nuclear

markers (Djernæs and Murienne 2022).

514   The results of our UCE analysis also conflict with the evolutionary relationships

inferred from our mtPCG data set, which united Nocticolidae with Anaplectidae. The

instability of the position of Nocticolidae in our analyses of mitochondrial genomes could be

due to the long stem branches leading to the two clades of the family. In our analysis, there

was also a long stem branch leading to Anaplectidae because of minimal taxon sampling in

the family. Consequently, long-branch attraction could have occurred in the maximum-

likelihood analysis, even though this phenomenon is less pronounced when using model-

based phylogenetic methods (Swofford et al. 2001; Kolaczkowski and Thornton 2009; Kück

et al. 2012). Furthermore, our tests of model adequacy suggested a high risk of biased

inferences according to the consistency index, suggesting that among-lineage rate variation

could be responsible for the incongruence between our data sets. These results highlight that

extreme among-lineage rate variation can still have substantial impacts on model-based

methods of statistical phylogenetic inference.

Causes of Increased Substitution Rates in Nocticolidae

The causes of the increased mitochondrial and nuclear rates in Nocticolidae are yet to be

determined. Previous studies have suggested that the rate acceleration might be the result of

small population sizes and repeated bottlenecks associated with the colonization of caves, or

relaxed selection associated with the loss or reduction of characters no longer required in

subterranean environments (Bourguignon et al. 2018). Increased mitochondrial substitution

rates have been associated with the loss of flight in a number of insect lineages (Mitterboeck

and Adamowicz 2013). However, our results suggest that an increase in substitution rate

occurred early in the evolution of Nocticolidae when the ancestor had not yet become

restricted to the subterranean habitat, and presumably still possessed fully developed wings.

Similar patterns have been seen using a smaller number of nocticolids (Legendre et al. 2015;

Li and Huang 2020). Previous studies have placed the subfamily Latindiinae as the sister

lineage to Nocticolidae (Legendre et al. 2015; Wang et al. 2017; Li and Huang 2020; Liu et al.

2023), which also exhibits increased substitution rates despite not being associated with

subterranean habitats, suggesting that habitat type is unlikely to be the sole driver of

evolutionary rate acceleration.

One common factor across all studied lineages of Nocticolidae, including epigean and

subterranean taxa, is their reduced body size compared with other cockroaches. Small body

size is associated with higher substitution rates in a number of animals, presumably because

of its covariation with shorter generations, higher metabolic rate, and increased fecundity,

none of which has been studied in Nocticolidae (Bromham 2009; Thomas et al. 2010). In

cockroaches, there is a positive relationship between body size and clutch size, although this

might be offset by increased frequency of clutches in smaller-bodied species (Djernæs et al.

2020). Other groups of cockroaches with small body sizes have also shown greater

phylogenetic root-to-tip distances, including the subfamily Latindiinae, and a distantly related

genus of minute myrmecophilous cockroaches, *Attaphila* (Djernæs et al. 2020). The effects of

life-history traits in elevating substitution rates are more likely to be seen in mitochondrial

genomes, which have smaller effective population sizes than nuclear genomes.

We found evidence of increased substitution rates for nocticolids in the UCE data set,

although less pronounced than the increases found in mitochondrial genomes. The UCE loci

are likely to be under strong selective constraints because many are found in exonic regions

in insects (Zhang et al. 2019; Hellemans et al. 2022)(Zhang et al. 2019; Hellemans et al.

2022). Our finding of an increased rate in the nuclear DNA of nocticolids stands in contrast

with the results of a previous analysis of a transcriptome data set, which found broadly

similar root-to-tip distances among nocticolids and other taxa (Evangelista et al. 2019).

However, the transcriptome data set comprised the second codon sites from ~2000 protein-

coding genes conserved across a diverse group of insects, which are likely to be under strong

purifying selection.

Our results suggest that Nocticolidae is an old family with a crown age of ~115 Ma and most

recent common ancestor with Corydiidae in the Jurassic. This aligns with results from recent

molecular dating studies (Evangelista et al. 2019; Li and Huang 2020), and the existence of

~100 Myr old nocticolid fossils from Asia (Li and Huang 2020; Sendi et al. 2020). The

crown age of the family might be even older than inferred in this study, given that we did not

include nocticolid representatives from Africa and Madagascar.

The estimated timing of divergence between Nocticolidae A and B suggests that these

two lineages separated after the breakup of Pangaea (~200–150 Ma) (Seton et al. 2012). An

Asian origin for Nocticolidae A is suggested by the placement of Asian taxa and the nested

position of Australian mainland and Christmas Island taxa. Our date estimates suggest that

nocticolids arrived on the Australian mainland some time between 31 and 21 Ma, which

aligns with the collision of the Australian and Asian tectonic plates ~25 Ma. Around this time,

there was extensive floral and faunal exchange between Australia and South-East Asia

(Maekawa et al. 2003; Crayn et al. 2015). The estimated crown age of 42 Ma for

Nocticolidae B suggests its presence on the ancient Australia/New Guinea landmass prior to

its collision with the Asian plate to the north. However, because we believe our divergence

dates in Nocticolidae B could be slight overestimates, we cannot rule out the possibility of

dispersal into Australia around the same time as Nocticolidae A, especially considering the

close relationship inferred between *N. babindaensis* and a specimen from China (Wang et al.

2017). Furthermore, any biogeographic hypotheses should be considered tentative due to the

absence of African, Madagascan, and other Asian lineages in our data sets. Further analyses

involving additional taxa will be needed to test the biogeographic origins of Nocticolidae B

and of Nocticolidae.

Australia was largely covered in rainforest ~25–100 Ma (Byrne et al. 2011), which

would have favoured the movement of ancestral epigean nocticolids across the continent.

Periods of aridity from the Miocene onwards led to rainforest habitats receding to a few

refugia along the east coast of Australia (Byrne et al. 2011). These conditions might have

forced various ancestral nocticolid lineages to enter caves or other subterranean habitats, or to

develop an inquiline relationship with burrowing insects.

Nocticolid Systematics and Taxonomy

Our study is the first to include genetic data for a diverse range of Nocticolidae and

highlights the need for taxonomic revision in the family (Roth 1988). A recent morphological

analysis was unable to resolve relationships amongst the genera of Nocticolidae and

Corydiidae (Li and Huang 2020). By combining morphological data for all individuals of a

genus, the authors assumed the monophyly of each, despite widespread taxonomical

uncertainty. As with most troglomorphic taxa, regressive evolution is likely to have occurred

in parallel across Nocticolidae, owing to multiple independent transitions into caves.

Therefore, it is likely that the current taxonomy of the group, which is based on morphology,

does not accurately reflect true evolutionary relationships. Our analysis of UCE data yielded

an estimate of ~115 Ma for the crown age of *Nocticola*, making this genus very old and

potentially containing at least some of the other nocticolid genera. Our samples from

Christmas Island are likely to represent *Metanocticola christmasensis*, which would confirm

the non-monophyly of *Nocticola*, although we only sampled nymphs which are difficult to

identify (Roth 1999) (full discussion in Supplementary Material). Further genetic sampling of

other nocticolid genera is required to confirm that *Nocticola* is indeed polyphyletic.

Members of Nocticolidae A and B are likely to represent two distinct genera.

Representatives of the genus *Nocticola* have previously been split into two groups: the *uenoi-*

species group, in which the male terga are specialized; and the *simoni*-species group, in

which the male's terga are unspecialized (Roth 1988). This grouping does not align with the

phylogenetic relationships inferred here. Instead, tergal glands appear to have been derived

independently on multiple occasions, potentially during subterranean adaptation, as they are

not present in any epigean or inquiline taxa. Tergal specializations vary substantially in

Nocticola, with some taxa (e.g., *N. australiensis* and *N.* sp. (Carpentaria Cave)) possessing a

simple sclerotized indent on the fourth tergum, while others have complex structures

involving multiple tergal segments (e.g., *N. currani*, *N. ueoni*, and *N.* sp. (Kimberley 175_1)).

There are two morphological characters that align with the relationships inferred

using molecular data and are good candidate synapomorphies. One potential synapomorphy

of members of Nocticolidae A constitutes a group of large sclerotized spines on the ventral

surface of segments 4–6 of the female cerci (Roth 1988; Trotter et al. 2017). This has been

observed in all available females of Nocticolidae A (Supplementary Table S1), including

Nocticola sp. (Malaysia) and our Christmas Island samples, which suggests that the trait

could be ancestral in this clade (Mari Fujita, personal communication). Importantly, this

character was not observed in any of the females in Nocticolidae B. This character could

provide some guidance for the placement of species without genetic data (see Supplementary

Table S1). Unfortunately, no females have been sampled from most of the other nocticolid

genera, but females in *Speleoblatta* and *Alluaudellina* do not have this morphology (Chopard

1932; Vidlička et al. 2003, 2017). *Alluaudellina* is likely to be a synonym of *Nocticola*

(Chopard 1932; Roth 1988) and the lack of female cerci spinals suggests that African

nocticolids are not closely related to Nocticolidae A, although genetic analysis will be

required to confirm this.

Previous studies have found striking differences in male genitalia between closely

related species in *Nocticola* (Trotter et al. 2017). One genital character that aligns with our

molecular results is the shape of L3d (left phallomere 3 dorsal). In three representatives of

Nocticolidae B (*N.* sp. (Kimberley 175_1), *N. currani*, and *N. flabella*), L3d is long, curved,

and spear-like (Roth 1991; Trotter et al. 2017). Contrastingly, in five representatives of

Nocticolidae A (*N. cockingi*, *N. quartermainei*, *N.* sp. (Kimberley 107_1), *N. australiensis*,

and *N.* sp. (Glennie Tablelands)), L3d is rounder, complex, and shovel-like (Trotter et al.

2017). The spear-like L3d present in the African species *N. clavata*, *N. scytala*, *N. wliensis*,

and *A. cavernicola* (Chopard 1932; Andersen and Kjaerandsen 1995) appears to be similar to

the morphology seen in Nocticolidae B, although modern, high quality genitalia drawings (eg.

Lucañas and Lit 2016; Trotter et al. 2017; Lucañas and Maosheng 2023) are required to

confirm this.

It is also possible that unsampled nocticolids represent sister lineages to the two

studied here, potentially with a close relationship to the undescribed sample from an

unknown location which we found to be the sister lineage to the rest of Nocticolidae.

Unfortunately, the type species for the genus, *Nocticola simoni*, cannot be grouped into either

clade because of its basic description (Bolivar 1892). Therefore, re-examination and

generation of genomic data for the type taxon is required to identify whether Nocticolidae A

or B should be redescribed as *Nocticola*, and which of these should be described as a new

genus.

Conclusions

In performing the largest phylogenomic and molecular dating analysis of nocticolid

cockroaches, we have uncovered an extraordinary level of genetic diversity in the family.

Nocticolids have experienced remarkably high substitution rates compared with other

cockroaches, especially in mitochondrial protein-coding genes. This extreme lineage-specific

rate acceleration is likely to have misled phylogenetic inference using maximum-likelihood

and Bayesian methods. Our phylogenetic analysis of UCE loci has resolved the deep

relationships among the major cockroach families, in uniting Corydiidae with Nocticolidae

and placing Anaplectidae as the sister lineage to the rest of Blattoidea. Within Nocticolidae,

we identify two highly divergent clades that separated ~115 Ma despite both containing

representatives of the cryptic genus *Nocticola*, suggesting that the genus requires taxonomic

attention.

Our results suggest that some of the widely used relaxed-clock models are unable to

account for large disparities in substitution rates among lineages, with potentially negative

impacts on estimates of divergence times. In this regard, localized accelerations of rates can

be taken into account using models such as the flexible local clock. This model has not been

widely used, possibly because it requires that the number and phylogenetic placements of

local relaxed clocks be chosen a priori. However, unusual patterns of among-lineage rate

heterogeneity call for the use of more complex clock models. Modelling the variation in

molecular rates more accurately will allow more reliable reconstructions of evolutionary

timescales and phylogenetic relationships.

SUPPLEMENTARY MATERIAL

Data available from the Dryad Digital Repository: http://dx.doi.org/10.5061/dryad.[NNNN]

FUNDING

This project was funded by the Linnean Society of NSW and in part by a Discovery Project

from the Australian Research Council (DP220103265). The Yim Family Foundation

provided TGLK with generous financial support.

AUTHOR CONTRIBUTIONS

TGLK, JW, SYWH, and NL conceived and designed the study. Personal collections

were provided by JW and sample collection was undertaken by TGLK, JW, and NL. NJT and

JMM provided museum and environmental survey samples. SH and TB designed and

generated UCE data. SYWH provided input into phylogenetic and molecular dating analyses.

TGLK completed the wet lab work, analysed the molecular data, performed data analyses,

and drafted the manuscript with assistance from SYWH and NL. All authors approved the

manuscript prior to submission.

ACKNOWLEDGEMENTS

We thank Bruce Gray for help with collecting *Nocticola* specimens, Mari Fujita for

correspondence regarding the morphology of *Nocticola* sp. (Malaysia), Maosheng Foo

(LKCNHMS), Jiří Patoka, and Martin Bláha for providing specimens, and Kyle Ewart and

Yi-Kai Tea for guidance on data analyses. We also thank the late Fred Stone for his lifelong

contribution to furthering our understanding of subterranean fauna. The authors acknowledge

the Sydney Informatics Hub and the use of the University of Sydney's high-performance

computing cluster, Artemis. We thank an anonymous reviewer and the Editors for valuable

input   on   an   earlier   version   of   this   manuscript.

REFERENCES

Abascal F., Zardoya R., Telford M.J. 2010. TranslatorX: Multiple alignment of nucleotide

sequences guided by amino acid translations. Nucleic Acids Res. 38:7–13.

Aberer A.J., Kobert K., Stamatakis A. 2014. Exabayes: Massively parallel bayesian tree

inference for the whole-genome era. Mol. Biol. Evol. 31:2553–2556.

Andersen T., Kjaerandsen J. 1995. Three new species of Nocticola Bolivar from Ghana, West

Africa (Blattaria: Nocticolidae). J. African Zool. 109:377–385.

Asahina S. 1974. The Cavernicolous Cockroaches of the Ryukyu Islands. Mem. Natl. Sci.

Museum Tokyo. 7:145–156.

Bolivar I. 1892. Voyage de M. E. Simno aux iles Philippines (mars et avril 1890). Etudes sur

les Arthropodes cavernicoles de l'ile de Luzon. Orthopteres. Ann. la Société Entomol.

Fr. 61:29–34.

Bouckaert R., Vaughan T.G., Barido-Sottani J., Duchêne S., Fourment M., Gavryushkina A.,

Heled J., Jones G., Kühnert D., De Maio N., Matschiner M., Mendes F.K., Müller N.F.,

Ogilvie H.A., Du Plessis L., Popinga A., Rambaut A., Rasmussen D., Siveroni I.,

Suchard M.A., Wu C.H., Xie D., Zhang C., Stadler T., Drummond A.J. 2019. BEAST

2.5: An advanced software platform for Bayesian evolutionary analysis. PLoS Comput.

Biol. 15:1–28.

Bourguignon T., Tang Q., Ho S.Y.W., Juna F., Wang Z., Arab D.A., Cameron S.L., Walker

J., Rentz D., Evans T.A., Lo N. 2018. Transoceanic dispersal and plate tectonics shaped

global cockroach distributions: Evidence from mitochondrial phylogenomics. Mol. Biol.

Evol. 35:970–983.

Bromham L. 2009. Why do species vary in their rate of molecular evolution? Biol. Lett.

5:401–404.

Bromham L., Cowman P.F., Lanfear R. 2013. Parasitic plants have increased rates of

molecular evolution across all three genomes. BMC Evol. Biol. 13:126.

Bui Q.M., Schmidt H.A., Chernomor O., Schrempf D., Woodhams M.D., Von Haeseler A.,

Lanfear R. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic

Inference in the Genomic Era. Mol. Biol. Evol. 37:1530–1534.

Byrne M., Steane D.A., Joseph L., Yeates D.K., Jordan G.J., Crayn D., Aplin K., Cantrill D.J.,

Cook L.G., Crisp M.D., Keogh J.S., Melville J., Moritz C., Porch N., Sniderman J.M.K.,

Sunnucks P., Weston P.H. 2011. Decline of a biome: Evolution, contraction,

fragmentation, extinction and invasion of the Australian mesic zone biota. J. Biogeogr.

38:1635–1656.

Chernomor O., Von Haeseler A., Minh B.Q. 2016. Terrace Aware Data Structure for

Phylogenomic Inference from Supermatrices. Syst. Biol. 65:997–1008.

Chopard L. 1932. Un cas de microphtalmie liée à l'atrophie des ailes chez une blatte

cavernicole. Livre du Centen. la Société Entomol. Fr. 485:485–496.

Crayn D.M., Costion C., Harrington M.G. 2015. The Sahul-Sunda floristic exchange: Dated

molecular phylogenies document Cenozoic intercontinental dispersal dynamics. J.

Biogeogr. 42:11–24.

Crisp M.D., Hardy N.B., Cook L.G. 2014. Clock model makes a large difference to age

estimates of long-stemmed clades with no internal calibration: A test using Australian

grasstrees. BMC Evol. Biol. 14:1–17.

Dierckxsens N., Mardulyn P., Smits G. 2017. NOVOPlasty: De novo assembly of organelle

genomes from whole genome data. Nucleic Acids Res. 45:e18.

Djernæs M., Klass K.D., Eggleton P. 2015. Identifying possible sister groups of

Cryptocercidae+Isoptera: A combined molecular and morphological phylogeny of

Dictyoptera. Mol. Phylogenet. Evol. 84:284–303.

Djernæs M., Klass K.D., Picker M.D., Damgaard J. 2012. Phylogeny of cockroaches (Insecta,

Dictyoptera, Blattodea), with placement of aberrant taxa and exploration of out-group

sampling. Syst. Entomol. 37:65–83.

Djernæs M., Murienne J. 2022. Phylogeny of Blattoidea (Dictyoptera: Blattodea) with a

revised classification of Blattidae. Arthropod Syst. Phylogeny. 80:209–228.

Djernæs M., Varadínova Z.K., Kotyk M., Eulitz U., Klass K.-D. 2020. Phylogeny and Life

History Evolution of Blaberoidea (Blattodea). Arthropod Syst. Phylogeny. 78:29–67.

Donath A., Jühling F., Al-Arab M., Bernhart S.H., Reinhardt F., Stadler P.F., Middendorf M.,

Bernt M. 2019. Improved annotation of protein-coding genes boundaries in metazoan

mitochondrial genomes. Nucleic Acids Res. 47:10543–10552.

Dornburg A., Brandley M.C., McGowen M.R., Near T.J. 2012. Relaxed clocks and

inferences of heterogeneous patterns of nucleotide substitution and divergence time

estimates across whales and dolphins (Mammalia: Cetacea). Mol. Biol. Evol. 29:721–

736.

Drummond A.J., Ho S.Y.W., Phillips M.J., Rambaut A. 2006. Relaxed phylogenetics and

dating with confidence. PLoS Biol. 4:e88.

Drummond A.J., Suchard M.A. 2010. Bayesian random local clocks, or one rate to rule them

all. BMC Biol. 8:114.

Duchêne D.A., Duchêne S., Ho S.Y.W. 2018. PhyloMAd: Efficient assessment of

phylogenomic model adequacy. Bioinformatics. 34:2300–2301.

Duchêne D.A., Mather N., Van Der Wal C., Ho S.Y.W. 2022. Excluding Loci with

Substitution Saturation Improves Inferences from Phylogenomic Data. Syst. Biol.

71:676–689.

Duchêne D.A., Tong K.J., Foster C.S.P., Duchêne S., Lanfear R., Ho S.Y.W. 2020. Linking

branch lengths across sets of loci provides the highest statistical support for

phylogenetic inference. Mol. Biol. Evol. 37:1202–1210.

Duchêne S., Lanfear R., Ho S.Y.W. 2014. The impact of calibration and clock-model choice

on molecular estimates of divergence times. Mol. Phylogenet. Evol. 78:277–289.

Evangelista D.A., Wipfler B., Béthoux O., Donath A., Fujita M., Kohli M.K., Legendre F.,

Liu S., Machida R., Misof B., Peters R.S., Podsiadlowski L., Rust J., Schuette K.,

Tollenaar W., Ware J.L., Wappler T., Zhou X., Meusemann K., Simon S. 2019. An

integrative phylogenomic approach illuminates the evolutionary history of cockroaches

and termites (Blattodea). Proc. R. Soc. B. 286:20182076.

Faircloth B.C. 2016. PHYLUCE is a software package for the analysis of conserved genomic

loci. Bioinformatics. 32:786–788.

Fernando W. 1957. New species of insects from Ceylon (I). Ceylon J. Sci. (Biological Sci.

1:157–167.

Fisher A.A., Ji X., Nishimura A., Lemey P., Suchard M.A. 2021. Shrinkage-based random

local clocks with scalable inference. arXiv, 10.48550/arXiv.2105.07119.

Fourment M., Darling A.E. 2018. Local and relaxed clocks: The best of both worlds. PeerJ.

2018:e5140.

Gan H.M., Tan M.H., Lee Y.P., Schultz M.B., Horwitz P., Burnham Q., Austin C.M. 2018.

More evolution underground: Accelerated mitochondrial substitution rate in Australian

burrowing freshwater crayfishes (Decapoda: Parastacidae). Mol. Phylogenet. Evol.

118:88–98.

Hellemans S., Wang M., Hasegawa N., Šobotník J., Scheffrahn R.H., Bourguignon T. 2022.

Using ultraconserved elements to reconstruct the termite tree of life. Mol. Phylogenet.

Evol. 173:107520.

Ho S.Y.W. 2020. The molecular clock and evolutionary rates across the tree of life. In: Ho

S.Y.W., editor. The Molecular Evolutionary Clock: Theory and Practice. Springer,

Cham. p. 3–23.

Ho S.Y.W., Duchêne S. 2014. Molecular-clock methods for estimating evolutionary rates and

timescales. Mol. Ecol. 23:5947–5965.

Ho S.Y.W., Duchêne S., Duchêne D. 2015. Simulating and detecting autocorrelation of

molecular evolutionary rates among lineages. Mol. Ecol. Resour. 15:688–696.

Inward D., Beccaloni G., Eggleton P. 2007. Death of an order: A comprehensive molecular

phylogenetic study confirms that termites are eusocial cockroaches. Biol. Lett. 3:331–

335.

Jarvis E.D., Mirarab S., Aberer A.J., Li B., Houde P., Li C., Ho S.Y.W., Faircloth B.C.,

Nabholz B., Howard J.T., Suh A., Weber C.C., da Fonseca R.R., Li J., Zhang F., Li H.,

Zhou L., Narula N., Liu L., Ganapathy G., Boussau B., Bayzid M.S., Zavidovych V.,

Subramanian S., Gabaldón T., Capella-Gutiérrez S., Huerta-Cepas J., Rekepalli B.,

Munch K., Schierup M., Lindow B., Warren W.C., Ray D., Green R.E., Bruford M.W.,

Zhan X., Dixon A., Li S., Li N., Huang Y., Derryberry E.P., Bertelsen M.F., Sheldon

F.H., Brumfield R.T., Mello C. V, Lovell P. V, Wirthlin M., Schneider M.P.C.,

Prosdocimi F., Samaniego J.A., Velazquez A.M.V., Alfaro-Núñez A., Campos P.F.,

Petersen B., Sicheritz-Ponten T., Pas A., Bailey T., Scofield P., Bunce M., Lambert

D.M., Zhou Q., Perelman P., Driskell A.C., Shapiro B., Xiong Z., Zeng Y., Liu S., Li Z.,

Liu B., Wu K., Xiao J., Yinqi X., Zheng Q., Zhang Y., Yang H., Wang J., Smeds L.,

Rheindt F.E., Braun M., Fjeldsa J., Orlando L., Barker F.K., Jønsson K.A., Johnson W.,

Koepfli K.-P., O'Brien S., Haussler D., Ryder O.A., Rahbek C., Willerslev E., Graves

G.R., Glenn T.C., McCormack J., Burt D., Ellegren H., Alström P., Edwards S. V,

Stamatakis A., Mindell D.P., Cracraft J., Braun E.L., Warnow T., Jun W., Gilbert

M.T.P., Zhang G. 2014. Whole-genome analyses resolve early branches in the tree of

life of modern birds. Science. 346:1320–1331.

Kaltenpoth M., Corneli P.S., Dunn D.M., Weiss R.B., Strohm E., Seger J. 2012. Accelerated

evolution of mitochondrial but not nuclear genomes of hymenoptera: New evidence

from crabronid wasps. PLoS One. 7:e32826.

Kalyaanamoorthy S., Minh B.Q., Wong T.K.F., Von Haeseler A., Jermiin L.S. 2017.

ModelFinder: Fast model selection for accurate phylogenetic estimates. Nat. Methods.

14:587–589.

Karny H. 1924. Beiträge zur Malayischen Orthopteren fauna. Treubia. 5:12–19.

Katoh K., Toh H. 2008. Recent developments in the MAFFT multiple sequence alignment

program. Brief. Bioinform. 9:286–298.

Kolaczkowski B., Thornton J.W. 2004. Performance of maximum parsimony and likelihood

phylogenetics when evolution is heterogenous. Nature. 431:980–984.

Kolaczkowski B., Thornton J.W. 2009. Long-branch attraction bias and inconsistency in

bayesian phylogenetics. PLoS One. 4:e7891.

Kück P., Mayer C., Wägele J.W., Misof B. 2012. Long branch effects distort maximum

likelihood phylogenies in simulations despite selection of the correct model. PLoS One.

7:31–33.

Lanfear R., Calcott B., Ho S.Y.W., Guindon S. 2012. PartitionFinder: Combined selection of

partitioning schemes and substitution models for phylogenetic analyses. Mol. Biol. Evol.

29:1695–1701.

Legendre F., Nel A., Svenson G.J., Robillard T., Pellens R., Grandcolas P. 2015. Phylogeny

of dictyoptera: Dating the origin of cockroaches, praying mantises and termites with

molecular data and controlled fossil evidence. PLoS One. 10:1–27.

Legendre F., Whiting M.F., Bordereau C., Cancello E.M., Evans T.A., Grandcolas P. 2008.

The phylogeny of termites (Dictyoptera: Isoptera) based on mitochondrial and nuclear

markers: Implications for the evolution of the worker and pseudergate castes, and

foraging behaviors. Mol. Phylogenet. Evol. 48:615–627.

Lemaire B., Huysmans S., Smets E., Merckx V. 2011. Rate accelerations in nuclear 18S

rDNA of mycoheterotrophic and parasitic angiosperms. J. Plant Res. 124:561–576.

Lepage T., Bryant D., Philippe H., Lartillot N. 2007. A general comparison of relaxed

molecular clock models. Mol. Biol. Evol. 24:2669–2680.

Lepage T., Lawi S., Tupper P., Bryant D. 2006. Continuous and tractable models for the

variation of evolutionary rates. Math. Biosci. 199:216–233.

Li X.R., Huang D. 2020. A new mid-Cretaceous cockroach of stem Nocticolidae and

reestimating the age of Corydioidea (Dictyoptera: Blattodea). Cretac. Res. 106:104202.

Linder M., Britton T., Sennblad B. 2011. Evaluation of bayesian models of substitution rate

evolution-parental guidance versus mutual independence. Syst. Biol. 60:329–342.

Liu J., Zhang J., Han W., Wang Y., He S., Wang Z. 2023. Advances in the understanding of

Blattodea evolution: Insights from phylotranscriptomics and spermathecae. Mol.

Phylogenet. Evol. 182:107753.

Lo N., Beninati T., Stone F., Walker J., Sacchi L. 2007. Cockroaches that lack

Blattabacterium endosymbionts: The phylogenetically divergent genus Nocticola. Biol.

Lett. 3:327–330.

Lucañas C.C., Lit I.L. 2016. Cockroaches (Insecta, Blattodea) from caves of Polillo Island

(Philippines), with description of a new species. Subterr. Biol. 19:51–64.

Lucañas C.C., Maosheng F. 2023. A new macropterous Nocticola Bolivar , 1892

(Blattodea□ : Nocticolidae) from Singapore. J. Asia. Pac. Entomol. 26:102062.

Maekawa K., Lo N., Rose H.A., Matsumoto T. 2003. The evolution of soil-burrowing

cockroaches (Blattaria: Blaberidae) from wood-burrowing ancestors following an

invasion of the latter from Asia into Australia. Proc. R. Soc. B Biol. Sci. 270:1301–1307.

Maturana Russel P., Brewer B.J., Klaere S., Bouckaert R.R. 2019. Model Selection and

Parameter Inference in Phylogenetics Using Nested Sampling. Syst. Biol. 68:219–233.

Mitterboeck F.T., Adamowicz S.J. 2013. Flight Loss Linked to Faster Molecular Evolution in

Insects. Proc. R. Soc. B. 280.

Nurk S., Meleshko D., Korobeynikov A., Pevzner P.A. 2017. metaSPAdes: a new versatile

metagenomic assembler. Genome Res. 27:824–834.

Rannala B., Yang Z. 2007. Inferring speciation tunes under an episodic molecular clock. Syst.

Biol. 56:453–466.

dos Reis M., Gunnell G.F., Barba-Montoya J., Wilkins A., Yang Z., Yoder A.D. 2018. Using

phylogenomic data to explore the effects of relaxed clocks and calibration strategies on

divergence time estimation: Primates as a test case. Syst. Biol. 67:594–615.

dos Reis M., Yang Z. 2011. Approximate likelihood calculation on a phylogeny for Bayesian

Estimation of Divergence Times. Mol. Biol. Evol. 28:2161–2172.

Roch S., Nute M., Warnow T. 2019. Long-Branch Attraction in Species Tree Estimation:

Inconsistency of Partitioned Likelihood and Topology-Based Summary Methods. Syst.

Biol. 68:281–297.

Roth L.M. 1988. Some cavernicolous and epigean cockroaches with six new species, and a

discussion of the Nocticolidae (Dictyoptera: Blattaria). Rev. suisse Zool. 95:297–321.

Roth L.M. 1991. A new cave-dwelling cockroach from Western Australia (Blattaria:

Nocticolidae). Rec. West. Aust. Museum. 15:17–21.

Roth L.M. 1999. New cockroach species, redescriptions, and records, mostly from Australia,

and a description of Metanocticola christmasensis gen. nov., sp. nov., from Christmas

Island (Blattaria). Rec. West. Aust. Museum. 19:327–364.

Roth L.M., McGavin G.C. 1994. Two new species of nocticolidae (Dictyoptera: Blattaria)

and a rediagnosis of the cavernicolous genus spelaeoblatta bolivar. J. Nat. Hist.

28:1319–1326.

Sanderson M.J. 2002. Estimating absolute rates of molecular evolution and divergence times:

A penalized likelihood approach. Mol. Biol. Evol. 19:101–109.

Sendi H., Vršanský P., Podstrelená L., Hinkelman J., Kúdelová T., Kúdela M., Vidlička Ľ.,

Ren X., Quicke D.L.J.J., Hinkelman J., Kúdelová T., Kúdela M., Vidlička Ľ., Ren X.,

Quicke D.L.J.J. 2020. Nocticolid cockroaches are the only known dinosaur age cave

survivors. Gondwana Res. 82:288–298.

Seton M., Müller R.D., Zahirovic S., Gaina C., Torsvik T., Shephard G., Talsma A., Gurnis

M., Turner M., Maus S., Chandler M. 2012. Global continental and ocean basin

reconstructions since 200Ma. Earth-Science Rev. 113:212–270.

Smith S.A., Brown J.W., Walker J.F. 2018. So many genes, so little time: A practical

approach to divergence-time estimation in the genomic era. PLoS One. 13:e0197433.

Susko E. 2015. Bayesian long branch attraction bias and corrections. Syst. Biol. 64:243–255.

Swofford D.L., Waddell P.J., Huelsenbeck J.P., Foster P.G., Lewis P.O., Rogers J.S.,

Swofford D.L., Waddell P.J., Huelsenbeck J.P., Foster P.G., Lewis P., Rogers J.S. 2001.

Bias in Phylogenetic Estimation and Its Relevance to the Choice between Parsimony

and Likelihood Methods. Syst. Biol. 50:525–539.

Thomas J.A., Welch J.J., Lanfear R., Bromham L. 2010. A generation time effect on the rate

of molecular evolution in invertebrates. Mol. Biol. Evol. 27:1173–1180.

Thorne J.L., Kishino H., Painter I.S. 1998. Estimating the rate of evolution of the rate of

molecular evolution. Mol. Biol. Evol. 15:1647–1657.

Trotter A.J., McRae J.M., Main D.C., Finston T.L. 2017. Speciation in fractured rock

landforms: Towards understanding the diversity of subterranean cockroaches

(Dictyoptera: Nocticolidae: Nocticola) in Western Australia. Zootaxa. 4250:143–170.

Vankan M., Ho S.Y.W., Duchêne D.A. 2021. Evolutionary Rate Variation among Lineages

in Gene Trees has a Negative Impact on Species-Tree Inference. Syst. Biol. 71:490–500.

Vidlička L., Vršanský P., Kúdelová T., Kúdela M., Deharveng L., Hain M. 2017. New genus

934 and species of cavernicolous cockroach (Blattaria, Nocticolidae) from Vietnam. Zootaxa.

935 4232:361–375.

936 Vidlička Ľ., Vršanský P., Shcherbakov D.E. 2003. Two new troglobitic cockroach species of

937 the genus spelaeoblatta (blattaria: Nocticolidae) from North Thailand. J. Nat. Hist.

938 37:107–114.

939 Wang Z., Shi Y., Qiu Z., Che Y., Lo N. 2017. Reconstructing the phylogeny of Blattodea:

940 Robust support for interfamilial relationships and major clades. Sci. Rep. 7:3903.

941 Yang Z. 2007. PAML 4: Phylogenetic analysis by maximum likelihood. Mol. Biol. Evol.

942 24:1586–1591.

943 Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: Polynomial time species

944 tree reconstruction from partially resolved gene trees. BMC Bioinformatics. 19:15–30.

945 Zhang Y.M., Williams J.L., Lucky A. 2019. Understanding UCEs: A Comprehensive Primer

946 on Using Ultraconserved Elements for Arthropod Phylogenomics. Insect Syst. Divers.

947 3:1–12.

FIGURE CAPTIONS

FIGURE CAPTIONS

FIGURE 1. Evolutionary timescale of Blattodea, with Mantodea and Phasmatodea as

outgroup taxa. Divergence times were estimated based on UCE loci, using a Bayesian

phylogenomic approach with an uncorrelated relaxed clock in MCMCTree. The fixed tree

topology used in this analysis was inferred using maximum-likelihood and Bayesian analyses

of UCE loci. Node support from these analyses is represented by squares: outline colours

indicate values of the SH-like approximate likelihood-ratio test (SH-alrt), fill colours indicate

ultrafast bootstrap (UFb) values. Nodes with SH-alrt and UFb > 99 are not labelled. Posterior

probabilities were 1.0 for all nodes, with the exception of the node labelled with a star, which

had a different topology. Insert shows the diverse morphology present in Nocticolidae: a)

female *Nocticola australiensis* (Royal Arch Cave) with highly reduced eyes (photograph by

Braxton Jones); b) blind female *Nocticola baumi* with elongated appendages (photograph by

Martin Bláha); and c) male *Nocticola pheromosa* with fully developed eyes and wings

(photograph by Maimon Hussin).

FIGURE 2. Phylogeny of Nocticolidae. Maximum-likelihood tree estimated using

concatenated UCE loci, second codon sites of 13 mitochondrial protein-coding genes, and

mitochondrial *16S*. There are two major clades in Nocticolidae, A (numbers in black) and B

(numbers in grey). Node support is represented by squares: outline colours indicate values of

the SH-like approximate likelihood-ratio test (SH-alrt), fill colours indicate ultrafast bootstrap

(UFb) values. Nodes with SH-alrt and UFb = 100 are not labelled. Colours represent the

geographical location of taxa from the Australian mainland (blue), South-East Asia (green),

Christmas Island (orange), and New Guinea (purple).

FIGURE 3. Accelerated mitochondrial substitution rates in Nocticolidae compared with other

families in Blattodea (cockroaches). (a) Maximum-likelihood tree of Blattodea inferred using

the second codon sites of the 13 mitochondrial protein-coding genes. (b) Phylogenetic root-

to-tip distances for the concatenated mitochondrial protein-coding genes and the

concatenated UCE loci. Root-to-tip distances were calculated for each of the major cockroach

families, with Nocticolidae split into its two major clades, A and B. Root-to-tip distances

were normalized by dividing the mean for each family by the mean root-to-tip distance across

all taxa.

FIGURE 4. Estimates of divergence times and substitution rates using a range of molecular-

clock models. (a) Dates for crown Nocticolidae, crown Nocticolidae A, and crown

Nocticolidae B estimated from mitochondrial protein-coding genes using a strict clock

(strict), uncorrelated lognormal relaxed clock (UCLN), uncorrelated exponential relaxed

clock (UCEXP), random local clock (RLC), flexible local clock (FLC), autocorrelated

relaxed clock (AC), and independent lognormal relaxed clock (ILN). Date estimates for the

same nodes using our UCE data set are also presented for comparison using an autocorrelated

relaxed clock (UCE-AC) and independent lognormal relaxed clock (UCE-ILN). Circles

indicate posterior means and bars indicate 95% credibility intervals. b) Estimates of

mitochondrial substitution rates in Nocticolidae compared with other families in Blattodea

(cockroaches). Violin plots represent the mean substitution rate (substitutions per site per

Myr) for all terminal and internal branches within Nocticolidae (including stem) and outside

Nocticolidae (non-Nocticolidae). Rates were estimated using concatenated mitochondrial

protein-coding genes.