APPROXIMATIONS TO THE SOLUTION OF THE KUSHNER-STRATONOVICH EQUATION FOR THE STOCHASTIC CHEMOSTAT[*]

JOSÉ AUGUSTO FONTENELE MAGALHÃES[†], MUHAMMAD FUADY EMZIR[‡], AND FRANCESCO CORONA[§]

Abstract. In order to characterise the dynamics of a biochemical system such as the chemostat, we consider a differential description of the evolution of its state under environmental fluctuations. We present solutions to the filtering problem for a chemostat subjected to geometric Brownian motion. Under this modelling assumption, our best knowledge about the state of the system is given by its distribution in time, given the distribution of the initial state. Such a function solves a deterministic partial differential equation, the Kolmogorov forward equation. In this paper, however, we refine our knowledge about the state of the chemostat when additional information about the system is available in the form of measurements. More formally, we are interested in obtaining the distribution of the state conditional on measurements as the solution to a non-linear stochastic partial integral differential equation, the Kushner-Stratonovich equation. For the chemostat, this solution is not available in closed form, and it must be approximated. We present approximations to the solution to the Kushner-Stratonovich equation based on methods for partial differential equations. We compare the solution with a linearisation method and with a classical sequential Monte Carlo method known as the bootstrap particle filter.

Key words. stochastic chemostat; state estimation, non-linear filtering, Fokker-Planck equation, Kushner-Stratonovich equation

MSC codes. 92D25, 60G35, 60H15, 62M20, 65C05

1. Introduction. The aim of filtering is to estimate the current state of an evolving dynamical system, customarily modelled by a stochastic process, denoted by X and called the signal process. The signal process cannot be measured directly, but only via a related process Y, termed the observation process. The aim is to compute π_t, the conditional distribution of the signal X_t, at the current time t given the observation data accumulated up to this time. More precisely, we compute the conditional distribution of the signal X_t given the observation filtration $\mathcal{F}_t^Y = \sigma\{Y_s : s \leq t\}$. It is possible to formalise this by defining a stochastic process $\pi = (\pi_t, t \geq 0)$ describing this conditional distribution. Such process is the solution to a non-linear stochastic partial integral differential equation, the Kushner-Stratonovich equation. The exact form of the process π can be derived for a specific class of problems, for instance, those with a linearly evolving observation process [3, 21]. Even in this scenario, however, the signal is only allowed to have a non-linear drift under some restrictive conditions (see Beneš condition [3]). In general, π is an infinite-dimensional stochastic process and finding it in closed-form is a non-trivial task due to the inherent non-linearities of the system. In this case, the solution must be approximated.

Here, our goal is to approximate the conditional distribution π_t for a process performed in a chemostat in order to learn more about the system. The chemostat was the major player in the use of continuous culture systems to characterise certain

kinds of microbes in the mid-20th century, prior to when molecular biology and thence genomics took centre stage instead [6, 17]. Being a realistic and thoroughly tested continuous culture system, it has regained some interest in this post-genomic era. For instance, chemostat-grown cultures lead to reproducible and homogeneous results, which are necessary to identify the roles genes play in the biology of organisms [18].

We focus on estimating the state of a chemostat to understand its operation and to devise strategies for optimal operation. Here, we consider two substrate-controlled growth models: a basic model (Monod kinetics) and a more complex model, where an excessive level of the substrate can inhibit the growth (Haldane kinetics). We consider a differential description of the substrate and biomass concentrations in a chemostat under environmental fluctuations [19]; in particular, chemostats subjected to geometric Brownian motion. Our best knowledge about the state of the system is given by its conditional distribution in time, given the distribution of the initial state. Such a function solves a deterministic partial differential equation (PDE), the Kolmogorov forward equation. Our focus is to refine our knowledge about the state of the chemostat when additional information about the system is available in the form of measurements. To describe how measurements and states are related, we augment the dynamics with an observation model in which the measurements are also subjected to Brownian motion. When measurements are collected, our estimate of the state is given by the conditional distribution π_t, now given the distribution of the initial state and the collected measurements.

For non-linear systems, the classical method for finding π_t is the extended Kalman filter (EKF), which linearises certain non-linearities in the model so that the traditional Kalman-Bucy filter (KBF, [21]) can be applied. A less common approach consists of solving an associated stochastic partial differential equation (SPDE), the Zakai equation. Its solution is an unnormalised density function, given the distribution of the initial state and the collected measurements. In this approach, the solution is also unavailable in closed form for our system, and it must be approximated numerically [2]. Despite being an SPDE, classical PDEs methods can still be applied. These methods are successful in low dimensions of the state space but cannot be applied to high-dimensional problems. This is because the number of finite elements required for an accurate approximation goes up exponentially with the dimension of the signal.

Among other numerical methods for solving the filtering problem, particle filters (PFs, also known as sequential Monte Carlo methods) are among the most popular ones. In this approximation, particles (also called samples) firstly evolve according to a desirable transition kernel and then, they interact selectively in accordance with their likelihood with respect to the measurements.

In the context of the stochastic chemostat, the existing state-of-the-art comprises approximations to the first two moments of the distribution π_t [16, 31]. We propose to advance beyond that by presenting results on the approximation to the entire distribution π_t, from which higher-moments could be derived. We focus on the approximation with methods for PDEs. In this work, the available observations are a transformed and noisy version of the signal. In particular, we observe the flow rate of outgoing biogas produced inside the chemostat, and not the biomass and substrate concentrations directly. We present alternative scenarios in the supplementary material, in which noisy observations of biomass and/or substrate concentrations are available. After presenting how the solution to π_t looks like, we evaluate the method for PDEs by using the Hellinger distance. More specifically, we quantify the similarity between the resulting approximation and the solution approximated via sequential Monte Carlo. We also present how the EKF performs under the same scenario.

This paper is structured as follows: we begin Section 2 by introducing some basic
notation and the equations governing the dynamics of a chemostat. The rest of the
section is devoted to the exposition of two scenarios, one with Monod kinetics and
the other with Haldane kinetics. In Section 3, we present a detailed description of
the stochastic chemostat; in particular, our goal is to showcase the prior probabilistic
knowledge about the chemostat given some initial condition of the system. In Sec-
tion 4, we have additional information in the form of measurements. In Sections 5
and 6, we present the filtering equations; we explore the problem when observations
are corrupted by Gaussian noise, we introduce some approaches used to solve the
filtering problem and we conclude the paper by comparing them.

2. Chemostat equations. The chemostat is a bioreactor in which one or more
populations of microorganisms grow in a nutrient medium consisting of a cocktail
of various molecules. Here, we call biomass the set of growing microorganisms and
substrate the cocktail of various molecules. Let b denote the concentration of biomass
and s the concentration of substrate. The following reaction takes place to produce
one unit of biomass and γ units of methane (biogas) from κ units of substrate, with
reaction rate $\rho(\cdot)$:

(2.1) $$\kappa s \xrightarrow{\rho(\cdot)} b + \gamma CH_4.$$

Let $F_{in}(t)$ denote the amount of media flowing into a vessel of volume $V(t)$ per
hour, and $F_{out}(t)$ denote the outgoing amount departing from the same vessel per
hour. Here, the focus is on bioreactor processes performed at continuous mode, where
at a specific time t, $F_{in}(t) = F_{out}(t) = F(t)$ and $V(t) = V$. Consequently, the dilution
rate is given by $D(t) = F(t)/V$. The ordinary differential model for the growth of a
single species in such a chemostat is given by

(2.2) $$\frac{db(t)}{dt} = (\mu(\cdot) - D(t))b(t) + D(t)b_{in}(t), \frac{ds(t)}{dt} = D(t)(s_{in}(t) - s(t)) - \kappa\mu(\cdot)b(t),$$

with initial concentrations b_0 and s_0 and incoming concentrations $b_{in}(t)$ and $s_{in}(t)$.
Here, μ is the "specific growth velocity", a function which models the growth of an
organism population and which guarantees that the reaction rate is zero in the absence
of biomass. See Subsection SM1.1 of the supplementary material for the derivation
from a mass balance equation.

In this work, we consider the case of no incoming flow of biomass, thus $b_{in}(t) =$
0, $t \geq 0$. We work with the minimal model [17], where we assume that (i) $\mu(\cdot)$ is a
function of the substrate only, with $\mu(s) > 0$ for $s > 0$ and $\mu(0) = 0$, and that (ii)
the yield coefficient κ is constant. These assumptions are to be interpreted in the
following way: (i) the velocity - or kinetics - of the reaction in Eq. (2.1) does not
change as the biomass concentration changes and (ii) temperature and pressure are
kept constant, thus κ is also constant.

Now, let $x(t)$ denote the state of the chemostat at instant t, that is $x(t) =$
$(b(t), s(t))$. We begin with an ordinary differential equation (ODE) on the signal $x(t)$
to describe the dynamics of the minimal system:

(2.3) $$d\underbrace{\begin{bmatrix} b(t) \\ s(t) \end{bmatrix}}_{x(t)} = \underbrace{\begin{bmatrix} (\mu(s(t)) - D(t))b(t) \\ D(t)(s_{in}(t) - s(t)) - \kappa\mu(s(t))b(t) \end{bmatrix}}_{f(t,x(t),u(t)|\theta)} dt, \qquad \underbrace{\begin{bmatrix} b(0) \\ s(0) \end{bmatrix}}_{x(0)} = \underbrace{\begin{bmatrix} b_0 \\ s_0 \end{bmatrix}}_{x_0},$$

where f is a function parametrised by a vector θ of parameters, and $u(t)$ is input at
instant t, which for the minimal model refers to $u(t) = (D(t), s_{in}(t))$. If the state of

the system has a known starting position, say point x_0 at $t = 0$, and $u(t)$ is given, the
function $x(t)$, also called a solution to Eq. (2.3), is then completely determined in the
future, i.e. we know $x(t)$ for all $t > 0$.
In this paper, we consider chemostat cultures with a single growth-limiting sub-
strate. We show two cases for which different types of organisms are being cultivated,
each with its own growth function.
Case 1 - Monotonic growth μ - Monod type
This first case consists of the Monod relation of growth [28], which assumes that
the growth rate is zero when there is no substrate and tends to an upper limit when
the substrate is in great excess:

(2.4)
$$\mu(s) = \frac{\mu_{max}s}{K_s + s}.$$

Here, μ depends on the maximum growth μ_{max} and the half-saturation coefficient K_s.
Case 2 - Non-monotonic growth μ - Haldane type
Some experiments indicate that excessive substrate may inhibit the growth of a
microbial population in the biomass [1]. To consider such scenario, this case works
with the following growth function:

(2.5)
$$\mu(s) = \frac{\mu_0 s}{K_s + s + I_s s^2}.$$

152 Here, μ depends on the growth parameter μ_0, half-saturation coefficient K_s and in-
153 hibitor coefficient I_s.
154   Let us now look at the solution to Eq. (2.3), given the parameters from Table 1.

(●) stable steady states (●) unstable steady state

Fig. 1: Left: trajectories from the deterministic model with growth function of the Monod type. Right:
trajectories from the deterministic model with growth function of the Haldane type.

Consider equilibrium pairs (b_e, s_e) for which the right-hand side of Eq. (2.3) is
zero. Case 1 has the equilibrium point $E_1^{(1)} = (9.966, 0.345)$. Furthermore, $E_1^{(1)}$ is a
locally exponentially stable equilibrium [17, Section 2.1.2]. Case 2 has a trivial equilib-
rium point $E_2^{(1)} = (0, s_{in})$. In addition, it has interior equilibria $E_2^{(2)} = (0.859, 1.141)$
and $E_2^{(3)} = (1.912, 0.088)$. It is possible to verify that one of the two interior equilibria
is a stable nodal point and the other is a saddle point [17, Section 2.1.3].

3. Continuous-time dynamic models.

Table 1: Model parameters and inputs for Eq. (2.3).

	Model parameters		Inputs	
	Specific growth	κ	$D(t) = D\,(V = 100\text{L})$ (h^{-1})	$s_{in}(t) = s_{in}$ (mgL^{-1})
Case 1	$\frac{0.3s}{10+s}$	10	0.01	100.0
Case 2	$\frac{5s}{0.5+s+5s^2}$	1	0.7	2.0

3.1. Stochastic differential equations.

3.1.1. Definition. Now let us consider the probability space $(\Omega, \mathcal{F}, \mathbb{P})$. A solution $x = (x(t), t \geq 0)$ to an ODE becomes a stochastic process $X = (X_t, t \geq 0)$ the moment we introduce some randomness by (i) randomizing the initial condition and (ii) introducing a stochastic driving process, e.g. Brownian motion. This procedure results in a class of differential equations known as stochastic differential equations (SDEs), whose solutions involve ordinary and stochastic integrals.

We consider integration with respect to Brownian motion and its properties only. Here, the differential equations are called Itô stochastic differential equations, and the stochastic integrals are called Itô integrals.

DEFINITION 3.1. *A stochastic process $B = (B_t, t \geq 0)$ is called a d-dimensional (standard) Brownian motion if the following conditions are satisfied:*

1. *It starts at zero, i.e. $B_0 = 0$ almost surely;*
2. *$B_t(\omega)_{\omega \in \Omega}$ has independent increments, so that $\forall v \geq 0$, the random variable $B_{t+v} - B_t$ is independent of the σ-algebra generated by $(B_s, 0 \leq s \leq t)$;*
3. *For all $0 \leq s < t$, the random variable $B_t - B_s$ is normally distributed with mean 0 and variance $(t - s)$, i.e. $B_t - B_s \sim \mathcal{N}(0, (t-s) \cdot I_{d \times d})$, where $I_{d \times d}$ stands for identity matrix of size d;*
4. *It has almost surely continuous sample paths.*

If we let the Brownian motion $B^x = (B_t^x, t \leq 0)$ drive the dynamics seen in Eq. (2.3), the following SDE emerges:

$$(3.1) \qquad dX_t = f(t, X_t, u(t)|\theta)dt + g(t, X_t, u(t)|\theta)dB_t^x, \quad X_0 = \xi,$$

where $\xi = \xi(\omega)_{\omega \in \Omega}$ is a random variable on Ω, and the deterministic functions f and g are the coefficients of the equation. For notation simplicity, we will omit the reference to the vector θ of parameters. We assume a constant input $u(t)$ and autonomous functions f and g, such that Eq. (3.1) can be rewritten as

$$(3.2a) \qquad dX_t = f(X_t)dt + g(X_t)dB_t^x, X_0 = \xi,$$

or equivalently in coordinate form

$$(3.2b) \qquad dX_t^{d_x} = f_{d_x}(X_t)dt + \sum_{m=1}^{M} g_{d_x m}(X_t)dB_t^{x,m}, \quad X_0^{d_x} = \xi^{d_x}, \, d_x = 1, \ldots, D_x,$$

where $B^x = (B^{x,1}, \ldots, B^{x,M})$ is an M-dimensional standard Brownian motion ($M \geq 1$). Here, the state process and the coefficients of the equation have the mapping

$$X : [0, \infty) \times \Omega \to \mathbb{R}^{D_x}, \quad f : \mathbb{R}^{D_x} \to \mathbb{R}^{D_x}, \quad g : \mathbb{R}^{D_x} \to \mathbb{R}^{D_x} \times \mathbb{R}^M.$$

The SDEs in (3.2) are equivalent to the following stochastic integral equation:

(3.3) $$X_t = X_0 + \int_0^t f(X_s)ds + \int_0^t g(X_s)dB_s^x, \quad t \geq 0,$$

where the first integral on the right-hand side is a Riemann integral, and the second
one is an Itô integral. Here, f and g satisfy the Lipschitz condition for existence and
uniqueness of solutions. In order to demonstrate challenges concerning SDE existence
and uniqueness, we now discuss what we mean by solutions of Eq. (3.2) and Eq. (3.3).

3.1.2. Existence and uniqueness. The goal of this section is to find conditions
on the coefficients f and g which guarantee the existence and uniqueness of solutions.
To build a notion of time into our probability space, we specify which sub-σ-
algebra of information in $\mathcal{F}$ can be gathered by time t. We consider the increasing
sequence $\{\mathcal{F}_t\}$ of σ-algebras of $\mathcal{F}$ for which for $t, s \in \mathbb{R}_+$ with $t \leq s$, we have $\mathcal{F}_t \subseteq$
$\mathcal{F}_s \subseteq \mathcal{F}$. The quadruple $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0,\infty)}, \mathbb{P})$ forms our filtered probability space, on
which we can now define an M-dimensional $\mathcal{F}_t$-Brownian motion B_t^x.
The solution of an SDE is usually described in the sense of strong or weak solu-
tions. Firstly, we proceed to detail what we mean by *strong solutions*.

DEFINITION 3.2 (Strong solutions [10, Section 10.4]). *Let $B^x = (B_t^x, t \geq 0)$ be a*
Brownian motion and ξ be a random variable on $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0,\infty)}, \mathbb{P})$ independent of
B^x. A stochastic process X is a strong solution of a stochastic differential equation if
• *X is adapted to the filtration $\{\mathcal{F}_t\}$;*
• *$\mathbb{P}$-almost surely $X_0 = \xi$;*
• *$\mathbb{P}$-almost surely $\int_0^t |f_{d_x}(X_s)| + (g_{dx,m}(X_s))^2 \, ds < \infty$ holds for all $t \in [0,\infty)$,*
$d_x = 1, \dots, D_x$ and $m = 1, \dots, M$;
• *$\mathbb{P}$-almost surely X solves Eq. (3.2).*

As usual, we assume that the sigma-algebra $\mathcal{F}_t$ is completed with all the $\mathbb{P}$-null
sets in Ω [10, Section 10.4]. This definition means that if we are given a probability
space which carries B^x and ξ, the solution X must be adapted to the generated
filtration $\{\mathcal{F}_t\}$; that is, the path $\{X_t\}_{t \geq 0}$ is completely characterized by the path of
the underlying driving process (Brownian motion) and the initial condition.
The condition of being adapted to the generated filtration is sometimes too strict
(the famous example of Tanaka's equation, found in, for example, [34]). One can find
pairs (X, B^x) which are solution to the SDE (3.2) instead of simply seeking X. Such
pairs are called *weak solutions*. In what follows, we consider mostly the case of strong
solutions of Itô stochastic differential equations.

THEOREM 3.3 (Existence and uniqueness [27, Theorem 2.2]). *Suppose that ξ is a*
random variable on $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0,\infty)}, \mathbb{P})$, for which ξ is independent of the Brownian
motion B^x also defined on this space. Suppose that
• *$\xi \in \mathcal{L}^2(\mathbb{P})$, i.e. ξ is square-integrable under the probability measure $\mathbb{P}$;*
• *f, g satisfy the Lipschitz continuous and linear growth conditions in x.*
Then there exists a solution $X = (X_t, t \geq 0)$ to the associated stochastic differential
equation (3.2) satisfying the initial condition ξ. Moreover, the solution is unique and
satisfies $X \in \mathcal{L}^2(\mathbb{T} \times \mathbb{P})$, where $\mathbb{T}$ is the Lebesgue measure on $[0,\infty)$.

EXAMPLE 3.4 (Geometric Brownian motion). *Consider the unidimensional case,*
for which $X_t : [0,\infty) \times \Omega \mapsto \mathbb{R}$. Let $f(X_t) = 0$ and $g(X_t) = \omega_x X_t$, where $\omega_x > 0$:

(3.4) $$dX_t = \omega_x X_t dB_t^x,$$

with initial condition $X_0 = \xi \equiv 1$. Now define a function $V(x) \overset{\Delta}{=} \ln(x)$. We apply Itô's formula [20] on the process $V(X)$, resulting in

$$d(\ln(X_t)) = \omega_x dB_t^x - \frac{1}{2}\omega_x^2 t, \qquad t \geq 0.$$

Since $B_0 = 0$, we further get $X_t = X_0 \exp\left(\omega_x B_t^x - \frac{1}{2}\omega_x^2 t\right)$. For this SDE, both existence and uniqueness properties hold. Moreover, if the initial condition is constrained to positive real numbers, all the values for paths $\{X_t\}_{t \geq 0}$ will follow the same property.

Now assume that the concentration of an element (e.g. biomass) changes within a time period by a random amount. Similarly to what happens in Example 3.4, assume that this random amount is a function of the current concentration itself. To give an idea of the driving process added to Eq. (2.3), we consider an SDE with a driving process similar to the geometric Brownian motion. We will see that the conditions for both existence and uniqueness properties have to be slightly modified to prove existence and uniqueness.

The chemostat model considered in [19] has the following dynamic behaviour:

$$(3.5) \qquad d\underbrace{\begin{bmatrix} b_t \\ s_t \end{bmatrix}}_{X_t} = \underbrace{\begin{bmatrix} (\mu(s_t) - D)b_t \\ D(s_{in} - s_t) - \kappa\mu(s_t)b_t \end{bmatrix}}_{f(X_t)} dt + \underbrace{\begin{bmatrix} \omega_b b_t & 0 \\ 0 & \omega_s s_t \end{bmatrix}}_{g(X_t)} d\underbrace{\begin{bmatrix} B_t^b \\ B_t^s \end{bmatrix}}_{B_t^x},$$

with $X_0 \equiv (b_0, s_0)$. Now, we see how the solution to the model in Eq. (3.5) looks like for Case 1 and Case 2. The noise intensities are $\omega_x = (\omega_b = 0.05, \omega_s = 0.05)$ for Case 1 and $\omega_x = (\omega_b = 0.075, \omega_s = 0.075)$ for Case 2.

(●) stable steady states (●) unstable steady state
·········· ODE baseline ——— stochastic realisations ——— target realisations

Fig. 2: Trajectories from the stochastic model with growth function of the Monod type (left) and of the Haldane type (right). The dashed lines in the background show the trajectories from the deterministic model, as in Fig. 1. Here, a target realisation denotes a simulated trajectory which will be estimated from noisy measurements in Section 5.

Fig. 2 (left) suggests that processes X move around the trajectories of the deterministic chemostat model, with small fluctuations. Here, $D < \mu(s_{in})$. Under the deterministic model, this condition implies that the microbial population forming the biomass survives regardless of the initial concentrations of substrate and

biomass [17]. In Fig. 2 (right), processes X also move around the trajectories of the deterministic chemostat model, but in order to avoid "near" extinction of the population around $E_2^{(1)} = (0, s_{in})$, we would have to choose X_0 appropriately.

THEOREM 3.5. *For any initial condition $X_0 \in (\mathbb{R}^2)_+^*$, i.e. any $\{(b_0, s_0) \in \mathbb{R}^2 | b_0 > 0, s_0 > 0\}$, there exists a unique strong solution $X = (X_t, t \geq 0)$ of the system in Eq. (3.5) with growth functions from Eq. (2.4) or Eq. (2.5). More-over, the solution remains in $(\mathbb{R}^2)_+^*$ with probability one, i.e. $X_t \in (\mathbb{R}^2)_+^*$ for $t \geq 0$ almost surely.*

Proof. See Subsection SM1.3 in the supplementary material.

Thm. 3.5 assures the positivity of the solution for all $t \geq 0$. However, we have not yet ensured persistence of the microbial population, i.e. that the system admits a unique stationary distribution which is also ergodic. In the context of the chemostat, persistence was discussed in [33], where sufficient conditions for ergodicity were imposed on s_{in}, D, ω_b and ω_s.

Let ζ be an integrable function with respect to the measure μ, an invariant measure of the process X. If the system is ergodic, then $\mathbb{P}\left(\frac{1}{t} \int_0^t \zeta(X_s) ds \to \int \zeta(y)\mu(dy)\right) = 1$ [5]. When we let time remove the stochas-ticity, the time-average of ζ converges almost surely (if the limit exists) to the ex-pectation of ζ at an arbitrary time t. Define the time average limit and the Monte Carlo approximation to the expectation limit by $\langle\zeta\rangle_{\infty,1}^{tim}$ and $\langle\zeta\rangle_{t,\infty}^{ens}$, respectively. Let $X_t^{(n)}$ denote the n-th realisation of X_t, then define

$$(3.6) \quad \langle\zeta\rangle_{\infty,1}^{tim} \triangleq \lim_{t\to\infty} \frac{1}{t} \int_0^t \zeta(X_s) \, ds, \quad \langle\zeta\rangle_{t,\infty}^{ens} \triangleq \lim_{N\to\infty} \frac{1}{N} \sum_{n=1}^{N} \zeta(X_t^n), X_t^{(n)} \sim \mu.$$

Fig. 3: Estimates of the expectation and the time-average of ζ, where $\zeta(x) = 0.5b - 0.1s$. The estimates are based on 30 experiments. Shaded regions are delimited by two standard deviations above and below the mean. For the Monte Carlo approximation, we used an ensemble of $N = 512$ samples of X_t when computing the sample averages (note that $N = 1$ when computing the time averages). The initial concentrations are $(b_0, s_0) = (1.0\,\mathrm{mgL}^{-1}, 1.0\,\mathrm{mgL}^{-1})$ for Case 1 (Monod kinetics) and $(b_0, s_0) = (1.65\,\mathrm{mgL}^{-1}, 2.5\,\mathrm{mgL}^{-1})$ for Case 2 (Haldane kinetics).

Fig. 3 suggests that ergodicity holds for the system in Case 1, as shown by the commutativity in $\langle\zeta\rangle_{\infty,1}^{tim}$ and $\langle\zeta\rangle_{t,\infty}^{ens}$. In fact, this is confirmed by the fact that the conditions for ergodicity outlined in [33] hold. For Case 2, however, some trajectories of the microbial population go towards the washout node $E_2^{(3)}$, instead of the interior node $E_2^{(1)}$ (as in Fig. 2). Moreover, they do not escape this attractor in finite time, as comfirmed by the non-commutativity in the limits.

3.2. The Markov property and the Kolmogorov forward equation. In this section, we use the theory of Markov processes to study the properties of the SDE (3.5), given that its solution, if it exists, satisfies the Markov property. We will see that this allows us to apply the Kolmogorov equations (in particular the forward, or Fokker-Planck, equation) from the theory of Markov process into the SDE setting. This can then be used to express expectations of functions of the solution to Eq. (3.5) in terms of certain non-random PDEs.

DEFINITION 3.6 (Markov process). *Let the process $X = (X_t, \, t \geq 0)$ be measurable under the filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0, \infty)}, \mathbb{P})$. X is called an $\mathcal{F}_t$-Markov process if we have*

$$(3.7) \qquad \mathbb{E}[\phi(X_t)] \, = \, \mathbb{E}[\phi(X_t)|X_\tau] \, =: \, \phi_{t-\tau}(X_\tau),$$

for all $t \geq \tau$ and all bounded measurable functions ϕ. Here, $\mathbb{E}[\cdot]$ denotes expectation.

THEOREM 3.7 (Markov property and SDEs [12, Theorem 17.2.3]). *Assume existence and uniqueness of the solution to the SDE in Eq. (3.2). The solution X that satisfies such SDE is an $\mathcal{F}_t$-Markov process.*

The intuitive content behind Thm. 3.7 is clear. Firstly, suppose that the process X generated by the SDE in Eq. (3.2) begins at time zero, and that we watch it up to time τ. Based on this information, we want to compute the conditional expectation $\phi_{t-\tau}(X_\tau)$, where $t \geq \tau$. Then we can pretend that the process is starting at time τ at its current position, generate the solution to the SDE corresponding to this initial condition, and compute the expected value of $\phi(X_t)$ generated in this way.

For simplicity, we start by studying the random process $X = (X_t, \, t \geq 0)$ for $\tau = 0$, i.e. given only the $\mathcal{F}_0$-measurable D_x-dimensional random variable ξ. Moreover, rather than studying the path of this random process itself, we can look at how the non-random function ϕ_t varies with t.

Let $p(t, \cdot)$ denote the density with respect to Lebesgue measure for the probability measure $\mathbb{P}$ at time t. We describe the process in terms of a transition kernel $p(t, x|0, x_0) \overset{\Delta}{=} \mathbb{P}(X_t = x | X_0 = x_0), t \geq 0$. This allows us to compute the probability of having the process X at a point $x \in \mathbb{R}^{D_x}$ at instant t conditioned on starting at $x_0 \in \mathbb{R}^{D_x}$ at time $t_0 = 0$. We then proceed to use the property shown in Def. 3.6.

$$\mathbb{E}[\phi(X_t)] = \mathbb{E}[\mathbb{E}[\phi(X_t)|\mathcal{F}_0]] = \mathbb{E}[\mathbb{E}[\phi(X_t)|X_0]]$$

$$= \int_{\mathbb{R}^{D_x}} \int_{\mathbb{R}^{D_x}} \phi(x) p(t, x|0, x_0) p(0, x_0) dx_0 dx$$

$$(3.8) \qquad = \int_{\mathbb{R}^{D_x}} \phi(x) p(t, x) dx.$$

Now, we find the time evolution of the probability density $p(t, x)$ by differentiating both sides of equation Eq. (3.8) with respect to time,

$$(3.9) \qquad \frac{d}{dt} \mathbb{E}[\phi(X_t)] \, = \, \int_{\mathbb{R}^{D_x}} \phi(x) \frac{\partial}{\partial t} p(t, x) dx \, = \, \int_{\mathbb{R}^{D_x}} \phi(x) \mathcal{A}^* p(t, x) dx,$$

where we introduced $\mathcal{A}^*$ to describe the time evolution of the probability density.

The following proposition is adapted from [29, Chapter 4].

PROPOSITION 3.8 (The Kolmogorov forward equation (KFE)). *Assume that the $\mathcal{F}_t$-Markov process $X = (X_t, \, t \geq 0)$ is solution to the SDE (3.2). Suppose that the*

density $p(t, x)$ exists and is $\mathcal{C}^1$ in t and $\mathcal{C}^2$ in x, then $p(t, x)$ satisfies the Kolmogorov forward equation, also called the Fokker-Planck equation:

$$(3.10) \qquad \frac{\partial}{\partial t} p(t, x) = \mathcal{A}^* p(t, x), \quad p(0, x) = p_0(x),$$

where the form of the adjoint operator $\mathcal{A}^*$ is

$$(3.11) \quad \mathcal{A}^* p(t, x) = -\sum_{d_x=1}^{D_x} \frac{\partial}{\partial x_{d_x}} [f_{d_x}(x) p(t, x)] + \sum_{d_x=1}^{D_x} \sum_{d_x'=1}^{D_x} \frac{\partial^2}{\partial x_{d_x} \partial x_{d_x'}} [G_{d_x d_x'}(x) p(t, x)].$$

Here, $G_{d_x d_x'}(x) = \frac{1}{2} \sum_{m=1}^{M} g_{d_x m}(x) g_{d_x' m}(x)$ and the initial condition $p_0(x)$ is given.

At this point, our best knowledge about the state of the system is given by the distribution whose probability density solves the KFE, given the distribution of its initial state. This motivates us to study the model in Eq. (3.5) through its associated KFE, as in [8]. For instance, one can find the solution given the distribution of initial concentrations, and then verify how likely the state of a system is to be within regions around $E_2^{(1)}$ or $E_2^{(3)}$. We present the solution to the KFE for Cases 1 and 2 in Figs. 4 and 5, respectively. We solve the PDE by following similar steps as in [8]: we make use of an upwind differencing scheme for the space discretization and of an implicit scheme for the time discretization.

$t = 0\text{h}$ $t = 2\text{h}$ $t = 4\text{h}$ $t = 8\text{h}$ $t = 16\text{h}$

Fig. 4: Case 1 - solution to the KFE for the growth function of Monod type. Initial condition is $X_0 \sim \mathcal{N}([1, 1]^\top, 0.05^2 I_{2\times2})$. The discretization in space is in the domain $(0, 5] \times (0, 5]$, with a count of 256^2 finite volumes. Marginal density functions are represented by marginal curves. Each marginal curve has been normalized by its maximum value for the sake of illustration.

$t = 0\text{h}$ $t = 2\text{h}$ $t = 4\text{h}$ $t = 8\text{h}$ $t = 16\text{h}$

Fig. 5: Case 2 - solution to the KFE for the growth function of Haldane type. Initial condition is $X_0 \sim \mathcal{N}([1.65, 2.5]^\top, 0.05^2 I_{2\times2})$. The discretization in space is in the domain $(0, 3] \times (0, 3]$, with a count of 512^2 finite volumes. Again, marginal density functions are represented by marginal curves.

So far, we have worked with a model describing the dynamic behaviour of the system. The prior probabilistic knowledge about how the state of the system evolves was formally embedded by the adjoint operator $\mathcal{A}^*$ in the Kolmogorov forward equation. Our focus, however, is to refine our knowledge about the state when information about the system is available in the form of measurements. In the following section, we resort to an observation model to describe how the signal gets corrupted. Generally, the state signal cannot be measured directly (e.g. only certain components of a

vector-valued process are observed), and the measurements are also subjected to er-
ror. To describe how measurements and states are related, we combine the dynamics
with an observation model in which the error is also stochastic.

4. Observation models. Let $0 = t_0 < t_1 < \cdots < t_n \ldots$ be a uniform partition
of the interval $[t_0, \infty)$. We let discrete-time measurements z be available from the
system, in the form of

(4.1) $$z_t = h(X_t) + k(t)v_t, \quad t \in \{t_0, t_1 \ldots\}.$$

Here, $k : [0, \infty) \to \mathbb{R}^{D_y} \times \mathbb{R}^{D_y}$ denotes noise intensity, v_t denotes white Gaussian noise
and $h : \mathbb{R}^{D_x} \to \mathbb{R}^{D_y}$ is the observation function. We simplify again our notation by
omitting the dependency on the vector θ of parameters and on the input $u(t)$.
Instead of proceeding with the model in Eq. (4.1), we will work with an integrated
form of the observations; we use an observation model for a process $Y = (Y_t, t \geq 0)$,
where this process represents the integral of the actually observed process. To explain
this, we begin by defining a D_y-dimensional $\mathcal{F}_t$-Brownian motion B^y independent of
B^x. We then introduce a process Y which obeys

(4.2) $$dY_t = h(X_t)dt + k(t)dB_t^y, \quad Y_0 = 0.$$

Here, Y_0 is considered to be identically zero (this does not mean that measurements
at time 0 are zero-valued, but rather that there is no information available initially).
Notice that Y as modelled by Eq. (4.2) is the integral of the observed process,
so that in discrete-time one really observes z rather than Y. We can think of v_t in
Eq. (4.1) as the hypothetical time derivative of the Brownian motion in Eq. (4.2) at
time t, and that v has zero mean and covariance $\mathbb{E}(v_s v_t) = \delta(t - s)$, where $\delta(\cdot)$ is the
Dirac's delta generalized function. By working with the integrated form, we obtain a
mathematical model in which we can disregard the need to work with $\delta(\cdot)$ and instead
use the machinery of stochastic calculus used in the description of the signal model.

In Section 2, we introduced two settings for the dynamics governing the state of
the chemostat: Case 1 with Monod kinetics and Case 2 with Haldane kinetics.
We now additionally explore an observation model in which the sensor transforms
the state variables in one-dimensional measurements. We assume that we are given
the flow rate of biogas produced by the biomass after substrate consumption. This
setting is denoted by scenario A.
Let Q denote the flow of biogas production. The flow Q, being proportional to
the microbial activity [4, 30], can be written as $Q(s, b) = \gamma\mu(s)b$, where γ is the
yield coefficient for CH_4 in Eq. (2.1). The integrated process considered is

(4.3) $$d\left[y_t^A\right] = \underbrace{\left[Q(s,b)\right]}_{h_A(x_t)} dt + \underbrace{\left[\omega_A\right]}_{k_A(t)} d\left[B_t^{y^A}\right].$$

We combine Cases 1 and 2 with the observation model in scenario A to form the
experiments described in Table 2. During a sub-interval of interest $t \in [0, \delta_u]$ for
which the input $u(t)$ is held constant, we simulate trajectories from processes X
and Y with time discretization δ_x and δ_y, respectively. We simulate the trajectories
for $\delta_u = 16$ hours. We assume that we are able to collect the measurements $\{y_t^A\}$
at high-frequency ($\delta_y = \delta_x$) or at low-frequency ($\delta_y \gg \delta_x$). Again, the initial
concentrations are $(b_0, s_0) = (1.0\,\mathrm{mgL}^{-1}, 1.0\,\mathrm{mgL}^{-1})$ for Case 1 and $(b_0, s_0) = (1.65\,\mathrm{mgL}^{-1}, 2.5\,\mathrm{mgL}^{-1})$ for Case 2.

Table 2: Additional model parameters and inputs for differential equations of X and Y.

	Model parameters		Simulation parameters	
	Observation function	ω_A	δ_x (h)	δ_y (h)
1A - high-frequency	$27.5\mu(s)b$	0.01	0.01	0.01
1A - low-frequency	$27.5\mu(s)b$	$0.01\left(\frac{\delta_y}{\delta_x}\right)^{0.5}$	0.01	2.0
2A - high-frequency	$2.75\mu(s)b$	0.01	0.005	0.005
2A - low-frequency	$2.75\mu(s)b$	$0.01\left(\frac{\delta_y}{\delta_x}\right)^{0.5}$	0.005	2.0

Fig. 6: Measurements simulated from the trajectories highlighted in Fig. 2.

5. The filtering problem. To ensure theoretical developments, we assume that $X_0 \in \mathcal{L}^2(\mathbb{P})$ (remember that we need this assumption to determine the random process $X = (X_t, t \geq 0)$ uniquely, according to Thm. 3.3).

In state-space models, the Bayesian methodology of computing posterior distributions of a latent state based on a history of noisy measurements is known as filtering and smoothing. The aim of solving a filtering problem is to determine the conditional distribution of the signal X_t given the observation σ-algebra $\mathcal{F}_t^Y = \sigma\{Y_s : s \leq t\}$. We formalize this by defining a stochastic process $\pi = (\pi_t, t \geq 0)$ describing this conditional distribution. If we assume that such distribution has a density p_t with respect to Lebesgue measure, then

$$(5.1) \qquad \pi_t(A) \triangleq \mathbb{P}\left(X_t \in A | \mathcal{F}_t^Y\right), \; \pi_t(\phi) = \mathbb{E}_{\mathbb{P}}\left[\phi(X_t)|\mathcal{F}_t^Y\right] = \int_{\mathbb{R}^{D_x}} \phi(x)p_t(x)dx.$$

Here, A is an arbitrary set in the σ-algebra $\mathcal{B}([0,\infty)\otimes\mathbb{R}^{D_x})$, ϕ is an arbitrary function of the state and $\mathbb{E}_{\mathbb{P}}$ denotes expectation under the density p_t. The first equation takes values in the space of functions from $[0,\infty) \times \mathbb{R}^{D_x}$ into $[0,1]$. The second equation allows us to get desirable information about the state through a transformation ϕ, assuming that the expectation exists (i.e. $\int |\phi(x)|p_t(x)\,dx < \infty$). Whether π_t has

a density with respect to a reference measure (e.g. Lebesgue measure) is rigorously
discussed in [2]. We proceed by assuming that we can prove the existence of p_t.

5.1. Filtering equations. We continue working on $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \in [0,\infty)}, \mathbb{P})$. Our
goal is to compute π_t on the basis of the observations $\{Y_s : s \leq t\}$. We begin by
considering the signal process with evolution in time described by Eq. (3.2), and also
the observation model given by Eq. (4.2). We rewrite here both equations:

353
$$dX_t = f(X_t)dt + g(X_t)dB_t^x, \; X_0 = \xi; \quad dY_t = h(X_t)dt + k(t)dB_t^y, \; Y_0 = 0.$$

A detailed and formal derivation of the filtering equations is available in [2, 32].
Here, the derivation is briefly outlined, following similar steps as in [23]. Among
the possible ways of deducing the evolution equation for π_t, there are two commonly
used approaches, namely the change of measure method and the innovation process
method. We present the change of probability measure method. For the second
approach, see [2, Section 3.7].

5.1.1. Changes of measure. If we have two such measures $\mathbb{P}$ and $\mathbb{Q}$, then $\mathbb{P}$
is called *absolutely continuous* with respect to $\mathbb{Q}$ if every nullset of $\mathbb{Q}$ is a null set of
$\mathbb{P}$. Moreover, $\mathbb{P}$ and $\mathbb{Q}$ are called *equivalent* if they have the nullsets. In other words,
if A denotes an event, and $\mathbb{P}(A)$ denotes its probability, then equivalence means that
$\mathbb{Q}(A) = 0$ if and only if $\mathbb{P}(A) = 0$.
Changing the measure allows us to compute expectations of a measurable function
$\phi(x)$ with respect to a measure $\mathbb{Q}$, which were originally expressed with respect to
measure $\mathbb{P}$. To see this, consider the two measures $\mathbb{P}$ and $\mathbb{Q}$ for some real-valued
random variable X, and write them in terms of their densities p, q (again, with respect
to Lebesgue measure). We then have

(5.2)
$$\mathbb{E}_{\mathbb{P}}[\phi(X)] = \int \phi(x)p(x)\,dx = \int \frac{p(x)}{q(x)}\phi(x)q(x)\,dx = \mathbb{E}_{\mathbb{Q}}[L(X)\phi(X)],$$

where we introduced the likelihood ratio $L(x) \triangleq \frac{p(x)}{q(x)}$. Changing the measure proves
to be useful whenever expectations under $\mathbb{Q}$ are easier to compute than under $\mathbb{P}$.
If $\mathbb{P}$ is a probability measure and we have a collection of processes (X and Y), the
measure $\mathbb{P}_t$ is the restriction of $\mathbb{P}$ to all events that can be described in terms of the
behaviour of $\{X_s\}$ and $\{Y_s\}$ for $0 \leq s \leq t$. In other words, $\mathbb{P}_t$ denotes the restriction
of the measure $\mathbb{P}$ to $[0, t] \times \mathbb{P}$. If $\mathbb{P}$ and $\mathbb{Q}$ are equivalent, also their restrictions $\mathbb{P}_t$
and $\mathbb{Q}_t$ are equivalent. By using the Radon–Nikodym theorem [22, Theorem 10.6],
we know that a random variable L_t exists, for which

(5.3)
$$\mathbb{E}_{\mathbb{P}}[\phi(X_t)] = \mathbb{E}_{\mathbb{Q}}[L_t\phi(X_t)],$$

380 where $L_t \triangleq \frac{d\mathbb{P}_t}{d\mathbb{Q}_t}$ is called the Radon–Nikodym derivative of $\mathbb{P}_t$ with respect to $\mathbb{Q}_t$.
381   As we previously mentioned, we aim at computing the conditional expectation
382 for some function ϕ. In terms of a reference probability measure $\mathbb{Q}$, the conditional
383 expectation is written as

384   (5.4)
$$\mathbb{E}_{\mathbb{P}}\left[\phi(X_t)|\mathcal{F}_t^Y\right] = \frac{\mathbb{E}_{\mathbb{Q}}\left[\phi(X_t)\,L_t \mid \mathcal{F}_t^Y\right]}{\mathbb{E}_{\mathbb{Q}}\left[L_t \mid \mathcal{F}_t^Y\right]} =: \frac{\rho_t(\phi)}{\rho_t(1)}.$$

Eq. (5.4) is known as a Bayes' formula for stochastic processes or *Kallianpur-*
Striebel formula. Here, we have defined the estimate $\rho_t(\phi) = \mathbb{E}_{\mathbb{Q}}\left[\phi(X_t)\,L_t \mid \mathcal{F}_t^Y\right]$.

The hope is that we can pick a reference measure $\mathbb{Q}$ such that both L_t and q are
simple enough to make $\mathbb{E}_\mathbb{P}\left[\phi(X_t)|\mathcal{F}_t^Y\right]$ more tractable to compute via Eq. (5.4) than
via an integration over the measure $\mathbb{P}$. For instance, some simplification might be
achieved by switching from a model with measure $\mathbb{P}$ in which X and Y are coupled,
i.e. statistically dependent, to a model with measure $\mathbb{Q}$ where they are independent
(while preserving the distribution of X_t). For our filtering problem, we will choose a
reference measure $\mathbb{Q}$ such that the path of the observations $\{Y_{0:t}\}$ (or equivalently the
set of the increments $\{dY_{0:t}\}$) becomes independent of the path of the state process
$\{X_{0:t}\}$, i.e. $q(t, X_{0:t}, dY_{0:t}) = p(t, X_{0:t})q(t, dY_{0:t})$.

For the time being, we will concentrate not on $\pi_t(\phi)$, but on its unnormalised
counter-part $\rho_t(\phi)$. Our next step is to find an explicit expression for $\rho_t(\phi)$. We start
by finding the expression for L_t, and proceed by applying Itô's rule to the process
$\phi(X_t)L_t$, then we compute the conditional expectation of this expression. Once we
have accomplished this, the remainder is another application of Itô's rule, which gives
us an expression for the normalized quantity $\pi_t(\phi)$.

5.1.2. Filtering for observations corrupted by Gaussian noise. For the
derivations in this section, we operate under some technical conditions: (i) $k(t)$ is
invertible for all $t \in [0, \infty)$, and $k(t), k^{-1}(t)$ are locally bounded; (ii) the solution to the
SDE in Eq. (3.2) exists and is unique; (iii) h has linear growth. See Subsection SM1.4
for detailed calculation steps.

We chose a reference measure $\mathbb{Q}$ such that the path of the observations $\{dY_{0:t}\}$
becomes independent of the path of the state process $\{X_{0:t}\}$, and yet the law of X is
the same under $\mathbb{P}$ as it is under $\mathbb{Q}$. By definition, let us see what the likelihood ratio
L_t has to look like for this measure change:

(5.5)　　$L_t = \dfrac{d\mathbb{P}_t}{d\mathbb{Q}_t} = \dfrac{p(t, X_{0:t}, dY_{0:t})}{q(t, X_{0:t}, dY_{0:t})} = \dfrac{p(t, dY_{0:t}|X_{0:t})p(t, X_{0:t})}{p(t, X_{0:t})q(t, dY_{0:t})} = \dfrac{p(t, dY_{0:t}|X_{0:t})}{q(t, dY_{0:t})}.$

The observation model in Eq. (4.2) assumes that increments dY_t are independently
distributed given the state X_t, and we know that $dY_t|X_t \sim \mathcal{N}(dY_t; h(X_t)dt, kk^\top(t)dt)$.
Furthermore, we choose $q(t, dY_t)$ as the density of $\mathcal{N}(dY_t; 0, kk^\top(t)dt)$. Thus, the
Radon-Nikodym derivative L_t can be written as

(5.6)　　$L_t = \exp\left[\displaystyle\int_0^t (k^{-1}(s)h(X_s))^\top k^{-1}(s)dY_s - \dfrac{1}{2}\int_0^t \|k^{-1}(s)h(X_s)\|^2 ds\right].$

Remark 5.1. An even more general model arises when we consider that the ob-
servation equation is driven by a signal dependent noise; that is, $k(t, X_t)dB_t^y$ instead
of $k(t)dB_t^y$. The filtering problem with this type of observation model can be then
converted into a classical one (signal independent observation noise) via a suitably
chosen stochastic flow mapping. We refer the interested reader to [13].

Eq. (5.6), along with Itô's formula, leads to the following proposition:

PROPOSITION 5.2 (The Zakai equation). *Let ϕ be $\mathcal{C}^2$ (twice continuously differ-*
entiable) and suppose that ϕ and all its derivatives are bounded. Then we can write

(5.7)　　　　$\rho_t(\phi) = \rho_0(\phi) + \displaystyle\int_0^t \rho_s(\mathcal{A}^*\phi)ds + \int_0^t \rho_s(k^{-1}(s)h\phi)^\top d\tilde{Y}_s,$

where $\rho_0(\phi) = \mathbb{E}_\mathbb{Q}\left[\phi(X_0)L_0|\mathcal{F}_0^Y\right]$, $h\phi = h(x)\phi(x)$, $\mathcal{A}^*$ is the operator in Eq. (3.9),
and the process $\tilde{Y} = (\tilde{Y}_t, t \geq 0)$ is the rescaled process $(\tilde{Y}_t = k^{-1}(t)Y_t, t \geq 0)$.

If we apply integration by parts to the Zakai equation, we obtain

(5.8) $$dq_t(x) = \mathcal{A}^* q_t(x) dt + q_t(x)(k^{-1}(t)h(x))^\top d\tilde{Y}_t.$$

Now that we have an equation for $\rho_t(\phi)$, the equation for $\pi_t(\phi)$ results from
another application of Itô's formula, this time on $\rho_t(\phi)/\rho_t(1)$, and of the Kallianpur-
Striebel formula introduced previously.

PROPOSITION 5.3 (The Kushner-Stratonovich equation). *Let ϕ be $\mathcal{C}^2$ and sup-*
pose that ϕ and all its derivatives are bounded. Then we can write

(5.9)
$$\pi_t(\phi) = \pi_0(\phi) + \int_0^t \pi_s(\mathcal{A}^*\phi)\, ds + $$
$$\int_0^t \pi_s(k^{-1}(s)h\phi) - \pi_s(\phi)\pi_s(k^{-1}(s)h))^\top (d\tilde{Y}_s - \pi_s(k^{-1}(s)h)\, ds),$$

where $\pi_0(\phi) = \mathbb{E}_\mathbb{Q}\left[\phi(X_0) L_0 | \mathcal{F}_0^Y\right] = \mathbb{E}_\mathbb{P}\left[\phi(X_0)\right]$ and $h\phi = h(x)\phi(x)$.

If we apply integration by parts to the Kushner-Stratonovich equation, we obtain

(5.10) $$dp_t(x) = \mathcal{A}^* p_t(x)\, dt + p_t(x)\left(k^{-1}(t)(h\phi - \pi_t(h))\right)^\top (d\tilde{Y}_t - \pi_t(k^{-1}(t)h)\, dt).$$

We have now obtained a stochastic integral expression for $\pi_t(\phi)$ (the Kushner-
Stratonovich equation) and for the evolution of p_t.

5.2. Approximations to the solution of the Kushner-Stratonovich equa-
tion. This section contains an overview of two classes of methods used to approximate
the solution to the filtering problem. For each class, we give a brief description of
the ideas behind the methods and state some related results. Note that so far we
have only presented the filter $\pi_t(\phi)$ on an unspecified set in the Borel σ-algebra of
$\mathcal{C}(\mathbb{R}_+, \mathbb{R}^{D_y})$ of measure one, as well as its assumed density p_t with respect to Lebesgue
measure. We have not yet shown what the filtered estimated $\pi_t(\phi)$ would be for a
given observation sample path $z \in \mathcal{C}([0, T], \mathbb{R}^{D_y})$. Typically, the observation data is
recorded at discrete times, and only these data are made available and used. Con-
sidering the practical implementation of a non-linear filter, we present the notion of
pathwise filtering as introduced in [11], and studied by many authors, as in [15] and
[14].

L_t in Eq. (5.6) is defined in terms of a stochastic integral with the observations
as the integrator. However, since the integrand is independent of the observations,
we can reverse the role of processes X and Y. Instead of using the observations as
the integrator, we compute L_t with an integral for every process sample path $\{X_{0:T}\}$,
provided that $h(X_t)$ is a semimartingale. Through an integration by parts:

(5.11) $$\int_0^T \tilde{h}^\top(X_s) d\tilde{Y}_s = \tilde{h}^\top(X_T)\tilde{Y}_T - \int_0^T \tilde{Y}_s^\top d\tilde{h}(X_s) \quad \text{almost surely.}$$

Here, the right-hand side can be interpreted pathwise with respect to the observations.
The pathwise Kallianpur-Striebel formula reads as follows:

(5.12) $$\hat{L}_T = \exp\left[\tilde{h}^\top(X_T)\tilde{Y}_T - \int_0^T \tilde{Y}_s^\top d\tilde{h}(X_s) - \frac{1}{2}\int_0^T \|\tilde{h}(X_s)\|^2 ds\right],$$

where $\hat{L}_T$ stands for L_T computed with respect to a fixed observation path. Conse-
quently, the SPDEs (5.8) and (5.10) reduce to PDEs with random coefficients.

5.2.1. Finite-differences approach + Splitting-up approximation. The following exposition follows closely the description of the method given in [7]. Let $0 = t_0 < t_1 < \cdots < t_n \ldots$ be a uniform partition of the interval $[0, \infty)$ with time step $\Delta = t_n - t_{n-1}$. If the set of observations $\{z_i, 1 \leq i \leq n\}$ is available, the density p_{t_n} will be approximated by p_n^Δ, where the transition from p_{n-1}^Δ to p_n^Δ follows a splitting-up approximation [24, 25].

Fig. 7: Splitting-up approximation for Case 1A - low-frequency.

Fig. 8: Splitting-up approximation for Case 2A - low-frequency.

The first step, called the *prediction* step, consists in solving the Fokker-Planck equation for the time interval $[t_{n-1}, t_n]$. We rewrite the PDE here,

$$(5.13) \qquad \frac{\partial p_t^n(x)}{\partial t} = \mathcal{A}^* p_t(x), \qquad p_{t_{n-1}}^n = p_{n-1}^\Delta,$$

and we denote the prior estimate by $\bar{p}_n^\Delta \triangleq p_{t_n}$.

475   The second step, called the *correction* step, uses the new observation z_n to up-
476 date $\bar{p}_n^\Delta$. We use the approximation from pathwise filtering to the Radon-Nikodym
477 derivative (Eq. (5.12)). We compute $p_n^\Delta(x)$ via $p_n^\Delta(x) \triangleq c_n \zeta_n^\Delta(x)\bar{p}_n^\Delta$, where c_n is a
478 normalization constant chosen such that $\int_{\mathbb{R}^{D_x}} p_n^\Delta(x)dx = 1$ and

479   (5.14)
$$\zeta_n^\Delta(x) \triangleq \exp\left(\tilde{h}^\top(x)\tilde{z}_n - \frac{1}{2}\Delta\|\tilde{h}(x)\|^2\right),$$

Here, $\tilde{z}_n = k^{-1}(t_n)z_n$ and $\tilde{h}(x) = k^{-1}(t_n)h(x)$.

5.2.2. Particle methods. We now switch to particle filtering, a technique used
to approximate the solution to the filtering problem by a finite number of samples
from the conditional distribution π_t. The following exposition is adapted from [2].
Let $V_j, j = 1, \ldots, N$, be N mutually independent stochastic processes, all in-
dependent of Y, each of them being a solution to Eq. (3.2). The pairs (V_j, Y) are
identically distributed and have the same distribution as the pair (X, Y) under $\mathbb{Q}$.
Also, let $w_j, j = 1, \ldots, N$, be the following exponential martingales

(5.15)
$$w_j(t) = 1 + \int_0^t w_j(s)(k^{-1}(s)h(V_j(s)))^\top k^{-1}(s)dY_s, \quad t \geq 0,$$

489 that is, for $t \geq 0$
(5.16)
490
$$w_j(t) = \exp\left(\int_0^t (k^{-1}(s)h(V_j(s)))^\top k^{-1}(s)dY_s - \frac{1}{2}\int_0^t \|k^{-1}(s)h(V_j(s))\|^2 ds\right).$$

491 As a result, the triples $(V_j, w_j, Y), j = 1, \ldots, N$, are identically distributed and have
492 the same distribution as the triple (X, L, Y) under $\mathbb{Q}$. Lastly, let $\rho^N = \{\rho_t^N, t \geq 0\}$
493 and $\pi^N = \{\pi_t^N, t \geq 0\}$ be the following sequences of measure-valued processes

494   (5.17)
$$\rho_t^N \triangleq \frac{1}{N}\sum_{j=1}^N w_j(t)\delta_{V_j(t)}, \quad \pi_t^N \triangleq \frac{\rho_t^N}{\rho_t^N(1)} = \sum_{j=1}^N \bar{w}_j^N(t)\delta_{V_j(t)},$$

where the normalised weights $\bar{w}_j(t)$ are given by $\bar{w}_j(t) = \left(\sum_{j'=1}^N w_{j'}(t)\right)^{-1} w_j(t), j' =$
$1, \ldots, N, t \geq 0$. That is, ρ_t^N is the empirical measure of N (random) particles with
positions $V_j(t)$ and weights $w_j(t), j = 1, \ldots, N$, and π_t^N is its normalized version.
Under certain conditions, it is possible to show both that (i) $\rho_t^N(\phi)$ converges in
expectation to the unnormalised conditional estimator $\rho_t(\phi)$ with rate of convergence
$N^{-\frac{1}{2}}$ and (ii) $\pi_t^N(\phi)$ converges almost surely with a rate slightly lower than $N^{-\frac{1}{2}}$
[2]. Additionally, $\pi_t^N(\phi)$ converges in expectation to $\pi_t(\phi)$, provided the integrability
condition on $\rho_t^{-1}(1)$. Recall the martingale term L introduced in Eq. (5.5). The
validity of these results requires the existence of higher moments of L. In the scenario
where we cannot assure that the higher moments of the martingale L exist, and we
only know that L is $\mathbb{Q}$-integrable. In this case, $\pi_t^N(\phi)$ still converges almost surely to
$\pi_t(\phi)$ for a fixed observation path $t \mapsto Y_t$ due to the strong law of large numbers [2].

Let us revisit the case outlined in Table 2 and Fig. 6. Without observations, the
probabilities of the hidden state variables evolved according to the Fokker-Planck
equation, as illustrated in Fig. 4 and Fig. 5. We then considered an observation
model for which the solution to each of the corresponding Kushner-Stratonovich

equation had to be approximated. Now, we firstly find the solution via finite-differences and a splitting-up approximation. We then apply a sequential Monte Carlo method (the continuous-time bootstrap particle filter) and a linearisation method (the extended Kalman filter). Fig. 9 and 10 show the results for a fixed instant t. Algorithms are available in the supplementary material (Section SM2).

We analyze the similarity among the approximated solutions by computing the Hellinger distance $\mathcal{H}$ between two probability distributions, say F_1 and F_2. With respect to Lebesgue measure, let f_1 and f_2 denote the density functions of the distributions F_1 and F_2, respectively. The square of the Hellinger distance is then defined as $\mathcal{H}^2(f_1, f_2) = \frac{1}{2} \int \left(\sqrt{f_1(x)} - \sqrt{f_2(x)} \right)^2 dx$. Note how this distance function is symmetric and is zero-valued when the compared densities are equal.

Fig. 9: Case 1A - approximated solution to the Kushner-Stratonovich equation at $t = 16$h with the finite differences and the splitting-up approximation for PDEs (KSE·PDE), with the bootstrap particle filter (KSE·BPF) and with the EKF (KSE·EKF). For the PDE method, the discretization in space is in the domain $(0, 5.0] \times (0, 5.0]$. We consider a refinement of the grid of a total count of 256^2 finite volumes. As for the BPF, we consider $N = 2.5 \times 10^6$. In white, we show the trajectory from which the measurements were simulated. The same trajectory appears in Fig. 2.

Fig. 10: Case 2A - approximated solution to the Kushner-Stratonovich equation at $t = 8$h with the finite differences and the splitting-up approximation for PDEs (KSE·PDE), with the bootstrap particle filter (KSE·BPF) and with the EKF (KSE·EKF). For the PDE method, the discretization in space is in the domain $(0, 3.0] \times (0, 3.0]$. We consider a refinement of the grid of a total count of 256^2 finite volumes. As for the BPF, $N = 2.5 \times 10^6$. Note how, differently from the solution to the KFE, we move more probability mass towards the washout on account of the measurements.

In Fig. 11, we plot the square of the Hellinger distance between the conditional distribution given by different setups of the method for PDEs and the approximation given by the BPF. It is clear that the discretization error decreases with a finer refinement of the grid. We also include the square of the Hellinger distance between the approximation with the EKF and with the BPF. The EKF fails to capture higher moments of the distribution, hence its divergence from a sequential Monte Carlo. Non-zero skewness and excess kurtosis result from having both a non-linear dynamic process and a non-linear measurement process with respect to the state variables. This can be seen in the Section SM3 of the supplementary material, where we plot the shape of the conditional distribution for different t's.

Fig. 11: Case 1A - Squared Hellinger distances between the distributions obtained by approximating the solution to the Kushner-Stratonovich equation with the extended Kalman filter and with methods for partial differential equations using different grid refinements. We consider three scenarios for the refinement of the grid, each containing a total count of $32^2, 128^2, 512^2$ finite volumes, respectively. As for the particle filter, we consider $N = 2.5 \times 10^6$.

Fig. 12: Case 2A - Squared Hellinger distances between the distributions obtained by approximating the solution to the Kushner-Stratonovich equation with the extended Kalman filter and with methods for partial differential equations using different grid refinements. We consider three scenarios for the refinement of the grid, each containing a total count of $32^2, 128^2, 512^2$ finite volumes, respectively. As for the particle filter, we consider $N = 2.5 \times 10^6$.

In Fig. 12, we again plot the square of the Hellinger distance between the con-

ditional distribution given by different setups of the method for PDEs and the approximation given by the BPF, now for Case 2A. We again include the square of the Hellinger distance between the approximations with the EKF and the BPF. Here, the EKF suffers in the highly non-linear region of the state-space, namely when the state is near the saddle point, $t > 4$h.

Note that the grid with 512^2 cells fails to give a perfect match between the method for PDEs and the particle method. At $t = 8h$, for instance, the Hellinger distance between both distributions increases, even though both methods approximate the same updating equation. The approximations with methods for PDEs or with the EKF are naturally smooth, whereas for such fine grid, estimates based on the particles' paths degenerate. This issue is similar to the so-called sample impoverishment, which is a reduction in the number of distinct sample values when the process model noise variance is very low, such that samples will hardly diversify and after a few iterations all particles will collapse into a single point in the state space. There are different solutions for this problem, the simplest one to implement being an increase in the number of particles. This, however, leads to unreasonable computational load. Other solutions are more complicated to implement and, unlike the BPF, they might need to be tailored to the problem in hand. See [9] and [26] for more methods to prevent sample impoverishment.

6. Concluding remarks. Whether it be an approximation to the Kushner-Stratonovich equation with particle methods or an approximation to the Zakai equation with methods for partial differential equations, no restrictions are imposed on the shape of the distribution. Therefore, these two approximation methods are often more accurate than linearization methods.

In this paper, we presented a scenario where persistence of the microorganisms forming the biomass is ensured under the deterministic model, but not under the stochastic one (Case 2). The corresponding filtering distribution π_t helped to identify the actual state of the system. π_t can be used, for instance, to design a control strategy on the input variables $D(t)$ and $s_{in}(t)$ in order prevent microorganism extinction.

We assumed that measurements of biogas flow rate were available, at either high- or low-frequency. We denoted this setup by scenario A. In the supplementary material, we additionally present results for Cases 1 and 2 under three other scenarios: in B, we are given noisy measurements of the substrate concentration at low-frequency; in C, we are given noisy measurements of the biomass concentration at low-frequency; in D, we are given noisy measurements of both biomass and substrate concentrations at low-frequency. We refer the reader to Section SM3 of the supplementary material for a comparison of the approximations outlined in this work under all four scenarios.

REFERENCES

[1] J. F. ANDREWS, *A mathematical model for the continuous culture of microorganisms utilizing inhibitory substrates*, Biotechnology and bioengineering., 10 (1968), https://doi.org/10.1002/bit.260100602.

[2] A. BAIN AND D. CRISAN, *Fundamentals of Stochastic Filtering*, Stochastic Modelling and Applied Probability, Springer New York, 2008, https://doi.org/10.1007/978-0-387-76896-0.

[3] V. E. BENEŠ, *Exact finite-dimensional filters for certain diffusions with nonlinear drift*, Stochastics, 5 (1981), pp. 65–92, https://doi.org/10.1080/17442508108833174.

[4] O. BERNARD, Z. HADJ-SADOK, D. DOCHAIN, A. GENOVESI, AND J.-P. STEYER, *Dynamical model development and parameter identification for anaerobic wastewater treatment process*, Biotechnology and bioengineering, 75 (2001), https://doi.org/10.1002/bit.10036.

[5] G. D. Birkhoff, *Proof of the ergodic theorem*, 1931, https://doi.org/10.1073/pnas.17.2.656.

[6] A. T. Bull, *The renaissance of continuous culture in the post-genomics age*, Journal of Industrial Microbiology and Biotechnology, 37 (2010), https://doi.org/10.1007/s10295-010-0816-4.

[7] Z. Cai, F. L. Gland, and H. Zhang, *An adaptive local grid refinement method for nonlinear filtering*. Research report 2679, INRIA, 1995, https://hal.inria.fr/inria-00074012/document.

[8] F. Campillo, M. Joannides, and I. Larramendy-Valverde, *Approximation of the fokker–planck equation of the stochastic chemostat*, Mathematics and Computers in Simulation, 99 (2014), pp. 37–53, https://doi.org/10.1016/j.matcom.2013.04.012.

[9] J. Carpenter, P. Cliffordy, and P. Fearnhead, *An improved particle filter for nonlinear problems*, IEE Proc. Radar, Sonar Navig., 146 (1999), pp. 2–7, https://doi.org/10.1049/ip-rsn:19990255.

[10] K. Chung and R. Williams, *Introduction to Stochastic Integration*, Modern Birkhäuser Classics, Springer New York, 2014.

[11] J. M. C. Clark, *The design of robust approximations to the stochastic differential equations of nonlinear filtering*, 1978, https://doi.org/10.1007/978-94-011-7577-7_41.

[12] S. Cohen and R. Elliott, *Stochastic Calculus and Applications*, Probability and Its Applications, Springer New York, 2015, https://doi.org/10.1007/978-1-4939-2867-5.

[13] D. Crisan, M. Kouritzin, and J. Xiong, *Nonlinear filtering with signal dependent observation noise*, Electronic Journal of Probability, 14 (2009), https://doi.org/10.1214/EJP.v14-687.

[14] D. Crisan, T. G. Kurtz, and S. Ortiz-Latorre, *Particle representation for the solution of the filtering problem. application to the error expansion of filtering discretizations*, Journal of Stochastic Analysis, 2 (2021), https://doi.org/10.31390/josa.2.3.15.

[15] M. H. A. Davis, *A pathwise solution of the equations of nonlinear filtering*, Theory of Probability & Its Applications, 27 (1982), https://doi.org/10.1137/1127017.

[16] O. Hadj-Abdelkader and A. Hadj-Abdelkader, *Estimation of substrate and biomass concentrations in a chemostat using an extended kalman filter*, International Journal of Bioautomation, 23 (2019), pp. 215–232, https://doi.org/10.7546/ijba.2019.23.2.000551.

[17] J. Harmand, C. Lobry, A. Rapaport, and T. Sari, *The Chemostat: Mathematical Theory of Microorganism Cultures*, John Wiley and Sons, ISTE, 2017, https://doi.org/10.1002/9781119437215.

[18] P. A. Hoskisson and G. Hobbs, *Continuous culture–making a comeback?*, Microbiology, 151 (2005), https://doi.org/10.1099/mic.0.27924-0.

[19] L. Imhof and S. Walcher, *Exclusion and persistence in deterministic and stochastic chemostat models*, Journal of Differential Equations, 217 (2005), https://doi.org/10.1016/j.jde.2005.06.017.

[20] K. Ito, *Stochastic integral*, Proceedings of the Imperial Academy, 20 (1944), https://doi.org/10.3792/pia/1195572786.

[21] R. E. Kálmán and R. S. Bucy, *New results in linear filtering and prediction theory*, Journal of Basic Engineering, 83 (1961), https://doi.org/10.1115/1.3658902.

[22] F. Klebaner, *Introduction to Stochastic Calculus with Applications*, Introduction to Stochastic Calculus with Applications, Imperial College Press, 2005.

[23] A. Kutschireiter, S. C. Surace, and J.-P. Pfister, *The hitchhiker's guide to nonlinear filtering*, Journal of Mathematical Psychology, 94 (2020), https://doi.org/10.1016/j.jmp.2019.102307.

[24] F. Le Gland, *Time discretization of nonlinear filtering equations*, Dec. 1989, https://doi.org/10.1109/CDC.1989.70650.

[25] F. LeGland, *Splitting-up approximation for SPDE's and SDE's with application to nonlinear filtering*, 1992, https://doi.org/10.1007/BFb0007332.

[26] T. Li, S. Sun, T. P. Sattar, and J. M. Corchado, *Fight sample degeneracy and impoverishment in particle filters: A review of intelligent approaches*, Expert Systems with Applications, 41 (2014), https://doi.org/10.1016/j.eswa.2013.12.031.

[27] X. Mao, *Stochastic differential equations and applications*, 2nd ed., 2007, https://doi.org/10.1533/9780857099402.

[28] J. Monod, *La technique de culture continue: theorie et applications*, 1950, https://doi.org/10.1016/B978-0-12-460482-7.50023-3.

[29] I. Nobuyuki and S. Watanabe, *Stochastic Differential Equations and Diffusion Processes*, North-Holland Pub. Co.; Kodansha; Sole Distributors for the U.S.A. and Canada, 1981, https://doi.org/10.1016/S0924-6509(08)70217-4.

[30] P. Renard, D. Dochain, G. Bastin, H. P. Naveau, and J. Nyns, *Adaptive control of anaerobic digestion processes—a pilot-scale application*, Biotechnology and Bioengineering, 31 (1988), https://doi.org/10.1002/bit.260310402.

[31] P. Sinner, M. Stiegler, O. Goldbeck, G. M. Seibold, C. Herwig, and J. Kager, *Online estimation of changing metabolic capacities in continuous Corynebacterium glutamicum cultivations growing on a complex sugar mixture*, Biotechnology and Bioengineering, 119 (2021), pp. 575–590, https://doi.org/10.1002/bit.28001.

[32] R. van Handel, *Filtering, stability, and robustness*, PhD thesis, 2007, https://doi.org/10.7907/4p53-1h42.

[33] L. Wang and D. Jiang, *A note on the stationary distribution of the stochastic chemostat model with general response functions*, Applied Mathematics Letters, 73 (2017), pp. 22–28, https://doi.org/10.1016/j.aml.2017.04.029.

[34] A. K. Zvonkin, *A transformation of the phase space of a diffusion process that removes the drift*, Mathematics of the USSR-Sbornik, 22 (1974), pp. 129–149, https://doi.org/10.1070/sm1974v022n01abeh001689.