High-throughput sequencing of macaque basolateral amygdala projections reveals dissociable connectional motifs with frontal cortex

Authors: Zachary R Zeisler[1], Liza London[1], William G Janssen[1,2], J Megan Fredericks[1], Catherine Elorette[1], Atsushi Fujimoto[1], Huiqing Zhan[5], Brian E Russ[1,3,4], Roger L Clem[1], Patrick R Hof[1], Frederic M Stoll[1], and Peter H Rudebeck[1]*

Author affiliations:

1. Nash Family Department of Neuroscience and Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, One Gustave L. Levy Place, New York, NY 10029
2. Microscopy and Advanced Bioimaging CoRE, Icahn School of Medicine at Mount Sinai, One Gustave L. Levy Place, New York, NY 10029
3. Center for Biomedical Imaging and Neuromodulation, Nathan Kline Institute, 140 Old Orangeburg Road, 10 Orangeburg, NY 10962
4. Department of Psychiatry, New York University at Langone, One, 8 Park Ave, New York, NY 10016
5. Cold Spring Harbor Laboratory, 1 Bungtown Rd, Syosset, NY 11791

*Corresponding author

Correspondence should be addressed to:

Dr. Peter Rudebeck – peter.rudebeck@mssm.edu
Nash Family Department of Neuroscience
Icahn School of Medicine at Mount Sinai
One Gustave L. Levy Place
New York, NY 10029

Author Contributions: ZRZ, RLC, PRH, FMS, and PHR conceived the project. ZRZ and PHR designed and carried out the experiments, performed the analyses, and wrote the paper. LL performed the stereology analysis, WGJ assisted in tissue preparation, JMF assisted in surgery, CE and AF collected the fMRI scans, HZ performed the RNA extraction and sequencing, and BER performed the fMRI preprocessing and analysis. RLC, PRH, and FMS provided vital feedback on analysis approaches throughout. All authors approved the final version of the paper.

Acknowledgements: ZRZ, RLC, PRH, and PHR are supported by a grant from the BRAIN initiative (R34NS122050). We thank Allison Sowa of the ISMMS Microscopy and Advanced Bioimaging CoRE for help with tissue preparation, members of the Rudebeck Lab for assistance with surgical procedures, and the MAPseq Core at CSHL for assistance in tissue processing and sequencing. We also thank Anthony Zador and Alex Vaughan for advice and encouragement and Elisabeth Murray for comments on an earlier version of the manuscript.

Abstract

The basolateral amygdala (BLA) projects widely across the macaque frontal cortex[1–4], and amygdalo-frontal projections are critical for optimal emotional responding[5] and decision-making[6]. Yet, little is known about the single-neuron architecture of these projections: namely, whether single BLA neurons project to multiple parts of the frontal cortex. Here, we use MAPseq[7] to determine the projection patterns of over 3000 macaque BLA neurons. We found that one-third of BLA neurons have two or more distinct targets in parts of frontal cortex and of subcortical structures. Further, we reveal non-random structure within these branching patterns such that neurons with four targets are more frequently observed than those with two or three, indicative of widespread networks. Consequently, these multi-target single neurons form distinct networks within medial and ventral frontal cortex consistent with their known functions in regulating mood and decision-making. Additionally, we show that branching patterns of single neurons shape functional networks in the brain as assessed by fMRI-based functional connectivity. These results provide a neuroanatomical basis for the role of the BLA in coordinating brain-wide responses to valent stimuli[8] and highlight the importance of high-resolution neuroanatomical data for understanding functional networks in the brain.

Introduction

Basolateral amygdala (BLA) is essential for adaptive emotional responding in humans

and animals[5,9,10]. In humans, dysfunction within or damage to the circuits that connect through

the amygdala are theorized to be the cause of numerous psychiatric disorders, including autism

spectrum disorder, post-traumatic stress disorder, and schizophrenia[11–13]. Based on decades of

tract-tracing studies in macaques, we know that the primate BLA projects widely across the

brain, sending connections primarily to ventral and medial parts of the frontal lobe, as well as

the temporal and occipital cortices, thalamus, and striatum[1–4]. These diverse and widespread

connections, especially those to the frontal lobe, are central to accounts of how BLA in humans

coordinates learning about and responding to different emotionally salient events[14,15].

Despite the appreciation that the BLA plays a central role in coordinating activity across

large networks to guide emotional behavior, the anatomical organization of single neuron

connections from this area are largely unknown. One possibility is that single BLA neurons

project to only one specific target, transmitting information to downstream targets in dedicated

pathways. Such an organizing principle or connectional motif would align closely with the BLA's

known role in model-based behaviors and the processing of sensory-specific stimuli through

interaction with distinct parts of frontal cortex[16,17]. An alternative is that single neurons in BLA

branch to many different areas, such that activity can be efficiently coordinated across

distributed networks of areas. This organization fits with the BLA's role in more general aspects

of motivation and response invigoration to approach or avoid salient stimuli that are

characteristic of model-free behaviors[8].

These two potential connectional motifs of single BLA neurons – specific vs branching –

are not necessarily mutually exclusive. However, at present, the extent to which either motif

best characterizes the projections of single BLA neurons or indeed populations of BLA neurons

is not known, in part because gold-standard tract-tracing approaches are either too coarse to

detect the projection patterns of individual neurons[18,19] or because the available single-axon

tracing techniques do not scale practically to non-human primates[20]. To surmount these issues,

we optimized and refined a high-throughput sequencing approach, multiplexed analysis of

projections by sequencing (MAPseq[7]) in macaque monkeys. MAPseq uses barcoded mRNA

technology[21] to map the connections of individual neurons at scale. Because of their potential

importance in psychiatric disorders[11], we focused on projections from BLA to frontal cortex,

striatum, anterior temporal lobe, and mediodorsal nucleus of the thalamus (MD), a part of

thalamus that receives input from both frontal cortex and amygdala[1]. Using this approach, we

found that individual BLA neurons project widely in frontal cortex; about half of the neurons

that leave amygdala project to more than one target area. The pattern of these branching

projections was not random, such that the connections of single neurons were organized into

distinct and reproducible connection motifs. Notably, BLA projections to posterior parts of

frontal cortex were highly specific, whereas those to more anterior parts of frontal cortex,

especially ventral frontal cortex, were more likely to branch to multiple areas.

Results

Optimization of MAPseq in macaques

MAPseq[7] relies on an engineered sindbis virus that infects neurons with unique RNA

sequences, referred to as barcodes[21]. Following viral expression these barcodes are conjugated

to a nonfunctional presynaptic protein and undergo anterograde axonal transport. Thus, by

dissecting and sequencing samples from injection and target brain areas, the projection

patterns of single neurons can be determined. MAPseq thereby complements and extends

traditional neuroanatomical approaches, as it provides both information about bulk projection

patterns from one area to another as well as the connection patterns of single neurons.

Further, it permits simultaneous analysis of projections to many target areas within the same

animal, allowing single-neuron branching to be discerned.

We performed bilateral, MRI-guided stereotactic injections of barcoded sindbis virus

into the BLA of two rhesus macaques (**Figure 1A**, see Extended Data Figure 1 for more detail);

10 to 12 injections of 400 nl each were placed throughout the lateral, basal, and accessory basal

nuclei to control viral spread. Following perfusion, brains were extracted, the hemispheres

separated and sectioned. The BLA and target areas in frontal cortex, striatum, entorhinal

cortex, hippocampus, and MD were dissected according to gray/white matter boundaries as

well as sulcal landmarks (**Figure 1B**). Simple qPCR on extracted mRNA recovered significantly

more barcode in amygdala sites near the injection compared to target sites (OLS regression, t(2)

= 46.43, *p* < 0.0001) and more in target sites compared to control sites in cerebellum (t(2) =

4.77, p < 0.0001; **Figure 1C**). High-throughput next-generation sequencing was then conducted

on extracted mRNA from BLA and target areas. Counts of unique barcodes from the extracted

RNA were normalized and a threshold applied to control for spurious sequencing results[7]

(Extended Data Figure 2). The combined thresholded barcode counts from the four

hemispheres were then analyzed together (see Extended Data Figures 3 and 4 for analyses of

each hemisphere separately and comparisons between hemispheres, respectively).

In total, we recovered 3,115 unique barcodes in samples dissected from BLA across the

four hemispheres. This yield of barcode counts is similar in level to that recovered in mice per

injection[21–24], indicating that MAPseq[7] was working as intended in macaques. To determine the

projection targets of putative single neurons from BLA, we then determined whether each

barcode found in amygdala was present in any of the target areas after collapsing across all

samples in a target area. Approximately one-third of the unique barcodes – which can be

interpreted to represent single neurons – projected only to other sites within amygdala, which

we refer to as zero-target. Another third projected to only one of the target areas we collected

– we refer to these neurons as having *specific* projections – and the remaining third had two or

more targets outside of amygdala. We refer to these neurons with multiple targets as having

branching projections (**Figure 1D**).

The overall proportion of barcode recovered in each target mirrored known connections

of BLA in macaques (**Figure 1E**). Accordingly, the highest amount of barcode was recovered in

entorhinal cortex and nucleus accumbens (NAcc), two areas which have previously been

identified as receiving dense projections from BLA[25]. A lower amount of barcode was recovered

from hippocampus and agranular insula (AI), which are also well-documented as receiving

strong projections from BLA[4,26], followed by subcallosal (scACC) and dorsal ACC (dACC).

Notably, relatively similar amounts of barcode were recovered from all other target areas in

frontal cortex and striatum, which again matches known gradients of BLA connections across

the frontal lobe[4,27]. The close alignment between our findings and prior tract-tracing supports

the validity of MAPseq in quantifying coarse area-to-area projections in macaques.

Next, we looked at the probability that a barcode found in one target structure was also

found in another target structure (**Figure 1F**). This approach allowed us to begin to ascertain

the degree to which connections of single BLA neurons are either specific or branching. For

instance, neurons that project to NAcc have a high probability of also projecting to AI. By

contrast, single neurons that project to AI are unlikely to project to either entorhinal cortex or

hippocampus. Similar patterns of non-overlapping projections are also apparent in BLA

projections to frontal cortex; neurons that project to medial orbitofrontal cortex (mOFC) are

unlikely to project to ventrolateral prefrontal cortex (vlPFC). As can be seen in by the

representative examples in **Figure 1G**, evidence for both specific and branching projection

patterns can be observed at the level of single neurons, as well. Notably, our results correspond

closely with the findings of two studies that investigated the patterns of BLA neuron branching

to frontal cortex and thalamus. One by Sharma, Fudge and colleagues[28] identified a subset of

amygdala neurons that project to adjacent parts of medial frontal areas 25/14 (which we refer

to as scACC and mOFC, respectively) and 32/24 (which encompasses our perigenual ACC

[pgACC] and dACC, respectively). We also identified neurons that projected to both areas,

although we identified a slightly higher proportion of neurons with this branching pattern

(between 7 and 20% depending on the animal) of them than previously reported (between 7

and 10% depending on the nucleus) – likely because our dissected areas encompassed larger

portions of medial frontal cortex. A study by Timbie and Barbas[29] found that two largely non-

overlapping populations of amygdala neurons projected to either MD or posterior OFC/AI, a

result highly similar to what we found (**Figure 1F**). This lack of branching as discerned by our

barcode analysis is an important negative finding; in additional to observing the expected

pattern of *branching projections* to certain targets, we have also observed the expected pattern

of *specific projections* to other targets. Taken together the above analyses provide novel

evidence that BLA neurons often project to multiple targets in frontal cortex, arguing against

specific targeting as the dominant connectivity principle, while also reproducing the known

gross connectivity of BLA – altogether confirming the validity of MAPseq in macaques.

Figure 1: MAPseq of macaque BLA. **A**) Schematic of injection approach. **B**) Example coronal sections showing dissection targets in striatum as well as frontal and temporal lobes. Ento, entorhinal; HC, hippocampus (not shown); AI, anterior insula cortex; mOFC, medial orbitofrontal cortex; lOFC, lateral orbitofrontal cortex; vlPFC, ventrolateral prefrontal cortex; scACC, subcallosal anterior cingulate cortex; pgACC, perigenual ACC; dACC, dorsal ACC; PMd, dorsal premotor cortex; NAcc, nucleus accumbens; MD, mediodorsal thalamus (also not shown). Anterior-posterior levels are in mm relative to the interaural plane. **C**) Normalized barcode amount relative to peak barcode amount in injection site across other locations in amygdala (blue), all target structures (green; OLS regression vs amygdala: t(2) = 46.43, *p* < 0.0001), and control areas (red; vs targets: t(2) = 4.77, *p* < 0.001). **D**) Proportion of unique barcodes in no targets outside of amygdala (green), one (orange), two (blue), or more than two targets outside of amygdala (red). **E**) Proportion of barcodes in each of the target and control areas; dashed line indicates the proportion of barcode found in control sites. **F**) Conditional probability that a barcode in area A is also found in area B. Note that the order of indexing here alters the probability such that P(A|B) can be greater than P(B|A). **G**) Normalized proportion of barcodes across all the potential target areas and control areas for 10 unique barcodes (colored lines).

Quantitative analysis of branching connectional motifs of single BLA neurons

The prior analysis on the branching of neurons to two targets, while revealing, does not

capture the full set of connections that single neurons make to multiple target areas – one of

the major strengths of MAPseq over standard tract-tracing approaches (**Figure 1G**).

Consequently, we focused our next analysis on the nearly 1,300 BLA neurons with branching

projections to multiple locations in frontal cortex, temporal cortex, striatum and thalamus

(**Figure 2A**). While all target areas received the majority of their input from branching neurons,

entorhinal cortex was found to receive the highest proportion of specific input from BLA (**Figure**

2B, z-test for proportions: entorhinal vs hippocampus [next highest], z = 10.67, p < 0.0001).

Next, to determine whether any branching motifs were over- or under-represented compared

to chance, we built a null distribution based on the overall barcode distribution (**Figure 2C**).

Assuming total independence for branching, the probability that a neuron projects to both

areas A and B can be computed as the product of the independent probabilities of projecting to

area A and area B[24]. By comparing the actual counts for each motif with this null distribution,

we identified only one over-represented bifurcating motif: neurons that project to both NAcc

and AI (binomial test, p = 0.003). This motif was found alongside predominantly under-

represented bi- and trifurcations, many of which included projection motifs encompassing

some combination of NAcc, entorhinal cortex, and hippocampus (**Figure 2D and 2E**). That

branching to only two of these areas happens less frequently than expected is somewhat

surprising considering these were the three areas most likely to receive projections from BLA in

our data (**Figure 1E**); however, these data are in line with small-world theories of brain network

organization[30]. By contrast, axon branching to four different targets were far more likely to be

over-represented compared to that same chance distribution (**Figure 2F**). In other words, there

are fewer bifurcations than expected based solely on the proportions of barcodes found in each

target area, while there were more of these quadfurcations than expected. There were no

significantly over- or under-represented motifs with five or more targets. Thus, these results

indicate that single BLA neurons demonstrate a high degree of branching, being more likely to

strongly innervate four distinct targets in the target areas sampled over two or three.

Importantly, this analysis also reveals that the observed branching motifs are not simply a

product of the distribution of barcodes; rather, branching of single BLA neurons appears to be

highly structured.

Figure 2: Single neuron analysis of branching projections from BLA. A) K-means clustering of 1,300 neurons that project to more than one area outside of amygdala (k=12). **B)** Proportion of projections from BLA to frontal cortex areas, striatum, and mediodorsal thalamus that are from specific (crosses) or branching projections (filled circles). **C)** Logic of null distribution computation for branching motifs. **D)** Observed (blue) and expected (green) counts of neurons with projections to multiple areas (top). Specific over (red) or under-represented (blue) branching motifs by area (bottom). **E)** Volcano plot of probability of all possible branching motifs with 2 (cream), 3 (grey), and 4 (turquoise) target areas. Dashed line marks the level of statistical significance. **F)** Proportion of significantly over- (red) and under- (blue) represented 2-, 3- and 4- target area branching motifs.

Single neuron projection networks within frontal cortex

With the appreciation that the projections of single neurons in BLA are highly likely to

branch to multiple areas we sought to understand how these projections are organized. Here

we separately focused on the pattens of single BLA projections to the medial and ventral frontal

cortex. We took this approach because BLA projections to these areas are thought to be

functionally distinct. Interaction between BLA and medial frontal cortex is heavily linked to

defensive threat conditioning in animals and anxiety-related disorders such as PTSD in

humans[31]. Projections from BLA to ventral frontal cortex are, by contrast, more frequently

associated with reward-guided behaviors[6] and dysfunction in these circuits is linked to addictive

disorders[32].

Medial frontal cortex

We identified 405 BLA neurons that project to either medial frontal areas scACC (area

25 as defined by Carmichael and Price[33]), pgACC (area 32), or dACC (area 24) (**Figure 3A**).

Although these areas are densely interconnected[18], we found marked differences in the

structure of BLA input that they receive. First, over half of the BLA neurons that projected to

medial frontal cortex exclusively targeted scACC, whereas a third targeted only dACC (130/405;

z-test for proportions, $z = 6.38$, $p < 0.0001$); even fewer BLA projections to medial frontal cortex

were specific to pgACC (55/405; $z = 6.28$, $p < 0.0\ 001$) (**Figure 3B**). Indeed, the majority of BLA

neurons targeting pgACC also projected to the other parts of medial frontal cortex, while a

smaller proportion of dACC-projecting neurons branched within medial frontal cortex ($z = 5.37$,

$p < 0.0001$); scACC-projecting neurons were least likely to branch ($z = 3.17$, $p = 0.0015$). Thus,

within medial frontal cortex there is a hierarchy of specific vs branching BLA connections where

pgACC receives the least specific BLA input and scACC receives the most.

As we noted earlier, BLA neurons tend to have more branching than specific projections

(**Figure 2C** and **E**); within this branching, we found additional structure among projections that

targeted single medial frontal areas. Neurons that project to scACC or dACC were more likely to

have four targets than neurons which projected to pgACC, which were dominated by two- and

three- target neurons (**Figure 3C,** permutation test). These results suggest that amygdala inputs

to pgACC, while frequently shared among other cingulate areas, are not shared as frequently

outside of medial frontal cortex.

Within the projections from BLA to medial frontal cortex there were notable differences

in the areas that these single neurons also targeted, indicative of different networks. Whereas

the BLA neurons projecting to each medial area had similar proportions of bifurcations to

hippocampus and entorhinal cortex (**Figure 3D** and **E**), neurons that projected to scACC were far

more likely to also project to NAcc than the other areas (z-test for proportions: scACC vs pgACC:

$z = 5.85$, $p < 0.0001$; scACC vs dACC: $z = 6.33$, $p < 0.0001$). This projection motif is evident at the

level of individual neurons, and these cells were also highly likely to connect to AI (**Figure 3E**).

dACC-projecting BLA neurons, however, were more likely to also project to lateral OFC (lOFC;

dACC vs pgACC: $z = 2.72$, $p = 0.0097$; dACC vs scACC: $z = 3.80$, $p < 0.001$) and vlPFC (dACC vs

pgACC: $z = 2.04$, $p = 0.061$; dACC vs scACC: $z = 3.52$, $p < 0.001$) on the ventral surface of the

frontal lobe, providing further anatomical support for the role of the dACC in valuation through

its interactions with more ventral areas[6,34]. In summary, single BLA neurons targeting medial

frontal areas appeared to target distinct networks; those targeting scACC were largely

constrained to this area and primarily sent bifurcations to NAcc and AI in the posterior ventral

frontal cortex, consistent with the known roles of these areas in regulating mood[35]. By contrast,

those targeting dACC also innervated parts of ventral frontal cortex including those linked to

Carmichael and Price's visceromotor network[18], consistent with its role in value-based decision-

making[6,34].

Figure 3: BLA specific projections to medial frontal cortex. A) Schematic showing the three populations of medial frontal cortex-projecting neurons analyzed here: BLA neurons targeting scACC (blue), pgACC (red), dACC (green). D and V refer to the dorsal and ventral directions, respectively; R and C designate rostral and caudal. **B**) Venn diagram illustrating projections to each medial frontal cortex area that also branch to the other two; pgACC-projecting neurons are more likely to branch than dACC-projecting neurons (z-test for proportions, z=3.17, p=0.0015), which are more likely to branch than scACC-projecting neurons (z=2.27, p=0.023). **C**) Degrees of branching for each medial frontal cortex-projecting population; dashed bars indicate the mean of 1000 shuffles of the data, downsampled for equal numbers of neurons from each population; error bars indicate 95% confidence intervals. **D**) Likelihood of medial frontal cortex-projecting neurons projecting to non-medial frontal cortex targets. **E**) Single-neuron projection patterns; shading indicates number of targets.

Ventral Frontal Cortex Networks

We conducted a similar analysis on the 627 BLA neurons that projected to areas on the

ventral surface of frontal cortex, including AI, mOFC, lOFC, and vlPFC (**Figure 4A**). Of those BLA

neurons that projected to ventral frontal cortex, almost two thirds (65%) of BLA neurons solely

targeted AI, whereas only 10% solely targeted vlPFC; within ventral frontal cortex, these areas

received the highest and lowest proportions of specific projections, respectively (z-test for

proportions: z = 19.85, *p* < 0.0001, **Figure 4B**). Indeed, the majority of BLA neurons projecting to

vlPFC also targeted other parts of the ventral frontal cortex and were not specific to this area.

BLA neurons projecting to mOFC and lOFC had similar proportions of cells projecting in a

specific or branching manner. Thus, BLA projections to AI are more specific compared to those

directed to more anterior regions (AI vs mOFC: z = 4.43, *p* < 0.0001), with vlPFC receiving the

fewest specific projections from BLA (vlPFC vs mOFC: z = 2.48, *p* = 0.013).

Similar to medial frontal cortex, BLA neurons projecting to ventral frontal cortex were

more likely to send branching as opposed to specific projections. AI-, lOFC-, and vlPFC-

projecting neurons were most likely to send 2-target projections (**Figure 4C**, permutation test),

whereas mOFC-projecting neurons tended to innervate either one or four targets. AI- and

vlPFC-projecting neurons were least likely to branch to four distinct targets, suggesting that

these two areas receive the most highly specialized input from BLA.

When we limited our analyses to BLA neurons that only projected to one of the four

ventral frontal areas without branching between them, BLA neurons that project to AI exhibited

the strongest projections to NAcc compared to other areas in ventral frontal cortex (**Figure 4C**

and **E;** Fisher's exact test: AI vs mOFC, *p* = 0.0057; AI vs lOFC, *p* = 0.00019; AI vs vlPFC, *p* <

0.0001). By contrast, vlPFC- and lOFC- projecting neurons were most likely to also project to

entorhinal cortex compared to mOFC and AI (**Figure 4C;** lOFC vs mOFC, *p* = 0.043; lOFC vs AI, *p* <

0.0001; vlPFC vs AI, *p* = 0.00051). Somewhat unexpectedly, BLA neurons with projections to

mOFC were more likely to project to dorsal premotor cortex (PMd; mOFC vs AI, *p* < 0.00001;

mOFC vs lOFC, *p* = 0.028) and other more medial areas of the frontal lobe such as the pgACC

(**Figure 4C** and **E;** mOFC vs AI, *p* = 0.019; mOFC vs lOFC, *p* = 0.038) than were BLA neurons

projecting to other ventral frontal cortex regions. These patterns of projections indicate the BLA

neurons targeting the ventral frontal cortex form highly-structured networks; those targeting AI

are less likely also to project to other ventral frontal areas. When they do branch to other areas,

they primarily innervate other areas in the posterior frontal lobe (i.e. scACC) and NAcc. By

contrast, BLA projections to more anterior parts of ventral frontal cortex, especially vlPFC,

branch more both within frontal cortex and also to other areas. This finding that vlPFC-

projecting neurons connect so broadly aligns closely with the known role of the BLA-vlPFC

circuit in representing and updating model-free stimulus-outcome associations[36], information

that is likely to be shared among other ventral frontal areas to inform reward-guided choice

behavior[37].

Taken together, the patterns of projections strongly indicate the existence of BLA

neurons that target distinct networks in the medial and ventral frontal cortex. Those BLA

neurons that target scACC and AI in the posterior frontal cortex preferentially innervate NAcc.

By contrast, BLA neurons that target to more anterior medial and ventral frontal areas appear

to be part of more distributed networks of areas. This extends our understanding of the

organization of BLA-to-frontal cortex networks revealing distinct connectional motifs that may

in part explain the pervasive influence of BLA on frontal cortex processing.

Figure 4: BLA specific projections to ventral frontal cortex. A) Schematic showing the four populations of ventral frontal cortex-projecting neurons analyzed here: BLA neurons targeting AI (yellow), mOFC (blue), lOFC (red), and vlPFC (green). A and P designate anterior and posterior directions, respectively; M and L refer to medial and lateral. **B**) Venn diagram illustrating projections to each area that also branch to the other three; AI-projecting neurons were least likely to branch to the other areas (z-test for proportions, z=4.425, p<0.0001), while vlPFC-projecting neurons were most likely to branch within ventral FC (z=2.476, p=0.0133).
C) Degrees of branching for each ventral frontal cortex-projecting population; dashed bars indicate the mean of 1000 shuffles of the data, downsampled for equal numbers of neurons from each population; error bars indicate 95% confidence intervals. **D**) Likelihood of ventral frontal cortex-projecting neurons projecting to non-ventral frontal cortex targets. **E**) Single-neuron projection patterns; shading indicates number of targets.

Retrograde tracing validation of specific and branching BLA projections to ventral frontal cortex

Next, we sought to assess the most distinct BLA projection patterns revealed by MAPseq using standard retrograde viral tracing. In a single macaque, MRI-guided stereotactic injections of retro-AAV2 coding for mCherry and EGFP fluorophores were injected into NAcc and a lateral subregion of AI in one hemisphere. Injections were targeted to posterior-lateral lOFC (area 13m as defined by Carmichael and Price[33]) and posterior-medial vlPFC (area 12o) in the other hemisphere (**Extended Data Figure 5A and B**). As cross-hemispheric BLA connections are negligible[27], we were able to analyze the two hemispheres separately. We then conducted unbiased stereological counting of neurons in BLA that were either single or double labeled with each fluorophore in each hemisphere[38]. Thus, we were able to compare the connectivity profiles of single amygdala neurons found in specific subregions of frontal cortex.

For BLA neurons projecting to AI, we found a high degree of correspondence in the proportion of branching neurons between stereology and MAPseq estimates (z-test for proportions $p > 0.05$, Extended Data Figure 5C); for NAcc-projecting neurons, however, our MAPseq and stereology results did not agree, as stereological estimates of specific projections were higher than MAPseq ($p < 0.0001$). For lOFC and vlPFC, too, the numbers of BLA neurons that projected specifically to one or branched to both areas was different to what we found with MAPseq ($p < 0.001$, Extended Data Figure 5D). This was primarily because stereological estimates of BLA projections to vlPFC almost entirely overlapped with those targeting lOFC. The difference in estimates of specific versus branching BLA projections between retrograde tracing and MAPseq results is not unexpected[22]. This is because the injections of retro-AAV into lOFC and vlPFC do not cover the full extent of these cortical regions. By contrast, MAPseq estimates of specific and branching projections are based on BLA projections to the whole of lOFC and vlPFC that extend far beyond the extent of the areas target with retro-AAV. The difference in estimates of the branching and specific projections between techniques is, however, revealing as it indicates that projections to this subregion of area 12 are even less specific than the analysis of connectivity of the larger vlPFC would indicate.

Branching and specific projections shape fMRI functional connectivity

Anatomical connections in the brain constrain functional networks[39,40]. Given the

reproducible patterns of specific and branching projections from BLA that we identified, we

sought to determine if multi-area connectional motifs identified by MAPseq could be identified

at the level of fMRI functional connectivity (FC). If such patterns were identifiable in FC this

would indicate that the unique anatomical networks identified with MAPseq are related to and

in fact shape functional networks in the brain. We analyzed a dataset of resting state fMRI

scans from six rhesus monkeys[41] (**Figure 5A**), with single seed voxels in amygdala and target

voxels in the same areas used in the MAPseq experiments (Extended Data Figure 6). After

computing and z-transforming FC values for each amygdala voxel with each target voxel, we

binarized the FC signal by setting a threshold at 70% of each voxels' maximum connectivity.

Practically, this meant that if FC between an amygdala voxel and any target area was above the

threshold, we counted that amygdala voxel as being *functionally connected* to that target area.

We then counted the number of connections to each target area and normalized those counts

by the size of each target area. This approach meant that single amygdala voxels could be

designated as being functionally connected to multiple parts of frontal cortex, striatum,

temporal lobe and thalamus, following a similar approach to MAPseq.

Despite fMRI-derived FC operating over a much larger spatial scale than standard

anatomical approaches and having no directional sensitivity, we were still able to identify a

largely similar pattern of relative FC between amygdala and our target areas as we had found

with MAPseq (compare **Figures 1E** and **5C**). Notably, the pattern also generally matches prior

tract-tracing results as there was strong connectivity between amygdala and entorhinal cortex,

hippocampus, NAcc, AI, scACC as well as MD with lower connectivity to lOFC and pgACC[42]. For a

number of areas, functional connectivity with amygdala was higher than expected based on

MAPseq and tract-tracing data, most notably PMd and mOFC (**Figure 5C**). This result is not

necessarily surprising because FC measures not only reflect unidirectional connections from

one area to another; rather, strong functional connection between two areas can be driven by

strong reciprocal connections or strong coupling with a third area with which both areas

strongly interact[43].

Next to look for patterns of network-level connectivity we assessed the FC of amygdala

voxels with high connectivity to areas in medial frontal cortex (**Fig. 5C**) or ventral frontal cortex

(**Fig. 5D**). This is essentially the same analysis depicted in **Figures 3E** and **4E**. Notably, we were

able to identify many of the prominent connectivity motifs that we observed with MAPseq. For

instance, voxels with a high likelihood of being connected to AI neurons also had higher

connectivity with NAcc (**Figure 5D**). Similarly, dACC-connected neurons also had a high

likelihood of being connected to vlPFC (**Figure 5C**) and if a voxel showed high FC with any of the

medial and ventral frontal cortex areas it had uniformly low connectivity with MD (**Figures 5C**

and **D**). This latter point demonstrates the correspondence between MAPseq, tract-tracing and

fMRI[42].

Despite being able to discern these broad connectional motifs, some key differences

between medial and ventral areas were obscured, such that fMRI estimates of multi-region

connectivity estimated highly similar connectivity profiles for all three medial frontal cortex

areas. This difference between fMRI and MAPseq analyses is likely explained by the dense and

bidirectional interconnections within medial frontal cortex[18], inputs from other structures

received by this part of frontal cortex most notably from hippocampus and neuromodulatory

systems[27,44], and the low resolution of fMRI compared to individual neurons. Taken together,

that major anatomical features identified by MAPseq were also identifiable at the level of fMRI

FC likely reflects their significance in shaping brain-wide activity patterns organized into

functional networks.

Figure 5: MRI functional connectivity replicates some anatomical features. A) Surface projection of right amygdala ROI functional connectivity to right hemisphere targets, averaged across six animals. Compass directions refer to dorsal (D)/ventral (V) and posterior (P)/anterior (A) directions; sulci labelled are: principal sulcus (ps), lateral orbital sulcus (los), arcuate sulcus (arc), superior temporal sulcus (sts), lunate sulcus (ls). **B**) Functional connectivity strength across areas. Colored bars indicate number of amygdala voxels functionally connected to each target area, corrected for the number of voxels in each target area; dotted bars with 95% confidence intervals indicate random samples of the same number of voxels from each area as the smallest target area (scACC). **C**) Likelihood of medial FC-projecting voxels being functionally connected to non-medial FC targets. **D**) Likelihood of ventral FC-projecting voxels being functionally connected to non-ventral FC targets.

318   Discussion

We have optimized and validated MAPseq for single-neuron connectivity mapping in

macaque monkeys, opening new avenues of research into the single-cell structural connections

of the non-human primate brain. Using this approach, we successfully determined the

projection patterns of over 3,000 single neurons from BLA to frontal cortex, striatum, parts of

the temporal lobe, and MD. Notably, the bulk patterns of connections identified here are in

close alignment with previous reports of BLA projections based on traditional tract-tracing

techniques[45,1,2,25,3,26,29] (**Figures 1 and 2**). Overall, we found that single BLA neurons branch

extensively, with branching to four distinct targets being more likely to be over-represented

compared to chance than branching to two or three targets (**Figure 2**). Within these patterns of

branching, we identified distinct connectional motifs; BLA projections to posterior parts of the

medial and ventral frontal cortex were highly specific and less likely to branch to other areas. By

contrast, projections to more anterior or dorsal areas were much more likely to branch to other

brain areas and exhibited unique projection profiles (**Figures 3 and 4**). We also identified broad

similarities and slight differences between this anatomical connectivity and more standard

tract-tracing techniques (**Extended Data Figure 5**) and functional connectivity identified by fMRI

(**Figure 5**). These findings begin to reveal the unique patterns of projections of single BLA

neurons, connections that are heavily implicated in the control of affect and that become

dysfunctional in psychiatric disorders[11].

It is well documented that BLA input to frontal cortex is strongest to posterior regions

and weaker to more anterior regions[2,4]. On top of this, we found that in both medial and

ventral frontal cortex, BLA neuron branching is highest to the most anterior areas. For instance,

BLA neurons projecting to pgACC showed the highest degree of branching to other areas in

medial frontal cortex, whereas neurons projecting to the more posterior scACC branched the

least (**Figure 3B**). A generally similar pattern was seen in ventral frontal cortex with the

exception of vlPFC, which received the highest degree of branching projections (**Figure 4**). This

lack of specific BLA projections to vlPFC may be related to its role in model-free as opposed to

model-based behaviors; lesions of vlPFC do not impact reinforcer devaluation that depends on

specific sensory information about an outcome but do impact outcome-independent

probabilistic learning[36]. Thus, the patterns of specific and branching connections from single

BLA neurons may serve a functional role in relaying distinct sensory information or providing a

salience signal to invigorate responding, respectively. In medial frontal cortex, while so-called

ventromedial PFC has been generally implicated in affective regulation, and by extension

anxiety and mood disorders[46], more recent evidence suggests that particular subregions may

play distinct roles in affect[47–49]. The highly specific and segregated inputs to scACC and pgACC

identified here potentially provide a neuroanatomical basis for their opposing roles in affective

responding[35].

Appreciating the unique features of BLA projections to frontal cortex is potentially

critical for understanding the basis of a number of amygdala-linked psychiatric disorders[11]. For

example, obsessive-compulsive disorder is associated with dysfunction in basal ganglia-OFC

circuitry regulating valuation as well as basal ganglia-dACC circuits involved in action selection[50].

Here we identified a population of dACC-projecting BLA neurons that also preferentially

targeted parts of ventral frontal cortex including lOFC (**Figure 3**). Our findings therefore provide

a potential anatomical basis through which dysfunction in a single, small population of BLA

neurons could influence a distributed network of areas. Determining the distinct functions of

the different BLA projection motifs identified here as well as their molecular signatures[23,51] has

the potential to bring network-level understanding to basic and translational neuroscience and

might provide a more biologically-realistic basis for the construction of neural network

architectures[52].

References

1. Aggleton, J. P. & Mishkin, M. Projections of the amygdala to the thalamus in the cynomolgus monkey. *J. Comp. Neurol.* **222**, 56–68 (1984).

2. Amaral, D. G. & Price, J. L. Amygdalo-cortical projections in the monkey (Macaca fascicularis). *Journal of Comparative Neurology* **230**, 465–496 (1984).

3. Ghashghaei, H. T. & Barbas, H. Pathways for emotion: interactions of prefrontal and anterior temporal pathways in the amygdala of the rhesus monkey. *Neuroscience* **115**, 1261–1279 (2002).

4. Ghashghaei, H. T., Hilgetag, C. C. & Barbas, H. Sequence of information processing for emotions based on the anatomic dialogue between prefrontal cortex and amygdala. *NeuroImage* **34**, 905–923 (2007).

5. Bucy, P. C. & Klüver, H. An anatomical investigation of the temporal lobe in the monkey (Macaca mulatta). *Journal of Comparative Neurology* **103**, 151–251 (1955).

6. Rudebeck, P. H., Ripple, J. A., Mitz, A. R., Averbeck, B. B. & Murray, E. A. Amygdala Contributions to Stimulus–Reward Encoding in the Macaque Medial and Orbital Frontal Cortex during Learning. *J. Neurosci.* **37**, 2186–2202 (2017).

7. Kebschull, J. M. *et al.* High-Throughput Mapping of Single-Neuron Projections by Sequencing of Barcoded RNA. *Neuron* **91**, 975–987 (2016).

8. Davis, M. & Whalen, P. J. The amygdala: vigilance and emotion. *Mol Psychiatry* **6**, 13–34 (2001).

9. Horvath, F. E. Effects of basolateral amygdalectomy on three types of avoidance behavior in cats. *Journal of Comparative and Physiological Psychology* **56**, 380–389 (1963).

10. Sengupta, A. *et al.* Basolateral Amygdala Neurons Maintain Aversive Emotional Salience. *J. Neurosci.* **38**, 3001–3012 (2018).

11. Schumann, C. M., Bauman, M. D. & Amaral, D. G. Abnormal structure or function of the amygdala is a common component of neurodevelopmental disorders. *Neuropsychologia* **49**, 745–759 (2011).

12. Brady, R. O. *et al.* State dependent cortico-amygdala circuit dysfunction in bipolar disorder. *Journal of Affective Disorders* **201**, 79–87 (2016).

13. Marusak, H. A. *et al.* You say 'prefrontal cortex' and I say 'anterior cingulate': meta-analysis
of spatial overlap in amygdala-to-prefrontal connectivity and internalizing symptomology.
Transl Psychiatry **6**, e944–e944 (2016).

14. Yin, S., Liu, Y., Petro, N. M., Keil, A. & Ding, M. Amygdala Adaptation and Temporal
Dynamics of the Salience Network in Conditioned Fear: A Single-Trial fMRI Study. *eNeuro* **5**,
(2018).

15. Murray, E. A. & Fellows, L. K. Prefrontal cortex interactions with the amygdala in primates.
Neuropsychopharmacol. **47**, 163–179 (2022).

16. Johnson, A. W., Gallagher, M. & Holland, P. C. The Basolateral Amygdala Is Critical to the
Expression of Pavlovian and Instrumental Outcome-Specific Reinforcer Devaluation Effects.
J. Neurosci. **29**, 696–704 (2009).

17. Rhodes, S. E. V. & Murray, E. A. Differential Effects of Amygdala, Orbital Prefrontal Cortex,
and Prelimbic Cortex Lesions on Goal-Directed Behavior in Rhesus Macaques. *J. Neurosci.*
33, 3380–3389 (2013).

18. Carmichael, S. T. & Price, J. L. Connectional networks within the orbital and medial
prefrontal cortex of macaque monkeys. *Journal of Comparative Neurology* **371**, 179–207
(1996).

19. Rockland, K. S. Axon Collaterals and Brain States. *Front. Syst. Neurosci.* **12**, (2018).

20. Rockland, K. S. What we can learn from the complex architecture of single axons. *Brain*
Struct Funct **225**, 1327–1347 (2020).

21. Kebschull, J. M. & Zador, A. M. Cellular barcoding: lineage tracing, screening and beyond.
Nat Methods **15**, 871–879 (2018).

22. Han, Y. *et al.* The logic of single-cell projections from visual cortex. *Nature* **556**, 51–56
(2018).

23. Chen, X. *et al.* High-Throughput Mapping of Long-Range Neuronal Projection Using In Situ
Sequencing. *Cell* **179**, 772-786.e19 (2019).

24. Gergues, M. M. *et al.* Circuit and molecular architecture of a ventral hippocampal network.
Nature Neuroscience 1–9 (2020) doi:10.1038/s41593-020-0705-8.

25. Russchen, F. T., Bakst, I., Amaral, D. G. & Price, J. L. The amygdalostriatal projections in the
monkey. An anterograde tracing study. *Brain Research* **329**, 241–257 (1985).

26. Höistad, M. & Barbas, H. Sequence of information processing for emotions through
pathways linking temporal and insular cortices with the amygdala. *Neuroimage* **40**, 1016–
1033 (2008).

27. Aggleton, J. P., Wright, N. F., Rosene, D. L. & Saunders, R. C. Complementary Patterns of
Direct Amygdala and Hippocampal Projections to the Macaque Prefrontal Cortex. *Cereb*
Cortex **25**, 4351–4373 (2015).

28. Sharma, K. K., Kelly, E. A., Pfeifer, C. W. & Fudge, J. L. Translating Fear Circuitry: Amygdala
Projections to Subgenual and Perigenual Anterior Cingulate in the Macaque. *Cerebral*
Cortex bhz106 (2019) doi:10.1093/cercor/bhz106.

29. Timbie, C. & Barbas, H. Pathways for Emotions: Specializations in the Amygdalar,
Mediodorsal Thalamic, and Posterior Orbitofrontal Network. *J. Neurosci.* **35**, 11976–11987
(2015).

30. Bassett, D. S. & Bullmore, E. T. Small-World Brain Networks Revisited. *Neuroscientist* **23**,
499–516 (2017).

31. Park, J. & Moghaddam, B. Impact of anxiety on prefrontal cortex encoding of cognitive
flexibility. *Neuroscience* **345**, 193–202 (2017).

32. Goldstein, R. Z. & Volkow, N. D. Drug Addiction and Its Underlying Neurobiological Basis:
Neuroimaging Evidence for the Involvement of the Frontal Cortex. *AJP* **159**, 1642–1652
(2002).

33. Carmichael, S. T. & Price, J. L. Architectonic subdivision of the orbital and medial prefrontal
cortex in the macaque monkey. *Journal of Comparative Neurology* **346**, 366–402 (1994).

34. Monosov, I. E. & Rushworth, M. F. S. Interactions between ventrolateral prefrontal and
anterior cingulate cortex during learning and behavioural change. *Neuropsychopharmacol.*
47, 196–210 (2022).

35. Wallis, C. U., Cardinal, R. N., Alexander, L., Roberts, A. C. & Clarke, H. F. Opposing roles of
primate areas 25 and 32 and their putative rodent homologs in the regulation of negative
emotion. *PNAS* **114**, E4075–E4084 (2017).

36. Rudebeck, P. H., Saunders, R. C., Lundgren, D. A. & Murray, E. A. Specialized

Representations of Value in the Orbital and Ventrolateral Prefrontal Cortex: Desirability

versus Availability of Outcomes. *Neuron* **95**, 1208-1220.e5 (2017).

37. Padoa-Schioppa, C. & Conen, K. E. Orbitofrontal Cortex: A Neural Circuit for Economic

Decisions. *Neuron* **96**, 736–754 (2017).

38. West, M. J., Slomianka, L. & Gundersen, H. J. G. Unbiased stereological estimation of the

total number of neurons in the subdivisions of the rat hippocampus using the optical

fractionator. *The Anatomical Record* **231**, 482–497 (1991).

39. Haber, S. N., Liu, H., Seidlitz, J. & Bullmore, E. Prefrontal connectomics: from anatomy to

human imaging. *Neuropsychopharmacol.* 1–21 (2021) doi:10.1038/s41386-021-01156-6.

40. Trambaiolli, L. R. *et al.* Anatomical and functional connectivity support the existence of a

salience network node within the caudal ventrolateral prefrontal cortex. *eLife* **11**, e76334

(2022).

41. Fujimoto, A. *et al.* Resting-State fMRI-Based Screening of Deschloroclozapine in Rhesus

Macaques Predicts Dosage-Dependent Behavioral Effects. *J. Neurosci.* **42**, 5705–5716

(2022).

42. Timbie, C., García-Cabezas, M. Á., Zikopoulos, B. & Barbas, H. Organization of primate

amygdalar–thalamic pathways for emotions. *PLoS Biol* **18**, e3000639 (2020).

43. van den Heuvel, M. P. & Hulshoff Pol, H. E. Exploring the brain network: A review on

resting-state fMRI functional connectivity. *European Neuropsychopharmacology* **20**, 519–

534 (2010).

44. Ghashghaei, H. T. & Barbas, H. Neural interaction between the basal forebrain and

functionally distinct prefrontal cortices in the rhesus monkey. *Neuroscience* **103**, 593–614

(2001).

45. Avendan˜o, C., Price, J. L. & Amaral, D. G. Evidence for an amygdaloid projection to

premotor cortex but not to motor cortex in the monkey. *Brain Research* **264**, 111–117

(1983).

46. Hiser, J. & Koenigs, M. The multifaceted role of ventromedial prefrontal cortex in emotion,

decision-making, social cognition, and psychopathology. *Biol Psychiatry* **83**, 638–647 (2018).

47. Myers-Schulz, B. & Koenigs, M. Functional anatomy of ventromedial prefrontal cortex:

implications for mood and anxiety disorders. *Mol Psychiatry* **17**, 132–141 (2012).

48. Alexander, L. *et al.* Over-activation of primate subgenual cingulate cortex enhances the

cardiovascular, behavioral and neural responses to threat. *Nature Communications* **11**,

5386 (2020).

49. Alexander, L., Wood, C. M. & Roberts, A. C. The ventromedial prefrontal cortex and emotion

regulation: lost in translation? *The Journal of Physiology* **n/a**,.

50. Ahmari, S. E. & Dougherty, D. D. Dissecting Ocd Circuits: From Animal Models to Targeted

Treatments. *Depression and Anxiety* **32**, 550–562 (2015).

51. Huang, L. *et al.* BRICseq Bridges Brain-wide Interregional Connectivity to Neural Activity and

Gene Expression in Single Animals. *Cell* **182**, 177-188.e27 (2020).

52. Zador, A. M. A critique of pure learning and what artificial neural networks can learn from

animal brains. *Nature Communications* **10**, 3770 (2019).

53. Cox, R. W. AFNI: Software for Analysis and Visualization of Functional Magnetic Resonance

Neuroimages. *Computers and Biomedical Research* **29**, 162–173 (1996).

54. Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image

analysis. *Nat Methods* **9**, 671–675 (2012).

55. Saunders, R. C., Aigner, T. G. & Frank, J. A. Magnetic resonance imaging of the rhesus

monkey brain: use for stereotactic neurosurgery. *Exp Brain Res* **81**, 443–446 (1990).

56. Harris, C. R. *et al.* Array programming with NumPy. *Nature* **585**, 357–362 (2020).

57. Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *Journal of Machine Learning*

Research **12**, 2825–2830 (2011).

58. Hunter, J. D. Matplotlib: A 2D Graphics Environment. *Computing in Science & Engineering* **9**,

90–95 (2007).

59. Waskom, M. L. seaborn: statistical data visualization. *Journal of Open Source Software* **6**,

3021 (2021).

60. Reveley, C. *et al.* Three-Dimensional Digital Template Atlas of the Macaque Brain. *Cereb*

Cortex **27**, 4463–4477 (2017).

61. Jung, B. *et al.* A comprehensive macaque fMRI pipeline and hierarchical atlas. *NeuroImage*

235, 117997 (2021).

62. Saleem, K. S. *et al.* High-resolution mapping and digital atlas of subcortical regions in the

macaque monkey based on matched MAP-MRI and histology. *NeuroImage* **245**, 118759

(2021).

63. Hartig, R. *et al.* The Subcortical Atlas of the Rhesus Macaque (SARM) for neuroimaging.

NeuroImage **235**, 117996 (2021).

64. Etzel, J. A., Zacks, J. M. & Braver, T. S. Searchlight analysis: Promise, pitfalls, and potential.

NeuroImage **78**, 261–269 (2013).

Methods

Subjects

Two male rhesus monkeys (*Macaca mulatta*) and one male long-tailed macaque (*Macaca fascicularis*), 8-9 years of age, were used for our experiments, weighing between 10 and 15 kg. The two rhesus macaques were used in MAPseq experiments whereas the long-tailed macaque was used in retro-AAV tracing experiments. Animals were housed individually and kept on a 12-hour light/dark cycle. Food was provided daily with water *ad libitum*. Environmental enrichment was provided daily, in the form of play objects or small food items. All procedures were approved by the Icahn School of Medicine IACUC and were carried out in accordance with NIH standards for work involving non-human primates.

Virus prep

Modified sindbis virus for MAPseq was obtained from the MAPseq Core Facility at Cold Spring Harbor Laboratory[7]. The viral library used in this study had a diversity of 20,000,000 unique barcodes. Retro-AAV2 coding for mCherry (pAAV2-hSyn-mCherry, Addgene #114472, 2 x 10^{13} GC/ml) and green fluorescent protein (pAAV2-hSyn-eGFP, Addgene #50465, 2.2 x 10^{13} GC/ml) under the human synapsin promoter were obtained from Addgene. All viruses were stored at -80°C and aliquots were thawed over wet ice immediately prior to injection.

Surgery and perfusion

Sindbis virus injections

For each animal, T1-weighted MRIs were obtained on a Skyra 3T scanner (Siemens) for surgical targeting. Animals were anesthetized with 5 mg/kg ketamine and 0.015 mg/kg dexmedetomidine; anesthesia was maintained with isoflurane as needed for the duration of the scan. Animals were scanned using an MRI-compatible stereotaxic frame (Jerry-Rig, Inc.). 3-4 images were obtained per scan, which were subsequently averaged together using the AFNI software suite (NIH)[53]. Then, stereotactic coordinates for the BLA could be computed using ImageJ (NIH)[54]. Two injection tracks were planned to target the middle of the amygdala's

anterior/posterior extent, equally spaced in the medial/lateral plane. One anterior injection

track was also planned 1.5 mm in front of the middle of the amygdala, centered in the

medial/lateral plane. Within each track, 4-5 injections 1.5 mm apart were planned in the

dorsal/ventral plane to cover the entire extent of the basal and accessory amygdala nuclei.

After allowing approximately one week to elapse after the MRI, anesthesia was induced

using ketamine and dexmedetomidine and maintained with isoflurane as described above.

Surgery was performed under aseptic conditions, using the toothmarker method[55] to place the

animals in the same 3D position as the MRI. The skin, fascia, and muscles were retracted, and

holes were drilled in the skull at each injection location using a surgical drill, widened if

necessary, using rongeurs. Small dural incisions permitted Hamilton syringes access to the brain

surface. 0.4 μl of virus was delivered at each injection site at a rate of 0.2 μl per minute, after

which the needle was left in place for at least 3 minutes before reaching for the next site.

Injections proceeded from the deepest site to the most superficial. The post-injection wait

period was extended to at least 5 minutes at the top of each injection track before removing

needles from the brain and proceeding to the next track.

After all injections were completed, the muscles, fascia, and skin were closed in

anatomical layers. Following surgery, the animal was closely monitored in his home cage until

normal behavior resumed. Postoperative treatment included buprenorphine (0.01 mg/kg, i.m.,

every 8 h) and meloxicam (0.2 mg/kg, i.m., every 24 h), based on attending veterinary guidance,

as well as cefazolin (25 mg/kg, i.m., every 24 h) and dexamethasone sodium phosphate (0.4 –1

mg/kg, every 12–24 h) on a descending dose schedule.

Perfusion took place 67-72 hours after the final injection series. After being terminally

anesthetized with ketamine/dexmedetomidine, the animal was prepared for RNAse-free

perfusion. All tools were cleaned with RNAseZap solution (Fisher), and all solutions were

prepared using RNAse-free reagents. The animal was perfused transcardially with ice-cold 1%

paraformaldehyde (PFA; Electron Microscopy Science) in phosphate-buffered saline (PBS;

Invitrogen) for approximately two minutes, followed by 4% PFA in PBS for approximately 18

minutes. Breathing was supplemented by manual ventilation until access to the heart was
obtained.

Following brain extraction, the brain was placed in 4% PFA briefly before dissection and
blocking as follows. After the cerebellum was removed, the brain was separated into
hemispheres, the temporal lobes were dissected, and the remaining brain was cut into two
blocks using a cryostat blade: one coronal cut was performed posterior to the central sulcus,
separating the frontal and anterior parietal lobes from the remaining caudal portions of the
brain – ensuring that the thalamus remained in the anterior block. Brain blocks were then post-
fixed in 4% PFA for 48 hours. After post-fix, blocks were frozen slowly over dry ice before being
stored at -80°C until sectioning.

Retro-AAV2 injections

Using the same MRI-guided stereotactic approach as above we targeted retro-AAV
injections to the NAcc and AI in the left hemisphere and lOFC and vlPFC in the right hemisphere
of a single monkey. Again, prior to surgery the animal was anesthetized, scanned at 3 T to
obtain structural images, and injection targets were planned based on these scans. On the day
of surgery, the animal was deeply anesthetized and injections were made during an aseptic
neurosurgery. The skin, fascia, and muscles were opened in anatomical layers, burr holes were
drilled over target locations and the dura opened. Injection syringes were then lowered into the
brain and injections of retro-AAV2 were then made into the targets. At each location we
injected 10 μl of virus, at a rate of 0.5 μl per minute. At the conclusion of each set of injections
the needle was left in place for 5 minutes to allow the virus to diffuse. After all injections were
completed, the muscles, fascia, and skin were closed in anatomical layers. Following surgery,
the animal was closely monitored in their home cage until normal behavior resumed.
Postoperative treatment included buprenorphine (0.01 mg/kg, i.m., every 8 h) and meloxicam
(0.2 mg/kg, i.m., every 24 h), based on attending veterinary guidance, as well as cefazolin (25
mg/kg, i.m., every 24 h) and dexamethasone sodium phosphate (0.4 –1 mg/kg, every 12–24 h)
on a descending dose schedule.

Eight weeks after surgery, the animals were perfused transcardially with ice-cold 1% PFA

(Electron Microscopy Sciences) in PBS (Invitrogen) for approximately two minutes, followed by

4% PFA in PBS for approximately 18 minutes. The brains were then extracted, postfixed for 24 h

in 4% PFA at 4°C and cryoprotected in 10% glycerol in PBS for 24 h, followed by 20% glycerol in

PBS for another 24 h. The brains were then blocked with one coronal cut posterior to the

thalamus, and the two resulting blocks were frozen in isopentane before storage at -80°C. The

brain was then sectioned in the coronal plane on a freezing stage sliding microtome (Leica) at

40 μm in a 1:8 series. Tissue sets were stored either in PBS with 0.1% sodium azide (Sigma) at

4°C or in cryoprotectant comprised of glycerol, ethylene glycol, PBS, and distilled water

(30/30/10/30 v/v/v/v, respectively) at -80°C.

Sectioning and dissection

From brains injected with sindbis virus, tissue was sectioned at 200 μm on a Leica 3050S

cryostat that had been cleaned with RNAseZap prior to use. Sections were collected over dry ice

and stored at -80°C prior to dissection. Cortical areas were then dissected according to sulcal

landmarks over dry ice. The areas that were collected and our operational definitions of their

boundaries can be found in this table.

Brain Region	Abbreviation	Brodmann's Areas	A/P boundaries	M/L boundaries
Amygdala	n/a		All nuclei across entire A/P extent	
Entorhinal cortex	ento		From the anterior tip to a point adjacent to the mid-hippocampus.	
Hippocampus	HC		Anterior tip to the middle of hippocampus	
Medial orbitofrontal cortex	mOFC	11m, 13a/b, 14	From emergence of medial orbital sulcus to its disappearance	Between rostral sulcus and medial orbital sulcus

Lateral orbito-frontal cortex	lOFC	11l, 12m/r, 13m/l	From emergence of lateral orbital sulcus to its disappearance	Between medial and lateral orbital sulci
Ventrolateral prefrontal cortex	vlPFC	45, 12o, 12l	From emergence of lateral orbital sulcus to its disappearance	Between lateral orbital sulcus and midpoint of laterally extending white matter
Dorsal anterior cingulate cortex	dACC	24	From emergence of cingulate sulcus until area 25 disappears from medial wall/emergence of septum.	Dorsal and ventral banks of cingulate sulcus
Subcallosal anterior cingulate cortex	scACC	25		Medial surface, ventral to corpus callosum, medial/dorsal to medial orbital sulcus
Perigenual anterior cingulate cortex	pgACC	32/24	Anterior to corpus callosum	Cortex on medial surface of the frontal lobe between the rostral sulcus and the cingulate sulcus
Dorsal premotor cortex	PMd	6	Posterior to emergence of arcuate sulcus	Between arcuate and central sulci on dorsal surface
Caudate	n/a			
Putamen	n/a			
Nucleus Accumbens	NAcc		Posterior to emergence of circular sulcus,	ventromedial portion of striatum

			forming dorsal aspect of insula	
Agranular Insula	AI		Orbital surface after disappearance of medial and lateral orbital sulci	From medial corner of orbital surface to where fundus of lateral sulcus used to be
Insula	n/a		From anterior commissure meeting midline	Only cortical surface in depth of lateral fissure
Medial dorsal thalamus	MD		Begins at mid-hippocampus level	Midline nuclei dorsal to midbrain

Samples from each area were combined across 3 sections in the anterior/posterior plane into

1.5-ml Eppendorf tubes, which were stored at -80°C prior to shipping frozen on dry ice for

sequencing.

mRNA sequencing and preprocessing

Sequencing of MAPseq projections was performed by the MAPseq Core Facility at CSHL

as described in Kebschull et al. 2016[7]. Briefly, sections were homogenized and treated with

protease before RNA extraction. Total RNA was extracted using an established Trizol-based

protocol. RNA quality was verified on Bioanalyzer and bulk amounts of barcodes were

examined by qPCR and compared with the amount of housekeeping gene prior to sequencing.

Barcode RNA was reverse transcribed into cDNA, and a known amount of RNA (spike-in

sequence) was added to each sample during reverse transcription. Barcode cDNA was then

double-stranded, PCR amplified to produce a sequencing library, and the purified barcode

library was then submitted for a paired-end36 run on an Illumina NextSeq machine.

Preprocessing was performed by CSHL using MATLAB (Mathworks) to create a barcode

matrix with size (n samples x n barcodes). Corresponding barcode sequences and spike-in

counts are also extracted to allow for normalization and assessment of duplicates.

Filtering and analysis

Filtering was performed in Python 3.9[56–59] using a modification of the publicly available

normBCmat.m script[7]; specific analyses can be found in the code available on Github. Briefly,

for each animal, barcodes are filtered, then the raw barcode counts are divided by the spike-in

counts for normalization; barcodes survive filtering if the max barcode amount is found in the

injection sites (amygdala), the max count is greater than 20, and if any other sample (whether

within amygdala or outside) has greater than 5 barcode counts. This adaptation from the

original filtering allows barcodes which send their strongest projections only within amygdala to

survive filtering.

The resulting barcode matrix was then binarized and collapsed within brain regions: if a

barcode was found in any of the multiple samples for each brain area, then we counted that

neuron as projecting to that brain area. This collapsing and binarization allowed us to combine

the results from the four sequencing runs: each of the two hemispheres from one animal, and

one hemisphere sequenced twice from the other animal. Only unique barcodes from the re-

sequencing were included in analyses; duplicates were removed based on their barcode

sequences. This filtered, binarized, and collapsed matrix combined across the four hemispheres

was the basis for all analyses in this study.

Population summary

Projection proportions were calculated by summing the barcodes found in one brain

area and dividing by the total number of barcodes in the dataset; chance level was defined as

the barcode proportion found in control sites collected from cerebellum. The number of targets

for each neuron is calculated by summing the number of areas in which individual barcodes

were found and subtracting one, to count the number of non-amygdala targets for each

neuron; thus, a neuron which projects from one amygdala sample to another was defined as

having zero targets.

Conditional probabilities

To calculate the conditional probability of a neuron projecting to two areas, A and B, we

first found all of the cells that projected to area A (irrespective of their other targets). Then, we

found the subset of those cells which also project to area B. Thus, the conditional probability of

B|A is the number of cells that project to both A and B divided by the number of cells that

project to A.

K-means clustering

We performed k-means clustering using *scikit-learn* Python function. The optimal

number of clusters was determined using the elbow method, in which we plotted the number

of clusters against the within-cluster sum of squares. We found k= 12 to be optimal. We then

sorted the projection matrices by k-means cluster.

Over- and under-represented motifs

To assess whether any branching motifs were under- or over- represented compared to

chance, we first constructed a null distribution based on the overall proportion of barcode

found in each area. We assumed that the probability of a neuron branching to two areas, A and

B, would be the product of the independent probabilities of projecting to A and B: $p(A\&B) =$

$p(A) * p(B)$. This approach was taken for bi-, tri-, and quadrifurcations. We did not pursue

analysis of neurons with five or more targets, as there were fewer than 100 neurons with each

number of targets, leading to insufficient sample size on which to perform count-based

statistics.

Then, we compared our actual barcode counts to the expected null distribution using a

binomial probability test (*scipy.stats.binomtest*). Resulting p-values were FDR-corrected within

each branching degree, and effect sizes were calculated as $\log_2(\frac{observed}{expected})$.

Network analysis

To compare projections within the medial and ventral frontal cortex, we first isolated

projections to only one, not multiple, of the areas. We defined these networks by comparing

areas on the orbital surface (mOFC, lOFC, vlPFC, AI) to one another and areas on the medial

surface (dACC, pgACC, scACC). First, we computed the degree of overlap within these networks

by preparing Venn diagrams of projections specific to the areas and which branched between

them. The proportion of branching within each area was compared using z-tests for

proportions; p-values were adjusted using FDR correction.

We then focused on the projections which were specific while excluding any neurons

which projected to multiple areas in the same network. Projection strength to other areas were

calculated as described above and compared using pairwise z-tests for proportions and Fisher's

exact tests, corrected for multiple comparisons. Degrees of branching were calculated as

before, and branching was compared between brain areas using a permutation test in which

area labels were shuffled 1000x to generate a null distribution; distributions were then

compared using Chi-squared tests. Clustered projection heatmaps were constructed using

seaborn.clustermap with Ward's distance.

Comparison of neurons across populations

To understand whether projection motifs were reproducible across hemispheres and

across animals, we compared each of our independent sequencing runs to simulated neurons

from a uniform distribution (Extended Data Figure 4). 250 neurons were randomly sampled

from each of the sequencing runs and compared to other runs and to the simulated neurons

using cosine distance.

Stereology

Tissue for stereology was mounted from PBS onto gelatin coated slides and mounted

and coverslipped using Vectashield Vibrance Antifade aqueous mounting medium (Vector

Labs). Slides were stored in a lightproof slide box at 4°C to prevent fluorophore fading during

analysis.

An adjacent series was histochemically stained for acetylcholinesterase (AChE) for more

reliable identification of amygdala boundaries. Tissue was incubated overnight at 4°C in a

solution containing 0.68% sodium acetate (Thermo Scientific), 0.1% copper (II) sulfate (Thermo

Scientific), 0.12% glycine (Thermo Scientific), 0.12% acetylthiocholine iodide (TCI America), and

0.003% ethopropazine (Sigma Aldrich). The following morning, sections were rinsed 3x in PBS

for 5 minutes each, transferred to a solution of 0.1 M acetic acid (LabChem) with 1% sodium

sulfide (Thermo Scientific) for 1-2 minutes, and then rinsed again 3x with PBS before mounting

as above.

Stereology was performed using a Zeiss Apotome.2 microscope equipped with a Q-

Imaging digital camera, motorized stage, and Stereo Investigator software (MBF Bioscience). A

total of 12 sections were used for stereology, evenly distributed to cover the anterior-posterior

extent of the amygdala. The borders of the amygdala were identified using a 5x objective on

the AChE-stained sections, then ROI contours were realigned with the unstained sections for

stereological analysis. EGFP-, mCherry-, and double-labelled cells were counted based on the

soma as the counting target; the optical fractionator probe was used for stereological

estimation as described in West et al., 1991[38]. Neurons were counted under a 10x objective,

with a counting frame of size 150 x 150 x 15 μm; a 5-μm guard was applied to the dorsal aspect

of each section and a 20-μm guard to the ventral side. Counting frames were arranged in a

670.8 x 670.8 μm grid for systematic-random sampling. Results of the stereological analysis

were compared using z-tests for proportions.

fMRI analysis

Functional connectivity analyses were conducted on a previously published resting-state

functional MRI dataset[41]. We re-analyzed a subset of the scans, including data from 6 rhesus

macaques (5 males), focusing only on the control and pre-injection scans. After performing

standard preprocessing and warping all the brains to a standard template, masks for target

areas were drawn according to the areal boundaries used for MAPseq utilizing the D99,

CHARM, and SARM atlases as a basis[60–63]. Then, functional connectivity was computed between

each amygdala voxel and each target voxel in the ipsilateral hemisphere[64]. Correlations were z-

transformed, and all animals' data was combined into one large dataset after computing

within-animal z-scores. This way, we could analyze all of the amygdala voxels from each animal

without averaging across animals. The following analyses utilized the voxel-wise connectivity

matrix. Whole brain functional connectivity for the amygdala was calculated by averaging the

time-series within the anatomically defined amygdala and then correlating the average time

series with every voxel within the brain, for each animal, before averaging across animals.

The continuous measure of functional connectivity was binarized as follows. For each

amygdala voxel, the peak target connectivity was found; other target voxels which had

connectivity z-scores within 70% of that maximum were said to be connected, while those

under that threshold were said to not be connected. That 70% threshold allows about 5% of the

target voxels to survive filtering (**Supplemental Figure 6**). Then, as in MAPseq, we collapsed the

data across voxels within each target area, such that if one voxel in a target area survived

filtering, we concluded that amygdala was 'functionally connected' to that area. Then, we

followed a similar analysis pipeline as for the MAPseq data; computing projection strengths by

counting the proportion of voxels in each area to have survived filtering, computing conditional

probabilities, counting the number of 'targets' for each voxel, and performing k-means

clustering. We also performed the same network analysis by filtering for voxels connected to

areas in ventral or medial frontal cortex.

Data and Code Availability

Data and code can be made available upon reasonable request by reaching out to

peter.rudebeck@mssm.edu and zach.zeisler@icahn.mssm.edu.

Extended Data Figures

Extended Data Figure 1: Anatomical verification. A) Representative MRI images showing anterior (top) and middle (bottom) injection targets within amygdala. Vertical lines indicate intended injection tracks, while horizontal lines indicate injection depths along those tracks. **B**) Locations of injections for individual animals (colors); anterior injection on the left, middle on the right. Example tissue sections shown below with the extent of the amygdala surrounded by the dotted line (ento refers to entorhinal cortex, rs rhinal sulcus, sts superior temporal sulcus).

Extended Data Figure 2. Filtering parameters do not dramatically change recovered barcodes.
A) Number of barcodes surviving filtering for different source thresholds (color of line) and
target thresholds (x-axis). Sufficient source threshold eliminates majority of noise. **B**) Effect of
thresholding on number of projection targets per neuron. Each plot is one source threshold,
while colored lines reflect different target thresholds. The shape of the distributions is lightly
flattened by increasing projection threshold, while again, source threshold is responsible for
most of the noise.

Extended Data Figure 3: MAPseq is consistent across animals. A) Overall barcode distribution across areas. Colored bars represent the mean across all 4 hemispheres sequenced, error bars are standard deviation, and individual points reflect counts within each hemisphere's data separately. **B**) Number of targets for each neuron across hemispheres – roughly equal proportions of 1-target and 2+-target neurons, with most variance observed in proportion of 0-target neurons. **C**) K-means clustered branching projections, labelled by hemisphere. Note that most clusters are comprised of neurons from multiple hemispheres.

Extended Data Figure 4: Comparison across hemispheres. Density plot of cosine distance between two actual samples (red/blue) and simulated neurons from a uniform distribution (green). **A**) Monkey 1 neurons (both hemispheres) as a basis are more similar to monkey 2's neurons (Kolmogorov-Smirnov test, D = 0.12, p = 0.81) than the simulated neurons (D = 0.52, p < 0.0001. **B**) Same for monkey 2 as a basis. **C**) Within monkey 1, the two hemispheres are more similar to each other (D = 0.18, p = 0.27) than the random neurons (D = 0.43, p < 0.0001). **D**) Within monkey 2, the two sequencing runs were more similar to each other (D = 0.22, p = 0.12) than the simulated neurons (D = 0.48, p < 0.0001).

Extended Data Figure 5: Stereological confirmation of branching motifs. A) Top: injection strategy plotted on atlas sections (distances from intra-aural plane in mm); red and green lines refer to mCherry and EGFP retro-AAVs. Bottom: photomicrographs of actual injections sites. **B)** Example retrograde labelling in amygdala (shown in atlas on right). The bottom image is comprised of approximately 4x4 tiled 10x magnified images. Double labelled cells are non-exhaustively labelled with arrows. **C)** Stereology results (left) from left hemisphere injections compared to MAPseq results (right). **D)** Same for right hemisphere injections.

Extended Data Figure 6: fMRI analysis. A) ROIs used for fMRI analysis shown on NMT atlas slices (hippocampus and MD not shown). **B)** Proportion of voxels determined to be 'functionally connected' decreases with increasing threshold (the threshold was set at 70% because ~5% of voxels survive filtering). B) Binarized and collapsed MRI connectivity. Number labels refer to individual animals, while L and R refer to left and right hemispheres, respectively; only ipsilateral connections were assessed.