Main Manuscript for:

Cyclic and pseudo-cyclic electron pathways play antagonistic roles during nitrogen deficiency in *Chlamydomonas reinhardtii*

Ousmane Dao[1] [*], Adrien Burlacot[2,3], Felix Buchert[4], Marie Bertrand[1], Pascaline Auroy[1], Carolyne Stoffel[2], Jacob Irby[2], Michael Hippler[4,5], Gilles Peltier[1], Yonghua Li-Beisson[1] [*]

[1]Aix Marseille Univ, CEA, CNRS, Institute of Biosciences and Biotechnology of Aix Marseille, BIAM, CEA Cadarache, Saint Paul-Lez-Durance, France
[2]Department of Plant Biology, The Carnegie Institution for Science, Stanford, CA, 94305 USA
[3]Department of Biology, Stanford University, Stanford, CA, 94305, USA
[4]Institute of Plant Biology and Biotechnology, University of Münster, 48143 Münster, Germany
[5]Institute of Plant Science and Resources, Okayama University, Kurashiki 710-0046, Japan

*Correspondence to: Yonghua Li-Beisson (yonghua.li@cea.fr) or Ousmane Dao (ousmanedao17@yahoo.fr)

ORCID IDs: 0000-0002-7040-5770 (O.D.), 0000-0001-7434-6416 (A.B.), 0000-0001-6704-6634 (F.B.), 0000-0001-5098-1554 (M.B), 0000⎵0002⎵3376⎵6550 (P.A.), 0000-0002-4078-6250 (C.S.), 0000-0001-9670-6101 (M.H.), 0000-0002-2226-3931 (G.P.), 0000-0003-1064-1816 (Y.L.-B.)

Competing Interest Statement: There are no conflicts of interest.

Classification: Biological Sciences: Plant Biology

Keywords: Nitrogen deficiency; Flavodiiron proteins; Cyclic electron flow; Triacylglycerol; Photosynthetic regulation

This PDF file includes:

Main Text
Figures 1 to 5

Abstract (250 words)

Nitrogen (N) scarcity is a frequently encountered situation that constrains global biomass productivity. In response to N deficiency, cell division stops and photosynthetic electron transfer is downregulated, while carbon storage is enhanced. However, the molecular mechanism downregulating photosynthesis during N deficiency and its relationship with carbon storage are not fully understood. The Proton Gradient Regulator-like 1 (PGRL1) controlling cyclic electron flow (CEF) and Flavodiiron proteins involved in pseudo-(CEF) are major players in the acclimation of photosynthesis. To determine the role of PGRL1 or FLV in photosynthesis under N deficiency, we measured photosynthetic electron transfer, oxygen gas exchange and carbon storage in *Chlamydomonas pgrl1* and *flvB* knockout mutants. Under N deficiency, *pgrl1* maintains higher net photosynthesis and O_2 photoreduction rates, while *flvB* shows a similar response compared to control strains. Cytochrome b_6f and PSI are maintained at a higher abundance in *pgrl1*. The photosynthetic activity of *flvB* and *pgrl1 flvB* double mutants decreases in response to N deficiency similar to the control strains. Furthermore, the preservation of photosynthetic activity in *pgrl1* is accompanied by an increased accumulation of triacylglycerol depending on the genetic background. Taken together, our results suggest that in the absence of PGRL1-controlled CEF, FLV-mediated PCEF maintains net photosynthesis at a high level and that CEF and PCEF play antagonistic roles during N deficiency. It further illustrates how nutrient status and genetic makeup of a strain can affect the regulation of photosynthetic energy conversion in relation to carbon storage and provides new strategies for improving lipid productivity in algae.

Significance statement

Nitrogen (N) deficiency, an often-encountered phenomenon in nature, results in growth arrest, downregulation of photosynthesis and massive carbon storage in microalgae. However, more mechanistic insights involved in tuning photosynthetic electron transfer during N deficiency are required. Here, we provide evidence that a well-conserved protein in chlorophytes, the Proton Gradient Regulator-like 1 (PGRL1), is a key regulator of photosynthesis during N deficiency. In its absence, cells exhibited sustained photosynthesis thanks to the Flavodiiron (FLV) proteins. We propose that both PGRL1 and FLV, by having antagonistic roles in N deficiency, manage the redox landscape, carbon storage and biomass production. Our work revolves around the current paradigm of photosynthesis regulation during N deficiency and provides a new framework for improving biomass production and carbon storage in microalgae for biotechnological purposes.

Introduction

Nitrogen (N) deficiency is one of the most harsh environmental situation that constrains global primary biomass productivity in all ecosystems (1–3). Under N deficiency, cell division stops and photosynthetic CO_2 assimilation is downregulated, the carbon and energy being used to synthesize starch and triacylgcerols (TAGs) (4–7). The downregulation of photosynthetic electron transfer (PET) reactions, together with a re-routing of the excess reducing power towards carbon storage, prevents the over-production of reactive oxygen species, thus ensuring cell fitness (8–12).

Due to the variability in their natural habitat, microalgae must constantly adjust the photosynthetic conversion of energy to match the metabolic demand and have therefore developed a set of regulatory mechanisms to fine-tune electron transfer reactions. During photosynthesis, energy in the form of ATP and NADPH is mostly produced by the linear electron flow (LEF) (13). The balance of ATP and NADPH is essential for optimal CO_2 capture and metabolism to which cyclic electron flow (CEF) and pseudo-cyclic electron flow (PCEF) play a critical role (14–18). Two pathways of CEF around PSI have been described in the green microalga *Chlamydomonas reinhardtii* (*Chlamydomonas* hereafter), one involving the type II NADPH dehydrogenase (NDA2) (19), and the other being controlled by the proton gradient regulator 5 (PGR5)/PGR-like 1 (PGRL1) proteins (20).

Recent work in *Chlamydomonas* provided evidence that PGR5 and PGRL1 are involved in CEF (20–22). Hereby PGR5 is required for efficient stromal electron intake into the cytochrome b_6f complex (Cyt b_6f) (23). Its deletion thereby strongly disturbs the Mitchellian Q cycle (23). Also, deletion of PGRL1 impacts CEF in *Chlamydomonas* (20, 21). Despite the fact that PGRL1 has been implicated in plastoquinone reduction during CEF (24), it appears that PGRL1 is rather important for PGR5 expression and protein stability, as in the absence of PGRL1, PGR5 is strongly diminished, mimicking PGR5 dependent phenotypes (21, 25, 26).

Both CEF pathways reduce the plastoquinone (PQ) pool by using either NADPH or ferredoxin (Fd) as electron donor respectively (19, 24, 27). CEF has been shown to contribute to the acidification of thylakoid lumen by generating an extra proton motive force (*pmf*) in addition to the one produced by LEF (19, 20, 28, 29). PCEF mediated by Flavodiiron proteins (FLVs), by transferring electrons toward O_2 at the acceptor side of PSI, also contributes to the establishment of the *pmf* and therefore to the lumen acidification (30–32). The *pmf* is used to either (1) produce ATP (33), (2) trigger light energy quenching via a low lumenal pH (34–36), (3) convert HCO_3^- into CO_2 thanks to carbonic anhydrase in the lumen (32) or (4) repress electron transfer at the level the Cyt b_6f complex through the photosynthetic control triggered by the low lumenal pH (28, 37–40). Both CEF and PCEF have been shown to be critical under various conditions of light, CO_2 availability or sulfur deficiency by playing a synergistic role (14, 31, 32, 41, 42).

Despite the importance of CEF and PCEF in response to dynamic environments, little is known about their role during N deficiency. The role of NDA2-involved CEF have been recently addressed during CO_2-limiting photoautotrophic N deficiency and the knockdown of NDA2 was shown to impair the establishment of non-photochemical quenching (29). The role of PGRL1/PGR5-controlled CEF has been investigated under mixotrophic N deficiency and the lack of PGRL1 was shown to decrease the rate of CEF and TAG production (43). However, the role of PGRL1 during photoautotrophic N deficiency and the possible bioenergetic interactions between carbon/energy sinks (TAG and starch) and CEF or PCEF pathways that

generate ATP has not yet been explored so far. Considering that massive carbon reallocation occurs under N deficiency (4), the bioenergetics governing this reallocation are likely critical. Fully understanding photosynthesis regulatory pathways and how they affect biomass production and reserve formation is needed towards engineering photosynthesis and carbon storage in conditions of nutrient deficiency.

Here, we evaluated the contribution of PGRL1/PGR5-controlled CEF in the regulation of photosynthesis during N deficiency and evaluate its metabolic consequences on carbon storage in *Chlamydomonas* cells grown in photoautotrophic conditions under non-limiting CO_2 concentrations (using 1% CO_2-enriched air), conditions which favor the accumulation of carbon reserves (7, 44, 45). By monitoring photosynthetic activity based on chlorophyll fluorescence and O_2 exchange rate measurements, we observed high net photosynthetic activity in PGRL1-deficient strains under N deficiency compared to the control strains. Furthermore, the lack of PGRL1 in 137AH genetic background resulted in an over-accumulation of TAGs. Those effects being suppressed in double mutants deficient in both PGRL1 and FLVB. We conclude that FLVs, by maintaining a strong PCEF activity in the *pgrl1* mutants, catalyze high photosynthetic rates. Finally, we discuss how modulating photosynthetic electron flow could constitute an efficient strategy to improve photosynthesis and eventually boost further TAG production under N deficiency.

Results
Photosynthetic activity sustained longer in *pgrl1* under N deficiency
To assess the role of the PGRL1/PGR5-controlled CEF in the regulation of photosynthesis during photoautotrophic N deficiency, we simultaneously monitored chlorophyll fluorescence and O_2 exchange following a dark-light-dark transition in N-replete and N-deprived cells. We compared the photosynthesis efficiency of a PGRL1-deficient strain (*pgrl1$_{137AH}$*) with its control wild-type strain (137AH) and a complemented line (20) (**Fig. 1 and Supplemental Fig. S1 and S2**). Under N replete conditions, both chlorophyll fluorescence and O_2 exchange patterns were mostly similar in *pgrl1$_{137AH}$* and control cells (**Fig. 1*A, C, E and F* and Supplemental Fig. S1 *B-C***). After 2 days of N deficiency, the PSII operating yield measured in the light decreased by about 75% in the wild-type and only by 35% in *pgrl1* (**Fig. 1A, B, and E; Supplemental Fig. S1*C* and *E***). Conversely, *pgrl1$_{137AH}$* showed twice higher net O_2 evolution and light-dependent O_2 uptake as compared to the wild-type (**Fig. 1*D, F***). As previously reported (5, 46)**,** the dark O_2 consumption rate was stimulated in the 137AH strain during N deficiency, this effect being reduced in *pgrl1$_{137AH}$* (**Fig. 1*D*; Supplemental Fig. S2*B***). By using *pgrl1$_{137AH}$* complemented line *pgrl1::PGRL1*-2 (20), we noticed a full recovery of all the observed phenotypes (**Fig. 1 *E-G;* Supplemental Fig. S1 and S2**). To further strengthen the phenotype-genotype relationship, we generated an additional *pgrl1$_{CC125}$* mutant in the CC125 wild-type genetic background using CRISPR-Cas9. In line with our observation in the 137AH background **(Fig. 1*E-F*)**, we detected a sustained retention of photosynthetic PSII yield and higher net O_2 evolution in the *pgrl1$_{CC125}$* during N deficiency compared to CC125 (**Supplemental Fig. S3**). The higher O_2 consumption under N deficiency in the dark-adapted 137AH strain from **Fig. 1*D*** was independent of PGRL1 in the CC125 background (**Supplemental Fig. S3 *D-E***). The preservation of photosynthetic activity in *pgrl1* suggests that PGRL1-controlled CEF contributes to repressing the PET reactions under photoautotrophic N deficiency.

The PGRL1 modulates the PSI donor and acceptor side during N deficiency

The PGRL1/PGR5-controlled CEF is a determinant of the downstream PSI electron fate which prompted us to monitor P700, the primary PSI donor (**Fig. 2; Supplemental Fig. S4**). P700 forms three populations and only the first two are photo-oxidizable: 1. PSI donor shortage generates $P700^+$ in the environmental light, 2. PSI yield quantifies $P700^-$ during environmental acclimation that converts within the ms range to $P700^+$ upon a saturating light pulse, and 3. PSI acceptor side limitation represents a redox-inactive P700 pool (see Materials and Methods). In the absence of PGRL1/PGR5-controlled CEF the redox-inactive P700 pool is enhanced under CO_2-limiting conditions (47). This was also observed in the 137AH genetic background when *pgrl1* was grown photoautotrophically under 1% CO_2 (**Fig. 2A, B**). The failure to produce $P700^+$ in the light gradually developed within 0.5 s and persisted throughout the illumination period (**Fig. 2A, B**). However, *pgrl1* did not display lower PSI yields, typically detected in ambient CO_2 (48), since the diminished $P700^+$ was due to lower donor side limitation (**Fig. 2B**). The P700 differences in *pgrl1* relied on PSII activity, pointing to a LEF/CEF entanglement. N depletion further increased the $P700^+$ pool in a strictly donor side-dependent fashion in both wild-type and mutant. Nevertheless, *pgrl1* displayed twice the PSI acceptor side limitation when PSII was active. Moreover, the mutant showed a delayed $P700^+$ formation within the first 200 ms of light when PSII was inhibited (**Fig. 2A**). The latter feature was also displayed in N-deficient CC125 under DCMU conditions (**Supplemental Fig. S4A**). This genetic background showed identical photo-oxidizable P700 pools between wild-type and mutant in the presence of N, with *pgrl1* producing slightly higher acceptor side limitation (**Supplemental Fig. S4B**). N depletion came at the expense of PSI yields and *pgrl1* failed to alleviate the PSI acceptor side through sufficient donor side induction. These results suggest that PGRL1-controlled CEF facilitates the onset of photosynthetic control during photoautotrophic N deficiency by inducing a donor side limitation at PSI to alleviate its acceptor side redox pressure.

FLV-mediated O_2 photoreduction drives photosynthesis in *pgrl1* under N deficiency

Because light-dependent O_2 uptake can lead to the generation of extra ATP and support CO_2 fixation but at various levels depending on the mechanism involved (18), we sought to test the nature of the light-dependent O_2 uptake mechanism activated in *pgrl1* mutants during N deficiency (**Fig. 1D**). FLVs have been reported to be a major O_2 uptake mechanism in the light (31, 32, 49). *pgrl1 flvB* double mutants impaired in both PGRL1 and FLVB have been recently generated by crossing the *pgrl1_{137AH}* with a FLVB deficient strain (*flvB21*) as described in (32). Therefore, we used the parental lines as control strains (*flvB-21* and *pgrl1_{137AH}*) as well as sibling strains from the progeny of the crossing (WT1 and WT3) harboring both FLV and PGRL1 proteins (**Supplemental Fig. S1A**). The two mutants (*pgrl1 flvB-3* and *pgrl1 flvB-5)* and control sibling (WT1 and WT3) strains used throughout the manuscript have been chosen for having similar maximal photosynthesis under N-replete conditions (V_{Max}) (18, 32). For all the experiments, we kept the *pgrl1_{137AH}* alongside with its control 137AH and *pgrl1::PGRL1-2*; *flvB* mutants (*flvB21*, *flvB208* and *flvB308*) alongside with their control CC4533 and finally *pgrl1 flvB-3* and *pgrl1 flvB-5* alongside WT1 and WT3 as control. We evaluated steady-state O_2 exchange rate in *pgrl1_{137AH}*, *flvB* mutants as well as in the *pgrl1 flvB* double mutants after 2 days of N deficiency (**Fig. 3**). The light-dependent O_2 uptake was highly increased in *pgrl1_{137AH}* but strongly

impaired in *flvB* and *pgrl1 flvB* mutants (**Fig. 3A; Supplemental Fig. S2**) which
mirrors what has been reported for N-replete conditions in the presence of
atmospheric CO_2 level (31, 32). Interestingly, while the gross O_2 uptake, the gross O_2
and the net O_2 evolution measured during steady-state photosynthesis remained high
in *pgrl1$_{137AH}$*, no difference was detected between *flvB*, *pgrl1 flvB* mutants and their
respective control lines (**Figure 3B-D and Supplemental Fig. S2**). Taken together,
these data show that the absence of FLV-mediated O_2 photoreduction (in *pgrl1 flvB*
mutants) suppresses the effect of N deficiency observed in *pgrl1$_{137AH}$* strain. We
conclude that FLV-mediated O_2 photoreduction, by maintaining PET reactions in
pgrl1, allows net photosynthesis to be maintained at a high-level during N deficiency.
239
Cyt b_6f and PSI subunits accumulate to higher levels in *pgrl1* under N
deficiency
To gain insights into the mechanisms behind the high photosynthetic activity in *pgrl1*
during N deficiency, we compared the relative abundance of representative catalytic
core subunits from photosynthetic complexes in the *pgrl1$_{137AH}$* and the 137AH wild-
type during N deficiency (**Fig. 4A and D**). Additionally, we also investigated potential
changes in mitochondrial electron transport chain by probing Complex II (Cox IIB).
We observed a sustained retention of the PSI subunit PsaD and Cyt *f* subunit of Cyt
b_6f in *pgrl1$_{137AH}$* compared to the 137AH (**Fig. 4A and D**). Higher amounts of PsaD
and Cyt *f* were observed in *pgrl1$_{137AH}$* under N replete without any functional effect on
PSII yield and Net O_2 evolution (**Fig. 1E-F; 4A and D**). As for the stimulated dark O_2
consumption **(Fig. 1D)**, the amount of mitochondrial respiratory chains component
Cox IIB was increased in the 137AH whereas it remains stable in the *pgrl1$_{137AH}$*
during N deficiency (**Fig. 4A**). It is worth noting that the hallmark of autophagy, i.e.,
ATG8 was barely detectable in the *pgrl1$_{137AH}$* whereas it was highly induced in the
137AH under N deficiency (**Fig. 4A, D**). All the other proteins tested accumulated to a
similar amount in *pgrl1$_{137AH}$* and 137AH (**Fig. 4A**). Note that higher amounts of Cyt *f*
were also observed in CRISPR-generated *pgrl1$_{CC125}$* mutant as compared to CC125
in response to N deficiency (**Supplemental Fig. S3E**). We conclude from these
experiments during N deficiency that the higher photosynthetic activity observed in
mutants impaired in PGRL1 might result from a lower decrease in the amounts of PSI
and Cyt b_6f complexes.
To test whether the retention of the PSI and Cyt b_6f observed in the *pgrl1$_{137AH}$*
are consequences of the existence of a FLV activity, similar immunoblot analyses
were performed in the *flvB-21* and *pgrl1 flvB-3* mutants. Consistent to the
photosynthetic activity, immunoblot analyses showed a similar accumulation of tested
proteins in the *flvB-21* and *prgl1 flvB-3* compared to controls with the exception of
lower PsaD and ATG8 in *prgl1 flvB-3* (**Fig. 4 B-C and E-F**).
268
Defects in CEF and PCEF altered carbon storage during N deficiency
Emerging literature suggests that alterations in energy management pathways affect
not only biomass production but also its composition (16, 17, 44, 50). Starch and
TAGs are major forms of carbon storage in *Chlamydomonas* during N deficiency.
Since the *pgrl1$_{137AH}$* mutant shows a stronger net photosynthetic rate, we measured
its ability to accumulate storage compounds. Starch accumulation was similar in
pgrl1$_{137AH}$ and control lines under both N replete and deficiency (**Fig. 5A;**
Supplemental Fig. S5A and $S6A$). In contrast, twice more TAG accumulation was
observed in *pgrl1$_{137AH}$* compared to the control lines under N deficiency whereas no

difference was observed under N-replete condition (**Fig. 5*D*; Supplemental Fig. S5*B***
and S6*D*). Lipid droplet imaging confirmed the accumulation of higher TAG amounts
(**Supplemental Fig. S5*D***). Intriguingly, the *PGRL1* knockout mutant did not induce
TAGs or starch accumulation in response to N deficiency in the CC125 genetic
background (**Supplemental Fig. S6 *G-J***), which may be related to the fact that
photosynthesis preservation (PSII yield and net O_2 evolution) was lower in the
pgrl1$_{CC125}$ background as in the *pgrl1$_{137AH}$* background (**Supplemental Fig. S3**) thus
limiting reserve accumulation, or indicating the existence of a metabolic limitation for
storage compound accumulation in the CC125 background.
Under N replete, the *flvB* single mutants accumulated higher amounts of starch than
the CC4533 wild-type (**Supplemental Fig. S6*B***) but accumulated lower amounts of
TAG under N deficiency (**Fig. 5*E***). The double mutants *pgrl1$_{137AH}$ flvB* accumulated
similar amounts of TAG and starch as compared to control strains either in N replete
or N deprived conditions (**Fig. 5*C, F*; Supplemental Fig. S6*C, F***).
We then evaluated to which extent observed changes in photosynthesis affect
biomass accumulation, by following changes in cellular volume as a proxy for
biomass accumulation. After 6 days of N deficiency, the cell volume increased twice
as fast in the *pgrl1$_{137AH}$* compared to the control lines (**Supplemental Fig. S5*C***). We
conclude from the above experiments that the higher photosynthetic activity in
pgrl1$_{137AH}$ results in increased biomass and TAG productivity under N deficiency.
Altogether, our data suggested that during phototrophic N deficiency, photosynthetic
electron flow mediated by FLVs could trigger higher TAG accumulation while PGRL1
tends to constrain it, likely via a photosynthetic control-related process or specific
protease pathways that target Cyt b_6f. We propose that both PGRL1 (CEF) and FLVs
(PCEF), by having an antagonistic role during N deficiency, manage the redox
landscape, carbon storage and biomass production. It is worth pointing out that this
modification of metabolism as a consequence of redox management depends not
only on genetic background (137AH versus CC125) but also on the environmental
state (+N versus –N).
307
Discussion

The ability of microalgae to coordinate their energy conversion (from light to chemical
energy) to meet the metabolic demand is crucial for their survival in a constantly
fluctuating environment. Mechanisms involved in photosynthesis regulations have
been abundantly studied in response to light or CO_2 levels, but not much is known
during nutrient deficiency when a massive re-orientation of metabolic pathways
occurs. In the absence of N, the major cellular energy sinks (cell division, protein
biosynthesis and photosynthetic CO_2 fixation) are restricted whereas the carbon and
energy are stored as TAGs or starch. Moreover, the PET reactions are
downregulated under N deficiency. However, we do not know whether the
downregulation of PET is a mean to match cells energy status with the limited
metabolic demand. In other words, how the metabolism accommodates when
photosynthesis remains high during N deficiency is not well understood. In this study,
we demonstrated that PGRL1, which controls a CEF pathway, tunes PET during N
deficiency. This tuning favors the photosynthetic control mechanism via Cyt b_6f to
limit PSI donor availability (see below for details) and might affect energy-transducing
enzyme ratios at least in chloroplasts. Lack of PGRL1 resulted in sustained
photosynthetic activity up to 2 days of N deficiency. The misregulation of the
PGRL1/PGR5-controlled CEF is notorious for decreasing the photosynthetic control
efficiency and re-routing of electrons into alternative acceptors such as H_2 production

by Hydrogenases or O_2 photoreduction by FLVs (14, 20, 50). Here, we have further
shown that the higher FLVs-mediated PCEF in *pgrl1* mutants channels excess
electrons toward O_2 under N deficiency. PCEF thus counteracts PSI over-reduction,
likely facilitating ATP production and at the same time keeping PET active.

PGRL1/PGR5-controlled CEF contributes to photosynthetic control during N
deficiency
The photosynthetic control refers to mechanisms that restrict the PET reactions
mostly occurring in response to environmental fluctuations and is typically achieved
on the level of the Cyt b_6f upon acidification of the thylakoid lumen (37, 38, 52, 53).
The PGRL1-PGR5-controlled CEF increases lumen acidification efficiency and has
been shown to contribute to the photosynthetic control in higher plants (26, 28, 54) as
well as in microalgae (20–23, 51). The induction of the photosynthetic control results
in PSI donor side limitation in the light which is typically determined *in vivo* via P700$^+$
optical readouts (55–57). Our results showed higher acceptor side limitation (at the
expense of donor side limitation) in *pgrl1* mutants (**Fig. 2 and Supplemental Fig. S4**)
indicating that they are affected in the induction of the photosynthetic control. The
decrease in Cyt b_6f abundance could be an additional cellular strategy to further
restrict the PET toward PSI (higher donor side limitation in wild-type, **Fig. 2 and**
Supplemental Fig. S4). While the wild-type successfully relaxes the redox pressure
on PSI and degrades Cyt b_6f under N deficiency, *pgrl1* mutants sustained PET and
failed to degrade the Cyt b_6f. This dysfunctioning of the photosynthetic control in
pgrl1 (high acceptor side limitation and Cyt b_6f abundance; **Fig. 2 and 4*A***), promotes
the recruitment of stromal electron carriers such as FLVs and TAG biosynthesis (in
137AH background). Similar observations were reported under sulphur deprivation
where deletion of PGRL1 or PGR5 resulted in sustained linear electron flow toward
H_2 production in *Chlamydomonas* (20, 51). Our work shows that impairing the
accumulation of PGRL1 removes a bottleneck of photosynthetic electron flow during
N deficiency under non-limiting CO_2 conditions (**Fig. 1-3; Supplemental Fig. S1-3**).
This characteristic of photosynthetic control makes PGRL1/PGR5-controlled CEF a
promising target for improving photosynthetic yield under nutrients deficiency.
Indeed, the control of photosynthesis under N deficiency has been seen as safety
mechanisms protecting cells from phototoxicity (28, 53, 54). Here, we show that
thanks to the presence of FLVs, photosynthesis could be improved by removing
PGRL1-controlled CEF during N deficiency under non-limiting CO_2 conditions.

FLVs become predominant in the absence of PGRL1/PGR5-controlled CEF
So far, a compensation mechanism between PGRL1 and FLVs has been reported
during algal adaptation to light or low-CO_2 conditions (14, 18, 32). Under high light or
low-CO_2 conditions, the increased activity of FLVs in the PGRL1-deficient strain does
not improve the net photosynthesis despite its strong efficiency in generating ATP
(18, 58). Similar compensation has also been observed in higher plants where
orthologous expression of FLVs rescues the *pgr5* mutants phenotype without further
improving photosynthesis (54, 59) although some levels of increased biomass were
measured in wild-type Arabidopsis expressing FLVs under light fluctuations thanks to
the protective role of FLVs in these conditions (60). In contrast, our results suggest
that under N deficiency, FLVs and PGRL1 have an antagonistic role. Indeed, the
increased activity of FLVs in the PGRL1-deficient strains resulted in higher net
photosynthesis (**Fig. 3*D*; Supplemental Fig. S2**), which shows that rather than

compensating each other, PGRL1 is indirectly controlling the activity of FLVs by
limiting the maximal electron flow capacity. Our observation of the sustained
accumulation of Cyt b_6f in the *pgrl1* but not in *pgrl1 flvB* double mutants under N
deficiency **(Fig. 4 A and *D*)** could be attributed to the activity of FLVs-mediated PCEF
draining electrons from the photosynthetic chain and indirectly preventing Cyt b_6f
from degradation. The latter process is catalysed in *pgrl1 flvB* upon PCEF shortage.
The presence of strong FLVs activity in *pgrl1* thereby allows maintaining high PET
while generating ATP for CO_2 fixation and reserve formation.
Because CEF and PCEF are dominant pathways for supplying extra ATP in
addition to LEF in the chloroplast, we were expecting that the removal of both PGRL1
and FLVs would have more severe consequences on cell physiology and metabolism
as it was observed when CO_2 is limiting (32). Instead, we observed similar
photosynthetic activity and TAG biosynthesis in the *pgrl1 flvB* double mutants as their
control strains (**Fig. 3 and 5**), i.e. the additional removal of FLVs suppressed the
phenotype of PGRL1 deficiency. A third pathway generating the energy required for
photosynthetic CO_2 fixation and ensuring redox dissipation could be operating in the
double mutant *pgrl1 flvB*. Both the NDA2-dependent CEF (19, 29) or a chloroplast-
mitochondria electron flow (CMEF) (18) could be good candidates. NDA2 protein
level was shown to increase during air photoautotrophic N deficiency under
atmospheric CO_2 level and the CEF rate in *Chlamydomonas* was decreased by 50%
in *nda2* mutants (29). However, we observed a similar protein level of NDA2 in the
pgrl1 flvB as their control lines (**Fig. 4*C***). Nevertheless, the protein level might not
always correlate with the activity and other regulatory mechanisms (e.g.,
phosphorylation or redox regulation) can modulate enzyme activity. Additionally, the
stimulated mitochondrial respiration rates as well as increased in Cox IIB protein level
during N-deficiency in all the strains (except *pgrl1$_{137AH}$*) (**Fig. 4; Supplemental Fig.
S3**) points to a strong activity of CMEF (18, 32), which might compensate for CEF
and PCEF deficiency in *pgrl1 flvB* N-deprived cells. CMEF may have two distinct
roles, either to supply additional ATP or favour redox dissipation. In the context of
generating extra ATP, the Cox IIB pathways could take over the AOX1 alternative
oxidase pathway because of its efficiency in generating ATP (18). Further
investigation would be required to identify the mechanisms generating ATP in the
absence of PGRL1 and FLVs. Altogether, we conclude that FLVs maintain high
photosynthetic activity under N deficiency in the absence of PGRL1 by channeling
excess electrons toward O_2 meanwhile generating ATP for CO_2 and downstream
metabolic pathways (TAG production).

Relationship between cellular redox landscape and carbon storage
A major biotechnological challenge in algal domestication for biofuel is the tradeoff
between growth and lipid productivity. In *Chlamydomonas* as in many other
microalgae, starch and TAG massively accumulate but mostly under stress
conditions in particular N deficiency when cell division stops and productivity is
impaired. Considerable efforts have focused on the study of the molecular
mechanisms behind the onset of reserve accumulation by monitoring omics
responses to a stress (5, 61–64), or focused on specific steps of fatty acid and TAG
biosynthesis, which have resulted in some limited improvement in productivity (65,
66). Improving productivity requires a better understanding of the crosstalk between
photosynthetic carbon fixation, environmental signals and the redox balance, which
all govern reserve accumulation (67–69). Here by studying mutants affected in CEF

and PCEF, we explored the relationships between the cellular redox status and carbon storage.

The increased accumulation of TAG but not starch observed in the PGRL1-deficient strain (137AH background) under N deficiency (**Fig. 5 *A* and *B***) is a consequence of continuous production of NADPH and ATP through PCEF-dependent PET. In line with our finding, the *Chlamydomonas pgd1* mutants (e.g. *Plastid galactoglycerolipid degradation 1)* with reduced LEF rate (less ATP and NADPH) are shown to produce less TAG under N starvation (11, 70). The report that the *pgrl1* mutant made less TAG than wild-type under mixotrophic conditions (71) is not surprising. It is well known that the bioenergetics of *Chlamydomonas* under photoautotrophic conditions differ from mixotrophic conditions (72–74). The presence of acetate can drastically affect cellular bioenergetics levels i.e. its uptake consumes ATP and its metabolism produces NADH, therefore further favoring oil synthesis (72, 73, 75, 76). TAGs accumulation was different in the CC125 background where the *pgrl1$_{CC125}$* mutant made similar amounts of TAGs as its background strain (**Supplemental Fig. S6*J***). This seems not surprising when we consider that TAG accumulation is a metabolic consequence of changes in chloroplast redox state and in photosynthetic performance. The preservation of PSII yield and net O_2 evolution was less dramatic in the CC125 background than in the 137AH background, which could be one of the reasons behind strain-dependent phenotype. This further reflects the fact that oil content is a complex trait, and it is the consequence of the interaction between genetic makeup, its metabolic flexibility and the capacity of the extent of a given cell's response to environmental changes.

In contrast to *pgrl1$_{137AH}$*, the *flvB* mutants accumulated lower amounts of TAGs under N deficiency, and no difference in TAG was observed in the *pgrl1 flvB* double mutants, consisting again with the profile in photosynthetic performance and with their antagonistic roles. The *flvB* mutants accumulated higher amount of starch under N sufficient condition though. To conclude, this work further points to the importance of the role redox management on carbon allocation and storage, and that this effect is dependent on many factors including the genetic background, the trophic style, as well as the nutrient status.

Materials and methods
Growth conditions and strains
The *pgrl1$_{137AH}$* (from the background 137AH) with its complementing strain (*pgrl1::PGRL1*-2), *flvB* mutants from the background CC-4533 and the *pgrl1 flvB* double mutants were previously described (20, 31, 32). *pgrl1$_{CC125}$* mutants were generated in the CC125 background using CRISPR-Cas9 (see Supplemental Materials and Methods). All the information about the strains used are reported in **Supplemental Table 1**. Cells were routinely cultivated in an incubation shaker (INFORS Multitron pro) maintained at 25°C, with 120 rpm shaking and constant illumination at 50 µmol m^{-2} s^{-1}. Fluorescent tubes delivering white light enriched in red wavelength supplied lightings in the INFORS. All experiments were performed under photoautotrophic condition with minimum medium with or without nitrogen source (MM and MM-N) buffered with 20 mM MOPS at pH 7.2 in air enriched with 1% CO_2. Due to cell aggregation in the 137AH background (notably the *pgrl1* mutant) that prevent accurate cell counting, total cellular volume was measured using a Multisizer 4 Coulter counter (Beckman Coulter) and the different strains were diluted to reach a similar cellular concentration before N deficiency experiments.

Measurement of chlorophyll fluorescence using a Pulse Amplitude Modulation (PAM)

Chlorophyll fluorescence was measured using a PAM fluorimeter (Dual-PAM 100, Walz GmbH, Effeltrich, Germany) on the MIMS chamber as described in (49) using green actinic light (1250 µmol photon m^{-2} s^{-1}, green LEDs). Red saturating flashes (8,000 µmol photons m^{-2} s^{-1}, 600 ms) were delivered to measure the maximum fluorescence (FM) every 30 s (before and upon actinic light exposure). The maximum PSII quantum yield was calculated as *Fv/Fm= (Fm-Fo)/Fm* where F_0 is the basal fluorescence obtained with the measuring light and Fm the fluorescence emitted after saturating pulse (77). PSII operating yield (ΦPSII) was calculated as ΦPSII= $(F_M'-Fs)/F_M'$ with F_M' the fluorescence value after saturating pulse, Fs the stationary fluorescence during actinic light exposure.

O_2 exchange measurement using Membrane Inlet Mass Spectrometry (MIMS)

O_2 exchanges were measured in the presence of $[^{18}O]$-enriched O_2 using a water-jacketed, thermoregulated (25°C) reaction vessel coupled to a mass spectrometer (model Prima ΔB; Thermo Electronics) through a membrane inlet system (78). The cell suspension (1.5 mL) was placed in the reaction vessel and bicarbonate (10 mM final concentration) was added to reach a saturating CO_2 concentration. One hundred microliters of $[^{18}O]$-enriched O_2 (99% $^{18}O_2$ isotope content; Euriso-Top) was bubbled at the top of the suspension just before vessel closure and gas exchange measurements. O_2 exchanges were measured during a 3-min period in the dark, then the suspension was illuminated at 1250 µmol photons m^{-2} s^{-1} for 5 min using green LEDs followed by 3-min in the dark. Isotopic O_2 species $[^{18}O^{18}O]$ (m/e = 36), $[^{18}O^{16}O]$ (m/e = 34), and $[^{16}O^{16}O]$ (m/e = 32) were monitored, and O_2 exchange rates were determined (78). Argon gas was used to correct O_2 exchange measured by the spectrometer as described in (78).

Redox state measurements of P700

The PSI primary electron donor, were obtained from optical signals (705 nm – 740 nm) using a Joliot-type spectrophotometer (JTS-150, Spectrologix USA), described in detail elsewhere (23, 79, 80). Harvested cells (2 min, 4000 rpm, 22°C) were resuspended in cuvettes to comparable densities in growth medium containing 20% Ficol (*w/v*). Samples were subjected to 2.5-s alterations of dark/light (490 µmol photon m^{-2} s^{-1}, 630-nm LEDs), leading to a partial P700 pre-oxidation during the light period (donor side limitation). Additional fractions of photo-oxidizable P700 (PSI yield) were obtained by a 25 ms saturating pulse before the dark period, revealing non-oxidizable P700 (acceptor side limitation) after comparison to fully oxidized P700 when measured in the presence of 20 µM 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU).

All other methods are described in ***SI Materials and Methods.***

Acknowledgments

O.D. thanks The French Atomic Energy and Alternative Energy Commission (CEA) for a PhD scholarship. G.P. and Y.L.-B. acknowledge the continuous financial support of CEA (LD-power, CO2Storage). A.B. acknowledges the support from the Carnegie Institution for Science. We thank Bertrand Legeret for maintaining the HelioBiotec lipidomics platform and thank Mallaury Cabanel for assistance in performing immunoblot. We also acknowledge the ZoOM microscopy facility. M.H.

526 (DFG Research Unit FOR 5573) and F.B. (BU 3426/3-1) acknowledge Deutsche
527 Forschungsgemeinschaft for funding.
528

Author's Contributions:
Y.L-B., G.P., A.B., and O.D. conceived the study. Y.L-B., G.P. and A.B. provided
supervision. O.D. performed most of the experiments. P.A., M.B. and O.D. carried out
biochemical experiments. M.B. and O.D. performed starch and lipid analysis. G.P.
supervised the MIMS experiments. C.S. and J.I. generated the CRISPR-mediated
mutant lines under the supervision of A.B.. F.B. and M.H. performed and analyzed
the P700 measurements. O.D. drafted the manuscript with contributions from A.B.,
G.P, F.B., M.H. and Y.L-B.
537

Figures
Figure 1. *pgrl1* showed sustained photosynthesis after 2 d of N deficiency.
A-B, Chlorophyll fluorescence was measured using a Dual-PAM during the dark-
light-dark transition from N-replete (**A**) and N-deficient (**B**) conditions.
C-D, O_2 exchange rates were measured using a MIMS during the dark-light-dark
transition from N-replete (**C**) and N-deficient (**D**) conditions. Net O_2 evolution (green)
was calculated as gross O_2 evolution (blue) – gross O_2 uptake (red).
E, PSII quantum yield before and after 2 d of N deficiency measured using green
actinic light (1250 µmol photon m^{-2} s^{-1}, green LEDs) and calculated as ΦPSII= (F_M'-
Fs)/ F_M' with F_M' the fluorescence value after saturating pulse, Fs the stationary
fluorescence during actinic light exposure.
F, The Net O_2 evolution before and after 2 d of N deficiency in the wild-type 137AH,
pgrl1$_{137AH}$ and the complemented line *pgrl1::PGRL1-2*.
G, Immunoblot analysis of PGRL1 accumulation in the wild-type 137AH, *pgrl1* and
the complemented line. L30 antibody was used as a positive control. Coomassie blue
staining was used as the loading control. Protein samples were obtained from 2 d N-
starved cells.
N-replete wild-type 137AH, *pgrl1*$_{137AH}$, and *pgrl1::PGRL1-2* cells cultivated
photoautotrophically with 1% CO_2 in air under continuous light (50 µmol photons m^{-2}
s^{-1}) were transferred into N-free media for 2 d prior to measurements. Shown are an
average of at least three biological replicates ± SD. Asterisks represent statistically
significant differences compared to the control 137AH (* $p<0.05$, ** $p<0.01$ and ***
$p<0.001$) using one-way ANOVA.
561

Figure 2. PSI oxidation is facilitated by PGRL1 under N deficiency
The influence of PGRL1 on the redox state of the primary PSI donor, P700, was
monitored in the genetic background 137AH **(A-B)** and CC125 **(Supplemental Fig.**
S4). Panel A shows raw kinetics (red/yellow/black bars: 490/3000/0 µmol photons
m^{-2} s^{-1}) in the presence and absence of PSII inhibitor DCMU (means ± SD of 4
biological replicates). Panel B quantifies the corresponding P700 pools at the end of
the 2.5-s light period (letters indicate parameter-specific significances using one-way
ANOVA/Fisher-LSD, $p<0.05$). At the time of quantification, P700 remained photo-
oxidizable by a saturating pulse (PSI yield), was redox inactive (no acceptors) or was
pre-oxidized (no donors). N-replete cells of wild-type 137AH and *pgrl1*$_{137AH}$ were
cultivated photoautotrophically with 1% CO_2 in air under continuous light (50 µmol
photons m^{-2} s^{-1}), and then transferred into N-free media for 24 h prior to
measurements.
575

Figure 3. The photosynthesis in *pgrl1* is driven by FLV-mediated O_2 photoreduction under N deficiency.

A, Light-dependent O_2 uptake attributed to FLV activity and calculated as the difference between the O_2 uptake during the 1st min of illumination and dark O_2 uptake as shown by the **Supplemental Figure 2**.

B, Gross O_2 uptake measured between 5-7min as shown by the **Supplemental Figure 2**.

C, Gross O_2 evolution measured between 5-7min as shown by the **Supplemental Figure 2**.

D, Net O_2 evolution measured between 5-7min as shown by the **Supplemental Figure 2**.

O_2 exchange rates were measured using a MIMS in the presence of [^{18}O]-enriched O_2. Net O_2 evolution (green) was calculated as gross O_2 evolution (blue) – gross O_2 uptake (red). Total proteins were extracted from N-replete and 2 days N-deprived cells and samples were loaded at equal total proteins amounts as shown on Coomassie blue staining. Cells cultivated photoautotrophically with 1% CO_2 in air under continuous light of 50 µmol photons $m^{-2} s^{-1}$ were transferred into N-free media for 2 days prior to measurements or sampling for immunoblot. Shown are an average of at least three biological replicates ± SD. Asterisks represent statistically significant difference compared to the wild-type strains (* $p<0.05$, ** $p<0.01$, *** $p<0.001$ and **** $p<0.0001$) using one-way ANOVA.

Figure 4. *pgrl1* accumulated more cytochrome Cyt b_6f and PsaD during N deficiency.

A-C, Immunodetection of photosynthetic proteins in the *pgrl1$_{137AH}$*, *flvB-21* and *pgrl1 flvB-3* as compared to their controls during N deficiency. Cells were harvested at 0, 8, 22, 30 and 50 hours of N deficiency. Samples were loaded at equal total protein amounts as shown by Coomassie blue staining. Shown are representative images of three biological replicates. The α-tubulin was used as housekeeping protein whereas ATG8 and the chloroplast 50S ribosomal large subunit L30 are controls of N depletion (81, 82).

D-F, Histogrammes showing the abundance Cyt *f*, PsaD and ATG8 proteins in the *pgrl1$_{137AH}$*, *flvB-21* and *pgrl1 flvB-3*during N deficiency in the wild-type cells. Shown represent the ratio of mutants over control strains. Data are average of at least three biological replicates.

Figure 5. *pgrl1* over accumulated TAGs under N deficiency

Starch **(A-C)** and TAG (**D-F**) quantification in N-deprived cells of *pgrl1$_{137AH}$*, *flvB* and *pgrl1 flvB* mutants respectively. Cells were cultivated photoautotrophically with 1% CO_2 in air under continuous light of 50 µmol photons $m^{-2} s^{-1}$. For N deficiency, cells were transferred into N-free media for 2 d prior to sampling for starch and TAG. Shown are an average of at least three biological replicates ± SD. Asterisks represent statistically significant difference comparing mutants with their control strains (* $p<0.05$, ** $p<0.01$ and *** $p<0.001$) using one-way ANOVA.

Reference

1. P. M. Vitousek, R. W. Howarth, Nitrogen limitation on land and in the sea: How can it occur? *Biogeochemistry* **13**, 87–115 (1991).

2. D. S. LeBauer, K. K. Treseder, Nitrogen limitation of net primary productivity in terrestrial ecosystems is globally distributed. *Ecology* **89**, 371–379 (2008).

3. E. Du, *et al.*, Global patterns of terrestrial nitrogen and phosphorus limitation. *Nat. Geosci.* **13**, 221–226 (2020).

4. M. Siaut, *et al.*, Oil accumulation in the model green alga Chlamydomonas reinhardtii: characterization, variability between common laboratory strains and relationship with starch reserves. *BMC Biotechnology* **11**, 7 (2011).

5. S. Schmollinger, *et al.*, Nitrogen-Sparing Mechanisms in Chlamydomonas Affect the Transcriptome, the Proteome, and Photosynthetic Metabolism. *The Plant Cell* **26**, 1410–1435 (2014).

6. M. T. Juergens, *et al.*, The Regulation of Photosynthetic Structure and Function during Nitrogen Deprivation in Chlamydomonas reinhardtii. *Plant Physiology* **167**, 558–573 (2015).

7. M. Schulz-Raffelt, *et al.*, Hyper-accumulation of starch and oil in a Chlamydomonas mutant affected in a plant-specific DYRK kinase. *Biotechnology for Biofuels* **9**, 55 (2016).

8. Y.-M. Zhang, H. Chen, C.-L. He, Q. Wang, Nitrogen Starvation Induced Oxidative Stress in an Oil-Producing Green Alga Chlorella sorokiniana C3. *PLOS ONE* **8**, e69225 (2013).

9. J.-J. Park, *et al.*, The response of Chlamydomonas reinhardtii to nitrogen deprivation: a systems biology analysis. *The Plant Journal* **81**, 611–624 (2015).

10. M. Gargouri, P. D. Bates, J.-J. Park, H. Kirchhoff, D. R. Gang, Functional photosystem I maintains proper energy balance during nitrogen depletion in Chlamydomonas reinhardtii, promoting triacylglycerol accumulation. *Biotechnology for Biofuels* **10**, 89 (2017).

11. Z.-Y. Du, *et al.*, Galactoglycerolipid Lipase PGD1 Is Involved in Thylakoid Membrane Remodeling in Response to Adverse Environmental Conditions in Chlamydomonas. *The Plant Cell* **30**, 447–465 (2018).

12. Q.-G. Tran, *et al.*, Impairment of starch biosynthesis results in elevated oxidative stress and autophagy activity in Chlamydomonas reinhardtii. *Sci Rep* **9**, 9856 (2019).

13. J. F. Allen, Cyclic, pseudocyclic and noncyclic photophosphorylation: new links in the chain. *Trends in Plant Science* **8**, 15–19 (2003).

14. K.-V. Dang, *et al.*, Combined Increases in Mitochondrial Cooperation and Oxygen Photoreduction Compensate for Deficiency in Cyclic Electron Flow in Chlamydomonas reinhardtii. *The Plant Cell* **26**, 3036–3050 (2014).

15. B. Bailleul, *et al.*, Energetic coupling between plastids and mitochondria drives CO_2 assimilation in diatoms. *Nature* **524**, 366–369 (2015).

16. A. Burlacot, G. Peltier, Y. Li-Beisson, Subcellular Energetics and Carbon Storage in Chlamydomonas. *Cells* **8**, 1154 (2019).

17. S. Saroussi, *et al.*, Alternative outlets for sustaining photosynthetic electron transport during dark-to-light transitions. *Proc Natl Acad Sci USA* **116**, 11518–11527 (2019).

18. G. Peltier, *et al.*, Algal CO2 capture is powered by alternative electron pathways of photosynthesis. 2023.08.08.552514 (2023).

19. C. Desplats, *et al.*, Characterization of Nda2, a Plastoquinone-reducing Type II NAD(P)H Dehydrogenase in Chlamydomonas Chloroplasts. *J. Biol. Chem.* **284**, 4148–4157 (2009).

20. D. Tolleter, *et al.*, Control of Hydrogen Photoproduction by the Proton Gradient Generated by Cyclic Electron Flow in Chlamydomonas reinhardtii[W]. *Plant Cell* **23**, 2619–2630 (2011).

21. D. Petroutsos, *et al.*, PGRL1 Participates in Iron-induced Remodeling of the Photosynthetic Apparatus and in Energy Metabolism in Chlamydomonas reinhardtii*. *Journal of Biological Chemistry* **284**, 32770–32781 (2009).

22. X. Johnson, *et al.*, Proton Gradient Regulation 5-Mediated Cyclic Electron Flow under ATP- or Redox-Limited Conditions: A Study of ΔATPase pgr5 and ΔrbcL pgr5 Mutants in the Green Alga Chlamydomonas reinhardtii1[C][W]. *Plant Physiol* **165**, 438–452 (2014).

23. F. Buchert, L. Mosebach, P. Gäbelein, M. Hippler, PGR5 is required for efficient Q cycle in the
cytochrome b6f complex during cyclic electron flow. *Biochemical Journal* **477**, 1631–1650
(2020).
24. A. P. Hertle, *et al.*, PGRL1 Is the Elusive Ferredoxin-Plastoquinone Reductase in Photosynthetic
Cyclic Electron Flow. *Molecular Cell* **49**, 511–523 (2013).
25. T. Rühle, *et al.*, PGRL2 triggers degradation of PGR5 in the absence of PGRL1. *Nat Commun* **12**,
3941 (2021).
26. G. DalCorso, *et al.*, A complex containing PGRL1 and PGR5 is involved in the switch between
linear and cyclic electron flow in Arabidopsis. *Cell* **132**, 273–285 (2008).
27. K. Tagawa, H. Y. Tsujimoto, D. I. Arnon, ROLE OF CHLOROPLAST FERREDOXIN IN THE ENERGY
CONVERSION PROCESS OF PHOTOSYNTHESIS. *Proc Natl Acad Sci U S A* **49**, 567–572 (1963).
28. Y. Munekage, *et al.*, PGR5 Is Involved in Cyclic Electron Flow around Photosystem I and Is
Essential for Photoprotection in Arabidopsis. *Cell* **110**, 361–371 (2002).
29. S. I. Saroussi, T. M. Wittkopp, A. R. Grossman, The Type II NADPH Dehydrogenase Facilitates
Cyclic Electron Flow, Energy-Dependent Quenching, and Chlororespiratory Metabolism during
Acclimation of Chlamydomonas reinhardtii to Nitrogen Deprivation. *Plant Physiology* **170**,
1975–1988 (2016).
30. C. Gerotto, *et al.*, Flavodiiron proteins act as safety valve for electrons in Physcomitrella patens.
PNAS **113**, 12322–12327 (2016).
31. F. Chaux, *et al.*, Flavodiiron Proteins Promote Fast and Transient O2 Photoreduction in
Chlamydomonas. *Plant Physiology* **174**, 1825–1836 (2017).
32. A. Burlacot, *et al.*, Alternative photosynthesis pathways drive the algal CO2-concentrating
mechanism. *Nature*, 1–6 (2022).
33. J. F. Allen, Photosynthesis of ATP—Electrons, Proton Pumps, Rotors, and Poise. *Cell* **110**, 273–
276 (2002).
34. G. Peers, *et al.*, An ancient light-harvesting protein is critical for the regulation of algal
photosynthesis. *Nature* **462**, 518–521 (2009).
35. G. Bonente, *et al.*, Analysis of LhcSR3, a Protein Essential for Feedback De-Excitation in the
Green Alga Chlamydomonas reinhardtii. *PLOS Biology* **9**, e1000577 (2011).
36. E. Erickson, S. Wakao, K. K. Niyogi, Light stress and photoprotection in Chlamydomonas
reinhardtii. *The Plant Journal* **82**, 449–465 (2015).
37. H. H. Stiehl, H. T. Witt, Quantitative treatment of the function of plastoquinone in
phostosynthesis. *Z Naturforsch B* **24**, 1588–1598 (1969).
38. C. Foyer, R. Furbank, J. Harbinson, P. Horton, The mechanisms contributing to photosynthetic
control of electron transport by carbon assimilation in leaves. *Photosynth Res* **25**, 83–100
(1990).
39. Y. Munekage, *et al.*, Cytochrome b6f mutation specifically affects thermal dissipation of
absorbed light energy in Arabidopsis. *The Plant Journal* **28**, 351–359 (2001).
40. L. A. Malone, M. S. Proctor, A. Hitchcock, C. N. Hunter, M. P. Johnson, Cytochrome b6f –
Orchestrator of photosynthetic electron transfer. *Biochimica et Biophysica Acta (BBA) -
Bioenergetics* **1862**, 148380 (2021).
41. M. Jokel, *et al.*, Chlamydomonas Flavodiiron Proteins Facilitate Acclimation to Anoxia During
Sulfur Deprivation. *Plant and Cell Physiology* **56**, 1598–1607 (2015).
42. M. Jokel, X. Johnson, G. Peltier, E.-M. Aro, Y. Allahverdiyeva, Hunting the main player enabling
Chlamydomonas reinhardtii growth under fluctuating light. *The Plant Journal* **94**, 822–835
(2018).
43. H. Chen, *et al.*, Ca2+-regulated cyclic electron flow supplies ATP for nitrogen starvation-induced
lipid biosynthesis in green alga. *Sci Rep* **5**, 15117 (2015).
44. F. Kong, *et al.*, Interorganelle Communication: Peroxisomal MALATE DEHYDROGENASE2
Connects Lipid Catabolism to Photosynthesis through Redox Coupling in Chlamydomonas. *The
Plant Cell* **30**, 1824–1847 (2018).

45. S. Wu, *et al.*, Elevated CO2 improves both lipid accumulation and growth rate in the glucose-6-
phosphate dehydrogenase engineered Phaeodactylum tricornutum. *Microb Cell Fact* **18** (2019).

46. G. Peltier, G. W. Schmidt, Chlororespiration: an adaptation to nitrogen deficiency in
Chlamydomonas reinhardtii. *Proceedings of the National Academy of Sciences* **88**, 4791–4795
(1991).

47. F. Chaux, *et al.*, PGRL1 and LHCSR3 Compensate for Each Other in Controlling Photosynthesis
and Avoiding Photosystem I Photoinhibition during High Light Acclimation of Chlamydomonas
Cells. *Mol Plant* **10**, 216–218 (2017).

48. G. DalCorso, *et al.*, A complex containing PGRL1 and PGR5 is involved in the switch between
linear and cyclic electron flow in Arabidopsis. *Cell* **132**, 273–285 (2008).

49. A. Burlacot, *et al.*, Flavodiiron-Mediated O2 Photoreduction Links H2 Production with CO2
Fixation during the Anaerobic Induction of Photosynthesis. *Plant Physiol.* **177**, 1639–1649
(2018).

50. S. Saroussi, E. Sanz-Luque, R. G. Kim, A. R. Grossman, Nutrient scavenging and energy
management: acclimation responses in nitrogen and sulfur deprived Chlamydomonas. *Current
Opinion in Plant Biology* **39**, 114–122 (2017).

51. J. Steinbeck, *et al.*, Deletion of Proton Gradient Regulation 5 (PGR5) and PGR5-Like 1 (PGRL1)
proteins promote sustainable light-driven hydrogen production in Chlamydomonas reinhardtii
due to increased PSII activity under sulfur deprivation. *Front Plant Sci* **6**, 892 (2015).

52. J. E. Johnson, J. A. Berry, The role of Cytochrome b6f in the control of steady-state
photosynthesis: a conceptual and quantitative model. *Photosynth Res* **148**, 101–136 (2021).

53. S. Saroussi, *et al.*, Restricting electron flow at cytochrome b6f when downstream electron
acceptors are severely limited. *Plant Physiology*, kiad185 (2023).

54. H. Yamamoto, T. Shikanai, PGR5-Dependent Cyclic Electron Flow Protects Photosystem I under
Fluctuating Light at Donor and Acceptor Sides. *Plant Physiology* **179**, 588–600 (2019).

55. J.-F. Penzler, *et al.*, Commonalities and specialties in photosynthetic functions of PROTON
GRADIENT REGULATION5 variants in Arabidopsis. *Plant Physiol* **190**, 1866–1882 (2022).

56. Q. Zhou, H. Yamamoto, T. Shikanai, Distinct contribution of two cyclic electron transport
pathways to P700 oxidation. *Plant Physiology* **192**, 326–341 (2023).

57. H. Yamamoto, T. Shikanai, PGR5-Dependent Cyclic Electron Flow Protects Photosystem I under
Fluctuating Light at Donor and Acceptor Sides. *Plant Physiology* **179**, 588–600 (2019).

58. A. Burlacot, Quantifying the roles of algal photosynthetic electron pathways: a milestone
towards photosynthetic robustness. *New Phytologist* **n/a**.

59. S. Wada, *et al.*, Flavodiiron Protein Substitutes for Cyclic Electron Flow without Competing CO2
Assimilation in Rice. *Plant Physiology* **176**, 1509–1518 (2018).

60. L. Basso, K. Sakoda, R. Kobayashi, W. Yamori, T. Shikanai, Flavodiiron proteins enhance the rate
of CO2 assimilation in Arabidopsis under fluctuating light intensity. *Plant Physiology*, kiac064
(2022).

61. R. Miller, *et al.*, Changes in Transcript Abundance in Chlamydomonas reinhardtii following
Nitrogen Deprivation Predict Diversion of Metabolism1[W][OA]. *Plant Physiol* **154**, 1737–1752
(2010).

62. I. K. Blaby, *et al.*, Systems-level analysis of nitrogen starvation-induced modifications of carbon
metabolism in a Chlamydomonas reinhardtii starchless mutant. *Plant Cell* **25**, 4305–4323
(2013).

63. K. Zienkiewicz, Z.-Y. Du, W. Ma, K. Vollheyde, C. Benning, Stress-induced neutral lipid
biosynthesis in microalgae — Molecular, cellular and physiological insights. *Biochimica et
Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids* **1861**, 1269–1281 (2016).

64. T. Takeuchi, C. Benning, Nitrogen-dependent coordination of cell cycle, quiescence and TAG
accumulation in Chlamydomonas. *Biotechnology for Biofuels* **12**, 292 (2019).

65. F. Kong, Y. Yamaoka, T. Ohama, Y. Lee, Y. Li-Beisson, Molecular Genetic Tools and Emerging
Synthetic Biology Strategies to Increase Cellular Oil Content in Chlamydomonas reinhardtii.
Plant Cell Physiol **60**, 1184–1196 (2019).

66. Y. Li-Beisson, J. J. Thelen, E. Fedosejevs, J. L. Harwood, The lipid biochemistry of eukaryotic
algae. *Progress in Lipid Research* **74**, 31–68 (2019).
67. P. Geigenberger, A. Kolbe, A. Tiessen, Redox regulation of carbon storage and partitioning in
response to light and sugars. *Journal of Experimental Botany* **56**, 1469–1479 (2005).
68. L. Michelet, *et al.*, Redox regulation of the Calvin–Benson cycle: something old, something new.
Front Plant Sci **4** (2013).
69. P. Geigenberger, A. R. Fernie, Metabolic Control of Redox and Redox Control of Metabolism in
Plants. *Antioxid Redox Signal* **21**, 1389–1421 (2014).
70. X. Li, *et al.*, A Galactoglycerolipid Lipase Is Required for Triacylglycerol Accumulation and
Survival Following Nitrogen Deprivation in Chlamydomonas reinhardtii. *The Plant Cell* **24**, 4670–
4686 (2012).
71. H. Chen, *et al.*, Ca(2+)-regulated cyclic electron flow supplies ATP for nitrogen starvation-
induced lipid biosynthesis in green alga. *Sci Rep* **5**, 15117 (2015).
72. X. Johnson, J. Alric, Interaction between Starch Breakdown, Acetate Assimilation, and
Photosynthetic Cyclic Electron Flow in Chlamydomonas reinhardtii. *J. Biol. Chem.* **287**, 26445–
26452 (2012).
73. X. Johnson, J. Alric, Central Carbon Metabolism and Electron Transport in Chlamydomonas
reinhardtii: Metabolic Constraints for Carbon Partitioning between Oil and Starch. *Eukaryot Cell*
12, 776–793 (2013).
74. M. Saint-Sorny, P. Brzezowski, S. Arrivault, J. Alric, X. Johnson, Interactions Between Carbon
Metabolism and Photosynthetic Electron Transport in a Chlamydomonas reinhardtii Mutant
Without CO2 Fixation by RuBisCO. *Frontiers in Plant Science* **13** (2022).
75. C. Goodson, R. Roth, Z. T. Wang, U. Goodenough, Structural correlates of cytoplasmic and
chloroplast lipid body synthesis in Chlamydomonas reinhardtii and stimulation of lipid body
production with acetate boost. *Eukaryot Cell* **10**, 1592–1606 (2011).
76. U. Goodenough, *et al.*, The path to triacylglyceride obesity in the sta6 strain of Chlamydomonas
reinhardtii. *Eukaryot Cell* **13**, 591–613 (2014).
77. K. Maxwell, G. N. Johnson, Chlorophyll fluorescence—a practical guide. *J Exp Bot* **51**, 659–668
(2000).
78. A. Burlacot, F. Burlacot, Y. Li-Beisson, G. Peltier, Membrane Inlet Mass Spectrometry: A
Powerful Tool for Algal Research. *Front Plant Sci* **11**, 1302 (2020).
79. F. Buchert, M. Scholz, M. Hippler, Electron transfer via cytochrome b 6 f complex displays
sensitivity to Antimycin A upon STT7 kinase activation. *Biochemical Journal* **479** (2022).
80. C. Klughammer, U. Schreiber, An improved method, using saturating light pulses, for the
determination of photosystem I quantum yield via P700+-absorbance changes at 830 nm.
Planta **192**, 261–268 (1994).
81. S. Upadhyaya, S. Agrawal, A. Gorakshakar, B. J. Rao, TOR kinase activity in Chlamydomonas
reinhardtii is modulated by cellular metabolic states. *FEBS Letters* **594**, 3122–3141 (2020).
82. J. L. Crespo, M. E. Pérez-Pérez, "Monitoring Autophagic Flux in the Model Single-Celled
Microalga Chlamydomonas reinhardtii" in *Plant Proteostasis: Methods and Protocols*, Methods
in Molecular Biology., L. M. Lois, M. Trujillo, Eds. (Springer US, 2023), pp. 123–134.
820