A robust normalized local filter to estimate occupancy directly from cryo-EM maps

Bjoern O. Forsberg [1,2,✉], **Pranav N.M. Shah** [2], **Alister Burt** [3]

[1] Department of Physiology and Pharmacology, Karolinska Institute, 171 77 Stockholm, Sweden. ; [2] Division of Structural Biology, University of Oxford, OX3 7BN Oxford, UK; [3] MRC Laboratory of Molecular Biology, Cambridge CB2 0QH, UK.

✉ **For correspondence:**
bjorn.forsberg@ki.se (B.O.F.)

Present address: Division of Structural Biology, University of Oxford, OX3 7BN Oxford, United Kingdom

Data availability: All EMDB data entries used for development of OccuPy are listed in the supplementary information. The software is publicly available at github.com/bforsbe/OccuPy, and pypi.org/project/OccuPy/. Instructions, tutorials, and measures for transparent reproducibility are hosted on occupy.readthedocs.io.

Funding: The present study was funded by the Swedish Research Council grant 2020-06413 and Cancer Research UK grant C20724/A26752.

Competing interests: The authors declare no competing interests.

Abstract

Cryo electron microscopy (cryo-EM) is used by biological research to visualize biomolecular complexes in 3D, but the heterogeneity of cryo-EM reconstructions is not easily estimated. Current processing paradigms nevertheless exert great effort to reduce flexibility and heterogeneity to improve the quality of the reconstruction. Clustering algorithms are typically employed to identify populations of data with reduced variability, but lack assessment of remaining heterogeneity. We have developed a fast and simple algorithm based on spatial filtering to estimate the heterogeneity of a reconstruction. In the absence of flexibility, this estimate approximates macromolecular component occupancy. We show that our implementation can derive reliable input parameters automatically, that the resulting estimate is accurate, and the reconstruction can be modified accordingly to emulate altered constituent occupancy. We demonstrate the utility of this method for cryo-EM map interpretation, particle-image signal subtraction, and its potential to promote divergence of clusters during conventionally employed maximum-likelihood classification methods.

Introduction

Protein, DNA and other molecular polymers sustain the fundamental processes of life, and structural biology is the study of their functions and interactions. Cryo-electron microscopy (cryo-EM) aims to visualize them by aligning and averaging noisy images of many individual macromolecules [1–3], producing a 3D scalar field known as a map or reconstruction. The reconstruction represents the local density or scattering potential of the atoms that make up the imaged macromolecule, but other representations also exist [4–9]. The ability to confidently deduce the molecular structure from the reconstruction crucially depends on its local quality, which varies due to variability or heterogeneity among the images used to build the reconstruction [10]. Methods for isolating homogeneous subsets through e.g. clustering is therefore employed [11–15], and commonly utilizes gradient descent methods [14, 16, 17]. Methods that parameterise the data have also been developed [7, 18–21]. The ability to separate data computationally by any method affords cryo-EM an unprecedented capacity to analyze the experimental particle distribution [22, 23], but this ability can be sensitive to user input. Existing alignment and clustering methods e.g. require a good initial (input) reconstruction to converge with good fidelity, since it provides sufficiently accurate es-

timates of image data parameters. However, lasting and potentially detrimental reference bias or *over-fitting*[24] may occur. This can distort the reconstruction and lead to misinterpretation or false features[25], and is exacerbated by the low signal-to-noise ratio (SNR) of cryo-EM data. So-called ab-initio 3D reconstructions can now be made without user input bias [14, 26, 27], but it is important to note that this nonetheless incurs downstream reference bias [24]. Current methods thus implicitly balance useful reference bias that permits convergence of the alignment against minimizing reinforcement of any reference bias arising from either spurious correlations in the data or the initial reconstruction. The imperative to minimize over-fitting has lead to methods that avoid detrimental reference bias [28–30], and some which implicitly isolates useful reference bias[31, 32] in a supervised fashion. Higher quality data and more robust reconstruction methods has also lead to more frequent iterative updates to the reconstruction than previously, based on smaller subsets of the data. This has implicitly increased reference bias within the reconstruction procedure by giving decent methods higher inertia. It is thus appropriate and more prudent than ever to investigate if and how unsupervised methods could improve its outcomes through supplementing relevant reference bias based on the heterogeneity of the reconstruction. However, until now no method has been published to estimate the heterogeneity of the reconstruction, which states could be inferred to exist in the data based on the final reconstruction and its heterogeneity, nor which states can be extrapolated from the initial reference during clustering. Recently established methods assign a measure of heterogeneity to the clusters [23, 33, 34], to attain more efficient clustering [35], and systematically choose an appropriate number of clusters [22]. Convergence of conventionally employed clustering is however largely subjective, and heterogeneity in the data may remain unresolved despite the apparent convergence of a given clustering algorithm[36]. As a result, biologically relevant differences may not be apparent across established clusters, and local reconstructions may suffer from undue incoherent averaging.

The present work formalizes the notion that cryo-EM reconstructions contain local information about latent heterogeneity[37]. Heterogeneity leads to local attenuation of the reconstructed density, a property we refer to as local scale. We provide OccuPy as a tool to estimate this local scale at all points within a reconstruction. OccuPy estimates local scale differences of arbitrary origin, e.g. due to flexibility and/or partial occupancy. OccuPy does however provide a method which reduces the influence of the former, to better approximate macromolecular occupancy. The local scale in occupancy mode can then further be used to modify the input to emulating reconstructions expected from more homogeneous image data. Since reference bias confines convergence, such modification is a justified approach to improve existing algorithms through weak biasing during clustering [34, 38]. OccuPy requires only a reconstruction as input and runs in seconds without the need for GPUs or HPC infrastructure. The approach is thus compatible with conventional iterative refinement and clustering tools. In this work, we establish the necessary formalism and tools to aid visual analysis of cryo-EM reconstructions and estimate local heterogeneity. A GUI is provided for ease of use, expanding the toolkit for reconstruction analysis available to cryo-EM researchers.

Results

Local scale is accurately estimated against synthetic data

To evaluate if the local scale can accurately estimate contrast degradation, we utilized simulated data with predefined, induced contrast degradation. A molecular model of malate dehydrogenase (PDB-1uxi, Fig. 1**a**) was altered by decreasing the occupancy of chain A, leaving all other atoms unchanged. Maps were generated using the theoretical electron scattering factors implemented in Gemmi (https://gemmi.readthedocs.io), and the local scale was estimated from such maps. Where occupancy-mode was used, a the low-pass filter used to neutralizes resolution-based differential contrast used a 8Å cutoff. As evident in Fig. 1**b**, contrast degradation due to partial occupancy is accurately estimated in the absence of flexibility. Systematic investigation of estimated scale at

atom positions however shows that when resolution is homogeneous, the local scale is more accurate than occupancy mode(1**e**). Either method appears robust to global changes in resolution(Fig. S1**c-d**). Because small-molecule (ligand) occupancy is of interest to quantify and enrich in biological structures, we investigated if OccuPy can do so accurately with a small enough filter window to segment the ligand from its binding pocket. We set the occupancy of all atoms of one NAD cofactor in PDB-1uxi to 0.4, leaving all other atoms at full occupancy. The density generated clearly shows that this co-factor is not visible at the same threshold as other elements (Fig. 1**c**). The same co-factor becomes evident following amplification, without being unduly exaggerated (Fig. 1**d**). Systematic analysis of the estimated scale of NAD however indicates that it is over-estimated at lower occupancy(Fig. 1**f**), due to the influence of neighboring regions under the dilating effect of the max-value filter used to estimate local scale.

Figure 1. Local scale estimation with ground truth. a, Malate dehydrogenase dimer PDB-1uxi, with chains colored individually. Two NAD cofactors are also shown in ball-and-stick representation. The occupancy of atoms in chain A was reduced and the local scale estimated from synthetic maps generated from this atomic model. Maps were also amplified with a power of 30. **b**, Amplified maps, annotated by the occupancy of chain A, and colored according to the estimated local scale. A grey-scale volume slice of the input is also shown (insets). **c**, The density generated by reducing the occupancy of atoms in one NAD co-factor to 0.4. **d**, The density in panel c, following amplification based on occupancy-mode local scale, also colored according to the estimated scale. **e**, Average local scale map value at atom positions of chain A, as function of occupancy. Shaded area represents the standard deviation. The ashed line denotes perfect estimation. **f**, as panel e, but for the NAD co-factors. **g**, as panel e, but as a function of the b-factor of atoms, at occupancy 0.5.

Finally, atomic B-factors were altered in chain A, and their occupancy reset to 100%, emulating lower resolution at full occupancy. Local scale reflects the decreased resolution, and as expected exaggerates local density if used for amplification (Fig. S1**e**). Occupancy-mode local scale suffers less of this effect (Fig. S1**f**). Systematic analysis shows that local scale has a strong dependence on the b-factor value, and that the occupancy-mode local scale has a much weaker such dependence(Fig. 1**g**). Taken together, OccuPy is able to estimate their local scale in a meaningful way, but variations in resolution pose a challenge to accurate estimation. Occupancy mode does decrease the dependence on resolution but the theoretically derived values of τ_n in combination with the default size of the normalizing region W here lead to under-estimation of local occupancy, which may require manual tuning.

Modification by local scale emulates homogeneous data

The present work has devised methods to modify a reconstruction proportionally to the estimated local scale, as exemplified in Fig. 1**d**. More generally, these modifications will decrease macromolecular heterogeneity by either increasing or decreasing regions with intermediate occupancy-mode local scale. Consequently, this makes it a natural method to modify the reference bias used to establish conditions of convergence in numerous current cryo-EM processing tools. We therefore evaluate how amplification, attenuation and differential sigmoid modification of local scale manifest in reconstructions with features encountered during cryo-EM processing, including heterogeneity, flexibility, and amorphous regions. First, EMD-14085 displays partial constituent occupancy and limited flexibility. As a result, we might expect the occupancy-mode local scale to be accurate but under-estimate macromolecular occupancy (Fig. 2**a**). In line with this, amplification restores low-occupancy components, but also amplifies some inherent noise. In part, this noise is elevated in an apparently spherical region which was presumably used a as a mask for classification to establish EMD-14085. The use of wholly amplified reconstructions for enforcing reference bias should thus be used with caution or under due noise reduction, through e.g. subsequent low-pass filtering. Conversely, attenuation acts conservatively at underestimated scale, and does not suffer any detrimental noise amplification effects. Sigmoid modification with a tuned pivot value achieves amplification without undue noise amplification, but omits components with very low occupancy.

Figure 2. Modification of reconstructed density by local scale a, EMD-14085 shown as a central slice, and iso-surface representation colored by estimated local scale. Further, it was amplified ($\gamma = 4$), attenuated ($\gamma = 2$), and sigmoid-modified ($\gamma = 5, \mu = 0.22$). **b**, EMD-3061, as in panel a, which sigmoid pivot $\mu = 0.5$. **c**, EMD-31466, as in panel a, which sigmoid pivot $\mu = 0.25$.

Second, we apply modifications to EMD-3061 (Fig. 2**b**), which has a region of detergent surrounding a transmembrane protein[39], since such regions are conventionally subject to signal subtraction[31] to reduce their influence on particle alignment and classification. Their resolution is ill-defined [40]; their physical extent can be determined with quantifiable accuracy, but any internal structure is effectively infinitely poor. Further, such a region is expected to have full occupancy since de-solvation of the transmembrane region is crucial for its structure, but its amorphous na-

ture results in incoherent averaging that reduces local scale. This is also observed; the detergent micelle displays reduced local scale. Because local scale does not represent occupancy in this case, it can not be compensated by direct filtering to emulate decreased heterogeneity. However, regions that display reduced local scale due to incoherent averaging may still be suitably modified for visualization and induced reference bias. Amplification in this case leads to grave exaggeration of local mass since local scale is severely underestimated due to resolution effects, even in occupancy-mode. This also displaces the reconstruction grey-scale outside the expected range, which further reduces its fidelity as the expected reconstruction from more homogeneous data. This again advocates that amplified reconstruction may be unsuitable for direct interpretation or use without further considerations. In the case of EMD-3061, attenuation and sigmoid modification curiously also cause undue modification, since no part of the input data is expected to be without a detergent micelle. The utility of such modification is however evident from its use in existing protocols for signal subtraction to reduce the influence of regions which cannot be coherently aligned, permitting structured regions to be better resolved. In this capacity, attenuation and sigmoid modification both appear well suited, signifying a direct way to weight reconstruction data by an objectively determined local property. To corroborate that such an approach is more broadly applicable to e.g. macromolecular flexibility, we subject EMD-31466 to the same analysis (Fig. 2c. This displays the same tendencies as EMD-3061, showing that local scale can be used to (de-)emphasize local regions of reconstructions in an automated or semi-automated manner using an objectively estimated attribute.

Variations in local mass can be accommodated

OccuPy is not equipped to consider the expected average density, the mass or scattering potential of constituent atoms, or a physical image formation model. Instead, it assumes that all regions with equal resolution and full occupancy will produce an identical voxel intensity distribution. This does not hold for atoms of unequal mass, which could potentially be estimated at a different scale. To evaluate how robust OccuPy is to such situations, we first consider the case of a nucleosome protein-DNA complex (EMD-32148), since the phosphate-rich backbone could lead to under-estimated scale of protein components. This does not seem to be the case (Fig. 3a). Next, we examined a high-resolution (1.22Å) reconstruction of apoferritin (EMD-11638). First, the region $\mathbf{W}$ used to determine full scale and normalize the scale estimate was reduced to a single pixel, which makes the local scale estimate sensitive to single pixels with high values, e.g. at high-mass atoms. Methionine side-chain sulfur atoms are thus estimated at higher scale than surrounding protein (Fig. 3b). The default size of the region $\mathbf{W}$ (Eq. (2), methods) efficiently diminishes the influence of such heavy atoms by considering the max-value distribution of the highest contrast region found (Fig. 3c-d), unless it dominates an entire region $\mathbf{W}$. This nevertheless emphasizes that full scale (or occupancy) is a relative term in the absence of true underlying mass, here defined based on the size of $\mathbf{W}$ and the percentile τ. A smaller region $\mathbf{W}$ allows local mass to define full scale. Conversely, a larger $\mathbf{W}$ will reduce the influence of local mass differences. However, setting $\mathbf{W}$ too large may cause systematic overestimation of the scale since no region of this size will be uniform at full scale. The default size of $\mathbf{W}$ in OccuPy is chosen to consider the size granularity of biomolecular complexes typically reconstructed by cryo-EM, while neglecting individual atoms. This can be adjusted by the user. To further examine the potential pitfalls of the method, we also estimate the scale of respiratory complex I (EMD-13611), which contains a number of FeS clusters(Fig. 3e). Naturally, these FeS clusters are estimated at full scale. In spite of this, the protein content is only slightly under-estimated(Fig. 3f). A decreased value of τ can further compensate for this (Fig. 3g), and the size of the region W may be increased to define full scale (Fig. 3h). Both these parameter adjustments reduce the influence of high local values, but the latter offers a direct interpretation as redefining the granularity of the estimate through the size of the reference region $\mathbf{W}$.

Figure 3. Local scale estimation under variation of atomic mass. a, EMD-32148 shows no difference in estimated scale of DNA relative to protein. **b**, local scale of EMD-11638 without normalizing region **W** shows sensitivity to local mass. **c**, local scale of EMD-11638 with default parameters. **d**, occupancy-mode local scale of EMD-11638 with default parameters. **e**, Maximum-value projection of the EMD-13611, showing FeS-clusters. **f**, local scale of EMD-13611 with default parameters. **g**, local scale of EMD-13611 with larger normalizing region **W**. **h**, local scale of EMD-13611 with reduced percentile τ.

Macromolecular flexibility can be partially accommodated

Local resolution variability within published reconstructions is common and primarily due to flexibility [41]. We therefore evaluate OccuPy against a a reconstruction of a flexible helical assembly (F-actin EMD-30171) for which negligible variation in occupancy is expected (Fig. 4**a**). In line with expectation, the local scale correlates with decreased resolution further from the box center(Fig. 4**b-c**). The local scale in occupancy mode is less affected, but still indicates decreased occupancy, which is not expected. The estimated occupancy can be validated by performing full amplification (power $\gamma = 30$), followed by a low-pass filtration. Doing so for EMD-30171 reveals that mass has been exaggerated in such regions(Fig. 4**d**), indicating that the occupancy was strictly underestimated. This can not be ameliorated by increasing the low-pass cutoff, in line with expectation (see methods). To contrast these findings, EMD-12104, exhibits partial occupancy but negligible flexibility, in which case low-pass filtration of the amplified map shows no emphasis on the regions estimated at low occupancy scale(Fig. 4**h**), indicating accurate occupancy estimation and appropriate amplification. To further illustrate the fidelity of the local scale estimate as compared to existing methods, EMD-13015 was used. The local resolution varies as estimated by ResMap [42] and RELION [13] (Fig. 5**a-b**). The local scale reproduces the relative contrast estimation (Fig. 5**c**). In occupancy-mode, the local scale is more homogeneous, indicating that these differences can be attributed largely to flexibility (Fig. 5**d**). The occupancy estimated by LocOccupancy [43](Fig. 5**e**) however correlates strongly to the estimated resolution.

Local scale can be estimated from sharpened maps

Maps are typically post-processed after reconstruction to maximize information and fidelity. Global B-factor estimation and compensation is most common [44], but local filtering [45–47] and machine-learning [48] are also employed. To investigate if post-processing introduces or obscures expected features to the detriment of a faithful scale estimate, we modify EMD-3943 by common post-processing methods and then estimate local scale, since this reconstruction contains differential resolution in its subunits and partial occupancy of a bound recycling factor (RRF). First, the local scale estimate of a map sharpened by a global B-factor exhibits a larger range (Fig. S2**d**), since

Figure 4. Local scale estimates in the presence of flexibility. a, Input map (EMD-30171). **b-c**, Amplified map, colored according to estimated local scale and occupancy-mode local scale, respectively. Panels **a-c** are shown at the same threshold value. **d**, Low-pass filtration of the amplified map, showing exaggerated mass in low-scale regions. **e**, The PDB-7b9k atomic model, with the catalytic trans-acetylase domain (CTD) in white and a lipoyl domain (LD) shown in black. **f**, The cryo-EM reconstruction EMD-12104, showing apparent partial occupancy with negligible flexibility. **g**, The amplified reconstruction, utilizing and colored by the occupancy-mode local scale. **h**, Low-pass filtration of the amplified map.

Figure 5. Comparison of local resolution and occupancy estimation methods. a, The local resolution of EMD-13015 was estimated using ResMap [42] in the range 2.3-4.3, with a resolution step of 0.25Å. **b**, The local resolution was estimated using RELION [13] with a spacing of 10Å. **c-d**, The local scale and and occupancy-mode local scale estimated using OccuPy. **e**, The occupancy estimated using LocOccupancy [43], using a resolution range of 2.5-6Å.

---

local contrast is increased in proportion to SNR. The local scale estimate in occupancy mode is however similar to that estimated from the unmodified map. While OccuPy is not intended to be used on post-processed maps, it nonetheless appears permissible. Next, a local-resolution filtered reconstruction displays decreased local scale in the subunit at lower estimated resolution. This reconstruction is however highly similar to the original map, both in terms of the local scale estimate and that in occupancy mode(Fig. S2**g-i**). Finally, a reconstruction 'enhanced' by deepEMhancer [48] shows a very uniform full scale, apart from the RRF, which is likely at a lower occupancy. Indeed the RRF is lower in occupancy-mode as well. Curiously, the subunit occupancy scale is inverted with respect to the unmodified input map (Fig. S2**k**), indicating that DeepEMhancer results in alterations in local mass dependent on the local resolution. Based on this, we surmise that reconstruction post-processed by machine-learning methods are not suited for use in OccuPy without further

prior validation.

Improved robustness of confidence estimate through solvent definition

OccuPy estimates a solvent model and subsequent confidence map to avoid solvent noise amplification. In some cases this solvent model is incorrectly estimated due to unexpected solvent characteristics. In this event, an additional input mask can be provided to limit the regions of the input map considered when determining the solvent model. We denote this a solvent definition, since it does not mask the output. We illustrate its use on the map of the asymmetric unit of a viral capsid re-framed such that the solvent volume is only 22% of the cubic map volume (42% of the map radius sphere) (Fig. S3**a**). The capsid interior also contains a disordered component with low variance but higher mean than the solvent. An accurate solvent definition (Fig. S3**b**) that excludes all protein content and capsid interior results in a single Gaussian solvent peak and accurate confidence. If the viral capsid spike and interior is not excluded by the solvent definition (Fig. S3**c**), the solvent model and confidence is still accurately estimated. (Fig. S3**f**). This demonstrates that the solvent definition does not strictly enforce what is amplified, permitting map modification outside the provided solvent definition.

Discussion

This work demonstrates that local scale can be used to quantify macromolecular occupancy in cryo-EM reconstructions without a molecular model, and used as a meaningful means to modify them. However, accurate estimation of local scale may require parameter tuning response to reconstruction characteristics, which could lead to user confirmation bias. LocOccupancy [43] is the only other method designed to approach quantification of compositional variation, meriting a direct comparison. LocOccupancy requires a resolution range to be specified which in some sense dictates the granularity of the estimation, whereas OccuPy instead estimates local properties with a minimal kernel given the resolution of the reconstruction, and regulates the granularity of the occupancy estimate through the normalization region $\mathbf{W}$. Both LocOccupancy and OccuPy are also dependent on a percentile cutoff, which signifies different characteristics in each implementation. LocOccupancy sets this value to 0.25, signifying the top percentile that defines full occupancy in some sense, whereas OccuPy automatically sets a theoretically optimal value to minimize the probability of both over- and under-estimating the occupancy. In further comparison, LocOccupancy naturally maps each region to the [0, 1]-range, whereas OccuPy instead normalizes by a value lower than the global maximum and clamps the estimate to the [0, 1]-range. A clear benefit of LocOccupancy is that it effectively marginalizes the occupancy estimate over the desired resolution range, while OccuPy makes no such provision and estimates occupancy-mode local scale under the assumption that variations in local resolution have been neutralized. Despite this apparent disadvantage, Fig. 5 suggests that OccuPys occupancy-mode is able to disregard resolution-dependent contrast degradation better. Fig 5 also shows that the local scale is an accurate estimate of relative local quality, which is tantamount to resolution. Resolution is however a contested term in cryo-EM [40, 49]. By consensus, the spatial frequency at which the global Fourier shell correlation (FSC) drops below a given significance [44] is quoted as the best resolution at which the reconstruction can be reliably interpreted. Local metrics also permit variations to be quantified under the term "resolution". As discussed elsewhere [50] these measures are not identical, and the term resolution is thus not well defined. It is however clear that resolution correlates positively with data amount and quality, and how coherently it can be averaged. This in turn principally depends on macromolecular flexibility and occupancy, which mirrors that of the local scale estimated here. In this sense, the OccuPy local scale does constitute a true estimate of relative local resolution. However OccuPy assumes that the density originates from identical point scatters. Due to their variation in mass and occupancy, the local scale is not a universal estimator of resolution. When resolution becomes poorer than the physical spacing of the point source of scattering, their environment also influences the scale

estimate, as shown in Fig. S1. Bearing these points in mind we conclude that the local scale is an accurate estimate of the relative local resolution, but that this estimate is dependent on properties that e.g. FSC-based resolution estimates are independent of.

Taken together, we find OccuPy to be a reliable tool available to quantitatively estimate macromolecular occupancy within cryo-EM reconstructions, and the only tool able to modify this occupancy in a meaningful way. Through its GUI, users can directly adjust estimation parameters such as input low-pass frequency, kernel size, and normalization tile-size, and visualize the results. The solvent model can also be directly evaluated, and the optional input solvent definition constructed. From this interface, further evaluation is also facilitated by invoking UCSF ChimeraX [51] with a command-script that is part of the default output. This will also display complementary visualizations to evaluate the results. OccuPy may also be used to improve signal subtraction by providing accurate subtraction masks, and its capacity to accurately estimate local scale with minimal user input suggests that this capacity could be employed in iterative refinement procedures. In order for OccuPy to remain a viable step in iterative optimization strategies for cryo-EM data clustering, the scale estimation and modification must be comparable in speed to the gradient methods it will complement. Comparing the spatial filtering approach to existing methods of local property estimation shows that it completes in a few seconds for any reasonably sized input (Table S1). Other available methods of local property estimation are clearly infeasible for the purpose of assessing cluster heterogeneity within an iterative clustering algorithm. The local scale established here can also be considered a highly accurate estimator of local properties of the reconstruction, which can be used to visualize and modify reconstructions in a meaningful way. OccuPy is thus fit for the purpose of improving current optimization procedures, where it could be used with a weak power to bias reference-based alignment and clustering. This stands to be validated in practice. OccuPy nonetheless represents the first incarnation of local spatial processing of constituent heterogeneity cryo-EM reconstructions, and stands to be developed further to benefit future reconstruction analysis and refinement algorithms broadly.

Methods & Materials

A spatial filter to estimate local scale

The best-resolved region in a cryo-EM reconstruction displays the highest contrast. We axiomatically define the images used to make the reconstruction to be completely homogeneous with respect to this region. Globally homogeneous input data would thus result in a theoretically ideal cryo-EM reconstruction F_{ideal}. In F_{ideal} all non-solvent regions exhibit identical local scale. In practice, contrast is attenuated through local flexibility and/or partial occupancy. The observed map F can thus be considered F_{ideal} degraded by the local scale S:

$$F = F_{\text{ideal}} \cdot S \tag{1}$$

where $< \cdot >$ denotes pixel-wise multiplication and $\{S | S_i \in [0, 1] \forall i\}$. S is thus a normalized estimate of the of the local signal strength. We estimates S using a windowed max-value filter over the local neighborhood $\mathbf{V}_i$ of each pixel i, i.e. by spatial filtering. This measures the width of the voxel value distribution sampled within $\mathbf{V}_i$, which signifies signal above noise in cryo-EM reconstructions. A max-value filter is fast, insensitive to the inclusion of solvent, and robust to noise even for very small window sizes. It is also capable of preserving sharp transitions, with respect to a morphological dilation.

To normalize the estimated scale at each pixel i, we have to determine the maximal expected value given the finite sampling in regions around i. To do so, we first subdivide the reconstruction into non-exhaustive regions $\mathbf{W}_j$. A percentile filter with parameter τ is used for robustness to high-value outliers, with due consideration to follow. The region j with the largest such value is used to determine "full scale" at a given percentile τ, and is used as a denominator to normalize the scale

estimate:

$$\hat{S}_i = \frac{\max\limits_{i \in \mathbf{V}_i}(F_i)}{\max\limits_{\{\mathbf{W}_j\}}\left(\mathrm{per}_\tau\limits_{i \in \mathbf{W}_j}(F_i)\right)}\Big\lceil_{[0,1]} \tag{2}$$

where we have defined a function $\lceil$ which clamps values to a specified interval

$$x\Big\lceil_{[a,b]} \equiv \max\big(a, \min(x, b)\big) \tag{3}$$

since the max-value reduction over $\mathbf{V}_i$ may exceed that of the percentile τ in the full-scale region found. The established procedure permits the intensity distribution of a small region to define full scale/contrast, and does not enforce any specific portion of the reconstruction to be assigned any nominal scale value. It also obviates the need for masking areas of interest. The set of regions $\{\mathbf{W}_j\}$ need not be exhaustive since the denominator is globally defined. Occupy thus utilizes a sparse set of j regions $\{\mathbf{W}_j\}$, evenly distributed across the reconstruction. By default, 8000 (20^3) regions of 1728 (12^3) voxels are used, which represents a fundamental granularity of biomolecular components that works in a broad range of test-cases. This is a tunable parameter in the present implementation.

We now derive a reasonable choice for the percentile τ. We first note that as long as sufficiently many regions $\mathbf{W}$ are sampled, $\hat{S}_i$ may be an overestimate of S_i by at most $1 - \tau$. On the other hand, $\hat{S}_i$ would instead be an underestimate if the number of elements $n_{\mathbf{V}}$ within $\mathbf{V}$ is unlikely to have sampled as high as the percentile given by τ. As a compromise, we seek the percentile τ_n of the distribution that equals the confidence that the maximum value of n samples is also τ_n. Considering the reconstruction values in a local region to be a random variable X, we solve

$$G_{Y_n}(x) = 1 - G_X(x) \tag{4}$$

where G is the cumulative distribution function (CDF) and Y is the maximum value distribution

$$Y_n = \max\{X_1, ..., X_n\} \tag{5}$$

However, the CDF of Y_n can be simplified as

$$G_{Y_n}(x) = P(Y_n \leq x) = \prod_{i=1}^{n} P(X_i \leq x) = G_X(x)^n \tag{6}$$

Consequently, we can rewrite Eq. (4) as

$$G_X(x)^n = 1 - G_X(x) \tag{7}$$

from which we see that τ_n is simply the only positive real root to the polynomial

$$x^n + x - 1 = 0 \tag{8}$$

This result is independent of the underlying distribution of X. However, the CDF of Eq. (6) is only separable under the assumption that adjacent voxels are independent. By setting τ in Eq. (2) to τ_n as dependent on the kernel size $n_{\mathbf{V}}$, we are thus guaranteed that S_i at most over-estimated by $1 - \tau_n$, and under-estimated with a probability $1 - \tau_n$. By solving Eq. (8) we find that $\tau_n \geq 0.9$ for $n_{\mathbf{V}} \geq 27$. This places narrow error bounds on the scale estimate for any realistic kernel size. In reality, the voxels sampled within V are not independent, which reduces the effective sampling number compared to $n_{\mathbf{V}}$, such that τ should reasonably be set lower than τ_n. Additionally, a region $\mathbf{W}$ is assumed to exist that has homogeneous and full scale. When this is not the case, the normalizing value in the denominator of Eq. (2) may be increased due to e.g. high-mass atoms in the full-scale region, or conversely decrease due to inclusion of solvent. This will lead to systematic under- and over-estimation of local scale, respectively.

Finally, we note that in OccuPy, the kernel size k (voxels along each dimension) is automatically calculated in resolution dependent manner as the smallest odd integer larger than

$$k = 2 \cdot r/d \tag{9}$$

where r is the applied input low-pass filter or resolution of the input reconstruction and d the input voxel size.

Establishing an occupancy-mode

We first note that $\hat{F}_{\text{ideal}}$ can be found as an estimate of F_{ideal} as

$$\hat{F}_{\text{ideal}} = F \cdot \hat{S}^{-1} \tag{10}$$

The instability of inverse filtering at low values of S is handled by a complementary confidence estimation to follow. From Eq. (10), it is clear that S constitutes a spatial filter capable of modifying the estimated macromolecular occupancy without masking or segmentation. The local scale S is however a measure of contrast attenuation, which correlates with both resolution (peak broadening) and occupancy (peak reduction). Modification of a reconstruction through the use of S as a spatial filter is however only appropriate to compensate or further exaggerate peak reduction due to occupancy, not resolution. To omit resolution-dependent effects, we employ the simple procedure of estimating the local scale from a low-pass filtered copy of the input reconstruction. We term this occupancy-mode local scale. The low-pass procedure achieves resolution-dependent attenuation by the same magnitude at all points in the reconstruction. All regions estimated must thus be affected by the low-pass filter, lest regions better resolved should be estimated at higher scale by virtue of higher resolution. In addition, we implicitly assume that local density values do not suffer differential influence by peak broadening of nearby values. This is violated e.g. comparing points internal to the protein core to those near solvent. This violation becomes more severe at lower resolution, so that any omission of resolution-dependent effects is countered by detrimental convolution of local contrast. The occupancy-mode local scale will thus be difficult to establish faithfully by the low-pass filter approach when the reconstruction displays large variations in local resolution due to flexibility. The general local scale is however accurate with arbitrary internal variation in local contrast.

γ-modification of estimated scale

To achieve attenuation or amplification or partial occupancy, Occupy implements a proportional and inverse power scaling of the estimated occupancy-mode scale S by a power γ analogous to conventional γ-correction.

$$F_{+\gamma} = F \cdot S_{+\gamma} = F \cdot S^{1/\gamma - 1} \ \left(= \hat{F}_{\text{ideal}} \cdot S^{1/\gamma} \right) \tag{11}$$

$$F_{-\gamma} = F \cdot S_{-\gamma} = F \cdot S^{\gamma - 1} \ \left(= \hat{F}_{\text{ideal}} \cdot S^{\gamma} \right) \tag{12}$$

Amplification in this context is thus attenuation of $\hat{F}_{\text{ideal}}$ by $\hat{S}^{1/\gamma}$, signifying less heterogeneity than what was estimated from the input data. Attenuation is conversely application of more heterogeneity than $\hat{S}$. For consistency, the present implementation only permits $\gamma \geq 1$. $\hat{F}_{\text{ideal}}$ corresponds to $F_{+\infty}$, a direct inverse filter of the input by $\hat{S}$. γ-modification is illustrated in Fig. S5**a** exemplified in Fig. **??d-e**.

Attenuation of a reconstruction can be directly applied to attain precise and automatic masks for particle subtraction workflows [31]. Conventionally, a mask M is provided where each voxel value $m \in [0, 1]$ determines the retention of image values:

$$I_{\text{subtracted}} = I - \mathbf{P}_{\phi} \left(F \cdot (1 - M) \right) \tag{13}$$

where $\mathbf{P}$ is the projection operator and ϕ is the alignment of image I. Conventionally the mask is constructed by manual volume segmentation and user manipulation. Here, the estimated scale S and the desired output scale S' can be used to formulate an optimal mask M:

$$\hat{F}_{\text{ideal}} \cdot S' = \hat{F}_{\text{ideal}} \cdot S - (1 - M) \cdot \hat{F}_{\text{ideal}} \cdot S \Leftrightarrow \tag{14}$$

$$M = S'/S \tag{15}$$

M is on the interval $[0, 1]$ as long as $S' < S$, i.e. when the desired output scale S' is lower than the input scale S. This signifies strict attenuation, which is reasonable as components are to be subtracted. Occupy provides a simple interface to create such a mask, with any necessary adjustments.

Sigmoid modification

OccuPy also implements a sigmoid modification, which attenuates components below specified local scale value, but also amplifies components above the same value. Like γ-modification, sigmoid modification is dependent on the power γ. Again, $\gamma = 1$ signifies no change, and increasing values result in increased modification. The additional parameter μ signifies the threshold scale value that remains unmodified and is thus denoted the pivot value. Formally, the scale S is altered as

$$S_{\gamma,\mu} = \left(1 + \left(\frac{S \cdot (1 - \mu_s)}{\mu_s \cdot (1 - S)} \right)^{-\gamma} \right)^{-1} \tag{16}$$

where

$$\mu_s = \left(\left(\frac{1 - \mu}{\mu} \right)^{1 - \frac{1}{\gamma}} + 1 \right)^{-1} \tag{17}$$

The sigmoid modification is thus formulated as

$$F_{\gamma,\mu} = F \cdot S_{\gamma,\mu} \cdot S^{-1} \ \left(= \hat{F}_{\text{ideal}} \cdot S_{\gamma,\mu} \right) \tag{18}$$

Like attenuation, sigmoid modification can be used to construct a mask for particle subtraction according to Eq. (15), under the provision that the sigmoid mapping is adjusted so that values above μ remain unmodified. Sigmoid modification is illustrated in Fig. S5**b-c** and exemplified in Fig. **??f**.

Suppression of solvent modification

Amplification is a form of inverse filtering, which is sensitive to noise. OccuPy therefore estimates a solvent confidence map to avoid amplification of solvent. While local signal-to-noise ratio (SNR) is a reasonable estimate of the confidence in the reconstructed voxel value, it is too strict for the purposes here. More leniently, we establish the confidence as the relative probability of observing a given voxel value in content over solvent. To do so, we determine a solvent model Θ as a Gaussian fit to the main peak of the reconstruction histogram, much like previous methods [52], resulting in a confidence map for each voxel. This is exemplified in Fig. **??c**. OccuPy does not identify solvent regions prior to this fit, but instead relies on the assumption that the majority of the reconstruction volume is composed of solvent and has a pronounced peak in the image histogram. If not, the solvent variance is typically over-estimated, leading to decreased confidence of low-scale regions. To permit more accurate fitting in these cases, a solvent definition can be supplied in the form of a mask which covers the non-solvent regions of the input reconstruction. This is not employed as a mask, but instead delineates the regions omitted when fitting the Gaussian solvent model. Consequently, it does not restrict the confidence map and permits scale modification outside the provided solvent definition. This is show in Fig. S3.

Formally, the confidence C computed as the ratio of the probability that each voxel pertains to the solvent model or not:

$$C_i = \frac{P(A_i \in [F_i - \delta/2, F_i + \delta/2]|F)}{P(F_i|\Theta)} \tag{19}$$

where δ is the bin with of the reconstruction histogram. The spatial filter C_i essentially constitutes a soft solvent mask that is applied to suppress solvent, without segmenting the reconstruction or enforcing a hard threshold.

$$Q_{\pm\gamma} = F_{\pm\gamma} \cdot C \tag{20}$$

$$Q_{\mu,\nu} = F_{\mu,\nu} \cdot C \tag{21}$$

Retention of solvent

OccuPy intends to modify reconstructions to mimic the expectation if the input data were more homogeneous. Solvent should thus not be excluded as described by Eq. (20). OccuPy therefore utilizes the inverse confidence to retain the original solvent background.

$$R_{\pm\gamma} = Q_{\pm\gamma} + F \cdot (1 - C) \tag{22}$$

$$R_{\gamma,\mu} = Q_{\gamma,\mu} + F \cdot (1 - C) \tag{23}$$

Further to this, attenuated noise is compensated in proportion to the attenuation:

$$T_{-\gamma} = R_{-\gamma} + N \cdot (S - S^\gamma) \cdot C \tag{24}$$

where N is noise generated to have the same distribution and spectral properties as the solvent of the input reconstruction.

References

1. Sigworth, F. J., Doerschuk, P. C., Carazo, J. M. & Scheres, S. H. W. *An introduction to maximum-likelihood methods in cryo-EM* 1st ed. **C,** 263–294 (Elsevier Inc., 2010).
2. Danev, R., Yanagisawa, H. & Kikkawa, M. Cryo-Electron Microscopy Methodology : Current Aspects and Future Directions. *Trends Biochem. Sci.* **44,** 837–848 (2019).
3. *Single-particle Cryo-EM of Biological Macromolecules* 1st ed. (eds Glaeser, R. M., Nogales, E. & Chiu, W.) (Biophysical Society IOP Series, 2021).
4. Kawabata, T. Gaussian-input Gaussian mixture model for representing density maps and atomic models. *J. Struct. Biol.* **203,** 1–16 (2018).
5. Donati, L., Nilchian, M., Sorzano, C. O. S. & Unser, M. Fast multiscale reconstruction for Cryo-EM. *J. Struct. Biol.* **204,** 543–554 (2018).
6. Bonomi, M. *et al.* Bayesian Weighing of Electron Cryo-Microscopy Data for Integrative Structural Modeling. *Structure* **27,** 175–188 (2019).
7. Zhong, E. D., Bepler, T., Berger, B. & Davis, J. H. CryoDRGN: reconstruction of heterogeneous cryo-EM structures using neural networks. *Nat. Methods* **18,** 176–185 (2021).
8. Ranno, N. & Si, D. Neural representations of cryo-EM maps and a graph-based interpretation. *BMC Bioinformatics* **23,** 1–19 (2022).
9. Urzhumtsev, A. G. & Lunin, V. Y. Analytic representation of inhomogeneous-resolution maps of three- dimensional scalar fields. *bioRxiv,* 1–20 (2022).
10. Lyumkis, D. Challenges and opportunities in cryo-EM single-particle analysis. *J. Biol. Chem.* **294,** 5181–5197 (2019).
11. Tang, G. *et al.* EMAN2: An extensible image processing suite for electron microscopy. *J. Struct. Biol.* **157,** 38–46 (2007).
12. Lyumkis, D., Brilot, A. F., Theobald, D. L. & Grigorieff, N. Likelihood-based classification of cryo-EM images using FREALIGN. *J. Struct. Biol.* **183,** 377–388 (2008).

13. Scheres, S. H. W. RELION: Implementation of a Bayesian approach to cryo-EM structure determination. *J. Struct. Biol.* **180,** 519–530 (2012).
14. Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. CryoSPARC: Algorithms for rapid unsupervised cryo-EM structure determination. *Nat. Methods* **14,** 290–296 (2017).
15. Singer, A. & Sigworth, F. J. Computational Methods for Single-Particle Electron Cryomicroscopy. *Annu. Rev. ofBiomedical Data Sci.* **3,** 163–190 (2020).
16. Hu, M. *et al.* A particle-filter framework for robust cryo-EM 3D reconstruction. *Nat. Methods* **15,** 1083–1089 (2018).
17. Kimanius, D., Dong, L., Sharov, G., Nakane, T. & Scheres, S. H. W. New tools for automated cryo-EM single-particle analysis in RELION-4.0. *Biochem. J.* **478,** 4169–4185 (2021).
18. Moscovich, A., Halevi, A., Andén, J. & Singer, A. Cryo-EM reconstruction of continuous heterogeneity by Laplacian spectral volumes. *Inverse Probl.* **36,** 1–31 (2020).
19. Chen, M. & Ludtke, S. J. Deep learning-based mixed-dimensional Gaussian mixture model for characterizing variability in cryo-EM. *Nat. Methods* **18,** 930–936 (2021).
20. Barreto, J. G. *et al.* A Bayesian approach to extracting free-energy profiles from cryo-electron microscopy experiments. *Sci. Rep.* **11,** 1–15 (2021).
21. Kinman, L. F., Powell, B. M., Zhong, E. D., Berger, B. & Davis, J. H. Uncovering structural ensembles from single particle cryo-EM data using cryoDRGN. *bioRxiv* (2022).
22. Zhou, Y., Moscovich, A. & Bartesaghi, A. Data-driven determination of number of discrete conformations in single-particle cryo-EM. *Comput. Methods Programs Biomed.* **221,** 106892 (2022).
23. Rabuck-gibbons, J. N., Lyumkis, D. & Williamson, J. R. Quantitative mining of compositional heterogeneity in cryo-EM datasets of ribosome assembly intermediates. *Structure* **30,** 498–509 (2022).
24. Sorzano, C. O. *et al.* On bias, variance, overfitting, gold standard and consensus in single-particle analysis by cryo-electron microscopy. *Acta Crystallogr. Sect. D Struct. Biol.* **78,** 410–423 (2022).
25. Henderson, R. Avoiding the pitfalls of single particle cryo-electron microscopy : Einstein from noise. *PNAS* **110,** 18037–18041 (2013).
26. Elmlund, D. & Elmlund, H. SIMPLE: Software for ab initio reconstruction of heterogeneous single-particles. *J. Struct. Biol.* **180,** 420–427 (2012).
27. Zivanov, J. *et al.* New tools for automated high-resolution cryo-EM structure determination in RELION-3. *Elife* **7;e42166,** 1–22 (2018).
28. Chen, S. *et al.* High-resolution noise substitution to measure overfitting and validate resolution in 3D structure determination by single particle electron cryomicroscopy. *Ultramicroscopy* **135,** 24–35 (2013).
29. Ramlaul, K., Palmer, C. M., Nakane, T. & Aylett, C. H. S. Mitigating local over-fitting during single particle reconstruction with SIDESPLITTER. *J. Struct. Biol.* **211,** 107545 (2020).
30. Punjani, A., Zhang, H. & Fleet, D. J. Non-uniform refinement: adaptive regularization improves single-particle cryo-EM reconstruction. *Nat. Methods* **17,** 1214–1221 (2020).
31. Bai, X. C., Rajendra, E., Yang, G., Shi, Y. & Scheres, S. H. Sampling the conformational space of the catalytic subunit of human g-secretase. *Elife* **4:e11182,** 1–19 (2015).
32. Nakane, T., Kimanius, D., Lindahl, E. & Scheres, S. H. Characterisation of molecular motions in cryo-EM single-particle data by multi-body refinement in RELION. *Elife* **7,** e36861 (2018).
33. Aizenbud, Y. & Shkolnisky, Y. A max-cut approach to heterogeneity in cryo-electron microscopy. *J. Math. Anal. Appl.* **479,** 1004–1029 (2019).
34. Yin, S., Zhang, B., Yang, Y., Huang, Y. & Shen, H.-b. Clustering Enhancement of Noisy Cryo-Electron Microscopy Single- Particle Images with a Network Structural Similarity Metric. *J. Chem. Inf. Model.* **59,** 1658–1667 (2019).
35. Zhou, Y., Moscovich, A., Bendory, T. & Bartesaghi, A. Unsupervised particle sorting for high-resolution single-particle cryo-EM. *Inverse Probl.* **36,** 1–17 (2020).

36. Forsberg, B., Aibara, S., Howard, R. J., Mortezaei, N. & Lindahl, E. Arrangement and symmetry of the fungal E3BP-containing core of the pyruvate dehydrogenase complex. *Nat. Commun.* **11,** 1–10 (2020).

37. Matsumoto, S. *et al.* Extraction of protein dynamics information from cryo-EM maps using deep learning. *Nat. Mach. Intell.* **3,** 153–160 (2021).

38. Lei, H. & Yang, Y. CDAE: A Cascade of Denoising Autoencoders for Noise Reduction in the Clustering of Single-Particle Cryo-EM Images. *Front. Genet.* **11,** 1–9 (2021).

39. Piper, S. J., Johnson, R. M., Wootten, D. & Sexton, P. M. Membranes under the Magnetic Lens : A Dive into the Diverse World of Membrane Protein Structures Using Cryo-EM. *Chem. Rev.* **122,** 13989–14017 (2022).

40. Liao, H. Y. & Frank, J. Definition and Estimation of Resolution in Single-Particle Reconstructions. *Structure* **18,** 768–775 (2010).

41. Palamini, M., Canciani, A. & Forneris, F. Identifying and Visualizing Macromolecular Flexibility in Structural Biology. *Front. Mol. Biosci.* **3,** 1–17 (2016).

42. Kucukelbir, A., Sigworth, F. J. & Tagare, H. D. Quantifying the local resolution of cryo-EM density maps. *Nat. Methods* **11,** 63–65 (2014).

43. Kaur, S. *et al.* Local computational methods to improve the interpretability and analysis of cryo-EM maps. *Nat. Commun.* **12,** 1–12 (2021).

44. Rosenthal, P. B. & Henderson, R. Optimal determination of particle orientation, absolute hand, and contrast loss in single-particle electron cryomicroscopy. *J. Mol. Biol.* **333,** 721–745 (2003).

45. Jakobi, A. J., Wilmanns, M. & Sachse, C. Model-based local density sharpening of cryo-EM maps. *Elife* **6:e27131,** 1–26 (2017).

46. Vargas, J., Gómez-Edrero, J. A., Quiroga, J. A. & Alonso, J. Enhancement of Cryo-EM maps by a multiscale tubular filter. *Opt. Express* **30,** 4515–4527 (2022).

47. Bharadwaj, A. & Jakobi, A. J. Electron scattering properties of biological macromolecules and their use for cryo-EM map sharpening. *Faraday Discuss.* (2022).

48. Sanchez-garcia, R. *et al.* DeepEMhancer: a deep learning solution for cryo-EM volume post-processing. *Commun. Biol.* **874,** 1–8 (2021).

49. Cardone, G., Heymann, J. B. & Steven, A. C. One number does not fit all : Mapping local variations in resolution in cryo-EM reconstructions. *J. Struct. Biol.* **184,** 226–236 (2013).

50. Vilas, J. L., Heymann, J. B., Tagare, H. D., Carazo, J. M. & Sorzano, C. O. S. ScienceDirect Local resolution estimates of cryoEM reconstructions. *Curr. Opin. Struct. Biol.* **64,** 74–78 (2020).

51. Pettersen, E. F. *et al.* UCSF ChimeraX : Structure visualization for researchers , educators , and developers. *Protein Sci.* **8,** 70–82 (2021).

52. Beckers, M. & Sachse, C. Thresholding of cryo-EM density maps by false discovery rate control. *IUCrJ* **6,** 18–33 (2019).

53. Burnley, T., Palmer, C. M. & Winn, M. Recent developments in the CCP-EM software suite research papers. *Acta Crystallogr. Sect. D Struct. Biol.* **73,** 469–477 (2017).

54. Harris, C. R. *et al.* Array programming with NumPy. *Nature* **585,** 357–362 (2020).

55. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat. Methods* **17,** 261–272 (2020).

56. Hunter, B. J. D. Matplotlib: A 2D graphics environment. *Comput. Sci. Eng.* **9,** 90–95 (2007).

Acknowledgments

This preprint was created using the LaPreprint template by Mikkel Roald-Arbøl. The authors are grateful to those who provided feedback and functionality testing, including Dari Kimanius, Marta Carroni and Loic Carrique. The authors are grateful to Jesse Hopkins at BioCAT for help with software packaging.

Author contributions

Role	Contribution	Authors
Conceptualization	Conceived ideas or experimental design	1,2
Methodology	Devised method or theory	1
Software	Development and implementation	1,2,3
Investigation	Performed experiments/data collection	1
Validation	Data analysis and interpretation	1
Writing	Provided revisions to scientific content of the manuscript	1,2,3
Writing	Provided other revisions to the manuscript	1,2,3
Funding	Provided funding	1
Supervision	Principal investigator	1

Table 1. Author contributions. Defined with reference to The Company of Biologists.
1. B.O.F. **2**, P.N.M.S. **3**, A.B.

Competing interests

The authors declare no competing interests.

Funding

The present study was funded by the Swedish Research Council grant 2020-06413 and Cancer Research UK grant C20724/A26752.

Data availability

All EMDB data entries used for development of OccuPy are listed in the supplementary information. The software is publicly available at github.com/bforsbe/OccuPy, and pypi.org/project/OccuPy/. Instructions, tutorials, and measures for transparent reproducibility are hosted on occupy.readthedocs.io.

Supplementary

Map	Size	ResMap	RELION	LocOcc.	OccuPy
EMD-33442	200	172	6'300	480	4
EMD-12104	300	572	79'660	828	6
EMD-7770	338	335	25'770	1'092	10
EMD-13015	420	886	449'260	696	11

Table S1. The execution time of local property estimation by ResMap [42], RELION [13], and LocOccupancy [48]. Reported time is in CPU-seconds, as RELION and LocOccupancy execute with process-level parallelism.

Supplementary Figures

Supplementary Figure S1. Evaluation of local scale against synthetic data. All shown densities are fully amplified and colored according to the indicated local scale. **a**, The occupancy field of all atoms in chain A of the malate dehydrogenase dimer PDB-1uxi structure file was altered to reflect the indicated occupancy. Chain B, and the two nicotinamide adenine-dinucleotide (NAD) cofactors were left at full occupancy (1.0). Theoretical densities at 8Å resolution were generated from the atomic model using the program gemmi. **b**, Occupancy-mode local scale at various theoretical occupancies and 8Å resolution density. **c**, Local scale at fixed partial occupancy of chain A, at various generated map resolutions. **d**, Occupancy-mode as in panel **c**. **e**, Local scale after altering B-factors of chain A as indicated, leaving occupancy of all atoms at 100%. **f**, Occupancy-mode as in panel **e**.

Supplementary Figure S2. Local scale of modified maps. All views are cutaway. **a-b**, EMD-3943 colored according to the estimated local scale and occupancy-mode local scale, respectively. **c**, EMD-3943 sigmoid-modified with power $\nu = 5$, and pivot $\mu = 0.2$, using the estimated occupancy-mode scale. **d-f**, as panels **a-c** following global B-factor sharpening in RELION [13]. **g-i**, as panels **a-c** following local FSC-filtering in RELION [13]. **j-l**, as panels **a-c** following sharpening by the neural-net based DeepEMhancer [48].

Supplementary Figure S3. The use of a solvent definition to aid solvent model estimation. a, The reconstruction of an asymmetric unit of a viral capsid is re-framed within the original reconstruction, with a very small fractional volume of solvent, leading to an overestimated solvent variance. The solvent model is shown in green, the confidence in red, and the histogram of the data used to establish the solvent model is shown in black. The histogram of the input map is shown in gray. **b**, An accurate solvent definition and the resulting solvent model. **c**, A less accurate solvent definition. **d**, The estimated confidence without a solvent definition. **e**, The estimated confidence with the accurate solvent definition in panel **b**. **f**, The estimated confidence with the less accurate solvent definition in panel **c**.

Supplementary Figure S4. Radial cutoff kernels used in OccuPy. The kernels are numbered by kernel size and order. The number of sampling points within the kernel is plotted. Further to this, the value of τ_n is plotted for a number of sampling size n. Manually altering automatically derived parameters may cause less spherical kernels in some cases, e.g. a 5x5x5.

Supplementary Figure S5. Mapping of modified scale. **a**, Amplification and attenuation of estimated map scale by the power coefficient γ. **b**, Sigmoid modification, at variable power γ but fixed pivot μ. **c**, Sigmoid modification, at fixed power γ and variable pivot μ. $S_{\gamma,\mu}$

Supplementary Figure S6. The OccuPy GUI. The GUI is written in PyQt5, using the python libraries mrcfile [53], numpy [54], scipy [55], and matplotlib [56]. The user may open and visualize the input map, modify the automatically generated kernel settings, and estimate local scale. The scale, solvent model, and confidence can be directly inspected, and settings adjusted. Modifications can also be previewed in real-time, and generated with additional output control. An exhaustive log documents processing parameters is also maintained. Annotated visualization of estimated scale and modified maps is facilitated through a command script for UCSF ChimeraX [51].

EMDB entries used for development

Entries in bold are used in the paper. One of the deposited half maps was used for all entries where this was available, since deposited "main" maps can be post-processed, masked, and/or a composite reconstruction without disclosure, at the discretion of the depositor.

EMD-0056	EMD-11077	EMD-13163	EMD-22855	EMD-26477
EMD-0355	EMD-11266	EMD-13241	EMD-22959	EMD-26513
EMD-0452	EMD-11268	EMD-13378	EMD-23006	**EMD-30171**
EMD-0667	EMD-11432	**EMD-13611**	EMD-23015	EMD-30547
EMD-0910	EMD-11576	EMD-13619	EMD-23017	EMD-30675
EMD-3061	**EMD-11638**	EMD-13620	EMD-23082	EMD-31213
EMD-3943	**EMD-12104**	EMD-13886	EMD-23096	EMD-31432
EMD-2984	EMD-12112	EMD-13961	EMD-23099	**EMD-31466**
EMD-4058	EMD-12197	EMD-13991	EMD-23114	EMD-32121
EMD-4470	EMD-12202	**EMD-14085**	EMD-23121	EMD-32126
EMD-5529	EMD-12204	EMD-14122	EMD-23658	**EMD-32148**
EMD-6840	EMD-12220	EMD-14336	EMD-24095	EMD-32593
EMD-7770	EMD-12286	EMD-14369	EMD-24176	EMD-32612
EMD-8194	EMD-12341	EMD-14428	EMD-24536	EMD-32875
EMD-8332	EMD-12870	EMD-20546	EMD-24717	EMD-33707
EMD-8333	EMD-12885	EMD-21485	EMD-24823	EMD-34028
EMD-9108	**EMD-13015**	EMD-21847	EMD-24976	
EMD-09215	EMD-13033	EMD-22074	EMD-25121	
EMD-10160	EMD-13078	EMD-22075	EMD-25170	
EMD-10978	EMD-13162	EMD-22358	EMD-26357	