Evolution of two gene networks underlying adaptation to drought stress in the wild

tomato *Solanum chilense*

Kai Wei*[1], Saida Sharifova*[2], Xiaoyun Zhao[1], Neelima Sinha[3], Hokuto Nakayama[4], Aurélien

Tellier[#1], Gustavo A Silva-Arias[#1,5]

[1]Professorship for Population Genetics, Department of Life Science Systems, School of Life

Sciences, Technical University of Munich, Liesel-Beckmann Strasse 2, 85354 Freising,

Germany

[2]Department of Life Sciences, Graduate School of Science, Art and Technology, Khazar

University, Mahsati 41, AZ1096, Baku, Azerbaijan

[3]Department of Plant Biology, University of California Davis, One Shields Avenue, Davis, CA

95616, USA

[4]Department of Biological Sciences, The University of Tokyo, 113-0033, Tokyo, Japan

[5]Instituto de Ciencias Naturales, Facultad de Ciencias, Universidad Nacional de Colombia,

Sede Bogotá, Bogotá, Colombia

* these authors contributed equally

[#]Correspondence email: aurelien.tellier@tum.de, gasilvaa@unal.edu.co

Abstract

Drought stress is a key factor limiting plant growth and the colonization of arid habitats by

plants. Here, we study the evolution of gene expression response to drought stress in a wild

tomato, *Solanum chilense* naturally occurring around the Atacama Desert in South America.

We conduct a transcriptome analysis of plants under standard and drought experimental

conditions to understand the evolution of drought-response gene networks. We identify two

main regulatory networks corresponding to two typical drought-responsive strategies: cell

cycle and fundamental metabolic processes. We estimate the age of the genes in these

networks and the age of the gene expression network, revealing that the metabolic network

has a younger origin and more variable transcriptome than the cell-cycle network. Combining

with analyses of population genetics, we found that a higher proportion of the metabolic

network genes show signatures of recent positive selection underlying recent adaptation

within *S. chilense,* while the cell-cycle network appears of ancient origin and is more

conserved. For both networks, however, we find that genes showing older age of selective

sweeps are the more connected in the network. Adaptation to southern arid habitats over the

last 50,000 years occurred in *S. chilense* by adaptive changes core genes with substantial

network rewiring and subsequently by smaller changes at peripheral genes.

Introduction

Drought stress is one of the major environmental constraints negatively influencing plant development and preventing plant growth, resulting in decreased yield in agriculture and as a constraining factor for colonization of arid or hyper-arid habitats (Ciais et al. 2005; Juenger 2013). Plants respond to water-insufficiency through multiple strategies underpinned by various physiological and developmental processes, such as storage of internal water to avoid tissue damage and tolerance (endurance) to drought stress to maintain the growth process (Basu et al. 2016). These strategies involve many biological functions such as increasing the metabolic activity of some tissues, i.e. root water uptake and closing stomata, or activation of metabolic pathways including phytohormone signaling, antioxidant and metabolite production in order to regulate osmotic processes (Rodrigues et al. 2019). Drought response involves numerous quantitative and polygenic traits acting in (complex) gene co-expression networks (GCN). To improve crops and predict the evolutionary responses of plant species under the current and predicted global water deficits, it is thus of interest to pinpoint and decipher the evolutionary history of the relevant GCNs underpinning the adaptation of wild plants to arid or hyper-arid habitats (Gehan et al. 2015).

Comparative transcriptomics involving the inference of gene co-expression patterns show that many GCNs are conserved through the tree of life (Stuart et al. 2003; Gerstein et al. 2014; Zarrineh et al. 2014; Crow et al. 2022). Moreover, phylogenetic and developmental studies have demonstrated that many physiological, structural, and regulatory innovations to cope drought stress have arisen throughout the history of plants, many of them even predating the emergence of land plants (Jill Harrison 2017; de Vries et al. 2018; de Vries and Archibald 2018; Mustafin et al. 2019; Wang et al. 2020; Bowles et al. 2021). Several conserved GCNs can be observed in fundamental biological processes such as protein metabolism, cell cycle, and photosynthesis and well as key traits such wood formation (Stuart et al. 2003; Ficklin and Feltus 2011; Zinkgraf et al. 2020).

A key question in functional and evolutionary genomics is thus to link GCN evolution

and (relatively) short-scale evolutionary processes such as adaptation and

population/species divergence in order to assess the relative importance of contingency,

exaptation and evolution of novel genes (duplication, neofunctionalization) allowing

colonization of novel habitats. Two main hypotheses are formulated. First, highly conserved

sub-networks (so-called hubs or kernels) evolve under strong purifying selection to ensure

the functionality of the GCNs (Papakostas et al. 2014; Josephs et al. 2017; Mähler et al.

2017; Masalia et al. 2017), so that genetic variation is only found at (less connected) genes

at the periphery of the GCNs that may be the target of positive natural selection (Flowers et

al. 2007; Kim et al. 2007; Luisi et al. 2015; Erwin 2020). However, this argument is likely

because the novel habitats may not differ much from the original one, so that only minor

adjustments in the GCNs are enough to provide adaptation. This is also in line with so-called

developmental systems drift (DSD; True and Haag 2001), that predicts GCN rewiring only

occurs in 'flexible' (sub-)modules with the accumulation of neutral variation that keep the

network function intact until a new viable function (phenotype or developmental pathway)

appears. Second, despite the general belief that genes with higher connectivity evolve at a

slower rate, there is also evidence that changes at central genes (with higher connectivity)

can be responsible for the short-term response to selection (Jovelin and Phillips 2009; Luisi

et al. 2015) and promote rewiring of the GCN (Koubkova-Yu et al. 2018). Thus, highly

connected genes may be targets of positive selection during environmental change, *e.g.*

adaptation to novel habitats, even though these genes experience purifying selection in

stable environments (Hämälä et al. 2020). Indeed, if the second hypothesis is correct, we

expect a correlation between the age of positive selection and the connectivity of a gene in a

network, but no correlation under the first hypothesis.

To test these hypotheses, we reveal the selective forces (positive versus purifying

selection) acting on different components of the networks (hub vs peripheral genes) across

species/lineages adapted to contrasting conditions, and correlate the signals of positive

selection with gene connectivity in the wild tomato species *Solanum chilense*. Wild tomatoes

are a model of interest as their diversification is accompanied by the exploration of wide

environmental gradients along the Pacific coast of South America (from tropical to subtropical,

coastal to high mountain, and wet to extremely dry regions; Nakazato et al. 2010; Haak et al.

2014). In addition, the infra-specific diversification within *S. chilense* resulted in several

lineages with strong environmental differentiation (Raduski and Igić 2021; Wei et al. 2023).

Populations of *S. chilense* are challenged by prolonged drought, with the most severe

drought conditions occurring in the southern part of the range. Wild relative tomato species

such *S. chilense*, *S. sitiens* and *S. pennellii* become well-established systems to study

tolerance strategies to survive in extreme environments (Bolger et al. 2014; Martínez et al.

2014; Tapia et al. 2016; Kashyap et al. 2020; Blanchard-Gros et al. 2021; Molitor et al. 2021;

Barrera-Ayala et al. 2023). In a previous study, we assayed for evidences of positive

selection in 30 fully sequenced genomes of *S. chilense* to identify candidate genes

underpinning adaptation along the species range. We found genes with putative functions

related to root hair development and cell homeostasis as being likely involved in drought

stress tolerance (Wei et al. 2023). However, to date, most research in *S. chilense* has

focused on the evolution of a few genes potentially involved in abiotic stress response

(Fischer et al. 2011; Mboup et al. 2012; Fischer et al. 2013; Böndel et al. 2015; Nosenko et al.

2016; Böndel et al. 2018), and we still lack information regarding the evolutionary

mechanisms driving drought tolerance in this species.

Our aim is to study the GCN evolution underpinning *S. chilense* adaptation to arid

habitats. We identify drought stress responsive gene regulatory networks combining multiple

analyses of transcriptome data of *S. chilense* and focus on two networks involved in cell-

cycle and metabolic processes. Furthermore, we infer the evolutionary processes at these

two networks across three different evolutionary levels (tree of life/plants, species and

population) by computation of transcriptome indices to explore the evolutionary age and

sequence divergence of the drought responsive transcriptome. We then analyze the

emergence of adaptive variation in the identified drought-responsive genes of these networks

and the association to gene connectivity.

Results

Drought experiments and transcriptome analyses

Plants of *S. chilense* growing either under well-watered or a moderate water stress regimes
(hereafter, control and drought) showed clear morphological differences by the time of tissue
collection. Plant growth and ramification was boosted in well-watered group while plants
under drought were smaller and slow-growing. Hence, on the day 12, newly expanded leaf
and shoot apices were collected for the expression analysis of stress-responsive genes and
four biological replicates were used for all RNA-Seq experiments from each tissue type.

We analyze short-read transcriptome data from 16 libraries aligned to the reference
genome of *S. chilense* (Dataset S1). A total of 27,832 genes are identified to be expressed in
the 16 libraries (Dataset S2), of which 1,536 genes are uniquely expressed in drought
condition and 1,767 genes in control condition (Dataset S2). A principal component analysis
(PCA) based on the gene expression profiles reveals consistent clustering primarily
associated with the experimental conditions (control and drought) and secondarily to the
developmental stages (leaf and shoot apex) (Figure 1A). PC1 accounts for 28.17% of the
expression variability and separates the libraries from the two experimental conditions,
indicating transcriptome remodeling between drought and control conditions. Libraries from
different developmental stages are separated along the PC2 axis (accounting for 18.24% of
the variance), supporting tissue age transcriptome specificity. Consistently, the transcriptome
similarity analysis between libraries reveals that the watering conditions explain the major
differences between treatments (Figure 1B). Hierarchical clustering also revealed that the
transcriptomes were grouped mainly according to water deficit intensity, rather than by tissue
type (Figure 1C), demonstrating the predominant effect of the stress response in
transcriptome remodeling. Therefore, we thereafter focus on comparing the transcriptome
profiles of the drought and control experimental conditions.

Identification of gene networks involved in drought stress

We identified gene networks involved in drought response in *S. chilense* based on differential

expression analysis and weighted gene co-expression network analysis (WGCNA). First,

three sets of differential expression genes (DEGs) are identified from three drought/control

comparison groups (full data set, only leaf and only shoot apex tissues) (Figure 2A; Dataset

S3; log2FoldChange ≥ 1, FDR *P* ≤ 0.001). A total of 4,905 DEGs are obtained in three

comparison groups, of which 2,484 DEGs (1,235 up-regulated and 1,249 down-regulated in

drought transcriptome) are shared in three comparison groups (Figure 2B). We deduce that

these shared DEGs correspond to a core functionally drought-responsive network.

In construction of gene co-expression networks (GCNs), we do not directly used DEGs

in WGCNA as suggested by the developer of WGCNA, because DEGs are invalid for

assumption of the scale-free topology. Therefore, a set of 16,181 genes after filtering from all

expressed genes were used in WGCNA (see methods), and clustered into seven co-

expression modules named after different colors. The module sizes range from 183 up to

5,364 genes (Figure 2C, Dataset S4). Among the identified co-expression modules, the blue

module (3,852 genes) shows significantly positive correlation with control condition and

negative correlation with drought condition (Figure 2C, Kendall's test, *P* = 2.2e-11). In

contrast, the turquoise module (5,364 genes) is significantly positively correlated with drought

condition and negatively correlated with control condition (Figure 2C, Kendall's test, *P* =

2.34e-13). In addition, the genes within blue and turquoise modules are observed to show

higher connectivity than other modules (Figure S1, Kolmogorov-Smirnov test on connectivity

measure, *P* = 2.41e-17), indicating higher interaction and closer correspondence in biological

process among genes within each module in response to water deprivation.

We next check the overlap between 2,484 DEGs and co-expression modules to

confirm that blue and turquoise modules are associated with drought stress in *S. chilense*

(Table S1). DEGs share far more genes with the blue and turquoise modules than with other

co-expression modules. Almost all shared DEGs (2,302 genes out of 2,484) are found in the

blue and turquoise modules. This confirms that blue and turquoise modules are two sets of

co-expressed drought stress responsive genes. The overlapping DEGs and module genes

are extracted to constitute the now refined two high-confidence subsets of the blue and

turquoise modules and comprising 1,223 and 1,079 genes, respectively.

To independently support regulatory relationships among genes identified in the two co-

expression networks, we identify transcription factors (TFs) and transcription factor binding

sites (TFBSs) for the two subsets of genes. Therefore, we extract the genes that can bind to

one another (Table S2) from the two high-confidence subsets, which we hereafter name as

sub-blue (686 genes) and sub-turquoise (948 genes), respectively (Dataset S5). These

results show that genes in the sub-blue and sub-turquoise networks not only show specific

co-expression patterns, but they show also predicted interact between the TFs and TFBSs.

Subsequently, the co-expression network is reconstructed using the same steps for the set of

genes of the sub-blue and sub-turquoise networks. Higher connectivity is observed in the

sub-turquoise network (Figure S2, Kolmogorov-Smirnov test on connectivity measure, P =

0.002), suggesting a closer regulatory relationship among genes in the sub-turquoise than in

the sub-blue network.

To verify the robustness of two drought-responsive network (sub-blue and sub-

turquoise) across different tomato species (and thus the generality of our results), we

employed also the same pipeline to construct GCNs combined to transcriptomic data of *S.

pennelliii* (PRJEB5809; Bolger et al. 2014) and *S. lycopersicum* (PRJNA812356; Yang et al.

2022) (Dataset S1). Those transcriptomes also exhibit difference due to water conditions in

the analysis of PCA, correlations and hierarchical clustering (Figure S3). We first observe

that the 1,837 (74%) genes in the DEG sets based on only *S. chilense* overlap to the

combined DEG set of *S. pennellii* and *S. lycopersicum* (Figure S4A and S4B). The GCNs

from combined transcriptome profiles consistently show two networks as we find in *S.

chilense* alone (Figure 2C and S4C). We find that 576 (84%) and 778 (82%) genes overlap

to the sub-blue and sub-turquoise networks, respectively (Figure S4D and S4E). In addition,

our two drought-responsive networks (sub-blue and sub-turquoise) also overlap with

previous study based on drought transcriptomes of *S. lycopersicum* (Nicolas et al. 2022).

Nearly 60% of drought-responsive genes in the two networks are observed as response to

water stress in *S. lycopersicum*. These overlap rates suggest that our two drought-

responsive networks are present and perform similar functions in different tomato species.

We therefore concentrate on these two networks: sub-blue (686 genes) and sub-turquoise

(948 genes).

Functional enrichment analysis of drought-responsive GCNs

We assess whether the two identified gene networks (sub-blue and sub-turquoise) show

functional differences. The gene ontology (GO) enrichment reveals that sub-blue network is

significantly enriched ($P < 0.05$) in cell cycle and regulation biological processes, including

replication and modification of genetic information, ribosome production and assembly,

cytoskeleton organization, among others (Figure 3A; Table S3). Conversely, the sub-

turquoise network is enriched in biological processes related to response of physiological

and metabolic processes to water shortage and heat, including some metabolic processes,

signal pathways, changes of stomata and cuticle, amongst other processes (Figure 3A; Table

S3). These functional differences suggest that genes in the two sub-networks are activated

and expressed in different cellular compartments. Consistent with the mentioned biological

process, the sub-blue network genes are mainly enriched in cellular components in the

nucleus, including nucleolus, chromosome, nuclear envelope, and ribosome (Figure 3B;

Table S4). These cellular components are at the center of cell division processes. On the

other hand, the sub-turquoise network is enriched in cellular components related to

metabolism processes, such as complexes and membrane structures in the cell (Figure 3B;

Table S4). Many studies have indicated that modulation in the cell cycle and fundamental

metabolism are two main strategies in response to drought stress (Gupta et al. 2020; Yang et

al. 2021; Nicolas et al. 2022). We focus, thereafter, on these two sub-networks and from now

on, the sub-blue network is referred to as the *cell-cycle network* and the sub-turquoise as the

metabolic network.

Evolutionary age of drought-responsive transcriptome in *S. chilense*

To generate a comprehensive understanding of the emergence of the identified drought-

responsive GCNs, we estimate the transcriptome ages of the identified cell cycle and

fundamental metabolism networks. For that, we build phylostratigraphic profiles for all genes

of the two GCNs, summarizing the gene emergence in 18 stages of plant evolution or

phylostrata (PS): PS1 representing the emergence of oldest genes (at the time of the first

cellular organisms) to PS18 for the most recent genes (*i.e.* present only in *S. chilense*). The

PS18 shares no homologue genes with any other species in the nr (non-redundant protein)

databases of NCBI (Figure 4A and 4B, Dataset S6). Most genes in the two analyzed GCNs

(76.79% in metabolic network and 65.45% in cell-cycle network) are assigned to three main

PS: Cellular organisms (PS1), Land plants (Embryophyta; PS5) and Flowering plants

(Magnoliopsida; PS8) (Figure 4A). This suggests that the two drought-responsive GCNs we

identify have an ancient origin and the components are fairly conserved across the tree of

life/plants. Therefore, many drought-responsive pathways likely emerged during the

colonization of land by plants (PS5), but many others could derive from exaptation processes

from GCNs involved in the core cell process (PS1) or reproductive organ differentiation of

flowering plants (PS8). Interestingly, the cell-cycle network shows older origin ages (with

more genes (43.73%) assigned to the PS1-3), while the metabolic network presents a larger

proportion (48.52%) of genes originating in PS8 (Figure 4A and 4B). Under drought

conditions, we also find that cell-cycle network genes of almost all PS ages are down-

regulated, while genes of the metabolic network are up-regulated (Figure S5).

Furthermore, we estimate the age of cell-cycle and metabolic GCNs using the

transcriptome age index (TAI). We do not find a significant difference of TAI between control

and drought samples based on 1,000 randomly selected genes from non-drought responsive

genes (Figure S6A; Kolmogorov-Smirnov test, P = 0.34), while in cell-cycle and metabolic

networks, the mean evolutionary ages of the transcriptomes are significantly different

between drought and control conditions (Figure 4C; Kolmogorov-Smirnov test, P = 0.03). The

TAI profile would be expected to be a flat horizontal line if genes' ages remain constant

across the transcriptomes. In addition, a higher TAI value implies that evolutionary younger

genes are preferentially expressed at the corresponding condition/developmental stage. We

observe higher TAI in drought samples, supporting that the drought-responsive genes exhibit

a younger transcriptome age than genes expressed under control conditions. Moreover, TAI

of the metabolic GCN is significantly higher than the cell-cycle (Figure 4C; Kolmogorov-

Smirnov test, P = 12.51e-7), supporting the previous result that transcriptome ages of the

genes in the cell-cycle are older than in the metabolic GCNs.

The contributions of the different PS to the TAI profiles also show notable patterns

between the cell-cycle and metabolic GCNs (Figure 4D and 4E). On one hand, early

divergent genes (PS1 to PS7) show more constant transcriptome age in all conditions and

the genes with ages in PS1, PS5 and PS8 appeared as remarkably important in two GCNs.

On the other hand, late-emerging genes (PS8 to PS18) contribute increasingly with their age

to the differential expression patterns between control and drought samples, indicating that

younger drought-responsive genes are differentially expressed under drought stress in both

GCNs (as observed in Domazet-Lošo and Tautz 2010; Piasecka et al. 2013). Remarkably,

the youngest genes in PS18 (only found in *S. chilense*), also present a higher contribution in

the metabolic GCN, suggesting that these genes are involved in either speciation or local

adaptation of *S. chilense* to drought conditions. Note that younger genes (PS9 to PS18) in

the cell-cycle GCN hardly contribute to the TAI profile (Figure 4D and 4E).

Divergence of the drought tolerance transcriptome in *S. chilense*

To drill down into the evaluation of the drought-response mechanisms at the species level,

we calculate the TDI index, which represents the mean sequence divergence of a

transcriptome. A total of 10 divergence strata (DS) are constructed based on the sequence

divergence between genes of *S. chilense* and *S. pennellii* by computing the Ka/Ks ratio

(Figure 5A; Figure S7; Dataset S6). The distributions of the Ka/Ks ratio per gene for both

GCNs indicate the action of purifying selection, which confirms the conservation of most of

drought-responsive genes at the species level. Consistent with the phylostratigraphic

patterns, the purifying selection signals in the cell-cycle GCN (Ka/Ks = 0.279 ± 0.333) are

higher than in the metabolic GCN (Ka/Ks = 0.329 ± 0.331) (Kolmogorov-Smirnov test, P =

2.34e-11; Figure 5A; Table S5). In addition, higher TDI values are observed in the drought

samples (Figure 5B) suggesting that the expressed genes we identify in the two GCNs

exhibit a more conserved transcriptome profile under control condition compared to drought

condition (Kolmogorov-Smirnov test, P = 0.004). No significant difference is found between

control and drought samples based on 1,000 random genes (Kolmogorov-Smirnov test, P =

0.17; Figure S6B). This result supports that different selective pressures act on *S. chilense*

GCNs across conditions. In accordance with the TAI results, the transcriptome of the

metabolic GCN appears to exhibit a higher transcriptome divergence than the cell-cycle GCN

(Figure 5B; Kolmogorov-Smirnov test, P = 2.25e-7). Moreover, the low TDI in the cell-cycle

GCN and larger TDI differences between drought and control transcriptomes also suggest

that regulation of the cell-cycle is likely an ancestral (older) strategy of stress response, not

involved in the speciation process. The transcriptome of the cell-cycle GCN may have been

evolving and changing in older times, and reached a conserved structure in recent times.

Conversely, changes of metabolic pathways and rewiring of the metabolic GCN may appear

to be more pronounced and/or common in recent times.

The contributions of the low divergence DS classes (low Ka/Ks in DS1 to DS5) in the

cell-cycle GCN (~ 50% of the genes) are larger than in the metabolic GCN (DS1 to DS5

about 30%), especially in DS1 (lowest Ka/Ks ratio; Figure 5C and 5D). This indicates that

purifying selection is acting on genes of the cell-cycle GCN, possibly constraining further

changes. In contrast, the metabolic network genes show about 70% contributions in high DS

(higher Ka/Ks ratio in DS6 to DS10), especially in DS10 (highest Ka/Ks ratio), indicating that

genes in the metabolic network evolve under weaker purifying selection and that recent

evolutionary changes occurred.

As a summary, from the deep phylogenetic to the species level, the TAI profile of the

cell-cycle network is mainly composed of older phylostrata (PS1 to PS8), while new genes

contribute about 20% to the TAI profile of the metabolic network (Figure 4D and 4E). This

indicates that the gene expression levels of the cell-cycle network have likely been optimized

and fixed early on during evolution, while being maybe also involved in other functional

pathways than drought response (Harrison et al. 2012). TDI profiles support this claim:

conserved genes do contribute more to the TDI profiles in cell-cycle networks and show

adaptive changes in expression for drought response (higher TDI difference between control

and drought transcriptomes in cell-cycle network, Figure 5B). In contrast, drought-responsive

genes in metabolism network appear more variable in their expression in response to

drought stress, because this strategy may be linked to an initial response to severe water

scarcity (Dubois and Inzé 2020).

Population genetics analysis of drought-responsive networks

We also study the selective forces acting on the identified drought-responsive gene networks

at the population level. Using full genome sequences of six *S. chilense* populations

(C_LA1963, C_LA3111, C_LA2931, SC_LA2932, SC_LA4107, and SH_LA4330; five plants

each) recently reported in Wei et al. (2023), aligned to the reference genome of *S. chilense*,

we identify 45,208,263 high-quality single-nucleotide variants (SNPs), in which 111,606 SNPs

are found in genes of the cell-cycle GCN and 167,334 SNPs in genes of the metabolic GCN.

We first compare population structure between the whole-genome data and drought-

responsive genes (Figure S8). The results corroborate the genetic structure revealed in Wei

et al. (2023) (Figure S8A and S8C). However, the structure exhibited by drought genes

shows stronger differentiation among populations than the WGS data (especially for

clustering of populations of the central region and SH_LA4330). Moreover, the strong

differences from WGS data between the two south coastal populations (SC_LA2932 and

SC_LA4107) is attenuated when analyzing SNPs from the drought-responsive genes (Figure

S8B and S8D).

We find that the mean nucleotide diversity (π) per gene does not differ between the two

GCNs (Figure S9A; Table S5; Kolmogorov-Smirnov test, P = 0.15). In addition, the π values

of the promoter regions (here 2kb upstream of the transcription initiation site) are significantly

higher than those of the gene (coding) regions (Figure S9A; Table S5; Kolmogorov-Smirnov

test, P = 0.03). This result suggests that the selective constraints in promoter regions may be

more relaxed, which could in part explain why certain transcription factors are able to bind to

multiple genes in the GCNs. (Table S2). TFs are indeed conserved at the coding sequence

level, especially at the functional domains, but higher amount of polymorphism of TF binding

sites in the promoter can be indicative of complex and diverse regulation, for example in

response to stressful conditions (Spivakov 2014; Sato et al. 2016). Albeit, there is no

difference in the nucleotide diversity at the promoter regions between the two GCNs (Figure

S9A; Table S5). Furthermore, the genes for the metabolic GCN show lower Tajima's D values

than those of the cell-cycle GCN (Figure S9B; Table S5; Kolmogorov-Smirnov test, P = 0.04),

suggesting recent positive selection pressure in the metabolic GCN. We find weak correlation

between Tajima's D and Ka/Ks ratio for the cell-cycle GCN and absence of correlation for the

metabolic GCN (Figure S10A and S10B). As a negative correlation between Tajima's D and

Ka/Ks ratio is indicative of recent positive selection, our results suggest the possibility of

recent positive selection acting on multiple genes within the metabolic GCN (Figure S9B;

Table S5).

We further find significant, but opposite, correlations between π or Tajima's D and the

contributions of the different DS for the two GCNs (Figure S10A and S10B). In the cell-cycle

GCN, the contributions of different DS have significant positive correlation with π and

Tajima's D (Figure S10A and S10C). This indicates that DS of high contribution to TDI

profiles show high nucleotide diversity (and positive Tajima's D), meaning that older genes

are under stronger purifying selection than younger genes in this network because the

sequence divergence of cell-cycle genes occurred at old time periods. In contrast, a negative

correlation is observed between the contribution of each DS and π or Tajima's D in the

metabolic network (Figure S10B and S10D). Hence, DS with high contribution show low

nucleotide diversity and low Tajima's D, especially DS10. Therefore, it appears likely that the

metabolic genes, likely recently evolved, may be under positive selection underpinning the

recent evolution of the drought response transcriptome.

Drought-responsive genes under positive selection promote adaptive evolution in

response to drought stress

Genetic drift or changes in selective pressure is one of the main factors that contribute to

gene-expression variation (Koenig et al. 2013). To investigate drought-responsive genes that

have potentially undergone a shift in selection regime, we search for overlap between genes

of two drought-response GCNs studied here and our previously identified 799 candidate

genes under positive selection in six populations of *S. chilense* (Wei et al. 2023). We find 74

and 126 drought-responsive genes in the cell-cycle and metabolic networks, respectively in

the list of candidate genes under positive selection (Figure 6A; Table S6). These genes

exhibit the typical characteristics of positively selected genes with low π and Tajima's D

(Table S5). This indicates that drought stress is likely an important driver of adaptation and

these drought-response genes may play key roles for colonization of new arid habitats.

Similar numbers of drought-responsive genes likely under positive selection are observed

across different populations of *S. chilense* encompassing different parts of the range, except

for SH_LA4330 (Wei et al. 2023). The number of candidate genes belonging to the metabolic

or cell-cycle GCNs is similar in the three central populations (C_LA1963, C_LA3111 and

C_LA2931) (Figure 6A; Table S6). The most recent diverged highland population

(SH_LA4330) contains the largest number of positively selected drought-responsive genes

(Figure 6A; Table S6) with a similar proportion of genes from both networks. Noticeably, in

the two south-coast populations (SC_LA2932 and SC_LA4107) a large majority of genes

under positive selection belong to the metabolic GCN (showing absence of cell-cycle genes

in population SC_LA2932, Figure 6A; Table S6).

Previous studies have demonstrated that adaptive genes are pleiotropic and proposed

functional connectivity between networks related to different quantitative traits (Wagner et al.

2007; Erwin and Davidson 2009; Hämälä et al. 2020). To address the role that (putatively)

positively selected genes may play within the drought-responsive networks, we compare the

connectivity of these genes in the two networks (Figure 6B; Table S7). In the metabolic

network, the connectivity of positively selected genes (0.55 ± 0.10) is significantly higher than

other drought-responsive genes (0.44 ± 0.12) (Figure S11A; Kolmogorov-Smirnov test, P =

0.017), but we do not observe such significant difference for the cell-cycle network (Figure

S11A; Kolmogorov-Smirnov test, P = 0.43). Furthermore, the connectivity of positively

selected genes of the metabolic network is much higher than those from the cell-cycle

network in six populations (Figure 6B; Table S7; Kolmogorov-Smirnov test, P = 0.007). These

results suggest that highly connected (likely more pleiotropic) genes in the metabolic GCN

may have facilitated the recent colonization of new habitats (Hämälä et al. 2020) during the

divergence process of *S. chilense*. In contrast, the connectivity of positively selected genes in

the cell-cycle network is significantly lower (Figure S11A). Therefore, we suggest that the two

networks underwent different evolutionary selective pressures during the range expansion of

S. chilense.

Finally, we compare the age of the selective sweep at the candidate genes of the two

GCNs based on the results in Wei et al. (2022). We find that sweep ages at the cell-cycle

genes are slightly younger than at those of the metabolic network, especially in the three

highland populations (C_LA2931, C_LA3111 and SH_LA4330; Figure S11B and S11C; Table

S7). This supports that drought adaptation is an important mechanism underlying the recent

(re)colonization of highland habitats (Raduski and Igić 2021; Wei et al. 2023). Interestingly,

we find significantly positive correlation between the age of the sweep and gene connectivity

for both GCNs and across all six populations (Figure 6C). Figure 6D and 6E provide the

visualizations of two networks and exhibit the relationship between sweep age and

connectivity (depicting weighted connection strength greater than 0.65 between any two

genes). In other words, it appears that selective sweeps occur first at more connected genes

and, subsequently at less connected genes, during the history of colonization/adaptation of

new arid habitats. To our knowledge, this is the first report of a correlation between the age of

a selective sweep and the connectivity of genes in a network. To obtain more evidence to

support this inference, we also calculate the tMRCA (time to most recent common ancestor)

to estimate the age of drought-responsive genes based on allele frequency of SNPs. The

positive correlation between tMRCA of drought-responsive genes under the positive selection

and connectivity is also supported (Pearson's cor=0.69, P = 2.47e-5), consistent with the

correlation with sweep age. Moreover, the low correlation (Pearson's cor=0.31, P = 0.14) is

observed between tMRCA of other (outside of sweep regions) drought-responsive genes and

connectivity. This supports the hypothesis of polygenic adaptation in GCNs where the

positive selection acts first on core genes (with high connectivity and more pleiotropic) of

networks, and subsequently on the peripheral genes (less connectivity and less pleiotropic).

These positively selected genes ultimately regulate the expression of other genes in the

network.

Discussion

In this study, we identify two drought-responsive GCNs by analyzing gene expression profiles

of plants growing under control and drought conditions. Two GCNs involved in cell-cycle and

metabolic biological processes are detected and their structural relevance are supported by

TF/TFBS predictions. These networks represent two different strategies for drought response

(Farooq et al. 2009; Danilevskaya et al. 2019). We then demonstrate that the cell-cycle

network is evolutionary older and more conserved than the metabolic network. Despite the

ancient history of these two GCNs, we further show that both GCNs also contribute to

different extents to contemporary processes of adaptation to drought conditions when *S.*

chilense colonizes new arid habitats around the Atacama desert. The joint analyses of

genomic and transcriptomic data indicates that 1) at the transcriptome level, metabolic GCN

present a higher evolvability especially with younger selection events linked to response to

new environments, 2) cell-cycle GCN is less evolvable, and 3) both networks still present

signals of evolution under positive selection in core elements of the GCN, while peripheral

genes of the network can be involved in adaptation at later stages of the colonization

processes.

Drought tolerance is mediated by regulation in cell proliferation and metabolism

When roughly defining the organ development into cell proliferation and differentiation,

water deficit appears to be a limiting factor for both processes (Alves and Setter 2004;

Verelst et al. 2013). Drought stress reduces the activity of the cell cycle and thus slows down

the growth and development of plants. The down-regulated genes we find in the cell-cycle

network also indicate that genes related to cell cycle are suppressed by drought stress

possibly to restrict the cell division in *S.chilense.* Reduction of cell number due to mild

drought stress is also found in *A. thaliana* (Skirycz and Inzé 2010). This means that the cell-

cycle response to drought may be very general and indirect. However, our speculations are

mainly based on the aboveground tissues of *S. chilense*. Conversely, the changes of

fundamental metabolic activity may be a faster and a flexible drought-responsive strategy

presumably related to acclimation (Harb et al. 2010). Plant water shortage is first reflected in

changes in metabolic processes, such as accelerating the catabolism of macromolecules in

order to regulate the penetration of tissues, to maintain physiological water balance, or

slowing down metabolism to reduce energy and water consumption (Reddy et al. 2004;

Gupta et al. 2020). In addition, the signaling pathways related to the metabolic gene network

are also demonstrated to be a response to drought stress, for example, the abscisic acid

(ABA) signaling pathway regulates the response to dehydration and optimizes water

utilization (Harb et al. 2010; Wilkinson and Davies 2010). Although these two GCNs

correspond to two different strategies of drought response, they are not isolated, but interact

with one another in a time-dependent manner. Water deprivation and heat first change the

metabolic processes leading to stomata closure, which leads then to cell cycle network to be

affected under long-term lack of water. In return, the increased or decreased cell cycle gene

expression affects the further physiology and metabolism of the plant (Gupta et al. 2020).

Indeed, drought-responsive strategies regulating the cell cycle appear to be activated later

than metabolism processes, as glucose metabolism rapidly follows drought stress, whereas

the accumulation of amino acids which is a crucial part of the cell cycle response starts at a

later time in response to drought (Fàbregas and Fernie 2019).

Rewiring of ancient GCNs drives recent adaptation to dry environments

The phylostratigraphic analysis supports that the majority of drought-responsive genes in *S.*

chilense evolved during the early to middle stages of plant evolutionary history, which is in

agreement with the time of origin of multiple abiotic response genes in *Arabidopsis thaliana*

(Mustafin et al. 2019). This reinforces that the emergence of drought-responsive genes

coincides with the time periods of divergence among major plant groups (land- and flowering

plants), which are marked by frequent whole genome duplication events that trigger gene

family expansions, gene neo- and sub-functionalization, and genome reorganization

processes (Wang et al. 2012; Clark and Donoghue 2018). These genomic processes likely

contributed to the enrichment of drought-responsive GCNs. For instance, fundamental

morphological traits involved in drought responses, such as stomata, are present in the

ancestral land plants. However, stomatal genes existed prior to the divergence of land plants

and underwent multiple duplications during the course of evolution. Additionally, their

response to environmental cues, such as humidity, light, CO_2, and ABA, is widely distributed

and may be ancestral to land plants (Clark et al. 2022). Therefore, we propose that our two

drought-responsive networks were primarily established during or shortly after the

divergence of land plants and have subsequently undergone expansion. This highlights the

crucial role of ancestral genomic processes in shaping the genetic mechanisms that underlie

plant adaptation to drought.

Previous studies show that TAI and TDI profiles across embryogenesis, seed

germination and transition to flowering in *A. thaliana* exhibit a 'hourglass pattern' (older and

conserved transcriptomes are preferentially active at the mid-development stages; Quint et al.

2012; Drost et al. 2016). However, our TAI/TDI profiles for the two developmental stages

remain stable under the same conditions (Figures 4C and 5B). The similar TAI/TDI between

developmental stages (Figure 4C and 5B) that we obtained is certainly because our analyses

focused on two modules (co-expressed genes) highly correlated to the differential expression

between drought and control conditions (Figure 2D; Table S1). Therefore, developmental

stage-specific response genes are underrepresented in the two analyzed networks. However,

increased TAI/TDI values under drought conditions suggest that stress response

transcriptomes are composed of relatively more recently diverged genes, and therefore are

more evolvable. We suggest that this inference needs to be verified in other stress

responsive transcriptomes (salt, heat, cold, etc.). We then speculate, that although abiotic

stress response regulatory networks are mostly composed of highly ancient and conserved

elements across species (Chen and Zhu 2004), networks retain the ability to change

expression patterns to respond rapidly to environmental changes or to explore new

ecological niches. Moreover, given the pleiotropic nature of the abiotic stress-response traits,

we can expect shared patterns of evolution (at the constitutive and expression components)

of the networks for different stress conditions (and possible trade-offs between traits and

GCNs).

Extensive network rewiring in relatively recent and short time-frames have been found

in maize and tomato in response to domestication (Swanson-Wagner et al. 2012; Koenig et

al. 2013). It is therefore not surprising to find signs of adaptive variation in core elements of

rather conserved regulatory networks related to the colonization processes of new (here arid)

habitats. The genetic (and morphological) divergence of the *S. chilense* marginal southern

populations, southern coastal and highland, is recent but strong (Raduski and Igić 2021). It is

congruent with theoretical results showing that gene networks with higher mutation sensitivity

can facilitate local adaptation, present increasing variance in gene expression and underlie

accelerated range expansion processes across abiotic environmental gradients (Deshpande

and Fronhofer 2022). Complementarily, our empirical approach shows the existence of two

regulatory networks with different evolutionary trends, one being more conserved than the

other and exhibiting different gene expression responses. One GCN would exhibit a faster

and more variable response (metabolic), while the other a later (delayed) but more

constitutive response (cell-cycle) to drought. Despite the differences in gene age and

variation between the networks, our results show that both GCNs have undergone sufficient

changes leading to their rewiring during the divergent process of colonization of *S. chilense*

around the Atacama. Nevertheless, genes in the metabolic network show more recent

evolution, with new genes members appearing in *S. chilense*, concomitantly with more

variable expression in the drought transcriptome.

These drought-responsive genes to *S. chilense* likely facilitated the adaptation of this

species to unique arid (up to hyper-arid) habitats, especially when colonizing the southern

part of the range. Indeed, population structure based on SNPs indicates that drought-

responsive genes reflected adaptation/colonization to arid habitats in *S. chilense* (Figure S8).

Importantly, we found about 200 drought-responsive genes previously identified as candidate

genes under positive selection (*i.e.* located within sweep regions in Wei et al. 2023). This

confirms that drought stress is an important driver of ecological divergence in *S. chilense.* We

finally provide some indirect evidence that changes at central genes (with higher connectivity)

can be responsible for the short-term response to selection (Jovelin and Phillips 2009; Luisi

et al. 2015) and promote rewiring of the gene network (Koubkova-Yu et al. 2018). Thus,

highly connected genes may be targets of positive selection during the first phase of the

environmental change or colonization to contrasting environments, and may be keys for

'piggybacking', defined as the change in gene expression of a focal gene driving phenotypic

change. Altogether, our results on the age-dependent adaptive role of genes with different

network connectivity (and possible pleiotropic effects) provide another line of evidence

supporting the view that molecular evolution follows an adaptive walk and are

complementary to the recent study by (Moutinho et al. 2022).

Limitations and further work

A limitation of our gene expression study is that our transcriptomic analyses are based on

individuals from a single location (near the putative region of origin of the species; Wei et al.

2022), while variability in gene expression and phenotypic response has been observed

between different populations (Mboup et al. 2012; Fischer et al. 2013; Nosenko et al. 2016).

Further expression studies including plants from multiple locations would be useful to verify

that the identified GCNs are also present and expressed in other populations and study the

possible variation in the most southern populations. More evidence based on multiple

populations is needed to confirm the 'piggybacking' phenomenon of gene expression in *S.*

chilense. Additional support on the variability of transcriptome evolution across populations

as well as long read sequencing of more genomes will be beneficial in assessing the role of

gene duplication and gene deletion yielding the evolution of the gene networks. Such studies

would also allow the analysis of evolution of adaptive gene networks and polygenic selection

occurring for complex traits such as drought tolerance. Finally, more detailed studies with a

larger sample size from the field will help to discover other gene networks and their

interactions related to abiotic stress and the evolution of the species. A detailed discussion of

the potential biases associated with the use of multiplied accessions at TGRC (Tomato

Genetics Resource Center, UC Davis, USA) compared to samples from natural populations

is found in Wei et al. (2022). Sampling and experimental work in the field would improve the

resolution of transcriptome and genomic studies, in order to assess phenotypic differences

between organs and stages of development and thus extend the knowledge to other relevant

characteristics such as secondary metabolism, which is known to have relevant influence on

biotic and abiotic interactions (Mes et al. 2008; Bolger et al. 2014; Tapia et al. 2022).

Material and methods

Plant material and drought stress experiment

Seeds of *S. chilense* accession LA1963 were acquired from Tomato Genetics Resource Center (TGRC), University of California at Davis. Seeds were soaked in 50% household bleach (2.7% sodium hypochlorite) for 30 minutes and rinsed thoroughly with water according to instructions provided by TGRC. The rinsed seeds were sown into pots containing sterilized soil with perlite and sand (1:2) and grown under controlled conditions (22C day/20C night, 16h light/8h dark photoperiod). On the 24th day after sowing, all plants were randomly distributed into two groups and watered with a sufficient volume to reach the bottom of containers (30-40 ml). The first group of plants were maintained under normal watering condition, watered with a sufficient volume of water (50-55 ml) on 4, 7 and 11 days after start of the experiment (day 24). A moderate water stress regime was imposed to second group of plants by stopping irrigation for 7 days followed by re-watering with 25 ml of water. On day 12, newly expanded leaf (1-1.5 cm length) and shoot apices with immediately surrounding leaf primordia (shoot apices and P1-P5 leaf primordia) from each group were dissected carefully using razor blades and immediately grounded into fine powder in liquid nitrogen for RNA extraction. Four biological replicates were used for all RNA-Seq experiments from each tissue type. Each replicate of leaf and shoot apex samples included the pooled tissues from five and six plants, respectively.

RNA extraction and cDNA library construction

Libraries were constructed and named as follows: leaves under control (optimal watering) condition (CL-A to D), shoot apices under control condition (CSA-E to H), leaves under drought condition (DL-I to L), and shoot apices under drought condition (DSA-M to P). Tissues were lysed using zircon beads in Lysate Binding Buffer containing Sodium Dodecyl

Sulfate. mRNA was isolated from 200 µl of lysate per sample with streptavidin coated

magnetic beads for indexed non-strand specific RNA-Seq library preparation according to the

method described by (Kumar et al. 2012). 1 µl of 12.5 µM of 5-prime biotinylated polyT

oligonucleotide and streptavidin-coated magnetic beads were used to capture mRNA and

isolate captured mRNAs from the lysate, respectively. Equal amount of mRNA of each

experimental group were used to construct 16 libraries. For library construction the rapid

version of Kumar et al. (2012) RNA-sequencing method (Townsley et al. 2015) was used.

Each sample was barcoded using standard Illumina adaptors 1-16 to allow up to 16 samples

to be pooled in one lane of sequencing on Illumina HiSeq4000. The libraries were eluted

from the pellet with 10 µl 10 mM Tris pH 8.0 and pooled as described by Kumar et al. (2012).

Quantification and quality assessment of resulting libraries were performed on Fragment

Analyzer (FGL_DNF-474-2- HS NGS Fragment 1-6000bp.mthds) and sequenced using the

Illumina HiSeq 4000 platform to generate 100 bp single-end reads at the Vincent J. Coates

Genomic Sequencing Facility at UC Berkeley.

Transcriptome and genome data processing and mapping

For transcriptome data, the adapters were removed from raw reads by two consecutive

rounds using BBDuk in BBTools v38.90 (Bushnell 2014). Two sets of parameters were used

in two rounds respectively: first round 'ktrim=r k=21 mink=11 hdist=2 tpe tbo minlength=21

trimpolya=4'; second round 'ktrim=r k=19 mink=9 hdist=1 tpe tbo minlength=21 trimpolya=4'.

Then Low-quality reads were also removed with BBDuk using parameters 'k=31 hdist=1

qtrim=lr trimq=10 maq=12 minlength=21 maxns=5 ziplevel=5'. The clean reads of each

sample were mapped to the *S. chilense* reference genome (Silva-Arias et al. submited) using

BBMap in BBTools. The SAM files were then converted and sorted to BAM files using

Samtools v1.11 (Wysoker et al. 2009). The number of reads were mapped to each gene

were counted via featureCounts v2.0.1 in each sample (Liao et al. 2014). To eliminate the

differences between samples, the gene expression level was normalized using the TPM

(Transcripts Per Kilobase Million) method (Wagner et al. 2012).

The relationships among transcriptome samples were evaluated using the TPM values.

The correlation coefficient between two samples was calculated to evaluate repeatability

between samples using Pearson's test. Principal component analysis (PCA) was performed

using the *prcomp()* function in R (R Core Team 2020) based on TPM values .

Identification of differentially expressed genes and gene co-expression analysis

Differential expression analysis of groups among the different conditions and tissues was

performed using the DESeq2 R package (Love et al. 2014). The raw read counts were

inputted to detect Differential Expressed Genes (DEGs). The *P*-value ≤ 0.001, the absolute

value of log2FoldChange ≥ 1 and a false discovery rate (FDR) adjusted $P \le 0.001$ were

classified as differentially expressed genes.

To identify the gene co-expression networks, weighted gene correlation network

analysis (WGCNA) was constructed using TPM values to identify specific modules of co-

expressed genes associated with drought stress (Langfelder and Horvath 2008). We first

checked for genes and samples with too many missing values using *goodSamplesGenes()*

function in WGCNA R package. We then removed the offending genes (the last statement

returns 'FALSE'). To construct an approximate scale-free network, a soft thresholding power

of five was used to calculate adjacency matrix for a signed co-expression network.

Topological overlap matrix (TOM) and dynamic-cut tree algorithm were used to extract

network modules. We used a minimum module size of 30 genes for the initial network

construction and merged similar modules exhibiting > 75% similarity. To discover modules of

significantly drought-related, module eigengenes were used to calculate correlation with

samples with different conditions. The visualization of networks were created using

Cytoscape v3.8.2 (Su et al. 2014).

Identification of transcript factor families and transcript factor binding sites

The protein sequences were obtained from the reference genome and annotation 'gff' file

with GffRead (Pertea and Pertea 2020), and were used to identify TF families using online

tool PlantTFDB v5.0 (Guo et al. 2007). Furthermore, the upstream 2000 bp sequences of the

transcription start sites (TSS) were extracted as the gene promoter from the reference

genome to detect TFBS. The TFBS dataset of relative species *S. pennellii* was also

downloaded   from   Plant   Transcriptional   Regulatory   Map   (PlantRegMap,

http://plantregmap.gao-lab.org/) as background of TFBS identification (Tian et al. 2020).

Then, the TFBS of *S. chilense* was identified using FIMO program in motif-based sequence

analysis tools MEME Suit v5.3.2 (Bailey et al. 2015). The TFBS was extracted with $p < 1e\text{-}5$

and $q < 0.01$.

Gene ontology (GO) analysis

We first constructed the dataset of assigned GO terms for all genes used protein sequence

by PANTHER v16.0 (Mi et al. 2021). Then, the GO enrichment analysis of drought-

responsive genes was performed using clusterProfilter v3.14.2 (Yu et al. 2012). Benjamini–

Hochberg method was used to calibrate *P* value, and the significant GO terms were selected

with *P*-value below to 0.05.

Construction of phylostratigraphic map

We performed phylostratigraphic analysis based on the following steps. First, the phylostrata

(PS) was defined according to the full linkage of *S. chilense* from NCBI taxonomy database.

The similar PS was merged and finally 18 PS were generated (Figure 4A). Second, the

protein sequences were blast to a database of non-redundant (nr) proteins downloaded from

NCBI (https://ftp.ncbi.nlm.nih.gov/blast/db/) with a minimum length of 30 amino acids and an

E-value below 10^{-6} using blastp v2.9.0 (Camacho et al. 2009). Third, each gene was

assigned to its PS by the following criterion: if no blast hit or only one hit of *S. chilense* with

an E-value below 10^{-6} was identified, we assigned the gene to the youngest PS18. When

multiple blast hits were identified, we computed lowest common ancestor (LCA) for multiple

hits using TaxonKit v0.8.0 (Shen and Ren 2021) and then assigned LCA to specific PS.

Construction of divergence map

We performed divergence stratigraphy analysis to construct sequence divergence map of *S. chilense* using function *divergence_stratigraphy()* of R package 'orthologr' (Drost et al. 2015) following four steps: 1) the coding sequences for each gene of *S. chilense* and *S. pennellii* (NCBI assembly SPENNV200) were extracted from their reference and annotation files. 2) We identified orthologous gene pairs of both species by choosing the best blast hit for each gene using blastp. We only considered a gene pair orthologous when the best hit has an E-value below 10^{-6}, the gene pair is considered orthologous; otherwise, it is discarded. 3) Codon alignments of the orthologous gene pairs were performed using PAL2NAL (Suyama et al. 2006). Then, Ka/Ks values of the codon alignments were calculated using Comeron's method (Comeron 1995). And 4) all genes were sorted according to Ka/Ks values into discrete deciles, which are called divergence stratum (DS).

Estimation of transcriptome age index and transcriptome divergence index

The TAI is computed based on phylostratigraphy and expression profile, which assign each gene to different phylogenetic ages by identification of homologous sequences in other species (Domazet-Lošo et al. 2007). The evolutionary age of each gene was quantified combining its PS and expression level to obtain weighted evolutionary age. Finally, weighted ages of all genes are averaged to yield TAI, which is defined as the mean evolutionary age of a transcriptome (Domazet-Lošo and Tautz 2010). A lower value of TAI describes an older mean evolutionary age, whereas a higher value of TAI denotes a younger mean evolutionary age and implies that evolutionary younger genes are preferentially expressed in the corresponding sample or condition (Domazet-Lošo and Tautz 2010; Piasecka et al. 2013). The TDI represents the mean sequence divergence of a transcriptome quantified by divergence strata (DS) and gene expression profile (Quint et al. 2012). The genes are assigned to different DS and then weighted by their expression level to yield the TDI. A lower value of TDI describes a more conserved transcriptome (in terms of sequence dissimilarity),

whereas a higher value of TDI denotes a more variable transcriptome. Here, we calculate TAI

and TDI profiles in different samples using *PlotSignature()* function of the myTAI R package.

Population genetics analysis and detection of positive selection on drought-

responsive genes

Whole-genome sequence data from six populations *S. chilense* (five individuals each)

previously analyzed in (Wei et al. 2022; BioProject PRJEB47577) were used to calculate

population genetics statistics for coding and promoter region sequences for all genes

identified in the GCNs. Single nucleotide variants (SNPs) based on the short-read alignment

to the new reference genome for *S. chilense* (Silva-Arias et al. submitted) using the same

methods in Wei et al. (2022). Population genetics statistics namely, nucleotide diversity (π)

and Tajima's D were calculated with ANGSD v0.937 (Korneliussen et al. 2014) over gene and

promoter regions. These statistics first were calculated at per site in gene and promoter

regions, and then we used a R script (https://gitlab.lrz.de/population_genetics/s.chilense-

drought-transcriptome) to obtain statistics in each gene and promoter regions. PCA on SNP

data from 30 whole genomes was also performed using GCTA (v1.91.4; Yang et al. 2011).

The genetic structure inference was performed using ADMIXTURE v1.3.0 (Alexander et al.

2009).

Drought-responsive genes under positive selection were extracted by blast (e-value <

1e-6) between drought-responsive genes identified in this study and the genes located inside

sweep regions in our previous study using *S. pennellii* as the reference genome. We also

use the sweep ages obtained in Wei et al. (2022).

Estimation of allele age

We implemented in GEVA (Genealogical Estimation of Variant Age; Albers and McVean 2020)

to dating genomic variants in the drought-responsive genes. We generated input for GEVA

based on the recombination rate 3.24×10^{-9} per site per generation (based on the overall

recombination density in *S. lycopersicum* [1.41 cM/Mb] Anderson and Stack 2002; Nieri et al.

2017; and within the possible range of rates used in Wei et al. 2022). We used population

size (N_e) 20,000 and mutation rate 5.1 x 10^{-9} (Roselius et al. 2005; Wei et al. 2023), and then

relied on the recombination clock to estimate the age of alleles (tMRCA).

Supplementary material

Supplementary data are available at Molecular Biology and Evolution online.

Acknowledgements

KW was funded by the Chinese Scholarship Council. SS was funded by the Fulbright Visiting

Scholar Program of the U.S. Department of State and by German Academic Exchange

Service (DAAD) within the framework of Research Stays for University Academics and

Scientists. GAS-A was funded by the Technical University of Munich. AT acknowledges

funding from DFG (Deutscheforschungsgemeinschaft) Grant Number: 317616126 (TE809/7-

1). NS acknowledges funding from U.S. National Science Foundation (NSF/ IOS-1238243).

This work was supported by grants from JSPS KAKENHI (JP19K23742, JP20K06682,

JP20KK0340 to H.N.), and NSF PGRP IOS-1856749 and IOS-211980 to NS. The laboratory

work of H.N. was supported by a Grant-in-Aid for Scientific Research on Innovative Areas

(JP19H05672). We thank the Tomato Genetics Resource Center (TGRC) of the University of

California, Davis for generously providing us with the seeds of the accession included in this

study.

Data Availability

The raw sequencing RNA data is available in PRJDB15063 (*S. chilense*), PRJEB5809 (*S.*

pennellii) and PRJNA812356 (*S. lycopersicum*). The raw pair-end whole-genome sequencing

data can be accessed at the European Nucleotide Archive (ENA) project accession

PRJEB47577. All codes used in this study and other previously published genomic data are

available at the sources referenced. The code for implementing the analyses used in this

paper can be found on our GitLab repository:

https://gitlab.lrz.de/population_genetics/s.chilense-drought-transcriptome

References

Albers PK, McVean G. 2020. Dating genomic variants and shared ancestry in population-scale sequencing data. *PLOS Biol.* 18:e3000586.

Alexander DH, Novembre J, Lange K. 2009. Fast model-based estimation of ancestry in unrelated individuals. *Genome Res.* 19:1655–1664.

Alves AA, Setter TL. 2004. Response of cassava leaf area expansion to water deficit: cell proliferation, cell expansion and delayed development. *Ann. Bot.* 94:605–613.

Anderson L, Stack S. 2002. Meiotic recombination in plants. *Curr. Genomics* 3:507–525.

Bailey TL, Johnson J, Grant CE, Noble WS. 2015. The MEME suite. *Nucleic Acids Res.* 43:W39–W49.

Barrera-Ayala D, Tapia G, Ferrio JP. 2023. Leaf carbon and water isotopes correlate with leaf hydraulic traits in three *Solanum* species (*S. peruvianum, S. lycopersicum and S. chilense*). *Agriculture* 13:525.

Basu S, Ramegowda V, Kumar A, Pereira A. 2016. Plant adaptation to drought stress. *F1000Research* 5.

Blanchard-Gros R, Bigot S, Martinez J-P, Lutts S, Guerriero G, Quinet M. 2021. Comparison of drought and heat resistance strategies among six populations of *Solanum chilense* and two cultivars of *Solanum lycopersicum*. *Plants* 10:1720.

Bolger A, Scossa F, Bolger ME, Lanz C, Maumus F, Tohge T, Quesneville H, Alseekh S, Sørensen I, Lichtenstein G, et al. 2014. The genome of the stress-tolerant wild tomato species *Solanum pennellii. Nat. Genet.* 46:1034–1038.

Böndel KB, Lainer H, Nosenko T, Mboup M, Tellier A, Stephan W. 2015. North–south colonization associated with local adaptation of the wild tomato species *Solanum chilense. Mol. Biol. Evol.* 32:2932–2943.

Böndel KB, Nosenko T, Stephan W. 2018. Signatures of natural selection in abiotic stress-responsive genes of *Solanum chilense. R. Soc. Open Sci.* 5:171198.

Bowles AMC, Paps J, Bechtold U. 2021. Evolutionary origins of drought tolerance in Spermatophytes. *Front. Plant Sci.* 12.

Bushnell B. 2014. BBTools. A suite of bioinformatics tools used for DNA and RNA sequence data analysis. *DOE Jt. Genome Inst.* [Internet]. Available from: https://sourceforge.net/projects/bbmap/

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10:1–9.

Chen WJ, Zhu T. 2004. Networks of transcription factors with roles in environmental stress response. *Trends Plant Sci.* 9:591–596.

Ciais P, Reichstein M, Viovy N, Granier A, Ogée J, Allard V, Aubinet M, Buchmann N,
Bernhofer C, Carrara A, et al. 2005. Europe-wide reduction in primary productivity
caused by the heat and drought in 2003. *Nature* 437:529–533.

Clark JW, Donoghue PCJ. 2018. Whole-genome duplication and plant
macroevolution. *Trends Plant Sci.* 23:933–945.

Clark JW, Harris BJ, Hetherington AJ, Hurtado-Castaño N, Brench RA, Casson S,
Williams TA, Gray JE, Hetherington AM. 2022. The origin and evolution of stomata.
Curr. Biol. 32:R539–R553.

Crow M, Suresh H, Lee J, Gillis J. 2022. Coexpression reveals conserved gene
programs that co-vary with cell type across kingdoms. *Nucleic Acids Res.* 50:4302–
4314.

Danilevskaya ON, Yu G, Meng X, Xu J, Stephenson E, Estrada S, Chilakamarri S,
Zastrow-Hayes G, Thatcher S. 2019. Developmental and transcriptional responses of
maize to drought stress under field conditions. *Plant Direct* 3:e00129.

Deshpande JN, Fronhofer EA. 2022. Genetic architecture of dispersal and local
adaptation drives accelerating range expansions. *Proc. Natl. Acad. Sci.*
119:e2121858119.

Domazet-Lošo T, Brajković J, Tautz D. 2007. A phylostratigraphy approach to uncover
the genomic history of major adaptations in metazoan lineages. *Trends Genet.*
23:533–539.

Domazet-Lošo T, Tautz D. 2010. A phylogenetically based transcriptome age index
mirrors ontogenetic divergence patterns. *Nature* 468:815–818.

Drost H-G, Bellstädt J, Ó'Maoiléidigh DS, Silva AT, Gabel A, Weinholdt C, Ryan PT,
Dekkers BJW, Bentsink L, Hilhorst HWM, et al. 2016. Post-embryonic hourglass
patterns mark ontogenetic transitions in plant development. *Mol. Biol. Evol.* 33:1158–
1163.

Drost H-G, Gabel A, Grosse I, Quint M. 2015. Evidence for active maintenance of
phylotranscriptomic hourglass patterns in animal and plant embryogenesis. *Mol. Biol.*
Evol. 32:1221–1231.

Dubois M, Inzé D. 2020. Plant growth under suboptimal water conditions: early
responses and methods to study them.Mittler R, editor. *J. Exp. Bot.* 71:1706–1722.

Erwin DH. 2020. Chapter Thirteen - Evolutionary dynamics of gene regulation. In:
Peter IS, editor. Current Topics in Developmental Biology. Vol. 139. Gene Regulatory
Networks. Academic Press. p. 407–431. Available from:
https://www.sciencedirect.com/science/article/pii/S0070215320300326

Erwin DH, Davidson EH. 2009. The evolution of hierarchical gene regulatory
networks. *Nat. Rev. Genet.* 10:141–148.

Fàbregas N, Fernie AR. 2019. The metabolic response to drought. *J. Exp. Bot.*
70:1077–1085.

Farooq M, Wahid A, Kobayashi N, Fujita D, Basra S. 2009. Plant drought stress:
effects, mechanisms and management. *Sustain. Agric.*:153–188.

Ficklin SP, Feltus FA. 2011. Gene coexpression network alignment and conservation
of gene modules between two grass species: maize and rice. *Plant Physiol.*
156:1244–1256.

Fischer I, Camus-Kulandaivelu L, Allal F, Stephan W. 2011. Adaptation to drought in
two wild tomato species: the evolution of the *Asr* gene family. *New Phytol.* 190:1032–
1044.

Fischer I, Steige KA, Stephan W, Mboup M. 2013. Sequence evolution and
expression regulation of stress-responsive genes in natural populations of wild
tomato. *PLoS One* 8:e78182–e78182.

Flowers JM, Sezgin E, Kumagai S, Duvernell DD, Matzkin LM, Schmidt PS, Eanes
WF. 2007. Adaptive evolution of metabolic pathways in *Drosophila. Mol. Biol. Evol.*
24:1347–1354.

Gehan MA, Greenham K, Mockler TC, McClung CR. 2015. Transcriptional
networks—crops, clocks, and abiotic stress. *Curr. Opin. Plant Biol.* 24:39–46.

Gerstein MB, Rozowsky J, Yan K-K, Wang D, Cheng C, Brown JB, Davis CA, Hillier L,
Sisu C, Li JJ. 2014. Comparative analysis of the transcriptome across distant species.
Nature 512:445–448.

Guo A-Y, Chen X, Gao G, Zhang H, Zhu Q-H, Liu X-C, Zhong Y-F, Gu X, He K, Luo J.
2007. PlantTFDB: a comprehensive plant transcription factor database. *Nucleic Acids*
Res. 36:D966–D969.

Gupta A, Rico-Medina A, Caño-Delgado AI. 2020. The physiology of plant responses
to drought. *Science* 368:266–269.

Haak DC, Kostyun JL, Moyle LC. 2014. Merging ecology and genomics to dissect
diversity in wild tomatoes and their relatives. In: Landry CR, Aubin-Horth N, editors.
Ecological Genomics. Vol. 781. Dordrecht: Springer Netherlands. p. 273–298.
Available from: http://link.springer.com/10.1007/978-94-007-7347-9_14

Hämälä T, Gorton AJ, Moeller DA, Tiffin P. 2020. Pleiotropy facilitates local adaptation
to distant optima in common ragweed (*Ambrosia artemisiifolia*). *PLoS Genet.*
16:e1008707.

Harb A, Krishnan A, Ambavaram MMR, Pereira A. 2010. Molecular and physiological
analysis of drought stress in *Arabidopsis* reveals early responses leading to
acclimation in plant growth. *Plant Physiol.* 154:1254–1271.

Harrison PW, Wright AE, Mank JE. 2012. The evolution of gene expression and the
transcriptome–phenotype relationship. *Semin. Cell Dev. Biol.* 23:222–229.

Jill Harrison C. 2017. Development and genetics in the evolution of land plant body
plans. *Philos. Trans. R. Soc. B Biol. Sci.* 372:20150490.

Josephs EB, Wright SI, Stinchcombe JR, Schoen DJ. 2017. The relationship between
selection, network connectivity, and regulatory variation within a population of
Capsella grandiflora. *Genome Biol. Evol.* 9:1099–1109.

Jovelin R, Phillips PC. 2009. Evolutionary rates and centrality in the yeast gene
regulatory network. *Genome Biol.* 10:1–10.

Juenger TE. 2013. Natural variation and genetic constraints on drought tolerance.
Curr. Opin. Plant Biol. 16:274–281.

Kashyap SP, Prasanna HC, Kumari N, Mishra P, Singh B. 2020. Understanding salt
tolerance mechanism using transcriptome profiling and *de novo* assembly of wild
tomato *Solanum chilense*. *Sci. Rep.* 10:1–20.

Kim PM, Korbel JO, Gerstein MB. 2007. Positive selection at the protein network
periphery: evaluation in terms of structural constraints and cellular context. *Proc. Natl.*
Acad. Sci. 104:20274–20279.

Koenig D, Jiménez-Gómez JM, Kimura S, Fulop D, Chitwood DH, Headland LR,
Kumar R, Covington MF, Devisetty UK, Tat AV, et al. 2013. Comparative
transcriptomics reveals patterns of selection in domesticated and wild tomato. *Proc.*
Natl. Acad. Sci. 110:E2655–E2662.

Korneliussen TS, Albrechtsen A, Nielsen R. 2014. ANGSD: Analysis of Next
Generation Sequencing Data. *BMC Bioinformatics* 15:356.

Koubkova-Yu TC-T, Chao J-C, Leu J-Y. 2018. Heterologous Hsp90 promotes
phenotypic diversity through network evolution. *PLoS Biol.* 16:e2006450.

Kumar R, Ichihashi Y, Kimura S, Chitwood DH, Headland LR, Peng J, Maloof JN,
Sinha NR. 2012. A high-throughput method for Illumina RNA-Seq library preparation.
Front. Plant Sci. 3:202.

Langfelder P, Horvath S. 2008. WGCNA: an R package for weighted correlation
network analysis. *BMC Bioinformatics* 9:559.

Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program
for assigning sequence reads to genomic features. *Bioinformatics* 30:923–930.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15:550.

Luisi P, Alvarez-Ponce D, Pybus M, Fares MA, Bertranpetit J, Laayouni H. 2015.
Recent positive selection has acted on genes encoding proteins with more
interactions within the whole human interactome. *Genome Biol. Evol.* 7:1141–1154.

Mähler N, Wang J, Terebieniec BK, Ingvarsson PK, Street NR, Hvidsten TR. 2017.
Gene co-expression network connectivity is an important determinant of selective
constraint. *PLoS Genet.* 13:e1006402.

Martínez JP, Antúnez A, Araya H, Pertuzé R, Fuentes L, Lizana XC, Lutts S. 2014.
Salt stress differently affects growth, water status and antioxidant enzyme activities in

Solanum lycopersicum and its wild relative *Solanum chilense. Aust. J. Bot.* 62:359–
368.

Masalia RR, Bewick AJ, Burke JM. 2017. Connectivity in gene coexpression
networks negatively correlates with rates of molecular evolution in flowering plants.
PLoS One 12:e0182289.

Mboup M, Fischer I, Lainer H, Stephan W. 2012. Trans-species polymorphism and
allele-specific expression in the *CBF* gene family of wild tomatoes. *Mol. Biol. Evol.*
29:3641–3652.

Mes PJ, Boches P, Myers JR, Durst R. 2008. Characterization of tomatoes
expressing anthocyanin in the fruit. *J. Am. Soc. Hortic. Sci.* 133:262–269.

Mi H, Ebert D, Muruganujan A, Mills C, Albou L-P, Mushayamaha T, Thomas PD.
2021. PANTHER version 16: a revised family classification, tree-based classification
tool, enhancer regions and extensive API. *Nucleic Acids Res.* 49:D394–D403.

Molitor C, Kurowski TJ, Fidalgo de Almeida PM, Eerolla P, Spindlow DJ, Kashyap SP,
Singh B, Prasanna HC, Thompson AJ, Mohareb FR. 2021. *De novo* genome
assembly of *Solanum sitiens* reveals structural variation associated with drought and
salinity tolerance.Valencia A, editor. *Bioinformatics* 37:1941–1945.

Moutinho AF, Eyre-Walker A, Dutheil JY. 2022. Strong evidence for the adaptive walk
model of gene evolution in Drosophila and Arabidopsis. *PLOS Biol.* 20:e3001775.

Mustafin ZS, Zamyatin VI, Konstantinov DK, Doroshkov AV, Lashin SA, Afonnikov DA.
2019. Phylostratigraphic analysis shows the earliest origination of the abiotic stress
associated genes in *A. thaliana*. *Genes* 10:963.

Nakazato T, Warren DL, Moyle LC. 2010. Ecological and geographic modes of
species divergence in wild tomatoes. *Am. J. Bot.* 97:680–693.

Nicolas P, Shinozaki Y, Powell A, Philippe G, Snyder SI, Bao K, Zheng Y, Xu Y,
Courtney L, Vrebalov J, et al. 2022. Spatiotemporal dynamics of the tomato fruit
transcriptome under prolonged water stress. *Plant Physiol.* 190:2557–2578.

Nieri D, Di Donato A, Ercolano MR. 2017. Analysis of tomato meiotic recombination
profile reveals preferential chromosome positions for NB-LRR genes. *Euphytica*
213:206.

Nosenko T, Böndel KB, Kumpfmüller G, Stephan W. 2016. Adaptation to low
temperatures in the wild tomato species *Solanum chilense. Mol. Ecol.* 25:2853–2869.

Papakostas S, Vøllestad LA, Bruneaux M, Aykanat T, Vanoverbeke J, Ning M,
Primmer CR, Leder EH. 2014. Gene pleiotropy constrains gene expression changes
in fish adapted to different thermal conditions. *Nat. Commun.* 5:1–9.

Pertea G, Pertea M. 2020. GFF Utilities: GffRead and GffCompare. Available from:
https://f1000research.com/articles/9-304

Piasecka B, Lichocki P, Moretti S, Bergmann S, Robinson-Rechavi M. 2013. The
hourglass and the early conservation models—co-existing patterns of developmental
constraints in vertebrates. *PLoS Genet.* 9:e1003476.

Quint M, Drost H-G, Gabel A, Ullrich KK, Bönn M, Grosse I. 2012. A transcriptomic
hourglass in plant embryogenesis. *Nature* 490:98–101.

R Core Team. 2020. R: A language and environment for statistical computing.

Raduski AR, Igić B. 2021. Biosystematic studies on the status of *Solanum chilense*.
Am. J. Bot. 108:520–537.

Reddy AR, Chaitanya KV, Vivekanandan M. 2004. Drought-induced responses of
photosynthesis and antioxidant metabolism in higher plants. *J. Plant Physiol.*
161:1189–1202.

Rodrigues J, Inzé D, Nelissen H, Saibo NJ. 2019. Source–sink regulation in crops
under water deficit. *Trends Plant Sci.* 24:652–663.

Roselius K, Stephan W, Städler T. 2005. The relationship of nucleotide polymorphism,
recombination rate and selection in wild Tomato species. *Genetics* 171:753–763.

Sato MP, Makino T, Kawata M. 2016. Natural selection in a population of *Drosophila*
melanogaster explained by changes in gene expression caused by sequence
variation in core promoter regions. *BMC Evol. Biol.* 16:1–12.

Shen W, Ren H. 2021. TaxonKit: A practical and efficient NCBI taxonomy toolkit. *J.*
Genet. Genomics.

Skirycz A, Inzé D. 2010. More from less: plant growth under limited water. *Curr. Opin.*
Biotechnol. 21:197–203.

Spivakov M. 2014. Spurious transcription factor binding: non-functional or genetically
redundant? *Bioessays* 36:798–806.

Stuart JM, Segal E, Koller D, Kim SK. 2003. A gene-coexpression network for global
discovery of conserved genetic modules. *science* 302:249–255.

Su G, Morris JH, Demchak B, Bader GD. 2014. Biological network exploration with
Cytoscape 3. *Curr. Protoc. Bioinforma.* 47:8.13.1-8.13.24.

Swanson-Wagner R, Briskine R, Schaefer R, Hufford MB, Ross-Ibarra J, Myers CL,
Tiffin P, Springer NM. 2012. Reshaping of the maize transcriptome by domestication.
Proc. Natl. Acad. Sci. 109:11878–11883.

Tapia G, Castro M, Gaete-Eastman C, Figueroa CR. 2022. Regulation of anthocyanin
biosynthesis by drought and UV-B radiation in wild tomato (*Solanum peruvianum*)
fruit. *Antioxidants* 11:1639.

Tapia G, Méndez J, Inostroza L. 2016. Different combinations of morpho-
physiological traits are responsible for tolerance to drought in wild tomatoes *Solanum*
chilense and *Solanum peruvianum.* *Plant Biol.* 18:406–416.

Tian F, Yang D-C, Meng Y-Q, Jin J, Gao G. 2020. PlantRegMap: charting functional
regulatory maps in plants. *Nucleic Acids Res.* 48:D1104–D1113.

Townsley BT, Covington MF, Ichihashi Y, Zumstein K, Sinha NR. 2015. BrAD-seq:
Breath Adapter Directional sequencing: a streamlined, ultra-simple and fast library
preparation protocol for strand specific mRNA library construction. *Front. Plant Sci.* 6.

True JR, Haag ES. 2001. Developmental system drift and flexibility in evolutionary
trajectories. *Evol. Dev.* 3:109–119.

Verelst W, Bertolini E, De Bodt S, Vandepoele K, Demeulenaere M, Pè ME, Inzé D.
2013. Molecular and physiological analysis of growth-limiting drought stress in
Brachypodium distachyon leaves. *Mol. Plant* 6:311–322.

de Vries J, Archibald JM. 2018. Plant evolution: landmarks on the path to terrestrial
life. *New Phytol.* 217:1428–1434.

de Vries J, Curtis BA, Gould SB, Archibald JM. 2018. Embryophyte stress signaling
evolved in the algal progenitors of land plants. *Proc. Natl. Acad. Sci.* 115:E3471–
E3480.

Wagner GP, Kin K, Lynch VJ. 2012. Measurement of mRNA abundance using RNA-
seq data: RPKM measure is inconsistent among samples. *Theory Biosci* 131:281–
285.

Wagner GP, Pavlicev M, Cheverud JM. 2007. The road to modularity. *Nat. Rev.*
Genet. 8:921–931.

Wang S, Li L, Li H, Sahu SK, Wang H, Xu Y, Xian W, Song B, Liang H, Cheng S, et al.
2020. Genomes of early-diverging streptophyte algae shed light on plant
terrestrialization. *Nat. Plants* 6:95–106.

Wang Y, Wang X, Paterson AH. 2012. Genome and gene duplications and gene
expression divergence: a view from plants. *Ann. N. Y. Acad. Sci.* 1256:1–14.

Wei K, Silva-Arias GA, Tellier A. 2023. Selective sweeps linked to the colonization of
novel habitats and climatic changes in a wild tomato species. *New Phytol.* 237:1908–
1921.

Wilkinson S, Davies WJ. 2010. Drought, ozone, ABA and ethylene: new insights from
cell to plant to community. *Plant Cell Environ.* 33:510–525.

Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R. 2009. The
Sequence alignment/map (SAM) format and SAMtools. *Bioinformatics* 25:2078–2079.

Yang L, Bu S, Zhao S, Wang N, Xiao J, He F, Gao X. 2022. Transcriptome and
physiological analysis of increase in drought stress tolerance by melatonin in tomato.
PLOS ONE 17:e0267594.

Yang X, Lu M, Wang Yufei, Wang Yiran, Liu Z, Chen S. 2021. Response mechanism
of plants to drought stress. *Horticulturae* 7:50.

Yu G, Wang L-G, Han Y, He Q-Y. 2012. clusterProfiler: an R Package for Comparing
Biological Themes Among Gene Clusters. *OMICS J. Integr. Biol.* 16:284–287.

Zarrineh P, Sánchez-Rodríguez A, Hosseinkhan N, Narimani Z, Marchal K, Masoudi-
Nejad A. 2014. Genome-scale co-expression network comparison across *Escherichia*
coli and *Salmonella enterica* serovar *Typhimurium* reveals significant conservation at
the regulon level of local regulators despite their dissimilar lifestyles. *PLoS One*
9:e102871.

Zinkgraf M, Zhao S-T, Canning C, Gerttula S, Lu M-Z, Filkov V, Groover A. 2020.
Evolutionary network genomics of wood formation in a phylogenetic survey of
angiosperm forest trees. *New Phytol.* 228:1811–1823.

1030