A mathematical model for axonal transport of large cargo vesicles

Nizhum Rahman[1] and Dietmar B. Oelz[1]

[1] School of Mathematics and Physics, The University of Queensland, QLD, 4072, Australia.

January 18, 2023

Abstract: In this study, we consider axonal transport of large cargo vesicles characterised by transient expansion of the axon shaft. Our goal is to formulate a mathematical model which captures the dynamic mechanical interaction of such cargo vesicles with the membrane associated periodic cytoskeletal structure (MPS). It consists of regularly spaced actin rings that are transversal to the longitudinal direction of the axon and involved in the radial contraction of the axon. A system of force balance equations is formulated by which we describe the transversal rings as visco-elastic Kelvin-Voigt elements. In a homogenisation limit, we reformulate the model as a free boundary problem for the interaction of the submembranous MPS with the large vesicle. We derive a non-linear force-velocity relation as a quasi-steady state solution. Computationally we analyse the vesicle size dependence of the transport speed and use an asymptotic approximation to formulate it as a power law that can be tested experimentally.

Keywords: axonal transport; motor protein; actin rings; potential energy; obstacle problem; asymptotic approximation.

1 Introduction

Neurons are the foundational units of the nervous system. They are involved with information processing and transport along their long cellular extensions termed dendrites and axons. Tightly regulated axonal transport of multiple types of cargo such as synaptic vesicles, organelles and proteins packed into cargo vesicles is paramount for the functioning of the neuron. This involves anterograde axonal transport driven by molecular motors of the kinesin family and dynein-dependent retrograde transport [1, 2].

The axon shaft is characterised by a particular cytoskeletal organisation termed membrane associated periodic skeleton (MPS). It involves a sub-membraneous actin cortex and subcortical evenly spaced actin rings with a periodicity of $180 - 190$nm [3, 4]. They are made of a few long intertwined actin filaments forming actin braids [5]. The MPS stabilises the axon diameter [6–8]. Indeed it has been reported that F-Actin disassociation leads to the collapse of axons [9] whereas inhibition of the F-Actin capping protein α-adducin leads to axon diameter expansion [10]. Non-muscle myosin IIa and IIB are associated with the periodic actin rings [6] and involved in radial contractility [11].

The protein spectrin couples neighbouring rings through an elastic connection and contributes to the stability of the axon [6, 12]. A longitudinally aligned system of cytoskeletal elements involves F-actin trails, neurofilaments and microtubules [8]. Particularly the latter serve as tracks for the molecular motors involved in axonal transport.

The diameter of the axon shafts varies among different tissues and species and ranges from 0.1 µm to several microns [13]. Cargo sizes vary within a wide range which includes mitochondria (0.75–3µm) [14], autophagosomes (0.5–1.5µm) [15] and endosomes (50nm–1µm) [16]. Thus, cargo sizes are comparable to the size of various CNS axons, often exceeding it. Radial axonal contractility in combination with transient expansion of axon calibre facilitates the passage of large cargoes. When a large cargo passes through the axon, the axon acts like an elastic material avoiding the further delay of cargo passage. Subsequent radial contraction of the transversal actin rings is myosin dependent[11]. Yet, whether the actin rings are open loops, or closed rings (see figure 1) is an open question [6].

Figure 1: Actin rings (sketched according to the closed ring hypothesis [6]) are wrapped around the circumference of the axon with equi-distant spacing. Vesicles of various sizes are moving through the axon.

The speed of axonal transport varies significantly between slow axonal transport (< 0.1 µm/s) and fast axonal transport (> 0.5 µm/s) [17]. For large cargo particles, whose diameter exceeds the diameter of the axon shaft, it has recently been reported that the speed of retrograde axonal transport is inversely correlated with the diameter of the cargo particles, i.e. large-diameter cargo moving slower than smaller-diameter cargo [11].

One possible explanation is size-dependent friction on the cargo originating from the mechanical interaction with the axonal cortex [11]. This hypothesis is also supported by a simulation study suggesting that the effective viscosity between organelles and the axon wall at a small distance can become large enough to impede cargo transport[18].

Another potential mechanism, which might explain the slower speed of transport of large cargo particles, is tight spatial alignment of microtubules in small calibre axons which could stabilise the pool of molecular motors engaged in active transport. When the axon is stretched by large calibre cargo distances between parallel microtubules might be larger and impede likely reattachment of molecular motor proteins after detachment [18].

Despite these theories, it is an open question how axonal radial contractility regulates cargo transport [11]. In particular, the observation that in myosin inhibited cells trajectories of large cargo retrograde transport visualised through kymographs appear more random and even exhibit frequent backtracking [11] seems to contradict the friction hypothesis, but could be supported by the microtubule alignment hypothesis. Another potential explanation could originate from the direct mechanical interaction of the contractile actin rings. In this study, it is our goal to develop the modelling framework to investigate this interaction between the transversal actin rings and large cargo particles.

To this end, we formulate a detailed mathematical model which couples the active and passive mechanical properties of the actin rings modelled as visco-elastic Kelvin-Voigt elements. We start by formulating the model in a time-discrete setting with a fixed spacing of the actin rings. In a second step, we formally pass to a continuum limit of both time and actin ring

spacing, both in section 2. In the following sections, we process the model further. In section 3 we non-dimensionalise. The resulting mathematical model resembles the classical obstacle problem [19] which is a model problem in the study of free boundary problems. In this study, the free boundary is the interface where the cargo vesicle touches the axonal cortex. We consider quasi-stationary solutions of this problem from which we derive the force-velocity relation for the transport of large cargos in section 4. In section 5 we study the dependence of transport speed on vesicle size numerically, and we use asymptotic approximation to formulate the decrease of transport speed for large vesicle sizes as a power law valid for vesicle sizes which are close to the axon size.

2 Mathematical Model

Our goal is to develop a mathematical model which relates the forces, especially those exerted by the transversal actin rings and their associated myosin motor proteins, to the velocity of large cargo vesicles undergoing axonal transport.

To introduce the model we regard the axon as a one-dimensional structure decorated with concentric rings located at equi-distant positions $y_j \in \mathbb{R}$ (figures 1 and 2). Here we use $j = 1...M$ as an index for actin rings assuming – for notational convenience – the presence of M actin rings in total.

The time-dependent degrees of freedom of the model are the position $x = x(t) \in \mathbb{R}$ of a large cargo vesicle with radius R_v and the radii of actin rings written as $r_j = r_j(t) > 0$ where $t \geq 0$ refers to time.

Figure 2: As the cargo vesicle (green) is moving through the axon it extends the radii of the actin rings.

We construct the model as a system of force-balance equations taking into account the following forces acting on the cargo vesicle and the actin rings.

We assume that the cargo vesicle is pulled by the given force F, which, depending on direction, may be positive or negative, and that through its interaction with the cytoplasm the cargo vesicle is exposed to the drag friction F_{friction}, which we model as linear drag with coefficient $\xi > 0$,

$$F_{\text{friction}} = -\xi \, \dot{x}(t) \, ,$$

where $\dot{x}$ is the notation we use for the velocity of the cargo vesicle. Note that when modelling the pulling force exerted by molecular motors [20, 21] through an affine force-velocity relation, the parameters F and ξ will be determined by stall force and free moving velocity of the molecular motors.

We model the actin rings as visco-elastic Kelvin-Voigt elements with equilibrium radius R_a (figure 2). This implies that the elastic and viscous forces exerted by every ring are in parallel.

Its elastic component, which in (figure 2) is visualised by a spring, is given by

$$F_{\text{elastic}} = -\kappa \left(r_j(t) - R_a \right) ,$$

where $\kappa > 0$ is the constant of elasticity. Note that this implies that we are modelling myosin motors indirectly as drivers of radial contraction, namely through the equilibrium radius of actin rings R_a, and through their stretching stiffness κ.

The viscous resistance to changes in actin ring size – in figure 2 visualised by a dashpot – is given by

$$F_{\text{viscous}} = -\chi \, \dot{r}_j(t) ,$$

where χ is the coefficient of viscosity. It models the internal resistance of the actin rings to shortening and stretching causing stretching and realignment of actin fibers and cross-linker proteins as well as transient unbinding from crosslinker proteins and myosin motors.

We assume that the cargo vesicle and the actin rings are mechanically interacting: when the size of the vesicle is close to the radii of the rings, the vesicle would push the actin rings apart so it can pass. The resulting force is modelled as a reactive force resulting from the impenetrability of vesicles and rings.

To formulate a closed system of equations we introduce the following variational formulation of the model, in which every force term above is represented by a corresponding conservative or dissipative energy, and in which the constraint is represented by a penalising potential. We compute the time-discrete solution x^n, r_j^n for every index of time $n = 1, \ldots$ with timesteps Δt as a minimiser,

$$\left(x^n, (r_j^n)_{j=1\ldots M} \right) = \text{argmin} \, E[x^{n-1}, (r_j^{n-1})_{j=1..M}](x, (r_j)_{j=1\ldots M}) \tag{1}$$

of an energy functional with respect to the configuration of the system, i.e. actin ring radii and vesicle position. The functional is also parametrised by the configuration of the system at the previous point in time written as x^n, $(r_j^n)_{j=1\ldots M}$. Its components are

$$E[x^{n-1}, (r_j^{n-1})_{j=1..M}](x, (r_j)_{j=1..M}) = E_{\text{ext}} + E_{\text{friction}} + E_{\text{elastic}} + E_{\text{viscous}} + E_{\text{impenetrability}} . \tag{2}$$

The components of the energy functional corresponding to conservative and dissipative forces are

$$E_{\text{ext}} = -F \, x ,$$

$$E_{\text{friction}} = \xi \frac{(x - x^{n-1})^2}{2 \, \Delta t} ,$$

$$E_{\text{elastic}} = \kappa \sum_{j=1}^{M} \frac{(r_j - R_a)^2}{2} ,$$

$$E_{\text{viscous}} = \chi \sum_{j=1}^{M} \frac{(r_j - r_j^{n-1})^2}{2 \, \Delta t} ,$$

which are chosen in a way such that the negative variation of these energies evaluated at x^n, $(r_j^n))$ corresponds to the forces listed above. For dissipative forces this holds in the limit when $\Delta t \to 0$, e.g. $F_{\text{viscous}} = -\chi \lim_{\Delta t \to 0} \frac{r_j^n - r_j^{n-1}}{\Delta t}$.

The penalising potential which enforces the constraint that the actin rings do not penetrate the vesicle is given by

$$E_{\text{impenetrability}} = \sum_{j: \, |x - y_j| < R_v} \frac{1}{2\delta} \left(h(x - y_j) - r_j \right)_+^2 , \tag{3}$$

where δ is a small parameter implying that the coefficient in front of the penalising potential is large. In (3) we restrict the sum to those actin rings which are sufficiently close for interaction with the vesicle and we use the subscript $(.)_+$ as a short notation for the positive part of what is inside the bracket, i.e. $(x)_+ = x$ if $x > 0$, otherwise it is zero. The respective term contributes to the sum (3) if the cross-sectional radius h of the vesicle at the position of the ring is larger than the radius of that actin ring. It is computed as a function of the distance between ring and vesicle position, $h = h(x - y_j)$ according to (see figure 2)

$$h(d) = \begin{cases} 0 & \text{if} \quad |d| \geq R_v , \\ \sqrt{R_v^2 - d^2} & \text{if} \quad |d| < R_v . \end{cases} \tag{4}$$

This finalises the formulation of the mathematical model as a variational scheme in the time-discrete setting.

Next we will derive the equations of motion. The solution to the variational scheme is characterised by $0 = \frac{d}{dx}E(x^n, r_1^n, \ldots, r_M^n)$, which implies

$$0 = \xi \frac{x^n - x^{n-1}}{\Delta t} - F + \frac{1}{\delta} \sum_{j: \, |x-y_j| < R_v} (h(x^n - y_j) - r_j^n)_+ \, h'(x^n - y_j) , \tag{5}$$

and by $0 = \frac{d}{dr_j}E(x^n, r_1^n, \ldots, r_M^n)$ for every $j = 1 \ldots M$, which yields

$$0 = \kappa \, (r_j^n - R_a) + \chi \, \frac{r_j^n - r_j^{n-1}}{\Delta t} - \frac{1}{\delta} \, (h(x^n - y_j) - r_j^n)_+ . \tag{6}$$

In the limit when the time steps are small, i.e. $dt \to 0$, we write $\frac{x^n - x^{n-1}}{dt} \to \dot{x}$ and $\frac{r_j^n - r_j^{n-1}}{dt} \to \dot{r}_j$ and the system (5), (6) reads

$$\xi \, \dot{x} = F - \sum_{j: \, |x-y_j| < R_v} \frac{1}{\Delta y} \frac{1}{\delta} \, (h(x^n - y_j) - r_j^n)_+ \, h'(x^n - y_j) \, \Delta y ,$$
$$\frac{1}{\Delta y}\chi \, \dot{r}_j = \frac{1}{\Delta y}\kappa \, (R_a - r_j) + \frac{1}{\Delta y} \frac{1}{\delta} \, (h(x^n - y_j) - r_j^n)_+ . \tag{7}$$

In the formal limit when the penalisation of the impenetrability constraint becomes large and the spacing of the actin rings small, $\delta \to 0$ and $\Delta y \to 0$, we define the Lagrange multiplier formally as a limit,

$$\frac{1}{\Delta y} \frac{1}{\delta}(h(x - y) - r(t))_+ \to \lambda(t, y) , \tag{8}$$

which is a force density corresponding to the interaction force between vesicle and actin rings. We also identify $\bar{\chi}$ with the limit of $\chi/\Delta y$ and $\bar{\kappa}$ with the limit of $\kappa/\Delta y$. Note that the vesicle position $x = x(t)$ is a dependent variable whereas $y \in \mathbb{R}$ is an independent variable of our model. We get the following system of two equations for the functions $x = x(t)$ (vesicle position), $r = r(t, y)$ (actin ring/axon radius) and $\lambda = \lambda(t, y)$ (interaction force density), see figure 3,

Figure 3: The degrees of freedom of the continuum model (9)-(11): The cargo vesicle at position $x(t)$ is touching the axon with radius $r = r(t, y)$ along the interval (A, B). The radius of the segment of the vesicle which is perpendicular to the axon is written as $h(x(t), y)$.

$$\xi \, \dot{x}(t) \;=\; F - \int_{A(t)}^{B(t)} \lambda(t, y) \, h'(x - y) \, dy \;, \tag{9}$$

$$\bar{\chi} \, \dot{r} \;=\; \bar{\kappa} \, (R_a - r) + \lambda \;, \tag{10}$$

$$r(t, y) \;\geq\; h(x(t) - y) \;. \tag{11}$$

Here A and B define the interval (A, B) where the axon - represented by the transversal actin rings – touches the vesicle, i.e. where (11) is an equality. This equality determines the interaction force density $\lambda = \lambda(t, x)$ within the interval $(A, B) \subset \mathbb{R}$, outside of which λ is zero.

Figure 4: Derivation of the free boundary condition at the rear of the vesicle (here moving from right to left).

Note that an additional equation is derived as follows: In figure 4 which is sketched in a regime where both time and the spacing of actin rings are discrete, we consider the boundary

point at B and assume that it moves to the left, i.e. $\dot{B} < 0$. We consider the actin ring at y_j which at time n is immediately to the right of B^n and which at the previous point in time $n-1$ is immediately to the left of the interface point. This suggests that the force balance equation according to (6) is given by

$$0 = \kappa(r_j^n - R_a) + \chi \frac{r_j^n - r_j^{n-1}}{\Delta t} \,,$$

where the penalisation term for the interaction of vesicle and axon is zero. Since at the previous point in time $n-1$ this actin ring does touch the vesicle, we replace r_j^{n-1} by $h(x^{n-1} - y_j)$. In the homogenisation procedure by which $\Delta t \to 0$ and $\Delta y \to 0$ we may place the ring j arbitrarily close to B. By continuity we replace r_j^n by $r^n(B) = h(x^n - B)$ obtaining $0 = \kappa(r_j^n - R_a) + \chi \frac{h(x^n - B^n) - h(x^{n-1} - B^n)}{\Delta t}$. In the aforementioned limit we find that

$$0 = \bar{\kappa} \left(r(t, B) - R_a \right) + \bar{\chi} \, h'(x(t) - B(t)) \, \dot{x}(t) \,.$$

On the other hand, the limit of (10) evaluated at a sequence of points which converges to $B(t)$ from the right implies that $0 = \bar{\kappa}(r(t,B) - R_a) + \bar{\chi}\dot{r}(B(t))$. Therefore $h'(x(t) - B(t))\dot{x}(t) = \dot{r}(B(t))$.

Note that the opposite assumption, i.e. $\dot{B} > 0$, does not imply such a condition. For a fibre that would be immediately left from $B(t)$, though at any previous point in time right of $B(t)$, the force balance equation (6) would have a non-zero interaction term. The same equation taken from away of the vesicle would not involve this term $0 = \bar{\kappa} \left(r(t,B) - R_a \right) + \bar{\chi} \, \dot{r}(B(t))$ and would therefore not allow to draw a conclusion about the time derivatives.

Finally, the same arguments applied the the left interface point assuming $\dot{A}(t) > 0$ suggests that

$$\dot{r}(t, A) = h'(x(t) - A(t)) \, \dot{x}(t) \,. \tag{12}$$

3 Nondimensionalization

To facilitate the further analysis of the model (9)-(11) we non-dimensionalise introducing the re-scaled independent and dependent variables $\tilde{x} = \frac{x}{R_a}$, $\tilde{y} = \frac{y}{R_a}$, $\tilde{r} = \frac{r}{R_a}$, $\tilde{A} = \frac{A}{R_a}$, $\tilde{B} = \frac{B}{R_a}$, $\tilde{R}_v = \frac{R_v}{R_a}$ and $\tilde{t} = \frac{t}{t_c}$, where t_c is the reference scale for time which we will determine further below. The system (9)-(11) becomes

$$\xi \, \dot{\tilde{x}}(t) = \frac{t_c}{R_a} F - \frac{t_c}{R_a} R_a \int_{\tilde{A}(t)}^{\tilde{B}(t)} \lambda(t_c \tilde{t}, R_a \tilde{y}) \, h'(\tilde{x}(t) - \tilde{y}) \, d\tilde{y} \,,$$

$$\dot{\tilde{r}} = \frac{t_c}{R_a \bar{\chi}} \bar{\kappa} \, R_a (1 - \tilde{r}) + \frac{t_c}{R_a \bar{\chi}} \lambda(t_c \tilde{t}, R_a \tilde{y}) \,,$$

$$R_a \, \tilde{r}(\tilde{t}, \tilde{y}) \geq R_a \, h(\tilde{x}(t) - \tilde{y}) \,,$$

where we used that $h'(R_a \tilde{d}) = \tilde{h}'(\tilde{d})$ ($\tilde{h}'$ is defined by (4) for the scaled vesicle radius $\tilde{R}_v$). We determine t_c such that $\frac{\bar{\kappa} \, t_c}{\bar{\chi}} = \tilde{\kappa} = 1$, i.e. $t_c = \frac{\bar{\chi}}{\bar{\kappa}} = \frac{\chi}{\kappa}$. Let $\tilde{\xi} = \frac{\xi}{R_a \bar{\chi}}$, $\tilde{\lambda}(\tilde{t}, \tilde{y}) = \frac{t_c}{R_a \, \bar{\chi}} \lambda(\tilde{t} \, t_c, \tilde{y} \, R_a)$ and $\tilde{F} = \frac{F \, t_c}{R_a^2 \, \bar{\chi}} = \frac{F}{R_a^2 \, \bar{\kappa}}$. We obtain

$$\begin{cases} \xi \, \dot{x}(t) = F - \int_{A(t)}^{B(t)} \lambda(t, y) \, h'(x(t) - y) \, dy \,, \\ \quad \dot{r} = 1 - r + \lambda \,, \\ r(t, y) \geq h(x(t) - y) \,, \end{cases} \tag{13}$$

Description	Symbol	Value	References
Radius of cargo vesicle	R_v	0.3 µm	[11]
Equilibrium radius of axon	R_a	0.2 µm	[11, 13]
Pulling force	F	15 pN	estimated
Spacing between the actin rings	Δy	0.18 µm	[3]
Drag friction	ξ	10 pN µm^{-1} sec	estimated
Stretching stiffness of actin rings	κ	30 pN µm^{-1}	estimated
Viscosity of actin rings	χ	5 pN µm^{-1} sec	estimated

Table 1: List of parameters values

whee we have omitted the tildes again. Note that the interaction force vanishes, $\lambda \equiv 0$, where the axon does not touch the vesicle, i.e. where $r(t, y) > h(x(t) - y)$.

The parameters of the model are listed in table 1. We estimate that the cargo velocity V is of the order of magnitude 1.5 µm/sec [11] and therefore we determine the drag friction coefficient as $\xi \approx F/V = 10$ pN sec/µm having in mind that this refers to a cargo vesicle the size of which does not exceed the axon diameter. Furthermore, with the parameters listed in table 1, we obtain the dimensionless parameters $\tilde{F} = \frac{F\Delta y}{R_a^2 \kappa} = 2.25$ and $\tilde{\xi} = \frac{\xi \Delta y}{R_a \chi} = 1.8$.

4 Force-velocity relation

In this section, it is our goal to determine the force-velocity relation predicted by our model as the quasi-stationary solution characterised by a constant vesicle velocity V which we assume to satisfy $V > 0$. This implies that $x(t) = Vt$ as well as $A(t) = \bar{A} + Vt$ and $B(t) = \bar{B} + Vt$, where we expect $\bar{A} < 0 < \bar{B}$ (see figure 5).

The ring/axon radius r as well as the interaction force/Lagrange multiplier λ in this scenario will be simply advected with speed V and have the structure of travelling wave solutions $r(t, y) = \bar{r}(y - Vt)$ and $\lambda(t, y) = \bar{\lambda}(y - Vt)$, where $\bar{r} = \bar{r}(\eta)$ and $\bar{\lambda} = \bar{\lambda}(\eta)$ only depend on the travelling wave variable $\eta = y - Vt$.

With this ansatz – omitting the bar-notation again – the system (13) becomes

$$\begin{cases} \xi V = F - \int_A^B \lambda(\eta) \, h'(-\eta) \, d\eta \,, \\ -V \, r' = 1 - r + \lambda \,, \\ r(\eta) \geq h(\eta) \,. \end{cases} \tag{14}$$

where we used $h(-\eta) = h(\eta)$. In addition, since $\dot{A} = V > 0$, the free boundary condition (12) translated into travelling wave variables holds, i.e. $r'(A) = h'(A)$. A numerical solution corresponding to the travelling wave solutions is sketched in figure 5.

We solve the equation $-V \, r' = 1 - r$ on the interval $(-\infty, A)$ where the axon and the vesicle do not touch and find the general solution $r(y) = C_1 \, e^{y/V} + 1$.

Now we consider the interface point at A where axon and vesicle touch. It therefore holds that $r(A) = h(A)$ which allows to infer the value of C_1. We find that

$$r(y) = e^{\frac{1}{V}(y - A)} \left(\sqrt{R_v^2 - A^2} - 1 \right) + 1 \,.$$

The solution of $r(y)$ between points A and B is the height of the vesicle h. We make the modelling assumption that as the vesicle moves it progresses into an axon which is initially not

Figure 5: Travelling wave solution with vesicle positioned at $x = 0$ for the dimensionless parameters $F = 2.25$ and $\xi = 1.8$ corresponding to the physical parameters listed in table 1. The cargo (red) touches the axon (blue) along the interval $[A, B]$.

stretched. The solution r to the right of B is therefore constant and equal to the axon radius $R_a = 1$ (after nondimensionalisation). This implies

$$r(y) = \begin{cases} e^{(y-A)/V} \left(\sqrt{R_v^2 - A^2} - 1 \right) + 1 & -\infty < y < A \,, \\ \sqrt{R_v^2 - y^2} & A \le y \le B \,, \\ 1 & B < y < \infty \,, \end{cases} \tag{15}$$

which still depends on A. The point B is to the right of $x = 0$ and defined by $R_v^2 = 1 + B^2$, i.e.

$$B = \sqrt{R_v^2 - 1} \,. \tag{16}$$

We finally use that $h'(A) = r'(A)$ to compute A as a function of V. Using (15) this equation reads $\sqrt{R_v^2 - A^2} - 1 + \frac{A V}{\sqrt{R_v^2 - A^2}} = 0$ which implies that

$$V[A] = \frac{1}{A} \left(\sqrt{R_v^2 - A^2} - R_v^2 \right) + A \,. \tag{17}$$

Note that the function $V(A) : [-\sqrt{R_v^2 - 1}, 0] \mapsto (0, \infty)$ is monotonically increasing since $V'[A] = \frac{R_v^2}{A^2} \left(1 - 1/\sqrt{R_v^2 - A^2} \right) + 1 \ge 1$ and convex since $V''[A] = \frac{A^2 R_v^2 + (\sqrt{R_v^2 - A^2} - 1)(R_v^2 - A^2) 2 R_v^2}{-A^3 (R_v^2 - A^2) \sqrt{R_v^2 - A^2}} > 0$ where we use $\sqrt{R_v^2 - A^2} > 1$.

In addition to $V(-\sqrt{R_v^2 - 1}) = 0$ and $\lim_{A \to 0} V(A) = +\infty$ we also find that $V'(-\sqrt{R_v^2 - 1}) = 1$ and $\lim_{A \to 0} V'(A) = +\infty$. As a consequence, taking the reciprocal values of the derivatives, we find that $A(0) = -\sqrt{R^2 - 1}$, $\lim_{V \to \infty} A(V) = 0$, $A'(0) = 1$ and $\lim_{V \to \infty} A'(V) = 0$.

We also consider A as a function of V, namely the inverse of (17). As the inverse of $V(A)$, it is monotonically increasing and concave. Note that in principle (17) can be inverted using the solution formulas for quartic equations. The result, however, is not of practical value, we therefore plot the graph of (20) numerically in figure 6.

Finally, to find the vesicle speed V, we compute the density of repulsive force λ using the second equation of the system (14), which implies that $\lambda = r - 1 - V r'$. This implies that the Lagrange multiplier λ is given by

$$\lambda(y) = \begin{cases} 0 & -\infty < y \ge A \,, \\ \frac{V y}{\sqrt{R_v^2 - y^2}} + \sqrt{R_v^2 - y^2} - 1 & A < y \le B \,, \\ 0 & B < y < \infty \,. \end{cases} \tag{18}$$

Figure 6: The position of the point A where the axon detaches from the vesicle as a function of the vesicle velocity V for various vesicle radii.

Figure 7: Interaction force λ and local traction $f(y) = \lambda(y)h'(-y)$ for V=0 (A) and for the numerical example in figure 5 (B) where $V = 1.02$, $A = -0.54$, $B = 1.12$.

In figure 7 we show the profile of the interaction force λ as well as the traction $f := \lambda(y)h'(-y)$. The traction is an odd function if $V = 0$ (figure 7A) and doesn't generate any net traction on the vesicle since its integral in (13) vanishes. For $V > 0$ significant traction is generated along the front part of the vesicle (figure 7B).

Note that as a consequence of (17) which is just $\lambda(A) = 0$ it holds that $\lambda > 0$ on $(A, B]$ which confirms that same conclusion drawn from the constructive definition of λ in (8).

We substitute (18) in the first equation of (14) keeping in mind that A depends on V through the inverse of (17),

$$\xi V = F + \int_A^B (h(y) - 1 - V h'(y)) h'(y) dy = \tag{19}$$

$$= F + \left[\frac{1}{2} h(y)^2 - h(y) \right]_A^B - V \int_A^B h'(y)^2 dy =$$

$$= F + \left[\frac{1}{2} (R_v^2 - y^2) - h(y) \right]_A^B - V \int_A^B h'(y)^2 dy =$$

$$= F + \left[-\frac{y^2}{2} - h(y) - V \left(\frac{1}{2} R_v \log \frac{R_v + y}{R_v - y} - y \right) \right]_A^B$$

where we used that $h^2(y) = R_v^2 - y^2$ and the derivative of h' which we obtain differentiating (4).

For the right endpoint of the interval $[A, B]$ we substitute (16) and use (17) obtaining

$$\xi V = F + \left(-\frac{R_v^2 - 1}{2} - 1 + V \left(\sqrt{R_v^2 - 1} - \frac{1}{2} R_v \log \frac{R_v + \sqrt{R_v^2 - 1}}{R_v - \sqrt{R_v^2 - 1}} \right) \right) - $$
$$ - \left[-\frac{A^2}{2} - \sqrt{R_v^2 - A^2} + V \left(A - \frac{1}{2} R_v \log \frac{R_v + A}{R_v - A} \right) \right]_A ,$$

which, after rewriting, becomes

$$F = \xi V - V \sqrt{R_v^2 - 1} + \frac{1}{2} V R_v \left(\log \frac{R_v - A}{R_v + A} + \log \frac{R_v + \sqrt{R_v^2 - 1}}{R_v - \sqrt{R_v^2 - 1}} \right) - \frac{R_v^2 - 1}{2} + \frac{A^2}{2} . \quad (20)$$

Figure 8: Force-velocity relation for $R_v = 1.64$ and $\xi = 0$.

Note that A is taken here as a function of V, namely the inverse of (17). Since the solution formulas for quartic equations are unpractical, we plot the graph of (20) numerically (see figure 8) interpreting it as force-velocity relation for the axonal transport of vesicles with (scaled) radius R_v.

In the supplementary section B we confirm the validity of the equation (20) numerically.

Note that $A(0) = -B$ and therefore $F(0) = 0$. Taking derivatives in (19) and (20) we obtain the following expressions for the derivative $F'(V)$ which is the marginal force to accelerate the vesicle (keep in mind that this is a model for over-damped dynamics),

$$F'[V] = \xi + \int_A^B h'^2 \, dy \, , \quad (21)$$

which implies that F is monotonically increasing since $F' > 0$ and it is also concave since $F''(V) = -h'(A)^2 < 0$. Substituting $B = \sqrt{R_v^2 - 1}$ we obtain

$$F' = \xi + \left[\frac{1}{2} R_v \log \frac{R_v + y}{R_v - y} - y \right]_A^B = \quad (22)$$

$$= \xi - \sqrt{R_v^2 - 1} + A + \frac{1}{2} R_v \left(-\log \frac{R_v + A}{R_v - A} + \log \frac{R_v + \sqrt{R_v^2 - 1}}{R_v - \sqrt{R_v^2 - 1}} \right) \, ,$$

which can only be evaluated numerically. At the endpoints, i.e. for $V = 0$ and in the limit where $V \to \infty$ we can find, though, explicit values using $A(0) = -\sqrt{R^2 - 1}$ and $\lim_{V \to \infty} A(V) = 0$, namely

$$F'[0] = \xi + G_0 \, ,$$

$$\lim_{V \to \infty} F'[V] = \xi + \frac{1}{2} G_0 \, ,$$

where

$$G_0 = R_v \log \frac{R_v + \sqrt{R_v^2 - 1}}{R_v - \sqrt{R_v^2 - 1}} - 2 \sqrt{R_v^2 - 1} \, , \quad (23)$$

which is non-negative because $F' > 0$ for all $\xi \geq 0$. Not taking into account linear drag with coefficient ξ exerted on the vesicle through the crowded cytoplasm, the marginal friction $F'[V]$ decreases by a factor 2 as the velocity increases (Figure 8).

5 Vesicle size dependence of transport speed

In this section, it is our goal to understand how the vesicle speed predicted by our model depends on the vesicle size R_v. We will evaluate this dependence numerically and use asymptotic expansion to compute an approximate solution when the vesicle radius R_v is similar to the axon radius R_a. Note that after non-dimensionalisation the axon radius is 1. To facilitate the asymptotic analysis we therefore write the vesicle radius as $R_v = 1 + \varepsilon^2$ and consider V_ε as the solution of the system (20), (17). Numerical solutions of this system for fixed parameters F and ξ illustrate that the predicted vesicle speed decreases with larger vesicle diameter (Figure 9) in line with experimental observations[11].

Figure 9: The velocity decreases as the cargo diameter increases. Compared to the reference in red, the blue and green graphs correspond to fixed ξ with different values of force F, while the pink and red graphs represent fixed force F with various values for (scaled) drag friction ξ.

For the asymptotic analysis we start with the interface points where the axon wall detaches from the vesicle. In the modified notation the interface point in front of the vesicle is at

$$B = \sqrt{R_v^2 - 1} = \varepsilon \sqrt{2 + \varepsilon^2} \ . \tag{24}$$

The interface point at the rear of the vesicle, on the other hand, is determined by equation (17) which relates it to the velocity V. Using the modified notation, this equation reads

$$V = \frac{1}{A} \left(\sqrt{(1 + \varepsilon^2)^2 - A^2} - (1 + \varepsilon^2)^2 \right) + A \ . \tag{25}$$

Using the ansatz $A \sim A_1 \varepsilon + A_2 \varepsilon^2 + ...$ we find that the expansion up to order 5 in ε is given by

$$A \sim -\frac{1}{V} \varepsilon^2 + \frac{1 - 2 V^2}{2 V^3} \varepsilon^4 + O(\varepsilon^6) \tag{26}$$

Note that this expansion is not valid in regimes where V is small as it yields un-physically small or large values of the position A where the vesicle detaches from the axon.

We now turn to equation (20) to compute an approximate solution for the vesicle transport velocity. We couple this equation to the expansion (26) and use the ansatz $V \sim V_0 + V_1\varepsilon + V_2\varepsilon^2 + V_3\varepsilon^3 + ...$ Equating coefficients of equal powers of ε yields the asymptotic solution (detailed in supplementary section A)

$$V \sim \frac{F}{\xi} - \frac{2\sqrt{2}F\varepsilon^3}{3\xi^2} - \frac{\varepsilon^4}{2\xi} + O\left(\varepsilon^5\right) . \tag{27}$$

In the same way as the expansion of A in (26), the expansion of V is not valid when ξ is small in which case the expansion would predict negative velocity. In figure 10 we compare the numerical solution for the vesicle velocity to the asymptotic approximation given by the leading order terms up to order 3 in (27).

Figure 10: Asymptotics for cargo velocity as a function of cargo size for $\xi = 1$, $F = 2$.

Finally we rewrite the asymptotic solution (27) up to third order in ε, $\tilde{V} \approx \frac{\tilde{F}}{\tilde{\xi}}\left(1 - \frac{2\sqrt{2}\varepsilon^3}{3\tilde{\xi}}\right)$, in dimensional form substituting the definitions of the dimensionless independent and dependent variables, $\tilde{V} = V\frac{\chi}{R_a \kappa}$, $\tilde{F} = F\frac{\Delta y}{R_a^2 \kappa}$, $\tilde{\xi} = \xi\frac{\Delta y}{R_a \chi}$ and $\varepsilon^2 = \tilde{R}_v - 1 = \frac{R_v}{R_a} - 1$ from section 3. We find that

$$V \approx \frac{F}{\xi}\left(1 - \frac{2\sqrt{2}}{3}\left(\frac{R_v}{R_a} - 1\right)^{3/2}\frac{R_a}{\Delta y}\frac{\chi}{\xi}\right), \tag{28}$$

which quantifies the decrease of transport speed as a function of vesicle radius R_v in the regime when the vesicle radius is close to the axon radius and drag friction is not too small.

6 Discussion

In this study, we establish a mathematical model for the interaction of the membrane associated periodic skeleton (MPS) in axons with large cargo vesicles undergoing axonal transport. The model accounts for transient swelling caused by the excess size of the cargo and for the mechanical resistance this imposes on its motion. The resulting model is a free boundary problem and resembles the classical obstacle problem.

We consider quasi-stationary solutions and reformulate them as force-velocity relation for the transport speed of the cargo. It predicts that when neglecting drag, the marginal force - the slope of the force-velocity relation - decays by a factor of 2 starting from when the vesicle is at rest, which can be tested experimentally.

We also compute - numerically and asymptotically as closed-form expression - the size dependence of the transport speed. It predicts that the velocity decays for larger vesicles

which is in line with recent observations [11]. It also predicts that the transport speed as a function of vesicle size follows a power law with exponent 3/2 which is amenable to experimental verification.

In its present form, the model is designed to be a foundation for future computational studies of axonal transport through stochastic simulation of molecular motors dynein and kinesin involved in cargo transport. Some additional aspects of the mechanics involved in the axonal transport of large vesicles will most likely be of interest and should therefore be incorporated into the model. One is friction between the axon wall and vesicles taking into account both normal forces and shear forces which has already been suggested as a potential explanation of the size-dependence of the vesicle speed [11]. Such friction could be either caused by direct interaction or mediated by cytoplasm viscosity [18]. Another striking aspect of axonal mechanics is the coupling between neighbouring actin rings which is mediated by spectrin and sustains axial tension [22], therefore coupling radial and axial tension [23]. Incorporating spectrin into the present model would most likely introduce 2nd order derivatives into the continuum model (13) and lead to different free boundary conditions.

Acknowledgements

The authors acknowledge funding from the Australian Research Council (ARC) Discovery Program (grant number DP180102956), awarded to D. B. O. An RTP scholarship funded by the University of Queensland (UQ), awarded to N. R.

Conflict of Interest

The authors declare that there is no conflict of interest.

A Asymptotic expansion of transport velocity

Starting with the first equation in (19), we have the following integral equation

$$V = \frac{F}{\xi} + \frac{1}{\xi} \int_A^B \left(h(y) - 1 - V\, h'(y) \right) h'(y) dy \ .$$

Now we substitute h and its derivative given in (4) obtaining

$$0 = V - \frac{F}{\xi} - \frac{1}{\xi} \int_A^B -\frac{y \left(\frac{yV}{\sqrt{R_v^2 - y^2}} + \sqrt{R_v^2 - y^2} - 1 \right)}{\sqrt{R_v^2 - y^2}} dy \ .$$

Coupling this to the asymptotic expansions of A and B given in (26) and (24) we conclude that

$$0 = V - \frac{F}{\xi} - \frac{1}{\xi} \left[\frac{1}{2}V \left(\varepsilon^2 + 1 \right) \log \left(\frac{-y + \varepsilon^2 + 1}{y + \varepsilon^2 + 1} \right) + Vy - \sqrt{\left(\varepsilon^2 + 1 \right)^2 - y^2} - y^2 \right]_A^B , \quad (29)$$

where we used $R_v = 1 + \varepsilon^2$. Finally, substitute the asymptotic expansion for V into (29),

$$\left(V_0 - \frac{F}{\xi} \right) + V_1 \varepsilon + V_2 \varepsilon^2 + \varepsilon^3 \left(\frac{2\sqrt{2}V_m}{3\xi} + V_3 \right) + \frac{\varepsilon^4 \left(6\xi V_4 + 4\sqrt{2}V_1 + 3 \right)}{6\xi}$$

$$+ \varepsilon^5 \left(-\frac{V_m}{15\sqrt{2}\xi} + \frac{2\sqrt{2}V_2}{3\xi} + V_5 \right) + O(\varepsilon)^6 = 0 \ .$$

Equating coefficients of equal powers of ε we obtain

$$V_0 = \frac{F}{\xi}, \ V_1 = 0, \ V_2 = 0, \ V_3 = -\frac{2\sqrt{2}F}{3\xi^2}, \ V_4 = -\frac{1}{2\xi} \text{ and } V_5 = \frac{F}{15\sqrt{2}\xi^2} \ .$$

B   Numerical test of quasi-steady state approximation

In this supplementary section we numerically test the validity of the quasi-steady state approximation (20). To this end we solve the time discrete model (5)-(6) numerically, extract the cargo velocity and compare it to the result obtained from numerically solving (20). The numerical results coincide (figure 11).

Figure 11: The velocity obtained through solving the discrete system (5)-(6) coincides with the velocity computed numerically using the quasi-steady state approximation of the continuum model (20) for $\xi = 1$, $F = 2$.

References

[1] P. Guedes-Dias and E. L. F. Holzbaur, "Axonal transport: Driving synaptic function," *Science (American Association for the Advancement of Science)*, vol. 366, no. 6462, p. 199, 2019.

[2] P. D. Chowdary, D. L. Che, K. Zhang, and B. Cui, "Retrograde ngf axonal transport—motor coordination in the unidirectional motility regime," *Biophysical journal*, vol. 108, no. 11, pp. 2691–2703, 2015.

[3] K. Xu, G. Zhong, and X. Zhuang, "Actin, spectrin, and associated proteins form a periodic cytoskeletal structure in axons," *Science (American Association for the Advancement of Science)*, vol. 339, no. 6118, pp. 452–456, 2013.

[4] B. Han, R. Zhou, C. Xia, and X. Zhuang, "Structural organization of the actin-spectrin–based membrane skeleton in dendrites and soma of neurons," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 114, no. 32, pp. E6678–E6685, 2017, cited By 68.

[5] S. Vassilopoulos, S. Gibaud, A. Jimenez, G. Caillol, and C. Leterrier, "Ultrastructure of the axonal periodic scaffold reveals a braid-like organization of actin rings," *Nature Communications*, vol. 10, no. 1, 2019, cited By 41.

[6] X. Pan, Y. Zhou, P. Hotulainen, F. Meunier, and T. Wang, "The axonal radial contractility: Structural basis underlying a new form of neural plasticity," *BioEssays*, vol. 43, no. 8, 2021, cited By 0.

[7] R. Bar-Ziv, T. Tlusty, E. Moses, S. Safran, and A. Bershadsky, "Pearling in cells: A clue to understanding cell shape," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 96, no. 18, pp. 10 140–10 145, 1999, cited By 127.

[8] A. Prokop, "Cytoskeletal organization of axons in vertebrates and invertebrates," *Journal of Cell Biology*, vol. 219, no. 7, 05 2020.

[9] A. Datar, J. Ameeramja, A. Bhat, R. Srivastava, A. Mishra, R. Bernal, J. Prost, A. Callan-Jones, and P. Pullarkat, "The roles of microtubules and membrane tension in axonal beading, retraction, and atrophy," *Biophysical Journal*, vol. 117, no. 5, pp. 880–891, 2019, cited By 17.

[10] S. Leite, P. Sampaio, V. Sousa, J. Nogueira-Rodrigues, R. Pinto-Costa, L. Peters, P. Brites, and M. Sousa, "The actin-binding protein α-adducin is required for maintaining axon diameter," *Cell Reports*, vol. 15, no. 3, pp. 490–498, 2016.

[11] T. Wang, W. Li, S. Martin, A. Papadopulos, M. Joensuu, C. Liu, A. Jiang, G. Shamsollahi, R. Amor, V. Lanoue, P. Padmanabhan, and F. A. Meunier, "Radial contractility of actomyosin rings facilitates axonal trafficking and structural stability," *The Journal of cell biology*, vol. 219, no. 5, 2020.

[12] M. Hammarlund, E. Jorgensen, and M. Bastiani, "Axons break in animals lacking β-spectrin," *Journal of Cell Biology*, vol. 176, no. 3, pp. 269–275, 2007, cited By 155.

[13] D. Liewald, R. Miller, N. Logothetis, H.-J. Wagner, and A. Schüz, "Distribution of axon diameters in cortical white matter: an electron-microscopic study on three human brains and a macaque," *Biological Cybernetics*, vol. 108, no. 5, pp. 541–557, 2014, cited By 151.

[14] H. M. McBride, M. Neuspiel, and S. Wasiak, "Mitochondria: More than just a power-house," *Current biology*, vol. 16, no. 14, pp. R551–R560, 2006.

[15] N. Mizushima, Y. Ohsumi, and T. Yoshimori, "Autophagosome formation in mammalian cells," *Cell structure and function*, vol. 27, no. 6, pp. 421–429, 2002.

[16] A. L. Altick, L. M. Baryshnikova, T. Q. Vu, and C. S. von Bartheld, "Quantitative analysis of multivesicular bodies (mvbs) in the hypoglossal nerve: Evidence that neurotrophic factors do not use mvbs for retrograde axonal transport," *Journal of comparative neurology (1911)*, vol. 514, no. 6, pp. 641–657, 2009.

[17] S. Roy, "Seeing the unseen: The hidden world of slow axonal transport," *The Neuroscientist (Baltimore, Md.)*, vol. 20, no. 1, pp. 71–81, 2014.

[18] J. Wortman, U. Shrestha, D. Barry, M. Garcia, S. Gross, and C. Yu, "Axonal transport: How high microtubule density can compensate for boundary effects in small-caliber axons," *Biophysical Journal*, vol. 106, no. 4, pp. 813–823, 2014, cited By 21.

[19] D. Danielli, "An overview of the obstacle problem," *Notices of the American Mathematical Society*, vol. 67, no. 10, pp. 1487–1497, 2020.

[20] M. E. Fisher and A. B. Kolomeisky, "The force exerted by a molecular motor," *Proceedings of the National Academy of Sciences - PNAS*, vol. 96, no. 12, pp. 6597–6602, 1999.

[21] A. Kunwar, S. K. Tripathy, J. Xu, M. K. Mattson, P. Anand, R. Sigua, M. Vershinin, R. J. McKenney, C. C. Yu, A. Mogilner, and S. P. Gross, "Mechanical stochastic tug-of-war models cannot explain bidirectional lipid-droplet transport," *Proceedings of the National Academy of Sciences - PNAS*, vol. 108, no. 47, pp. 18 960–18 965, 2011.

[22] N. Unsain, F. D. Stefani, and A. Cáceres, "The actin/spectrin membrane-associated periodic skeleton in neurons," *Frontiers in Synaptic Neuroscience*, vol. 10, 2018.

[23] A. Fan, A. Tofangchi, M. Kandel, G. Popescu, and T. Saif, "Coupled circumferential and axial tension driven by actin and myosin influences in vivo axon diameter," *Sci Rep*, vol. 7, no. 1, p. 14188, 10 2017.