# Hyperparasitism and the evolution of parasite virulence

Jason Wood[1,*] and Ben Ashby[1,2,3]

1. Department of Mathematical Sciences, University of Bath.

2. Department of Mathematics, Simon Fraser University.

3. The Pacific Institute on Pathogens, Pandemics and Society (PIPPS), Simon Fraser University,

Burnaby, BC

**Keywords:** hyperparasitism, parasitism, virulence, evolution, biocontrol

**Author contributions:** JW and BA designed the study together and contributed equally to the writing

of the manuscript. JW carried out the analysis.

*Corresponding author: jrw78@bath.ac.uk

ABSTRACT

Hyperparasites (species which parasitise other parasites) are common in natural populations,

affecting many parasitic taxa, including: eukaryotic parasites; bacterial and fungal pathogens.

Hyperparasitism is therefore likely to shape the ecology and evolution of many host-parasite

systems, representing a promising method for biocontrol (e.g., treating antimicrobial resistant

infections). However, the eco-evolutionary consequences of hyperparasitism have received little

attention. We use a host-parasite-hyperparasite model to explore how introducing a hyperparasite

drives the evolution of parasite virulence, and what impact this has on the host population. We

show when the introduction of a hyperparasite selects for higher or lower parasite virulence, and

the changes in virulence experienced by the host population. Crucially, we show that variation in the

direct effects of hyperparasites on virulence and transmission, and the probability of co-

transmission, can lead to a previously unseen hysteresis effect, whereby small shifts in hyperparasite

characteristics can lead to sudden shifts in parasite virulence. We also show that hyperparasites can

induce diversification in parasite virulence, leading to the coexistence of high and low virulence

strains. Our results show hyperparasites can have dramatic effects on the evolution of parasite

virulence, and that the use of hyperparasites in biocontrol should be approached with caution.

 INTRODUCTION

Hyperparasitism, wherein parasitic organisms are themselves parasitised by another species, is

ubiquitous in the natural world. Hyperparasitism has been observed across many taxa, including

bacterial (Greer 2005; Ashelford, Day, and Fry 2003) and fungal (Mikhailov, Simdyanov, and Aleoshin

2016) pathogens, parasitic worms (Mohan et al. 2020) and other eukaryotic parasites (Wendling et

al. 2017). For example, many human bacterial pathogens are parasitised by bacteriophages,

including *E. Coli* (e.g., phage O157:H7 (Munns et al. 2015)); *Salmonella typhimurium* (e.g., phage

$\phi$AB2 (Berchieri, Lovell, and Barrow 1991)); and *Klebsiella pneumoniae* (e.g., phage B5055 (Chhibber,

Kaur, and Kumari 2008)). Hyperparasitism is also common among fungal plant pathogens, such as

the powdery mildew fungus *Podosphaera plantaginis* with the hyperparasitic fungus *Ampelomyces*

spp. (Parratt and Laine 2018), or the chestnut blight fungus *Cryphonectria parasitica* (Nuss 2005)

which can be infected by *Cryphonecria hypovirus 1* (Parratt and Laine 2016).

Hyperparasites are understood to play a major role in the ecology of parasites, and have been

shown to influence both the early epidemic dynamics and the overwintering success (Parratt and

Laine 2018) of plant pathogens. They have also been implicated as a driver of seasonal epidemics of

cholera (Faruque et al. 2005). In addition to ecological effects on the parasite, hyperparasites can

have significant effects on virulence, either inducing hypovirulence (a reduction in the virulence

experienced by the host, e.g., by reducing the population size of the parasite) or hypervirulence (an

increase in the virulence experienced by the host, e.g., by causing the parasite to release toxins or by

introducing virulence factors). Hyperparasite-induced changes in virulence have been observed in

many bacterial pathogens. For example, a phage protein enhances motility (and hence virulence) of

*E. coli* (Kakkanat et al. 2017); the temperate phage PHB09 reduces the virulence of *Bordetella*

*bronchiseptica* both *in vivo* and *in vitro* (Chen et al. 2020); the phage CTX$\phi$ encodes the cholera toxin

within *Vibrio cholerae* (Waldor and Mekalanos 1996); and the phage $\phi$CDHM1 interferes with

quorum sensing in *Clostridium difficile* (Hargreaves, Kropinski, and Clokie 2014).

Due to their ecological and virulence-mediating effects, hyperparasites long have been considered as

possible sources of biological control (biocontrol) for many infectious diseases (Holtappels et al.

2021; Obradovic et al. 2004), including in the agricultural and food industries and in the treatment of

chronic or antimicrobial resistant infections in humans (Gordillo Altamirano and Barr 2019). Indeed,

phage therapy has long been used as an alternative to antibiotics in some countries (Schooley et al.

2017; Ferriol-González and Domingo-Calap 2021). However, little is known about the evolutionary

implications of hyperparasitism for parasites, and so their use as agents of biocontrol in novel

settings could lead to unexpected outcomes for important traits such as parasite transmission and

virulence. For instance, (Prospero et al. 2021) speculated that the introduction of hyperparasites

which reduce parasite virulence (hypovirulence) might select for higher virulence in chestnut blight

fungus, which has been observed experimentally (Bryner and Rigling 2012).

Understanding the interplay of host, parasite and hyperparasite ecological and evolutionary

dynamics is crucial not only to the development of novel agents of biocontrol, but also for

understanding their role in natural ecosystems. Theoretical studies of hyperparasites are rare and

have in the past mainly focussed on their ecological consequences (Taylor et al. 1998; A. Yu.

Morozov, Robin, and Franc 2007). Yet recently there has been renewed theoretical interest in

hyperparasitism in an evolutionary context, in the form of theoretical models of hyperparasite

evolution (Northrup et al. 2021) and parasite-hyperparasite coevolution (Sandhu et al. 2021). Two

findings are of particular interest. First, Sandhu et al. (2021), who considered parasite-hyperparasite

coevolution, observed that the introduction of hyperparasites always increases parasite virulence,

but in almost all scenarios decreases average host mortality. Second, Northrup et al. (2021)

observed that when hyperparasites are more readily co-transmitted with evolutionarily static

parasites, this selects for less harm by the hyperparasite due to an increased link between its fitness

and parasite transmission (similar to virulence evolution in vertically transmitted parasites). These

findings highlight the importance of understanding hyperparasitism in both an ecological and an

evolutionary context.

Here, we further investigate the eco-evolutionary consequences of introducing a hyperparasite for

parasite virulence and the resulting net impact on the host population. We show how the

introduction of hyperparasites causes changes in parasite virulence and whether this leads to a net

positive or negative effect on the host population depending on the relative transmission rate and

virulence of hyperparasitised parasites. We also show how small changes in the effects of the

hyperparasite on parasite transmission and virulence can lead to large shifts in the evolution of

virulence due to hysteresis. Finally, we show that the hyperparasite can induce diversification in

parasite virulence, leading to the coexistence of a relatively high and low virulence strains.

METHODS

*MODEL DESCRIPTION*

We consider a well-mixed population of asexual hosts, parasites, and hyperparasites, where $S$ is the

density of uninfected (susceptible) hosts, $I$ is the density of hosts only infected by the parasite

(parasitised) and $H$ is the density of hosts infected by both the parasite and the hyperparasite

(hyperparasitised). Hosts reproduce at a baseline per-capita rate $b$, subject to density-dependent

crowding $qN$, with $q > 0$ and $N = S + I + H$. Parasite transmission is density-dependent, with

parasitised and hyperparasitised hosts having parasite transmission rates of $\beta$ and $\eta\beta$ to susceptible

hosts, respectively, where $\beta$ is the baseline transmission rate, $\eta > 1$ implies "hypertransmission"

(the hyperparasite increases parasite transmissibility) and $\eta < 1$ implies "hypotransmission" (the

hyperparasite decreases parasite transmissibility). Hyperparasites are co-transmitted with parasites

to susceptible hosts with probability $\rho$. Hyperparasite infection of parasitised hosts is also density-

dependent, with transmission rate $\sigma$. All hosts experience a natural mortality rate $d$, with parasitised

and hyperparasitised hosts experiencing additional mortality due to disease at rates $\alpha$ and $\lambda\alpha$,

respectively. Thus, when $\lambda < 1$ the hyperparasite induces hypovirulence (the hyperparasite

decreases the disease-associated mortality rate) and when $\lambda > 1$ the hyperparasite induces

hypervirulence (the hyperparasite increases the disease-associated mortality rate). Both parasitised

and hyperparasitised hosts recover from infection at rate $\gamma$, with no lasting immunity (see Table 1

for a full summary of model parameters and their default values for the analysis).

The ecological dynamics for a monomorphic population are described by the following three

ordinary differential equations:

$$\dot{S} = (b - qN)N - (\beta I + \eta\beta H + d)S + \gamma(I + H)$$
$$\dot{I} = \big(\beta S - \sigma H - (d + \alpha + \gamma)\big)I + (1 - \rho)\eta\beta SH \qquad\qquad (1)$$
$$\dot{H} = \big(\rho\eta\beta S + \sigma I - (d + \lambda\alpha + \gamma)\big)H$$

| Parameter | Description | Default value |
|:---:|:---:|:---:|
| $b$ | Natural birth rate of hosts | 2.0 |
| $c$ | Curvature parameter for transmission-virulence trade-off | $-60$ |
| $d$ | Natural mortality rate of hosts | 0.5 |
| $q$ | Strength of density dependent competition | 0.1 |
| $\alpha$ | Parasite virulence | n/a |

| | | |
|---|---|---|
| $\alpha_{max}$ | Maximum parasite virulence | 5.0 |
| $\beta$ | Parasite transmission rate | n/a |
| $\beta_0$ | Scale factor in virulence-transmission trade-off | $\sqrt{5}$ |
| $\beta_s$ | Non-linear transmission scale factor | 0.5 |
| $\beta_l$ | Linear transmission scale factor | 1.0 |
| $\gamma$ | Host recovery rate | 0.5 |
| $\eta$ | Hyperparasite transmission modifier | n/a |
| $\lambda$ | Hyperparasite virulence modifier | n/a |
| $\rho$ | Probability of hyperparasite co-transmission | 0.5 |
| $\sigma$ | Hyperparasite transmission rate | 4.0 |

123
*Table 1: Model parameters and default values*

124 We explore the evolution of parasite virulence under a standard transmission-virulence trade-off

125 with diminishing returns, such that $\beta = \beta(\alpha)$, $\beta'(\alpha) > 0$ and $\beta''(\alpha) < 0$. We primarily consider the

126 following simple trade-off between virulence and transmission,

$$\beta(\alpha) = \beta_{max}\sqrt{\frac{\alpha}{\alpha_{max}}} = \beta_0\sqrt{\alpha} \qquad (2)$$

127 where virulence increases quadratically with the transmission rate of the parasite. To check the

128 generality of our results we consider a second trade-off,

$$\beta(\alpha) = \beta_s\frac{1 - e^{c\left(\frac{\alpha}{\alpha_{max}}\right)}}{1 - e^c} + \beta_l\alpha \qquad (3)$$

which allows us to explore trade-offs with different curvatures.

We assume that mutations are sufficiently rare so that the ecological dynamics in Equation 1 reach a

stable endemic equilibrium (with either the host and parasite, or all three species present; see

*Supplementary materials*) before a new mutant arises, and that mutations have small phenotypic

effects. Note that since the endemic equilibrium of Equation 1 is analytically intractable, in practice

we find the endemic equilibrium and verify its stability numerically. The invasion dynamics of a rare

mutant parasite with transmission rate $\beta_m$ and virulence $\alpha_m$ are then,

$$\dot{I_m} = \left(\beta_m \hat{S} - \sigma \hat{H} - (d + \alpha_m + \gamma)\right) I_m + (1 - \rho)\eta \beta_m \hat{S} H_m,$$

$$\dot{H_m} = \left(\rho \eta \beta_m \hat{S} - (d + \lambda \alpha_m + \gamma)\right) H_m + \sigma I_m \hat{H}.$$

(4)

138      where $\hat{K}$ indicates the equilibrium density for population $K \in \{S, H\}$. The invasion fitness $w(\alpha_m)$ is

139      given by the largest eigenvalue of the next generation matrix (Hurford, Cownden, and Day 2010)

140      (see *Supplementary materials*),

$$\begin{bmatrix} \dfrac{\beta_m \hat{S}}{\sigma \hat{H} + d + \alpha_m + \gamma} + \dfrac{(1 - \rho)\eta \sigma \beta_m \hat{S} \hat{H}}{(\sigma \hat{H} + d + \alpha_m + \gamma)(d + \lambda \alpha_m + \gamma)} & \dfrac{(1 - \rho)\eta \beta_m \hat{S}}{d + \lambda \alpha_m + \gamma} \\ \dfrac{\rho \eta \sigma \beta_m \hat{S} \hat{H}}{(\sigma \hat{H} + d + \alpha_m + \gamma)(d + \lambda \alpha_m + \gamma)} & \dfrac{\rho \eta \beta_m \hat{S}}{d + \lambda \alpha_m + \gamma} \end{bmatrix}$$

which we omit here for the sake of brevity. We derive the fitness gradient $F(\alpha) = \left.\dfrac{\partial w}{\partial \alpha_m}\right|_{\alpha_m = \alpha}$ and

hence calculate singular strategies, $\alpha^*$, that satisfy $F(\alpha^*) = 0$. To distinguish between direct ($\lambda$) and

evolutionary effects of the hyperparasite on virulence we refer to $\alpha^*$ as intrinsic virulence. We

determine the evolutionary stability of a singular strategy by the sign of $E(\alpha^*) = \left.\dfrac{\partial^2 w}{\partial \alpha_m^2}\right|_{\alpha_m = \alpha = \alpha^*}$

(negative values indicate evolutionary stability), and convergence stability by numerically

approximating the derivative $M(\alpha^*) = \left.\dfrac{\partial^2 w}{\partial \alpha_m \partial \alpha}\right|_{\alpha_m = \alpha = \alpha^*}$ and checking if the inequality $E(\alpha^*) +$

$M(\alpha^*) < 0$ holds. If a singular strategy is both evolutionarily and convergence stable then it is a

continuously stable strategy (an evolutionary attractor); if it is convergence stable but not

evolutionarily stable then it is an evolutionary branching point; and if it is convergence unstable then

it is an evolutionary repeller.

 *SIMULATIONS*

We complement our numerical analysis with simulations of the evolutionary dynamics of our

system, which relax the adaptive dynamics assumptions of continuous traits and separate ecological

and evolutionary timescales. To perform our simulations, we first create a discretised trait space for

the parasite. We then initialise the hyperparasite-free system at its eco-evolutionary attractor, and

introduce hyperparasites at an arbitrarily low population density. We use a fourth order Runga-Kutta

method to solve the ordinary differential equations over a long time period, stopping when the

populations have relatively small changes in size, or the time threshold has been reached.

After removing phenotypes that fall below an arbitrary threshold, we choose one of the extant

parasite phenotypes (using a weighted probability based on parasite density) and introduce a rare

mutant a small phenotypic distance away at a low frequency. We then use this new population as

the initial condition for our ordinary differential equation system, which we again evaluate using a

fourth order Runga-Kutta method. All code used to produce the figures within this paper is available

within the Supplementary Material and on GitHub (GitHub – JasonRWood/Wood_Ashby_2023).

*MEASURING THE IMPACT OF THE HYPERPARASITE ON THE HOST POPULATION*

We assume that the hyperparasite is introduced into a well-established host-parasite system, such

that the parasite is initially at its continuously stable strategy $\alpha_0$ (see *Supplementary materials*) and

the system is at equilibrium. In addition to exploring the evolutionary implications for parasite

virulence we also consider the ecological implications for the host. Specifically, we consider two

metrics to encapsulate the impact on the host population following the introduction of the

hyperparasite, namely: the impact on the host population size $\Delta_N(\alpha^*) = \frac{\widehat{N}(\alpha^*)}{N_0}$, and the impact on

the average mortality rate of infected individuals $\Delta_M(\alpha^*) = \frac{\alpha^*(\hat{I}(\alpha^*) + \lambda\widehat{H}(\alpha^*))}{\alpha_0(\hat{I}(\alpha^*) + \widehat{H}(\alpha^*))}$, where $N_0$ corresponds

to the steady state of the host population before the introduction of the hyperparasite (with

ancestral virulence $\alpha_0$), and $\widehat{N}(\alpha^*), \widehat{I}(\alpha^*)$ and $\widehat{H}(\alpha^*)$ are the steady states following the introduction

of the hyperparasite evaluated at a continuously stable strategy, $\alpha^*$. For brevity, we refer to $\Delta_M(\alpha^*)$

as the relative average virulence. When $\Delta_M(\alpha^*) > 1$ the hyperparasite leads to an increase in the

average virulence experienced by infected hosts, whereas when $\Delta_M(\alpha^*) < 1$ the average virulence

experienced by infected hosts decreases.

## RESULTS

*HYPERPARASITE-INDUCED SHIFTS IN VIRULENCE EVOLUTION*

Following the introduction of the hyperparasite, we see pronounced shifts in the evolution of

virulence depending on the extent to which hyperparasites modify parasite transmission ($\eta$) and/or

virulence ($\lambda$), and the extent to which co-transmission of the hyperparasite ($\rho$) occurs. When the

hyperparasite induces hypovirulence (a direct reduction in the disease-associated mortality rate, $\lambda <$

1), the parasite always experiences selection for higher intrinsic virulence (Fig. 1A). However, when

the hyperparasite either has no direct effect on virulence (when $\lambda = 1$; Fig. 1B) or induces

hypervirulence (when $\lambda > 1$; Fig. 1C) this is no longer the case. When the hyperparasite strongly

reduces parasite transmission ($\eta \ll 1$), the parasite evolves higher intrinsic virulence, and when the

hyperparasite increases transmission ($\eta > 1$), intrinsic virulence either remains virtually unchanged

($\lambda = 1$; Fig. 1B) or evolves to lower levels ($\lambda > 1$; Fig. 1C). Yet at intermediate values of $\eta < 1$

(hypotransmission), the behaviour of the system becomes more complicated, as there is a bistable

region which becomes more prominent as the virulence modifier ($\lambda$) or probability of co-

transmission ($\rho$) increase (Fig. 1B-C).

The bistable region means that the system has two evolutionary attractors, and therefore small

changes in the underlying ecology (e.g., direct effects of the hyperparasite on parasite virulence or

transmission) may cause sudden shifts between high and low virulence states (Fig. 1A-C). Crucially,

these shifts in virulence may be difficult to reverse as the switching points towards higher or lower

intrinsic virulence are not the same (i.e., there is a "hysteresis effect"; Fig. 2). For example, suppose

the hyperparasite causes a hypervirulence ($\lambda > 1$; Fig. 1C), co-transmits with high probability ($\rho \approx$

1), and reduces the transmissibility of the parasite ($\eta = 0.5$). Selection would then favour a

reduction in intrinsic virulence relative to the absence of the hyperparasite ($\alpha^* < \alpha_0$) (Fig. 2). A

small reduction in the transmission modifier below a critical threshold, $\eta_c^1$, would suddenly shift

selection for increased intrinsic virulence ($\alpha^* > \alpha_0$), but a reversion in the transmission modifier to

its initial value ($\eta > \eta_c^1$) would not lead to a drop in intrinsic virulence until a second critical

threshold, $\eta > \eta_c^2 > \eta_c^1$ is breached (Fig. 2). Thus, relatively small changes in the effects of the

hyperparasite on parasite transmission can lead to large changes in the evolution of virulence that

may be difficult to reverse.

The key driver of the bistable region (and hence the hysteresis effect) is hyperparasite prevalence.

The upper virulence branch (Fig. 1A-C) corresponds to relatively low hyperparasite prevalence (Fig.

1D-F), whereas the lower branch corresponds to relatively high prevalence. While there is a

numerical feedback between parasite virulence and hyperparasite prevalence, one can understand

the link between the two (and hence the hysteresis effect) when viewed through the lens of life-

history strategies. Specifically, higher intrinsic virulence is an adaptation to shift reproductive output

(transmission) before the parasite is likely to be infected. For example, when the hyperparasite fully

prevents parasite transmission ($\eta = 0$), hyperparasitised hosts make no contribution to parasite

fitness and so high intrinsic virulence and transmissibility is advantageous because all transmission

must occur before the parasite is infected. Similarly, when the hyperparasite significantly reduces

but does not prevent parasite transmission ($0 < \eta \ll 1$), most onward transmission occurs before

being hyperparasitised. Yet this strategy of shifting reproductive output (transmission) earlier in the

parasite's typical lifespan is only advantageous provided hyperparasite prevalence is relatively low

so that sufficient transmission can occur before encountering the hyperparasite.

If the hyperparasite is sufficiently common (prevalence increases with the transmission modifier ($\eta$)

and probability of co-transmission ($\rho$)), then the parasite does not reap the benefits of high

transmission before being hyperparasitised. Thus, intrinsic virulence shifts to the lower branch of the

bistable region where most parasites are hyperparasitised and there is little change in average

virulence ($\Delta_M(\alpha^*) \approx \frac{\lambda \alpha^*}{\alpha_0} \approx 1$). The bistable region therefore exists due to a positive feedback loop

between hyperparasite prevalence and parasite virulence (e.g., higher initial hyperparasite

prevalence selects for lower intrinsic virulence, which in turn increases hyperparasite prevalence).

Note that the key driver of intrinsic virulence along the lower branch appears to be the direct effects

of the hyperparasite on virulence ($\lambda$). Intuitively, if most parasites are hyperparasitised, then $\eta$

(hypo/hyper- transmission) and $\rho$ (co-transmission) are close to simple scaling factors for parasite

fitness, whereas $\lambda$ (hypo/hyper-virulence) affects the parasite's infectious period and so is the key

driver of selection along this branch.

*Figure 1: Evolutionary consequences for parasite virulence (A-C) (solid: evolutionary attractors, dashed: evolutionary*

*repellers),the prevalence of the hyperparasite (D-E), relative average mortality (G-I) and relative population size (J-L), as*

*the direct effects of the hyperparasite on parasite transmission (hypotransmission: $\eta < 1$; hypertransmission: $\eta > 1$) and*

*virulence ($\lambda$) vary. Evolutionary attractors are both convergence stable and evolutionarily stable, and are therefore*

*continuously stable strategies (CSSs). Left hand column - hypovirulence ($\lambda < 1$); Central column - no effect on virulence*

*($\lambda = 1$); Right hand column - hypervirulence ($\lambda > 1$). All panels contain three sets of curves showing the evolutionary*

*endpoint as the probability of hyperparasite co-transmission varies: $\rho = 0.1$ (blue), $\rho = 0.5$ (orange), $\rho = 0.9$ (green). The*

*start and end of each continuous set of evolutionary attractors are shown with empty and filled shapes. The black line*

*indicates the ancestral state prior to the introduction of the hyperparasite.*

*Figure 2: Evolutionary simulations showing the hysteresis effect observed in Fig. 1 due to variation in the direct effects of*

*the hyperparasite on parasite transmission, $\eta$. A relatively small shift (blue) from an initial value of $\eta$ (black) leads to a sharp*

*increase in intrinsic virulence. However, a reversion to the initial value of $\eta$ (black) does not lead to a return to lower*

*intrinsic virulence. Instead, intrinsic virulence does not change significantly until a critical threshold is reached (red).*

*Parameters as in Table 1 except $\lambda = 2.0, \rho = 0.9$. Included inset figure replicates Fig. 1C when $\rho = 0.9$, coloured dots*

*indicated predicted $\alpha$ for respective value of $\eta$.*

Although the parasite may evolve higher or lower intrinsic virulence following the introduction of

the hyperparasite, whether this has a net positive or negative effect for the host population depends

on not only the new level of virulence, but also the effects on parasite prevalence, the prevalence of

hyperparasites, and the effects of hyperparasitism on virulence. We therefore consider the

ecological consequences of the hyperparasite on the host following evolutionary shifts in parasite

virulence.

Intuitively, when the hyperparasite selects for higher intrinsic virulence ($\alpha^* > \alpha_0$) while not

inducing hypovirulence ($\lambda \geq 1$), relative average virulence is always higher (Fig. 1H-I), and the

resulting host population size is lower (Fig. 1K-L). This is especially pronounced when

hyperparasitism strongly reduces transmission ($\eta \ll 1$). However, the net impact on hosts is less

straightforward when the hyperparasite induces hypovirulence while selecting for higher intrinsic

virulence (Fig. 1G, J), or induces hypervirulence while selecting for lower intrinsic virulence (Fig. 1I,

L). For example, the latter scenario can lead to little effect on the mortality rate of infected hosts

(Fig. 1I) but a marked increase in the host population size (Fig. 1L).

Crucially, the bistability of intrinsic virulence has a striking effect on the host population, with

relatively small shifts in the direct effects of hyperparasites on transmission ($\eta$) or on the probability

of co-transmission ($\rho$) causing substantial changes in relative average virulence (Fig. 1H-I) and the

host population size (Fig. 1K-L). Typically, infected hosts either experience a significant increase in

relative average virulence (for sufficiently low values of $\eta$), or there is little net effect. This suggests

that the introduction of a hyperparasite rarely improves the outcome of infection on average, even if

it is beneficial at the population level by reducing the proportion of the population that is infected.

We also see that as the probability of coinfections ($\rho$) increases, the effects of introducing the

hyperparasites on the relative average virulence and host population size typically grow stronger.

Note that while relative average virulence is close to one for sufficiently large values of $\eta$ in Fig. 1G-I,

it does not tend to one (in fact, all the curves pass through 1 before the point $\eta = 2$; see also Fig. S2,

S5).

*HYPERPARASITE-INDUCED DIVERSIFICATION IN PARASITE VIRULENCE*

In the absence of the hyperparasite, parasite virulence always evolves to a continuously stable

strategy due to the trade-off with transmission, which results in diminishing returns for the parasite.

The introduction of the hyperparasite either shifts the continuously stable strategy to a new value

(as in Fig. 1) or causes the parasite population to diversify through evolutionary branching into two

strains, one with relatively high intrinsic virulence which is less hyperparasitised (green branch in Fig.

3) and the other with relatively low intrinsic virulence which is more hyperparasitised (orange

branch in Fig. 3). Evolutionary branching occurs when a singular strategy is convergence stable, but

is not evolutionarily stable. This does not occur for the quadratic trade-off in Equation 2 as the

curvature of the trade-off is too strong. We therefore explored an alternative trade-off (Equation 3)

where the curvature is weaker. Note that the second trade-off produces qualitatively similar results

to those discussed above (Fig. S2).

The more virulent "short-lived" strain essentially prioritises earlier reproductive output

(transmission), infecting as many hosts as possible before being itself infected by the hyperparasite

and dying soon after. The less virulent "long-lived" strain instead prioritises transmission while

hyperparasitised. The contrast in strategies can be seen in Fig. 3B-C where the hyperparasite is more

commonly associated with the less virulent strain. In general, branching is most likely to occur when

the hyperparasite induces hypervirulence ($\lambda > 1$) and strong hypotransmission ($\eta \ll 1$), and the

probability of co-transmission is not too high ($\rho \ll 1$) (Fig. 4).

However, even though one of the parasite strains is intrinsically less virulent than the other, both are

more virulent than the ancestral strain in the absence of the hyperparasite (Fig. 3A). The more

virulent strain is rarely infected by the hyperparasite (Fig. 3D), as the host either dies prior to

encountering the hyperparasite, or experiences hypervirulence and so dies shortly after being

hyperparasitised. Despite the parasite branching into two distinct strategies that are both

intrinsically more virulent than the ancestral state, the overall impact on host and parasite

population sizes is minimal (Fig. 3B-D). Additionally, following the introduction of the hyperparasite,

there is initially a drastic reduction in host population density, with a further but much smaller

decrease in following branching.

*Figure 3. Evolutionary branching in parasite virulence following the introduction of the hyperparasite. Parasite virulence in*

*the absence of the hyperparasite is shown by the dashed line in (A). The parasite branches into relatively high (green) and*

*low (orange) virulence strains. (B-E) Densities of the host, parasite, hyperparasite and total populations: total (blue), high-*

*virulence strain (green), low-virulence strain (orange). The density of the host, parasite and total populations in the*

absence of the hyperparasite are plotted with a dashed black line. *Parameters as in Table 1 except* $d = 0.1, \sigma = 0.4, \rho =$

$0.5, \eta = 0.15, \lambda = 1.0, \gamma = 0.1.$

*Figure 4. Regions where evolutionary branching occurs (yellow) for the parasite following the introduction of the*

*hyperparasite, as the direct effects on transmission ($\eta$) and virulence ($\lambda$) vary. Parameters as in Table 1 except $d =$*

$0.1, \sigma = 0.4, \gamma = 0.1$.

DISCUSSION

Hyperparasites are abundant in nature and are promising sources of biological control in industry

(e.g., food production) and public health (e.g., phage therapy to combat antimicrobial resistance).

Yet the eco-evolutionary effects of hyperparasitism are poorly understood. In this study, we have

theoretically explored the impact of hyperparasites on the evolution of parasite virulence and the

consequences for the host population. We have shown that the introduction of hyperparasites can

lead to either higher or lower levels of intrinsic virulence depending on the direct effects of the

hyperparasitism. Crucially, we have shown that relatively small changes in the ecological effects of

hyperparasites can cause large shifts in both evolved and ecologically realised virulence, which can

be difficult to reverse due to hysteresis. Although a hyperparasite may cause selection for higher

intrinsic virulence, we have shown that this is not necessarily negative for the host population

overall, as the hyperparasite can suppress parasite prevalence and may mitigate virulence in

infected hosts. Finally, our model shows that the introduction of a hyperparasite can induce

diversification in the parasite population, with the potential for both strains to evolve to higher

levels of intrinsic virulence than in the absence of the hyperparasite. Overall, our results suggest that

the introduction of hyperparasites can have a strong impact on the evolution of parasite virulence

and that the nature of the outcome depends crucially on how the hyperparasite directly affects

parasite virulence and transmission.

Higher intrinsic virulence typically evolves when the hyperparasite dramatically reduces the

transmission rate of the parasite ($\eta \ll 1$) and/or the hyperparasite induces hypovirulence ($\lambda < 1$),

yet the reasons differ. Strong hypotransmission selects for higher intrinsic virulence ($\alpha^*$) because

hyperparasitism results in a significant loss in new infections, and so parasites experience selection

to infect as many hosts as possible prior to being hyperparasitised. In contrast, when the

hyperparasite induces hypovirulence ($\lambda < 1$), selection favours an increase in intrinsic virulence

because hyperparasitism effectively reduces the costs of higher virulence. This can be trivially

understood by considering a hyperparasite that completely mitigates virulence ($\lambda = 0$), which

greatly benefits the parasite by extending the infectious period. Conversely, when the hyperparasite

causes hypervirulence ($\lambda > 1$) it can select for lower intrinsic virulence provided there is not also

strong reduction in transmission. Note that we can also conclude that selection for higher intrinsic

virulence is driven by hyperparasite prevalence rather than being an adaptation to counteract the

direct effects of the hyperparasite on transmission ($\eta$), as higher intrinsic virulence evolves even

when the hyperparasite fully prevents onwards transmission ($\eta = 0$).

The hysteresis effect in our model suggests that not only might relatively small changes in

hyperparasite traits cause large evolutionary changes in the parasite, but that these might be

difficult to reverse. Hysteresis arises because for certain regions of parameter space the system is

bistable, meaning that the parasite can evolve to have relatively high or low intrinsic virulence

depending on the initial level of virulence. The bistability can be understood in terms of a positive

feedback loop between hyperparasite prevalence and parasite virulence, with high prevalence

selecting for low virulence, and vice versa. When the hyperparasite causes a significant reduction in

transmission ($\eta \ll 1$) it is very costly to the parasite and so selection favours higher intrinsic

virulence as this shifts most transmission to before being hyperparasitised (upper branch in Fig. 1A-

C). The hyperparasite is therefore at relatively low prevalence (Fig. 1D-F). As the effects of the

hyperparasite on parasite transmission weaken ($\eta$ increases), the hyperparasite may become more

common if it can co-transmit well with the parasite (high $\rho$), resulting in a gradual increase in

hyperparasite prevalence (Fig. 1D-F). Eventually a threshold is reached where the risk of

hyperparasitism is sufficiently high and the cost sufficiently low that selection favours a large

reduction in intrinsic virulence (lower branch in Fig. 1A-C), which feeds back to further increase

hyperparasite prevalence (Fig. 1D-F). This positive feedback, resulting in high hyperparasite

prevalence, is crucial to the hysteresis effect. If the hyperparasite once again reduces parasite

transmission more strongly ($\eta$ decreases), the system does not return to the high virulence state

because hyperparasite prevalence remains high (Fig. 1D-F). As $\eta$ continues to fall so too does

hyperparasite prevalence, until eventually the risk of hyperparasitism is sufficiently low and the cost

sufficiently high that selection favours a large increase in intrinsic virulence (upper branch in Fig. 1A-

C).

The hysteresis effect has especially important implications for biocontrol, as a relatively small change

in the hyperparasite can have a large effect on the evolution of virulence and on the host

population. It might be possible to reverse sudden shifts in virulence by removing the biocontrol for

a period of time, as selection would then push virulence towards the ancestral state, but the rate at

which this occurs will be system-specific. While hysteresis is well documented in the ecological

literature (e.g., the spruce budworm (Ludwig, Jones, and Holling 1978)), few examples exist in

evolutionary models (Prado et al. 2009). As an exception, Prado *et al.* 2009 observed hysteresis

when considering the evolution of host sociality and pathogen virulence within contact networks.

Prado *et al.* observed that the cycling behaviour they see within their system, has hysteresis-like

behaviour where selection for or against host sociality does not occur until the parasite passes

critical thresholds.

Our model reveals that the introduction of a hyperparasite can cause disruptive selection leading to

diversification into relatively high and low virulence parasite strains (although the "low" virulence

strain may still be more virulent than the ancestral strain, as in Fig. 3A). Branching typically requires

trade-offs that are close to linear (Bowers et al. 2005), such as the trade-off presented in Equation 3,

as strongly diminishing returns have a balancing effect on selection. Branching also typically requires

the hyperparasite to directly reduce the transmission rate of the parasite, which facilitates the

existence of high and low virulence phenotypes by creating distinct ecological niches. The high

virulence phenotype is rarely infected by the hyperparasite, favouring a "live fast, die young"

strategy, while the less virulent phenotype has a longer infectious period to mitigate the burden of

the hyperparasite.

Our study is closely related to previous theoretical explorations of evolution in host-parasite-

hyperparasite systems (Sandhu et al. 2021; Northrup et al. 2021). Sandhu et al (2021) also explored

the effects of hyperparasitism on the evolution of parasite virulence, but in contrast to our study

found that the introduction of a hyperparasite always selects for increased virulence and generally

reduces the average mortality rate of the host. Furthermore, Sandhu et al (2021) did not observe a

hysteresis effect, nor did they find diversification in virulence. The differences in our results are likely

due to several crucial differences in our assumptions. Here, we modelled both hypo- and hyper-

virulence and transmission, whereas Sandhu et al (2021) focused on hypo- virulence and

transmission. When we restrict our parameter space to hypo- virulence ($\lambda < 1$) and transmission

($\eta < 1$) we regain the key finding from Sandhu et al (2021) that higher virulence always evolves (Fig.

1A). However, this does not hold when there is hypervirulence ($\lambda > 1$; Fig. 1C). The hysteresis effect

produced by the bistable region is also more prominent when there is hypervirulence, which may

explain why it has not previously been observed. Recovery is possible for many infectious diseases

and moreover, one would not expect the hyperparasite to always co-transmit with the parasite ($\rho <$

1). When we remove recovery from our model ($\gamma = 0$), and assume that the hyperparasite is always

co-transmitted with the parasite ($\rho = 1$), as in Sandhu et al (2021), the bistable region is still

present (Fig. S3), and so these differences in model assumptions do not explain the lack of bistability

in Sandhu et al (2021). Notably, Sandhu et al (2021) allowed superinfection (replacement of parasite

strains) mediated by the hyperparasite and assumed that the transmissibility of the hyperparasite

($\sigma$) was linked to the infectivity of the parasite, such that more infectious parasites also generated

more infections by hyperparasites. These effects were not present in our model. As the bistable

region in our model represents a change in parasite strategies due to contrasts in hyperparasite

prevalence, with more virulent parasites evolving when hyperparasite prevalence is lower, the

positive relationship between parasite transmission/virulence and hyperparasite infectivity may

explain the lack of bistability in Sandhu et al (2021).

Although models of hyperparasitism are relatively rare, especially in an evolutionary context, some

of our key findings are mirrored in models of other tripartite systems. For example, multiple studies

have found that introducing an additional species to a host-parasite system can lead to evolutionary

branching in the host (Best 2018) or the parasite (Kisdi, Geritz, and Boldin 2013; Best 2018; Smith

and Ashby 2022; Morozov and Best 2012) populations. In a related study that also has relevance to

biocontrol, Smith and Ashby (2022) explore how the introduction of a tolerance-conferring defensive

symbiont affects the evolution of parasite virulence. They show that even if the defensive symbiont

is initially beneficial to the host population, in the long-term it is costly because it always selects for

higher parasite virulence. More generally, recent theory has highlighted how biocontrol agents can

lead to unintuitive ecological and evolutionary outcomes (Ashby and King 2017; Rafaluk-Mohr et al.

2018; King and Bonsall 2017; Nelson and May 2020). Our results similarly emphasise the complex

eco-evolutionary outcomes that can arise following the introduction of a hyperparasite, with

potentially disastrous consequences for the host population.

In established host-parasite-hyperparasite systems, it is difficult to separate the ecological and

evolutionary consequences of the hyperparasite. Few empirical studies have therefore explored the

effects of hyperparasitism on parasite evolution (Parratt and Laine 2016), and those that do often

focus on consequences for antimicrobial resistance (Chan et al. 2016; Burmeister et al. 2020), the

acquisition of virulence factors from hyperparasites (Miao and Miller 1999), or pleiotropic effects on

virulence due to selection for resistance against hyperparasitism (Castledine et al. 2022). For

example, (Evans et al. 2010) showed that strains of the bacteria *Erwinia carotovora ssp. atroseptica*

*(Eca)* resistant to the hyperparasite $\phi$AT1 were less likely to produce rot in potato tubers; Casteldine

et al. showed that evolution of phage resistance in *Pseudomonas aeruginosa* coincided with the loss

of virulence in vitro and in vivo; and (Scanlan and Buckling 2012) showed that coevolution between

the bacteria *P. fluourescens* and a lytic phage ($\phi$2) selects for a mucoid phenotype, which is a

virulence factor in both lung infections of cystic fibrosis patients and in plant infections. Clearly, the

available empirical evidence suggests that hyperparasites can indeed have significant evolutionary

effects on parasite virulence, although the precise effects may depend on pleiotropy between

resistance and virulence. Our model did not consider resistance to hyperparasitism but

understanding how pleiotropy with resistance affects virulence evolution is a critical direction for

future theoretical work (Sandhu et al. 2021). Additionally, were the hyperparasite to die out at any

point, the parasite would always return to the ancestral level of virulence prior to the introduction of

the hyperparasite (Fig. S4).

We focused our investigation on the evolution of virulence, but the evolutionary dynamics of the

hyperparasite are also likely to be important (Sandhu et al. 2021; Northrup et al. 2021), especially for

driving additional eco-evolutionary feedback loops (Ashby et al. 2019) . However, in certain cases the

hyperparasite might behave as if it is evolutionarily static (e.g., the repeated application of a

particular biocontrol to an agricultural crop), in which case the model analysed here will be

especially applicable. Still, future theoretical work should consider how our findings are affected by

coevolution with the parasite and/or the host (Buckingham and Ashby 2022) , and whether the

hysteresis in our model could lead to fluctuating selection as in Prado et al. (2009). Additionally, the

joint dynamics of resistance and virulence evolution deserve further scrutiny, which could be

explored using a resource-allocation model where the parasite can either allocate resources to

transmission or defence, with the overall resource "budget" depending on the level of virulence.

Overall, we have shown that the introduction of hyperparasites can have dramatic effects on the

evolutionary dynamics of parasite virulence, and when this leads to selection for higher or lower

virulence, or to evolutionary branching, with the outcome critically dependent on the within-host

effects of the hyperparasite on transmission and virulence, and on the probability of co-

transmission. We have also shown how relatively small changes in hyperparasite traits can have

dramatic consequences for virulence evolution, which has important implications for the use of

hyperparasites as agents of biocontrol.

ACKNOWLEDGEMENTS

JW is supported by a scholarship from the EPSRC Centre for Doctoral Training in Statistical Applied
Mathematics at Bath (SAMBa), under the project EP/L015684/1. BA is supported by the Natural
Environment Research Council (grant numbers NE/N014979/1 and NE/V003909/1). We acknowledge
the support of the Natural Sciences and Engineering Research Council of Canada (NSERC). Nous
remercions le Conseil de recherches en sciences naturelles et en génie du Canada (CRSNG) de son
soutien.

CONFLICTS OF INTEREST

The authors declare they have no conflicts of interest.

REFERENCES

Ashby, Ben, Ryosuke Iritani, Alex Best, Andy White, and Mike Boots. 2019. 'Understanding the Role
of Eco-Evolutionary Feedbacks in Host-Parasite Coevolution'. *Journal of Theoretical Biology* 464
(March): 115–25. https://doi.org/10.1016/j.jtbi.2018.12.031.

Ashby, Ben, and Kayla C. King. 2017. 'Friendly Foes: The Evolution of Host Protection by a Parasite:
EVOLUTION OF HOST PROTECTION BY A PARASITE'. *Evolution Letters* 1 (4): 211–21.
https://doi.org/10.1002/evl3.19.

Ashelford, Kevin E., Martin J. Day, and John C. Fry. 2003. 'Elevated Abundance of Bacteriophage
Infecting Bacteria in Soil'. *Applied and Environmental Microbiology* 69 (1): 285–89.
https://doi.org/10.1128/AEM.69.1.285-289.2003.

Berchieri, A., M.A. Lovell, and P.A. Barrow. 1991. 'The Activity in the Chicken Alimentary Tract of
Bacteriophages Lytic for Salmonella Typhimurium'. *Research in Microbiology* 142 (5): 541–49.
https://doi.org/10.1016/0923-2508(91)90187-F.

Best, Alex. 2018. 'Host–Pathogen Coevolution in the Presence of Predators: Fluctuating Selection
and Ecological Feedbacks'. *Proceedings of the Royal Society B: Biological Sciences* 285 (1885):
20180928. https://doi.org/10.1098/rspb.2018.0928.

Bowers, Roger G., Andrew Hoyle, Andrew White, and Michael Boots. 2005. 'The Geometric Theory of
Adaptive Evolution: Trade-off and Invasion Plots'. *Journal of Theoretical Biology* 233 (3): 363–77.
https://doi.org/10.1016/j.jtbi.2004.10.017.

Bryner, Sarah Franziska, and Daniel Rigling. 2012. 'VIRULENCE NOT ONLY COSTS BUT ALSO BENEFITS
THE TRANSMISSION OF A FUNGAL VIRUS: VIRULENCE AND TRANSMISSIBILITY OF A FUNGAL VIRUS'.
*Evolution* 66 (8): 2540–50. https://doi.org/10.1111/j.1558-5646.2012.01637.x.

Buckingham, Lydia J., and Ben Ashby. 2022. 'Coevolutionary Theory of Hosts and Parasites'. *Journal*
*of Evolutionary Biology* 35 (2): 205–24. https://doi.org/10.1111/jeb.13981.

Burmeister, Alita R., Abigail Fortier, Carli Roush, Adam J. Lessing, Rose G. Bender, Roxanna
Barahman, Raeven Grant, Benjamin K. Chan, and Paul E. Turner. 2020. 'Pleiotropy Complicates a
Trade-off between Phage Resistance and Antibiotic Resistance'. *Proceedings of the National*
*Academy of Sciences* 117 (21): 11207–16. https://doi.org/10.1073/pnas.1919888117.

Castledine, Meaghan, Daniel Padfield, Pawel Sierocinski, Jesica Soria Pascual, Adam Hughes, Lotta
Mäkinen, Ville-Petri Friman, et al. 2022. 'Parallel Evolution of Pseudomonas Aeruginosa Phage
Resistance and Virulence Loss in Response to Phage Treatment in Vivo and in Vitro'. *ELife* 11
(February): e73679. https://doi.org/10.7554/eLife.73679.

Chan, Benjamin K., Mark Sistrom, John E. Wertz, Kaitlyn E. Kortright, Deepak Narayan, and Paul E.
Turner. 2016. 'Phage Selection Restores Antibiotic Sensitivity in MDR Pseudomonas Aeruginosa'.
*Scientific Reports* 6 (1): 26717. https://doi.org/10.1038/srep26717.

Chen, Yibao, Lan Yang, Dan Yang, Jiaoyang Song, Can Wang, Erchao Sun, Changqin Gu, et al. 2020.
'Specific Integration of Temperate Phage Decreases the Pathogenicity of Host Bacteria'. *Frontiers in*
*Cellular and Infection Microbiology* 10 (February): 14. https://doi.org/10.3389/fcimb.2020.00014.

Chhibber, Sanjay, Sandeep Kaur, and Seema Kumari. 2008. 'Therapeutic Potential of Bacteriophage
in Treating Klebsiella Pneumoniae B5055-Mediated Lobar Pneumonia in Mice'. *Journal of Medical*
*Microbiology* 57 (12): 1508–13. https://doi.org/10.1099/jmm.0.2008/002873-0.

Evans, T.J., A. Trauner, E. Komitopoulou, and G.P.C. Salmond. 2010. 'Exploitation of a New
Flagellatropic Phage of *Erwinia* for Positive Selection of Bacterial Mutants Attenuated in Plant
Virulence: Towards Phage Therapy'. *Journal of Applied Microbiology* 108 (2): 676–85.
https://doi.org/10.1111/j.1365-2672.2009.04462.x.

Faruque, Shah M., Iftekhar Bin Naser, M. Johirul Islam, A. S. G. Faruque, A. N. Ghosh, G. Balakrish
Nair, David A. Sack, and John J. Mekalanos. 2005. 'Seasonal Epidemics of Cholera Inversely Correlate
with the Prevalence of Environmental Cholera Phages'. *Proceedings of the National Academy of*
*Sciences* 102 (5): 1702–7. https://doi.org/10.1073/pnas.0408992102.

Ferriol-González, Celia, and Pilar Domingo-Calap. 2021. 'Phage Therapy in Livestock and Companion
Animals'. *Antibiotics* 10 (5): 559. https://doi.org/10.3390/antibiotics10050559.

Gordillo Altamirano, Fernando L., and Jeremy J. Barr. 2019. 'Phage Therapy in the Postantibiotic Era'.
*Clinical Microbiology Reviews* 32 (2): e00066-18. https://doi.org/10.1128/CMR.00066-18.

Greer, G. Gordon. 2005. 'Bacteriophage Control of Foodborne Bacteria†'. *Journal of Food Protection*
68 (5): 1102–11. https://doi.org/10.4315/0362-028X-68.5.1102.

Hargreaves, Katherine R., Andrew M. Kropinski, and Martha R. J. Clokie. 2014. 'What Does the
Talking?: Quorum Sensing Signalling Genes Discovered in a Bacteriophage Genome'. Edited by
Gunnar F. Kaufmann. *PLoS ONE* 9 (1): e85131. https://doi.org/10.1371/journal.pone.0085131.

Holtappels, Dominique, Kiandro Fortuna, Rob Lavigne, and Jeroen Wagemans. 2021. 'The Future of
Phage Biocontrol in Integrated Plant Protection for Sustainable Crop Production'. *Current Opinion in*
*Biotechnology* 68 (April): 60–71. https://doi.org/10.1016/j.copbio.2020.08.016.

Hurford, Amy, Daniel Cownden, and Troy Day. 2010. 'Next-Generation Tools for Evolutionary
Invasion Analyses'. *Journal of The Royal Society Interface* 7 (45): 561–71.
https://doi.org/10.1098/rsif.2009.0448.

Kakkanat, Asha, Minh-Duy Phan, Alvin W. Lo, Scott A. Beatson, and Mark A. Schembri. 2017. 'Novel
Genes Associated with Enhanced Motility of Escherichia Coli ST131'. Edited by Mark Isalan. *PLOS*
*ONE* 12 (5): e0176290. https://doi.org/10.1371/journal.pone.0176290.

King, Kayla C., and Michael B. Bonsall. 2017. 'The Evolutionary and Coevolutionary Consequences of
Defensive Microbes for Host-Parasite Interactions'. *BMC Evolutionary Biology* 17 (1): 190.
https://doi.org/10.1186/s12862-017-1030-z.

Kisdi, Eva, Stefan A.H. Geritz, and Barbara Boldin. 2013. 'Evolution of Pathogen Virulence under
Selective Predation: A Construction Method to Find Eco-Evolutionary Cycles'. *Journal of Theoretical*
*Biology* 339 (December): 140–50. https://doi.org/10.1016/j.jtbi.2013.05.023.

Ludwig, D., D. D. Jones, and C. S. Holling. 1978. 'Qualitative Analysis of Insect Outbreak Systems: The
Spruce Budworm and Forest'. *The Journal of Animal Ecology* 47 (1): 315.
https://doi.org/10.2307/3939.

Miao, Edward A., and Samuel I. Miller. 1999. 'Bacteriophages in the Evolution of Pathogen–Host
Interactions'. *Proceedings of the National Academy of Sciences* 96 (17): 9452–54.
https://doi.org/10.1073/pnas.96.17.9452.

Mikhailov, Kirill V., Timur G. Simdyanov, and Vladimir V. Aleoshin. 2016. 'Genomic Survey of a
Hyperparasitic Microsporidian *Amphiamblys* Sp. (Metchnikovellidae)'. *Genome Biology and*
*Evolution*, September, evw235. https://doi.org/10.1093/gbe/evw235.

Mohan, Sharad, K. Kiran Kumar, Vivek Sutar, Supradip Saha, Janet Rowe, and Keith G. Davies. 2020.
'Plant Root-Exudates Recruit Hyperparasitic Bacteria of Phytonematodes by Altered Cuticle Aging:
Implications for Biological Control Strategies'. *Frontiers in Plant Science* 11 (June): 763.
https://doi.org/10.3389/fpls.2020.00763.

Morozov, Andrew, and Alex Best. 2012. 'Predation on Infected Host Promotes Evolutionary
Branching of Virulence and Pathogens' Biodiversity'. *Journal of Theoretical Biology* 307 (August): 29–
36. https://doi.org/10.1016/j.jtbi.2012.04.023.

Morozov, Andrew Yu., Cécile Robin, and Alain Franc. 2007. 'A Simple Model for the Dynamics of a
Host–Parasite–Hyperparasite Interaction'. *Journal of Theoretical Biology* 249 (2): 246–53.
https://doi.org/10.1016/j.jtbi.2007.05.041.

Munns, Krysty D., L. Brent Selinger, Kim Stanford, Leluo Guan, Todd R. Callaway, and Tim A.
McAllister. 2015. 'Perspectives on Super-Shedding of *Escherichia Coli* O157:H7 by Cattle'. *Foodborne*
*Pathogens and Disease* 12 (2): 89–103. https://doi.org/10.1089/fpd.2014.1829.

Nelson, Paul, and Georgiana May. 2020. 'Defensive Symbiosis and the Evolution of Virulence'. *The*
*American Naturalist* 196 (3): 333–43. https://doi.org/10.1086/709962.

Northrup, Graham R, Steven R Parratt, Carly Rozins, Anna-Liisa Laine, and Mike Boots. 2021. 'The
Evolutionary Dynamics of Hyperparasites'. Preprint. Evolutionary Biology.
https://doi.org/10.1101/2021.12.01.470853.

Nuss, Donald L. 2005. 'Hypovirulence: Mycoviruses at the Fungal–Plant Interface'. *Nature Reviews*
*Microbiology* 3 (8): 632–42. https://doi.org/10.1038/nrmicro1206.

Obradovic, A., J. B. Jones, M. T. Momol, B. Balogh, and S. M. Olson. 2004. 'Management of Tomato
Bacterial Spot in the Field by Foliar Applications of Bacteriophages and SAR Inducers'. *Plant Disease*
88 (7): 736–40. https://doi.org/10.1094/PDIS.2004.88.7.736.

Parratt, Steven R, and Anna-Liisa Laine. 2016. 'The Role of Hyperparasitism in Microbial Pathogen
Ecology and Evolution'. *The ISME Journal* 10 (8): 1815–22. https://doi.org/10.1038/ismej.2015.247.

Parratt, Steven R., and Anna-Liisa Laine. 2018. 'Pathogen Dynamics under Both Bottom-up Host
Resistance and Top-down Hyperparasite Attack'. Edited by Ainhoa Magrach. *Journal of Applied*
*Ecology* 55 (6): 2976–85. https://doi.org/10.1111/1365-2664.13185.

Prado, Federico, Alyssa Sheih, Jevin D. West, and Benjamin Kerr. 2009. 'Coevolutionary Cycling of
Host Sociality and Pathogen Virulence in Contact Networks'. *Journal of Theoretical Biology* 261 (4):
561–69. https://doi.org/10.1016/j.jtbi.2009.08.022.

Prospero, Simone, Leticia Botella, Alberto Santini, and Cécile Robin. 2021. 'Biological Control of
Emerging Forest Diseases: How Can We Move from Dreams to Reality?' *Forest Ecology and*
*Management* 496 (September): 119377. https://doi.org/10.1016/j.foreco.2021.119377.

Rafaluk-Mohr, Charlotte, Ben Ashby, Dylan A. Dahan, and Kayla C. King. 2018. 'Mutual Fitness
Benefits Arise during Coevolution in a Nematode-Defensive Microbe Model'. *Evolution Letters* 2 (3):
246–56. https://doi.org/10.1002/evl3.58.

Sandhu, Simran K., Andrew Yu. Morozov, Robert D. Holt, and Michael Barfield. 2021. 'Revisiting the
Role of Hyperparasitism in the Evolution of Virulence'. *The American Naturalist* 197 (2): 216–35.
https://doi.org/10.1086/712351.

Scanlan, Pauline Deirdre, and Angus Buckling. 2012. 'Co-Evolution with Lytic Phage Selects for the
Mucoid Phenotype of Pseudomonas Fluorescens SBW25'. *The ISME Journal* 6 (6): 1148–58.
https://doi.org/10.1038/ismej.2011.174.

Schooley, Robert T., Biswajit Biswas, Jason J. Gill, Adriana Hernandez-Morales, Jacob Lancaster,
Lauren Lessor, Jeremy J. Barr, et al. 2017. 'Development and Use of Personalized Bacteriophage-
Based Therapeutic Cocktails To Treat a Patient with a Disseminated Resistant Acinetobacter
Baumannii Infection'. *Antimicrobial Agents and Chemotherapy* 61 (10): e00954-17.
https://doi.org/10.1128/AAC.00954-17.

Smith, Cameron A., and Ben Ashby. 2022. 'Tolerance-Conferring Defensive Symbionts and the
Evolution of Parasite Virulence'. Preprint. Evolutionary Biology.
https://doi.org/10.1101/2022.12.05.519091.

Taylor, Douglas R., Andrew M. Jarosz, Dennis W. Fulbright, and Richard E. Lenski. 1998. 'The
Acquisition of Hypovirulence in Host-Pathogen Systems with Three Trophic Levels'. *The American*
*Naturalist* 151 (4): 343–55. https://doi.org/10.1086/286123.

Waldor, Matthew K., and John J. Mekalanos. 1996. 'Lysogenic Conversion by a Filamentous Phage
Encoding Cholera Toxin'. *Science* 272 (5270): 1910–14.
https://doi.org/10.1126/science.272.5270.1910.

Wendling, Carolin C., Agnes Piecyk, Dominik Refardt, Cynthia Chibani, Robert Hertel, Heiko
Liesegang, Boyke Bunk, Jörg Overmann, and Olivia Roth. 2017. 'Tripartite Species Interaction:
Eukaryotic Hosts Suffer More from Phage Susceptible than from Phage Resistant Bacteria'. *BMC*
*Evolutionary Biology* 17 (1): 98. https://doi.org/10.1186/s12862-017-0930-2.