Title: Defining the cellular complexity of the zebrafish bipotential gonad

Authors: Michelle E. Kossack[1†], Lucy Tian[1], Kealyn Bowie[2], Jessica S. Plavicki[1*‡]

[1] Pathology and Laboratory Medicine Department, Brown University, Providence, RI

[2] Department of Ecology, Evolution, and Organismal Biology, Brown University, Providence, RI

* Corresponding Author: jessica_plavicki@brown.edu, 70 Ship St, Box G-E5, Providence, RI,

02903

[†] Supported by F32ES023650 and 5T32ES007272 both from NIEHS

[‡] Supported by NIEHS K99/R00 (ES023848), a CPVB Phase II COBRE (2PG20GM103652),

and an NIEHS ONES award (ES030109)

Running Title: cellular complexity of the bipotential gonad

Keywords: Zebrafish, bipotential gonad, endothelial, lymphatic, pericyte, macrophage,

reproduction, single-cell, vasculature

Summary Sentence

Delineating the complex cellular interactions between vascular and lymphatic endothelial cells,

pericytes, and macrophage in the bipotential gonad is essential for understanding the

differentiation and functioning of the mature gonad.

Abstract

Zebrafish are routinely used to model reproductive development, function, and disease, yet we

still lack a clear understanding of the fundamental steps that occur during early bipotential

gonad development, including when endothelial cells, pericytes, and macrophage cells arrive at

the bipotential gonad to support gonad growth and differentiation. Here, we use a combination

of transgenic reporters and single-cell sequencing analyses to define the arrival of different

critical cell types to the larval zebrafish gonad. We determined that blood initially reaches the

gonad via a vessel formed from the swim bladder artery, which we have termed the gonadal

artery. We find that vascular and lymphatic development occurs concurrently in the bipotential

zebrafish gonad and our data suggest that similar to what has been observed in developing

zebrafish embryos, lymphatic endothelial cells in the gonad may be derived from vascular

endothelial cells. We mined preexisting sequencing data sets to determine whether ovarian

pericytes had unique gene expression signatures. We identified 215 genes that were uniquely

expressed in ovarian pericytes that were not expressed in larval pericytes. Similar to what has

been shown in the mouse ovary, our data suggest that *pdgfrb*+ pericytes may support the

migration of endothelial tip cells during ovarian angiogenesis. Using a macrophage-driven

photoconvertible protein, we found that macrophage established a nascent resident population

as early as 12 dpf and can be observed removing cellular material during gonadal

differentiation. This foundational information demonstrates that the early bipotential gonad

contains complex cellular interactions, which likely shape the health and function of the mature,

differentiated gonad.

Graphical Abstract

Introduction:

The developmental basis of the adult disease hypothesis postulates that the origins of some diseases can be traced back to developmental changes, sometimes very subtle, that produce long-term cascading effects that ultimately alter organ function and health (1). Given their genetic similarity to humans, zebrafish have been extensively used to model early embryonic development (2). More recently zebrafish have also been used to model juvenile development and adult health, including gonad formation and fertility (3–5).

The zebrafish gonad undergoes the same developmental stages as mammals; however, the timeline is elongated. The protracted developmental timeline allows for a detailed examination of very early periods of gonadal development, which have been understudied and may be sensitive to genetic and environmental disruptions. In mice and rats, primordial germ cells form at 7 days post conception (dpc) and, between ~9.5 and ~11.5 dpc, migrate from the hindgut to the urogenital ridge (6–8). In mice, the primary sex-determining gene, *sex-determining region Y* (*Sry*), must be activated before 11 dpc to initiate the transition from a bipotential gonad to a testicular development program (9). Since the murine bipotential period only lasts <2 days, it can be difficult to investigate early sources of reproductive disease that could occur during this critical developmental stage.

In mice, a major area of research has been to delineate the role of support and steroidogenic cells in sex determination and gonad maturation. Comparatively less has been done to investigate the role of other critical cell types in the developing bipotential and differentiated mammalian gonad. Research indicates that additional cell types, including macrophage, endothelial cells, and pericytes, are present at early stages of gonad development. At E10.5 in the mouse, macrophage are found at the urogenital ridge, just prior to sex determination (10). Macrophage depletion has been shown to hinder vascular remodeling and morphogenesis in the fetal testis (10). Endothelial cells in XX and XY individuals are present by 10.5 dpc and form lumenized vessels that are perfused with blood (10,11). Vessel

formation is essential for testis development (12–14); however, considerably less is known

about the dynamics of vascular development in the ovary as well as how lymphatic and

perivascular cell development affects ovarian development (15). Recently, single-cell

sequencing in 10.5 dpc mouse gonad and mesonephros found many different cell types are

present at this early stage, including vascular and lymphatic endothelial cells, pericytes, and

macrophage (16,17).

In the zebrafish, germ cells are specified at 4 hours post-fertilization (18). However, the

primordial germ cells remain quiescent while migrating ventral of the hindgut, a process which

occurs from 1 day post-fertilization (dpf) to approximately 10 dpf (see (19) for a detailed review).

Somatic cells are shown in close association with primordial germ cells as early as 4 dpf (20).

By 11 dpf, the somatic cells have formed a bilayer around the germ cells and begin to express

sex-specific markers (21). From 10 dpf until approximately 20 dpf, most of the germ cells

undergo mitosis, initiate meiosis, and remain in a bipotential state. During the bipotential period,

while early-stage oocytes are forming, somatic cells surround the germ cells and the gonad

exhibits mosaic expression of testicular- and ovarian-specific genes (i.e. anti-Mullerian hormone

or aromatase, respectively). Early-stage oocytes in the bipotential period produce paracrine

signals that act on somatic cells to further support oocyte development (21–23). The process of

sex determination is initiated around 20 dpf. In the presence of sufficient oocyte-derived

signaling, the oocytes continue to mature and, ultimately, form an ovary with mature granulosa

and theca cells, which support ovarian health and function. In the absence of sufficient oocyte-

derived signaling, oocytes undergo apoptosis and the gonad will progress with a testicular

developmental program with Sertoli and Leydig cells supporting spermatogenesis (24,25). Thus,

during the bipotential phase of zebrafish gonad development, although oocytes are present, the

ultimate fate of the gonad is still undetermined. Unlike in the developing mouse testis and ovary,

there is no mesonephric tissue adjacent to zebrafish gonad to give rise to the gonadal

vasculature. There are still fundamental questions about the lineage of support cells in the

differentiated ovary and testis. It is not definitively known if the somatic cells present during the

bipotential phase are progenitors for mature support cells or if mature support cells of the adult

gonad are derived from a different origin.

While it is known that the mature zebrafish ovary and testis contain endothelial and

lymphatic vasculature as well as perivascular cells and macrophage (26), the developmental

timeline for the arrival of these cell types to the gonad has not previously been described. The

long bipotential period of gonadal development in zebrafish compared to mice provides an

excellent opportunity to thoroughly assess the arrival of these critical cell types prior to sex

determination. Using transgenic zebrafish with fluorescently expressed cellular markers, we

tracked the arrival of the endothelial and lymphatic vasculature as well as perivascular cells and

macrophage to the bipotential zebrafish gonad. We found that the bipotential gonad is more

cellularly complex than originally thought. This foundational information suggests that additional

cell types have the potential to contribute to gonad development, sex determination, and fertility.

Material and Methods

Zebrafish Rearing: Zebrafish (*Danio rerio*) were raised as described in Kossack et al. and

Westerfield (27,28). Larvae were placed in tanks at 5 dpf in 100mL rotifer culture, 400mL of fish

water static, and one drop of RG Complete (Reef Nutrition). From 5 to10 dpf, larvae were fed

GEMMA Micro 75 (Skretting Zebrafish©) twice a day and live rotifers once a day. At 10 dpf,

water flow was initiated as a slow drip with a rate of approximately one drop a second. From 10-

20 dpf, fish were fed GEMMA Micro 75 twice a day along with rotifers and artemia once a day.

From 21 to 28 dpf, the water flow rate was increased until a standard flow rate was established.

The fish were treated as adults at 90 dpf and fed GEMMA Micro 300 once a day.

Juvenile fish were euthanized according to procedures approved by the Institutional

Animal Care and Use Committee at Brown University and adhered to the National Institutes of

Health "Guide for the Care and Use of Laboratory Animals". In short, fish were placed in 0.04%

MS-222 solution for 10-15 mins followed by ice water for 20 mins at which point they were

decapitated and placed in 4% PFA overnight.

In establishing the developmental timeline, we measured the standard length of fish. The

standard length is correlated with developmental stage during these ages (29,30). The full list of

replicates, individuals, average length, and observations can be found in **Supplemental Table**

S1.

Transgenic lines utilized: To mark vascular endothelial cells, we used three transgenic lines,

Tg(fli1:nEGFP)^{y7}, Tg(kdrl:GFP) originally *Tg(flk1:GFP)^{la116},* and *Tg(kdrl:DsRed2)* (31–33).

Lymphatic endothelial cells were labeled with *Tg(mrc1a:EGFP)* and *Tg(-5.2lyve1b:DsRed)*

(34,35). Tg(piwil1:EGFP) originally *Tg(ziwi:EGFP)* marked germ cells, and

TgBAC(pdgfrb:EGFP) marked perivascular cells (36,37). In the study of macrophage, we

utilized *Tg(mpeg1:EGFP), Tg(mpeg1:Gal4FF)^{gl25},* and *Tg(UAS:Kaede)* (38,39). For readability

and simplicity, we will refer to these transgenic lines by the simplified notations inside the

parentheses, i.e *Tg(-5.2lyve1b:DsRed2)* is listed as *lyve1b:DsRed*.

Antibody Staining: Antibody staining for germ cell protein Vasa was performed as described in

Leerberg et al. (21) using anti-Vasa antibody from GeneTex (GTX1238306) at a working

concentration of 1:1000. From 10-15 dpf, all gonads were left intact in the body cavity with the

gut and excess anterior and posterior portions of the fish dissected away. In fish >15 dpf the

gonad was dissected out of the fish before staining. All secondary antibodies were used at

1:500 and are as follows: goat anti-rabbit IgG 488 (Thermo Fisher Scientific Cat# A-11034,

RRID:AB_2576217) and goat anti-rabbit IgG 633 (Thermo Fisher Scientific Cat# A-21071,

RRID:AB_2535732). After protein labeling, tissues (either whole gonad or body cavity) were

stained with Hoechst 33342 (Invitrogen H3570) at a concentration of 1:10,000 overnight,

dissected, and placed in Vectashield mounting media (Fisher Scientific, NC9265087) for

subsequent imaging.

Photoconversion and live imaging: Double transgenic line *mpeg1:GAL4;UAS:Kaede* fish were

identified at 3 dpf. At 12 dpf, fish were anesthetized in 0.02% MS222 for 1 minute then mounted

on a 35mm glass bottom microwell dish (MatTek, Part No. P35G-1.5-14-C) in 2% low-melting

temperature agarose (Fisher Scientific, bp1360-100) made in egg water (4 parts per thousand

salt). The mounted fish was surrounded in 0.02% MS-222 and placed on the confocal

microscope. Using 20x objective lens, the mid-section of the fish was exposed to full-power LED

405nm fluorescent light for 60 seconds to photoconvert the Kaede from green to red. The fish

was immediately removed from agarose and replaced in the tank. The tank was covered with

foil to protect the fish from additional light during transportation back to the fish facility. 24 hours

later, the fish were again anesthetized and mounted laterally in agarose. Using a 20x objective

lens, *piwil1:EGFP* expressing fish were identified from the previously photo-converted fish and

imaged.

Imaging: Images were collected on a Zeiss LSM 880 confocal microscope. Maximum intensity

projections, orthogonal slices, and colocalization analysis were performed with ZenBlack

(Zeiss). Tissues from 10 – 15 dpf zebrafish were mounted on Premium microscope slides

(Fisher Scientific, 12-544-7) with Premium cover glass (Fisher Scientific,12-548-B) held in place

by vacuum grease. Tissues >20 dpf were mounted in 1% low-melting agarose in a 35mm glass

bottom microwell dish. All raw imaging files are available (https://doi.org/10.26300/4ce5-jk07).

Sequencing Analysis: From Liu et al., (26) we obtained matrixes of gene and feature count.

These large matrixes gathered from raw single-cell sequencing data display individual cells on

one axis vs. counts of every gene detected on the other. From this data, we are able to perform

principal component analysis to group similar cells together, which is referred to as clustering.

For more details about the analysis see Seurat (40,41), and R code

(https://doi.org/10.26300/4ce5-jk07). Clustering of the ovarian cells was performed as described

in Liu et al. and cluster 5 was identified as the "Blood Vessel" cluster as described in the

publication (26). We isolated this group using the subset() function and looked at the expression

of vascular endothelial cell and lymphatic endothelial cell markers. We attempted sub-clustering

analysis, but there were insufficient cell numbers within this cluster alone to draw meaningful

information.

To generate the pericyte Venn diagram, we used publicly available data from Liu et al.

and Shih et al. (26,42). Briefly, both publications included lists of highly expressed genes

identified in all cell types. We, therefore, were able to download the list of genes expressed in

the cell types of interest, filter them for expression level, and compare the lists. From Liu et al.

(26), we downloaded the dataset of differentially expressed genes in "stromal cells" (from

Supplementary file 5). Based on Lui et al. ovarian stromal cells are cells that are not able to be

categorized as the following: germ, somatic, endothelial, or immune. Within ovarian stromal

cells, there were 6 sub-types of cells: interstitial, pericytes, vascular smooth muscle, stromal

progenitors, and ovarian cavity epithelial cells. From this data we created two groups, "ovarian

pericytes" which included genes highly expressed in the pericyte sub-cluster 2, and "ovarian

stromal cells" which included genes enriched in interstitial, vascular smooth muscle, stromal

progenitors, and ovarian cavity epithelial cells (sub-clusters 0, 1, 3, 4, 5, respectively). We

eliminated duplicates within each list to generate 1,831 genes in the ovarian stromal cell group

and 441 genes in the ovarian pericyte group.

Similarly, using publicly available data from Shih et al. (42), we downloaded the highly

expressed genes identified in all *pdgfrb*-expressing cells (from Table S2.1). As described in the

publication we filtered for expression level $\log_2 > 1$, this list of genes was defined as "*pdgfrb*+

cells". We performed the same methodology with Table S1.4, identified as "cluster 39", which

represents genes enriched in larval pericytes. We eliminated duplicates within each list, which

resulted in 2,084 genes in *pdgfrb*+ cells and 105 genes in cluster 39. Using the Multiple List

Comparator tool on molbiotools© (https://molbiotools.com/listcompare.php) we compared the

groups and created a 4-way venn diagram. List details are available in **Supplemental Table S2**.

Results

Vascular endothelial cells make initial contact with the anterior gonad

To establish a timeline of vascular development in the gonad, we used a double

transgenic line *kdrl:DsRed; fli1:nEGFP* that carries both the pan-endothelial marker *fli1:nEGFP*

as well as *kdrl:DsRed*, which marks vascular endothelial cells. Beginning at 10 dpf, we used a

qualitative scale to measure the distance between the gonad, marked by germ cells, and the

vascular endothelial cells. We defined the relationship between the cells using three categories,

"No Contact" (**Figure 1A**), "Proximal" (near, but not physically associated, **Figure 1B, Movie 1**),

or "Contact" (direct physical contact, **Figure 1D, Supplemental Figure S1, Movie 2**). Because

zebrafish do not have a coelomic epithelium, we defined the gonad as the cells within the

boundary created by the somatic cells. We found that in most individuals (n=15/17), contact

between vascular endothelial cells and germ cells occurred by approximately 20 dpf (**Figure**

2A), which corresponded to a standard length of ~7mm (**Figure 2B, Supplemental Table S1**).

We observed that vascular endothelial cells initially made contact with the anterior-most portion

of the developing gonad (**Figure 1E, Movies 1 and 2**). Prior to this study, it was not known

which vessels supplied blood to the bipotential gonad. We determined that at 20 dpf the blood

supply to the bipotential gonad originates from the swim bladder artery (**Figure 1H**). Posterior to

the anterior chamber of the swim bladder, the swim bladder artery divides, with two different

branches supplying blood to the right and left gonad (**Figure 1H**). We termed this artery the

gonadal artery.

At 40 dpf, the ovary and testis are morphologically distinct. The ovary is significantly

larger than the testis **(Figure 3A vs. 4A)**, however, both gonads are highly vascularized by this

time point **(Figure 3B and 4B)**. By 60 dpf, each maturing oocyte appears to be surrounded by a

vessel **(Figure 3E, Movie 3)**, while within the testis, vessels divide lobules of maturing sperm

(Figure 4E, Movie 4). Within the ovary, we found that vessels have distinct relationships with

different stage oocyte. Vessels come in direct contact with stage <1 oocyte, but are separated

by a layer of somatic cells between the endothelium and the oocytes at stages 2 and 3

(**Supplemental Figure S2**).

Specification of lymphatic endothelial cells in the gonad resembles larval lymphatic development

The role of the lymphatic vasculature in organogenesis and homeostasis has been

understudied in zebrafish (43). Recent single-cell sequencing of the zebrafish ovary found that

lymphatic endothelial cells are present in the ovary by 40 dpf (26); however, it is not known

when lymphatic cells invade the ovary and create functional networks within it. To determine

when vascular and lymphatic endothelial cells reach the ovary, we visualized the endothelium

by crossing canonical vascular endothelial reporter lines *kdrl:GFP* or *kdrl:DsRed* with

established lymphatic endothelial transgenic reporter lines *mrc1a:EGFP* or *lyve1b:DsRed* (34).

Using this double transgenic approach, we determined that lymphatic endothelial cells are

detected in the gonad between 15 and 20 dpf and are always present with vascular endothelial

cells (**Figure 2A,** 20 dpf, standard length of ~7mm, n=13/13). Environmental conditions and

genetic background will result in different growth rates; thus, the use of standard length is a

more accurate measure of developmental stage and supports the concurrent arrival of VEC and

LEC when the fish reach a length of ~7mm (**Figure 2B**).

It is unclear from our initial analysis of early gonadal development if vascular endothelial

cells give rise to lymphatic endothelial cells or if the lymphatic endothelium enters the gonad

independent of the vascular endothelium. During larval development, lymphatic endothelial cells

are derived from venous vascular endothelial cells (35,44–51). *flt4*, a marker of venous

endothelial cells and lymphatic endothelial cells, is required for initiating lymphatic endothelial

cell development. Expression of *flt4* is followed by the expression of *prox1a*, which specifies the

lymphatic fate in mammals (51–58). However, in other contexts, such as anal fin

vascularization, vascular endothelial cells are derived from lymphatic endothelial cells(57). To

gain insight into which mechanism might be utilized in the gonad, we mined previously available

single-cell sequencing from the 40 dpf zebrafish ovary (26). We sub-clustered the vasculature

("Blood Vessels") and looked for expression and co-localization of specific endothelial markers.

We found vascular endothelial markers *kdrl*, *flt1,* and pan-endothelial marker *fli1*, to be

expressed in the majority of cells (**Figure 5A**) whereas *lyve1b*, *flt4*, and *prox1a* were expressed

in a smaller subset of cells (**Figure 5B**). Next, we assessed whether *kdrl* and *lyve1b* were co-

expressed in this cluster and we found no overlap in expression (**Figure 5C**). However, *lyve1b*

was co-expressed with the venous marker *flt4*, suggesting that lymphatic development in the

gonad may occur in a manner similar to what has been observed in the larval endothelium (34).

We note that we were not able to examine the expression of *mrc1a* in the single-cell sequencing

data set because it was not detected in the original sequencing analysis.

To assess if lymphatic endothelial cells are derived from vascular endothelial cells in the

bipotential gonad, we used confocal microscopy to image 20 dpf gonads expressing transgenic

markers for both vascular endothelial cells and lymphatic endothelial cells (*kdrl:DsRed;*

mrc1a:EGFP). Our imaging revealed overlap in the *kdrl:DsRed* and *mrc1a:EGFP* expression

domains (**Figure 6A**). However, in most of the cells that expressed both genes, the level of

transgenic expression was not equal. We consistently observed expression of one marker being

dominant over the other in the maximum intensity projection (**Figure 6B and 6C**). Orthogonal

slices of the z-stacked images illustrate that the incongruent levels of expression between

markers are not an artifact of the maximum intensity projection (**Figure 6D**). Finally, we used

Zen Black to perform a co-localization analysis of the confocal images and found that although

the *mrc1a:*EGFP expression obscured the visibility of the *kdrl:DsRed*, there was 100% overlap

in signal (**Figure 6E,** Overlap coefficient = 0.97). Together this suggests that lymphatic

endothelial cells in the 20 dpf zebrafish gonad are derived from vascular endothelial cells, and

as lymphatic specific expression increases, vascular endothelial expression decreases.

Pdgfrb+ perivascular cells associate with vascular tip cells invading the bipotential gonad

Pericytes are a sub-type of perivascular cells that have, since their discovery, been

difficult to clearly distinguish from other perivascular cell types (59). In the developing zebrafish

embryo, pericytes were originally defined by their expression of platelet derived growth factor

receptor ß (*pdgfrb)*, their morphology, and their positional relationship to the vasculature.

Recent single-cell sequencing data from larval zebrafish as well as the imaging of *pdgfrb*:EGFP

zebrafish indicate that while pericytes tend to have high *pdgfrb* expression, they constitute only

a subset of the *pdgfrb*:EGFP+ cells (42). The subset of *pdgfrb*:EGFP+ cells with high *pdgfrb*

expression and low expression of other cell type markers was termed cluster 39 in a study by

Shih et al., using single-cell RNA sequencing to examine the signatures of gene expression in

larval zebrafish pericytes (42). Single-cell sequencing of the zebrafish ovary showed that

pdgfrb+ cells are found in a cluster of "stromal cells" defined as non-germ, non-somatic, non-

endothelial, and non-immune cells. Within the ovarian stromal cell cluster, ovarian pericytes had

the highest *pdgfrb* and *notch3* expression, known pericyte markers in larval zebrafish, and were

more uniquely defined by the expression of *plp1b* (26). We hypothesized that ovarian pericytes,

as defined by Liu et al., and the larval pericyte cluster (cluster 39) would have more similar

expression profiles than other *pdgfrb+* cells in the larvae or ovary. We compared expression

profiles of larval *pdgfrb*:EGFP cells and the larval pericytes (cluster 39) from Shih et al., with

ovarian stromal cells and ovarian pericytes from Liu et al. (26, 42). 215 genes were identified as

unique to ovarian pericytes (a complete list of overlaps in **Supplemental Table S2**). This finding

suggests that the 215 genes detected, including *plp1b*, may be unique to ovarian pericyte

function and not required at larval stages of pericyte development. Only 3 genes, *pcdh1b*,

ptenb, and *s1pr1* were found uniquely in ovarian pericytes and cluster 39 (**Figure 7**) *pcdh1b* and

s1pr1 are associated with cell adhesion (60,61) and are also both expressed in the follicle cell,

germline stem cell, and endothelial cell clusters. *ptenb* is expressed broadly in all of the ovarian

cell clusters and is a tumor suppressor gene known to control cell proliferation during

embryogenesis however its functions in the zebrafish ovary is unclear (62). *notch3,* along with

10 other genes, was expressed in all four cell groups.

As with all vascular beds, pericytes in the ovary are an essential part of the vascular unit,

creating stability for the constant vascular modeling that occurs as an inherent part of ovarian

function (as reviewed in (63)). Since *pdgfrb* is an imperfect marker of pericytes, we used the

double transgenic *pdgfrb:EGFP;kdrl:DsRed* to determine if *pdgfrb+* cells within the gonad were

also in contact with vessels. We found that oblong *pdgfrb+* cells are associated with vessels in

the developing gonad and based on their gene expression, position, and morphology, appeared

to be pericytes. The *pdgfrb+* presumptive pericytes arrived at the gonad along with the

endothelial cells (**Figure 2A,** 20 dpf, n=8/9). Confocal images revealed *pdgfrb+* cells in close

association with *kdrl+* vessels and extending tip cells (**Figure 8, Movie 5, Supplemental Figure**

S3), suggesting that pericytes may aid in the migration of vascular tip cells, which has been

observed during the vascularization of other tissues (37,64).

Macrophage arrive at the bipotential gonad prior to stromal cells types

Tissue resident macrophage play essential roles in organogenesis and organ

homeostasis and macrophage are known to play a substantial role in both ovary and testis

function in mice. However, the timeline of macrophage colonization of the zebrafish gonad

remains unclear (65–67). We found that while macrophage colonization of the gonad in most

individuals was concurrent with vascularization of the gonad, in some cases we observed

macrophage within the gonad prior to vascularization (**Figure 2A,** 12 dpf n=6/14, 3 replicates).

Notably, we found that at 20 dpf, when the gonad is undergoing sex determination, there are

more macrophage present, and they can be seen engulfing germ cells (**Figure 9B, 9C, Movie**

6). This suggests that macrophage may play an active role in gonad differentiation in zebrafish.

We asked if the macrophage that make initial contact with the gonad became resident

macrophage or if they were only transiently interacting with the germ cells. Given the superficial

position of the macrophage in contact with the early gonad (**Figure 9A**), we hypothesized that

macrophage found during the bipotential stage were not tissue-resident and the resident

population would not be established until after sex determination. To test this hypothesis, we

used a pan-macrophage reporter to drive the expression of Kaede *(mpeg1:GAL4; UAS:Kaede)*,

a photoconvertible fluorescent protein that changes from green to red following exposure to

405nm light. We exposed fish at 12 dpf to 405nm light in a defined region surrounding the

gonad (**Supplemental Figure S4**). As a result, macrophage in and near the gonad were

converted to red, leaving the remaining macrophage in the body green. 24 hours later we used

live imaging to observe the macrophage population within the gonad. The presence of only red

macrophage would indicate that no new macrophage had entered the gonad and suggest that

macrophage become resident prior to sexual differentiation. The presence of green macrophage

in addition to red macrophage would indicate either: (1) continued colonization of the gonad, (2)

the presence of transient populations of macrophage, or (3) newly divided macrophage from a

previously photoconverted cell although likely some red would remain from the mother cell. We

found that after 24 hours, a subset of macrophage were retained in the gonad and expressed

the red photoconverted kaede protein (**Movie 7 and 8**, **Figure 9E,** red arrowhead, n=5). We

also observed green macrophage in and in the proximity of the gonad (**Figure 9F,** green arrow).

Finally, several cells also were green and red mixed suggesting that they had some previous

conversion while generating new Kaede protein (**Figure 9D,** blue arrow). It is difficult to say

definitively where the double-labeled cells originated. Given the ratio of green to red in these

cells, it is possible that these cells were on the periphery of the conversion area and migrated to

the gonad while producing new Kaede protein. Together, these data suggest that at 12 dpf,

there is an early population of resident macrophage that cells within the bipotential gonad.

Discussion

Herein, we provide the first report of vascularization of the early zebrafish gonad, a

fundamental and understudied component of reproductive development and health. We found

that vascular endothelial cells and lymphatic endothelial cells are present in the ovary by 15 dpf.

In mice, previous studies have tracked the arrival of endothelial cells to the early gonad using

fluorescent imaging and single-cell sequencing at E10.5 (11,16,17). Based on single-cell

sequencing from Liu et al (26), we know that a diverse population of cells are present in the 40

dpf zebrafish ovary. Yet without a developmental timeline, we lack key spatial and temporal

information about gonadal development. Based on these previous studies, we can conclude that

there are parallels between the vascularization of the zebrafish and mouse bipotential phase of

the gonad. Although our investigation clearly indicates vascular and lymphatic endothelial cells

are present in the bipotential gonad, it does not evaluate when the vessels establish a functional

network. Future investigations will be required to determine when the vessels are lumenized

and are able to transport physiological signals and waste.

Prior mouse studies did not rigorously study both vascular and lymphatic endothelial cell

markers simultaneously in early development and, therefore, these studies do not allow us to

distinguish between endothelial cell types during gonad development (15,68,69). In adult mice,

vascular vs. lymphatic vessels have distinct domains in both the ovary and testis (11,69), but

how these domains are established in early development is unclear. In mice, Brennen et al. (11)

found that lymphatic endothelial cells, marked by Prox1[+/lacZ], are present in the mesonephros at

13.5 dpc and enter the gonad at 17.5 dpc in both sexes. However, using a transgenic Prox1-

EGFP mouse, Svingen et al. found that the ovary does not contain lymphatic endothelial cells

until after birth at post-natal day 10, while the testis was reported to contain lymphatic

endothelial cells by 17.5 dpc (69). Yet murine expression analysis using single-cell sequencing

shows evidence of *prox1* expression in the gonad at 10.5 dpc (16,17). Each of these studies

used different methods and criteria (lacZ, transgenic line, mRNA expression, respectively) to

determine the presence of lymphatic endothelial cells in the gonad using *Prox1*. Therefore, the

observed discrepancies could be due to differences in methodology or criteria. When comparing

the zebrafish and mice bipotential gonad, it becomes more difficult to assess the parallels

between lymphatic development in these two model species. Most murine studies use *Prox1* as

the lymphatic endothelial cell marker. In our study, we defined lymphatic endothelial cells to be

mrc1a or *lyve1b* expressing cells, because *prox1* is known to be expressed in a number of

different ovarian cell types and is not required for lymphatic cell specification in zebrafish

(49,54). Single-cell sequencing is able to detect extremely low expression of *Lyve1*, the

mammalian orthologs of *lyve1b*, in endothelial cells at 10.5 dpc, which suggests the earlier

timeline of lymphatic endothelial cell arrival to the gonad observed in zebrafish may also occur

in mice (17). It is well established that lymphatic vessels are not found within the testis, but

instead restricted to the outer tunicate; therefore, more research is needed to determine the

organization of lymphatics in relation to the adult zebrafish testis and the differences between

zebrafish and mouse sex-specific lymphatic development (10,69,70).

We also demonstrated that the *pdgfrb* expression domain within the ovary is not limited

to pericytes and likely includes other perivascular cells with different sub-functionalizations. We

show that *pdgfrb*+ cells extend between vessels within the gonad and are in close association

with migrating vascular tip cells, a phenomenon that has been observed in other developmental

contexts where vascular tip cells release *pdgf* ligands, which signal perivascular cells to follow

and stabilize the forming vessel (37,64,71).

Pericyte biology is still relatively nascent and *pdgfrb* has been routinely used as the

primary marker across species. In embryonic zebrafish, *pdgfrb* is required for the formation of

the intersegmental vessels and meningeal angiogenesis (72,73). In other species, it is known

that pericytes are essential for vascular stability and reconstruction of the ovarian vasculature

during follicle development and post-ovulation (74–77). Consistent with this finding, inhibition of

Pdgfrb in the rat ovary causes hemorrhaging (78). Together, these reports indicate a role of

pericytes in the function and establishment of the blood follicle barrier (79,80). When we

compared the gene expression profiles of *pdgrfb*+ cells with larval pericytes (cluster 39), ovarian

stromal cells, and ovarian pericytes, we found that 215 genes were unique to ovarian pericytes,

and only three genes were shared solely between ovarian pericytes and cluster 39. Of these,

we note that two were associated with cell adhesion (*pcdh1b* and *s1pr1*). *s1pr1* has been shown

to play critical roles in negatively regulating angiogenesis in zebrafish (60) and knockout of

S1pr1 in mice causes size-selective openings in the blood-brain barrier (61). Although it is *s1pr1*

is not specific to pericytes, as it is also expressed in other ovarian cells as well as in vascular

endothelial cells including hepatocellular carcinoma (81, 82), the shared expression of cell

adhesion genes in larval pericyte cluster (cluster 39) and ovarian pericytes supports the idea

that pericytes play a critical role in the blood follicle barrier development in zebrafish. It will be

necessary to understand how pericytes interact with the ovarian vasculature, particularly in

addressing ovarian disease. Both ovarian hyperstimulation syndrome and polycystic ovary

syndrome are associated with altered ovarian vasculature (83–85). Although it is well

established that pericytes perform a critical role in vascularization and angiogenesis, the role

that perivascular cells play in disease etiology remains unclear (reviewed in (84)). Zebrafish are

able to model some, but not all aspects of ovarian diseases such as PCOS (86), therefore the

discovery of additional cell types in the zebrafish bipotential gonad offers an opportunity to

investigate the role of these cells in the very early reproductive disease formation.

We discovered that macrophage are present in the early bipotential gonad and can be

seen engulfing germ cells during the bipotential phase. In mice, yolk-sac-derived macrophage

are responsible for clearing cellular debris from endothelial cells, Sertoli cells, and germ cells, as

well as for pruning vascular networks in the testis (10). In other contexts, macrophage are

known to release angiogenic cues that shape vascular development (87–91). Therefore, their

presence in the bipotential gonad prior to the arrival to vascular endothelial cells could suggest

that macrophage are be involved in directing vascularization of the gonad. While ovarian

resident macrophage are essential for folliculogenesis, ovulation, and the clearing of atretic

follicles (67,92,93) as well as cellular debris (66), prior to this study, macrophage phagocytosis

in the zebrafish ovary had not been observed. We also report that resident macrophage in the

gonad are present by 12 dpf and the population is dynamic during this developmental window

with new macrophage migrating into and out of the gonad. In mice, ovarian resident

macrophages can be divided into 5 different populations. Two of these initial subpopulations are

embryonically derived, one population from the yolk-sac, and the other from the fetal liver (66).

However, the majority of research studies have investigated resident macrophage function after

sex determination. The presence of macrophage in the zebrafish bipotential gonad suggests

that macrophage may also be present in the mammalian bipotential gonad and contribute to

important steps in sexual differentiation.

In conclusion, we found that vascular and lymphatic endothelial cells, pericytes, and

macrophage begin populating the bipotential gonad at 12 dpf. Our data suggest that vascular

endothelial cells may give rise to lymphatic endothelial cells in the gonad to together create a

complex network of endothelial cells by 20 dpf. However, it is still unclear when these vessels

become functional conduits for physiological signals and nutrients and additional loss of function

experiments are necessary to confirm this hypothesis. We found that *pdgfrb+* pericytes support

the migration of vascular tip cells and 215 genes are unique to ovarian pericytes compared to

larval pericytes. Finally, we found that tissue-resident macrophage can be found in the gonad as

early as 12 dpf and clear oocyte debris during sexual differentiation. Our foundational

information regarding vascular and lymphatic endothelial cell, perivascular, and macrophage

invasion into the biopotential gonad contributes to the understanding of the early bipotential

environment and how they may play a part in the etiology of reproductive disease.

Acknowledgments

We would like to thank Rachel Cyr for all her meticulous work caring and raising the fish. We

would also like to that the Weinstein and Draper labs for sharing their transgenic lines that made

this study possible. This work was made possible by funding from NIEHS to Dr. Michelle

Kossack F32ES023650 and 5T32ES007272. Dr. Jessica Plavicki was supported by an NIEHS

K99/R00 (ES023848), a CPVB Phase II COBRE (2PG20GM103652), and an NIEHS ONES

award (ES030109).

Data Availability

The data underlying this article are available in the Brown Digital Repository at:

https://doi.org/10.26300/4ce5-jk07

Conflict of Interest Statement

MEK, LT, KB, and JSP have no conflicts of interest to declare.

Author Contribution

MK: Conceptualization, methodology, investigation, writing – original draft, review, and editing,

visualization. LT and KB: Investigation, writing – review and editing. JSP: Resources, writing –

review and editing, supervision, project administration, fund acquisition.

References

1. De Boo HA, Harding JE. The developmental origins of adult disease (Barker) hypothesis. Aust N Z J Obstet Gynaecol. 2006 Feb 1;46(1):4–14.

2. Howe K, Clark MD, Torroja CF, Torrance J, Berthelot C, Muffato M, et al. The zebrafish reference genome sequence and its relationship to the human genome. Nature. 2013;496(7446):498–503.

3.   Baker TR, Peterson RE, Heideman W. Early Dioxin Exposure Causes Toxic Effects in Adult
Zebrafish. Toxicol Sci. 2013 Sep;135(1):241–50.

4.   King Heiden T, Carvan MJ, Hutz RJ. Inhibition of follicular development, vitellogenesis, and
serum 17β-estradiol concentrations in zebrafish following chronic, sublethal dietary
exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin. Toxicol Sci. 2006;90(2):490–9.

5.   King Heiden TC, Spitsbergen J, Heideman W, Peterson RE. Persistent adverse effects on
health and reproduction caused by exposure of zebrafish to 2,3,7,8-Tetrachlorodibenzo-p-
dioxin during early development and gonad differentiation. Toxicol Sci. 2009;109(1):75–87.

6.   Bendel–Stenzel M, Anderson R, Heasman J, Wylie C. The origin and migration of primordial
germ cells in the mouse. Semin Cell Dev Biol. 1998 Aug;9(4):393–400.

7.   McLaren A, Durcova-Hills G. Germ cells and pluripotent stem cells in the mouse. Reprod
Fertil Dev. 2001;13(8):661.

8.   Wilhelm D, Palmer S, Koopman P. Sex Determination and Gonadal Development in
Mammals. Physiol Rev. 2007 Jan 1;87(1):1–28.

9.   Hiramatsu R, Matoba S, Kanai-Azuma M, Tsunekawa N, Katoh-Fukui Y, Kurohmaru M, et
al. A critical time window of Sry action in gonadal sex determination in mice. Development.
2009;136(1):129–38.

10. DeFalco T, Bhattacharya I, Williams AV, Sams DM, Capel B. Yolk-sac-derived
macrophages regulate fetal testis vascularization and morphogenesis. Proc Natl Acad Sci.
2014;111(23):E2384–93.

11. Brennan J, Karl J, Capel B. Divergent Vascular Mechanisms Downstream of Sry Establish
the Arterial System in the XY Gonad. Dev Biol. 2002 Apr;244(2):418–28.

12. Cool J, DeFalco TJ, Capel B. Vascular-mesenchymal cross-talk through Vegf and Pdgf
drives organ patterning. Proc Natl Acad Sci. 2011;108(1):167–72.

13. Bott RC, McFee RM, Clopton DT, Toombs C, Cupp AS. Vascular Endothelial Growth Factor
and Kinase Domain Region Receptor Are Involved in Both Seminiferous Cord Formation
and Vascular Development During Testis Morphogenesis in the Rat1. Biol Reprod. 2006 Jul
1;75(1):56–67.

14. Combes AN, Wilhelm D, Davidson T, Dejana E, Harley V, Sinclair A, et al. Endothelial cell
migration directs testis cord formation. Dev Biol. 2009;326(1):112–20.

15. Bott RC, Clopton DT, Fuller AM, McFee RM, Lu N, McFee RM, et al. KDR-LacZ-expressing
cells are involved in ovarian and testis-specific vascular development, suggesting a role for
VEGFA in the regulation of this vasculature. Cell Tissue Res. 2010/09/17 ed. 2010
Oct;342(1):117–30.

16. Ademi H, Djari C, Mayère C, Neirijnck Y, Sararols P, Rands CM, et al. Deciphering the
origins and fates of steroidogenic lineages in the mouse testis. Cell Rep. 2022 Jun
14;39(11):110935.

17. Garcia-Alonso L, Lorenzi V, Mazzeo CI, Alves-Lopes JP, Roberts K, Sancho-Serra C, et al.
Single-cell roadmap of human gonadal development. Nature. 2022 Jul 1;607(7919):540–7.

18. Yoon C, Kawakami K, Hopkins N. Zebrafish vasa homologue RNA is localized to the
cleavage planes of 2- and 4-cell-stage embryos and is expressed in the primordial germ
cells. Development. 1997;124(16):3157–65.

19. Michelle E. Kossack, Bruce W. Draper. Genetic regulation of sex determination and
maintenance in zebrafish (Danio rerio). Curr Top Dev Biol. 2019;134:119–49.

20. Braat AK, Zandbergen T, Van De Water S, Goos HJTH, Zivkovic D. Characterization of
zebrafish primordial germ cells: Morphology and early distribution of vasa RNA. Dev Dyn.
1999;216(2):153–67.

21. Leerberg DM, Sano K, Draper BW. Fibroblast growth factor signaling is required for early
somatic gonad development in zebrafish. Mullins MC, editor. PLOS Genet. 2017 Sep
5;13(9):e1006993.

22. Dranow DB, Hu K, Bird AM, Lawry ST, Adams MT, Sanchez A, et al. Bmp15 Is an Oocyte-
Produced Signal Required for Maintenance of the Adult Female Sexual Phenotype in
Zebrafish. Mullins MC, editor. PLOS Genet. 2016 Sep 19;12(9):e1006323.

23. Wilson CA, High SK, McCluskey BM, Amores A, Yan YL, Titus TA, et al. Wild sex in
zebrafish: Loss of the natural sex determinant in domesticated strains. Vol. 198, Genetics.
2014. 1291–1308 p.

24. Dranow DB, Tucker RP, Draper BW. Germ cells are required to maintain a stable sexual
phenotype in adult zebrafish. Dev Biol. 2013 Apr;376(1):43–50.

25. Takahashi H. Juvenile hermaphroditism in the zebrafish, Brachydanio rerio. Bull Fac Fish
Hokkaido Univ. 1977;28(2):57–65.

26. Liu Y, Kossack ME, McFaul ME, Christensen LN, Siebert S, Wyatt SR, et al. Single-cell
transcriptome reveals insights into the development and function of the zebrafish ovary.
eLife. 2022 May 19;11:e76014.

27. Westerfield M. The zebrafish book: A guide for the laboratory use of zebrafish Danio
(Brachydanio) rerio. Eugene, OK: Westerfield, M; 2007.

28. Kossack ME, Manz KE, Martin NR, Pennell KD, Plavicki J. Environmentally relevant uptake,
elimination, and metabolic changes following early embryonic exposure to 2,3,7,8-
tetrachlorodibenzo-p-dioxin in zebrafish. Chemosphere. 2022;136723.

29. Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. Stages of embryonic
development of the zebrafish. Dev Dyn. 1995 Jul;203(3):253–310.

30. Singleman C, Holtzman NG. Growth and maturation in the zebrafish, Danio rerio: a staging
tool for teaching and research. Zebrafish. 2014/06/30 ed. 2014 Aug;11(4):396–406.

31. Roman BL, Pham VN, Lawson ND, Kulik M, Childs S, Lekven AC, et al. Disruption of acvrl1
increases endothelial cell number in zebrafish cranial vessels. Development.
2002;129(12):3009–19.

32. Choi J, Dong L, Ahn J, Dao D, Hammerschmidt M, Chen JN. FoxH1 negatively modulates
flk1 gene expression and vascular formation in zebrafish. Dev Biol. 2007 Apr
15;304(2):735–44.

33. Kikuchi K, Holdway JE, Major RJ, Blum N, Dahn RD, Begemann G, et al. Retinoic Acid
Production by Endocardium and Epicardium Is an Injury Response Essential for Zebrafish
Heart Regeneration. Dev Cell. 2011 Mar;20(3):397–404.

34. Jung HM, Castranova D, Swift MR, Pham VN, Venero Galanternik M, Isogai S, et al.
Development of the larval lymphatic system in zebrafish. Dev Camb. 2017;144(11):2070–
81.

35. Okuda KS, Astin JW, Misa JP, Flores MV, Crosier KE, Crosier PS. Lyve1 expression
reveals novel lymphatic vessels and new mechanisms for lymphatic vessel development in
zebrafish. Dev Camb. 2012;139(13):2381–91.

36. Leu DH, Draper BW. The ziwi promoter drives germline-specific gene expression in
zebrafish. Dev Dyn. 2010 Aug 24;239(10):2714–21.

37. Ando K, Fukuhara S, Izumi N, Nakajima H, Fukui H, Kelsh RN, et al. Clarification of mural
cell coverage of vascular endothelial cells by live imaging of zebrafish. Development. 2016
Jan 1;dev.132654.

38. Ellett F, Pase L, Hayman JW, Andrianopoulos A, Lieschke GJ. mpeg1 promoter transgenes
direct macrophage-lineage expression in zebrafish. Blood. 2011 Jan 27;117(4):e49–56.

39. Davison JM, Akitake CM, Goll MG, Rhee JM, Gosse N, Baier H, et al. Transactivation from
Gal4-VP16 transgenic insertions for tissue-specific cell labeling and ablation in zebrafish.
Dev Biol. 2007;304(2):811–24.

40. Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, et al. Integrated analysis
of multimodal single-cell data. Cell. 2021 Jun;184(13):3573-3587.e29.

41. Satija R, Farrell JA, Gennert D, Schier AF, Regev A. Spatial reconstruction of single-cell
gene expression data. Nat Biotechnol. 2015;33(5):495–502.

42. Shih YH, Portman D, Idrizi F, Grosse A, Lawson ND. Integrated molecular analysis identifies
a conserved pericyte gene signature in zebrafish. Development. 2021 Dec
7;148(23):dev200189.

43. Padberg Y, Schulte-Merker S, van Impel A. The lymphatic vasculature revisited—new
developments in the zebrafish. Methods Cell Biol. 2017;138:221–38.

44. Astin JW, Haggerty MJ, Okuda KS, Le Guen L, Misa JP, Tromp A, et al. Vegfd can
compensate for loss of Vegfc in zebrafish facial lymphatic sprouting. Development.
2014;141(13):2680–90.

45. Bussmann J, Bos FL, Urasaki A, Kawakami K, Duckers HJ, Schulte-Merker S. Arteries provide essential guidance cues for lymphatic endothelial cells in the zebrafish trunk. Development. 2010 Aug 15;137(16):2653–7.

46. Cha YR, Fujita M, Butler M, Isogai S, Kochhan E, Siekmann AF, et al. Chemokine signaling directs trunk lymphatic network formation along the preexisting blood vasculature. Dev Cell. 2012;22(4):824–36.

47. Hogan BM, Bos FL, Bussmann J, Witte M, Chi NC, Duckers HJ, et al. Ccbe1 is required for embryonic lymphangiogenesis and venous sprouting. Nat Genet. 2009;41(4):396–8.

48. Nicenboim J, Malkinson G, Lupo T, Asaf L, Sela Y, Mayseless O, et al. Lymphatic vessels arise from specialized angioblasts within a venous niche. Nature. 2015;522(7554):56–61.

49. van Impel A, Zhao Z, Hermkens DM, Roukens MG, Fischer JC, Peterson-Maduro J, et al. Divergence of zebrafish and mouse lymphatic cell fate specification pathways. Development. 2014;141(6):1228–38.

50. Yaniv K, Isogai S, Castranova D, Dye L, Hitomi J, Weinstein BM. Imaging the developing lymphatic system using the zebrafish. In Chichester; New York; John Wiley; 1999; 2007. p. 139.

51. Küchler AM, Gjini E, Peterson-Maduro J, Cancilla B, Wolburg H, Schulte-Merker S. Development of the zebrafish lymphatic system requires VEGFC signaling. Curr Biol. 2006;16(12):1244–8.

52. Kaipainen A, Korhonen J, Mustonen T, van Hinsbergh VW, Fang GH, Dumont D, et al. Expression of the fms-like tyrosine kinase 4 gene becomes restricted to lymphatic endothelium during development. Proc Natl Acad Sci. 1995 Apr 11;92(8):3566–70.

53. Karkkainen MJ, Haiko P, Sainio K, Partanen J, Taipale J, Petrova TV, et al. Vascular endothelial growth factor C is required for sprouting of the first lymphatic vessels from embryonic veins. Nat Immunol. 2004 Jan;5(1):74–80.

54. Koltowska K, Lagendijk AK, Pichol-Thievend C, Fischer JC, Francois M, Ober EA, et al. Vegfc Regulates Bipotential Precursor Division and Prox1 Expression to Promote Lymphatic Identity in Zebrafish. Cell Rep. 2015;13(9):1828–41.

55. Villefranc JA, Nicoli S, Bentley K, Jeltsch M, Zarkada G, Moore JC, et al. A truncation allele in vascular endothelial growth factor c reveals distinct modes of signaling during lymphatic and vascular development. Development. 2013 Apr 1;140(7):1497–506.

56. Ferrara N, Gerber HP, LeCouter J. The biology of VEGF and its receptors. Nat Med. 2003 Jun 1;9(6):669–76.

57. Das RN, Tevet Y, Safriel S, Han Y, Moshe N, Lambiase G, et al. Generation of specialized blood vessels via lymphatic transdifferentiation. Nature. 2022 Jun 16;606(7914):570–5.

58. Fong GH, Rossant J, Gertsenstein M, Breitman ML. Role of the Flt-1 receptor tyrosine kinase in regulating the assembly of vascular endothelium. Nature. 1995 Jul;376(6535):66–70.

59. Bahrami N, Childs SJ. Pericyte biology in zebrafish. Adv Exp Med Biol. 2018;1109:33–51.

60. Gaengel K, Niaudet C, Hagikura K, Laviña B, Muhl L, Hofmann JJ, et al. The Sphingosine-
1-Phosphate Receptor S1PR1 Restricts Sprouting Angiogenesis by Regulating the Interplay
between VE-Cadherin and VEGFR2. Dev Cell. 2012 Sep;23(3):587–99.

61. Yanagida K, Liu CH, Faraco G, Galvani S, Smith HK, Burg N, et al. Size-selective opening
of the blood–brain barrier by targeting endothelial sphingosine 1–phosphate receptor 1.
Proc Natl Acad Sci. 2017 Apr 25;114(17):4531–6.

62. Faucherre, A., Taylor, G., Overvoorde, J. *et al.* Zebrafish *pten* genes have overlapping and
non-redundant functions in tumorigenesis and embryonic development. *Oncogene* **27**,
1079–1086 (2008). https://doi.org/10.1038/sj.onc.1210730

63. Robinson RS, Woad KJ, Hammond AJ, Laird M, Hunter MG, Mann GE. Angiogenesis and
vascular function in the ovary. Reproduction. 2009 Dec;138(6):869–81.

64. Seynhaeve ALB, Oostinga D, van Haperen R, Eilken HM, Adams S, Adams RH, et al.
Spatiotemporal endothelial cell – pericyte association in tumors as shown by high resolution
4D intravital imaging. Sci Rep. 2018 Dec;8(1):9596.

65. Svingen T, Koopman P. Building the mammalian testis: origins, differentiation, and
assembly of the component cell populations. Genes Dev. 2013 Nov 15;27(22):2409–26.

66. Jokela H, Lokka E, Kiviranta M, Tyystjärvi S, Gerke H, Elima K, et al. Fetal-derived
macrophages persist and sequentially mature in ovaries after birth in mice. Eur J Immunol.
2020 Oct;50(10):1500–14.

67. Zhang Z, Huang L, Brayboy L. Macrophages: an indispensable piece of ovarian health. Biol
Reprod. 2021 Mar 11;104(3):527–38.

68. Brown HM, Robker RL, Russell DL. Development and Hormonal Regulation of the Ovarian
Lymphatic Vasculature. Endocrinology. 2010 Nov 1;151(11):5446–55.

69. Svingen T, François M, Wilhelm D, Koopman P. Three-Dimensional Imaging of Prox1-EGFP
Transgenic Mouse Gonads Reveals Divergent Modes of Lymphangiogenesis in the Testis
and Ovary. Mallo M, editor. PLoS ONE. 2012 Dec 20;7(12):e52620.

70. Hirai S, Naito M, Terayama H, Qu N, Kuerban M, Musha M, et al. The Origin of Lymphatic
Capillaries in Murine Testes. J Androl. 2012 Jul 8;33(4):745–51.

72. Armulik A, Abramsson A, Betsholtz C. Endothelial/pericyte interactions. Circ Res.
2005;97(6):512–23.

72. Wiens KM, Lee HL, Shimada H, Metcalf AE, Chao MY, Lien CL. Platelet-Derived Growth
Factor Receptor β Is Critical for Zebrafish Intersegmental Vessel Formation. Hendricks M,
editor. PLoS ONE. 2010 Jun 25;5(6):e11324.

73. Bower NI, Koltowska K, Pichol-Thievend C, Virshup I, Paterson S, Lagendijk AK, et al.
Mural lymphatic endothelial cells regulate meningeal angiogenesis in the zebrafish. Nat
Neurosci. 2017;20(6):774–83.

74. Redmer DA, Reynolds LP. Angiogenesis in the ovary. Rev Reprod. 1996;1(3):182–92.

75. Israely T, Dafni H, Granot D, Nevo N, Tsafriri A, Neeman M. Vascular remodeling and
angiogenesis in ectopic ovarian transplants: A crucial role of pericytes and vascular smooth
muscle cells in maintenance of ovarian grafts. Biol Reprod. 2003;68(6):2055–64.

76. Chiaverina G, di Blasio L, Monica V, Accardo M, Palmiero M, Peracino B, et al. Dynamic
Interplay between Pericytes and Endothelial Cells during Sprouting Angiogenesis. Cells.
2019;8(9):1–13.

77. Kofler N, Naiche LA, Zimmerman LD, Kitajewski JK. Inhibition of Jagged-Specific Notch
Activation Reduces Luteal Angiogenesis and Causes Luteal Hemorrhaging of Hormonally
Stimulated Ovaries. ACS Pharmacol Transl Sci. 2019;2(5):325–32.

78. Hall AP, Ashton S, Horner J, Wilson Z, Reens J, Richmond GHP, et al. PDGFR Inhibition
Results in Pericyte Depletion and Hemorrhage into the Corpus Luteum of the Rat Ovary.
Toxicol Pathol. 2016;44(1):98–111.

79. Hess KA, Chen L, Larsen WJ. The ovarian blood follicle barrier is both charge- and size-
selective in mice. Biol Reprod. 1998;58(3):705–11.

80. Siu MKY, Cheng CY. The Blood-Follicle Barrier (BFB) in disease and in ovarian function.
Adv Exp Med Biol. 2013;763:186–92.

81. Wang X, Qiu Z, Dong W, Yang Z, Wang J, Xu H, et al. S1PR1 induces metabolic
reprogramming of ceramide in vascular endothelial cells, affecting hepatocellular carcinoma
angiogenesis and progression. Cell Death Dis. 2022 Sep 6;13(9):768.

82. Hübner, Kathleen et al. "Wnt/β-catenin signaling regulates VE-cadherin-mediated
anastomosis of brain capillaries by counteracting S1pr1 signaling." *Nature
communications* vol. 9,1 4860. 19 Nov. 2018, doi:10.1038/s41467-018-07302-x

83. Pascuali N, Scotti L, Oubiña G, de Zúñiga I, Gomez Peña M, Pomilio C, et al. Platelet-
derived growth factor B restores vascular barrier integrity and diminishes permeability in
ovarian hyperstimulation syndrome. Mol Hum Reprod. 2020 Aug 1;26(8):585–600.

84. Di Pietro M, Pascuali N, Parborell F, Abramovich D. Ovarian angiogenesis in polycystic
ovary syndrome. Reproduction. 2018;155(5):R199–209.

85. Zaidi J, Campbell S, Pittrof R, Kyei-Mensah A, Shaker A, S.Jacobs H, et al. Contraception:
Ovarian stromal blood flow in women with polycystic ovaries—a possible new marker for
diagnosis? Hum Reprod. 1995 Aug 1;10(8):1992–6.

86. Yan YL, Desvignes T, Bremiller R, Wilson C, Dillon D, High S, et al. Gonadal soma controls
ovarian follicle proliferation through Gsdf in zebrafish. Dev Dyn. 2017 Nov;246(11):925–45.

87. Fantin A, Vieira JM, Gestri G, Denti L, Schwarz Q, Prykhozhij S, et al. Tissue macrophages
act as cellular chaperones for vascular anastomosis downstream of VEGF-mediated
endothelial tip cell induction. Blood. 2010 Aug 5;116(5):829–40.

88. Eubank TD, Galloway M, Montague CM, Waldman WJ, Marsh CB. M-CSF Induces Vascular
Endothelial Growth Factor Production and Angiogenic Activity From Human Monocytes 1. J
Immunol. 2003 Sep 1;171(5):2637–43.

89. Rymo SF, Gerhardt H, Wolfhagen Sand F, Lang R, Uv A, Betsholtz C. A Two-Way
Communication between Microglial Cells and Angiogenic Sprouts Regulates Angiogenesis
in Aortic Ring Cultures. PLOS ONE. 2011 Jan 10;6(1):e15846.

90. Martin P, Gurevich DB. Macrophage regulation of angiogenesis in health and disease. Spec
Issue Muscle Growth Regen Kathryn Wagner Spec Issue Role Macrophages Tissue
Homeost Benedicte Chazaud. 2021 Nov 1;119:101–10.

91. Sunderkötter C, Goebeler M, Schulze-Osthoff K, Bhardwaj R, Sorg C. Macrophage-derived
angiogenesis factors. Pharmacol Ther. 1991 Jan 1;51(2):195–216.

92. Wu R. Macrophage contributions to ovarian function. Hum Reprod Update. 2004 Mar
1;10(2):119–33.

93. Ono Y, Nagai M, Yoshino O, Koga K, Nawaz A, Hatta H, et al. CD11c+ M1-like
macrophages (MΦs) but not CD206+ M2-like MΦ are involved in folliculogenesis in mice
ovary. Sci Rep. 2018 Dec;8(1):8171.

Figure Legends:

Figure 1. Vascular endothelial cells in the bipotential gonad. (A-E) Representative images

showing the defined 3 categories for describing the relationship between vascular endothelial

cells and germ cells in the biopotential gonad: "No Contact" (**A**), "Proximal" (**B-C**; open

arrowhead, Movie 1), and "Contact" (**D-E**; red arrow, Movie 2). All images are oriented with

anterior (A) to the left and posterior (P) to the right. (**F**) The main vessel supplying blood to the

gonad, the gonadal artery (white arrowheads), originates from the swim bladder artery (red

arrowheads). (**G**) A schematic representing the GA and SBA in the context of the body cavity.

VEC = Vascular Endothelial Cell, AC = Anterior Chamber, GA = Gonadal Artery, SBA = Swim

Bladder Artery, PD = Pneumatic Duct, AC = Anterior Chamber, PC = Posterior Chamber. Scale

= 100 μm.

Figure 2. Vascular endothelial cells, lymphatic endothelial cells, and macrophage are

present in the gonad when fish reach a standard length of 6.5mm. (**A**) At each age,

individual fish from multiple spawns were collected to determine if there was contact between

the cell of interest and the germ cells. Vascular endothelial cells (red circle), lymphatic

endothelial cells (green triangle), pericytes (blue square), and sometimes macrophage (purple

diamonds) arrive concurrently to the gonad between 15 and 20 dpf. (**B**) Each fish was

measured prior to dissection to determine its length Separate cohorts of fish developed at

different rates due to environmental conditions which led to variability in the initial point of

contact. In juvenile fish, length can be used as a more accurate means of assessing

developmental progression as compared to age. Contact between endothelial and germ cells

occurred when the fish is approximately 6.5mm in length. The full list of replicates, individuals,

average length, and observations can be found in **Supplemental Table S1**.

Figure 3. Vascularization in the zebrafish ovary. (A-B') The two lobes of the zebrafish ovary

are highly vascularized by 40 dpf. **(B-B')** A high magnification of one ovary shows how large

arteries quickly divide into smaller vessels surrounding oocytes. **(C)** As oocytes mature at 60

dpf, they will each be surrounded by at least one vessel (see Movie 3). *kdrl* and *sox9b* are,

respectively, green and magenta in A, and B. *kdrl* is pseudo-colored black in A' and B'. *kdrl*,

Vasa, and DNA are, respectively, green, magenta, and grey in C. Scale = 100µm.

Figure 4. Vascularization in the zebrafish testis. (A-B') The testis is significantly smaller than

the ovary at 40 dpf, yet is also densely packed with vessels. **(B-B')** High magnification of

testicular vasculature at 40 dpf. **(C)** Vessels remain outside of the somatic cells of the 60 dpf

testis and surround lobules of maturing sperm (see Movie 4). *kdrl* and *sox9b* are, respectively,

green and magenta in A and B. *kdrl* is pseudo-colored black in A' and B'. *kdrl*, Vasa, and DNA

are, respectively, green, magenta, and grey in C. Scale = 100µm.

763 **Figure 5. Endothelial gene expression in 40 dpf ovary. (A)** Expression of vascular

764 endothelial cell genes *kdrl, fli1*, and *flt1* occurred in a majority of the endothelial cells. (**B**)

765 lymphatic endothelial genes *lyve1b*, *flt4*, and *prox1* were expressed in a smaller subset of cells.

766 (**C**) There was no overlap in expression of *kdrl* (red) and *lyve1b* (green) by single-cell

767 sequencing. (**D**) There was overlap in expression of *flt4* (red) and *lyve1b* (green) suggesting that

768 lymphatic endothelial cells originate canonically.

770 **Figure 6. *kdrl* and *mrc1a* are co-expressed in the 20 dpf ovary.** (**A**) Maximum intensity

771 projection of the 20 dpf gonad (germ cells = blue, gonad outlined) showing overlap in *kdrl*

772 (magenta) and *mrc1a* (green) expression. (**B**) Single channel showing *kdrl* expression alone

773 suggests that *kdrl* is expressed in all cells. (**C**) Single channel showing *mrc1a* expression is

774 inversely correlated with the strength of *kdrl* expression. (**D**) Orthogonal slice of the maximum

775 intensity projection shows that the expression of both markers is overlapping within cells. (**E**)

776 Co-localization analysis of the cells circled in red in (**F**) showing expression of *kdrl* and *mrc1a* is

777 highly correlated. Scale = 20 μm.

779 **Figure 7. The *pdgfrb+* expression domain encompasses multiple cell types in the ovary**.

780 Using previously published data from single-cell sequencing in the embryo, we compared

781 expression profiles of *pdgfrb+* cells (blue) to cluster-39 (pink), which is a subset of *pdgfrb+* cells

782 with the highest *pdgfrb* expression and low expression of other known cell type markers. We

783 further compared larval pericytes to 40 dpf ovarian stromal cells (yellow), which are defined as

784 non-germ, non-somatic, non-immune, and non-endothelial cells, and ovary-specific pericytes

785 (purple) using single-cell data from the zebrafish ovary. 215 genes, including *plp1b,* were found

786 to be unique to ovarian pericytes (purple only), and 3 genes were expressed only in cluster 39

787 as well as ovary-specific pericytes (pink and purple overlap). *notch3* was expressed in all four

788 cell groups (yellow, purple, pink, and blue overlap).

Figure 8. *pdgfrb* positive cells elongate in close association with vascular endothelial tip

cells. (**A**) Maximum intensity projection of all channels of the 20 dpf gonad show pericytes,

labeled with *pdgfrb:EGFP* (green). (**B**) Pericytes (pseudo-colored black) extend protrusions

(arrowheads) between two existing vessels. *pdgfrb:EGFP* cells have classic pericyte

morphology and are located at the division of vessels supporting the conclusion that they are

pericytes (arrowheads). (**C**) *kdrl:DsRed* vessels (pseudo-colored black) are associated with an

endothelial tip cell. (**D**) DNA (pseudo-colored black). See Movie 5. Scale = 20 μm.

Figure 9. Tissue-resident macrophage engulf cells during gonad differentiation. (**A**)

Macrophage were found in contact with germ cells as early as 12 dpf. (**B**) At 20 dpf,

macrophage are abundant in the gonad. (**C**) Macrophage (magenta) were observed removing

oocyte material (green) and DNA (white) from the former bipotential gonad (white arrowheads,

Movie 6). (**D**) *mpeg1:GAL4; UAS:Kaede* expressing macrophage were photoconverted from

green to red (pseudocolored magenta) at 12 dpf. 24 hours after conversion green and red

macrophage could be found in the gonad (blue arrow, outlined). (**E**) Some macrophage in the

gonad were only magenta (red arrowhead), indicating that they are resident to the gonad during

this period. (**F**) In the green channel, we can observe that some macrophage migrate to the

gonad within the 24 hour window post-photoconversion (green arrows). See Movie 7 and 8.

Scale listed.

Movie Legends:

Movie 1: 3D rendering of "Proximal" association between vascular endothelial cells and

germ cells. In the center of the plane, the vascular endothelial cells elongate towards the germ

cells at 12 dpf. DNA from either the somatic gonad or the body wall is present between the

vessel and germ cells. *kdrl* (vascular endothelial cells, green), Vasa (germ cell, magenta), DNA

(Blue).

Movie 2: 3D rendering of "Contact" between vascular endothelial cells and germ cells

from Figure 1D. At ~9 seconds the germ cells (green) contact with the vascular endothelial

cells (magenta) are in close view. At ~16 seconds, vessels running vertically are in view, these

exist in the body wall between somites. The anterior is to the right, and posterior is to the left of

the video. *kdrl* (vascular endothelial cells, magenta), *fli1* (vascular endothelial cells, nuclear

green), Vasa (germ cell, green).

Movie 3: Z-stacks through the 60 dpf ovary (Figure 3). *kdrl* (green), *sox9b* (magenta), DNA

(gray). As oocytes mature, they will each be surrounded by at least one vessel.

Movie 4: Z-stacks through the 60 dpf testis (Figure 4). *kdrl* (green), *sox9b* (magenta), DNA

(gray). Vessels remain outside of the somatic cells of the testis and surround lobules of

maturing sperm.

Movie 5: 3D rendering of pericytes association with endothelial tip cells (Figure 8).

Pericytes = *pdgfrb:EGFP* (green), vascular endothelial cells = *kdrl:DsRed* (magenta), DNA =

Hoechst (blue)

Movie 6: 3D rendering and Z-stacks of macrophage engulfing germ cell material (Figure

9C). Macrophage = *mpeg:Kaede* (magenta), germ cells = *piwil1:EGFP* (green), DNA = Hoechst

(blue).

Movie 7: Z-stacks of photoconverted macrophage (Figure 9C). 24 hours after conversion

green and red macrophage could be found in the gonad (blue arrow) while some macrophage in

the gonad were only magenta (red arrowhead). In the green channel, we can observe that some

macrophage migrate to the gonad within the 24 hour window post-photoconversion (green

arrows).

Movie 8: Movement of a macrophage at 13 dpf over one hour. Within one hour some

macrophage are highly dynamic (arrows) while others remain in place.

Supplemental Figure Legends:

Supplemental Figure S1: Single channel images of contact between germ cells and vascular

endothelial cells.

Supplemental Figure S2: Ovarian vasculature in sectioned tissue. (**A, C, E, G, I**) *kdrl*

expressing vessels (green) show the relationship between the vasculature and germ cells (*piwi*,

magenta) in the adult zebrafish ovary. **(B, D, F, H, J)**. Hoechst staining (DNA, green) marks the

somatic and endothelial cells in the ovary relative to the germ cells (*piwi*, magenta). Note that

the more immature oocytes are in indirect contact with the vasculature. Scale: 100 µm.

Supplemental Figure S3: All channels from Figure 8 in of the 20 dpf zebrafish gonad. A.

vascular endothelial cells= kdrl:DsRed (magenta, panel B), lymphatic endothelial cells =

mcr1a:EGFP (green, panel C), germ cells = Vasa (blue, panel D), DNA = Hoechst (white, panel

E).

Supplemental Figure S4: Photoconverted area of *mpeg:Kaede*. (A) at 12 dpf the total area

which was photo converted and the 10x view of the area pre- and post-conversion. **(B)** After 24

hours, at 13 dpf, the area of *mpeg:Kaede* has expanded and there are both green and red cells

present. All images are of different individuals to maximize survival during live imaging. The box

denotes the area in which the gonad is located in the body cavity. * denote the gut which is

auto-fluorescent both green and red because of the food.

Supplemental Table Legends:

Supplemental Table S1: Number of individuals counted in each category of cell type being

assessed.

Supplemental Table S2: Highly expressed genes in each pericyte category: larval pericytes

(Cluster 39) (Sheet 1), pdgfrb+ cells (Sheet 2), ovarian non-germ Cells (Sheet 3), ovary

pericytes (Sheet 4), and comparisons between each group for the Venn diagram (Sheet 5).

Figures:

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.

Figure 8.

Figure 9.

Supplemental Figures:

Supplemental Figure S1:

kdrl:DsRed = Magenta
fli1:nEGFP = Green nuclei
Vasa antibody staining = Green whole cell
DNA = White

RED CHANNEL
kdrl:DsRed
Pseudo-colored Magenta

GREEN CHANNEL
fli1:nEGFP = nuclei
Vasa antibody staining = whole cell

908 **Supplemental Figure S2:**

911 **Supplemental Figure S3:**

A

kdrl:DsRed
mrc1a:EGFP
Vasa
DNA

B

kdrl:DsRed

C

mrc1a:EGFP

D

Vasa

E

DNA

Supplemental Figure S4: