Treponema pallidum subsp. _pallidum_ with an Artificially Impaired TprK Antigenic

Variation System is Attenuated in the Rabbit Model of Syphilis

Emily Romeis[1], Nicole A. P. Lieberman[2], Barbara Molini[1], Lauren C. Tantalo[1], Benjamin

Chung[2], Quynh Phung[2], Carlos Avendaño[2], Anastassia Vorobieva[3,4], Alexander L. Greninger[2,5]

and Lorenzo Giacani[1,6*]

[1]Department of Medicine, Division of Allergy and Infectious Diseases, University of

Washington, Seattle, WA, USA

[2]Department of Laboratory Medicine and Pathology, University of Washington, Seattle, WA,

USA

[3]VIB-VUB Center for Structural Biology, VIB, Brussels, Belgium

[4]Structural Biology Brussels, Vrije Universiteit Brussel, Brussels, Belgium

[5]Vaccine and Infectious Disease Division, Fred Hutchinson Cancer Research Center, Seattle,

WA, USA

[6]Department of Global Health, University of Washington, Seattle, WA, USA

[*]Corresponding Author: giacal@u.washington.edu

Short title: Impairing the _T. pallidum_ TprK system

Abstract

Background. The TprK protein of the syphilis agent, *Treponema pallidum* subsp.
pallidum (*T. pallidum*), undergoes antigenic variation in seven discrete variable (V) regions via
non-reciprocal segmental gene conversion. These recombination events transfer information
from a repertoire of 53 silent chromosomal donor cassettes (DCs) into the single *tprK* expression
site to continually generate TprK variants. Several lines of research developed over the last two
decades support the theory that this mechanism is central to *T. pallidum*'s ability for immune
avoidance and persistence in the host. Structural and modeling data, for example, identify TprK
as an integral outer membrane porin with the V regions exposed on the pathogen's surface.
Furthermore, infection-induced antibodies preferentially target the V regions rather than the
predicted β-barrel scaffolding, and sequence variation abrogates the binding of antibodies
elicited by antigenically different V regions. Here, we engineered a *T. pallidum* strain to impair
its ability to vary TprK and assessed its virulence in the rabbit model of syphilis.

Principal findings. A suicide vector was transformed into the wild-type (WT) SS14 *T.
pallidum* isolate to eliminate 96% of its *tprK* DCs. The resulting SS14-DC^{KO} strain exhibited an
in vitro growth rate identical to the untransformed strain, supporting that the elimination of the
DCs did not affect strain viability in absence of immune pressure. In rabbits injected
intradermally with the SS14-DC^{KO} strain, generation of new TprK sequences was impaired, and
the animals developed attenuated lesions with a significantly reduced treponemal burden
compared to control animals. During infection, clearance of V region variants originally in the
inoculum mirrored the generation of antibodies to these variants, although no new variants were
generated in the SS14-DC^{KO} strain to overcome immune pressure. Naïve rabbits that received
lymph node extracts from animals infected with the SS14-DC^{KO} strain remained uninfected.

Conclusion. These data further support the critical role of TprK in *T. pallidum* virulence and persistence during infection.

Author Summary

Syphilis is still endemic in low- and middle-income countries, and it has been resurgent in high-income nations, including the U.S., for years. In endemic areas, there is still significant morbidity and mortality associated with this disease, particularly when its causative agent, the spirochete *Treponema pallidum* subsp. *pallidum* (*T. pallidum*) infects the fetus during pregnancy. Improving our understanding of syphilis pathogenesis and *T. pallidum* biology could help investigators devise better control strategies for this serious infection. Now that tools to genetically manipulate this pathogen are available, we can engineer *T. pallidum* strains lacking specific genes or genomic regions known (or believed) to be associated with virulence. This approach can shed light on the role of the ablated genes or sequences in disease development using loss-of-function strains. Here, we derived a knockout (KO) *T. pallidum* mutant (SS14-DC^{KO}) impaired in its ability to undergo antigenic variation of TprK, a protein that has long been hypothesized to be central in evasion of the host immune response and pathogen persistence during infection. When compared to the WT isolate, which is still capable of antigenic variation, the SS14-DC^{KO} strain is significantly attenuated in its ability to proliferate and to induce early disease manifestations in infected rabbits. Our results further support the importance of TprK antigenic variation in syphilis pathogenesis and pathogen persistence.

Introduction

Syphilis is a chronic sexually transmitted infection that still represents a concern for public health as it causes significant morbidity and mortality worldwide. The World Health

Organization (WHO) estimated that syphilis global incidence is between 5.6 to 11 million new

cases every year, while global prevalence ranges between 18 to 36 million cases (1, 2). Although

most cases occur in low- and middle-income countries where the disease is endemic, syphilis

rates have been on the rise for years in high-income nations, primarily among men who have sex

with men (MSM), even though the gap in incidence between MSM and the heterosexual

population is gradually reducing (3-8). In the US, the rate of early syphilis in 2020 (12.7 cases

per 100,000 population) represented a 504% increase compared to the rate reported in 2000 (2.1

cases per 100,000 population) (3). Furthermore, about 1.4 million pregnant women a year

acquire syphilis worldwide. Congenital syphilis (CS), resulting from the ability of *Treponema*

pallidum subsp. *pallidum* (*T. pallidum*) to colonize and cross the placenta, causes an estimated

305,000 annual cases of fetal loss or stillbirth and 215,000 infants born prematurely and/or with

clinical signs of infection (9-11).

In untreated subjects, primary and secondary syphilis generally manifests in distinct

clinical stages, followed by a period of asymptomatic latency that, in most cases, will last for the

patient's lifetime (12). Data preceding the introduction of penicillin for syphilis treatment

showed that in about 30% of patients, the disease would reactivate after years to decades from

the onset of latency and progress to its tertiary stage and its most serious manifestations (13).

Key to this decades-long persistence is the pathogen's ability to evade the robust host immune

response that develops during infection. This central feature of *T. pallidum* virulence has long

been attributed to antigenic variation of the TprK (Tp0897) protein (12). This putative integral

outer-membrane porin (14) harbors seven discrete variable regions (V1-V7), each predicted by

AlphaFold2 (https://AlphaFold.ebi.ac.uk/) to form an external loop at the host-pathogen interface

(Fig.1). Sequence variability within the V regions is generated through non-reciprocal segmental

gene conversion, a process that transfers information contained in 53 silent chromosomal donor

cassettes (DCs) into the *tprK* expression site (15, 16), located ~175 kbp away from the DCs.

Most of the DCs (51/53; 96%) are located downstream of the *tprD* gene (*tp0131*) at nucleotide

(nt) position 148,435-152,400, while the remaining two DCs are downstream of the *tp0136* gene

at position 159,499-159,576, based on the SS14 strain reference genome sequence

(NC_021508.1/CP004011.1) (17). In a previous study by Addetia *et al.* (17) it was estimated that

the total *tprK* diversity that could be generated by the combined usage of all DCs would range

between 10^{13} - 10^{15} full-length individual TprK sequences (17).

Aside from sequence variability and potential for generation of quintillions of unique

sequences, the role of TprK in immune evasion and pathogen persistence is supported by the

evidence that a) sequence diversity accumulates over time during experimental infection (18), b)

infection-induced antibodies preferentially target the TprK V regions rather than the predicted β-

barrel scaffolding of the protein (19), c) sequence variation abrogates binding of antibodies

raised to antigenically different V regions (20), d) TprK variants accumulate more rapidly in

immunocompetent hosts compared to immunosuppressed rabbits as the result of immune

selection (21, 22) and, lastly, e) TprK sequences found in disseminated secondary lesions in the

rabbit model differ from those in the original inoculum, supporting that secondary lesions

contain treponemes that avoided the initial round of immune clearance associated with the

resolution of the primary lesion(s) (23).

Recently, Romeis *et al*. (24) described a procedure that successfully allowed the

replacement of a specific *T. pallidum* chromosomal region with a kanamycin resistance (*kan*[R])

cassette. This approach uses a suicide vector with the *kan*[R] cloned between homology arms

matching the regions upstream and downstream of the sequence to be deleted. This work

provided the first experimental evidence that *T. pallidum* could be genetically altered and opened

the gateway for systematic derivation of KO mutants to assess their fitness and virulence *in vivo*

and shed light on the role of specific genes and or genomic regions in syphilis pathogenesis.

Although a *tprK*^{KO} strain could not be derived, to further explore the role of TprK in *T. pallidum*

persistence, we ablated the 51 DCs situated downstream of the *tprD* gene in the SS14 *T.*

pallidum strain, which resulted in an impaired TprK antigenic variation system. We then

evaluated disease progression and the pathogen's ability to proliferate in rabbits infected with the

mutant strain (renamed SS14-DC^{KO}) in comparison to animals infected with the WT strain, still

fully capable of varying TprK.

Results

SS14-DC^{KO} mutant treponemes are not impaired in their ability to grow *in vitro* and *in vivo*

in an immune-privileged site.

130   Treponemes transformed with the p*DC*arms-47p-*kan*^{R} suicide vector, designed to

eliminate the DCs downstream of *tprD* through a double cross-over recombination event, were

selected by propagation in selective media (Fig.2A). As a control, the parent SS14 strain was

also propagated in both normal and selective media in parallel to the transformed strain. Due to

the ~44 hours generation time of the SS14 strain, *T. pallidum* cells needed to be sub-cultured

every 14 days until week 13 post-transformation (Passage #8; Fig.2A), and weekly thereafter.

Transformed SS14-DC^{KO} treponemes had significantly recovered by 4 weeks post-

transformation (Fig.2A) at the time passage #3 was harvested, and approximately 10 treponemal

cells per microscopic field (corresponding to ~10^7 *T. pallidum* cells/ml) could be counted at this

time. The cell density of transformed treponemes increased by week 10 and remained steady

with harvests of ~10^8 *T. pallidum* cells/ml (Fig.2A). Wild-type parent SS14 cells propagated in

selective media could not be seen by dark-field microscopy (DFM) at any time and sub-culturing

was stopped at week 8 (Fig.2A). Propagation of the WT parent strain was halted at week 18

(Fig.2A). DNA extracted from treponemes harvested at passage #8 post-transformation was used

for qualitative amplifications (primers in Table 1) designed to confirm integration of the *tp0574*

promoter-*kan*^{R} sequence where the DC-containing region used to be, as well as lack of residual

transformation vector or DNA from the parent strain. Results (Fig.2B) showed that no

amplification occurred in treponemal cultures with primers specific for the suicide vector

backbone (subpanel #1), confirming lack of residual plasmid DNA. As expected, amplification

of the *tp0574* gene was positive for both the parent SS14 and the SS14-DC^{KO} strain templates

(subpanel #2), confirming the presence of treponemal DNA in both cultures. Furthermore,

amplification of the target region with the reverse primer annealing downstream of the right

homology arm (hence not cloned into the suicide vector) and a forward primer annealing to the

kan^R gene, respectively (subpanel #3), yielded positive amplification only with template from the

SS14-DC^{KO} strain, confirming that integration of the *tp0574* promoter-*kan*^{R} sequence occurred

where expected. Amplification with both primers outside of the homology arms yielded

amplicons of different sizes (subpanel #4; 2,982 bp for SS14-DC^{KO} and 5,980 bp for the parent

strain), confirming deletion of the DCs and lack of parent SS14 DNA in the mutant strain

culture. Lastly, amplification with primers specific for the *kan*^{R} only amplified the SS14-DC^{KO}

and the vector control template (subpanel #5).

Rabbit intratesticular (IT) inoculation with either the WT or the SS14-DC^{KO} strain was

also used to assess whether the mutant strain would be impaired in its ability to grow *in vivo* and

induce orchitis. The time to orchitis in WT- and SS14-DC^{KO}-infected animals was identical (21

days post-inoculation) and, when harvested, both rabbits yielded comparable treponemal

concentrations ($\sim$10^8/ml).

Altogether, these data confirmed that a) the *kan*^{R} gene was integrated into the SS14-

DC^{KO} strain genome in place of the targeted 51 DCs, that b) no residual suicide plasmid DNA or

parent treponemes contaminated the SS14-DC^{KO} culture, that c) 96% of the DCs are missing in

the mutant strain and lastly, that d) despite this deletion, SS14-DC^{KO} treponemes are not

impaired in their ability to grow *in vitro* and *in vivo* compared to the WT strain.

Table 1. Primers used in this study

Targets; application	Forward (F) and reverse (R) primer sequences (5'-3'), and probe (P) if applicable	Amplicon length (bp)	Sub-panel # in Fig.2B
pUC57 vector primers; sequencing, assessing plasmid carry-over	(F) TAAAACGACGGCCAGTGAAT (R) GACCATGATTACGCCAAGC	2,967	1
tp0574; qualitative PCR	(F) TGTGGCTCGTCTCATCATGA (R) CTGGGCCACTACCTTCGCAC	313	2
kan^R (F primer), right *DC* flanking region (R primer); qualitative PCR and sequencing (F)	(F) GAGCCATATTCAACGGGAGA (R) CTTCTCCTTCGCCCTCGAC	1,799	3
Left *DC* flanking region (F primer), Right *DC* flanking region (R primer); qualitative PCR	(F) CACAGAAGGGCAGCAGTACA (R) CTTCTCCTTCGCCCTCGAC	2,982 (SS14-DC^{KO}) or 5,980 (WT)	4
kan^R; qualitative PCR	(F) GAGCCATATTCAACGGGAGA (R) ATTCCGACTCGTCCAACATC	663	5
Left *DC* flanking region (F primer), *kan*^{R} (R primer); qualitative PCR and sequencing	(F) CACAGAAGGGCAGCAGTACA (R) ATTCCGACTCGTCCAACATC	1,846	
kan^R; qualitative PCR	(F) GAGCCATATTCAACGGGAGA (R) ATTCCGACTCGTCCAACATC	663	
tp0897 (tprK); RT-qPCR	(F) GAAAATCGCCTGTGCCCTAC (R) GGTTCCCCACGTTTAGTTAG	80	
tp0574; qPCR, RT-qPCR	(F) CAAGTACGAGGGGAACATCG (R) TGATCGCTGACAAGCTTAGG (P) 6FAM-CGGAGACTCTGATGGATGCTGCAGTT-NFQMGB[1]	132	
tp0127; ddPCR	(F) GCGTGTATGGAGAAGCATTG (R) CGCCTGATAGCCTATATCCAC (P) HEX-AGATATGGT/ZEN/ACTACGCGCCGC-3IABkFQ[2]	139	
kan^R; ddPCR	(F) CACTCAGGCGCAATCAC (R) CCAGACTTGTTCAACAGGC (P) FAM-	91	

	ACGGTTTGGTTGATGCGAGTGATTT-BkFQ[3]	
tp0001 (*dnaA*); ddPCR,	(F) CTCATGGAAATACTGCTCC (R) CGGATACAAAGTTCTCGAAG (P) FAM-AGCTTTCACCCCGACCTGAAC-BkFQ[3]	135
tp0574 promoter; sequencing	(F) AGCGGATCCTCCCAAAAAGA (R) GATTACACCTCCGTATAGAG	N/A
pUC57 vector primers; sequencing, assessing plasmid carry-over	(F) TAAAACGACGGCCAGTGAAT (R) GACCATGATTACGCCAAGC	2,967

172   [1]Minor groove binder nonfluorescent quencher
173   [2]3′ Iowa Black fluorescence quencher
174   [3]Black hole fluorescence quencher
175

Genome sequencing and droplet digital PCR confirm deletion of 96% of the DCs

and *tprK* expression in mutant treponemes.

At passage #2-5 (Week 2-6 post-transformation in Fig.2A), ahead of genome sequencing,

DNA extracted from SS14-DC^{KO} and parent SS14 cells was processed for quantitative droplet

digital PCR (ddPCR) to calculate copy number of the *kan*^{R} and *tp0127* (contained within the

eliminated region containing the DCs) genes, normalized to the copy number of *dnaA* gene

(*tp0001*; present in both parent and transformed treponemes). The normalized ratios of either

kan^R or *tp0127* were expected to be either zero or ~1.0 depending on the strain analyzed. When

the parent SS14 DNA was used as control, the *tp0127*:*dnaA* was near 1.0 over the four analyzed

passages (Fig.3A), while the *kan*^{R}:*dnaA* ratio was zero due to lack of the *kan*^{R} gene in this strain

(Fig.3A). In the SS14-DC^{KO} strain, the *kan*^{R}:*dnaA* ratio neared 1.0 when DNA from passage #3

and #4 was analyzed (Fig.3A), and was 1.0 for passage #5, while the *tp0127*:*dnaA* ratio was zero

(due to the absence of *tp0127*), with the exception of passage #2. This result was however

expected due to the presence of residual parent SS14 DNA, given that the culture had not been

passaged yet after transformation (Fig.3A). These data reiterated that a) integration of the *kan*^{R}

gene occurred in the DC locus and supported that no extra copies of the *kan*^{R} gene existed

outside or within the SS14-DC^{KO} strain genome. As previously discussed (24), the slightly <1.0

results in some of these assays are likely due to the proximity of the *dnaA* gene (nt position 4-

1,395 in the SS14 genome) to the chromosomal origin of replication.

To ascertain that DC deletion did not affect *tprK* transcription, cDNA generated from the

SS14-DC^{KO} and WT strain RNA, was used to assess transcription of the *tprK* (*tp0897*) and

tp0574 genes (used as housekeeping gene; primers in Table 1). Message quantification (Fig.3B)

showed that, like in other *T. pallidum* strains (25), *tprK* is poorly expressed in the SS14 strain

compared to *tp0574*, but its expression does not differ significantly between the WT and mutant

strains.

Whole genome sequencing (WGS) performed on SS14-DC^{KO} treponemes from passage

#9 confirmed the absence of reads for the *tprK* DCs in the SS14-DC^{KO} strain following assembly

to the SS14 reference genome (Fig.3C). Assembly using a reference genome where the DCs

were replaced *in silico* with the *kan*^{R} cassette (KO reference) was consistent with the

replacement of the 51 DCs downstream of *tprD* with the *tp0574* promoter-*kan*^{R} sequence

(Fig.3D). Because sequencing used enrichment probes specific for multiple WT *T. pallidum*

genomes, coverage of the *kan*^{R} gene (Fig.3D) was slightly lower than the average compared to

the rest of *T. pallidum* genome. A search for reads matching the suicide vector backbone yielded

no results, confirming lack of residual suicide vector DNA. These results further confirmed

elimination of 96% of the DCs (schematically represented in Fig.3E) in the SS14-DC^{KO} strain

and their replacement with the *kan*^{R} cassette. Furthermore, these data showed that despite this

deletion, mutant treponemes are not impaired in their ability to express *tprK* compared to the WT

strain. The WT parent strain propagated in parallel to the mutant strain was also sequenced as a

control. As expected, in the WT strain the DC locus is intact (Fig.4F), while a gap is visible

where the kan^R cassette is located when reads from the WT strain were assembled to the KO

reference (Fig. 4G).

Early lesion development is attenuated in rabbits infected with treponemes lacking most

DCs.

Differences in early lesion development were assessed in New Zealand White rabbits

infected intradermally (ID) in multiple sites on their shaved backs with either the WT or mutant

strain. Following infection of two groups of four rabbits each, animals were regularly monitored

to measure lesion diameter. Samples were collected every ~10 days for quantification of

treponemal burden (both biopsies and aspirates) and *tprK* deep sequencing (biopsies only).

Serum samples were collected weekly for syphilis serology using both a non-treponemal

(Venereal Disease Research Laboratory; VDRL) and treponemal (*Treponema pallidum* particle

agglutination; TPPA) test. The diameter of primary lesions in rabbits infected with the WT strain

became significantly larger ($p<0.0001$) compared to the lesions induced by the SS14-DC^{KO} strain

starting at day 9 post-inoculation (Fig.4A). Lesion size in SS14-DC^{KO}-infected rabbits plateaued

at day 50 post-inoculation with an average diameter of 7.0 mm. In control animals, lesion size

steadily increased throughout the experiment (Fig.4A) up to day 65 post-inoculation ($p>0.05$ for

lesion diameter values between day 65 and day 72 in WT-infected animals) with mean diameters

of 26.75 mm (day 65) and 30.3 mm (day 72), respectively. Pictures of the four animals still in the

study at day 72 post-inoculation (Fig.4B) show that in addition to being significantly enlarged,

all lesions in the control animals had progressed to an ulcerated state, with large central necrotic

areas. In animals infected with the SS14-DC^{KO} strain, ulceration was not evident. Circled areas

correspond to lesions that were not biopsied at the time the picture was taken.

In SS14-DC^{KO}-infected rabbits, treponemal burden is reduced and development of humoral immunity to infection is delayed.

Treponemal burden in lesion biopsies and needle aspirates was assessed by qPCR and DFM, respectively. In both cases (Fig.5A, B), significantly more treponemes were detected in samples from the control animals compared to those from rabbits infected with the SS14-DC^{KO} strain. Although qPCR (Fig.5A) detected *T. pallidum* in all tested samples, treponemal burden in lesions induced by the mutant strain decreased by one (day 9-37), two (day 65-79), or three orders of magnitude (day 37-58) between the control and test samples. As expected, treponemal burden decreased in lesions from control animals over time due to immune clearance (13). In contrast, the overall burden in lesions from rabbits infected with the SS14-DC^{KO} strain remained stably low (330.6 copies/reaction in average) in samples collected at day 37-79 post-inoculation. No treponemes were detected by DFM of needle aspirates collected from SS14-DC^{KO}-infected rabbits at any time point, while treponemes could be detected in lesions from WT-infected animals from day 37-72 post-inoculation (Fig.5B), suggesting that the treponemal burden in samples from test animals was always below the limit of detection of this assay unless rabbits were infected with the WT strain.

VDRL tests performed on serum samples collected over time showed that the WT-infected animals seroconverted earlier and reached higher titers in a shorter time compared to the SS14-DC^{KO} rabbits (Fig.5C). In rabbits infected with the mutant strain, only one rabbit had seroconverted by day 37 post-inoculation (1:2 titer). This titer, however, could not be reproduced at day 44 post-infection (Fig.5C). VDRL titers between the two rabbit groups remained significantly different from day 58 to the end of the experiment. TPPA titers were significantly

different between control and test rabbits only up to day 23 post-infection. Together, these data

support impairment of the SS14-DC^{KO} strain's ability to proliferate in dermal lesions, leading to

an overall lower treponemal burden and to a delayed development of VDRL and TPPA

antibodies.

Rabbits used as recipients of lymph nodes from SS14-DC^{KO}-infected rabbits did not develop

syphilis infection.

Immediately after euthanasia, popliteal lymph nodes were harvested from each of the ID-infected

rabbits and transferred via IT injection into a naïve rabbit to assess *T. pallidum* presence. A total of four

animals (two per group) were picked randomly and euthanized on day 44 post-infection. The remaining

rabbits were euthanized on day 79 post-infection. Rabbits were monitored weekly for orchitis

development and bled monthly to assess seroconversion, unless orchitis developed and was confirmed

by visualization of treponemes in testicular aspirates, at which point the animal was deemed positive and

euthanized. One animal that received lymph nodes from the control rabbits (Table 2) had developed

orchitis (confirmed by treponemal presence in a needle aspirate) at day 33 post-inoculation, while two

more in the same group developed orchitis (Also DFM-positive) at day 35 post-inoculation. The fourth

rabbit in the group developed orchitis at day 74 post-inoculation (Table 2). In contrast, none of the

recipients of lymph node extracts from rabbits infected with the SS14-DC^{KO} strain developed orchitis or

seroconverted during the 3-month observation period, supporting absence of treponemal cells in these

lymph node extracts at the time the RIT was performed.

287 **Table 2. Rabbit infectivity test results**

Infecting strain	RIT day	Donor Rabbit[1]	Time to positive orchitis/DFM[2]	Month 1 serology (day 30 post-RIT)		Month 2 serology (day 60 post-RIT)		Month 3 serology (day 90 post-RIT)	
				VDRL[3]	TPPA[4]	VDRL[3]	TPPA[4]	VDRL[3]	TPPA[4]
SS14-WT	Day 44 post-IDI	505	74 days	NR	NR	NP	NP		
		512	33 days	NP	NP				
	Day 79 post-IDI	510	35 days	R; 1:20	R; 1:1280				
		511	35 days	R; 1:4	R; 1:80				
SS14-DC^KO	Day 44 post-IDI	507	N/A	NR	NR	NR	NR	NR	NR
		503	N/A	NR	NR	NR	NR	NR	NR
	Day 79 post-IDI	501	N/A	NR	NR	NR	NR	NR	NR
		504	N/A	NR	NR	NR	NR	NR	NR

288 [1]Identification number of ID-infected donor rabbit. Naïve RIT rabbits received lymph-node from the donor rabbits.
289 [2]In rabbits with orchitis, treponemes were also detected in testicular needle aspirates using DFM.
290 [3]VDRL results (NR = Non-reactive; R = Reactive) and titers are shown. NP: Not performed
291 [4]TPPA results (NR = Non-reactive; R = Reactive) and titers are shown when measured. NP: Not performed.
292
293

294 DC^KO treponemes do not generate new TprK V region variants

Deep sequencing was performed to define TprK isogenicity and antigenic identity in the

T. pallidum inocula (for both the WT and SS14-DC^KO strains) used for rabbit ID infection and to

monitor evolution of diversity over time in progressing early lesions. Table 3 shows the most

abundant (>5% of the total for either strain) amino acid sequences for each V region for the WT

and SS14-DC^KO strains, while all the detected inoculum variants are reported in Table S1.

Table 3. TprK variability profile in the SS14 and SS14-DC^KO strains used for rabbit ID
inoculation.

V Region	Amino acid sequence	Wild-type SS14 strain Percent sequence	SS14-DC^KO strain Percent sequence
V1	GIASETGGAGALKH	96.06195	81.30797
V1	GIASEKNGGAQPLKH	Not detected	14.59834
V2	WEGKDSKGVVQAGANHSK	85.1821	40.83901
V2	WEGKSNTGAPAAGANHSK	7.328483	Not detected
V2	WEGKDSQGKAPAGANHSK	1.914901	25.66107
V2	WEGKSNTGVVQAGANHSK	1.042542	29.81366
V3	TLSGDYATARAGADDILWD	88.95463	65.03622

V3	TLSGGYATARAGADDILWD	5.402198	0.476627
V3	TLSGDYAQAAGAGADDILWD	Not detected	28.20839
V4	TDVGRKKDGAQGTV	96.91317	96.16694
V5	QASNVFQGVFLTDTTPMLQHDC	53.176	1.716312
V5	QASNVFQGVFLTTPMQKDDC[1]	26.09685	0.443286
V5	QASNVFQGVFLTTPMLQHDC[1]	12.33243	68.85708
V5	KASNVFKDVFLTNAMDMQTHDC	1.511896	12.80529
V5	KASNVFKDVFLTDTTPMQTHDC[1]	Not detected	10.32818
V6	PVHWKALAPAQPPARVDIY	49.2956	Not detected
V6	PVHWKALARAQPPARVDIY[1]	27.46295	Not detected
V6	PVHWKALARARPPVPAIY[1]	Not detected	75.82605
V7	YGGTNKKNDAAPAAPATKWKAEYCGYY	56.19608	41.41053
V7	YGGTNKKNDAAPATKWKAEYCGYY	38.03072	3.803587
V7	YGGTNKQAATKWKAEYCGYY	Not detected	46.71209

303 [1]Synthesized as peptides for ELISA.
304

Results showed that neither inoculum was isogenic for *tprK*. Among all V regions, however, V4

exhibited the highest degree of antigenic identity with a single variant representing 96.9% and

96.1% of all detected sequences in the SS14 and the SS14-DC^{KO} strains, respectively (Table 3

and Table S1), while one V1 variant accounted for 96.0% and 81.3% of all sequences found in

the WT and mutant strain, respectively. Both strains shared at least one highly represented

(>40%) V region sequence in V2, V3, and V7. Higher heterogeneity was found in V5 and V6

(Table 3 and Table S1) compared to all the other V sequences.

TprK profiling in lesion samples collected sequentially showed that generation of new

variants was impaired in SS14-DC^{KO} treponemes compared to WT treponemes for all V regions.

Heatmaps showing amino acid sequence variants detected at greater than 0.25% prevalence in

inoculum and biopsies are reported in Fig.6A-D for V1, V5-V7, respectively, and in Fig.S1-S3

for V2, V3, and V4. In V1, for example (Fig.6A), five non-inoculum variants could be detected

in rabbits infected with the WT strain, while no new variants were detected in animals infected

with the SS14-DC^{KO} strain. In V5 and V6 (Fig.6B, C) a total of 35 and 172 non-inoculum

variants, respectively, were detected at greater than 0.25% prevalence in rabbits infected with the

WT strain over the course of the experiment, but only 5 and 1 variants were detected in samples

from animals that received the mutant strain for the same V regions. All variants detected at all

time points are reported in Table S2. All eleven putative non-inoculum variants detected in

rabbits infected with the SS14-DC^{KO} strain could be detected in the inoculum at levels <0.25%

prevalence (Table S3). Taken together, these data support that elimination of the DCs virtually

stopped the generation of new TprK variants in the SS14-DC^{KO}. In addition, the two DCs

downstream of the *tp0136*, predicted to preferentially recombine into V7 (17), did not contribute

to generation of variability in the samples we analyzed from rabbits infected with the SS14-

DC^{KO}.

Development of humoral reactivity to V region variants is limited in SS14-DC^{KO}-infected

rabbits

Synthetic peptides based on V5 and V6 sequences found in the inoculum treponemes for

the WT and the SS14-DC^{KO} strains (Table 3) were used to perform ELISAs with sera from

infected rabbits to evaluate whether variation in V sequence abundance would correlate with the

development of a humoral immune response against these sequences. Results showed that in

rabbits infected with WT treponemes, selected V regions underwent an initial expansion

relatively early following treponemal ID injection compared to the inoculum (Fig.7; panels A-C).

Sequence abundance, however, decreased sharply over the following 2-3 weeks and remained

low throughout the remaining timepoints. Antibodies specific for these V regions could be

clearly detected (Figure 7, panels D-F) at day 35 post-infection and, in some instances also at day

28 post-infection, albeit only slightly above background and not for all samples. Reactivity to a

larger recombinant portion of TprK, containing two conserved loops (C1 and C2; Fig.1) was

used as a control to evaluate development of immunity to the TprK antigen. These results

support that the decrease in V region abundance is due to immune clearance of treponemes

carrying such sequences, as also shown in earlier studies (21). On the contrary, in rabbits

infected with the SS14-DC^{KO} strain (Figure 7, panels G-I) the relative abundance of the analyzed

V regions either remained virtually constant or tended to increase over the first 41 days post-

inoculation. No deep sequencing data could be uniformly obtained from SS14-DC^{KO} samples

past day 41, due to the low treponemal burden in these samples. Antibodies towards these

abundant V regions were detectable in sera from the animals infected with the KO strain only at

much later time points after infection and at a much lower level compared to the sera from the

WT-infected rabbits. Also the reactivity to the larger recombinant TprK fragment became

evident only toward the latest time points and never reached a level comparable to that of the

control animals (Fig.7, panels J-L). These results are consistent with a slow host response to the

pathogen in these rabbits due to the reduced treponemal burden at the injection site following

inoculation (Fig.5A/B) which, in turn, induced a host response less robust than by rabbits

infected with WT treponemes (Fig.7; as well as lesion size in Fig.4A, and TPPA and VDRL

results in Fig.5C/D).

Discussion

The significant impact of syphilis on public and global health warrants research efforts to

improve our understanding of the molecular mechanisms behind its pathogenesis. In turn, such

efforts could aid strategies for disease control and even vaccine development. The approach to

derive *T. pallidum* knockouts recently developed and validated by our groups (24) provides a

novel tool to study the role of the pathogen's known or putative virulence factors by comparing

loss-of-function mutants with their parent WT isolates. Even though we are still far from having

all the genetic tools needed to fully satisfy Koch's molecular postulates (26) for *T. pallidum*, this

approach is a significant step forward from to the use of heterologous bacterial systems

transformed to express *T. pallidum* genes and perform gain-of-function assays (27-29).

With the exception of an isolated study (30), all others that focused on TprK pointed at

this antigen as the chief player in pathogen immune evasion and survival during natural and

experimental infection (14-18, 20-23, 31-34). The TprK-encoding *tp0897* gene, therefore, was

considered a prime candidate for ablation. However, despite several independent attempts

performed with a pUC57-based suicide vector identical to the one used here (but with homology

arms specific for the *tprK* gene) and the same transformation/selection approach, a *tprK*^{KO} isolate

could not be derived. Although we cannot exclude that these experiments were not successful

due to the difficulties of recovering treponemes post-transformation, the hypothesis that *tprK*

could be an essential *T. pallidum* gene should not be dismissed. It is intriguing that, even when

deep sequencing of full-length *tprK* ORFs is performed (34), amplicons containing non-

functional *tprK* sequences (e.g., due to frameshifts or stop codons within the ORF) are not found

despite the heavy recombination activity occurring at this site. Aside from immune evasion,

therefore, TprK could also mediate a biological function needed for *T. pallidum* survival even *in*

vitro, where immune pressure is absent. Preliminary structural (14) and modeling data using

AlphaFold2 (Fig.1) support this antigen as a porin transporter, a function predicted for virtually

all of the Tpr proteins of *T. pallidum*, even though the nature of the transported substrate(s) is yet

to be determined. This study also demonstrated that the putative *tp0127* gene is non-essential in

T. pallidum. This small 229 codon ORF encodes a hypothetical protein with unknown function,

and the gene is reported as the member of a family of paralogs that includes *tp0314, tp0315,*

tp0346, tp0347, tp0479, *tp0617-0619*, *tp0697, and tp0698*, a group of short, annotated genes

encoding hypothetical proteins that could possibly perform the same function of *tp0127*. This

work, therefore, albeit minimally, contributes to defining the essential *T. pallidum* genome.

Despite the difficulty in deriving a *tprK*[KO] mutant, however, a strain lacking the ~4kbp

region containing 96% of the DCs was readily attained, and this deletion did not produce any

notable changes SS14-DC[KO] proliferation *in vitro* or following rabbit IT infection compared to

the parent strain. After the weeks needed for the mutant treponemes to recover post-

transformation, the *in vitro* growth curve of the SS14-DC[KO] strain overlapped that of its parent

SS14 isolate (Fig.2A), while similar time to orchitis and identical treponemal yields were

measured following rabbit infection in testes, an immune privileged organ. Altogether, these data

support that the SS14-DC[KO] strain is not impaired in its ability to proliferate in the absence of a

host immune response. Evidence that the *tprK* mRNA level (Fig.3B) is not affected *in vitro* by

DCs ablation in the mutant strain was expected due to the significant distance (~175 kbp)

between the *kan*[R] insertion site and the *tprK* expression site. This evidence further supports that

the attenuated dermal lesion phenotype was not due to impaired expression of a hypothetically

essential gene. More comprehensive RNA sequencing studies are ongoing to evaluate possible

differences in the transcriptome of these strains using treponemes propagated both *in vitro* and *in*

vivo.

Previous studies also determined that not all *T. pallidum* strains share the same repertoire

of DCs and that, even within the Nichols strain, there are lineages missing a portion of the DCs

contained in the region downstream of the *tprD* gene (15). Although the minimum number of

DCs necessary for *T. pallidum* to maintain its full virulence is still undetermined, our results

showed that that the two remaining DCs downstream of *tp0136* in the SS14-DC^KO strain were

not sufficient to rescue the attenuated phenotype. In our study, these two sites did not appear to

have contributed to the generation of new TprK variants, suggesting that their role in antigenic

variation might be marginal. Past studies that analyzed the DC molecular architecture to define

in which V region a given DC would preferentially recombine into (15, 16, 34), supported the

specificity of the two residual DCs in the SS14-DC^KO for V7. Although more studies are needed

to evaluate the importance of these isolated cassettes in generating new TprK variants, it would

be plausible to assume that limited generation of variability within one single V region would not

fully compensate for lack of variability in all remaining TprK V regions.

An apparent limitation of this study could be seen in the fact that rabbits in both groups

were not infected with strains completely isogenic for TprK but rather with a pathogen

population carrying very few dominant TprK variants and several low-prevalence variants

identified by deep sequencing. This experimental design, however, was intentional, as it allowed

us to appreciate how underrepresented variants can expand as the more abundant ones are

preferentially targeted and cleared during infection by the host response. We are nonetheless

deriving a SS14-DC^KO strain fully isogenic for TprK to evaluate the hypothesis that, upon ID

infection with this strain, early disease manifestations will be even more accentuated compared

to the present study. The remaining two DCs for *tprK* do not seem to contribute to diversity

generation according to our study, thereby removing the need to eliminate the genomic region

harboring these residual sequences. The sensitivity of the RIT assay, albeit recently questioned

(35), is known to be very high, with a 50% infectious dose of 23 spirochetes (36, 37). The lack of

seroconversion of RIT rabbits injected with lymph node extracts obtained at both day 44 and 79

from ID-infected rabbits is intriguing, particularly if we consider that full treponemal clearance

was not supported by qPCR on samples collected at both time points at which the RIT was

performed. However, one could hypothesize that the significant reduction of the treponemal

burden seen in dermal lesions would necessarily translate in an abatement of the number of

treponemes that had disseminated to the lymph nodes to a number below the sensitivity of the

RIT assay. In the future, we will also collect biopsy specimens for RNA extraction and

quantification as a surrogate of treponemal viability. In this context, an isogenic strain would

also help us understand more clearly whether the host response cleared the SS14-DC^{KO}

treponemes during infection upon performing the RIT assay. Regardless, evidence that the test

animals remained RIT negative, when all the control animals developed orchitis or

seroconverted, further supports the attenuation phenotype of the SS14-DC^{KO} strain.

A model for the attenuated phenotype exhibited by the mutant strain could be

hypothesized on the evidence that following rabbit ID infection, inoculum SS14-DC^{KO}

treponemes started to be cleared early after inoculation. By day 9, lesion development and

treponemal burden in the skin tissue biopsies started to diverge significantly between the test and

control rabbits (Fig.4A and Fig.5A). Within a few weeks, treponemal antibodies (including to

TprK) had started to develop (Fig.5D; Fig.7D-F) and led to clearance of WT treponemes

expressing the most abundant TprK inoculum sequences (Fig.7A-C). Nonetheless, wild-type

treponemes could still expand (Fig.5A) through the generation of new variants or the expansion

of low-frequency ones in the inoculum (Fig.6). In rabbits infected with the SS14-DC^{KO} strain,

however, treponemal burden increase was likely hindered by the lack of DCs, even though

underrepresented variants could still expand (Fig.6; Table S2) and maintain a very low pathogen

burden over time (Fig.5A). Abundant *tprK* inoculum sequences in the SS14-DC^{KO} treponemes

were virtually absent by day 41 post-inoculation (Fig.7G-I), even though the delayed immunity

that developed to the infection in these animals likely made the antibodies responsible for

clearance detectable only at later time points post-infection (Fig.7J-L). Future experiments will

evaluate whether immunosuppression of the animals prior to infection would rescue the

attenuation phenotype exhibited by the SS14-DC^{KO} strain.

Conclusions

We demonstrated that when genetically altered to impair its ability to vary the TprK

antigen, the *T. pallidum* SS14 strain induces attenuated disease manifestations and is unable to

proliferate in the rabbit model of the infection as effectively as the WT strain. This work further

demonstrates the importance of TprK for *T. pallidum* virulence and emphasizes how our newly

developed genetic engineering approach can contribute to shed light on the molecular

mechanisms behind syphilis pathogenesis.

Materials and Methods

Ethics statement

Male NZW rabbits (*Oryctolagus cuniculus*) ranging from 3.5-4.5 kg in weight were

purchased from Western Oregon Rabbit Company (Philomath, OR) and housed at the

Harborview Medical Center – Research and Training Building vivarium. Care was provided in

accordance with the Guide for the Care and Use of Laboratory Animals (38). All procedures

involving these animals were approved by the UW Institutional Animal Care and Use Committee

(IACUC; Protocol # 4243-01, PI: Lorenzo Giacani). Upon arrival, rabbits were allowed to

acclimate to the vivarium for several days, then bled and tested with both a treponemal

(*Treponema pallidum* particle agglutination, TPPA; Fujirebio, Tokyo, Japan) and a non-

treponemal (Venereal Disease Research Laboratory, VDRL; Becton Dickinson, Franklin Lakes,

NJ) test to confirm lack of previous or current infection with the agent of rabbit syphilis,

Treponema paraluiscuniculi. Both tests were performed according to the manufacturer`s

instructions. Only seronegative rabbits were used for experimental infection with the SS14-DC^{KO}

and SS14 parent strains (see paragraph below for rabbit infection).

TprK structure modeling

We used the ColabFold interface (39) to construct Multiple Sequence Alignments (MSA)

for the TprK query sequences by searching UniRef30 (40), Mgnify (41) and ColabFold sequence

databases with MMSeq2 (42). The MSA was used as input for structure prediction with

AlphaFold2 (43) using the default settings (template=False, amber relax=False, 3 recycles).

Increasing the number of recycles to 6 did not significantly improve the prediction. Visualization

was performed using PyMol software (https://pymol.org) (44).

Suicide vector construction

The pUC57 vector (Genscript, Piscataway, NJ) was engineered to carry the *kan*^{R} gene

downstream of the promoter and ribosomal binding site (RBS) of the *T. pallidum tp0574* gene

(encoding the Tp47 lipoprotein) as described by Weigel *et al.* (45). An 8-nt spacer separated the

RBS and the start codon of the *kan*^{R} gene. The *kan*^{R} gene sequence was derived from the Rts1

plasmid originally isolated from *Proteus vulgaris* (46) and chosen because the CDC STI

treatment guidelines do not recommend the use of kanamycin for syphilis therapy. Codon

optimization of the *kan*^{R} gene was shown not to be necessary in previous studies (24).

Two homology arms corresponding to the regions flanking the 51 DCs located downstream of

the *tprD* gene were cloned to the left and right of the *tp0574* promoter-*kan*[R] hybrid sequence in the

pUC57 vector, respectively. The upstream and downstream arms were 990 bp and 965 bp in length,

respectively, and corresponded to position 149,425-148,434 and 152,401-153,366 of the SS14 genome.

The construct was cloned between the EcoRV and KpnI sites of the pUC57 vector, in opposite

orientation compared to the *lac* promoter upstream of the polylinker. Prior to use, the insert underwent

Sanger sequencing to ensure sequence accuracy. The sequence of the insert is provided in File S1. This

construct was named p*DC*arms-47p-*kan*[R]. Primers annealing a) to the vector only, b) within the cloned

insert, and c) upstream of the DCs homology arms in the *T. pallidum* genome are reported in Table 1.

Previous work confirmed that sequences flanking the *bla* gene (encoding a β-lactamase) of the pUC57

did not carry sufficient homology to *T. pallidum* DNA to induce recombination and integration of the

bla gene into *T. pallidum* genome (24).

To obtain a highly concentrated, endotoxin-free plasmid preparation, the p*DC*arms-47p-*kan*[R] was

transformed into TOP10 *E. coli* cells (Thermo Fisher, Waltham, MA), which were then grown first in a

5-ml starter culture overnight, and then in 500 ml of LB media supplemented with 100 µg/ml of

ampicillin at 37□C. The plasmid was purified using the Endo-Free Plasmid Mega Kit (Qiagen,

Germantown, MD) according to the manufacturer's instructions. Following purification, plasmid

concentration was assessed using an ND-1000 spectrophotometer (Nanodrop Technologies,

Wilmington, NC). The purified vector was then aliquoted and stored at -20°C until use.

Source of *T. pallidum* for *in vitro* cultivation, transformation, and selection

The SS14 parent strain of *T. pallidum* used for *in vitro* propagation was obtained from a

frozen stock previously propagated via intratesticular (IT) infection in NZW rabbits as

previously reported (47). The SS14 strain was selected as a representative of the members of a

globally dominant cluster, known as the SS14 clade (48). *In vitro* strain co-culturing with rabbit

Sf1Ep cells was performed as per published protocols in a 6-well culture plate (Corning Inc,

Corning, NY) (49). Before the addition of the treponemal cells, Sf1Ep cells were incubated in a

5% CO_2 atmosphere in a HeraCell 150 incubator (Thermo Fisher). Following treponemal cell

inoculation, the microaerophilic atmosphere (MA; 1.5% O_2, 3.5% CO_2, and 95% N_2) necessary to

maintain treponemal viability was generated in a Heracell VIOS 160i tri-gas incubator (Thermo

Fisher).

For transformation, treponemes were first sub-cultured into the wells of a 24-well plate as

previously reported (24). Briefly, the day before treponemal inoculation, a 24-well plate was

seeded with $2x10^4$ rabbit Sf1Ep cells/well in 2.5 ml of culture media. The plates were then

incubated overnight in the HeraCell incubator. On the same day, TpCM-2 media was prepared

according to protocol and equilibrated overnight at 34°C in the tri-gas incubator. The following

day, cell culture media was removed from the 24-well plate, and cells were rinsed with TpCM-2

media equilibrated in the tri-gas incubator. Subsequently, each well was filled with 2.5 ml of

equilibrated TpCM-2 media, and the plate was transferred to the tri-gas incubator. To prepare the

treponemal inoculum, the Sf1Ep cells seeded the previous week with the parent SS14 cells were

trypsinized to allow the release and counting of spirochetes using an Optiphot 2 dark-field

microscope (DFM; Nikon, Melville, NY). Treponemal counting was performed according to

Lukehart and Marra (50). A total of ~$2.5x10^8$ treponemes were inoculated 24 hours after plating

the Sf1Ep. Following treponemal addition, the total volume of media in each well was brought to

2.5 ml. Two days following treponemal cell addition, 1 ml of old media was replaced with fresh

one. Four days after treponemal cell inoculation, the culture media was removed gently so as not

to disturb adherent treponemes. Media was replaced by 500 µl of transformation buffer (50 mM

CaCl$_2$, 10 mM Tris pH 7.4; equilibrated in MA) containing 15 ug total of p*DC*arms-47p-*kan*^{R}

vector. Cells were incubated in transformation buffer for 10 min at 34°C in the tri-gas incubator

and then washed twice with equilibrated TpCM-2 media to remove free plasmid from the culture

wells. Finally, 2.5 ml of fresh TpCM-2 equilibrated in tri-gas were added to the wells, and plates

were returned to the tri-gas incubator. The following day, concentrated tissue-culture grade

liquid kanamycin sulfate (Sigma-Aldrich, St. Louis, MO) was added to the appropriate wells to

reach a final concentration of 200 µg/ml. As a control, to confirm the treponemicidal activity of

kanamycin, WT treponemes were also incubated in fresh TpCM-2 media containing kanamycin.

Previous work (24) demonstrated lack of toxicity associated to CaCl$_2$ exposure for treponemes,

and a CaCl$_2$ only control was not performed here again. The parent SS14 strain, however, was

propagated in parallel to the transformed treponemes as a control. Kanamycin-containing TpCM-

2 media was exchanged weekly but treponemes were sub-cultured every two weeks as already

described (24) until they reached a density of ~10^7 cells/ml determined by DFM. At this point,

cultures were sub-cultured first into one well of a 6-well plate (seeded with 10^5 Sf1Ep cells on

the previous day) and then into all the wells of a 6-well plate at the following passage to further

expand the strain and minimize the chances of culture loss due to contamination. At selected

passages, residual treponemes were pelleted by centrifugation at 15,000 rpm for 10 min on a

tabletop centrifuge and resuspended in 1X DNA lysis buffer (10 mM Tris-HCl, 0.1 M EDTA,

0.5% SDS) for DNA extraction, or Trizol (Thermo Fisher) for RNA extraction. A WT SS14

strain with limited TprK variability was attained through *in vitro* cultivation of treponemes

following limiting dilution (49).

DNA and RNA extraction

DNA extraction from cultured SS14-DC^{KO} or WT strains propagated *in vitro* following the transformation procedure as well as from lesion biopsies (see below paragraph about rabbit infection and sample collection) was performed using the QIAamp DNA Mini Kit (Qiagen) or the Zymo Quick-DNA 96 Kit (Zymo Research, Irvine, CA), respectively, according to the manufacturer's instructions. Extracted DNA was stored at -80°C until use for qualitative/quantitative amplifications or genome sequencing (see below paragraphs) to evaluate successful insertion of the *kan*^{R} gene or to evaluate treponemal burden. Samples for RNA extraction were processed according to the Trizol reagent manual. Total RNA was treated with DNaseI according to the protocol provided with the TURBO DNA-free kit (Thermo Fisher). DNA-free RNA was checked for residual DNA contamination by qualitative amplification using primers specific for the *tp0574* gene (primers in Table 1) as already described (51). Reverse transcription (RT) of total RNA was performed using the High-Capacity cDNA Reverse Transcription kit (Thermo Fisher) with random hexamers according to the provided protocol. cDNA samples were stored at -80°C until use for qPCR to quantify the level of expression of the *tprK* gene normalized to the *tp0574* message (primers in Table 1) as previously reported (52).

Droplet digital PCR (ddPCR)

Droplet digital PCR assays were conducted to assess the ratio between the number of copies of the *kan*^{R} gene, the *tp0001* gene (*dnaA*), and an open reading frame (ORF; *tp0127*) encoding a 229 amino acid-long hypothetical protein with unknown function contained within the ablated DCs region. This short ORF was selected because it is unique to the ablated region, while primers annealing directly to the DCs sequences could potentially also recognize

sequences in the *tprK* expression site. Analyzed DNA was extracted from the SS14-DC^{KO} from

in vitro passages 2-5 post-transformation and from the parent SS14 strain propagated in parallel.

ddPCR was performed on a Bio-Rad QX100 system (Bio-Rad, Carlsbad, CA). Primers and probe

are listed in Table 1. Each reaction was performed using ddPCR Supermix for Probes (Bio-Rad)

with the final concentration of primers at 900 nM and probes at 250 nM in a total reaction

volume of 25 µl. Before amplification, template DNA was digested with 25 units of EcoRI (New

England Biolabs, Ipswich, MA). After droplet generation, droplets were transferred to a 96-well

PCR plate and amplified on a 2720 Thermal Cycler (Thermo Fisher) with the following cycling

parameters: 94°C for 10 min, followed by 40 cycles of 94°C for 30 s and 60°C for 1 min, and

98°C hold for 10 min. After amplification, the plate was transferred to QX200 droplet reader

(Bio-Rad). Results were analyzed using the QuantaSoft software (Bio-Rad).

***T. pallidum* whole-genome sequencing**

Whole-genome sequencing was performed following DNA extraction of SS14-DC^{KO}

cultured *in vitro* and harvested at passage #9 post-transformation. Pre-capture libraries were

prepared from up to 100 ng input DNA using the KAPA Hyperplus kit (Roche) and TruSeq

adapters and barcoded primers (Illumina), following the manufacturer's protocols, yielding an

average library size longer than 400 bp. Hybrid capture of *T. pallidum* genomic DNA was

performed overnight (>16 hours) using a custom IDT xGen panel designed against the reference

genome NC_010741, following the manufacturer's protocol. Short-read sequencing was

performed on an Illumina NextSeq 2000 with 150 bp paired-end reads, yielding 2.75 and 8.20

million raw reads for wild type and mutant strains, respectively. Reads were adapted and quality

trimmed using Trimmomatic v0.39 (53) and assembled to the *T. pallidum* SS14 reference

genome NC_021508.1, or NC_021508.1 with the DC region manually replaced by the kan^R

cassette, using Bowtie2 v2.4.1 (54) and deduplicated with Picard MarkDuplicates v2.23.3

(available at http://broadinstitute.github.io/picard), to an average coverage exceeding 177x for

the WT and 405x for the KO strain. Manual confirmation of expected coverage and junctions

was performed by visual inspection in Geneious Prime v2020.1.2 (55). Raw reads were

deposited to the NCBI Sequence Read Archive under BioProject PRJNA909291.

Qualitative and quantitative PCR

Samples harvested during routine propagation of the SS14-DC^{KO} and parent strains were

assessed for integration of the kan^R gene into the DC locus by using qualitative PCR. All primers

used in these assays and amplicon sizes are reported in Table 1. In the first amplification, the

sense primer targeted a region of the *T. pallidum* genome immediately upstream of the left

homology arm of the vector (and hence not cloned into p*DC*arms-47p-*kan*^{R}), while the antisense

primers targeted the kan^R gene, with the rationale that only a kan^R gene integrated into the DC

region would provide amplification. In the second PCR, the sense primer targeted the kan^R gene,

and the antisense primer targeted the genomic region downstream of the right homology arm of

the vector. A third PCR used both primers external to the homology arms to show a differential

in size when DNA from the transformed and parent strain, respectively, was used. Amplification

of the *tp0574* gene was used as positive amplification control, and amplification with pUC57

plasmid-specific primers was performed to confirm that no residual transformation plasmid

persisted in the SS14-DC^{KO} culture. Amplifications were performed using five microliters of

extracted DNA in 50 μl final volume containing 2.5 units of GoTaq polymerase (Promega,

Madison, WI), 200 μM of each dNTP, 1.5 mM of $MgCl_2$, and 400 nM of sense and antisense

primers. Cycling parameters were initial denaturation (94°C) and final extension (72°C) for 10

min each. Denaturation (94°C) and annealing (60°C) steps were carried on for 1 min each, while

the extension step (72°C) was carried out for 1 or 2 min depending on amplicon length. A total

of 40 cycles were performed in each reaction.

A TaqMan qPCR assay targeting the *tp0574* gene was instead used to quantify

treponemal load in samples extracted from rabbit lesion biopsies as previously described (56).

Primers/probe and amplicon size are reported in Table 1. Amplifications were run on a

QuantStudio 3 or 5 thermal cycler (Thermo Fisher) and results were analyzed using the

instrument software. Data were imported into Prism 8 (GraphPad Software, San Diego, CA) and

further analyzed to assess statistical significance of the values from the control rabbits and those

infected with the knockout strain using one-way ANOVA with the Dunnett test for correction of

multiple comparisons or Student's *t*-test, with significance set at $p<0.05$ in both cases.

TprK molecular profiling

TprK libraries were prepared similarly as previously described (17). Technical duplicates

were performed with thirty-five cycles of PCR in a 50 μL reaction with 0.4 mM TprK forward

and reverse primers (Table S4) using the 2x CloneAmp MasterMix (Takara) with a 62°C

annealing temperature. Biopsy DNA input was limited to 4 μL per reaction due to high host

DNA concentration inhibiting amplification, which represented a range of 75-1433320 genome

copies. For the parent and SS14-DC^{KO} inoculum propagated *in vitro*, 1e6 genome copies were

input. PCR products were run on a 1% agarose gel to confirm the presence of a band at

approximately 1.6 kb and cleaned with 0.6x Ampure beads (Beckman Coulter); samples with

fewer than 200 input genome copies were excluded from further analysis due to inconsistent

amplification. Two nanograms of PCR product were used in a two-fifths Nextera (Illumina)

library prep with 15 cycles of amplification. Low molecular weight products were removed with

0.6x Ampure beads, with an average size of approximately 350 bp. Libraries were quantified,

pooled, and run on a 2x150 paired end run on an Illumina Nextseq 2000. Raw reads were

adapter-trimmed and filtered for quality scores above Q20 using Trimmomatic v0.39 (53),

followed by merging the paired end reads using BBmerge (https://www.osti.gov/biblio/1241166)

in default parameters. Variable site sequences were identified using custom python and R scripts

(22), and those containing frameshifts or nonsense mutations in the translated sequences

removed. Replicate samples yielding a minimum of 2,000 reads in all V regions in both

replicates were kept for downstream analysis, corresponding to input merged reads ranging

between 89182–5746195. Only sequences with 5 or more raw reads per technical replicate were

retained per sample. Samples with a Pearson correlation coefficient lower than 0.95 between

replicates were removed, and the number of reads for each variable region in each technical

replicate were averaged. Sequences with less than 0.25% prevalence per V region per sample

were excluded from further analysis. Raw reads are available in NCBI BioProject

PRJNA909291.

Rabbit intradermal infection, lesion monitoring, and sample collection

Two groups of 4 animals each were infected intradermally (ID) on the same day on ten

sites/animal on their shaved backs with 10^6 viable *T. pallidum* cells. One group received the WT

SS14 strain, and the other group was infected with the SS14-DC^{KO} strain. Briefly, the day of the

ID infection, treponemes were harvested from culture plates as described above for routine

passages. Following a low speed-centrifugation for 10 min at 1,000 rpm in a R4702 centrifuge

(Eppendorf, Hauppauge, NY) to remove rabbit Sf1Ep cells, treponemal cells were enumerated

and checked for motility by DFM and diluted using sterile saline to a final concentration of 10^7

cells/ml. One hundred microliters of treponemal suspension were injected in each site on the

rabbit backs. Treponemal cells were evaluated again for motility using DFM after performing ID

inoculation to ensure that pathogen viability was unaltered at the end of the procedure.

Intradermally-infected rabbits were shaved daily post-inoculation to allow monitoring of

lesion progression and to facilitate collection of needle aspirates and lesion biopsies. Two

animals from each group were euthanized on day 44 post-inoculation to harvest popliteal lymph

nodes and perform an RIT (see paragraph below). The remaining four animals were monitored

for an additional 36 days, and then also euthanized to perform RIT.

Lesion biopsies and aspirates were collected at day 9, 22, 30, 37, 41 post-infection from

all animals, at day 44 post-infection from the four animals that were euthanized first, and at day

51, 58, 65, 72, and 79 from the remaining four animals. One biopsy per animal was obtained per

collection day. At day 44, however, biopsies were collected from all lesions (unless previously

biopsied) of the animals that were euthanized to perform the first round of RIT assays. Aspirates

were collected from approximately half of the lesions from each rabbit per collection day. Once

biopsied, a lesion was no longer measured or used to collect aspirates. Aspirates were obtained

using an Excel 1cc zero dead space tuberculin syringe with a 25Gx3/8" needle (Excel

international, Birmingham, AI) by inserting the needle laterally into the side of the developing

lesion. Once inserted, the plunger was drawn, and the needle moved back and forth five times

while holding the plunger. Lesion aspirates were resuspended in 60 μl of sterile saline, and 18 μl

of each suspension were examined by DFM to assess for the presence of treponemes and, in

case, enumerate them. *T. pallidum* cells were counted blindly in a total of 50 fields/sample. The

operator was blinded to the identity of the samples. Biopsies were minced using forceps and a

sterile scalpel, resuspended in 1X DNA lysis buffer. Serum was obtained weekly from each

animal to perform VDRL and TPPA tests to monitor development of humoral immunity in

response to infection. Statistical analyses were performed using one-way ANOVA with the

Dunnett correction of multiple comparisons or Student's *t*-test, with significance set at $p<0.05$.

Rabbit infectivity test (RIT)

Immediately after euthanasia, popliteal lymph nodes were harvested from rabbits to perform an

RIT as previously described (50). Following injection of minced popliteal lymph nodes in the animal's

left testis, RIT rabbits were monitored twice a week for orchitis development. If orchitis developed, to

ascertain that it was due to *T. pallidum* infection, testicular needle aspiration was performed as described

above to be analyzed by DFM. If orchitis was confirmed by finding treponemes in the sample, then the

animal was euthanized. As orchitis does not always develop in RIT animals, rabbits were also bled

monthly to assess seroconversion using VDRL and TPPA tests. If seroconversion occurred, the animal

was euthanized. If at the end of the 3-month observation period following IT injection, the rabbit had not

developed orchitis or seroconverted, then the RIT outcome was negative, and the animal was deemed

uninfected.

Rabbit intratesticular infection

To assess differences in *in vivo* growth between the parent and knockout strains studied

here, one rabbit was infected IT with the *T. pallidum* parent SS14 strain ($7x10^7$ cells/testis),

while a second animal was inoculated with the SS14-DC^{KO} strains IT ($6x10^7$ cells/testis). Both

animals were monitored daily to determine the time to orchitis. At peak orchitis, animals were

euthanized, testes were harvested, and content extracted to quantify treponemal burden by DFM.

Procedures were performed as previously described (50).

ELISA using V region synthetic peptides

Synthetic peptides were designed based upon the TprK profiling results for a subset of

animals infected with the SS14-DC^{KO}. A total of 5 peptides matching inoculum sequences for

both the WT and SS14-DC^{KO} strains (Table 3, labelled with a superscripted "1") were produced

by Genscript (Piscataway, NJ). Upon reception, lyophilized peptides were rehydrated in sterile

PBS (supplemented with DMSO or acetic-acid) to a stock solution of 4mg/ml. Reconstituted

peptides were stored at -80°C until use. For ELISA, peptides were further diluted to 10 μg/ml in

PBS, and 50 μl of working dilution (500 ng total) were used to coat the wells of a 96-well

Microwell Maxisorp flat-bottom plate (Thermo-Fisher, Waltham, MA) as previously described

(19). Briefly, plates were then washed three times with PBS+0.05% Tween-20 and blocked with

200 μl of 5% nonfat dry milk (NFM) in sterile PBS for three hours. At this step, NFM was used

to block no-antigen wells for background signal. Wells were then washed again as above and

primary antibody was applied immediately thereafter. Sera from infected animals were used at a

1:20 dilution. After overnight incubation with the primary antibody, plates were washed again as

above and 100 μl of a goat anti-rabbit IgG (H + L) alkaline phosphatase conjugate (Sigma, St.

Louis, MO) diluted 1:2,000 in PBS/1% NFM were added to each well and incubated for 1.5h at

room temperature. Plates were washed again and developed with 50 μl/well of 1 mg/ml para-

Nitrophenyl Phosphate (pNPP) substrate (Sigma), dissolved in glycine buffer (0.1M glycine, pH

10.4, 1mM $MgCl_2$ and 1 mM $ZnCl_2$) for 1 h according to the manufacturer`s instructions.

Absorbance was measured at OD_{405} using a Molecular Devices SpectraMax Plus microplate

reader (Molecular Devices, San Jose, CA). The mean of triplicate experimental wells minus the

mean of pre-infection serum wells tested with the same antigen was calculated for all

proteins/peptides tested. Plotted data represent the mean $\pm$ SEM for triplicate wells tested.

Results were analyzed using Student's *t*-test with significance set at $p \leq 0.05$.

Acknowledgments

This work was supported by NIAID grant number U19AI144133 (Project 2 and

Genomics and Isolation Core; Project 2 leader: L.G.; Core leaders: L.G. and A.L.G.; PI: Anna

Wald, University of Washington) and by Open Philanthropy (to L.G.). The content of this study

is solely the responsibility of the authors and does not necessarily represent the official views of

the National Institutes of Health or Open Philanthropy. The funders had no role in study design,

data collection, and analysis, decision to publish, or preparation of the manuscript.

Figure Legends

Figure 1. TprK (sequence AIP85979.1) is predicted by AlphaFold2 to fold into a 20-strands

beta-barrel structures with 10 surface-exposed loops. Variable loops (V1-V7) are highlighted in

colour. Conserved loops (C1-C3) are in white, and β-scaffolding are in light blue. OM: outer

membrane; PS: periplasmic space.

Figure 2. (**A**) Twenty-eight-week *in vitro* growth curve of transformed SS14 *T. pallidum* cells

(SS14-DC^{KO}) in selective media (solid orange circles), WT SS14 cells in selective media (solid

black triangles), and WT SS14 strain in non-selective media (solid blue squares). Propagation of

the parent strain for this experiment stopped at week 18. Parent treponemes propagated in

kanamycin-containing media were never countable by DFM. (**B; subpanels 1-5**) Qualitative

amplifications to assess kan^R integration. Primer combinations are reported in Table 1. Subpanels

show amplifications a) with vector-specific primers (subpanel #1, 2,967 bp amplicon), yielding

an amplicon only with suicide vector control DNA; b) with *tp0574* primers (subpanel #2, 313 bp

amplicon), positive in both parent and transformed treponemes; c) with primers annealing to the

kan^R gene and outside of the right homology arm in the *T. pallidum* genome (subpanels #3; 1,846

bp size amplicon), yielding a positive amplicon only with template from transformed

treponemes; d) with primers annealing upstream and downstream of the homology arms in the *T.*

pallidum genome (subpanel #4, 2,982 bp and 5,980 bp size amplicons), yielding a shorter

amplicon size in transformed treponemes due to deletion of the DCs); and primers annealing to

the kan^R gene (subpanel #5, 663 bp size amplicon). M: molecular size marker (bp); KO: SS14-

DC^{KO}, WT: parent WT SS14; NTC: No-template control, P: suicide vector (p*DC*arms-47p-kan^R)

plasmid DNA.

Figure 3. (A) Ratios between the kan^R, *dnaA (tp0001)*, and *tp0127* (contained within the deleted

DC) copy number determined by droplet digital PCR (ddPCR) on template DNA from the SS14-

DC^{KO} and parent SS14 strain harvested at passage #2-5 post-transformation. The kan^R:*dnaA* ratio

for the parent SS14 was zero at each passage. **(B)** Message quantification of *tprK* (normalized to

tp0574) in SS14-DC^{KO} and parent SS14 strain propagated *in vitro* showing comparable

expression of *tprK* between the control and transformed strain. Significance ($p<0.05$) was

assessed by performing Student's *t*-test. **(C-G)** Whole-genome sequencing of the SS14-DC^{KO}

and WT strains (passage #9) post-transformation. **(C)** SS14-DC^{KO} reads assembled to the WT

genome reference (NC_021508.1/CP004011.1) showing a gap where the DC locus previously

was. **(D)** SS14-DC^{KO} reads assembled to the SS14-DC^{KO} reference genome, where the DC locus

was replaced *in silico* with the *kan*^{R} cassette sequence showing reads aligning to the cassette

sequence. **(E)** Schematic of the 51 deleted DCs, color-coded based on their target V region and

the *kan*^{R} cassette (*tp0574* promoter- *kan*^{R}) that replaced the DCs. **(F)** Reads from the WT SS14

genome aligned to the SS14 reference genome showing the integrity of the DC locus. (**G**) Reads

from the WT genome aligned to the SS14-DC^{KO} reference genome showing a gap where the

kan^R cassette is located.

Figure 4. (A) Measurements of lesion diameter (mm) over time in rabbits infected with the SS14

strain and the SS14-DC^{KO} strain. Statistical analyses were performed using one-way ANOVA

with the Dunnett correction of multiple comparisons or Student's *t*-test, with significance set at

$p<0.05$, indicated by an asterisk. **(B)** Pictures of the animals still in the study at day 72 post-

infection showing large, ulcerated lesions in control rabbits (right panel), and significantly

smaller ($p<0.05$) lesions in animals infected with the knockout strain (left panel). Circled areas

represent lesions that were not yet biopsied at the time the picture was taken. Black ink dots near

the lesion area mark the injection sites.

Figure 5. (A) Treponemal burden in lesion biopsies collected from control and test animals

assessed by qPCR targeting the *tp0574* gene. The presence of *T. pallidum* DNA was confirmed

in all samples, but treponemal burden in control samples was uniformly higher. Statistical

analyses were performed using Student's *t*-test, with significance set at $p<0.05$ ($p<0.01$ is

indicated with two asterisks, $p<0.001$ is indicated with three asterisks). **(B)** Treponemal burden

in lesion biopsies collected from control and test animals assessed by DFM in lesion needle

aspirates. Treponemes could be detected only in samples from control rabbits. **(C)** Significant

delay in development of VDRL antibody titers in sera collected from animals infected with the

knockout strain compared to serum samples from control rabbits. Controls seroconverted earlier

and achieved significantly higher titers compared to test animals. Uninfected rabbit sera

(screened at animal arrival at the vivarium) pre-inoculation had no VDRL reactivity. **(D)** Initial

delay in development of treponemal antibodies (assessed using the TPPA test) in test animals

compared to control rabbits. Titer differences became not significant at day 30 post-inoculation.

NR: Nonreactive. In both cases significance ($p<0.05$) was assessed by performing Student's t-

test. Uninfected rabbit sera (screened at animal arrival at the vivarium) pre-inoculation had no

TPPA reactivity.

Figure 6. Heatmap of sequence diversity in V1 **(A)**, V5 **(B)**, V6 **(C)**, and V7 **(D)** regions

showing persistence of inoculum variants and generation of non-inoculum variants in samples

collected overtime from rabbits infected with the WT SS14 strain ("WT"-labeled samples, left

side of the map) and form rabbits infected with the SS14-DC^{KO} strain ("KO"-labeled samples,

right side of the map), separated by a bold vertical dashed line. In each map, inoculum sequences

for the WT strain are labeled as "WT Inoc B00 Day 00", where WT Inoc indicates WT inoculum

treponemes, B00 indicates that the sample was not obtained from a biopsy, and Day 00 indicates

the experiment's time 0. The same nomenclature was adopted for the SS14-DC^{KO} inoculum, with

the exception that KO replaced WT. Samples collected post-inoculation report rabbit number

(R#), biopsy number (B#), and day post-inoculation the sample was obtained (Day#). Light gray

dashed lines separate individual animals. Missing samples did not yield data.

Figure 7. Comparison between changes over time in sequence abundance of selected V5 and V6

regions present in the inoculum treponemes and the development of humoral immunity to the

same regions, assessed by ELISA performed with sera from infected rabbits and synthetic

peptides. Panels A-C (sequence abundance) and D-F (humoral response to the selected peptide)

pertain to rabbits infected with the WT strain, while panels G-I (sequence abundance) and J-L

(humoral response to the selected peptide) pertain to rabbits infected with the SS14-DC^{KO} strain.

Sequence abundance was based on the *tprK* deep sequencing results for all rabbits.

References

1.   WHO. Prevalence and incidence of selected sexually transmitted infections *Chlamydia*

trachomatis, Neisseria gonorrhoeae, syphilis and *Trichomonas vaginalis*: methods and results

used by WHO to generate 2005 estimates. World Health Organization, Geneva. 2011.

2.   Gerbase AC, Rowley JT, Mertens TE. Global epidemiology of sexually transmitted

diseases. The Lancet. 1998;351.

3.   CDC. 2018 Sexually Transmitted Disease Surveillance. Atlanta, GA: US Department of

Health and Human Services: Centers for Disease Control and Prevention. 2019.

4.   Savage EJ, Marsh K, Duffell S, Ison CA, Zaman A, Hughes G. Rapid increase in

gonorrhoea and syphilis diagnoses in England in 2011. Euro Surveill. 2012;17(29).

5.   Savage EJ, Hughes G, Ison C, Lowndes CM. Syphilis and gonorrhoea in men who have

sex with men: a European overview. Euro Surveill. 2009;14(47).

6.   Simms I, Fenton KA, Ashton M, Turner KM, Crawley-Boevey EE, Gorton R, et al. The

re-emergence of syphilis in the United Kingdom: the new epidemic phases. Sex Transm Dis.

2005;32(4):220-6.

7.   Tucker JD, Cohen MS. China's syphilis epidemic: epidemiology, proximate determinants

of spread, and control responses. Curr Opin Infect Dis. 2011;24(1):50-5.

8.   Jin F, Prestage GP, Kippax SC, Pell CM, Donovan BJ, Kaldor JM, et al. Epidemic

syphilis among homosexually active men in Sydney. Med J Aust. 2005;183(4):179-83.

9.   Goldenberg RL, Thompson C. The infectious origins of stillbirth. Am J Obstet Gynecol.

2003;189(3):861-73.

10.   Moline HR, Smith JF, Jr. The continuing threat of syphilis in pregnancy. Curr Opin

Obstet Gynecol. 2016;28(2):101-4.

11.   Wijesooriya NS, Rochat RW, Kamb ML, Turlapati P, Temmerman M, Broutet N, et al.

Global burden of maternal and congenital syphilis in 2008 and 2012: a health systems modelling

study. Lancet Glob Health. 2016;4(8):e525-33.

12.   Peeling RW, Mabey D, Kamb ML, Chen XS, Radolf JD, Benzaken AS. Syphilis. Nat

Rev Dis Primers. 2017;3:17073.

13.   LaFond RE, Lukehart SA. Biological basis for syphilis. Clin Microbiol Rev.

2006;19(1):29-49.

14.   Lian T, Zhang B, Giacani L, Kou C, Yang X, Zhang R, et al. Full-length TprK of

Treponema pallidum subsp. *pallidum* in lipid nanodiscs is a monomeric porin. Enzyme Microb

Technol. 2021;153:109897.

15.   Giacani L, Brandt SL, Puray-Chavez M, Brinck Reid T, Godornes C, Molini BJ, et al.

Comparative Investigation of the Genomic Regions Involved in Antigenic Variation of the TprK

Antigen among Treponemal Species, Subspecies, and Strains. J Bacteriol. 2012;194(16):4208-

25.

16.   Centurion-Lara A, LaFond RE, Hevner K, Godornes C, Molini BJ, Van Voorhis WC, et

al. Gene conversion: a mechanism for generation of heterogeneity in the *tprK* gene of

Treponema pallidum during infection. Molecular Microbiology. 2004;52(6):1579-96.

17.   Addetia A, Lin MJ, Phung Q, Xie H, Huang ML, Ciccarese G, et al. Estimation of Full-

Length TprK Diversity in *Treponema pallidum* subsp. *pallidum*. mBio. 2020;11(5).

18.   LaFond RE, Centurion-Lara A, Godornes C, Van Voorhis WC, Lukehart SA. TprK

sequence diversity accumulates during infection of rabbits with *Treponema pallidum* subsp.

pallidum Nichols strain. Infect Immun. 2006;74(3):1896-906.

19.   Morgan CA, Molini BJ, Lukehart SA, Van Voorhis WC. Segregation of B and T cell

epitopes of *Treponema pallidum* repeat protein K to variable and conserved regions during

experimental syphilis infection. J Immunol. 2002;169(2):952-7.

20.   LaFond RE, Molini BJ, Van Voorhis WC, Lukehart SA. Antigenic variation of TprK V

regions abrogates specific antibody binding in syphilis. Infect Immun. 2006;74(11):6244-51.

21.   Giacani L, Molini BJ, Kim EY, Godornes BC, Leader BT, Tantalo LC, et al. Antigenic

variation in *Treponema pallidum*: TprK sequence diversity accumulates in response to immune

pressure during experimental syphilis. J Immunol. 2010;184(7):3822-9.

22.   Lin MJ, Haynes AM, Addetia A, Lieberman NAP, Phung Q, Xie H, et al. Longitudinal

TprK profiling of in vivo and in vitro-propagated *Treponema pallidum* subsp. *pallidum* reveals

accumulation of antigenic variants in absence of immune pressure. PLoS Negl Trop Dis.

2021;15(9):e0009753.

23.   Reid TB, Molini BJ, Fernandez MC, Lukehart SA. Antigenic variation of TprK facilitates

development of secondary syphilis. Infect Immun. 2014;82(12):4959-67.

24.   Romeis E, Tantalo L, Lieberman N, Phung Q, Greninger A, Giacani L. Genetic

engineering of *Treponema pallidum* subsp. *pallidum*, the Syphilis Spirochete. PLoS Pathog.

2021;17(7):e1009612.

25.   Giacani L, Iverson-Cabral SL, King JC, Molini BJ, Lukehart SA, Centurion-Lara A.

Complete Genome Sequence of the *Treponema pallidum* subsp. *pallidum* Sea81-4 Strain.

Genome Announc. 2014;2(2).

26.   Falkow S. Molecular Koch's postulates applied to microbial pathogenicity. Rev Infect

Dis. 1988;10 Suppl 2:S274-6.

27.   Chi B, Chauhan S, Kuramitsu H. Development of a system for expressing heterologous

genes in the oral spirochete Treponema denticola and its use in expression of the *Treponema*

pallidum flaA gene. Infect Immun. 1999;67(7):3653-6.

28.   Chan K, Nasereddin T, Alter L, Centurion-Lara A, Giacani L, Parveen N. *Treponema*

pallidum Lipoprotein TP0435 Expressed in Borrelia burgdorferi Produces Multiple

Surface/Periplasmic Isoforms and mediates Adherence. Sci Rep. 2016;6:25593.

29.   Cameron CE, Kuroiwa JM, Yamada M, Francescutti T, Chi B, Kuramitsu HK.

Heterologous expression of the *Treponema pallidum l*aminin-binding adhesin Tp0751 in the

culturable spirochete *Treponema phagedenis*. J Bacteriol. 2008;190(7):2565-71.

30.   Hazlett KRO, Sellati TJ, Nguyen TT, Cox DL, Clawson ML, Caimano MJ, et al. The

TprK protein of *Treponema pallidum* is periplasmic and is not a target of opsonic antibody or

protective immunity. J Exp Med. 2001;193(9):1015-26.

31.   Stamm LV, Bergen HL. The sequence-variable, single-copy *tprK* gene of *Treponema*

pallidum Nichols strain UNC and street strain 14 encodes heterogeneous TprK proteins. Infect

Immun. 2000;68(11):6482-6.

32.   Centurion-Lara A, Godornes C, Castro C, Van Voorhis WC, Lukehart SA. The *tprK* gene

is heterogeneous among *Treponema pallidum* strains and has multiple alleles. Infect Immun.

2000;68(2):824-31.

33.   LaFond RE, Centurion-Lara A, Godornes C, Rompalo AM, Van Voorhis WC, Lukehart

SA. Sequence diversity of *Treponema pallidum* subsp. *pallidum tprK* in human syphilis lesions

and rabbit-propagated isolates. J Bacteriol. 2003;185(21):6262-8.

34.   Addetia A, Tantalo LC, Lin MJ, Xie H, Huang ML, Marra CM, et al. Comparative

genomics and full-length Tprk profiling of *Treponema pallidum* subsp. *pallidum* reinfection.

PLoS Negl Trop Dis. 2020;14(4):e0007921.

35.   Tong ML, Zhang HL, Zhu XZ, Fan JY, Gao K, Lin LR, et al. Re-evaluating the

sensitivity of the rabbit infectivity test for *Treponema pallidum* in modern era. Clin Chim Acta.

2017;464:136-41.

36.   Turner TB, Hollander DH. Biology of the Treponematoses. Geneva: World Health

Organization; 1957.

37.   Turner TB, Hardy PH, Newman B. Infectivity tests in syphilis. Br J Vener Dis.

1969;45(3):183-96.

38.   Janier M. Ceftriaxone is effective for treating patients with primary syphilis. Sex Transm

Dis. 1988;15(1):70.

39.   Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. ColabFold -

Making protein folding accessible to all. bioRxiv. 2021:2021.08.15.456425.

40.   Suzek BE, Wang Y, Huang H, McGarvey PB, Wu CH. UniRef clusters: a comprehensive

and scalable alternative for improving sequence similarity searches. Bioinformatics.

2015;31(6):926-32.

41.   Mitchell AL, Almeida A, Beracochea M, Boland M, Burgin J, Cochrane G, et al.

MGnify: the microbiome analysis resource in 2020. Nucleic Acids Res. 2020;48(D1):D570-d8.

42.   Steinegger M, Söding J. MMseqs2 enables sensitive protein sequence searching for the

analysis of massive data sets. Nat Biotechnol. 2017;35(11):1026-8.

43.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly

accurate protein structure prediction with AlphaFold. Nature. 2021;596(7873):583-9.

44.   Janson G, Zhang C, Prado MG, Paiardini A. PyMod 2.0: improvements in protein

sequence-structure analysis and homology modeling within PyMOL. Bioinformatics.

2017;33(3):444-6.

45.   Weigel LM, Brandt ME, Norgard MV. Analysis of the N-terminal region of the 47-

kilodalton integral membrane lipoprotein of *Treponema pallidum*. Infect Immun.

1992;60(4):1568-76.

46.   Murata T, Ohnishi M, Ara T, Kaneko J, Han CG, Li YF, et al. Complete nucleotide

sequence of plasmid Rts1: implications for evolution of large plasmid genomes. J Bacteriol.

2002;184(12):3194-202.

47.   Baker-Zander SA, Fohn MJ, Lukehart SA. Development of cellular immunity to

individual soluble antigens of *Treponema pallidum* during experimental syphilis. J Immunol.

1988;141(12):4363-9.

48.   Arora N, Schuenemann VJ, Jager G, Peltzer A, Seitz A, Herbig A, et al. Origin of

modern syphilis and emergence of a pandemic *Treponema pallidum* cluster. Nat Microbiol.

2016;2:16245.

49.   Edmondson DG, Hu B, Norris SJ. Long-Term In Vitro Culture of the Syphilis Spirochete

Treponema pallidum subsp. *pallidum*. mBio. 2018;9(3).

50. Lukehart SA, Marra CM. Isolation and laboratory maintenance of *Treponema pallidum*.

Curr Protoc Microbiol. 2007;Chapter 12:Unit 12A.1.

51. Giacani L, Hevner K, Centurion-Lara A. Gene organization and transcriptional analysis

of the *tprJ*, *tprI*, *tprG* and *tprF* loci in the Nichols and Sea 81-4 *Treponema pallidum* isolates. J

Bacteriol. 2005;187(17):6084-93.

52. Giacani L, Molini B, Godornes C, Barrett L, Van Voorhis WC, Centurion-Lara A, et al.

Quantitative analysis of *tpr* gene expression in *Treponema pallidum* isolates: differences among

isolates and correlation with T-cell responsiveness in experimental syphilis. Infect Immun.

2007;75(1):104-12.

53. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence

data. Bioinformatics. 2014;30(15):2114-20.

54. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.

2012;9(4):357-9.

55. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al. Geneious

Basic: an integrated and extendable desktop software platform for the organization and analysis

of sequence data. Bioinformatics. 2012;28(12):1647-9.

56. Lieberman NAP, Lin MJ, Xie H, Shrestha L, Nguyen T, Huang ML, et al. *Treponema*

pallidum genome sequencing from six continents reveals variability in vaccine candidate genes

and dominance of Nichols clade strains in Madagascar. PLoS Negl Trop Dis.

2021;15(12):e0010063.

1003

A

B

A tp0127:dnaA (white) kan^R:dnaA (black)

Copy number ratio — Parent SS14, Passage #: P2, P3, P4, P5; SS14-DC^KO, Passage #: P2, P3, P4, P5

B SS14-DC^KO (black), SS14 (white); $p > 0.05$; tprK mRNA copies/100 tp0574 copies

C KO genome vs WT reference; DC - containing region; Read Depth; Genome Position

D KO genome vs KO reference; kan^R; Read Depth; Genome Position

E V1 (red), V2 (blue), V3 (green), V4 (yellow), V5 (orange), V6 (purple), V7 (brown); TP0126d, TP0126c, TP0127, TP0128, TP0129, TP0130; p47p, kan^R

F WT genome vs WT reference; DC - containing region; Read Depth; Genome Position

G WT genome vs KO reference; kan^R; Read Depth; Genome Position

A

B

SS14-DC^KO –infected rabbits (day 72 PI)   SS14–infected rabbits (day 72 PI)

A SS14-infected rabbits
V6 peptide PVHWKALARAQPPARVDIY

B SS14-infected rabbits
V5 peptide QASNVFQGVFLTTPMQKDDC

C SS14-infected rabbits
V5 peptide QASNVFQGVFLTTPMLQHDC

D SS14-infected rabbits
Reactivity to V6 peptide PVHWKALARAQPPARVDIY
Reactivity to rTprK (aa 21-252)

E SS14-infected rabbits
Reactivity to V5 peptide QASNVFQGVFLTTPMQKDDC
Reactivity to rTprK (aa 21-252)

F SS14-infected rabbits
Reactivity to V5 peptide QASNVFQGVFLTTPMLQHDC
Reactivity to rTprK (aa 21-252)

G SS14-DC^{KO}-infected rabbits
V6 peptide PVHWKALARARPPVPAIY

H SS14-DC^{KO}-infected rabbits
V5 peptide KASNVFKDVFLTDTTPMQTHDC

I SS14-DC^{KO}-infected rabbits
V5 peptide QASNVFQGVFLTTPMLQHDC

J SS14-DC^{KO}-infected rabbits
Reactivity to V5 peptide PVHWKALARARPPVPAIY
Reactivity to rTprK (aa 21-252)

K SS14-DC^{KO}-infected rabbits
Reactivity to V5 peptide KASNVFKDVFLTDTTPMQTHDC
Reactivity to rTprK (aa 21-252)

L SS14-DC^{KO}-infected rabbits
Reactivity to V5 peptide QASNVFQGVFLTTPMLQHDC
Reactivity to rTprK (aa 21-252)