1
2
Comparison of gene set scoring methods for reproducible evaluation of multiple tuberculosis gene signatures

Xutao Wang, M.Sc[1,2]; Arthur VanValkenberg, Ph.D.[2], Aubrey R. Odom-Maybe[2], Jerrold

J. Ellner, M.D.[3]; Natasha S. Hochberg, M.D., M.P.H.[4,5]; Padmini Salgame, Ph.D[3];

Prasad Patil, Ph.D[1,*]; W. Evan Johnson, Ph.D[6,*]

Institutions:

[1]Department of Biostatistics, Boston University, Boston, MA, USA

[2]Division of Computational Biomedicine and Bioinformatics Program, Boston University,

Boston, MA, USA

[3]Department of Medicine, Center for Emerging Pathogens, Rutgers New Jersey Medical

School, Newark, NJ, USA

[4]Boston Medical Center, Boston, MA, USA

[5]Section of Infectious Diseases, Boston University School of Medicine, Boston, MA,

USA

[6]Division of Infectious Disease, Center for Data Science, Rutgers New Jersey Medical

School, Newark, NJ, USA

* WEJ and PP are co-senior authors

18
Corresponding Author: W. Evan Johnson, Ph.D., New Jersey Medical School, 225

Warren Street 450G, Newark, NJ, USA 07103. Email: w.evan.johnson@rutgers.edu

Authors Contribution: WEJ and PP conceived the study. XW conducted the

comparisons and analysis of TB signatures and generated the figures. All authors

contributed to the study design and guided the interpretation of the results. XW, PP and

WEJ wrote the manuscript and generated figures and results. All authors read and

approved the final manuscript.

Funding Sources: This work was supported by funds from the NIH (R01GM127430

and R21AI154387)

29
30   **Keywords:** Tuberculosis, genet set scoring methods, gene signatures, reproducibility

ABSTRACT

Rationale: Many blood-based transcriptional gene signatures for tuberculosis (TB) have been developed with potential use to diagnose disease, predict risk of progression from infection to disease, and monitor TB treatment outcomes. However, an unresolved issue is whether gene set enrichment analysis (GSEA) of the signature transcripts alone is sufficient for prediction and differentiation, or whether it is necessary to use the original statistical model created when the signature was derived. Intra-method comparison is complicated by the unavailability of original training data, missing details about the original trained model, and inadequate publicly-available software tools or source code implementing models. To facilitate these signatures' replicability and appropriate utilization in TB research, comprehensive comparisons between gene set scoring methods with cross-data validation of original model implementations are needed.

Objectives: We compared the performance of 19 TB gene signatures across 24 transcriptomic datasets using both re-rebuilt original models and gene set scoring methods to evaluate whether gene set scoring is a reasonable proxy to the performance of the original trained model. We have provided an open-access software implementation of the original models for all 19 signatures for future use.

Methods: We considered existing gene set scoring and machine learning methods, including ssGSEA, GSVA, PLAGE, Singscore, and Zscore, as alternative approaches to profile gene signature performance. The sample-size-weighted mean area under the curve (AUC) value was computed to measure each signature's performance across datasets. Correlation analysis and Wilcoxon paired tests were used to analyze the performance of enrichment methods with the original models.

Measurement and Main Results: For many signatures, the predictions from gene set scoring methods were highly correlated and statistically equivalent to the results given by the original diagnostic models. PLAGE outperformed all other gene scoring methods.

In some cases, PLAGE outperformed the original models when considering signatures'

weighted mean AUC values and the AUC results within individual studies.

Conclusion: Gene set enrichment scoring of existing blood-based biomarker gene sets

can distinguish patients with active TB disease from latent TB infection and other clinical

conditions with equivalent or improved accuracy compared to the original methods and

models. These data justify using gene set scoring methods of published TB gene

signatures for predicting TB risk and treatment outcomes, especially when original

models are difficult to apply or implement.

INTRODUCTION

Prior to the COVID-19 pandemic, tuberculosis (TB) was the leading infectious cause of

death worldwide (1, 2). Approximately 10 million people developed TB, and 1.4 million

died from the disease (1). Current diagnostic tests for TB disease include sputum acid-

fast bacilli (AFB) smear microscopy, rapid molecular tests, and culture-based

technologies (1). With the advent and widespread availability of nucleic acid

amplification tests (Xpert MTB/RIF), most cases of pulmonary TB can be diagnosed

quickly and accurately. Paucibacillary TB diagnosis (smear-negative pulmonary,

extrapulmonary, and pediatric TB) (3–6) and predicting treatment success/failure remain

difficult challenges. Furthermore, there are gaps in identifying individuals with quiescent

or percolating disease, and utilizing non-sputum-based technologies would facilitate

diagnosis for individuals unable to produce sputum (e.g., children). There is therefore an

urgent need for additional technologies that ensure high-quality, timely, effective testing

for people living with TB (1, 7).

Over the last ten years, multiple blood-based biomarkers using gene expression profiles

have been developed for TB. These signatures can distinguish active TB disease from

latent TB infection (LTBI) (8, 9), distinguish TB from other diseases (OD) (10–12),

predict progression from LTBI to active TB (13, 14), and assess treatment response

(15). These or other signatures may potentially meet target product profiles (TPP)

proposed by the World Health Organization (WHO) for point-of-care (POC) testing,

since some studies have shown that perturbation in the gene expression profile occurs

prior to sputum conversion (13, 14, 16, 17). Blood-based signatures could also identify

individuals for additional diagnostics or LTBI treatment. However, more research must

be done to establish the efficacy and reproducibility of using blood-based signatures in

the field, as shown by the CORTIS trial (18).

The common approach to building a gene signature is to identify a *gene set*, a subset of

the thousands of genes whose expression is measured for a patient cohort, using a

combination of domain knowledge and data-driven feature selection methods. The gene

set is then used to train a gene signature using a *prediction algorithm* such as

regression or some other machine learning approaches for deciding how to turn gene

expression measurements into an outcome prediction. Typically, biomarkers are *trained*

using one subset of data and then finalized or *fixed* before being applied for prediction

in new samples and cohorts. However, in the case of existing TB signatures, the

replicability of these biomarkers is inadequate, meaning that many of the original

publications did not give enough detail to replicate the published models. In addition,

nearly all of these publications failed to provide usable code to rerun the published

model on the study data—let alone apply the models to new datasets. Some of these

original models further lacked *reproducibility* of the performance and accuracy of TB

gene signatures—meaning that the signatures were *overfit* and thus experienced

significant reductions in performance in later observations compared to the training set

(10, 19, 20). Many factors likely contribute to this phenomenon, including overfitting to

encodes in the training data, incomplete validation of the biomarker, differences in study

population demographics or comorbidities, and possible inconsistencies due to

differences in the platforms used to measure gene expression (37, 46).

Several research teams have attempted to address issues of reproducibility in TB gene

expression biomarker research, either by rebuilding the original classification models (7,

19) or by using methods such as gene set enrichment analysis (GSEA) in place of the

original models (21). For example, Warsinske et al. (19) reconstructed the original

models for 16 TB gene signatures and compared gene signature accuracy in the

contexts of TB diagnostics and progression to disease. For some of their biomarkers,

the discovery dataset(s) used to train the gene signatures were unavailable, or the

proposed diagnostic models comprised the gene signatures were difficult to reproduce.

Other challenges included limited information about data preprocessing and

hyperparameter values used to build the model (22–24). Despite clear documentation of

their reconstruction of original models, Warsinske et al. (19) did not publish software

code for their work from scratch to use the biomarkers from their comparisons.

As a result of this deficit in replicability, we have endeavored to fill this gap. Our team

has recently developed and released the open-source TBSignatureProfiler software

toolkit, which provides a compilation of TB gene sets used from published biomarkers

and provides methods (21) to evaluate the performance of existing TB gene sets on

new datasets via the application of various scoring algorithms (25) (e.g., GSVA,

ssGSEA, PLAGE). However, while alternative methods, such as gene set scoring are

simpler to use compared to the original models, these methods have not been

established as reasonable approximations or alternatives to original model

performance.

To address the issues of reproducibility in reconstructing the discovery set, our study

uniformly evaluated the performance of 19 TB gene signatures across 24 datasets

using both original models and gene set scoring methods through the

TBSignatureProfiler platform. We systematically compared our results to previous

research (19) to the signature's ability to distinguish patients with active TB from other

clinical conditions. Our results illustrated that gene set scoring methods are effective

computational tools for profiling the performance of TB gene signatures and produced

similar accuracy compared to the original model, demonstrating alternative, accessible

ways to profile existing TB gene signatures. We also sought to facilitate the replicability

of TB gene signatures' application and comparative analysis. We therefore curated the

TB transcriptomic datasets used in this study and included the corresponding discovery

model for each gene signature in the TBSignatureProfiler R package, enabling the

reproducibility of all results.

METHODS

Gene signatures and original diagnostic models used for comparison

Nineteen existing TB gene signatures were selected for this study primarily based on

the results of Warsinske *et al.* (Table 1) (10, 12–14, 22–24, 26–32) to make a fair

comparison for the performance of these signatures. Sixteen gene signatures were

trained to predict active TB versus other clinical conditions. Three additional gene

signatures that predict progression from LTBI to active TB disease, Zak_RISK_16 (14),

Suliman_RISK_4 (13), and Leong_RISK_29 (32), were also included. Twenty-four

datasets (9–12, 16, 21, 27–29, 31, 33–48) that include subjects with active TB and other

clinical conditions have been selected for evaluation. These datasets were selected in

alignment with Warsinske *et al.* to systematically compare different gene scoring

methods and the original models (19). Twenty of these datasets were from microarray

studies; the other four consisted of RNA-sequencing data. A detailed description of the

transcriptomic studies included in this study can be found in the Supplementary

Material.

The reconstruction of each signature's original model followed the guidelines given by

Warsinske et al. (19), who reconstructed these models based on information from each

of the signature's original publications. Two training methods were used for TB gene

signature discovery: model-based and score-based approaches. The model-based

approaches use machine learning algorithms such as support vector machines (20) or

random forests (49) to reconstruct the biomarker's original model. The score-based

approaches primarily rely on the difference of sums or arithmetic means between the

upregulated genes and the downregulated genes within the gene set and therefore do

not require model retraining. Six of the 19 identified gene signatures used score-based

methods to distinguish active TB patients from other clinical conditions (Table 1). The

reconstructed models were validated against the performance reported in the original

publication using the AUC metric. To evaluate a biomarker's ability to identify patients

with active TB, the associated model was applied across the 24 curated transcriptomic

studies.

Gene set scoring methods

We used single sample GSEA (ssGSEA) (50), Gene Set Variation Analysis (GSVA)

(25), Pathway level analysis of gene expression (PLAGE) (51), Zscore (52), and

Singscore (unidirectional and bidirectional versions) (50, 53) to evaluate the accuracy of

TB gene signatures across 24 studies. Please see the Discussion section for a more

detailed comparison of these five gene set scoring methods. The evaluation of distinct

gene signatures using different gene set scoring methods and its original model was

performed by the runTBSigProfiler function from the TBSignatureProfiler R package

(21).

Gene signature splitting strategy

Some gene set scoring methods, including GSVA, ssGSEA, and Singscore, rank a

signature's genes against other genes in the dataset. Thus, gene sets with both

upregulated and downregulated genes do not usually score well with these methods—

they effectively counteract each other in the score computation. Most of the 19

signatures contain genes expressed strongly in both directions; thus, we proposed a

'signature splitting' strategy for gene signatures with equal to or more than ten genes to

overcome this limitation. Gene signatures with less than ten genes were excluded here

as their upregulated or downregulated gene subset was either empty or only contained

one gene. In such cases, signature splitting would have a decreased effect on these

gene sets compared to a large set.

For the ssGSEA and GSVA methods, we evaluated each signature's performance using

only the upregulated or the downregulated subsets of genes within the gene sets. For

gene signatures that did not require reconstruction of the training model, the

upregulated and downregulated genes were usually provided by the original publication.

If unavailable in the original publication, the upregulated or the downregulated genes

were identified by performing differential expression analysis on the datasets from which

the associated biomarkers were derived using the R package DESeq2 (54) for RNA-seq

data and limma (55) for microarray data. For the bidirectional implementation of

Singscore, we used the simpleScore function from the singscore R package (53) to

algorithmically combine the disease score for the upregulated and downregulated

genes.

Imputation of missing genes and removal of batch effects

Occasionally, various genes within gene sets were missing across transcriptomic

studies, mainly due to the inconsistent gene coverage across platforms or different

versions of the gene symbol naming systems across different sequencing platforms.

This might affect a biomarker's predictive accuracy in some datasets and across

biomarker methods. Two steps were taken to mitigate this issue. First, we updated gene

symbols for gene signatures and candidate datasets using the HGNChelper R package

(56) to reduce gene misidentification during the training and validation process. More

details on the matching of probe ID to gene symbol and missing gene information within

each dataset can be found in the Supplementary Material. Second, a k-nearest

neighbors (KNN) algorithm was applied to impute the missing gene expression values

based on the signature's discovery dataset(s) (57). This procedure may elicit potential

bias problems when large portions of genes within a signature are missing in some

datasets. To make a fair comparison, we applied this imputation procedure to all

biomarkers that used model-based diagnostic approaches across independent

datasets. Functionality *reference ComBat* (58, 59) was used to make generalized

comparisons of signatures across the independent studies. *reference ComBat* removes

batch effects between the training datasets and the testing datasets without changing

the reference dataset's gene expression values. With this functionality, the biomarker's

original model does not have to be retrained whenever the batch correction is

performed or when the model is applied to a new dataset, increasing efficiency when

evaluating biomarkers with model-based approaches.

Statistical Evaluation of Model Performance

The AUC value corresponding to sample scores against disease subtypes for each TB

gene signature was calculated for each dataset. The sample-size-weighted mean AUC

(weighted AUCs) was used to assess the overall performance of each gene set across

independent studies while excluding the discovery dataset(s) used to train the

corresponding signature (19). AUC results were calculated with the R package ROCit

(60) compute estimates. Each signature's 95% confidence interval for the weighted

mean AUC value was calculated using the bootstrap method with 10,000 bootstrap

resamples. The Wilcoxon paired test was applied to test whether AUCs given by

different methods were significantly different from each other.

Evaluating biomarker performance based on original models

The original models were inputted into the TBSignatureProfiler (21) platform, which

provides a computational profiling platform that allows researchers to evaluate multiple

TB gene signatures. In addition to the original models, users can access other scoring

methods and gene sets in the TBSignatureProfiler. The TBSignatureProfiler was used

to conduct all the analyses for this work, and the package can be downloaded from

GitHub (https://github.com/compbiomed/TBSignatureProfiler) or Bioconductor

(http://www.bioconductor.org/packages/release/bioc/html/TBSignatureProfiler.html).

Code for the analyses from this paper can be found at: https://github.com/xutao-

wang/Comparison-of-existing-tuberculosis-gene-signatures.

Comparing the performance of gene set scoring methods with original models

We noticed that there are several studies (i.e. GSE81746, GSE107331, GSE29536 etc.)

with a small sample sizes (n < 40) which only consisted of healthy controls (HCs) and

subjects with active TB. For these studies, most was generalizable to gene signatures

that could distinguish patients with active TB phenotype with high predictive accuracy;

however, such performance could not be generalized toward large populations or to

cohorts with heterogeneous disease types including LTBI. To produce a fair comparison

of gene signatures' performance using gene set scoring methods and their original

models, for each profiling method (including both gene set scoring methods and original

models) we selected datasets with a sample size greater than 40 and AUC estimates

greater than 0.80 across all TB biomarkers simultaneously excluding the corresponding

training study for each biomarker.

Several metrics were used to compare the performance of gene signatures as assessed

by different gene set scoring methods and their original models. For each TB gene

signature, the Spearman's rank correlation (ρ) was computed to measure the strength

of association of the prediction scores from a signature's original model and the different

gene set scoring methods. We then summarized correlation results, computing the

weighted Spearman's rank correlation (ρ_w), based on equation (1).

$$\rho_w = \frac{\sum_{i=1}^{k} \rho_i * n_i}{\sum_{i=1}^{k} n_i}$$

282   (1)

Moreover, the absolute AUC difference ($|\Delta AUC|$) between the original model and

various gene set scoring methods were computed as shown in equation 2.1 for each

selected dataset, with the weighted absolute AUC difference $|\Delta AUC|_w$ representing the

overall distribution pattern across all selected studies (equation 2.2). Additionally, the

Szymkiewicz–Simpson coefficient (61) or overlap coefficient (oc) as shown by equation

(3), was applied to evaluate the similarity of studies based on the results given by each

of the gene set scoring method and original model for all gene signatures.

$$|\Delta AUC_i| = |AUC_{GSEA,i} - AUC_{original\ model,i}|$$

291   (2.1)

292 $$|\Delta AUC|_w = \frac{\sum_{i=1}^{k} |\Delta AUC_i| * n_i}{\sum_{i=1}^{k} n_i}$$

293   (2.2)

294 $$overlap\ coefficient = \frac{\#\ of\ common\ datasets}{min(n_{GSEA}, n_{original\ model})}$$

295   (3)

RESULTS

Evaluation of discovery studies using training models from original publications

The list of signatures and information on their respective training data are presented in

Table S1, along with AUC values from their originating publication, Warsinske *et al.*, and

our results. The AUC values for our reconstructed models were nearly identical to the

results of the original publications, suggesting that the training models were accurately

reconstructed. Several gene signatures including Sweeney_OD_3, Maertzdorf_15,

Leong_24, Kaforou_27, Anderson_42, and Berry_393, when evaluated by ssGSEA, had

estimated AUC values above 0.9 (Table S1). These results suggest that ssGSEA is a

comparable signature profiling method, producing accurate results for some TB

signatures in differentiating the TB disease states (active TB vs. LTBI, HCs, or OD)

even if the evaluation studies were the signature's discovery datasets.

Performance of original models and gene set scoring methods for active TB

versus non-active TB

Figure 1 shows the AUC distribution of 24 independent TB studies across 19 TB gene

signatures based on the results from their original models. 13 of 16 signatures had

AUCs greater than 0.9 from their discovery dataset(s) (Figure 1). Each TB signature's

original model was applied across all the datasets, and their performance varied greatly

on a study-by-study basis (Figure 1). Notably, the Kaforou_OD_53, Kaforou_27,

Maertzdorf_15, and Sweeney_OD_3 signatures had consistently high AUC values

across different studies, with weighted AUCs of 0.83 for Kaforou_OD_53, 0.81 for

Kaforou_27, 0.82 for Maertzdorf_15, and 0.81 for Sweeney_OD_3 (Table 2). In

contrast, the performance of some signatures, such as Verhagen_10, was mixed across

studies (Figure 1). Verhagen_10 had a weighted mean AUC of 0.61 (Table 2),

performing well in some datasets (>0.9 AUC in GSE81746, GSE41055, and GSE29536)

but with poor performance in most of the remaining studies (<0.65 AUC in 16 out of 24

studies; Figure 1). Noted that Zak_RISK_16, Suliman_RISK_4, and

Leong_PREDICT_29 also performed poorly in these comparisons, but these are

signatures of disease progression and were not designed to distinguish between active

and LTBI (Figure 1).

Table 2 shows the weighted AUCs for each gene signature as computed using their

corresponding original model and five different gene set scoring methods. In this table,

AUC values in red denote results from our reimplemented models and gene set scoring

methods that outperformed the original model with differences greater than 0.05 units.

Values in blue represent cases where AUC values from the original model outperformed

other methods the methods by more than 0.05 units. The 0.05 threshold was chosen

mainly for visualization purposes, which suggested that there was a noticeable

difference in the weighted AUCs given by two different methods.

16 of 19 gene signatures had reimplemented weighted AUCs within 0.05 units

compared to Warsinske *et al*. (Table 2). However, our reimplemented models for

Verhagen_10 and Anderson_OD_51 outperformed the results from Warsinske *et al*.,

and our original model for Jacobsen_3 underperformed compared to Warsinske *et al*.

(Table 2). The weighted AUCs for Anderson_OD_51 computed by gene scoring

methods surpassed that of its original model but not the results given by ssGSEA,

although none of the results were statistically significant after adjusting for multiple

testing (p-value > 0.01; Table 2). For Berry_393, the weighted AUCs computed from its

original model underperformed all the five gene scoring methods; specifically, the

PLAGE AUC was 0.79 (0.75–0.84), which was significantly higher (p-value <= 0.01)

than that of the original model which had an AUC of 0.70 (0.66–0.74). Results from

Zak_RISK_16 given by its original model also underperformed five gene scoring

methods, and all the superior weighted AUCs were significant (p-value <= 1e-04) (Table

2). Additionally, the performance of each gene signature across individual datasets

using five gene set scoring methods was presented in Figure S1A-E.

Figure 2A shows the weighted mean AUC difference between the original model and

the corresponding five gene scoring methods across all gene sets. The outperformance

of the weighted AUCs given by PLAGE was consistent in most of the gene signatures

asides from Sambarey_HIV_10 and Anderson_42. The accompanying Wilcoxon signed-

rank test results indicated that the results from PLAGE were significantly different from

the results of the original models (p-value <= 0.01), whereas the results from the

remaining four gene scoring methods were statistically equivalent when compared to

the weighted AUCs given by the original model (p-value > 0.05; Figure 2A).

Accounting for gene expression direction

Table 3 shows the weighted mean AUCs from ssGSEA and GSVA after splitting the

genes into upregulated or downregulated subsets, along with the Singscore bidirectional

scoring results. There was significant improvements in the weighted AUCs for the

upregulated subsets of Berry_OD_86 based on the ssGSEA (AUC=0.78; 95% CI:0.75–

0.82) and GSVA (AUC=0.80; 95% CI: 0.76–0.84) results when compared to its original

models (p-value <= 1e-04 for both methods; Table 3). Similar improvement was

observed when using ssGSEA and GSVA for the upregulated subsets of Berry_393 (p-

value <= 0.01 for both methods). Interestingly, Anderson_OD_51 is the only gene

signature for which the weighted AUCs from GSVA (upregulated subsets: AUC = 0.81,

95% CI: 0.78-0.84; p-value = 0.03; downregulated subsets AUC = 0.77, 95% CI: 0.71-

0.82; p-value = 0.04) outperformed its original model (Table 3). Lastly, Berry_OD_86

and Berry_393 also outperformed their original models when was using Singscore

bidirectional scoring the signatures' (p-value <= 1e-3 for both gene sets; Table 3). Most

gene signatures had similar results when using the unidirectional and bidirectional

versions of Singscore, although Sambarey_HIV_10 and Anderson_OD_51 showed

significantly lower weighted AUCs using Singscore bidirectional scoring (p-value < 0.01

for both gene sets; Tables 2 and 3).

Figure 2B illustrates the weighted mean AUC differences between each gene scoring

method and original models across gene signatures. Again, the Wilcoxon signed-rank

test suggests that the weighted mean AUC results from all methods were statistically

equivalent (p-value > 0.05) to the results given by the original model, except for the

results computed from the signatures' down-regulated subset using ssGSEA (p-value =

0.036; Figure 2B). Additionally, Figure S2A-C shows the AUC distribution for the

up/down-regulated subset of each qualified gene signature across all datasets using the

gene set scoring methods ssGSEA, GSVA, and Singscore bidirectional scoring.

Finally, ridge plots created displayed the AUC distribution across 24 studies using

original models and all the five gene set scoring methods (Figure S3A-C). The gene

signatures were ranked in increasing order based on their median AUC values over the

individual training dataset used by the given original model. Signatures including

Kaforou_27, Kaforou_OD_53, and Sambarey_HIV_10 had statistically equivalent AUC

distributions for results obtained from their respective original models, PLAGE, and

ssGSEA with the upregulated subset (Kaforou_27 p-value = 0.54, Kaforou_OD_53 p-

value = 0.93, Sambarey_HIV_10 p-value = 0.88; Figure S3A).

Gene set scoring methods versus original models

Figure S4A-E shows the performance of 19 gene signatures after selection, as based on

the criteria described in the methods section. Each point represents a single study; the

y-axis gives Spearman's rank correlation of the predicted scores for the original model

and respective gene set scoring method, and the x-axis gives the AUC difference

between the results given by two different methods. As expected, Verhagen_10 had

poor performance when evaluated using its original model, Zscore, and Singscore.

None of the studies had AUCs greater than 0.8 after selection (Figure S4A-E).

GSE101705 was the only study selected, but it showed a distinct prediction pattern

given by ssGSEA/GSVA when compared to its original model ($0.18 <= |\Delta AUC| <= 0.21$;

$0.35 <= \rho <= 0.57$). Similar mediocre performance was observed for Suliman_RISK_4,

where GSE107994 was the only study selected by ssGSEA and its original model

($|\Delta AUC| = 0.061$; $\rho = 0.7$; Figure S4A), and no studies were selected using GSVA and

Zscore (Figure S4C-D). In addition, no studies had AUCs greater than 0.8 for

Leong_RISK_29 when evaluated using ssGSEA, GSVA, and Zscore (Figure S4A;

Figure S4C-D)

ssGSEA vs. Original Model

Kaforou_27, Maertzdorf_15, and Sweeney_OD_3 showed high diagnostic accuracy in

similar studies (oc for Kaforou_27 and Maertzdorf_15: = 1.00, oc for Sweeney_OD_3: =

0.86) with small AUC differences (p-value > 0.28 for three gene sets) given by ssGSEA

and their original models. Significantly positive Spearman's rank correlations of the

predicted scores were shown for Kaforou_27 (ρ_w = 0.78) and Sweeney_OD_3 (ρ_w =

0.86), and significantly negative correlations were produced for Maertzdorf_15 (ρ_w = -

0.81; p-value <= 1e-04 for three signatures; Figure S4A). Although more studies with

high AUCs were selected for Anderson_OD_51, Berry_393, Berry_OD_86,

Bloom_OD_144, Leong_24, and Zak_RISK_16 using ssGSEA, results from these six

signatures showed distinct predicted results when compared to their original models

(0.032 <= ρ_w <= 0.60; $|\Delta AUC|_w$ > 0.050; Figure S4A). Conversely, more datasets were

selected for Kaforou_OD_44, Kaforou_OD_53, LauxdaCosta_OD_3, and Maertzdorf_4

based on the results from these signatures' original models (0.070 <= ρ_w <= 0.72,

$|\Delta AUC|_w$ > 0.050; Figure S4A).

PLAGE vs. Original Model

For gene sets Kaforou_27, Kaforou_OD_53, LauxdaCosta_OD_3, Maertzdorf_4, and

Maertzdorf_15, PLAGE results were nearly equivalent to the results from their original

models (Figure S4B). The study overlap coefficient was 1.00 for all five gene signatures,

asides from Kaforou_OD_53, with an AUC value of 0.85 (Figure S4B). Specifically,

Kaforou_27, Kaforou_OD_53, and LauxdaCosta_OD_3 had significantly negative

Spearman's rank correlations for the predicted results given by two methods (-0.92 < ρ_w

< -0.82; p-value < 1e-04 for all three gene sets), with an absolute weighted AUC

difference less than 0.030 for all three gene sets. Two Maertzdorf signatures had strong

positive Spearman's rank correlations of the prediction scores and small AUC

differences (p-value for Maertzdorf_4: = 0.06, p-value for Maertzdorf_15: = 0.02), with

ρ_w of 0.88 for Maertzdorf_4 and of 0.94 for Maertzdorf_15 (p-value < 1e-05; Figure

S4B). Asides from these gene sets, 11 of 14 signatures (excluding Anderson_42,

Kaforou_OD_44, and Sambarey_HIV_10) had a higher number of studies with high

AUCs selected by PLAGE but showed large absolute AUC differences ($|\Delta AUC|_w$ >

0.080) and weak to moderate correlations (-0.68 <= ρ_w <= 0.62) of the profiling scores

when compared to the results from their original methods (Figure S4B).

GSVA vs. Original Model

When comparing the results given by GSVA and signatures' original models, more

studies were selected for Anderson_OD_51, Berry_393, Berry_OD_86,

Bloom_OD_144, and Zak_RISK_16 with weighted absolute AUC differences all greater

than 0.050 and weak Spearman's rank correlation (0.11 <= ρ_w <= 0.47) of the predicted

scores (Figure S4C). Similar to that of ssGSEA, the resulting scores from

Maertzdorf_15 (ρ_w: -0.78, $|\Delta AUC|_w$: 0.043) and Maertzdorf_4 (ρ_w: -0.63, $|\Delta AUC|_w$:

0.051) were negatively correlated with the results from the signatures' original model,

with a study overlap coefficient of 1.00 for both gene signatures (Figure S4C).

Zscore vs. Original Model

The performance of gene signatures using Zscore was similar to GSVA, for which more

datasets were selected for Anderson_OD_51, Berry_393, Berry_OD_86,

Bloom_OD_144, and Zak_RISK_16. However, a distinct prediction pattern was visible

when comparing these to their original methods (-0.0037 <= ρ_w <= 0.65, $|\Delta AUC|_w$ >

0.05 for all five gene sets; Figure S4D). Similar to the results from ssGSEA and GSVA,

Maertzdorf_15 (ρ_w = -0.85, $|\Delta AUC|_w$ = 0.044) and Maertzdorf_4 (ρ_w = -0.74, $|\Delta AUC|_w$ =

0.084) had negatively correlated prediction scores given using Zscore and their

respective original models (Figure S4D).

Singscore unidirectional scoring vs. Original Model

Although more studies with high AUCs were selected when assessing

Anderson_OD_51, Berry_393, Berry_OD_86, Bloom_OD_144, and Zak_RISK_16 with

Singscore unidirectional scoring, results from these signatures had weak to moderate

weighted Spearman's rank correlation (0.16 <= ρ_w <= 0.55) and a large AUC difference

($|\Delta AUC|_w$ >= 0.11 for all five gene sets; Figure S4E). Conversely, no datasets with high

AUCs were selected for Anderson_42 (ρ_w = -0.23, $|\Delta AUC|_w$ = 0.26) based on results

given by Singscore unidirectional scoring. Sweeney_OD_3 (ρ_w = 0.85, p-value < 1e-04;

$|\Delta AUC|_w$ = 0.031, p-value = 0.75) was the only gene set for whom results from

Singscore could act as a proxy to its original models, except for study GSE34608

(Figure S4E).

Gene signatures after splitting

Figure S5A-E shows the performance of 13 signatures as evaluated with their up/down

regulated subset of genes using ssGSEA, GSVA, and Singscore bidirectional scoring.

Notably, for the upregulated subset of Verhagen_10 , none of the datasets had AUCs

greater than 0.8, as given by ssGSEA and GSVA (Figure S5A-D). Study GSE101705

was the only selected study when evaluating the down-regulated subset of

Verhagen_10 using ssGSEA (ρ = 0.60, $|\Delta AUC|$ = 0.20), GSVA (ρ = 0.57, $|\Delta AUC|$ =

0.17), and Singscore bidirectional scoring (ρ = -0.63, $|\Delta AUC|$ = 0.18; Figure S5B-S5E).

However, it showed different prediction results than the original models. Additionally,

GSE107994 was selected for the downregulated subset of Verhagen_10 given by

GSVA (ρ = 0.44, $|\Delta AUC|$ = 0.19; Figure S5D).

ssGSEA with signature split vs. Original Model

When assessing gene signatures based on their upregulated subset, all selected

studies had positive Spearman's rank correlations of the predicted scores given by

ssGSEA and original models (Figure S5A). More studies with high AUCs were selected

for 10 out of 12 gene signatures (excluding Leong_RISK_29 and Maertzdorf_15) when

evaluated with their respective upregulated genes using ssGSEA. Results from subsets

of Kaforou_27 and Kaforou_OD_53 showed similar diagnostic features when compared

to their original models, with highly correlated prediction scores (ρ_w > 0.80 for both gene

sets; p-value <= 1e-04), AUC differences around 0 ($|\Delta AUC|_w$ < 0.030 for both gene sets;

p-value > 0.080), and a study overlap coefficient of 1.00 for both gene sets (Figure

S5A).

When gene signatures were evaluated using their downregulated subsets, only

Maertzdorf_15 had an absolute Spearman's rank correlation greater than 0.80 (ρ_w: -

0.89, p-value <= 1e-05) and a small AUC difference ($|\Delta AUC|_w$: 0.033, p-value = 0.010)

for results given by ssGSEA and its original model, with a study overlap coefficient of

1.00 (Figure S5B). Only Berry_393 (ρ_w: -0.62), Leong_24 (ρ_w: -0.64), and Verhagen_10

(ρ_w: 0.60) had a more studies with high AUCs using ssGSEA (0.052 < $|\Delta AUC|_w$ < 0.191;

Figure S5B).

GSVA with signature split vs. Original Model

Figure S5C shows that the predicted scores given by GSVA and original models are

positively correlated for most of the selected studies except GSE34608, for which the

study was negatively correlated when evaluated by the upregulated subsets of

Leong_24 (ρ = -0.11, $|\Delta AUC|$ = 0.25). Moreover, Kaforou_27 had the highest weighted

Spearman's rank correlation (ρ_w = 0.90, p-value < 1e-05) and the smallest AUC

difference ($|\Delta AUC|_w$ = 0.020, p-value = 0.68) among 12 gene sets when its upregulated

subset was assessed by GSVA, which presented an equivalent prediction pattern

compared to its original model (Figure S5C). The upregulated subset of Maertzdorf_15

was the only gene set whose original model results outperformed GSVA, which had a

greater number of studies with high AUCs from its original method (ρ_w = 0.68, $|\Delta AUC|_w$

= 0.094; Figure S5C).

When gene signatures were evaluated with their down-regulated subsets, the results

from most of the gene sets given by GSVA underperformed the results from their

original models. Specifically, nine of 13 gene sets had a greater number of studies with

high AUCs based on the results from their original model (Figure S5D). Notably, no

datasets with high AUCs were selected when the down-regulated subset of

Bloom_OD_144 and Kaforou_OD_44 were assessed by GSVA. GSE28623 was the

only study with high AUCs for the down-regulated subset of Anderson_42 (Figure S5D).

Maertzdorf_15's GSVA results, similar to those for the ssGSEA method, had the highest

absolute weighted Spearman's rank correlation (ρ_w = -0.88, p-value <= 1e-05) and the

smallest absolute AUC difference ($|\Delta AUC|_w$ = 0.03, p-value = 0.20) when compared to

its original model, with a study overlap coefficient of 1.00 (Figure S5D).

Singscore bidirectional scoring vs. Original Models

Most of the selected studies had positive correlations for the predicted scores as given

by Singscore bidirectional scoring and their original methods, except study GSE62525

from Bloom_OD_144 and GSE101705 from Verhagen_10 (Figure S5E). Both

Kaforou_27 and Kaforou_OD_53 had a weighted Spearman's rank correlation greater

than 0.85 (p-value < 1e-05 for both gene sets), absolute AUC difference smaller than

0.020 (p-value > 0.05 for both gene sets), and study overlap coefficient of 1.00, which

suggests that results for these gene signatures given by Singscore bidirectional scoring

could act as a proxy from the results given by signatures' original models (Figure S5E).

Signatures Anderson_OD_51, Kaforou_OD_44, Maertzdorf_15, and Sambarey_HIV_10

had a larger number of studies with high AUCs given by their original model, weighted

Spearman's rank correlations between 0.53 and 0.86, and absolute AUC differences

ranging from 0.021 to 0.16 (Figure S5E).

DISCUSSION

TB diagnostics is moving toward using blood-based biomarkers, but serious gaps

remain in the analyses of these data. Here we evaluated the performance of 19 TB

gene signatures in distinguishing active TB from other clinical conditions using the

original published model and five different gene set scoring methods based on 24

transcriptomic studies. These datasets represent real-world heterogeneity concerning

geographic regions, host and pathogen genetics, and clinical context (19). Our results

suggested that an original gene signature model's predictive ability can be improved or

recaptured using gene set scoring methods.

The five gene set scoring methods used here belong to a class of methods that

compute a gene set enrichment score for each sample using only the genes from a

signature. However, some differences between methods are present. Gene set scoring

methods, including ssGSEA, GSVA, Zscore, and Singscore are *single-sample* methods

that rank genes in each sample individually by comparing the ranks of the signature

genes with the ranks of non-signature genes in the sample. By design, using these

single-sample methods, the score for an individual sample is not affected by other

samples in the cohort. In contrast, in computing an individual sample score, PLAGE

implements singular value decomposition (SVD) on the standardized gene expression

profile of all subjects in the dataset (50, 51, 53) such that sample composition had a

large effect on the PLAGE scores. In addition, the original models for the gene

signatures Sweeney_OD_3, Kaforou (Kaforou_27, Kafotrou_OD_44, and

Kaforou_OD_53), and Anderson (Anderson_42 and Anderson_OD_51) could also be

characterized as single sample methods, which rely on the expression of upregulated

and downregulated subset of genes within gene sets (Table 1). These single-sample

methods were more likely to produce robust scores for individual subjects, especially in

studies with small sample sizes or heterogeneous disease subtypes. Multi-sample

methods were susceptible to sample composition (i.e., the number of samples for

different subtypes), which could introduce biases if the size of samples for different

disease subtypes changed (53). PLAGE and other multi-sample biomarker models

(e.g., random forest, linear discriminant analysis, and K-nearest neighbors) compute a

different score for an individual sample whenever a patient enters or leaves the test

population. This is known as "test set bias" (62), as a gene signature's performance

may be irreproducible when a patient population composition or size changes.

Therefore, one of the limitations of using these multi-sample methods for sample

scoring is that bias will be induced if researchers perform integration analysis by

merging datasets from different sources. In our study, the results given by PLAGE for

most of the signatures outperformed the rest of gene set scoring methods (Table 3;

Figure 2A). This is primarily because the gene signature selection process is usually

heavily based on prior knowledge and the activity levels of the selected genes are

biologically meaningful, which could be revealed using SVD-like methods, including

principal component analysis (51).

The weighted mean AUCs given by single-sample scoring methods were sometimes

lower than the AUC from the original model for some signatures (Table 3). For these

cases, the biomarker splitting strategy improved diagnostic ability when using the

single-sample scoring methods, ssGSEA, GSVA, and Singscore, consistent with

existing publications in other fields (50, 63). Further, we noted that the improvement of

weighted mean AUC values based on upregulated subsets was more dominant when

compared to the results from the downregulated subset (Table 3). This is likely because

upregulated genes are usually immune-related, such as FCGR1A/B, GBP5/6, C1QB,

SEPTIN4, and ANDKRD22 (21), which generate a clear signal in active TB and are

features of the immune response to the disease (64). The weighted AUCs from

Zak_RISK_16 were consistently greater than 0.80 for five gene set scoring methods

(Table 2), mainly due to the overexpression of all genes within Zak_RISK_16 relative to

its discovery dataset, with large number of genes being highly differentially expressed

from the recent Leicester clinical phenotype groups (64). Therefore, the choice of gene

set scoring method depends on the cohort composition and the differential expressed

genes within the gene signatures. Generally, it is ideal and efficient to implement gene

set scoring methods to evaluate the diagnostic accuracy of TB signatures, which we

have shown to achieve equivalent accuracy when compared to their original models.

The data preprocessing and training procedures are specialized and intractable for most

TB gene signature discovery cases, which contributes to low generalizability in some

biomarkers when evaluating their performance on independent datasets by

implementing the original classification model. Both Berry_393 and Berry_OD_86 gene

signatures used the K-nearest neighbor (KNN) algorithm, which demonstrated high

classification ability in their discovery studies (Table 2) but had poor results across

multiple studies (Table 3). KNN clustering worked well when gene expression values

were normalized to the median of each control group (10); however, performing such

normalization for transcriptomic datasets originating from different clinical conditions or

different platforms is unrealistic. KNN classification also assumes that similar inputs

share similar labels; however, data points tend to be close together in high dimensional

spaces (65). Since the Berry_393 and Berry_OD_86 consist of 393 and 86 transcripts

respectively, KNN lost predictive power in transcriptomic studies containing few

samples but large numbers of gene expression features.

On the other hand, the performance of Verhagen_10 was poor across independent

studies based on the results given by its original diagnostic model but had AUC values

equal to 1.00 in datasets GSE81746 and GSE41055 (Figure 1). GSE41055 is the

discovery study for Verhagen_10, and GSE81746 is a small study that only contains

two active TB diseases and four health control samples. In our comparison, 17 of 19

signatures could identify the active TB subtype with an AUC greater than 0.80 (Figure

1). The mediocre performance of Verhagen_10 across multiple studies is a sign of

overfitting, a common problem using random forest, which relies on optimizing the

tuning parameters (66). Since the original publication did not offer information about the

hyper-parameters, we could not reconstruct the diagnostic model in the same way.

Similar observations were made for Leong_RISK_29 (32), for which AUC results varied

from 0.5 and 1.00 across independent datasets (Figure 1). As Leong_RISK_29 was

designed to capture early host response to infection, it has not been generalized to

samples with clinical manifestations of the disease (32). In these and many other ways,

a gene signature's diagnostic accuracy may be underestimated by evaluating its

performance using its original model.

We also examined the gene signatures' predictive accuracy within individual studies

based on the results given by gene set scoring methods and the signatures' original

model. Results from PLAGE and ssGSEA for Kaforou_27 were highly correlated (ρ_w

from PLAGE: = -0.92, ρ_w from ssGSEA: = 0.78), with a small AUC difference ($|\Delta AUC|_w$

from PLAGE: = 0.030, $|\Delta AUC|_w$ from ssGSEA: = 0.032), and it had a study overlap

coefficient of 1.00 when compared to its original model (Figure S4A-B). This is primarily

because of the presence of 21 over-expressed genes within the Kaforou_27 gene set,

which dominates the performance of the gene signatures (Figure S5A). We identified a

similar situation for Maertzdorf_15, which had 12 down-regulated genes within its gene

set. Maertzdorf_15 and its down-regulated subsets had consistently high negative

Spearman's rank correlations (-0.89 <= ρ_w <= -0.78) for the prediction scores given by

its original model and gene set scoring methods except for PLAGE (ρ_w = 0.94) and

Singscore bidirectional scoring (ρ_w = 0.86; Figure S4 A-E; Figure S5B, D-E). The

outperformance of Sweeney_OD_3 in ssGSEA and Singscore unidirectional scoring

was largely due to the down-regulated gene *KLF2* within the Sweeney_OD_3 gene set.

It has a relatively lower summary effect of the standardized mean difference between

active tuberculosis and other clinical conditions from the biomarker's original discovery

datasets of Sweeney_OD_3; in contrast, the remaining two upregulated genes were

highly differentially expressed in the corresponding datasets (26).

Our study has several limitations. We only compared the performance of active TB

disease versus all other disease states regardless of how TB was diagnosed, and we

did not perform subgroup analysis stratified by different clinical conditions, such as age,

region, gender, or comorbidities. Additionally, some reconstructed signature models

may not be the same as those used in the original publication, as some of these

publications do not provide precise details to recreate the original training model.

Instead, we followed the details outlined in the previously published comparison of

Warsinske *et al.* (19). Due to different naming sequencing platforms for transcripts,

some genes within signatures were missing across multiple studies, and we could not

evaluate the biomarkers' diagnostic ability in those cases accurately. Although we used

KNN imputation to estimate the expression values for those missing genes in the

validation study, the imputation process may have potentially led to overfitting. Our

comparison would still be valid if we followed similar procedures to handle missing

values across independent studies.

In conclusion, we showed that gene set scoring methods are appropriate for evaluating

gene signature accuracy for comparing active TB disease versus controls. These

methods perform similarly when compared to the signatures' original models. Overall,

the performance of TB biomarkers across multiple studies was similar between the

PLAGE and the original models. The methods ssGSEA, GSVA, and Singscore can also

capture the diagnostic accuracy of gene signatures by taking the gene directional

information within gene sets into account. Given the challenges associated with

rebuilding or re-evaluating the signatures' original biomarker model(s), gene set scoring

methods could serve as a reliable alternative computational methodology to apply or

perform comparisons of TB biomarkers.

REFERNCES

1.   World Health Organization. *Global Tuberculosis Report 2020*. World Health

Organization; 2020.

2.   Pai M, Behr MA, Dowdy D, Dheda K, Divangahi M, Boehme CC, Ginsberg A,

Swaminathan S, Spigelman M, Getahun H, Menzies D, Raviglione M. Tuberculosis.

Nat Rev Dis Primers 2016;2:16076.

3.   Park JH, Choe J, Bae M, Choi S, Jung KH, Kim MJ, Chong YP, Lee S-O, Choi S-H,

Kim YS, Woo JH, Jo K-W, Shim TS, Kim MY, Kim S-H. Clinical Characteristics and

Radiologic Features of Immunocompromised Patients With Pauci-Bacillary

Pulmonary Tuberculosis Receiving Delayed Diagnosis and Treatment. *Open Forum*

Infect Dis 2019;6:ofz002.

4.   Swaminathan S, Ramachandran G. Challenges in childhood tuberculosis. *Clin*

Pharmacol Ther 2015;98:240–244.

5.   Sharma SK, Ryan H, Khaparde S, Sachdeva KS, Singh AD, Mohan A, Sarin R,

Paramasivan CN, Kumar P, Nischal N, Khatiwada S, Garner P, Tharyan P. Index-

TB guidelines: Guidelines on extrapulmonary tuberculosis for India. *Indian J Med*

Res 2017;145:448–463.

6.   Gaur M, Singh A, Sharma V, Tandon G, Bothra A, Vasudeva A, Kedia S, Khanna

A, Khanna V, Lohiya S, Varma-Basil M, Chaudhry A, Misra R, Singh Y. Diagnostic

performance of non-invasive, stool-based molecular assays in patients with

paucibacillary tuberculosis. *Sci Rep* 2020;10:7102.

7.   Gupta RK, Turner CT, Venturini C, Esmail H, Rangaka MX, Copas A, Lipman M,

Abubakar I, Noursadeghi M. Concise whole blood transcriptional signatures for

incipient tuberculosis: a systematic review and patient-level pooled meta-analysis.

Lancet Respir Med 2020;8:395–406.

8. Sloot R, Schim van der Loeff MF, van Zwet EW, Haks MC, Keizer ST, Scholing M,

Ottenhoff THM, Borgdorff MW, Joosten SA. Biomarkers Can Identify Pulmonary

Tuberculosis in HIV-infected Drug Users Months Prior to Clinical Diagnosis.

EBioMedicine 2015;2:172–179.

9. Esmail H, Lai RP, Lesosky M, Wilkinson KA, Graham CM, Horswell S, Coussens

AK, Barry CE 3rd, O'Garra A, Wilkinson RJ. Complement pathway gene activation

and rising circulating immune complexes characterize early disease in HIV-

associated tuberculosis. *Proc Natl Acad Sci U S A* 2018;115:E964–E973.

10. Berry MPR, Graham CM, McNab FW, Xu Z, Bloch SAA, Oni T, Wilkinson KA,

Banchereau R, Skinner J, Wilkinson RJ, Quinn C, Blankenship D, Dhawan R, Cush

JJ, Mejias A, Ramilo O, Kon OM, Pascual V, Banchereau J, Chaussabel D,

O'Garra A. An interferon-inducible neutrophil-driven blood transcriptional signature

in human tuberculosis. *Nature* 2010;466:973–977.

11. Walter ND, Miller MA, Vasquez J, Weiner M, Chapman A, Engle M, Higgins M,

Quinones AM, Rosselli V, Canono E, Yoon C, Cattamanchi A, Davis JL, Phang T,

Stearman RS, Datta G, Garcia BJ, Daley CL, Strong M, Kechris K, Fingerlin TE,

Reves R, Geraci MW. Blood Transcriptional Biomarkers for Active Tuberculosis

among Patients in the United States: a Case-Control Study with Systematic Cross-

Classifier Evaluation. *J Clin Microbiol* 2016;54:274–282.

12. Kaforou M, Wright VJ, Oni T, French N, Anderson ST, Bangani N, Banwell CM,

Brent AJ, Crampin AC, Dockrell HM, Eley B, Heyderman RS, Hibberd ML, Kern F,

Langford PR, Ling L, Mendelson M, Ottenhoff TH, Zgambo F, Wilkinson RJ, Coin

LJ, Levin M. Detection of tuberculosis in HIV-infected and -uninfected African adults

using whole blood RNA expression signatures: a case-control study. *PLoS Med*

2013;10:e1001538.

13. Suliman S, Thompson EG, Sutherland J, Weiner J 3rd, Ota MOC, Shankar S,

Penn-Nicholson A, Thiel B, Erasmus M, Maertzdorf J, Duffy FJ, Hill PC, Hughes

EJ, Stanley K, Downing K, Fisher ML, Valvo J, Parida SK, van der Spuy G, Tromp

G, Adetifa IMO, Donkor S, Howe R, Mayanja-Kizza H, Boom WH, Dockrell HM,

Ottenhoff THM, Hatherill M, Aderem A, *et al.* Four-Gene Pan-African Blood

Signature Predicts Progression to Tuberculosis. *Am J Respir Crit Care Med*

2018;197:1198–1208.

14. Zak DE, Penn-Nicholson A, Scriba TJ, Thompson E, Suliman S, Amon LM,

Mahomed H, Erasmus M, Whatney W, Hussey GD, Abrahams D, Kafaar F,

Hawkridge T, Verver S, Hughes EJ, Ota M, Sutherland J, Howe R, Dockrell HM,

Boom WH, Thiel B, Ottenhoff THM, Mayanja-Kizza H, Crampin AC, Downing K,

Hatherill M, Valvo J, Shankar S, Parida SK, *et al.* A blood RNA signature for

tuberculosis disease risk: a prospective cohort study. *Lancet* 2016;387:2312–2322.

15. Thompson EG, Du Y, Malherbe ST, Shankar S, Braun J, Valvo J, Ronacher K,

Tromp G, Tabb DL, Alland D, Shenai S, Via LE, Warwick J, Aderem A, Scriba TJ,

Winter J, Walzl G, Zak DE, Catalysis TB–Biomarker Consortium. Host blood RNA

signatures predict the outcome of tuberculosis treatment. *Tuberculosis*

2017;107:48–58.

16. Singhania A, Verma R, Graham CM, Lee J, Tran T, Richardson M, Lecine P,

Leissner P, Berry MPR, Wilkinson RJ, Kaiser K, Rodrigue M, Woltmann G, Haldar

P, O'Garra A. A modular transcriptional signature identifies phenotypic

heterogeneity of human tuberculosis infection. *Nat Commun* 2018;9:2308.

17. Roe J, Venturini C, Gupta RK, Gurry C, Chain BM, Sun Y, Southern J, Jackson C,

Lipman MC, Miller RF, Martineau AR, Abubakar I, Noursadeghi M. Blood

Transcriptomic Stratification of Short-term Risk in Contacts of Tuberculosis. *Clin*

Infect Dis 2020;70:731–737.

18. Scriba TJ, Fiore-Gartland A, Penn-Nicholson A, Mulenga H, Kimbung Mbandi S,

Borate B, Mendelsohn SC, Hadley K, Hikuam C, Kaskar M, Musvosvi M, Bilek N,

Self S, Sumner T, White RG, Erasmus M, Jaxa L, Raphela R, Innes C, Brumskine

W, Hiemstra A, Malherbe ST, Hassan-Moosa R, Tameris M, Walzl G, Naidoo K,

Churchyard G, Hatherill M, CORTIS-01 Study Team. Biomarker-guided

tuberculosis preventive therapy (CORTIS): a randomised controlled trial. *Lancet*

Infect Dis 2021;21:354–365.

19. Warsinske H, Vashisht R, Khatri P. Host-response-based gene signatures for

tuberculosis diagnosis: A systematic comparison of 16 signatures. *PLoS Med*

2019;16:e1002786.

20. Noble WS. What is a support vector machine? *Nat Biotechnol* 2006;24:1565–1567.

21. Johnson WE, Odom A, Cintron C, Muthaiah M, Knudsen S, Joseph N, Babu S,

Lakshminarayanan S, Jenkins DF, Zhao Y, Nankya E, Horsburgh CR, Roy G,

Ellner JJ, Sarkar S, Salgame P, Hochberg NS. Comparing Tuberculosis Gene

Signatures in Malnourished Individuals using the TBSignatureProfiler. *BMC Infect*

Dis 2020;

22. Laux da Costa L, Delcroix M, Dalla Costa ER, Prestes IV, Milano M, Francis SS,

Unis G, Silva DR, Riley LW, Rossetti MLR. A real-time PCR signature to

discriminate between tuberculosis and other pulmonary diseases. *Tuberculosis*

2015;95:421–425.

23. Jacobsen M, Repsilber D, Gutschmidt A, Neher A, Feldmann K, Mollenkopf HJ,

Ziegler A, Kaufmann SHE. Candidate biomarkers for discrimination between

infection and disease caused by Mycobacterium tuberculosis. *J Mol Med*

2007;85:613–621.

24. Sambarey A, Devaprasad A, Mohan A, Ahmed A, Nayak S, Swaminathan S,

D'Souza G, Jesuraj A, Dhar C, Babu S, Vyakarnam A, Chandra N. Unbiased

Identification of Blood-based Biomarkers for Pulmonary Tuberculosis by Modeling

and Mining Molecular Interaction Networks. *EBioMedicine* 2017;15:112–126.

25. Hänzelmann S, Castelo R, Guinney J. GSVA: gene set variation analysis for

microarray and RNA-seq data. *BMC Bioinformatics* 2013;14:7.

26. Sweeney TE, Braviak L, Tato CM, Khatri P. Genome-wide expression for diagnosis

of pulmonary tuberculosis: a multicohort analysis. *Lancet Respir Med* 2016;4:213–

224.

27. Anderson ST, Kaforou M, Brent AJ, Wright VJ, Banwell CM, Chagaluka G, Crampin

AC, Dockrell HM, French N, Hamilton MS, Hibberd ML, Kern F, Langford PR, Ling

L, Mlotha R, Ottenhoff THM, Pienaar S, Pillay V, Scott JAG, Twahir H, Wilkinson

RJ, Coin LJ, Heyderman RS, Levin M, Eley B. Diagnosis of childhood tuberculosis

and host RNA expression in Africa. *N Engl J Med* 2014;370:1712–1723.

28. Bloom CI, Graham CM, Berry MPR, Rozakeas F, Redford PS, Wang Y, Xu Z,

Wilkinson KA, Wilkinson RJ, Kendrick Y, Devouassoux G, Ferry T, Miyara M,

Bouvry D, Valeyre D, Gorochov G, Blankenship D, Saadatian M, Vanhems P,

Beynon H, Vancheeswaran R, Wickremasinghe M, Chaussabel D, Banchereau J,

Pascual V, Ho L-P, Lipman M, O'Garra A. Transcriptional blood signatures

distinguish pulmonary tuberculosis, pulmonary sarcoidosis, pneumonias and lung

cancers. *PLoS One* 2013;8:e70630.

29. Leong S, Zhao Y, Joseph NM, Hochberg NS, Sarkar S, Pleskunas J, Hom D,

Lakshminarayanan S, Horsburgh CR Jr, Roy G, Ellner JJ, Johnson WE, Salgame

P. Existing blood transcriptional classifiers accurately discriminate active

tuberculosis from latent infection in individuals from south India. *Tuberculosis*

2018;109:41–51.

30. Maertzdorf J, McEwen G, Weiner J 3rd, Tian S, Lader E, Schriek U, Mayanja-Kizza

H, Ota M, Kenneth J, Kaufmann SH. Concise gene signature for point-of-care

classification of tuberculosis. *EMBO Mol Med* 2016;8:86–95.

31. Verhagen LM, Zomer A, Maes M, Villalba JA, Del Nogal B, Eleveld M, van Hijum

SA, de Waard JH, Hermans PW. A predictive signature gene set for discriminating

active from latent tuberculosis in Warao Amerindian children. *BMC Genomics*

2013;14:74.

32. Leong S, Zhao Y, Ribeiro-Rodrigues R, Jones-López EC, Acuña-Villaorduña C,

Rodrigues PM, Palaci M, Alland D, Dietze R, Ellner JJ, Johnson WE. Cross-

validation of existing signatures and derivation of a novel 29-gene transcriptomic

signature predictive of progression to TB in a Brazilian cohort of household

contacts of pulmonary TB. *Tuberculosis* 2020;120:101898.

33. Banchereau R, Jordan-Villegas A, Ardura M, Mejias A, Baldwin N, Xu H, Saye E,

Rossello-Urgell J, Nguyen P, Blankenship D, Creech CB, Pascual V, Banchereau

J, Chaussabel D, Ramilo O. Host immune transcriptional profiles reflect the

variability in clinical disease manifestations in patients with Staphylococcus aureus

infections. *PLoS One* 2012;7:e34390.

34. Park J, Munagala I, Xu H, Blankenship D, Maffucci P, Chaussabel D, Banchereau

J, Pascual V, Cunningham-Rundles C. Interferon signature in the blood in

inflammatory common variable immune deficiency. *PLoS One* 2013;8:e74893.

35. Obermoser G, Presnell S, Domico K, Xu H, Wang Y, Anguiano E, Thompson-

Snipes L, Ranganathan R, Zeitner B, Bjork A, Anderson D, Speake C, Ruchaud E,

Skinner J, Alsina L, Sharma M, Dutartre H, Cepika A, Israelsson E, Nguyen P,

Nguyen Q-A, Harrod AC, Zurawski SM, Pascual V, Ueno H, Nepom GT, Quinn C,

Blankenship D, Palucka K, *et al.* Systems scale interactive exploration reveals

quantitative and qualitative differences in response to influenza and pneumococcal

vaccines. *Immunity* 2013;38:831–844.

36. VanValkenburg A, Kaipilyawar V, Sarkar S, Lakshminarayanan S, Cintron C,

Prakash Babu S, Knudsen S, Joseph NM, Horsburgh CR, Ellner JJ, Narasimhan

PB, Johnson WE, Hochberg NS, Salgame P. Malnutrition leads to increased

inflammation and expression of tuberculosis risk signatures in recently exposed

household contacts of pulmonary tuberculosis. *Front Immunol* 2022;13:1011166.

37. Maertzdorf J, Ota M, Repsilber D, Mollenkopf HJ, Weiner J, Hill PC, Kaufmann

SHE. Functional correlations of pathogenesis-driven gene expression signatures in

tuberculosis. *PLoS One* 2011;6:e26938.

38. Elliott TOJP, Owolabi O, Donkor S, Kampmann B, Hill PC, Ottenhoff THM, Haks

MC, Kaufmann SHE, Maertzdorf J, Sutherland JS. Dysregulation of apoptosis is a

risk factor for tuberculosis disease progression. *J Infect Dis* 2015;212:1469–1479.

39. Blankley S, Graham CM, Turner J, Berry MPR, Bloom CI, Xu Z, Pascual V,

Banchereau J, Chaussabel D, Breen R, Santis G, Blankenship DM, Lipman M,

O'Garra A. The transcriptional signature of active tuberculosis reflects symptom

status in extra-pulmonary and pulmonary tuberculosis. *PLoS One*

2016;11:e0162220.

40. Cai Y, Yang Q, Tang Y, Zhang M, Liu H, Zhang G, Deng Q, Huang J, Gao Z, Zhou

B, Feng CG, Chen X. Increased complement C1q level marks active disease in

human tuberculosis. *PLoS One* 2014;9:e92340.

41. Maertzdorf J, Weiner J 3rd, Mollenkopf H-J, TBornotTB Network, Bauer T, Prasse

A, Müller-Quernheim J, Kaufmann SHE. Common patterns and disease-related

signatures in tuberculosis and sarcoidosis. *Proc Natl Acad Sci U S A*

2012;109:7853–7858.

42. Ottenhoff THM, Dass RH, Yang N, Zhang MM, Wong HEE, Sahiratmadja E, Khor

CC, Alisjahbana B, van Crevel R, Marzuki S, Seielstad M, van de Vosse E, Hibberd

ML. Genome-wide expression profiling identifies type 1 interferon response

pathways in active tuberculosis. *PLoS One* 2012;7:e45839.

43. Dawany N, Showe LC, Kossenkov AV, Chang C, Ive P, Conradie F, Stevens W,

Sanne I, Azzoni L, Montaner LJ. Identification of a 251 gene expression signature

that can accurately detect M. tuberculosis in patients with and without HIV co-

infection. *PLoS One* 2014;9:e89925.

44. Marais S, Lai RPJ, Wilkinson KA, Meintjes G, O'Garra A, Wilkinson RJ.

Inflammasome activation underlying central nervous system deterioration in HIV-

associated tuberculosis. *J Infect Dis* 2017;215:677–686.

45.　de Araujo LS, Vaas LAI, Ribeiro-Alves M, Geffers R, Mello FCQ, de Almeida AS,

Moreira A da SR, Kritski AL, Lapa E Silva JR, Moraes MO, Pessler F, Saad MHF.

Transcriptomic biomarkers for tuberculosis: Evaluation of DOCK9. EPHA4, and

NPC2 mRNA expression in peripheral blood. *Front Microbiol* 2016;7:1586.

46.　Zak DE, Penn-Nicholson A, Scriba TJ, Thompson E, Suliman S, Amon LM,

Mahomed H, Erasmus M, Whatney W, Hussey GD, Abrahams D, Kafaar F,

Hawkridge T, Verver S, Hughes EJ, Ota M, Sutherland J, Howe R, Dockrell HM,

Boom WH, Thiel B, Ottenhoff THM, Mayanja-Kizza H, Crampin AC, Downing K,

Hatherill M, Valvo J, Shankar S, Parida SK, *et al.* A blood RNA signature for

tuberculosis disease risk: a prospective cohort study. *Lancet* 2016;387:2312–2322.

47.　Lee S-W, Wu LS-H, Huang G-M, Huang K-Y, Lee T-Y, Weng JT-Y. Gene

expression profiling identifies candidate biomarkers for active and latent

tuberculosis. *BMC Bioinformatics* 2016;17 Suppl 1:3.

48.　Tientcheu LD, Maertzdorf J, Weiner J, Adetifa IM, Mollenkopf H-J, Sutherland JS,

Donkor S, Kampmann B, Kaufmann SHE, Dockrell HM, Ota MO. Differential

transcriptomic and metabolic profiles of M. africanum- and M. tuberculosis-infected

patients after, but not before, drug treatment. *Genes Immun* 2015;16:347–355.

49.　Pavlov YL. *Random Forests.* Walter de Gruyter GmbH & Co KG; 2019.

50.　Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA,

Paulovich A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP. Gene set enrichment

analysis: a knowledge-based approach for interpreting genome-wide expression

profiles. *Proc Natl Acad Sci U S A* 2005;102:15545–15550.

51. Tomfohr J, Lu J, Kepler TB. Pathway level analysis of gene expression using

singular value decomposition. *BMC Bioinformatics* 2005;6:225.

52. Lee E, Chuang H-Y, Kim J-W, Ideker T, Lee D. Inferring pathway activity toward

precise disease classification. *PLoS Comput Biol* 2008;4:e1000217.

53. Foroutan M, Bhuva DD, Lyu R, Horan K, Cursons J, Davis MJ. Single sample

scoring of molecular phenotypes. *BMC Bioinformatics* 2018;19:404.

54. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol* 2014;15:550.

55. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. limma powers

differential expression analyses for RNA-sequencing and microarray studies.

Nucleic Acids Res 2015;43:e47–e47.

56. Oh S, Abdelnabi J, Al-Dulaimi R, Aggarwal A, Ramos M, Davis S, Riester M,

Waldron L. HGNChelper: identification and correction of invalid gene symbols for

human and mouse. *F1000Res* 2020;9:1493.

57. Hastie T, Tibshirani R, Narasimhan B, Chu G. *impute: Imputation for microarray*

data. 2020. at <10.18129/B9.bioc.impute>.

58. Zhang Y, Jenkins DF, Manimaran S, Johnson WE. Alternative empirical Bayes

models for adjusting for batch effects in genomic studies. *BMC Bioinformatics*

2018;19:262.

59. Johnson WE, Li C, Rabinovic A. Adjusting batch effects in microarray expression

data using empirical Bayes methods. *Biostatistics* 2007;8:118–127.

60. Khan MRA, Brandenburger T. *ROCit: Performance Assessment of Binary Classifier*

with Visualization. 2020. at <https://CRAN.R-project.org/package=ROCit>.

61. M.k V, K K. A survey on similarity measures in text mining. *Mach Learn Appl Int J*

2016;3:19–28.

62. Patil P, Bachant-Winner P-O, Haibe-Kains B, Leek JT. Test set bias affects

reproducibility of gene signatures. *Bioinformatics* 2015;31:2318–2323.

63. Barbie DA, Tamayo P, Boehm JS, Kim SY, Moody SE, Dunn IF, Schinzel AC,

Sandy P, Meylan E, Scholl C, Fröhling S, Chan EM, Sos ML, Michel K, Mermel C,

Silver SJ, Weir BA, Reiling JH, Sheng Q, Gupta PB, Wadlow RC, Le H, Hoersch S,

Wittner BS, Ramaswamy S, Livingston DM, Sabatini DM, Meyerson M, Thomas

RK, *et al.* Systematic RNA interference reveals that oncogenic KRAS-driven

cancers require TBK1. *Nature* 2009;462:108–112.

64. Tabone O, Verma R, Singhania A, Chakravarty P, Branchett WJ, Graham CM, Lee

J, Trang T, Reynier F, Leissner P, Kaiser K, Rodrigue M, Woltmann G, Haldar P,

O'Garra A. Blood transcriptomics reveal the evolution and resolution of the immune

response in tuberculosis. *J Exp Med* 2021;218:.

65. Tran TN, Wehrens R, Buydens LMC. KNN-kernel density-based clustering for high-

dimensional multivariate data. *Computational Statistics & Data Analysis* 2006;

66. Lulli A, Oneto L, Anguita D. Mining big data with random forests. *Cognit Comput*

2019;11:294–316.

Table 1. Summary of TB gene signatures compared in this study and their original training model.

Signature Name	Gene Signature Number	Comparison	Datasets	Original Model Description
Sweeney_OD_3	3	Active tuberculosis vs. (LTBI & HCs & OD)	GSE19491 & GSE42834 & GSE37250	Difference of geometric means between up and down-regulated genes
Jacobsen_3	3	Active tuberculosis vs. LTBI	GSE19491	Linear Discriminant Analysis
LauxdaCosta_OD_3	3	Active tuberculosis vs. OD	GSE42834	Random Forest
Maertzdorf_4	4	Active tuberculosis vs. HCs	GSE74092	Random Forest
Sambarey_HIV_10	10	Active tuberculosis vs. OD	GSE37250	Linear Discriminant Analysis
Verhagen_10	10	Active tuberculosis vs. (LTBI & HCs)	GSE41055	Random Forest
Maertzdorf_15	15	Active tuberculosis vs. HCs	GSE74092	Random Forest
Leong_24	24	Active tuberculosis vs. LTBI	GSE10175	Ridge Logistic Regression
Kaforou_27	27	Active tuberculosis vs. OD	GSE19491	Difference of arithmetic means between up and down-regulated genes
Anderson_42	42	Active tuberculosis vs. LTBI	GSE39940	Difference of sums between up and down-regulated genes
Kaforou_OD_44	44	Active tuberculosis vs. OD	GSE19491	Difference of arithmetic means between up and down-regulated genes
Anderson_OD_51	51	Active tuberculosis vs. OD	GSE39940	Difference of sums between up and down-regulated genes
Kaforou_OD_53	53	Active tuberculosis vs. OD	GSE19491	Difference of arithmetic means between up and down-regulated genes
Berry_OD_86	86	Active tuberculosis vs. OD	GSE19491	K-nearest neighbors algorithm
Bloom_OD_144	144	Active tuberculosis vs. (HCs & OD)	GSE42834	Support Vector Machines
Berry_393	393	Active tuberculosis vs. (LTBI & HCs)	GSE19491	K-nearest neighbors algorithm
Suliman_RISK_4	4	Incipient tuberculosis vs. HCs	GSE94438	Support Vector Machines (linear kernel, using paired ratio)
Zak_RISK_16	16	Incipient tuberculosis vs. HCs	GSE79362	Support Vector Machines (linear kernel)
Leong_RISK_29	29	Incipient tuberculosis vs. HCs	GSE79362	Lasso Logistic Regression

Figure 1. Heatmap of AUC distribution for each signature across 24 studies using original model. Grids with black borders were the discovery sets for each TB gene signature. Each row represents one signature. Signatures were clustered into different categories with respect to the TB subtypes that they aim to identify. The column of the heatmap representing the studies that were used in this paper. The datasets were rearranged in decreasing order based on their mean AUC values across all TB gene signatures.

Table 2. Weighted mean AUC and 95% CI for 19 gene signatures using original model and gene set scoring methods (ssGSEA, GSVA, PLAGE, Zscore, and Singscore) across 24 studies. Color-coded for AUC values that were 0.05 unit greater (red) than in Warsinske et al and were 0.05 less (blue) than Warsinske et al. (*: p <= 0.01, **: p <= 0.001, ***: p <= 0.0001 derived from Wilcoxon signed-rank test between the original model and corresponding gene set scoring methods; †: the original model results underperformed the Warsinske et al. with more than 0.05 unit; ‡: the original model results outperformed the Warsiske et al. with more than 0.05 unit).

Signature	Warsinske et al.	TBSignatureProfiler (Original Model)	TBSignatureProfiler (ssGSEA)	TBSignatureProfiler (GSVA)	TBSignatureProfiler (PLAGE)	TBSignatureProfiler (Zscore)	TBSignatureProfiler (Singscore)
LauxdaCosta_OD_3	0.76 (0.45–1.00)	0.80 (0.76-0.83)	0.82 (0.76-0.86)	0.79 (0.74-0.83)	0.83* (0.79-0.86)	0.78 (0.74-0.80)	0.82 (0.75-0.86)
Jacobsen_3	0.83 (0.69–0.98)	0.76† (0.71-0.80)	0.72 (0.68-0.76)	0.69 (0.64-0.73)	0.78 (0.75-0.81)	0.69 (0.65-0.73)	0.71 (0.67-0.75)
Sweeney_OD_3	0.85 (0.72–0.99)	0.81 (0.76-0.85)	0.82 (0.77-0.86)	0.75 (0.69-0.80)	0.82 (0.78-0.85)	0.77 (0.71-0.82)	0.82 (0.77-0.86)
Maertzdorf_4	0.79 (0.64–0.95)	0.80 (0.77-0.83)	0.70* (0.65-0.75)	0.73** (0.69-0.78)	0.81 (0.78-0.85)	0.72** (0.68-0.76)	0.66* (0.62-0.72)
Verhagen_10	0.54 (0.41–0.68)	0.61‡ (0.57-0.66)	0.57 (0.55-0.60)	0.59 (0.56-0.64)	0.65 (0.59-0.71)	0.62 (0.57-0.68)	0.58 (0.55-0.61)
Sambarey_HIV_10	0.82 (0.57–1.00)	0.83 (0.76-0.87)	0.80 (0.76-0.84)	0.80 (0.76-0.84)	0.76 (0.70-0.83)	0.75 (0.70-0.81)	0.79 (0.75-0.83)
Maertzdorf_15	0.79 (0.66–0.92)	0.82 (0.79-0.85)	0.83 (0.79-0.87)	0.82 (0.78-0.86)	0.83 (0.80-0.86)	0.79 (0.75-0.82)	0.79 (0.75-0.84)
Leong_24	0.75 (0.54–0.95)	0.72 (0.67-0.78)	0.73 (0.70-0.77)	0.61 (0.58-0.65)	0.73 (0.67-0.79)	0.61 (0.58-0.65)	0.73 (0.69-0.78)
Kaforou_27	0.83 (0.64–1.00)	0.81 (0.77-0.85)	0.82 (0.78-0.85)	0.79 (0.76-0.82)	0.83 (0.79-0.87)	0.79 (0.76-0.82)	0.78 (0.73-0.82)
Anderson_42	0.82 (0.66–0.97)	0.78 (0.73-0.83)	0.61** (0.58-0.66)	0.60** (0.58-0.64)	0.76 (0.70-0.82)	0.65** (0.59-0.73)	0.57** (0.53-0.61)
Kaforou_OD_44	0.78 (0.56–1.00)	0.76 (0.69-0.81)	0.67 (0.63-0.71)	0.72 (0.68-0.74)	0.80 (0.72-0.86)	0.70 (0.67-0.74)	0.70 (0.67-0.74)
Anderson_OD_51	0.58 (0.33–0.82)	0.71‡ (0.64-0.78)	0.75 (0.71-0.80)	0.81 (0.77-0.84)	0.79 (0.72-0.85)	0.75 (0.66-0.82)	0.77 (0.73-0.80)
Kaforou_OD_53	0.84 (0.70–0.99)	0.83 (0.78-0.87)	0.70** (0.66-0.75)	0.77** (0.74-0.80)	0.84 (0.80-0.87)	0.77** (0.73-0.80)	0.77** (0.73-0.81)
Berry_OD_86	0.69 (0.36–1.00)	0.69 (0.66-0.72)	0.71 (0.68-0.76)	0.74 (0.70-0.78)	0.75* (0.72-0.80)	0.73 (0.69-0.78)	0.73 (0.68-0.78)
Bloom_OD_144	0.74 (0.52–0.96)	0.70 (0.66-0.74)	0.77 (0.72-0.81)	0.76 (0.71-0.81)	0.71 (0.66-0.78)	0.70 (0.63-0.77)	0.76 (0.71-0.81)
Berry_393	0.71 (0.43–0.99)	0.70 (0.66-0.74)	0.78 (0.74-0.82)	0.77 (0.73-0.81)	0.79* (0.75-0.84)	0.77 (0.73-0.81)	0.79 (0.74-0.84)
Suliman_RISK_4	NA	0.62 (0.57-0.69)	0.62 (0.58-0.68)	0.55 (0.53-0.58)	0.74** (0.70-0.79)	0.60 (0.55-0.66)	0.61 (0.56-0.66)
Zak_RISK_16	NA	0.62 (0.56-0.70)	0.85*** (0.81-0.88)	0.84*** (0.80-0.88)	0.83*** (0.80-0.86)	0.83*** (0.79-0.86)	0.84*** (0.80-0.88)
Leong_RISK_29	NA	0.68 (0.65-0.73)	0.58* (0.56-0.61)	0.59* (0.56-0.61)	0.75 (0.70-0.80)	0.60 (0.56-0.64)	0.61* (0.57-0.65)

Table 3. Weighted mean AUC and 95% CI results for upregulated and downregulated subsets of gene signature using the ssGSEA, GSVA, and Singscore (bidirectional version). Color-coded for AUC values that were 0.05 greater (red) than the original model and were 0.05 less (blue) than the original model . (*: p <= 0.01, **: p <= 0.001, ***: p <= 0.0001 derived from Wilcoxon signed-rank test).

Signature	TBSignatureProfiler (Original Model)	TBSignatureProfiler (ssGSEA: upregulated)	TBSignatureProfiler (ssGSEA: downregulated)	TBSignatureProfiler (GSVA: upregulated)	TBSignatureProfiler (GSVA: downregulated)	TBSignatureProfiler (Singscore)
Verhagen_10	0.61 (0.57-0.66)	0.55 (0.54-0.58)	0.57 (0.55-0.62)	0.56 (0.53-0.60)	0.62 (0.58-0.67)	0.56 (0.54-0.59)
Sambarey_HIV_10	0.83 (0.76-0.87)	0.82 (0.78-0.85)	0.68 (0.64-0.73)	0.82 (0.78-0.85)	0.71 (0.65-0.76)	0.67^* (0.62-0.74)
Maertzdorf_15	0.82 (0.79-0.85)	0.71^* (0.67-0.76)	0.85 (0.81-0.88)	0.71^{**} (0.68-0.75)	0.84 (0.80-0.88)	0.80 (0.77-0.84)
Leong_24	0.72 (0.67-0.78)	0.80 (0.76-0.85)	0.73 (0.69-0.77)	0.76 (0.71-0.80)	0.73 (0.69-0.77)	0.80 (0.76-0.84)
Kaforou_27	0.81 (0.77-0.85)	0.84 (0.81-0.88)	0.72^{***} (0.68-0.76)	0.83 (0.80-0.87)	0.71^{***} (0.67-0.75)	0.82 (0.78-0.86)
Anderson_42	0.78 (0.73-0.83)	0.82 (0.78-0.86)	0.73^* (0.69-0.78)	0.81 (0.78-0.84)	0.72^* (0.68-0.76)	0.81 (0.77-0.85)
Kaforou_OD_44	0.76 (0.69-0.81)	0.81 (0.75-0.84)	0.65 (0.59-0.71)	0.80(0.74-0.84)	0.64 (0.59-0.70)	0.74 (0.67-0.80)
Anderson_OD_51	0.71 (0.64-0.78)	0.75 (0.70-0.80)	0.73 (0.70-0.76)	0.81 (0.78-0.84)	0.77 (0.71-0.82)	0.65^* (0.62-0.70)
Kaforou_OD_53	0.83 (0.78-0.87)	0.85 (0.81-0.88)	0.75^{***} (0.70-0.80)	0.84 (0.81-0.87)	0.76^{***} (0.70-0.80)	0.82 (0.78-0.86)
Berry_OD_86	0.69 (0.66-0.72)	0.78^{***} (0.75-0.82)	0.63 (0.60-0.68)	0.80^{***} (0.76-0.84)	0.70 (0.66-0.75)	0.78^{**} (0.74-0.82)
Bloom_OD_144	0.70 (0.66-0.74)	0.78 (0.74-0.83)	0.64 (0.60-0.68)	0.78 (0.73-0.83)	0.64^{***} (0.61-0.68)	0.77 (0.73-0.81)
Berry_393	0.70 (0.66-0.74)	0.80^* (0.76-0.85)	0.74 (0.70-0.79)	0.79^* (0.76-0.84)	0.74 (0.70-0.79)	0.79^{**} (0.75-0.84)
Leong_RISK_29	0.68 (0.65-0.73)	0.72 (0.69-0.77)	0.69 (0.64-0.75)	0.76 (0.72-0.80)	0.71 (0.66-0.75)	0.73^{**} (0.70-0.77)

Figure 2. Differences of the weighted mean AUC values between the original model versus gene set scoring methods. Each grid showed the difference of weighted mean AUC value between the use of the corresponding gene set scoring methods (ssGSEA, GSVA, PLAGE, and Singscore) and the use of original models. The results from the original model was used as the baseline. For all 19 TB gene signatures, the weighted mean AUC results computed by ssGSEA, Singscore, GSVA, and Zscore were statistically equivalent from the results given by the original model **(A)**. The weighted mean AUC results computed by ssGSEA and GSVA for the upregulated subsets of the gene signatures, Singscore (bidirectional scoring), and GSVA for the downregulated subsets of the gene signatures were statistically equivalent from the results given by the original model **(B)**. Red: the weighted mean AUC for gene scoring method outperformed the original model. Blue: the weighted mean AUC for gene scoring method underperformed the original model.
*: p-value < 0.05 derived from the Wilcoxon signed-rank test
**: p-value < 0.01 derived from Wilcoxon signed-rank test

Supplementary Table 1. Summary of AUC and 95% CI results from gene signature's discovery/training study. The original publication section presented the AUC and 95% CI for each signature from their publication. The Warsinske et al. subsection showed the results that have been re-evaluated using the original models functionality that had been implemented in the TBSignatureProfiler, and the ssGSEA section presented the performance of each gene signature using ssGSEA from TBSignatureProfiler (Courtesy to Warsinske et al. work; ATB: active tuberculosis).

Signature	Clinical Comparison	Dataset Compared	Original Publication AUC (95% CI)	Warsinske et al. AUC (95% CI)	TBSignatureProfiler AUC (95% CI) Warsinske et al.	TBSignatureProfiler AUC (95%CI) ssGSEA
Sweeney_OD_3	ATB vs. HCs	GSE19491✧	0.96 (0.94 - 0.98)	0.96 (0.94 - 0.98)	0.95 (0.91 - 0.99)	0.96 (0.91 - 0.99)
		GSE42834✧	1.00 (1.00-1.00)	1.00 (1.00-1.00)	1.00 (1.00 - 1.00)	0.92 (0.83 - 0.98)
	ATB vs. LTBI	GSE19491✧	0.93 (0.91 - 0.95)	0.93 (0.91-0.95)	0.92 (0.87 - 0.97)	0.93 (0.87 - 0.97)
		GSE37250	0.93 (0.91 - 0.94)	0.93 (0.91-0.94)	0.93 (0.90 - 0.95)	0.90 (0.86 - 0.93)
	ATB vs. OD	GSE19491✧	0.92 (0.89 - 0.94)	0.92 (0.89-0.94)	0.90 (0.84 - 0.95)	0.92 (0.88 - 0.97)
		GSE37250	0.87 (0.85 - 0.89)	0.87 (0.85-0.89)	0.87 (0.83 - 0.91)	0.83 (0.78 - 0.87)
		GSE42834✧	0.84 (0.80 - 0.88)	0.84 (0.80-0.88)	0.84 (0.79 - 0.90)	0.78 (0.67 - 0.85)
Jacobsen_3	ATB vs. LTBI	GSE19491✧	N/A	0.93 (0.91-0.95)	0.93 (0.88 - 0.97)	0.79 (0.71 - 0.86)
LauxdaCosta_OD_3	ATB vs. OD	GSE42834✧	0.95* (0.88-1.00*)	1.00 (1.00-1.00)	1.00 (1.00-1.00)	0.82 (0.76 - 0.88)
Maertzdorf_4	ATB vs. HCs	GSE74092	0.98 (0.96 - 1.00)	0.99 (0.99-1.00)	1.00 (1.00-1.00)	0.71 (0.64 - 0.79)
Sambarey_HIV_10	ATB vs. (LTBI & HCs & OD)	GSE37250*	N/A	0.89 (0.87-0.91)	0.89 (0.85 - 0.91)	0.79 (0.75 - 0.82)
Verhagen_10	ATB vs. (LTBI & HCs)	GSE41055	N/A	1.00 (1.00-1.00)	1.00 (1.00-1.00)	0.85 (0.66 - 0.97)
Maertzdorf_15	ATB vs. HC	GSE74092	0.99 (0.97 - 1.00)	1.00 (1.00-1.00)	1.00 (1.00 - 1.00)	0.95 (0.92 - 0.97)
Leong_24	ATB vs. LTBI	GSE101705	0.98 (0.98 - 0.98)	1.00 (1.00-1.00)	1.00 (1.00 - 1.00)	0.95 (0.87 - 1.00)
Kaforou_27	ATB vs. LTBI	GSE19491✧	0.98 (0.95 - 1.00)	0.95 (0.93-0.97)	0.96 (0.92 - 0.99)	0.95 (0.90 - 0.98)
Anderson_42	ATB vs. LTBI	GSE39940	0.98 (0.95 - 1.00)	0.97 (0.96-0.98)	0.98 (0.95 - 0.99)	0.93 (0.88 - 0.96)
Kaforou_OD_44	ATB vs. OD	GSE19491✧	0.95 (0.89 - 0.99)	0.91 (0.89-0.94)	0.93 (0.90 - 0.96)	0.69 (0.60 - 0.77)
Anderson_OD_51	ATB vs. OD	GSE39940	0.86 (0.77 - 0.94)	0.89 (0.87-0.91)	0.90 (0.87 - 0.94)	0.80 (0.75 - 0.85)
Kaforou_OD_53	ATB vs. (LTBI & HCs & OD)	GSE19491✧	N/A	0.92 (0.89-0.94)	0.92 (0.88 - 0.96)	0.71 (0.63 - 0.77)

Supplementary Table 1 (Continued)

Signature	Clinical Comparison	Dataset Compared	Original Publication	Warsinske et al.	TBSignatureProfiler	
			AUC (95% CI)	AUC (95% CI)	AUC (95% CI) Warsinske et al.	AUC (95%CI) ssGSEA
Berry_OD_86	ATB vs. LTBI	GSE19491✧	N/A	0.97 (0.96-0.99)	0.96 (0.92 - 0.98)	0.90 (0.85 - 0.95)
	ATB vs. HCs	GSE19491✧	N/A	1.00 (1.00-1.00)	0.99 (0.98 - 1.00)	0.89 (0.84 - 0.94)
Bloom_OD_144	ATB vs. (OD & HCs)	GSE42834✧	0.91 (Not available)	0.99 (0.98-1.00)	0.97 (0.94 - 1.00)	0.80 (0.71 - 0.87)
Berry_393	ATB vs. LTBI	GSE19491✧	N/A	0.97 (0.96-0.99)	0.96 (0.93 - 0.99)	0.94 (0.89 - 0.97)
	ATB vs. HCs	GSE19491✧	N/A	0.99 (0.98-1.00)	0.99 (0.97 - 1.00)	0.95 (0.91 - 0.98)
Leong_RISK_29	Progressor vs. Non-progressor	GSE79362 (Baseline)	N/A	N/A	0.96 (0.93 - 0.99)	0.56 (0.51 - 0.69)
Zak_RISK_16	Progressor vs. Non-progressor	GSE79362 (Total Time Period)	0.74 (0.73 - 0.76)	N/A	0.72 (0.61 - 0.82)	0.77 (0.62 - 0.87)
Suliman_RISK_4 (Site-Specific)	Progressor vs. Non-progressor	GSE94438	0.67 (0.57-0.77)	N/A	0.71 (0.65 - 0.77)	0.61 (0.55 - 0.67)

* Original training studies were not available for the signatures; ✧ Superseries of multiple datasets

Supplementary Figure 1. The distribution of AUC values for 19 TB gene signatures across 24 studies using ssGSEA **(A)**, GSVA **(B)**, PLAGE **(C)**, Zscore **(D)**, and Singscore unidirectional version **(E)**. Datasets had the same order as those from Figure 1.

Supplementary Figure 2. The distribution of AUC values for **upregulated and downregulated subsets** of 13 TB gene signatures across 24 studies using ssGSEA **(A)**, GSVA**(B)**, and Singscore bidirectional version **(C)**. Datasets had the same order as those from Figure 1.

Supplementary Figure 3. The distribution of AUC values for TB gene signatures across 24 studies using the original models and gene set scoring methods in the form of ridge plot. Gene signatures were ordered based on the median AUCs given by the original models. **(A)** Comparison of AUC distributions given by ssGSEA (evaluated on the upregulated subsets of the gene signatures), PLAGE, and the original models. **(B)** Comparison of AUC distributions given by five gene set scoring methods (ssGSEA, PLAGE, GSVA, Singscore, Zscore) and the original models. **(C)** Comparison of AUC distributions given by ssGSEA (evaluated on the upregulated and downregulated subsets of gene signatures), GSVA (evaluated on the upregulated and downregulated subsets of gene signatures), and the original models.

Supplementary Figure 4. Spearman's rank correlation versus AUC difference for studies with sample size larger than 40 and computed AUC values greater than 0.8 based on the results given by original models and ssGSEA **(A)**, original models and PLAGE **(B)**, original models and GSVA **(C)**, original models and Zscore **(D)**, original models and Singscore (unidirectional version) **(E)**. Overlapping points indicate the same dataset(s).

Supplementary Figure 5. Spearman's rank correlation versus AUC difference for studies with sample size larger than 40 and computed AUC greater than 0.8 based on results given by original models and ssGSEA (evaluated with upregulated subsets) **(A)**, original models and ssGSEA (evaluated with downregulated subsets) **(B)**, original models and GSVA (evaluated with upregulated subsets) **(C)**, original models and GSVA (evaluated with downregulated subsets) **(D)**, original models and Singscore (bidirectional version) **(E)**. Overlapping points indicate the same dataset(s).