Title: The scheduling of adolescence with Netrin-1 and UNC5C

Authors

Daniel Hoops[1,2], Robert F. Kyne[3], Samer Salameh[2,4], Elise Ewing[2,4], Alina T. He[2,4], Taylor Orsini[2,4],

Anais Durand[2,4], Christina Popescu[2,4], Janet M. Zhao[2,4],Kelcie C. Schatz[5],LiPing Li[5], Quinn E.

Carroll[5], Guofa Liu[6], Matthew J. Paul[3,5 #], Cecilia Flores[1,2,7,# *]

Affiliations

1 Department of Psychiatry, McGill University, Montréal, Quebec, Canada

2 Douglas Mental Health University Institute, Montréal, Quebec, Canada

3 Neuroscience Program, University at Buffalo, SUNY, New York, USA

4 Integrated Program in Neuroscience, McGill University, Montréal, QC, Canada

5 Department of Psychology, University at Buffalo, SUNY, New York, USA

6 Department of Biological Sciences, University of Toledo, Ohio, USA

7 Department of Neurology and Neurosurgery, McGill University, Montréal, Quebec, Canada

Co-senior authors

* Corresponding author C. Flores: cecilia.flores@mcgill.ca

Abstract

Dopamine axons are the only axons known to grow during adolescence. Here, using rodent

models, we examined how two proteins, Netrin-1 and its receptor, UNC5C, guide dopamine

axons towards the prefrontal cortex and shape behaviour. We demonstrate in mice (*Mus*

musculus) that dopamine axons reach the cortex through a transient gradient of Netrin-1

expressing cells – disrupting this gradient reroutes axons away from their target. Using a

seasonal model (Siberian hamsters; *Phodopus sungorus*) we find that mesocortical dopamine

development can be regulated by a natural environmental cue (daylength) in a sexually

dimorphic manner – delayed in males, but advanced in females. The timings of dopamine axon

growth and UNC5C expression are always phase-locked. Adolescence is an ill-defined,

transitional period; we pinpoint neurodevelopmental markers underlying this period.

Introduction

Adolescence is a critical developmental period involving dramatic changes in behaviour and

brain anatomy. The prefrontal cortex, the brain region responsible for our most complex

cognitive functions, is still establishing connections during this time (Gogtay et al., 2004;

Petanjek et al., 2011; Sowell et al., 2004). The trajectory of prefrontal cortex development in

adolescence determines the vulnerability or resilience of individuals to adolescent-onset

psychiatric diseases (Fuhrmann et al., 2015; Keshavan et al., 2014; Kessler et al., 2007, 2005;

Lee et al., 2014). The age at which this adolescent development occurs therefore represents a

critical window during which the brain is particularly susceptible to environmental influences.

Traditionally, the onset of adolescence is thought to coincide with puberty (Hollenstein and

Lougheed, 2013). In humans, the age of pubertal onset has been advancing throughout the

19th, 20th and 21st centuries, and environmental influences, such as nutrition, can pathologically

alter the age of puberty (Wolf and Long, 2016). However, it remains entirely unknown whether

the neural and cognitive maturational processes of adolescence can also be plastic. Here, for

the first time, we examine how the timing of adolescence is programmed in the brain, and

whether this timing can be plastic in response to a natural environmental cue, in parallel with

pubertal plasticity.

Dopamine innervation to the prefrontal cortex increases substantially across adolescence and

psychopathologies of adolescent origin prominently feature dopamine dysfunction. Evidence

continues to emerge that protracted dopamine innervation is a key neural process underlying

the cognitive and behavioural changes that characterize adolescence (Larsen and Luna, 2018).

The mesocorticolimbic dopamine system – which includes the prefrontal cortex – is unique

because not only are connections being formed and lost during adolescence, but there is also

long-distance displacement of dopamine axons between brain regions. At the onset of

adolescence, both mesolimbic and mesocortical dopamine axons innervate the nucleus

accumbens in rodents, but the mesocortical axons leave the accumbens and grow towards the

prefrontal cortex during adolescence and early adulthood (Hoops et al., 2018; Reynolds et al.,

2018a). This is the only known case of axons growing from one brain region to another so late

during development (Hoops and Flores, 2017).

The prolonged growth trajectory renders mesocortical dopamine axons particularly vulnerable

to disruption. Environmental insults during adolescence (e.g. drug abuse) alter the extent and

organization of dopamine innervation in the prefrontal cortex, leading to behavioural and

cognitive changes in mice throughout adulthood (Drzewiecki and Juraska, 2020; Hoops and

Flores, 2017; Reynolds and Flores, 2021). These changes often involve cognitive control, a

prefrontal function that develops in parallel with dopamine innervation to the cortex in

adolescence (Luna et al., 2015). Disruption of dopamine innervation frequently seems to result

in "immature" cognitive control persisting through adulthood (Larsen and Luna, 2018).

Here, we examine the guidance of growing dopamine axons to the prefrontal cortex, and its

timing. The guidance cue molecule Netrin-1, upon interacting with its receptor DCC, determines

which dopamine axons establish connections in the nucleus accumbens and which ones leave

this region to grow to the prefrontal cortex (Hoops and Flores, 2017; Reynolds and Flores,

2021). We hypothesized that the answers to *how* and *when* this extraordinary developmental

feat is achieved may also lie in the Netrin-1 signalling system.

Part 1: Netrin-1 "paves the way" for dopamine axons in adolescence

To identify the route by which dopamine axons grow from the nucleus accumbens to the

medial prefrontal cortex we visualized dopamine axons in the adult mouse forebrain. We

observed that dopamine axons medial to the nucleus accumbens form a distinct bundle

oriented dorsally towards the cortex (Figure 1A,B). Individual fibres can be seen crossing the

boundary between the nucleus accumbens shell and this bundle (Figure 1C). We hypothesized

that these are the fibres that grow to the prefrontal cortex during adolescence. If this is correct,

the number of dopamine axons in the bundle should continue to increase until adulthood. To

test this, we used a modified unbiased stereological approach (Kim et al., 2011) where axons

are counted only if they crossed the upper and lower bounds of a counting probe. We also

measured the average width of the dopamine axon bundle. We found, in both male and female

mice, that the density of dopamine axons does not change between adolescence (21 days old)

and adulthood (75 days old; Figure 1D). However, the width of the dopamine axon bundle does

change, increasing between adolescence and adulthood (Figure 1E). These results indicate that

the total number of dopamine axons passing through this bundle increases over adolescence

and that dopamine axons grow to the medial prefrontal cortex via this route.

Figure 1. A "pathway" of Netrin-1 expressing cells "paves the way" for dopamine (DA) axons growing from the

nucleus accumbens to the medial prefrontal cortex during adolescence. **A**, The brain regions containing the bundle

of dopamine fibres passing to the medial prefrontal cortex are highlighted in a line drawing of a coronal mouse

brain section derived from Paxinos and Franklin (Paxinos and Franklin, 2013). **B**, An image of a coronal section

through the forebrain of an adult mouse at low magnification (4x). Green fluorescence indicates immunostaining

for tyrosine hydroxylase (TH), a marker for dopamine. The smaller and larger white squares indicate the regions

enlarged in panel C and panels F & G, respectively. Scale bar = 500 µm. **C**, The nucleus accumbens (left) is densely

packed with dopamine axons (in green). Some of these axons can be observed extending from the nucleus

accumbens medially into a bundle of dopamine fibres (white arrows). Scale bar = 10 µm. **D**, Modified stereological

quantification revealed no significant difference in dopamine axon density in the fibre bundle between

adolescence (21 days old) and adulthood (75 days old). Mixed-effects ANOVA, effect of age: $F=1.53$, $p=0.22$; region

by age interaction: $F=1.44$, $p=0.49$. **E**, The average width of the fibre bundle increased significantly from

adolescence to adulthood, revealing that there is an increase in the total number of fibres passing to the medial

prefrontal cortex during this period. Mixed-effects ANOVA, effect of age: $F=9.45$, $p=0.0021$; region by age

interaction: $F=5.74$, $p=0.057$.   **F**, In order to quantify the Netrin-1 positive cells along the dopamine fibre bundle,

the bundle was contoured in each region, and a contour of equal area was placed medial to the fibre bundle as a

negative control. Scale bar = 200 µm. **G**, Using quantitative stereology, Netrin-1 positive cell density was

determined along and adjacent to the fibre bundle for each region. Red fluorescence indicates immunostaining for

Netrin-1. **H**, In adolescent mice there are more Netrin-1 positive cells along the dopamine fibre bundle ("TH+")

than medial to it ("TH-"). This is what we refer to as the "Netrin-1 pathway". Along the pathway, there is a

significant increase in Netrin-1 positive cell density in regions closer to the medial prefrontal cortex, the

innervation target. Mixed-effects ANOVA, effect of TH: $F=105$, $p<0.0001$. Effect of region: $F=9.74$, $p=0.021$. A post-

hoc Tukey Test revealed a difference ($p = 0.029$) between the densities of the lateral septum and infralimbic

cortex, but only within the dopamine fibre bundle. **I**, In adult mice the Netrin-1 pathway is maintained, however

there is no longer an increasing density of Netrin-1 positive cells towards the medial prefrontal cortex. Mixed-

effects ANOVA, effect of TH: $F=54.56$, $p<0.0001$. Effect of region: $F=1.22$, $p=0.75$. **J,** The virus injection location

within the mouse brain. A Netrin-1 knockdown virus, or a control virus, was injected into the dopamine fibre

bundle at the level of the dorsal peduncular cortex. **K,** Our experimental timeline: at the onset of adolescence a

Netrin-1 knockdown virus, or a control virus, was injected, and then in adulthood the mice were sacrificed and

stereological measurements taken. **L,** Dopamine varicosity density was quantified in the region below the injection

site, in the lateral septum, and in the region above the injection site, in the infralimbic cortex. There was a

significant decrease in dopamine varicosity density only in the infralimbic cortex. Mixed-effects ANOVA, virus by

region interaction: F=16.41, p<0.0001. **M,** The experimental set-up of the final (test) stage of the Go/No-Go

experimental protocol. A mouse that has previously learned to nose-poke for a reward in response to a visual cue

(illuminated nose-poke hole), must now inhibit this behaviour when the visual cue is paired with an auditory cue

(acoustic tone). **N,** Mice injected with the Netrin-1 knockdown virus show improved action impulsivity than

controls because they incur significantly fewer commission errors across the Go/No-go task. Mixed-effects ANOVA,

effect of day: F=68.32, p<0.0001. Day by virus interaction: F=9.00, p=0.0027. A sigmoidal curve is fit to each group

of mice to determine how the two groups differ. Points indicate group means and error bars show standard error

means. **O**, During the first days of Go/No-Go testing, both groups incur commission errors with high frequency,

but, the Netrin-1 knockdown group has fewer errors than the control group (t-test, t=5.18, p<0.0001. **P,** The ED50

– the inflection point in each sigmoidal curve – are not different between groups, indicating that all mice improve

their ability to inhibit their behavior at around the same time (T-test, t=0.97, p=0.35). **Q,** Mice microinfused with

the Netrin-1 knockdown virus incur substantially fewer commission errors in the last days of the Go/No-Go task

compared to mice injected with the control virus (T-test, t=12.38, p<0.0001). For all barplots, bars indicate group

means and error bars show standard error means.

Next, we focussed on Netrin-1, a secreted protein that acts as a guidance cue to growing axons

and is important for dopamine axon targeting in the nucleus accumbens in adolescence (Cuesta

et al., 2020). Using unbiased stereology, we quantified the number of Netrin-1 expressing cell

bodies along the dopamine axon route, and in an adjacent medial region as a control (Figure 1

F, G). We found that in adolescence there are more Netrin-1 positive neurons within the

dopamine axon route than adjacent to it. Furthermore, along the axon route the density of

Netrin-1 positive cells increases towards the medial prefrontal cortex, forming a dorsoventral

gradient (Figure 1H). In adulthood, there remains a higher density of Netrin-1 positive cells

along the dopamine route compared to the adjacent region, however the dorsoventral gradient

is no longer present (Figure 1I).

To determine if Netrin-1 along the dopamine axon route is necessary for axon navigation, we

silenced Netrin-1 expression in the dorsal peduncular cortex, the transition region between the

septum and the medial prefrontal cortex, at the onset of adolescence (Figure 1J,K). In

adulthood we quantified the number of dopamine axon terminals in the regions below and

above the injection site. Silencing Netrin-1 did not alter dopamine terminal density below the

injection, in the lateral septum (Figure 1L). In the infralimbic cortex, which is the first prefrontal

cortical region the axons reach after the injection site, terminal density was reduced in the

Netrin-1 knock-down group compared to controls (Figure 1L). The knock-down appears to erase

the Netrin-1 path to the prefrontal cortex, resulting in dopamine axons losing their way and

failing to reach their correct innervation target. We conclude that Netrin-1 expressing cells

"pave the way" for the mesocortical dopamine axons.

We next examined how the Netrin-1 path may be important for behaviour. The dopamine input

to the prefrontal cortex is a key factor in the transition from juvenile to adult behaviours that

occurs in adolescence. We hypothesized that cognitive processes involving mesocortical

dopamine function would be altered when these axons are misrouted in adolescence. To test

our hypothesis we used the Go/No-Go behavioural paradigm. This test quantifies inhibitory

control, which matures in parallel with the innervation of dopamine axons to the prefrontal

cortex in adolescence (Casey et al., 2008; Klune et al., 2021; Luna et al., 2015; Paus, 2005;

Reynolds and Flores, 2021; Spear, 2000) and it is impaired in adolescent-onset disorders like

depression and schizophrenia (Catts et al., 2013; Clementz et al., 2016; McTeague et al., 2016;

Millan et al., 2012).

At the onset of adolescence, we injected the Netrin-1 silencing, or a scrambled control virus,

bilaterally into the dorsal peduncular cortex; in adulthood we tested the mice in the Go/No-Go

task. This paradigm first involves discrimination learning and reaction time training

(Supplementary Figure 1), followed by a Go/No-Go test consisting of "Go" trials where mice

respond to a cue as previously trained and "No-Go" trials where mice must abstain from

responding to the cue (Figure 1M). Correct responses to both trial types are reinforced with a

food reward. We quantified the percent of "No-Go" trials where the mice incorrectly responded

to the cue ("Commission Errors") and the percent of "Go" trials where the mice correctly

responded ("Rewards" or "Hits"; Supplementary Figure 2). The ability of mice to respond

correctly overall to both trial types is quantified as the Correct Response Rate (Supplementary

Figure 3) (Cuesta et al., 2019; Reynolds et al., 2018a, 2018b; Vassilev et al., 2021).

Mice injected with the Netrin-1 silencing virus differed from controls in their performance

during "No-Go" trials. As the mice learn to withhold their responses over the course of the test,

the number of commission errors they made in No-Go trials decreased in a sigmoidal fashion

(Figure 1N). The upper and lower asymptotes of the sigmoidal curve quantify the number of

commission errors committed during early and late test days, respectively, while the inflection

point (ED50) indicates when mice start improving their ability to inhibit their behavior. Already

at the start of the Go/No part of the task, the Netrin-1 silencing group make slightly fewer

commission errors (Figure 1O) than control groups, although both groups begin to improve in

the No-Go task at around the same time. However, the Netrin-1 silencing group achieved a

substantially higher level of behavioural inhibition, quantified as a lower percentage of

commission errors in the last test days (Figure 1Q), indicating an improved ability to withhold

their behaviour on cue. These behavioural results demonstrate that the maturation of action

impulsivity is sensitive to the organization of the ventro-dorsal Netrin-1 path that guides

mesocortical dopamine axon growth. Deviations in this route lead to striking changes in the

cognitive development that is characteristic of adolescence.

Part 2: UNC5C expression coincides with the onset of adolescence

When axons leave the nucleus accumbens during adolescence, they follow a Netrin-1 "path"

through intermediate brain regions to reach their intended innervation target. However, only a

small subset of the dopamine axons that have reached the nucleus accumbens by early

adolescence leave; the vast majority stay and form connections in the accumbens throughout

life (Reynolds et al., 2018a). The "decision making" process of whether to "stay" (in the

accumbens) or "go" (to the cortex via the Netrin-1 path) happens during a narrow

developmental window at the onset of adolescence (Reynolds et al., 2018b). It remains

unknown how the timing of this process is determined.

Figure 2 The age of onset of UNC5C expression by dopamine axons in the nucleus accumbens of mice is sexually

dimorphic. Images are representative of observed immunofluorescence patterns in the nucleus accumbens core

(approx. location highlighted in the coronal mouse brain section Plate 19, modified from Paxinos & Franklin, 2013).

For each row, six individuals were sampled. In males (**A-B**), UNC5C expression on dopamine fibres in the nucleus

accumbens appears during adolescence. **A,** At the onset of adolescence (21 days old) dopamine fibres do not

express UNC5C. Scale bar = 10 µm. **B,** By adulthood (90 days old), dopamine fibres express UNC5C. In females (**C-**

E), UNC5C expression on dopamine fibres in the nucleus accumbens appears prior to adolescence. **C,** In juvenile

(15 day old) mice, there is no UNC5C expression on dopamine fibres. **D,** By adolescence, dopamine fibres express

UNC5C. **E,** In adulthood, dopamine fibres continue to express UNC5C.

In adolescence, dopamine neurons begin to express the repulsive Netrin-1 receptor UNC5C, and

reduction in UNC5C expression appears to cause growth of mesolimbic dopamine axons to the

prefrontal cortex (Auger et al., 2013; Manitt et al., 2010; Pokinko et al., 2015). Using

immunohistochemistry, we assessed the expression of UNC5C on nucleus accumbens dopamine

axons across development. In male mice, we found little expression of UNC5C on dopamine

axons at the onset of adolescence (Figure 2A), while we did find UNC5C expression on

dopamine axons in adults (Figure 2B). Remarkably, when we assessed this in females, we found

dopamine axons already expressing UNC5C in the nucleus accumbens at the onset of

adolescence (Figure 2D), similar to adult females (figure 2E), indicating that the onset of UNC5C

expression on dopamine axons in the nucleus accumbens is sexually dimorphic, with an earlier

emergence in females. We examined the nucleus accumbens in pre-adolescent female mice

and indeed found little UNC5C expression on dopamine axons (Figure 2C). The onset of UNC5C

expression in mesocorticolimbic dopamine axons is therefore peri-adolescent but occurs earlier

in females than in males, consistent with the earlier emergence of adolescence in female

rodents and the earlier onset of adolescence and puberty in humans (Wolf and Long, 2016).

Part 3: Environmental control of the timing of adolescence

We hypothesize that at the emergence of adolescence, UNC5C expression by dopamine axons

in the nucleus accumbens signals the initiation of the growth of dopamine axons to the

prefrontal cortex. We therefore examined whether the developmental timings of UNC5C

expression and dopamine innervation of the prefrontal cortex are similarly affected by an

environmental cue known to delay pubertal development in seasonal species.

Siberian hamsters (*Phodopus sungorus*) regulate many aspects of their behavior and physiology

to meet the changing environmental demands of seasonality (Paul et al., 2008; Stevenson et al.,

2017). In winter, they increase the thickness of their fur, exchange their brown summer coats

for white winter ones, and undergo a daily torpor to conserve energy (Scherbarth and

Steinlechner, 2010). In addition, adults suppress reproduction and juveniles delay puberty

(Pevet, 1988; Yellon and Goldman, 1984), including developmental changes in gonadotropin

releasing hormone neurons in the hypothalamus (Buchanan and Yellon, 1991; Heywood and

Yellon, 1997). This seasonal plasticity is regulated by long or short periods of daylight

(Heldmaier and Steinlechner, 1981; Hoffmann, 1978) and raises the possibility that aspects of

adolescent development are sensitive to these environmental cues. To our knowledge,

adaptive variation in the timing of adolescent neural development has never been recorded in

any animal.

Here, we tested whether day length regulates *when* dopamine axons grow to the cortex, and

whether the timing of UNC5C expression in the nucleus accumbens and adolescent changes in

exploratory behavior are similarly affected.

3.i The seasonality of adolescence

Male hamsters were examined at three ages: 15 days old (±1), 80 days old (±10), and 215 days

old (±20). We compared the density of the dopamine innervation to the medial prefrontal

cortex in male hamsters housed under lighting conditions that replicate summer daylengths

(long days, short nights) or winter daylengths (short days, long nights) (Figure 3A,B). We will

refer to these two groups as "summer hamsters" and "winter hamsters" to emphasize the

natural stimulus we are replicating in the laboratory environment. We confirmed that puberty

is delayed in male winter hamsters compared to summer hamsters in the present experiment

by measuring their gonadal weights across ages (Supplementary Figure 4).

In male summer hamsters, dopamine input density to the prefrontal cortex increases during

adolescence, after 15 days old and before 80 days old (Figure 3C), consistent with dopamine

axon growth in mice (Manitt et al., 2013, 2011; Reynolds et al., 2018a). Prefrontal cortex

dopamine innervation in summer hamsters continues to increase after 80 days old (Figure 3C).

In male winter hamsters, dopamine innervation to the prefrontal cortex is delayed until after 80

days, which coincides with their delayed pubertal onset (Figure 3D, Supplementary Figure 2B).

This is the first demonstration of an environmental cue shaping adolescent brain development.

Figure 3 Plasticity of adolescent development in male Siberian hamsters according to seasonal phenotype. All

results illustrated in this figure refer to results in male hamsters. **A,** Dopamine innervation was quantified in three

subregions of the medial prefrontal cortex, highlighted in blue. UNC5C expression was examined in the nucleus

accumbens, highlighted in red. Line drawing of a coronal section of the mouse brain was derived from Plate 19 of

Paxinos and Franlkin (Paxinos and Franklin, 2013). **B,** Hamsters were housed under either summer-mimicking long

days and short nights ("summer hamsters") or winter-mimicking short days and long nights ("winter hamsters"). **C,**

In male hamsters housed under a summer-mimicking daylength there is an increase in dopamine varicosity density

in the medial prefrontal cortex between 15 and 80 days old. Mixed-effects ANOVA, effect of age: F=9.6,

p=0.000255. Tukey Test, 15-80 days old (do): p=0.026; 80-215do: p<0.0001; 15-215do: p<0.0001. **D,** In male

hamsters housed under a winter-mimicking daylength there is no increase in dopamine varicosity density until

hamsters have reached 215 days of age. Mixed-effects ANOVA, effect of age: F=4.17, p=0.0205. Tukey Test, 15-

80do: p=0.54; 80-215do: p=0.0006; 15-215do: p=0.0085. **E,** At 15 days old, dopamine axons in the nucleus

accumbens of male summer-daylength hamsters largely do not express UNC5C. Scale bar = 20um (bottom right). **F-**

G, At 80 (**F**) and 215 (**G**) days old, dopamine axons in the nucleus accumbens express UNC5C. **H-I,** At 15 (**H**) and 80

(**I**) days old, dopamine axons in the nucleus accumbens of male winter hamsters largely do not express UNC5C. **J,**

By 215 days old there is UNC5C expression in dopamine axons in the nucleus accumbens of male winter hamsters.

E-J, Representative images of the nucleus accumbens shell, 6 individuals were examined per group. **K,** Male

hamsters house under a summer-mimicking daylength show an adolescent peak in exploratory behaviour in the

light/dark box apparatus. Instead, those raised under a winter-mimicking photoperiod show a steady increase in

exploratory behaviour over the same age range. Arrows indicate the ages at which exploratory behaviour peaks in

summer (orange) and winter (blue) hamsters. Polynomial regression, effect of season: F=3.551, p=0.00056. **L,** In

male hamsters, at 215 days of age, there is no difference in exploratory behaviour between hamsters raised under

summer and winter photoperiods. T-test, effect of season: t=0.975, p=0.341. For all barplots, bars indicate group

means and error bars show standard error means.

We then examined UNC5C expression by dopamine axons in the nucleus accumbens in male

summer and winter hamsters across age classes. UNC5C expression was apparent only after the

onset of adolescence in summer hamsters (Figure 3E,F,G), as observed in male mice. However,

UNC5C expression was delayed in male winter hamsters – this group did not show UNC5C

expression in dopamine axons in the nucleus accumbens until after 80 days old (Figure 3H,I,J).

This aligns with the delayed timing of mesocortical dopamine axon growth and pubertal onset

in male winter hamsters and demonstrates that the emergence of UNC5C is a marker of

adolescent onset in male mice.

A behaviour characteristic of adolescence is increased willingness to enter a novel environment,

called exploratory behaviour (Arrant et al., 2013; Lynn and Brown, 2009); to measure this we

used the light/dark test (Bourin and Hascoët, 2003). Time spent in the light compartment is

dopamine-dependent (Bahi and Dreyer, 2019; Gao and Cutler, 1993) and peaks in adolescence

(Arrant et al., 2013). We assessed the developmental profile of exploratory behavior in the

light/dark box test in summer and winter hamsters across adolescence. In male summer

hamsters, the exploratory behavior increases across adolescence, peaks around 50 days, then

subsequently declines (Figure 3K). However, the adolescent increase in exploratory behavior is

protracted in winter hamsters: across the age range examined we observe a gradual, consistent

increase in exploratory behaviour rather than a peak and decline.

We next assessed a cohort of 215-day old hamsters, for which both summer and winter male

hamsters have undergone puberty and exhibit high levels of dopamine innervation of the

prefrontal cortex (Figure 3C,D,G,J, Supplementary Figure 2B). In these hamsters, we find no

difference in exploratory behaviour between the male summer and winter groups (Figure 3L),

demonstrating that, after 80 days, exploratory behaviour begins to decline in male winter

hamsters and that by 215 days it has declined to the same level as in summer hamsters. Male

hamsters raised under summer-mimicking long days and winter-mimicking short days both

ultimately make the transition to the adult behavioral phenotype.

3.ii An extraordinary case of decoupling puberty and adolescence

In parallel with males, we conducted equivalent experiments in female hamsters (Figure 4A,B).

Under a summer-mimicking daylength, dopamine innervation to the medial prefrontal cortex

increases between 15 and 80 days old, similar to male summer hamsters (Figure 4C). There is

no further increase in innervation density after 80 days old, consistent with earlier adolescent

development in females observed in other rodent species (Juraska and Willing, 2017; Kopec et

al., 2018; Reynolds and Flores, 2021; Spear, 2000; Westbrook et al., 2018). We confirmed that

puberty is delayed in female winter hamsters compared to summer hamsters by measuring

their uterine weights across ages (Supplementary Figure 2A).

Figure 4 Plasticity of adolescent development in female Siberian hamsters according to seasonal phenotype. All

results illustrated in this figure refer to results in female hamsters. **A,** Dopamine innervation was quantified in

three subregions of the medial prefrontal cortex, highlighted here in blue. UNC5C expression was examined in the

nucleus accumbens, highlighted in red. Line drawing of a coronal section of the mouse brain was derived from

Paxinos and Franlkin (Paxinos and Franklin, 2013). **B,** Hamsters were housed under either a summer-mimicking or

winter-mimicking daylength. **C,** In female hamsters housed under a summer daylength dopamine varicosity density

in the medial prefrontal cortex increases between 15 and 80 days of age. Mixed-effects ANOVA, effect of age:

F=16.72, p<0.0001 **D,** In female hamsters housed under a winter daylength there is no increase in dopamine

varicosity density post-adolescence. Instead, there is a steep decline in density between 80 and 215 days of age.

Mixed-effects ANOVA, effect of age: F=12.33, p=0.000043. **E,** As our results in panel D were unexpected, we

replicated them with a second cohort of hamsters and found qualitatively identical results. Mixed-effects ANOVA,

effect of age: F=34.871, p<0.0001. **F,** To try and determine when dopamine varicosities innervate the medial

prefrontal cortex, we examined a cohort of 10- and 15-day-old hamsters. We found that varicosity density

increases in the medial prefrontal cortex during this time, indicating that dopamine innervation to the medial

prefrontal cortex is accelerated in female winter hamsters. Mixed-effects ANOVA, effect of age: F=5.05, p=0.03. **G-**

H, In 10- and 15-day-old female summer hamsters there is little UNC5C expression in nucleus accumbens

dopamine axons. Sample size: 4 (panel G) or 6 (panel H). **I-J,** By 80 days old (panel I), and continuing at 215 days old

(panel J), dopamine axons in the nucleus accumbens express UNC5C in female summer hamsters. Sample sizes: 6.

Scale bar = 20um (panel G bottom right). **K-N,** At all ages at which winter female hamsters were examined

dopamine axons in the nucleus accumbens express UNC5C in winter female hamsters. Sample sizes: 4 (panel K) or

6 (panels L-N). **O,** In female hamsters, those raised under summer and winter daylengths both show an increase in

exploratory behaviour over time. The winter hamsters peak later compared to the summer daylength hamsters.

Arrows indicate the ages at which exploratory behaviour peaks in summer (orange) and winter (blue) hamsters.

Polynomial regression, effect of season: F=3.305, p=0.00126. **P,** In female hamsters, at 215 days of age, there is no

difference in exploratory behaviour between hamsters raised under summer and winter photoperiods. T-test,

effect of season: t=0.309, p=0.76. For all barplots, bars indicate group means and error bars show standard error

means.

When housed under a winter-mimicking daylength, dopamine input density in the prefrontal

cortex of female hamsters is *not* delayed as in males, but rather reaches adult levels prior to 15

days old (Figure 4D). We replicated this unexpected finding in a separate, independent cohort

of female winter hamsters (Figure 4E). To our knowledge, this is the first time an intervention is

shown to accelerate adolescent cortical development.

We then measured dopamine axon density in female winter hamsters at two earlier ages: 10

and 15 days old. Dopamine innervation increases during this period (figure 4F), well before

normal adolescence and long before pubertal development. This is an extraordinary

phenomenon: a key marker of adolescent neurodevelopment is accelerated and dissociated

from puberty in female hamsters raised under winter-mimicking short days (Supplementary

Figure 2A).

The early increase in prefrontal cortex dopamine terminals in winter females is followed by a

dramatic reduction between 80 and 215 days old (Figure 4D,E). This overlaps with the delayed

timing of puberty in these females (Butler et al., 2007; Supplementary Figure 2A). Synaptic

pruning in the cortex is a well-known component of adolescent neural development across

species (Huttenlocher, 1984; Koss et al., 2013; Petanjek et al., 2011). Under normal conditions,

the effect of pruning on dopamine synapses is likely masked by the growth of new dopamine

axons to the prefrontal cortex (Manitt et al., 2013, 2011; Reynolds et al., 2018a). In the case of

female winter hamsters, we hypothesize that the growth of dopamine axons to the prefrontal

cortex occurs early while synaptic pruning, including of dopamine synapses, appears to occur

later. This leads to a remarkable dissociation between two cortical developmental processes

that are normally simultaneous, the behavioural implications of which are unclear.

If the developmental onset of UNC5C expression determines the timing of dopamine

innervation of the prefrontal cortex, then onset of UNC5C expression should also be advanced

in female winter hamsters. Hence, we examined UNC5C expression at the same ages as we

examined dopamine axon growth in female hamsters. At 10 and 15 days old, UNC5C

expression is present *only* in the winter hamsters (Figure 4G,H,K,L), but at 80 and 215 days old,

UNC5C expression is apparent in both summer and winter hamsters (Figure 4I,J,M,N).

We used the light/dark box test to examine potential exploratory behavioural implications of

the extraordinary developmental trajectory we observed in the prefrontal cortex of female

hamsters. In female summer and winter hamsters the adolescent increase and peak in

exploratory behaviour occurs between the ages of 15 and 80 days, as it does in summer

daylength males (Figure 4O). However, contrary to what we would expect, the peak in winter

females is delayed compared to summer females. When we assessed an independent cohort of

215-day-old female hamsters, we found no difference in exploratory behaviour between groups

(Figure 4R), indicating that, like males, female summer and winter hamsters both eventually

reach the same adult level of exploratory behaviour.

In both sexes, hamsters housed under a summer-mimicking daylength showed an adolescent

peak in exploratory behaviour at an age that we would predict based on results from other

rodents (Arrant et al., 2013; Pietropaolo et al., 2004; Tanaka, 2015). When raised under a

winter-mimicking daylength, hamsters of either sex show a protracted exploratory peak. In

males, it is delayed beyond 80 days old, but the delay is substantially less in females. For

dopamine and UNC5C neural markers, males delayed, whereas females advanced, adolescent

neural development. Based on the delayed developmental profile in exploratory behavior, we

predict that other adolescent neural changes are delayed in female winter hamsters. We

401 hypothesize that the advanced dopamine/UNC5C neural development lessens the delay in peak

402 exploratory behaviour of female winter hamsters compared to that of males.

403 **Conclusion**

404 Here we describe how the gradual growth of mesocortical dopamine axons marks adolescent

405 development, and how this process uses guidance cues and is sensitive to sex and environment.

406 Netrin-1 signalling provides the "stay-or-go" "decision making" conducted by dopamine axons

407 that innervate the nucleus accumbens at the onset of adolescence (Cuesta et al., 2020). UNC5C

408 expression by these dopamine axons marks the timing at which this decision is made. In mice,

409 UNC5C expression coincides with sex differences in both adolescent and pubertal development.

410 Females, which develop earlier, show earlier UNC5C expression in dopamine axons compared

411 to males.

412 In hamsters, behavioural and developmental shifts in response to environmental cues occur in

413 parallel with alterations in the timing of dopamine axon growth. As we show here, male

414 hamsters raised under a winter-mimicking daylength delay not only puberty, but also

415 adolescent dopamine and behavioural maturation. In contrast, female hamsters under identical

416 conditions delay puberty but accelerate dopamine axon growth, a key marker of adolescent

417 brain development. Behavioural shifts during adolescence appear to be delayed in these

418 females, but less substantially than in male hamsters. Notably, under all conditions, the

419 developmental timing of UNC5C expression corresponds to the timing of dopamine innervation

420 of the prefrontal cortex.

In both mice and hamsters, the emergence of UNC5C expression coincides with the onset of

dopamine axon growth to the prefrontal cortex, a key characteristic of the adolescent transition

period. While previously we have shown that the Netrin-1 signalling in the nucleus accumbens

is responsible for coordinating *whether* dopamine axons grow in adolescence (Reynolds and

Flores, 2021), here we propose that Netrin-1 signalling is also key to determining *how* and *when*

this marker of adolescence occurs.

Acknowledgments

We are grateful for the technical assistance of the Pathology Core, Centre for Phenogenomics,

Toronto, Canada, and in particular Milan Ganguly and Gregory Ossetchkine, for their excellent

histological assistance. Furthermore, we are grateful to Helen Cooper and Philip Vassilev for

their critical and insightful readings of our manuscript.

Author contributions:

Conceptualization: DH, MJP, CF

Data Curation: DH

Formal Analysis: DH

Investigation: DH, RFK, SS, EE, AH, TO, AD, MG, CP, JZ, KCS, LPL, QEC

Methodology: DH, MJP, CF

Resources: GL, MJP, CF

Supervision: DH, MJP, CF

Validation: DH

Visualization: DH

Writing—original draft: DH

Writing—review & editing: DH, MJP, CF

Funding:

National Institute on Drug Abuse grant R01DA037911 (CF)

Canadian Institutes of Health Research FRN: 156272 (CF)

Canadian Institutes of Health Research FRN: 170130 (CF)

National Science and Engineering Research Council of Canada RGPIN-2020-04703 (CF)

National Science Foundation IOS-1754878 (MJP)

National Science and Engineering Research Council of Canada PGSD3-415253-2012 (DH)

Quebec Nature and Technology Research Fund 208332 (DH)

National Science and Engineering Research Council of Canada PDF5171462018 (DH)

Competing interests: Authors declare that they have no competing interests.

Data and code availability: All data and code use in these analyses are available through the

Open Science Framework (DIO 10.17605/OSF.IO/DU3H4).

References

458
459

Arrant, A.E., Schramm-Sapyta, N.L., Kuhn, C.M., 2013. Use of the light/dark test for anxiety in
adult and adolescent male rats. Behav Brain Res 256, 119–127.
https://doi.org/10.1016/j.bbr.2013.05.035

Auger, M.L., Schmidt, E.R.E., Manitt, C., Dal-Bo, G., Pasterkamp, R.J., Flores, C., 2013. unc5c
haploinsufficient phenotype: striking similarities with the dcc haploinsufficiency model.
European Journal of Neuroscience 38, 2853–2863. https://doi.org/10.1111/ejn.12270

Bahi, A., Dreyer, J.-L., 2019. Dopamine transporter (DAT) knockdown in the nucleus accumbens
improves anxiety- and depression-related behaviors in adult mice. Behav Brain Res 359,
104–115. https://doi.org/10.1016/j.bbr.2018.10.028

Bourin, M., Hascoët, M., 2003. The mouse light/dark box test. Eur J Pharmacol 463, 55–65.
https://doi.org/10.1016/s0014-2999(03)01274-3

Buchanan, K.L., Yellon, S.M., 1991. Delayed Puberty in the Male Djungarian Hamster: Effect of
Short Photoperiod or Melatonin Treatment on the GnRH Neuronal System.
Neuroendocrinology 54, 96–102. https://doi.org/10.1159/000125857

Butler, M.P., Trumbull, J.J., Turner, K.W., Zucker, I., 2007. Timing of puberty and
synchronization of seasonal rhythms by simulated natural photoperiods in female Siberian
hamsters. Am J Physiology-regulatory Integr Comp Physiology 293, R413–R420.
https://doi.org/10.1152/ajpregu.00216.2007

Casey, B.J., Jones, R.M., Hare, T.A., 2008. The Adolescent Brain. Annals of the New York
Academy of Sciences 1124, 111–126. https://doi.org/10.1196/annals.1440.010

Catts, V.S., Fung, S.J., Long, L.E., Joshi, D., Vercammen, A., Allen, K.M., Fillman, S.G., Rothmond,
D.A., Sinclair, D., Tiwari, Y., Tsai, S.-Y., Weickert, T.W., Weickert, C.S., 2013. Rethinking
schizophrenia in the context of normal neurodevelopment. Front Cell Neurosci 7, 60.
https://doi.org/10.3389/fncel.2013.00060

Clementz, B.A., Sweeney, J.A., Hamm, J.P., Ivleva, E.I., Ethridge, L.E., Pearlson, G.D., Keshavan,
M.S., Tamminga, C.A., 2016. Identification of Distinct Psychosis Biotypes Using Brain-Based
Biomarkers. Am J Psychiat 173, 373–384. https://doi.org/10.1176/appi.ajp.2015.14091200

Cuesta, S., Nouel, D., Reynolds, L.M., Morgunova, A., Torres-Berrío, A., White, A., Hernandez,
G., Cooper, H.M., Flores, C., 2020. Dopamine Axon Targeting in the Nucleus Accumbens in
Adolescence Requires Netrin-1. Frontiers Cell Dev Biology 8, 487.
https://doi.org/10.3389/fcell.2020.00487

Cuesta, S., Restrepo-Lozano, J.-M., Popescu, C., He, S., Reynolds, L.M., Israel, S., Hernandez, G.,
Rais, R., Slusher, B.S., Flores, C., 2019. DCC-related developmental effects of abused- versus
therapeutic-like amphetamine doses in adolescence. Addiction Biology 24, e12791-11.
https://doi.org/10.1111/adb.12791

Drzewiecki, C.M., Juraska, J.M., 2020. The structural reorganization of the prefrontal cortex
during adolescence as a framework for vulnerability to the environment. Pharmacol
Biochem Be 199, 173044. https://doi.org/10.1016/j.pbb.2020.173044

Fuhrmann, D., Knoll, L.J., Blakemore, S.-J., 2015. Adolescence as a Sensitive Period of Brain
Development. Trends in Cognitive Sciences 19, 558–566.
https://doi.org/10.1016/j.tics.2015.07.008

Gao, B., Cutler, M.G., 1993. Effects of quinpirole on the behaviour shown by mice in the light-
dark box and during social interactions. Neuropharmacology 32, 93–100.
https://doi.org/10.1016/0028-3908(93)90134-o

Gogtay, N., Giedd, J.N., Lusk, L., Hayashi, K.M., Greenstein, D., Vaituzis, A.C., Nugent, T.F.,
Herman, D.H., Clasen, L.S., Toga, A.W., Rapoport, J.L., Thompson, P.M., 2004. Dynamic
mapping of human cortical development during childhood through early adulthood.
Proceedings of the National Academy of Sciences 101, 8174–8179.
https://doi.org/10.1073/pnas.0402680101

Heldmaier, G., Steinlechner, S., 1981. Seasonal pattern and energetics of short daily torpor in
the Djungarian hamster, Phodopus sungorus. Oecologia 48, 265–270.
https://doi.org/10.1007/bf00347975

Heywood, S.G., Yellon, S.M., 1997. Gonadotropin-Releasing Hormone Neural Projections to the
Systemic Vasculature during Sexual Maturation and Delayed Puberty in the Male Djungarian
Hamster. Biol Reprod 57, 873–878. https://doi.org/10.1095/biolreprod57.4.873

Hoffmann, K., 1978. Effects of short photoperiods on puberty, growth and moult in the
Djungarian hamster (Phodopus sungorus). Journal of Reproduction and Fertility 54, 29–35.
https://doi.org/10.1530/jrf.0.0540029

Hollenstein, T., Lougheed, J.P., 2013. Beyond Storm and Stress. Am Psychol 68, 444–454.
https://doi.org/10.1037/a0033586

Hoops, D., Flores, C., 2017. Making Dopamine Connections in Adolescence. Trends in
Neurosciences 1–11. https://doi.org/10.1016/j.tins.2017.09.004

Hoops, D., Reynolds, L.M., Restrepo-Lozano, J.-M., Flores, C., 2018. Dopamine Development in
the Mouse Orbital Prefrontal Cortex Is Protracted and Sensitive to Amphetamine in
Adolescence. eNeuro 5, ENEURO.0372-17.2017-9. https://doi.org/10.1523/eneuro.0372-
17.2017

Huttenlocher, P.R., 1984. Synapse elimination and plasticity in developing human cerebral
cortex. American journal of mental deficiency 88, 488–496.

Juraska, J.M., Willing, J., 2017. Pubertal onset as a critical transition for neural development and
cognition. Brain Research 1654, 87–94. https://doi.org/10.1016/j.brainres.2016.04.012

Keshavan, M.S., Giedd, J., Lau, J.Y.F., Lewis, D.A., Paus, T., 2014. Changes in the adolescent brain
and the pathophysiology of psychotic disorders. Lancet Psychiatry 1, 549–558.
https://doi.org/10.1016/s2215-0366(14)00081-9

Kessler, R.C., Amminger, G.P., Aguilar-Gaxiola, S., Alonso, J., Lee, S., Üstün, T.B., 2007. Age of
onset of mental disorders: a review of recent literature. Curr Opin Psychiatr 20, 359–
364. https://doi.org/10.1097/yco.0b013e32816ebc8c

Kessler, R.C., Berglund, P., Demler, O., Jin, R., Merikangas, K.R., Walters, E.E., 2005. Lifetime
    Prevalence and Age-of-Onset Distributions of DSM-IV Disorders in the National Comorbidity
    Survey Replication. Arch Gen Psychiat 62, 593–602.
    https://doi.org/10.1001/archpsyc.62.6.593

Kim, S.R., Chen, X., Oo, T.F., Kareva, T., Yarygina, O., Wang, C., During, M., Kholodilov, N., Burke,
    R.E., 2011. Dopaminergic pathway reconstruction by Akt/Rheb-induced axon regeneration.
    Annals of Neurology 70, 110–120. https://doi.org/10.1002/ana.22383

Klune, C.B., Jin, B., DeNardo, L.A., 2021. Linking mPFC circuit maturation to the developmental
    regulation of emotional memory and cognitive flexibility. Elife 10, e64567.
    https://doi.org/10.7554/elife.64567

Kopec, A.M., Smith, C.J., Ayre, N.R., Sweat, S.C., Bilbo, S.D., 2018. Microglial dopamine receptor
    elimination defines sex-specific nucleus accumbens development and social behavior in
    adolescent rats. Nat Commun 9, 3769. https://doi.org/10.1038/s41467-018-06118-z

Koss, W.A., Belden, C.E., Hristov, A.D., Juraska, J.M., 2013. Dendritic remodeling in the
    adolescent medial prefrontal cortex and the basolateral amygdala of male and female rats.
    Synapse 68, 61–72. https://doi.org/10.1002/syn.21716

Larsen, B., Luna, B., 2018. Adolescence as a neurobiological critical period for the development
    of higher-order cognition. Neurosci Biobehav Rev 94, 179–195.
    https://doi.org/10.1016/j.neubiorev.2018.09.005

Lee, F.S., Heimer, H., Giedd, J.N., Lein, E.S., Sestan, N., Weinberger, D.R., Casey, B.J., 2014.
    Mental health. Adolescent mental health-opportunity and obligation. Science 346, 547–549.
    https://doi.org/10.1126/science.1260497

Luna, B., Marek, S., Larsen, B., Tervo-Clemmens, B., Chahal, R., 2015. An integrative model of
    the maturation of cognitive control. Annual review of neuroscience 38, 151–170.
    https://doi.org/10.1146/annurev-neuro-071714-034054

Lynn, D.A., Brown, G.R., 2009. The ontogeny of exploratory behavior in male and female
    adolescent rats (Rattus norvegicus). Dev Psychobiol 51, 513–520.
    https://doi.org/10.1002/dev.20386

Manitt, C., Eng, C., Pokinko, M., Ryan, R.T., Torres-Berrío, A., Lopez, J.P., Yogendran, S.V.,
    Daubaras, M.J.J., Grant, A., Schmidt, E.R.E., Tronche, F., Krimpenfort, P., Cooper, H.M.,
    Pasterkamp, R.J., Kolb, B., Turecki, G., Wong, T.P., Nestler, E.J., Giros, B., Flores, C., 2013. dcc
    orchestrates the development of the prefrontal cortex during adolescence and is altered in
    psychiatric patients. Translational Psychiatry 3, e338-13.
    https://doi.org/10.1038/tp.2013.105

Manitt, C., Labelle-Dumais, C., Eng, C., Grant, A., Mimee, A., Stroh, T., Flores, C., 2010. Peri-
Pubertal Emergence of UNC-5 Homologue Expression by Dopamine Neurons in Rodents.
PLoS ONE 5, e11463-14. https://doi.org/10.1371/journal.pone.0011463

Manitt, C., Mimee, A., Eng, C., Pokinko, M., Stroh, T., Cooper, H.M., Kolb, B., Flores, C., 2011.
The Netrin Receptor DCC Is Required in the Pubertal Organization of Mesocortical Dopamine
Circuitry. J Neurosci 31, 8381–8394. https://doi.org/10.1523/jneurosci.0606-11.2011

McTeague, L.M., Goodkind, M.S., Etkin, A., 2016. Transdiagnostic impairment of cognitive
control in mental illness. J Psychiatr Res 83, 37–46.
https://doi.org/10.1016/j.jpsychires.2016.08.001

Millan, M.J., Agid, Y., Brüne, M., Bullmore, E.T., Carter, C.S., Clayton, N.S., Connor, R., Davis, S.,
Deakin, B., DeRubeis, R.J., Dubois, B., Geyer, M.A., Goodwin, G.M., Gorwood, P., Jay, T.M.,
Joëls, M., Mansuy, I.M., Meyer-Lindenberg, A., Murphy, D., Rolls, E., Saletu, B., Spedding, M.,
Sweeney, J., Whittington, M., Young, L.J., 2012. Cognitive dysfunction in psychiatric
disorders: characteristics, causes and the quest for improved therapy. Nat Rev Drug Discov
11, 141–168. https://doi.org/10.1038/nrd3628

Paus, T., 2005. Mapping brain maturation and cognitive development during adolescence.
Trends in Cognitive Sciences 9, 60–68. https://doi.org/10.1016/j.tics.2004.12.008

Paxinos, G., Franklin, K.B.J., 2013. The Mouse Brain in Stereotaxic Coordinates, 4th ed,
Academic Press. Academic Press.

Petanjek, Z., Judas, M., Simic, G., Rasin, M.R., Uylings, H.B.M., Rakic, P., Kostovic, I., 2011.
Extraordinary neoteny of synaptic spines in the human prefrontal cortex. Proceedings of the
National Academy of Sciences 108, 13281–13286.
https://doi.org/10.1073/pnas.1105108108

Pevet, P., 1988. The role of the pineal gland in the photoperiodic control of reproduction in
different hamster species. Reproduction Nutrition Développement 28, 443–458.
https://doi.org/10.1051/rnd:19880310

Pietropaolo, S., Branchi, I., Cirulli, F., Chiarotti, F., Aloe, L., Alleva, E., 2004. Long-term effects of
the periadolescent environment on exploratory activity and aggressive behaviour in mice:
social versus physical enrichment. Physiol Behav 81, 443–453.
https://doi.org/10.1016/j.physbeh.2004.02.022

Pokinko, M., Moquin, L., Torres-Berrío, A., Gratton, A., Flores, C., 2015. Resilience to
amphetamine in mouse models of netrin-1 haploinsufficiency: role of mesocortical
dopamine. Psychopharmacology 232, 3719–3729. https://doi.org/10.1007/s00213-015-
4032-9

Reynolds, L.M., Flores, C., 2021. Mesocorticolimbic Dopamine Pathways Across Adolescence:
Diversity in Development. Front Neural Circuit 15, 735625.
https://doi.org/10.3389/fncir.2021.735625

Reynolds, L.M., Pokinko, M., Torres-Berrío, A., Cuesta, S., Lambert, L.C., Pellitero, E.D.C.,
Wodzinski, M., Manitt, C., Krimpenfort, P., Kolb, B., Flores, C., 2018a. DCC Receptors Drive
Prefrontal Cortex Maturation by Determining Dopamine Axon Targeting in Adolescence.
Biological psychiatry 83, 181–192. https://doi.org/10.1016/j.biopsych.2017.06.009

Reynolds, L.M., Yetnikoff, L., Pokinko, M., Wodzinski, M., Epelbaum, J.G., Lambert, L.C.,
Cossette, M.-P., Arvanitogiannis, A., Flores, C., 2018b. Early Adolescence is a Critical Period
for the Maturation of Inhibitory Behavior. Cerebral cortex 29, 3676–3686.
https://doi.org/10.1093/cercor/bhy247

Scherbarth, F., Steinlechner, S., 2010. Endocrine mechanisms of seasonal adaptation in small
mammals: from early results to present understanding. J Comp Physiology B 180, 935–952.
https://doi.org/10.1007/s00360-010-0498-2

Sowell, E.R., Thompson, P.M., Toga, A.W., 2004. Mapping Changes in the Human Cortex
throughout the Span of Life. Neurosci 10, 372–392.
https://doi.org/10.1177/1073858404263960

Spear, L.P., 2000. The adolescent brain and age-related behavioral manifestations. Neurosci
Biobehav Rev 24, 417–463.

Tanaka, T., 2015. Sex differences in exploratory behaviour of laboratory CD-1 mice (Mus
musculus). Scandinavian Journal of Laboratory Animal Science 41, 1–9.

Vassilev, P., Pantoja-Urban, A.H., Giroux, M., Nouel, D., Hernandez, G., Orsini, T., Flores, C.,
2021. Unique effects of social defeat stress in adolescent male mice on the Netrin-1/DCC
pathway, prefrontal cortex dopamine and cognition (Social stress in adolescent vs. adult
male mice). Eneuro ENEURO.0045-21.2021. https://doi.org/10.1523/eneuro.0045-21.2021

Westbrook, S.R., Hankosky, E.R., Dwyer, M.R., Gulley, J.M., 2018. Age and Sex Differences in
Behavioral Flexibility, Sensitivity to Reward Value, and Risky Decision-Making. Behav
Neurosci 132, 75–87. https://doi.org/10.1037/bne0000235

Wolf, R.M., Long, D., 2016. Pubertal Development. Pediatr Rev 37, 292–300.
https://doi.org/10.1542/pir.2015-0065

Yellon, S.M., Goldman, B.D., 1984. Photoperiod Control of Reproductive Development in the
Male Djungarian Hamster (Phodopus sungorus)* | Endocrinology | Oxford Academic.
academic.oup.com.

High resolution figures and captions

Figure 1. A "pathway" of Netrin-1 expressing cells "paves the way" for dopamine (DA) axons growing from the

nucleus accumbens to the medial prefrontal cortex during adolescence. **A**, The brain regions containing the bundle

of dopamine fibres passing to the medial prefrontal cortex are highlighted in a line drawing of a coronal mouse

brain section derived from Paxinos and Franklin (Paxinos and Franklin, 2013). **B**, An image of a coronal section

through the forebrain of an adult mouse at low magnification (4x). Green fluorescence indicates immunostaining

for tyrosine hydroxylase (TH), a marker for dopamine. The smaller and larger white squares indicate the regions

enlarged in panel C and panels F & G, respectively. Scale bar = 500 µm. **C**, The nucleus accumbens (left) is densely

packed with dopamine axons (in green). Some of these axons can be observed extending from the nucleus

accumbens medially into a bundle of dopamine fibres (white arrows). Scale bar = 10 µm. **D**, Modified stereological

quantification revealed no significant difference in dopamine axon density in the fibre bundle between

adolescence (21 days old) and adulthood (75 days old). Mixed-effects ANOVA, effect of age: F=1.53, p=0.22; region

by age interaction: F=1.44, p=0.49. **E**, The average width of the fibre bundle increased significantly from

adolescence to adulthood, revealing that there is an increase in the total number of fibres passing to the medial

prefrontal cortex during this period. Mixed-effects ANOVA, effect of age: F=9.45, p=0.0021; region by age

interaction: F=5.74, p=0.057.   **F**, In order to quantify the Netrin-1 positive cells along the dopamine fibre bundle,

the bundle was contoured in each region, and a contour of equal area was placed medial to the fibre bundle as a

negative control. Scale bar = 200 µm. **G**, Using quantitative stereology, Netrin-1 positive cell density was

determined along and adjacent to the fibre bundle for each region. Red fluorescence indicates immunostaining for

Netrin-1. **H**, In adolescent mice there are more Netrin-1 positive cells along the dopamine fibre bundle ("TH+")

than medial to it ("TH-"). This is what we refer to as the "Netrin-1 pathway". Along the pathway, there is a

significant increase in Netrin-1 positive cell density in regions closer to the medial prefrontal cortex, the

innervation target. Mixed-effects ANOVA, effect of TH: F=105, p<0.0001. Effect of region: F=9.74, p=0.021. A post-

hoc Tukey Test revealed a difference (p = 0.029) between the densities of the lateral septum and infralimbic

cortex, but only within the dopamine fibre bundle. **I**, In adult mice the Netrin-1 pathway is maintained, however

there is no longer an increasing density of Netrin-1 positive cells towards the medial prefrontal cortex. Mixed-

effects ANOVA, effect of TH: F=54.56, p<0.0001. Effect of region: F=1.22, p=0.75. **J,** The virus injection location

within the mouse brain. A Netrin-1 knockdown virus, or a control virus, was injected into the dopamine fibre

bundle at the level of the dorsal peduncular cortex. **K,** Our experimental timeline: at the onset of adolescence a

Netrin-1 knockdown virus, or a control virus, was injected, and then in adulthood the mice were sacrificed and

stereological measurements taken. **L,** Dopamine varicosity density was quantified in the region below the injection

site, in the lateral septum, and in the region above the injection site, in the infralimbic cortex. There was a

significant decrease in dopamine varicosity density only in the infralimbic cortex. Mixed-effects ANOVA, virus by

region interaction: F=16.41, p<0.0001. **M,** The experimental set-up of the final (test) stage of the Go/No-Go

experimental protocol. A mouse that has previously learned to nose-poke for a reward in response to a visual cue

(illuminated nose-poke hole), must now inhibit this behaviour when the visual cue is paired with an auditory cue

(acoustic tone). **N,** Mice injected with the Netrin-1 knockdown virus show improved action impulsivity than

controls because they incur significantly fewer commission errors across the Go/No-go task. Mixed-effects ANOVA,

effect of day: F=68.32, p<0.0001. Day by virus interaction: F=9.00, p=0.0027. A sigmoidal curve is fit to each group

of mice to determine how the two groups differ. Points indicate group means and error bars show standard error

means. **O**, During the first days of Go/No-Go testing, both groups incur commission errors with high frequency,

but, the Netrin-1 knockdown group has fewer errors than the control group (t-test, t=5.18, p<0.0001. **P,** The ED50

– the inflection point in each sigmoidal curve – are not different between groups, indicating that all mice improve

their ability to inhibit their behavior at around the same time (T-test, t=0.97, p=0.35). **Q,** Mice microinfused with

the Netrin-1 knockdown virus incur substantially fewer commission errors in the last days of the Go/No-Go task

compared to mice injected with the control virus (T-test, t=12.38, p<0.0001). For all barplots, bars indicate group

means and error bars show standard error means.

Figure 2 The age of onset of UNC5C expression by dopamine axons in the nucleus accumbens of mice is sexually

dimorphic. Images are representative of observed immunofluorescence patterns in the nucleus accumbens core

(approx. location highlighted in the coronal mouse brain section Plate 19, modified from Paxinos & Franklin, 2013).

For each row, six individuals were sampled. In males (**A-B**), UNC5C expression on dopamine fibres in the nucleus

accumbens appears during adolescence. **A,** At the onset of adolescence (21 days old) dopamine fibres do not

express UNC5C. Scale bar = 10 μm. **B,** By adulthood (90 days old), dopamine fibres express UNC5C. In females (**C-**

E), UNC5C expression on dopamine fibres in the nucleus accumbens appears prior to adolescence. **C,** In juvenile

(15 day old) mice, there is no UNC5C expression on dopamine fibres. **D,** By adolescence, dopamine fibres express

UNC5C. **E,** In adulthood, dopamine fibres continue to express UNC5C.

A

Anterior Cingulate Cortex

Prelimbic Cortex

Infralimbic Cortex

Nucleus Accumbens

B Summer-like daylength: long photoperiod

10 hrs / 14 hrs

Winter-like daylength: short photoperiod

10 hrs / 14 hrs

C Male Hamsters Summer Daylength

15 Days Old
80 Days Old
215 Days Old

Tukey test by age:
15-80do: p=0.026
80-215do: p<0.0001
15-215do: p<0.0001

TH+ Varicosity Density (x10⁻³cm⁻¹)

Anterior Cingulate Cortex   Prelimbic Cortex   Infralimbic Cortex

Effect of age: F=9.6, p=0.0003

D Male Hamsters Winter Daylength

15 Days Old
80 Days Old
215 Days Old

Tukey test by age:
15-80do: p=0.54
80-215do: p=0.0006
15-215do: p=0.0085

TH+ Varicosity Density (x10⁻³cm⁻¹)

Anterior Cingulate Cortex   Prelimbic Cortex   Infralimbic Cortex

Effect of age: F=4.2, p=0.02

Summer

15 days old

E DA UNC5c Merge — NAcc

80 days old

F DA UNC5c Merge — NAcc

215 days old

G DA UNC5c Merge — NAcc

Winter

H DA UNC5c Merge — NAcc

I DA UNC5c Merge — NAcc

J DA UNC5c Merge — NAcc

K Exploratory Behaviour – Males

Summer
Winter

Time in Light Box (min)

Peak: ~50 days old
Peak: >80 days old

Age (days)

2nd degree effect of season: F=4.91, p<0.0001

L Old Males

F=0.98, p=0.34

Figure 3 Plasticity of adolescent development in male Siberian hamsters according to seasonal phenotype. All

results illustrated in this figure refer to results in male hamsters. **A,** Dopamine innervation was quantified in three

subregions of the medial prefrontal cortex, highlighted in blue. UNC5C expression was examined in the nucleus

accumbens, highlighted in red. Line drawing of a coronal section of the mouse brain was derived from Plate 19 of

Paxinos and Franlkin (Paxinos and Franklin, 2013). **B,** Hamsters were housed under either summer-mimicking long

days and short nights ("summer hamsters") or winter-mimicking short days and long nights ("winter hamsters"). **C,**

In male hamsters housed under a summer-mimicking daylength there is an increase in dopamine varicosity density

in the medial prefrontal cortex between 15 and 80 days old. Mixed-effects ANOVA, effect of age: F=9.6,

p=0.000255. Tukey Test, 15-80 days old (do): p=0.026; 80-215do: p<0.0001; 15-215do: p<0.0001. **D,** In male

hamsters housed under a winter-mimicking daylength there is no increase in dopamine varicosity density until

hamsters have reached 215 days of age. Mixed-effects ANOVA, effect of age: F=4.17, p=0.0205. Tukey Test, 15-

80do: p=0.54; 80-215do: p=0.0006; 15-215do: p=0.0085. **E,** At 15 days old, dopamine axons in the nucleus

accumbens of male summer-daylength hamsters largely do not express UNC5C. Scale bar = 20um (bottom right). **F-**

G, At 80 (**F**) and 215 (**G**) days old, dopamine axons in the nucleus accumbens express UNC5C. **H-I,** At 15 (**H**) and 80

(**I**) days old, dopamine axons in the nucleus accumbens of male winter hamsters largely do not express UNC5C. **J,**

By 215 days old there is UNC5C expression in dopamine axons in the nucleus accumbens of male winter hamsters.

E-J, Representative images of the nucleus accumbens shell, 6 individuals were examined per group. **K,** Male

hamsters house under a summer-mimicking daylength show an adolescent peak in exploratory behaviour in the

light/dark box apparatus. Instead, those raised under a winter-mimicking photoperiod show a steady increase in

exploratory behaviour over the same age range. Arrows indicate the ages at which exploratory behaviour peaks in

summer (orange) and winter (blue) hamsters. Polynomial regression, effect of season: F=3.551, p=0.00056. **L,** In

male hamsters, at 215 days of age, there is no difference in exploratory behaviour between hamsters raised under

summer and winter photoperiods. T-test, effect of season: t=0.975, p=0.341. For all barplots, bars indicate group

means and error bars show standard error means.

Figure 4 Plasticity of adolescent development in female Siberian hamsters according to seasonal phenotype. All

results illustrated in this figure refer to results in female hamsters. **A,** Dopamine innervation was quantified in

three subregions of the medial prefrontal cortex, highlighted here in blue. UNC5C expression was examined in the

nucleus accumbens, highlighted in red. Line drawing of a coronal section of the mouse brain was derived from

Paxinos and Franlkin (Paxinos and Franklin, 2013). **B,** Hamsters were housed under either a summer-mimicking or

winter-mimicking daylength. **C,** In female hamsters housed under a summer daylength dopamine varicosity density

in the medial prefrontal cortex increases between 15 and 80 days of age. Mixed-effects ANOVA, effect of age:

F=16.72, p<0.0001 **D,** In female hamsters housed under a winter daylength there is no increase in dopamine

varicosity density post-adolescence. Instead, there is a steep decline in density between 80 and 215 days of age.

Mixed-effects ANOVA, effect of age: F=12.33, p=0.000043. **E,** As our results in panel D were unexpected, we

replicated them with a second cohort of hamsters and found qualitatively identical results. Mixed-effects ANOVA,

effect of age: F=34.871, p<0.0001. **F,** To try and determine when dopamine varicosities innervate the medial

prefrontal cortex, we examined a cohort of 10- and 15-day-old hamsters. We found that varicosity density

increases in the medial prefrontal cortex during this time, indicating that dopamine innervation to the medial

prefrontal cortex is accelerated in female winter hamsters. Mixed-effects ANOVA, effect of age: F=5.05, p=0.03. **G-**

H, In 10- and 15-day-old female summer hamsters there is little UNC5C expression in nucleus accumbens

dopamine axons. Sample size: 4 (panel G) or 6 (panel H). **I-J,** By 80 days old (panel I), and continuing at 215 days old

(panel J), dopamine axons in the nucleus accumbens express UNC5C in female summer hamsters. Sample sizes: 6.

Scale bar = 20um (panel G bottom right). **K-N,** At all ages at which winter female hamsters were examined

dopamine axons in the nucleus accumbens express UNC5C in winter female hamsters. Sample sizes: 4 (panel K) or

6 (panels L-N). **O,** In female hamsters, those raised under summer and winter daylengths both show an increase in

exploratory behaviour over time. The winter hamsters peak later compared to the summer daylength hamsters.

Arrows indicate the ages at which exploratory behaviour peaks in summer (orange) and winter (blue) hamsters.

Polynomial regression, effect of season: F=3.305, p=0.00126. **P,** In female hamsters, at 215 days of age, there is no

difference in exploratory behaviour between hamsters raised under summer and winter photoperiods. T-test,

effect of season: t=0.309, p=0.76. For all barplots, bars indicate group means and error bars show standard error

means.