1

Decomposing drivers in avian insectivory: large-scale effects

of climate, habitat and bird diversity

Laura Schillé[1,*], Elena Valdés-Correcher[1], Frédéric Archaux[2], Flavius Bălăcenoiu[3,4], Mona Chor Bjørn[5],

Michal Bogdziewicz[6], Thomas Boivin[7], Manuela Branco[8], Thomas Damestoy[9], Maarten de Groot[10],

Jovan Dobrosavljević[11], Mihai-Leonard Duduman[12], Anne-Maïmiti Dulaurent[9], Samantha Green[13], Jan

Grünwald[14], Csaba Béla Eötvös[15], Maria Faticov[16], Pilar Fernandez-Conradi[7], Elisabeth Flury[17], David

Funosas[18], Andrea Galmán[19,20], Martin M. Gossner[21,22], Sofia Gripenberg[23], Lucian Grosu[12], Jonas

Hagge[24], Arndt Hampe[1], Deborah Harvey[25], Rick Houston[26], Rita Isenmann[27], Andreja Kavčič[10], Mikhail

V. Kozlov[28], Vojtech Lanta[29], Bénédicte Le Tilly[30], Carlos Lopez Vaamonde[31,32], Soumen Mallick[33], Elina

Mäntylä[28,34,35], Anders Mårell[2], Slobodan Milanović[11], Márton Molnár[36], Xoaquín Moreira[37], Valentin

Moser[38,39], Anna Mrazova[34,40], Dmitrii L. Musolin[41], Thomas Perot[2], Andrea Piotti[42], Anna V. Popova[43],

Andreas Prinzing[33], Ludmila Pukinskaya[43], Aurélien Sallé[45], Katerina Sam[34,40], Nickolay V. Sedikhin[44,46],

Tanja Shabarova[47], Ayco J. M. Tack[48], Rebecca Thomas[25], Karthik Thrikkadeeri[49], Dragoș Toma[3,4], Grete

Vaicaityte[50], Inge van Halder[1], Zulema Varela[51], Luc Barbaro[52,53,¤] & Bastien Castagneyrol[1,¤]

[¤] Luc Barbaro and Bastien Castagneyrol share last co-authorship

[1] BIOGECO, INRAE, University Bordeaux, Cestas, France

[2] EFNO, INRAE, Nogent-sur-Vernisson, France

[3] National Institute for Research and Development in Forestry « Marin Drăcea », Voluntari, Romania

[4] Romania Faculty of Silviculture and Forest Engineering, Transilvania University of Brașov, Romania

[5] Department of Geosciences and Natural Resource Management, Section Forest and Landscape

Ecology, University of Copenhagen, Denmark

[6] Forest Biology Center, Faculty of Biology, Adam Mickiewicz University in Poznan, Poland

[7] UR629, INRAE, Centre PACA, Avignon, France

[8] Centro de Estudos Florestais, Instituto Superior de Agronomia, Universidade de Lisboa, Lisbon,

Portugal

[9] UniLaSalle, AGHYLE, UP.2018.C101, Beauvais, France

[10] Slovenian Forestry Institute, Ljubljana, Slovenia

[11] Department of Forestry and Nature Protection, University of Belgrade Faculty of Forestry,

Belgrade, Serbia

[12] Forestry Faculty, Applied Ecology Lab, "Ştefan cel Mare" University of Suceava, Romania

[13] Centre for Agroecology, Water and Resilience, Coventry University, UK

[14] Institute for Environmental Studies, Faculty of Science, Charles University in Prague, Czech Republic

[15] University of Sopron, Forest Research Institute, Department of Forest Protection, Mátrafüred,

Hungary

[16] Université de Sherbrooke, 2500, boul. de l'Université, Sherbrooke, Canada

[17] Kinderlab Landquart, Landquart, Switzerland

[18] Laboratoire Ecologie Fonctionnelle et Environnement, University of Toulouse, CNRS, France

[19] Martin-Luther-Universität Halle-Wittenberg, Halle (Saale), Germany

[20] German Centre for Integrative Biodiversity Research (iDiv), Leipzig, Germany

[21] Forest Entomology, Swiss Federal Research Institute WSL, Birmensdorf, Switzerland

[22] Institute of Terrestrial Ecosystems, Department of Environmental Systems Science, ETH Zürich,

Switzerland

[23] School of Biological Sciences, University of Reading, UK

[24] Forest Nature Conservation, Northwest German Forest Research Institute, Hann. Münden,

Germany

[25] Royal Holloway University of London, Egham, UK

[26] Beckley School Oxford, UK

[27] Grimmelshausenschule, Renchen, Germany

[28] Department of Biology, University of Turku, Finland

[29] Department of Functional Ecology, Institute of Botany CAS, Průhonice, Czech Republic

[30] Collège Saint Joseph, Paimpol, France

[31] INRAE Centre Val de Loire, Unité Zoologie Forestière, Orléans, France

[32] IRBI, UMR 7261, CNRS-Université de Tours, France

[33] Centre National de la Recherche Scientifique, Université de Rennes 1, Research Unit UMR 6553,

Ecosystèmes Biodiversité Evolution (ECOBIO), Campus de Beaulieu, 35042, Rennes, France

[34] Institute of Entomology, Biology Centre of the Czech Academy of Sciences, Ceske Budejovice, Czech

Republic

[35] Faculty of Science, Department of Zoology, University of South Bohemia, Ceske Budejovice, Czech

Republic

[36] Magyar Madártani és Természetvédelmi Egyesület (Birdlife Hungary), Szuha, Hungary

[37] Misión Biológica de Galicia (MBG-CSIC), Apdo. 28, 36080 Pontevedra, Galicia, Spain

[38] Community Ecology, Swiss Federal Research Institute WSL, Birmensdorf, Switzerland

[39] Aquatic Ecology, Swiss Federal Institute of Aquatic Science and Technology Eawag, Dübendorf,

Switzerland

[40] University of South Bohemia, Ceske Budejovice, Czech Republic

[41] European and Mediterranean Plant Protection Organization, Paris, France

[42] Institute of Biosciences and BioResources, National Research Council of Italy, Sesto Fiorentino, Italy

[43] A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Moscow, Russia

[44] Saint Petersburg State Forest Technical University, Lisinsky Forest College, Saint Petersburg, Russia

[45] LBLGC, Université d'Orléans, France

[46] Zoological institute of Russian Academy of Science, Saint Petersburg, Russia

[47] Biology Centre CAS, Institute of Hydrobiology, Ceske Budejovice, Czech Republic

[48] Department of Ecology, Environment and Plant Sciences, Stockholm University, Sweden

[49] Nature Conservation Foundation, Bengaluru, India

[50] Lithuanian Centre of Non-Formal Youth Education, Vilnius, Lithuania

[51] CRETUS,Ecology Unit, Department of Functional Biology, Universidade de Santiago de Compostela,

Spain

[52] DYNAFOR, University of Toulouse, INRAE, France

[53] LTSER ZA Pyrenees Garonne, Auzeville-Tolosane, France

* Correspondance: *laura.schille@hotmail.fr*

Acknowledgements

This study was permitted by the financial support of the BNP Paribas Foundation through its Climate

& Biodiversity Initiative for the 'Tree bodyguards' citizen science project. FB and DT were supported

by the PN 23090102 and 34PFE./30.12.2021 'Increasing the institutional capacity and performance of

INCDS "Marin Drăcea" in the activity of RDI - CresPerfInst' funded by the Ministry of Research,

Innovation and Digitalization of Romania. MB was supported by the Forest Research Centre (CEF)

(UIDB/00239/2020) and the Laboratory for Sustainable Land Use and Ecosystem Services—TERRA

(LA/P/0092/2020) funded by FCT, Portugal. MdG and AK were supported by the core research group

"Forest biology, ecology and technology" (P4-0107) of the Slovenian Research Agency. MVK was

supported by the Academy of Finland (project 316182). VL was funded by the Czech Science

Foundation (project 23-07533S) and Academy of Sciences (RVO 67985939). EM and KS were supported

by the Grant Agency of the Czech Republic (19-28126X). KS and AM were supported by ERC StG BABE

805189. XM was supported by a grant from the Spanish National Research Council (2021AEP082) and

a grant from the Regional Government of Galicia (IN607A 2021/03). ZV was supported by a grant

awarded by the Autonomous Government of Galicia (Spain; Modalidade B-2019), and Maria Zambrano

program from the Spanish Ministry of Universities. LB was supported by fundings from LTSER ZA

Pyrenees Garonne. We thank the personnel of Centro Nazionale Carabinieri Biodiversità "Bosco

Fontana" for help in data collection. We thank Tarja Heinonen for help in field work in Berlin, Germany.

We thank Roman Hrdlička and Kari Mäntylä for bird voice identification. We thank E. L. Zvereva for

help in data collection.

Abstract

Aim

Climate is a major driver of large scale variability in biodiversity, as a likely result of more intense biotic

interactions under warmer conditions. This idea fuelled decades of research on plant-herbivore

interactions, but much less is known about higher-level trophic interactions. We addressed this

research gap by characterizing both bird diversity and avian predation along a climatic gradient at the

European scale.

Location

Europe.

Taxon

Insectivorous birds and pedunculate oaks.

Methods

We deployed plasticine caterpillars in 138 oak trees in 47 sites along a 19° latitudinal gradient in Europe

to quantify bird insectivory through predation attempts. In addition, we used passive acoustic

monitoring to (i) characterize the acoustic diversity of surrounding soundscapes; and (ii) infer both

taxonomic and functional diversity of insectivorous birds from recordings.

Results

The functional diversity of insectivorous birds increased with warmer climates. Bird predation

increased with forest cover and bird acoustic diversity but decreased with mean annual temperature

and functional richness of insectivorous birds. Contrary to our predictions, climatic clines in bird

predation attempts were not directly mediated by changes in insectivorous bird diversity or acoustic

diversity, but climate and habitat still had independent effects on predation attempts.

Main conclusions

Our study supports the hypothesis of an increase in the diversity of insectivorous birds towards warmer

climates, but refutes the idea that an increase in diversity would lead to more predation and advocates

for better accounting for activity and abundance of insectivorous birds when studying the large-scale

variation in insect-tree interactions.

Keywords: Acoustic diversity, Climatic gradient, Functional diversity, Insectivorous birds, Plasticine

caterpillars, Predation function

Résumé

Objectif

Le climat est l'un des principaux moteur de la variabilité à grande échelle de la biodiversité,

probablement en raison d'interactions biotiques plus intenses dans des conditions de temperature

plus élevées. Cette idée a alimenté des décennies de recherche sur les interactions plantes-herbivores,

mais on en sait beaucoup moins sur les interactions trophiques de niveau supérieur. Nous avons

comblé cette lacune en caractérisant à la fois la diversité des oiseaux et leur activité de prédation le

long d'un gradient climatique à l'échelle européenne.

Localisation

Europe.

Taxon

Oiseaux insectivores et chênes pédonculés.

Méthodes

Nous avons déployé des chenilles en pâte à modeler sur 138 chênes dans 47 sites le long d'un gradient

latitudinal de 19° en Europe pour quantifier l'insectivorie avienne par le biais de tentatives de

prédation. De plus, nous avons utilisé la surveillance acoustique passive pour (i) caractériser la diversité

acoustique des paysages sonores environnants ; et (ii) déduire à la fois la diversité taxonomique et

fonctionnelle des oiseaux insectivores à partir des enregistrements.

Résultats

La diversité fonctionnelle des oiseaux insectivores a augmenté sous des climats plus chauds. La

prédation avienne a augmenté avec la couverture forestière et la diversité acoustique des oiseaux,

mais a diminué avec la température annuelle moyenne et la richesse fonctionnelle des oiseaux

insectivores. Contrairement à nos prédictions, les clines climatiques dans les tentatives de prédation

avienne n'ont pas directement été médiées par des changements de diversité des oiseaux insectivores

ou de diversité acoustique, mais le climat et l'habitat ont tout de même eu des effets indépendants

sur les tentatives de prédation.

Conclusions principales

Notre étude confirme l'hypothèse d'une augmentation de la diversité des oiseaux insectivores vers des

climats plus chauds, mais réfute l'idée qu'une augmentation de la diversité conduirait à davantage de

predation. Elle plaide en faveur d'une meilleure prise en compte de l'activité et de l'abondance des

oiseaux insectivores lors de l'étude de la variation à grande échelle des interactions entre insectes et

arbres.

Mots-clés : Diversité acoustique, Gradient climatique, Diversité fonctionnelle, Oiseaux insectivores,

Chenilles en pâte à modeler, Fonction de prédation.

Introduction

Climate is a key driver of biotic interactions (Dobzhansky, 1950). A long held view in ecology posits that

warmer and more stable climatic conditions intensify biotic interactions and accelerates speciation

(MacArthur, 1984; Schemske, Mittelbach, Cornell, Sobel & Roy, 2009), which should result in large

scale positive correlations between biodiversity and biotic interactions. However appealing this idea

is, the generality of large scale climatic clines in biodiversity and biotic interactions as well as the

underlying causal links are still widely debated. Yet, insights into the controversy have been dominated

by studies on plant-insect interactions (Anstett, Chen & Johnson, 2016; Kozlov, Lanta, Zverev &

Zvereva, 2015). Biotic interactions involving higher trophic levels received much less attention. Yet,

insectivorous birds are among the predators contributing the most to the control of insect herbivores

in terrestrial ecosystems (van Bael et al., 2008; Sekercioglu, 2006) and therefore have consequences

on both the assembly of ecological communities and the functioning of ecosystems. The omission of

predation in theories linking large scale variability in climate with biodiversity therefore represents a

critical gap in knowledge that needs to be addressed.

Bird communities are highly responsive to climate, at both regional and continental scale. There is a

large body of literature demonstrating that several dimensions of bird diversity vary with climate,

including bird species richness, phylogenetic or functional diversity (Blackburn & Gaston, 1996; Willig,

Kaufman & Stevens, 2003). A well substantiated explanation is that niche opportunities increase with

increasing habitat heterogeneity under milder climatic conditions, which increases species coexistence

and ultimately species richness through functional complementarity (Hawkins, Diniz-Filho, Jaramillo &

Soeller, 2006). Yet, the biodiversity and ecosystem relationship theory predicts that the intensity of

avian predation generally increases with bird diversity (Bael et al., 2008; Otto, Berlow, Rank, Smiley &

Brose, 2008; Sinclair, Mduma & Brashares, 2003). Numerous studies support this theory and

demonstrate that bird functional diversity in particular --- that is the diversity, distribution and

complementarity of predator traits involved in predation --- is a good predictor of predation (Barbaro,

Giffard, Charbonnier, van Halder & Brockerhoff, 2014; Greenop, Woodcock, Wilby, Cook & Pywell,

2018; Philpott et al., 2009). It follows that variation in bird diversity along climatic gradients should be

mirrored by consistent variation in avian predation rates.

Local factors can however alter macroecological patterns (Ikin et al., 2014; Kissling, Sekercioglu & Jetz,

2012), by filtering the regional species pool (De la Mora, García-Ballinas & Philpott, 2015; Kleijn,

Rundlöf, Scheper, Smith & Tscharntke, 2011) and by influencing the behavior of organisms. Birds are a

taxon whose structural diversity heavily depends on local factors that provide niches and food

opportunities (Charbonnier et al., 2016). In this respect, multiscale forest cover proved to be a

particularly good predictor of avian biodiversity at different spatial scales, as bird foraging activity is

ultimately determined by vertical and horizontal habitat heterogeneity, which influences both where

prey can be found and caught, and where foraging birds can breed and hide from predators (Vickery

& Arlettaz, 2012). Thus, modeling the response of bird communities to large-scale bioclimatic drivers

as well as their role as predators would benefit from using a combination of habitat variables and biotic

predictors (Barbaro et al., 2019; Speakman et al., 2000). However, cross-continental studies exploring

the relationship between large scale climatic gradients and the strength of biotic interactions generally

ignore local factors, which may partly explain inconsistencies in their findings (but see Just, Dale, Long

& Frank, 2019).

A major challenge to analyze climatic clines in biotic interactions consists in simultaneously

characterizing changes in predator biodiversity and experimentally assessing the strength of predation,

while considering the effect of contrasting habitats. However, the recent development of passive

acoustic monitoring provides a standardized, low-cost and non-invasive approach for ecological

studies and biodiversity monitoring (Gibb, Browning, Glover-Kapfer, Jones & Börger, 2019). Acoustic

monitoring of a given habitat firstly allows the identification of bird species. Secondly, through the

calculation of acoustic diversity indices, the activity of the community as a whole can be characterised,

while also accounting for the importance of human disturbances (Gasc et al., 2013; Sueur, Farina, Gasc,

Pieretti & Pavoine, 2014). Should such indices consistently correlate with macro-scales biotic

interactions, ecoacoustics would be a promising complementary approach to existing methods in

macroecology and in functional ecology.

Here, we addressed the hypothesis of continental north-south clines on insectivorous bird community

diversity and their predation function , while controlling for local factors throughout the European

distribution range of the pedunculate oak (*Quercus robur* L., 1753), a major forest tree species.

Specifically, we predict the following (Fig. 1): (i) bird diversity (including bird acoustic diversity,

insectivorous bird species richness and functional diversity) and predation attempts increase with

warmer climates; (ii) bird predation attempts increase with bird acoustic activity, species richness and

greater functional diversity of insectivorous birds; (iii) bird diversity, acoustic activity and bird

predation attempts increase with increasing forest cover at both local (neighborhood) and larger

spatial scales; (iv) large-scale variability in bird predation attempts is driven by local changes in the

diversity and activity of birds. To test these predictions, we quantified bird predation attempts on

plasticine caterpillars and estimated bird species richness, functional diversity and activity through

simultaneous passive acoustic monitoring. We eventually tested the respective responses of these

variables and their relationships at the pan-European scale.

Figure 1: *Conceptual diagram of the predictions of this study and the relationships already established in the literature. Boxed*

elements written in bold correspond to the main categories of variables tested, they are not variables as such. Variables used

in models are shown in regular font. Where several variables described the same category (e.g. ACI, NDSI, BI, ADI, H, NPIC,

MultiAcou, all describing acoustic indices), we used multi-model comparisons to identify the best variable. Items framed in

black on a white background represent untested variables. Black arrows indicate relationships well supported by the literature

(see Fig.2 Sánchez-Giraldo, Correa Ayram & Daza, 2021). Our specific predictions are represented with grey arrows, solid and

dashed lines representing positive and negative (predicted) relationships. Numbers refer to predictions as stated in the main

text.

Materials and methods

Study area

We focused on the pedunculate oak, *Quercus robur,* which is one of the keystone deciduous tree

species in temperate European forests, where it is of high ecological, economic and symbolic

importance (Eaton, Caudullo, Oliveira & de Rigo, 2016). The species occurs from central Spain (39°N)

to southern Fennoscandia (62°N) and thus experiences a huge gradient of climatic conditions (Petit et

al., 2002). A widely diverse community of specialist and generalist herbivorous insects is associated

with this species throughout its distributional range (Southwood, Wint, Kennedy & Greenwood, 2005).

Between May and July 2021, we studied 138 trees in 47 sites across 17 European countries covering

most of the pedunculate oak geographic range (Fig. 2). The sites were chosen with the minimal

constraint of being located in a wooded area of at least 1 ha (Valdés-Correcher et al., 2021). We

randomly selected three mature oaks per site, with the exception of six sites (three sites with one tree,

one site with two trees and two sites with five trees, see Table S1.1 in Appendix S1 in Supporting

Information).

Figure 2: Locations of the 47 sites sampled in spring 2021. Average annual temperature (color scale) according to WorldClim

(Hijmans, Cameron, Parra, Jones & Jarvis, 2005) and Quercus robur distribution range are indicated.

Bird predation attempts

We measured bird predation attempts in the field by exposing a total of 40 plasticine caterpillars (20

plasticine caterpillars twice) on each individual oak. We made plasticine caterpillars of green plasticine,

mimicking common lepidopteran larvae (3 cm long, 0.5 cm diameter, see Low, Sam, McArthur, Posa &

Hochuli, 2014). We secured them on twigs with a 0.3 mm metallic wire. We attached five plasticine

caterpillars to each of four branches facing opposite directions (i.e., 20 caterpillars per tree) at about

2 m from the ground.

We installed the plasticine caterpillars six weeks after budburst in each study area, thus synchronizing

the study with local oak phenology. We removed the plasticine caterpillars after 15 days and installed

another set of 20 artificial caterpillars per tree for another 15 days. At the end of each exposure period

(which varied from 10 to 20 (mean ± SD: 14.5 ± 1.23) days due to weather conditions, we carefully

removed the plasticine caterpillars from branches, placed them into plastic vials and shipped them to

the project coordinator. Plasticine caterpillars from six sites were either lost or altered during shipping,

preventing the extraction of relevant data.

A single trained observer (EVC) screened the surface of plasticine caterpillars with a magnifying lens to

search for the presence of bill marks on clay surface (Low et al., 2014). As we were ultimately interested

in linking bird diversity with bird predation attempts, we did not consider marks left by arthropods and

mammals.

We defined *bird predation attempts index* as p / d, where p is the proportion of plasticine caterpillars

with at least one sign of attempted predation by birds and d is the number of days plasticine caterpillars

were exposed to predators in the field. We only considered as attacked those caterpillars that we

retrieved; missing caterpillars were not accounted for in the calculation of p. We calculated bird

predation attempts for each tree and survey period separately. Because other variables were defined

at site level (see below), we averaged bird predation attempts across trees and surveys in each site

(total: $n = 41$).

To assess the effect of temperature independently of other variables that could vary with latitude, we

also calculated a second bird predation attempts index by standardizing the predation attempts by

daylight duration in every site (see Fig. S2.1 in Appendix S2 for the calculation of this index). We ran

the statistical models in exactly the same way as for the non-standardized bird predation attempts and

the results are presented in Table S3.2 in Appendix S3.

Acoustic monitoring and related variables

We used passive acoustic monitoring to characterize both the overall acoustic diversity (i.e., the

soundscape surrounding target oaks) and the bird species richness and functional diversity associated

with oaks, targeting the canopy insectivores that are expected to play a functional predation role on

oak herbivores (Fig. 2). In each site, we randomly chose one oak among those used to measure bird

predation rates in which we installed an AudioMoth device (Hill et al., 2018) to record audible sounds

for 30 min every hour. Automated recording started the day we installed the first set of 20 plasticine

caterpillars in trees and lasted until batteries stopped providing enough energy. The recording settings

were the following: Recording period: *00.00-24.00 (UTC); Sample rate: 48 kHZ; Gain: Medium; Sleep*

duration: 1800 s, Recording duration: 1800 s.

In all 47 sites, Audiomoths were active on average (± S.D.) for 9 ± 3 days (range: 1-24), which

corresponded to 5920 h of recordings in total and from 70 to 335 (246 ± 65) 30 min continuous acoustic

samples per site. When Audiomoths ran out of battery, the recordings lasted less than 30 min

(between 1 and 56 recordings per site were affected).

We processed acoustic samples with functions in the "soundecology" v.1.3.3 (Villanueva-Rivera &

Pijanowski, 2018) and "seewave" v. 2.1.8 (Sueur, Aubin & Simonis, 2008) libraries in the R environment

version 4.1.2 (R Core Team, 2020), and a wrap-up function made available by A. Gasc in GitHub

(https://github.com/agasc/Soundscape-analysis-with-R). We first divided every acoustic sample

(regardless of its length) into non-overlapping 1 min samples. Then, we calculated the following six

acoustic diversity indices for each 1 min sample: the Acoustic Complexity Index (ACI), the Acoustic

Diversity Index (ADI), the Bioacoustic Index (BI), the total acoustic entropy (H), the Normalized

Difference Sound Index (NDSI), and the Number of frequency Peaks NPIC (Bradfer-Lawrence et al.,

2019; Sueur et al., 2014). We further aggregated these indices into a multiacoustic index (MultiAcou)

as proposed by Barbaro et al. (2022). We calculated the median of each acoustic index per day and

then averaged median values across days for each site separately. We proceeded like this because 24

h cycles summarize the acoustic activity and account for all possible sounds of a given day.

Furthermore, other studies have previously shown that median values of acoustic indices for a given

day are more representative than mean values of the acoustic activity because they are less sensitive

to extreme values (Barbaro et al., 2022; Dröge et al., 2021). This procedure resulted in one single value

of each acoustic diversity index per site.

Altogether, acoustic indices capture various dimensions of the soundscape such as: anthropophony,

technophony, bird activity, abundance of species richness (for more details, see Fig.1 and Sánchez-

Giraldo, Correa Ayram & Daza, 2021, Fig.2). They are not necessarily expected to reflect any particular

bird-related variable, but should rather be seen as an integrative component of the habitats in which

the experiment was conducted. How strongly these indices reflect peculiar components of biodiversity

or ecosystem properties is still unknown. We therefore opted for an open approach and considered

them all in further analyses, but used multi-model statistical inferences to identify the indices that

were the most strongly linked with the response variables of interest (see below).

Bird species richness and functional diversity

We used acoustic samples to identify birds based on their vocalizations (songs and calls) at the species

level, from which we further computed functional diversity indices (Fig. 3).

Data processing – For each site, we subsampled the 30 min samples corresponding to the songbird

morning chorus (*i.e.,* the period of maximum singing activity), which incidentally also corresponds to

the time of the day when anthropic sounds were of the lowest intensity. Specifically, we selected

sounds recorded within a period running from 30 min before sunrise to 3 h 30 min after sunrise. We

then split each 30 min sample into up to three 10 min sequences, from which we only retained those

recorded on Tuesday, Thursday, Saturday, and Sunday. We chose these days on purpose to balance

the differences in anthropogenic noises between working days and weekends. For each sound sample,

we displayed the corresponding spectrogram with the "seewave" library in the R environment (Sueur,

Aubin & Simonis, 2008). We visually sorted sound samples thanks to spectrograms and discarded

samples with noise from anthropogenic sources, rain, or wind, which can be recognized as very low

frequency noise on the spectrogram. We also discarded samples with noise of very high frequency

corresponding to cicada chirps. We then randomly selected one sound sample per site and per day,

with the exception of four sites for which the four samples only covered two to three days. In total, we

selected 188 samples of 10 min (i.e., 4 samples per site).

Figure 3: Methodological pathway used to identify bird species (in light green) and calculate acoustic indices (in dark green)

from automated recordings (see text for details)

Bird species identification – We distributed the samples among 21 expert ornithologists. Each expert performed aural bird species identifications from 4 (one site) to 52 samples (13 sites), primarily from her/his region of residence, for auditory acoustic detection of bird species. We established a presence/absence Site × Species matrix, from which we calculated species richness and functional diversity. It is important to note that, unlike typical point counting that is commonly done by ornithologists in the field, there is no possibility to determine the direction and distance at which birds are singing from audio recordings when using a single device for a given site. As a result, there is no standard method for determining whether or not two vocalizations of the same species at two different times come from one single individual or more, which prevents an accurate estimate of bird abundance. However, experienced ornithologists involved in this study consider that, given the territoriality of birds and the range of the recorders, it is unlikely that they recorded the vocalizations of several individuals of the same species. It therefore seems reasonable to assume that among-site differences in bird species richness were also representative of among-site differences in bird abundance.

Functional diversity – We defined 25 bird species as candidate insectivores for attacking plasticine caterpillars (Table S4.3 in Appendix S4) with those bird species meeting the following criteria: be insectivorous during the breeding season or likely to feed their offspring with insects, forage primarily in forested habitats, and are likely to use substrates such as lower branches or lower leaves of trees where caterpillars were attached to find their prey (Barbaro et al., 2021; Brambilla & Gatti, 2022). We calculated the functional diversity of these candidate insectivores by combining morphological, reproductive, behavioral and acoustic traits.

With the exception of acoustic traits, we extracted functional traits from different published sources, listed in Table S4.4 in Appendix S4. Specifically, we used three continuous traits: *body mass*, *mean clutch size* and *bill culmen length* (see Fig. 2 in Tobias et al., 2022) combined with four categorical traits:

foraging method (predominantly understory gleaner, ground gleaner, canopy gleaner), *diet*

(insectivores or mixed diet), *nest type* (open in shrub, open on ground, cavity or open in tree) and

migration (short migration, long migration or resident).

We derived acoustic traits calculations from the work of Krishnan & Tamma (2016). We first extracted

five pure recordings without sonic background for each of the 25 candidate insectivore species from

the online database Xeno-canto.org (Vellinga & Planque, 2015). We then calculated the *number of*

peaks (i.e., NPIC) in the audio signal (see § Acoustic diversity, above) as well as the *frequency of the*

maximum amplitude peaks for each vocal element using the "seewave" library (Sueur, Aubin &

Simonis, 2008) and averaged these frequencies for each species. Being based on song and call

frequency and complexity, these indices inform the adaptation of the vocal repertoire of these species

to their environment.

We summarized the information conveyed by the 9 traits categories into five indices representing

complementary dimensions of the functional diversity (FD) of a community (Mouillot, Graham,

Villéger, Mason & Bellwood, 2013): functional richness (FRic, i.e., convex hull volume of the functional

trait space summarized by a principal coordinates analysis), functional evenness (FEve, i.e., minimum

spanning tree measuring the regularity of trait abundance distribution within the functional space),

and functional divergence (FDiv, i.e., trait abundance distribution within the functional trait space

volume) (Villéger, Mason & Mouillot, 2008), as well as Rao's quadratic entropy (RaoQ, i.e., species

dispersion from the functional centroïd) (Botta-Dukát, 2005). These were calculated for each site with

the "dbFD" function of the "FD" library v.1.0.12 (Laliberté, Legendre & Shipley, 2014) in the R

environment.

Environmental data

Environmental data refer to local temperature and forest cover. We used the high 10-m resolution GIS

layers from the Copernicus open platform (Cover, 2018) to calculate forest cover for all European sites.

We manually calculated the percentage of forest cover for the two sites located outside Europe using

the "World imagery" layer of Arcgis ver. 10.2.3552. We calculated both the percentage of forest cover

in a 20-m (henceforth called *local* forest cover) and 200-m (*landscape* forest cover) buffer around the

sampled oaks. We chose two nested buffer sizes to better capture the complexity of habitat structure

on the diversity and acoustic activity of birds. Local forest cover is particularly important for estimating

bird occurrence probability (Melles, Glenn & Martin, 2003), whereas landscape forest cover is an

important predictor of bird community composition in urban areas (Rega-Brodsky & Nilon, 2017).

Moreover, both local and landscape habitat factors shape insect prey distribution (Barr, van Dijk,

Hylander & Tack, 2021). Preliminary analyses revealed that results were qualitatively the same using

10-, 20- or 50-m buffers as predictors of local forest cover and 200- or 500-m buffers as predictors of

landscape forest cover (see Table S5.5 in Appendix S5). Because other variables were defined at the

site level, we averaged the percentage of forest cover for the sampled trees per site and per buffer

size.

We extracted the mean annual temperature at each site from the WorldClim database (the spatial

resolution is ~86 km^2, Hijmans et al., 2005).

Statistical analyses

We analyzed 14 response variables in separate linear models (LMs) (Table S3.2 in Appendix S3): bird

predation attempts, species richness of the entire bird community and that of candidate insectivores,

functional diversity (each of the 4 indices) and acoustic diversity (each of the 7 indices). For each

response variable, we first built a full model including variables reflecting two components of the

environment: climate and local habitat. The general model equation was (Eq. 1):

$Y_i = \beta_0 + \beta_1 \times Forest_{20,\,i} + \beta_2 \times Forest_{200,\,i} + \beta_3 \times Climate_{\,i} + \varepsilon_i$ (1)

where *Y* is the response variable, β_0 the model intercept, β_{is} model coefficient parameters, *Forest$_{20}$* and

Forest$_{200}$ the effects of the local and landscape forest cover respectively, *Climate* the effect of mean

annual temperature and ε the residuals.

When modeling the response of bird predation attempts (Eq. 2), we added two more variables to the

model, being any of the seven acoustic diversity indices (*Acoustic diversity*, Eq. 2) and the species

richness or any of the four indices describing the functional diversity of candidate insectivores (*Bird*

diversity, Eq. 2):

$Y_i = \beta_0 + \beta_1 \times Forest_{20,i} + \beta_2 \times Forest_{200,i} + \beta_3 \times Climate_i +$

$\beta_4 \times Bird\ diversity_i + \beta_5 \times Acoustic\ diversity_i + \varepsilon_i$ (2)

It has to be noted that the including the acoustic component in the second set of models does not

imply any direct link between avian predation and acoustic diversity. By comparing models including

the acoustic diversity or not, we are asking whether residual variance can be explained by this

component while controlling for other sources of variation. If so, then acoustic diversity components

with non-null coefficients have to be considered as proxies of predation, i.e., relatively easily

measurable variables representative of unmeasured (or unknown) variables with a direct effect on

predation.

We used logarithmic transformations (for bird predation attempts and acoustic entropy (H) models)

or square rooted transformation (for species richness of the complete bird community) of some

response variables where appropriate to satisfy model assumptions. We scaled and centered every

continuous predictor prior to modeling to facilitate comparisons of their effect sizes, and made sure

that none of the explanatory variables were strongly correlated using the variance inflation factor (VIF)

(all VIFs < 5, the usual cutoff values used to check for multicollinearity issues (Miles, 2014)).

For each response variable, we ran the full model as well as every model nested within the full model

and then used Akaike's Information Criterion corrected for small sample size (AICc) to identify the most

effective model(s) fitting the data the best. We simultaneously selected the best variable describing

the diversity and acoustic component (variable selection) and the best set of variables describing the

variability of the response variable (model selection).

First, we ranked each model according to the difference in AICc between the given model and the

model with the lowest AICc (ΔAICc). Models within 2 ΔAICc units of the best model (i.e., the model

with the lowest AICc) are generally considered as likely (Burnham & Anderson, 2002). We computed

AICc weights for each model (w_i). w_i is interpreted as the probability of a given model being the best

model among the set of candidate models. Eventually, we calculated the relative variable importance

(RVI) as the sum of w_i of every model including this variable, which corresponds to the probability a

variable is included in the best model.

When several models competed with the best model (i.e., when multiple models were such that their

ΔAICc < 2), we applied a procedure of multimodel inference, building a consensus model including the

variables in the set of best models. We then averaged their effect sizes across all the models in the set

of best models, using the variable weight as a weighting parameter (i.e., model averaging). We

considered that a given predictor had a statistically significant effect on the response variable when its

confidence interval excluded zero.

We run all analyses in the R language environment (R Core Team, 2020) with libraries "MuMIn"

v.1.43.17 (Bartoń, 2020), "lme4" v. 1.1.27.1 (Bates, Mächler, Bolker & Walker, 2015).

Results

Bird acoustic diversity

Of the seven acoustic diversity indices (see Fig. S6.2 in Appendix S6 for correlation between indices), only Acoustic Diversity Index (ADI) and acoustic entropy (H) were significantly associated with any of the predictors tested, i.e., temperature, local forest cover and landscape forest cover (Table S3.2 in Appendix S3). ADI and H both increased with local forest cover (i.e., percentage of forest cover in a 20-m buffer around recorders). Landscape-scale forest cover (i.e., percentage of forest cover in a 200-m buffer around recorders) was the only other predictor retained in the set of competing models in a range of ΔAICc < 2 to explain acoustic entropy variation, but this predictor had little importance (RVI < 0.5) and its effect was not statistically significant (Fig. 5b; Table S3.2 in Appendix S3).

Bird species richness and functional diversity

We identified a total of 87 bird species, among which 25 were classified as candidate functional insectivores. Bird species richness varied from 8 to 23 species per recording site (mean ± SD: 15.2 ± 3.7, n = 47 sites) and richness of candidate insectivores from 2 to 9 species (5.7 ± 1.5). The null model was among models competing in a range of ΔAICc < 2 for both total species richness and candidate insectivores (Table S3.2 in Appendix S3).

Among the five bird functional diversity and species richness indices, only functional quadratic entropy (Rao's Q) characterizing species dispersion from the functional centroid was significantly influenced by the predictors tested (temperature, local and landscape forest cover, Table S3.2 in Appendix S3). Specifically, Rao's Q increased with increasing temperature (Fig. 4a and Fig. 5c). Other predictors retained in the set of competing models in a range of ΔAICc < 2 had little importance (RVI < 0.5) and were not significant (Fig. 5c; Table S3.2 in Appendix S3).

Figure 4: Scatter diagrams showing changes in (a) Rao's quadratic entropy (Rao's Q) and (b) predation attempts with mean annual temperature. These relationships were identified as significant in the linear models tested. A dot represents a site, the prediction line corresponds to a linear regression between the two variables and the gray bands represent the confidence intervals around this regression.

Figure 5: Effects of climate (described by the mean annual temperature) and habitat (percentage of forest cover at 20 or 200 m) on Acoustic Diversity Index (ADI) (a), Acoustic Entropy Index (H) (b), Rao's quadratic entropy (RaoQ) (c), bird predation attempts (d) and effects of acoustic (Bioacoustic Index), bird diversity (Functional Richness) on bird predation attempts (d). Circles and error bars represent standardized parameter estimates and corresponding 95% confidence intervals (CI), respectively. The vertical dashed line centered on zero represents the null hypothesis. Full and empty circles represent significant and non-significant effect sizes, respectively. Circle size is proportional to RVI. The colors correspond to the different components tested (acoustic, bird diversity, geography and habitat).

Bird predation attempts

Of the 4,860 exposed dummy caterpillars, 22.8% (*n* = 1,108) had bird bill marks. Model selection retained two models in the set of competing models in a range of ΔAICc < 2 (Table S3.2 in Appendix S3). Bird functional richness (FRic) (RVI = 1.00), bioacoustic index (BI) (RVI = 1.00) and temperature

(RVI=1.00) were selected in all models. Landscape forest cover (RVI = 0.66) was also selected in one of

the two best models.

Bird predation attempts decreased with increasing mean annual temperature. Bird predation attempts

further increased with bioacoustic index (BI), but decreased with bird functional richness (FRic) (Fig.

4b and Fig. 5d). This finding suggests that the acoustic component captures some features of the

habitat that influence predation attempts independently of bird functional diversity. Likewise, the fact

that temperature was selected a significant predictor of bird predation attempts suggests that climate

has an effect on predation that is not only mediated by its effect on bird communities.

The results were comparable when we incorporated latitudinal changes in diel phenology in the

calculation of predation attempts through the standardization with the daylight duration (see Table

S3.2 in Appendix S3).

Discussion

531
532

Our study confirms the well documented increase of bird diversity towards warmer regions, a pattern

supporting our initial assumption that avian predation would mirror this pattern. Yet, we found the

opposite – predation attempts decreased with increasing temperature – which dismissed our

prediction that bird diversity and avian predation rate should correlated positively across large

geographic gradients. An important result of our study is that even when the functional aspects of bird

communities were accounted for, a substantial amount of variability remained to be explained and

were only partially accounted for by climate- and habitat-related variables. Altogether, these findings

suggests that current theory should be re-assessed, which we discuss below speculating on the main

causes of deviation from theoretical expectations.

Functional diversity of insectivorous birds and bird predation attempts are both influenced by

climate, in opposite ways

In agreement with our first prediction (i, Fig. 1), we provide evidence for a significant positive

relationship between temperature and the functional diversity of insectivorous birds. Despite

substantial differences among functional diversity indices, this suggests that, in oak woods towards

the South of Europe, more functionally diverse assemblages of insectivorous birds are able to coexist

locally (Currie et al., 2004; Hillebrand, 2004; Willig et al., 2003). Of the multiple functional diversity

indices commonly used to describe ecological communities, it is noticeable that only the quadratic

entropy index responded positively to temperature, for it is a synthetic index that simultaneously takes

into account the richness, evenness, and divergence components of functional diversity (Mouillot et

al., 2013).

However, contrary to our predictions (i, Fig. 1), bird predation attempts decreased with increasing

temperature and were therefore inconsistently linked with bird functional diversity. More bird

predation attempts at lower temperatures could be due to longer daylight duration in spring

northwards, leading insectivorous birds to have more time per day to find their prey and thus allowing

high coexistence of predators during a period of unlimited resource availability (Speakman et al.,

2000). Alternatively, birds needing more energy in colder temperatures to thermoregulate may need

to feed more to maintain their metabolic activity (Caraco et al., 1990; Kendeigh, 1969; Steen, 1958;

Wansink & Tinbergen, 1994). Moreover, temperature remained an important, significant predictor of

bird predation attempts when we controlled for the duration of daylight (Table S3.2 in Appendix S3),

which further supports this explanation. However, we cannot exclude the possibility that the lower

predation rates at higher temperatures was due to lower prey detectability.

Bird predation attempts are partly predicted by bird functional and acoustic diversity

We predicted that bird predation attempts would increase with bird functional diversity (ii, Fig. 1), but

found no or opposite effects, depending on the functional index considered. Only functional

insectivore richness was negatively correlated to predation attempts. Negative relationships between

predation and predator functional diversity can arise from a combination of both intraguild predation

--- predators preying upon predators (Mooney et al., 2010) --- and intraguild competition (Houska

Tahadlova et al., 2022), although we could not tease them apart in the present study. An important

step forward would consist in testing whether predation patterns revealed using artificial prey are

representative of predation intensity as a whole (Zvereva & Kozlov, 2021). For example, functional

richness may be a proxy for dietary specialization in such a way that more functionally diverse predator

communities would seek more prey of which they are specialized on and thus predate less on artificial

caterpillars. It is also possible that a higher diversity of insectivorous birds in warmer regions was linked

to higher diversity and abundance of arthropod prey and foraging niches (Kissling et al., 2012) and

therefore to greater prey availability (Charbonnier et al., 2016). If so, then the pattern we observed

may merely be representative of the 'dilution' of bird attacks on artificial prey among more abundant

and diverse real prey (Zeuss, Brunzel & Brandl, 2017; Zvereva et al., 2019). However, the dynamics

between herbivore prey abundance and predation activity are complex. A higher abundance of real

herbivore prey could also lead to increased predation activity as demonstrated in studies such as

Singer, Singer, Farkas, Skorik & Mooney (2011), where the presence of abundant herbivorous prey was

found to drive higher predation rates by bird predators. This aligns with the notion that predator

populations respond to fluctuations in prey density (Salamolard, Butet, Leroux & Bretagnolle, 2000),

adjusting their foraging behavior to capitalize on available food resources. A follow-up of the present

study should therefore pay special attention to the real prey density pre-existing in each sampling site

where artificial prey are to be deployed as a standardized measure of predation rates across sites.

Our study further brings methodological insights into eco-acoustics. We found that among the various

acoustic indices tested, the Bioacoustic index was positively correlated with bird predation attempts.

Other indices were found to be acceptable proxies for bird diversity. For example, Towsey et al. (2014)

demonstrated that acoustic indices, in particular the Acoustic Entropy Index (H) (see also, Fig. S5.1 in

the supplementary information) can be a good indicator of bird species diversity. Whether acoustic

indices can be used as proxies for ecosystem functions is however still unknown. When dominated by

the song of a single species, acoustic indices can be used to infer the species' presence, behavior, and

potential impact on ecosystem processes (Smith et al., 2020). We show that this could also be the case

in more diverse communities. This is particularly relevant to studies on predation because bird songs

are territorial markers; if a species is recorded in a given site during the breeding season, it indicates

that it is probably feeding on that territory and can potentially affect predation rates. Bioacoustic Index

(BI) that was positively linked to predation attempts is an index linked to the abundance of singing

birds (Sueur et al., 2014). This finding is thus consistent with previous studies having reported positive

relationships between bird abundance and predation attempts on artificial prey (Roels, Porter &

Lindell, 2018; Sam, Koane & Novotny, 2015). It was known that acoustic diversity indices are

complementary to functional diversity indices and reflect both the composition of the avian

community and the nature of the 'acoustic' habitat (Gasc, Francomano, Dunning & Pijanowski, 2017;

Sugai, Silva, Ribeiro & Llusia, 2019). We further show that the acoustic monitoring of the habitat could

yield proxies for the intensity of bird predation on invertebrates. Our study is among the first to detect

significant relationships between acoustic indices and an ecosystem function measured independently

(Gasc et al., 2013), and the study opens pathways for new research on the link between functional and

acoustic ecology.

Local forest cover predicts bird acoustic activity, whereas landscape forest cover increases bird

predation

Acoustic diversity increased with closeness of canopy cover in the immediate neighborhood (20m

radius) of sampled trees (iii, Fig. 1). The most responsive indices were the acoustic diversity (ADI) and

the acoustic entropy (H). Both are especially suitable at predicting acoustic diversity across different

habitats under various ambient sound conditions (Fuller, Axel, Tucker & Gage, 2015; Machado, Aguiar

& Jones, 2017). The former is related to a greater regularity of the soundscape and the latter is related

to the amplitude between frequency bands and time. They therefore correspond to soundscapes

containing multiple vocalizing species (Sueur, Pavoine, et al., 2008; Villanueva-Rivera, Pijanowski,

Doucette & Pekin, 2011). Acoustic entropy is also known to respond significantly to local forest habitat

(Barbaro et al., 2022), which is generally a good predictor of bird occupancy probability (Morante-Filho,

Benchimol & Faria, 2021).

Bird predation attempts were best predicted by forest cover at the landscape level (Prediction (iii), Fig.

1). Indeed, it is likely that forest cover at the landscape level provides structural complexity with a

dense understorey and habitat heterogeneity that is both a source of food and niches for predatory

birds to exploit (Poch & Simonetti, 2013). As a result, forest cover at the landscape scale is often a key

predictor of avian insectivory in various study areas (Barbaro et al., 2014; González-Gómez, Estades &

Simonetti, 2006; Valdés-Correcher et al., 2021). This is also consistent with the results of Rega-Brodsky

& Nilon (2017) who found greater abundance of insectivorous birds in mosaic urban or rural landscapes

including a significant part of semi-natural wooded habitats, such as those we studied here.

Climate cline in avian predation seems independent from large-scale changes in the diversity and

acoustic activity of birds

We found no evidence that the relationship between climate and bird predation attempts was

mediated by changes in bird diversity or activity (iv, Fig. 1). On the contrary, climate and bird diversity

had independent and complementary effects on predation.

At the European scale, climate may directly drive both bird activity and abundance according to

available resources (Pennings & Silliman, 2005). Even changes in the abundance of a single, particularly

active, predator species along the European climatic gradient could explain the observed pattern

(Maas, Tscharntke, Saleh, Dwi Putra, Clough & Siriwardena, 2015; Philpott et al., 2009). For example,

the blue tit *Cyanistes caeruleus* and the great tit *Parus major* are typical and widespread canopy

insectivores of European oak forests and are particularly prone to predate herbivorous caterpillars

while showing considerable adaptive behavior to prey availability (Mols & Visser, 2002; Naef-Daenzer

& Keller, 1999). If the predation attempts on the plasticine caterpillars were to be predominantly due

to these species, then it would be their abundance and activity that would play a role in predation

attempts rather than the overall diversity of insectivores (Maas et al., 2015). Here, we based our

assessment of functional bird composition on candidate insectivore occurrences obtained from

standardized acoustic surveys, which on one hand insures that we have no observer, site, or

phenological biases on species occurrences, but on the other hand also makes it difficult to precisely

account for each species' abundance. Other complementary methods to assess the relative roles of

each individual bird species on predation rates should be deployed further to better account for actual

predatory bird abundance and activity, including DNA sampling (Garfinkel, Minor & Whelan, 2022),

camera traps (Martínez-Núñez et al., 2021) or species-specific bird surveys involving tape calls or

capture methods.

Conclusion

We found a positive association between temperature and bird functional diversity, and at the same

time, a negative relationship between temperature and avian predation. Our study therefore provides

partial support for the climatic clines in biodiversity hypothesis, but demonstrates that predation does

not follow the same pattern. As cross-continental studies exploring the large-scale relationship

between climate and the strength of biotic interactions generally ignore local factors, we argue that

characterizing the contrasting habitats of the study sites is a good way to circumvent some

inconsistencies in the results. We identify pre-existing real prey density and single key bird species

abundances as two particularly important variables deserving further attention. Furthermore,

predicting ecosystem services — here, potential pest regulation service — on a large scale by

standardized proxies such as acoustic ecology for predator diversity and plasticine caterpillars for

predation function seem to be good ways to reduce methodological biases and strengthen our

understanding of the macro-ecology of biotic interactions.

Data availability statement

For the moment there is an embargo on data and codes which will be lifted after open acceptance. Schille et al., 2023, « Data and codes for the article "Decomposing drivers in avian insectivory: large-scale effects of climate, habitat and bird diversity" », https://doi.org/10.57745/0E0JEA, Recherche Data Gouv.

Copyright statement

References

Anstett, D. N., Chen, W., & Johnson, M. T. J. (2016). Latitudinal Gradients in Induced and Constitutive Resistance against Herbivores. *Journal of Chemical Ecology*, *42*(8), 772-781. https://doi.org/10.1007/s10886-016-0735-6

Bael, S. A. V., Philpott, S. M., Greenberg, R., Bichier, P., Barber, N. A., Mooney, K. A., & Gruner, D. S. (2008). Birds as predators in tropical agroforestry systems. *Ecology*, *89*(4), 928-934. https://doi.org/10.1890/06-1976.1

Barbaro, L., Allan, E., Ampoorter, E., Castagneyrol, B., Charbonnier, Y., De Wandeler, H., Kerbiriou, C., Milligan, H. T., Vialatte, A., Carnol, M., Deconchat, M., De Smedt, P., Jactel, H., Koricheva, J., Le Viol, I., Muys, B., Scherer-Lorenzen, M., Verheyen, K., & van der Plas, F. (2019). Biotic predictors complement models of bat and bird responses to climate and tree diversity in European forests. *Proceedings of the Royal Society B: Biological Sciences*, *286*(1894), 20182193. https://doi.org/10.1098/rspb.2018.2193

Barbaro, L., Assandri, G., Brambilla, M., Castagneyrol, B., Froidevaux, J., Giffard, B., Pithon, J., Puig-

Montserrat, X., Torre, I., Calatayud, F., Gaüzère, P., Guenser, J., Macià-Valverde, F., Mary, S.,

Raison, L., Sirami, C., & Rusch, A. (2021). Organic management and landscape heterogeneity

combine to sustain multifunctional bird communities in European vineyards. *Journal of*

Applied Ecology, *58*(6), 1261-1271. https://doi.org/10.1111/1365-2664.13885

Barbaro, L., Giffard, B., Charbonnier, Y., van Halder, I., & Brockerhoff, E. G. (2014). Bird functional

diversity enhances insectivory at forest edges : A transcontinental experiment. *Diversity and*

Distributions, *20*(2), 149-159. https://doi.org/10.1111/ddi.12132

Barbaro, L., Sourdril, A., Froidevaux, J. S. P., Cauchoix, M., Calatayud, F., Deconchat, M., & Gasc, A.

(2022). Linking acoustic diversity to compositional and configurational heterogeneity in

mosaic landscapes. *Landscape Ecology*, *37*(4), 1125-1143. https://doi.org/10.1007/s10980-

021-01391-8

Barr, A. E., van Dijk, L. J. A., Hylander, K., & Tack, A. J. M. (2021). Local habitat factors and spatial

connectivity jointly shape an urban insect community. *Landscape and Urban Planning*, *214*,

104177. https://doi.org/10.1016/j.landurbplan.2021.104177

Bartoń, K. (2020). *MuMIn : Multi-model inference.* (R package version 1.43.17.) [Logiciel].

https://CRAN.Rproj ect.org/packa ge=MuMIn

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using

lme4. *Journal of Statistical Software*, *67*(1). https://doi.org/10.18637/jss.v067.i01

Blackburn, T. M., & Gaston, K. J. (1996). Spatial patterns in the species richness of birds in the New

World. *Ecography*, *19*(4), 369-376.

Botta-Dukát, Z. (2005). Rao's quadratic entropy as a measure of functional diversity based on

multiple traits. *Journal of vegetation science*, *16*(5), 533-540.

Bradfer-Lawrence, T., Gardner, N., Bunnefeld, L., Bunnefeld, N., Willis, S. G., & Dent, D. H. (2019).

Guidelines for the use of acoustic indices in environmental research. *Methods in Ecology and*

Evolution, *10*(10), 1796-1807.

Brambilla, M., & Gatti, F. (2022). No more silent (and uncoloured) springs in vineyards? Experimental

evidence for positive impact of alternate inter-row management on birds and butterflies.

Journal of Applied Ecology, 1365-2664.14229. https://doi.org/10.1111/1365-2664.14229

Burnham, K. P., & Anderson, D. R. (2002). *Model Selection and Multimodel Inference : A Practical*

Information Theoretic Approach, 2nd edn. (Springer).

Caraco, T., Blanckenhorn, W. U., Gregory, G. M., Newman, J. A., Recer, G. M., & Zwicker, S. M. (1990).

Risk-sensitivity : Ambient temperature affects foraging choice. *Animal Behaviour*, *39*(2),

338-345. https://doi.org/10.1016/S0003-3472(05)80879-6

Charbonnier, Y. M., Barbaro, L., Barnagaud, J.-Y., Ampoorter, E., Nezan, J., Verheyen, K., & Jactel, H.

(2016). Bat and bird diversity along independent gradients of latitude and tree composition

in European forests. *Oecologia*, *182*(2), 529-537. https://doi.org/10.1007/s00442-016-3671-9

Cover, C. L. (2018). *European Union, Copernicus Land Monitoring Service.* European Environment

Agency.

Currie, D. J., Mittelbach, G. G., Cornell, H. V., Field, R., Guégan, J.-F., Hawkins, B. A., Kaufman, D. M.,

Kerr, J. T., Oberdorff, T., & O'Brien, E. (2004). Predictions and tests of climate-based

hypotheses of broad-scale variation in taxonomic richness. *Ecology letters*, *7*(12), 1121-1134.

De la Mora, A., García-Ballinas, J. A., & Philpott, S. M. (2015). Local, landscape, and diversity drivers

of predation services provided by ants in a coffee landscape in Chiapas, Mexico. *Agriculture,*

Ecosystems & Environment, *201*, 83-91. https://doi.org/10.1016/j.agee.2014.11.006

Dobzhansky, T. (1950). Evolution in the tropics. *American scientist*, *38*(2), 209-221.

Dröge, S., Martin, D. A., Andriafanomezantsoa, R., Burivalova, Z., Fulgence, T. R., Osen, K.,

Rakotomalala, E., Schwab, D., Wurz, A., Richter, T., & Kreft, H. (2021). Listening to a changing

landscape : Acoustic indices reflect bird species richness and plot-scale vegetation structure

across different land-use types in north-eastern Madagascar. *Ecological Indicators*, *120*,

106929. https://doi.org/10.1016/j.ecolind.2020.106929

Eaton, E., Caudullo, G., Oliveira, S., & de Rigo, D. (2016). *Quercus robur and Quercus petraea*. 4.

Fuller, S., Axel, A. C., Tucker, D., & Gage, S. H. (2015). Connecting soundscape to landscape : Which

acoustic index best describes landscape configuration? *Ecological Indicators*, *58*, 207-215.

https://doi.org/10.1016/j.ecolind.2015.05.057

Garfinkel, M., Minor, E., & Whelan, C. J. (2022). Using faecal metabarcoding to examine consumption

of crop pests and beneficial arthropods in communities of generalist avian insectivores. *Ibis*,

164(1), 27-43. https://doi.org/10.1111/ibi.12994

Gasc, A., Francomano, D., Dunning, J. B., & Pijanowski, B. C. (2017). Future directions for soundscape

ecology : The importance of ornithological contributions. *The Auk*, *134*(1), 215-228.

https://doi.org/10.1642/AUK-16-124.1

Gasc, A., Sueur, J., Jiguet, F., Devictor, V., Grandcolas, P., Burrow, C., Depraetere, M., & Pavoine, S.

(2013). Assessing biodiversity with sound : Do acoustic diversity indices reflect phylogenetic

and functional diversities of bird communities? *Ecological Indicators*, *25*, 279-287.

https://doi.org/10.1016/j.ecolind.2012.10.009

Gibb, R., Browning, E., Glover-Kapfer, P., & Jones, K. E. (2019). Emerging opportunities and challenges

for passive acoustics in ecological assessment and monitoring. *Methods in Ecology and*

Evolution, *10*(2), 169-185. https://doi.org/10.1111/2041-210X.13101

González-Gómez, P. L., Estades, C. F., & Simonetti, J. A. (2006). Strengthened insectivory in a

temperate fragmented forest. *Oecologia*, *148*(1), 137-143. https://doi.org/10.1007/s00442-

005-0338-3

Greenop, A., Woodcock, B. A., Wilby, A., Cook, S. M., & Pywell, R. F. (2018). Functional diversity

positively affects prey suppression by invertebrate predators : A meta-analysis. *Ecology*,

99(8), 1771-1782. https://doi.org/10.1002/ecy.2378

Hawkins, B. A., Diniz-Filho, J. A. F., Jaramillo, C. A., & Soeller, S. A. (2006). Post-Eocene climate

change, niche conservatism, and the latitudinal diversity gradient of New World birds.

Journal of Biogeography, *33*(5), 770-780. https://doi.org/10.1111/j.1365-2699.2006.01452.x

Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high resolution

interpolated climate surfaces for global land areas. *International Journal of Climatology*,

25(15), 1965-1978. https://doi.org/10.1002/joc.1276

Hill, A. P., Prince, P., Piña Covarrubias, E., Doncaster, C. P., Snaddon, J. L., & Rogers, A. (2018).

AudioMoth : Evaluation of a smart open acoustic device for monitoring biodiversity and the

environment. *Methods in Ecology and Evolution*, *9*(5), 1199-1211.

https://doi.org/10.1111/2041-210X.12955

Hillebrand, H. (2004). On the Generality of the Latitudinal Diversity Gradient. *The American*

Naturalist, *163*(2), 192-211. https://doi.org/10.1086/381004

Houska Tahadlova, M., Mottl, O., Jorge, L. R., Koane, B., Novotny, V., & Sam, K. (2022). Trophic

cascades in tropical rainforests : Effects of vertebrate predator exclusion on arthropods and

plants in Papua New Guinea. *Biotropica*.

Ikin, K., Barton, P. S., Stirnemann, I. A., Stein, J. R., Michael, D., Crane, M., Okada, S., & Lindenmayer,

D. B. (2014). Multi-Scale Associations between Vegetation Cover and Woodland Bird

Communities across a Large Agricultural Region. *PLoS ONE*, *9*(5), e97029.

https://doi.org/10.1371/journal.pone.0097029

Just, M. G., Dale, A. G., Long, L. C., & Frank, S. D. (2019). Urbanization drives unique latitudinal

patterns of insect herbivory and tree condition. *Oikos*, *128*(7), 984-993.

https://doi.org/10.1111/oik.05874

Kendeigh, S. C. (1969). *Energy Responses of Birds to Their Thermal Environments*. *81*(4), 10.

Kissling, W. D., Sekercioglu, C. H., & Jetz, W. (2012). Bird dietary guild richness across latitudes,

environments and biogeographic regions : Guild richness in birds. *Global Ecology and*

Biogeography, *21*(3), 328-340. https://doi.org/10.1111/j.1466-8238.2011.00679.x

Kleijn, D., Rundlöf, M., Scheper, J., Smith, H. G., & Tscharntke, T. (2011). Does conservation on

farmland contribute to halting the biodiversity decline? *Trends in Ecology & Evolution*, *26*(9),

474-481. https://doi.org/10.1016/j.tree.2011.05.009

Kozlov, M. V., Lanta, V., Zverev, V., & Zvereva, E. L. (2015). Global patterns in background losses of

woody plant foliage to insects : Latitudinal patterns in insect herbivory. *Global Ecology and*

Biogeography, *24*(10), 1126-1135. https://doi.org/10.1111/geb.12347

Krishnan, A., & Tamma, K. (2016). Divergent morphological and acoustic traits in sympatric

communities of Asian barbets. *Royal Society Open Science*, *3*(8), 160117.

https://doi.org/10.1098/rsos.160117

Laliberté, E., Legendre, P., & Shipley, B. (2014). *Measuring Functional Diversity (FD) from Multiple*

Traits, and Other Tools for Functional Ecology : Package « FD » (1.0-12.1) [Logiciel].

Low, P. A., Sam, K., McArthur, C., Posa, M. R. C., & Hochuli, D. F. (2014). Determining predator

identity from attack marks left in model caterpillars : Guidelines for best practice.

Entomologia Experimentalis et Applicata, *152*(2), 120-126.

https://doi.org/10.1111/eea.12207

Maas, B., Tscharntke, T., Saleh, S., Dwi Putra, D., & Clough, Y. (2015). Avian species identity drives

predation success in tropical cacao agroforestry. *Journal of Applied Ecology*, *52*(3), 735-743.

https://doi.org/10.1111/1365-2664.12409

MacArthur, R. H. (1984). *Geographical ecology : Patterns in the distribution of species*. Princeton

University Press.

Machado, R. B., Aguiar, L., & Jones, G. (2017). Do acoustic indices reflect the characteristics of bird

communities in the savannas of Central Brazil? *Landscape and Urban Planning*, *162*, 36-43.

https://doi.org/10.1016/j.landurbplan.2017.01.014

Martínez-Núñez, C., Rey, P. J., Manzaneda, A. J., García, D., Tarifa, R., & Molina, J. L. (2021).

Insectivorous birds are not effective pest control agents in olive groves. *Basic and Applied*

Ecology, *56*, 270-280. https://doi.org/10.1016/j.baae.2021.08.006

Melles, S., Glenn, S. M., & Martin, K. (2003). Urban Bird Diversity and Landscape Complexity :

Species-environment Associations Along a Multiscale Habitat Gradient. *Conservation Ecology*,

7(1), art5. https://doi.org/10.5751/ES-00478-070105

Miles, J. (2014). Tolerance and variance inflation factor. *Wiley statsref: statistics reference online*.

Mols, C. M. M., & Visser, M. E. (2002). Great tits can reduce caterpillar damage in apple orchards :

Great tits reduce caterpillar damage. *Journal of Applied Ecology*, *39*(6), 888-899.

https://doi.org/10.1046/j.1365-2664.2002.00761.x

Mooney, K. A., Gruner, D. S., Barber, N. A., Van Bael, S. A., Philpott, S. M., & Greenberg, R. (2010).

Interactions among predators and the cascading effects of vertebrate insectivores on

arthropod communities and plants. *Proceedings of the National Academy of Sciences*,

107(16), 7335-7340. https://doi.org/10.1073/pnas.1001934107

Morante-Filho, J. C., Benchimol, M., & Faria, D. (2021). Landscape composition is the strongest

determinant of bird occupancy patterns in tropical forest patches. *Landscape Ecology*, *36*(1),

105-117. https://doi.org/10.1007/s10980-020-01121-6

Mouillot, D., Graham, N. A. J., Villéger, S., Mason, N. W. H., & Bellwood, D. R. (2013). A functional

approach reveals community responses to disturbances. *Trends in Ecology & Evolution*, *28*(3),

167-177. https://doi.org/10.1016/j.tree.2012.10.004

Naef-Daenzer, B., & Keller, L. F. (1999). The foraging performance of great and blue tits (Parus major

and P. caeruleus) in relation to caterpillar development, and its consequences for nestling

growth and fledging weight. *Journal of Animal Ecology*, *68*(4), 708-718.

Otto, S. B., Berlow, E. L., Rank, N. E., Smiley, J., & Brose, U. (2008). Predator diversity and identity

drive interaction strength and trophic cascades in a food web. *Ecology*, *89*(1), 134-144.

https://doi.org/10.1890/07-0066.1

Pennings, S. C., & Silliman, B. R. (2005). Linking biogeography and community ecology : Latitudinal

variation in plant–herbivore interaction strength. *Ecology*, *86*(9), 2310-2319.

https://doi.org/10.1890/04-1022

Petit, R. J., Brewer, S., Bordács, S., Burg, K., Cheddadi, R., Coart, E., Cottrell, J., Csaikl, U. M., van Dam,

B., & Deans, J. D. (2002). Identification of refugia and post-glacial colonisation routes of

European white oaks based on chloroplast DNA and fossil pollen evidence. *Forest ecology*

and management, *156*(1-3), 49-74.

Philpott, S. M., Soong, O., Lowenstein, J. H., Pulido, A. L., Lopez, D. T., Flynn, D. F. B., & DeClerck, F.

(2009). Functional richness and ecosystem services : Bird predation on arthropods in tropical

agroecosystems. *Ecological Applications*, *19*(7), 1858-1867. https://doi.org/10.1890/08-

1928.1

Poch, T. J., & Simonetti, J. A. (2013). Insectivory in Pinus radiata plantations with different degree of

structural complexity. *Forest Ecology and Management*, *304*, 132-136.

https://doi.org/10.1016/j.foreco.2013.04.044

R Core Team. (2020). *R: A Language and environment for statistical computing.* [Logiciel]. R

Foundation for Statistical Computing, Vienna, Austria. https://www.r-project.org/

Rega-Brodsky, C. C., & Nilon, C. H. (2017). Forest cover is important across multiple scales for bird

communities in vacant lots. *Urban Ecosystems*, *20*(3), 561-571.

https://doi.org/10.1007/s11252-016-0614-5

Roels, S. M., Porter, J. L., & Lindell, C. A. (2018). Predation pressure by birds and arthropods on

herbivorous insects affected by tropical forest restoration strategy. *Restoration Ecology*,

26(6), 1203-1211. https://doi.org/10.1111/rec.12693

Salamolard, M., Butet, A., Leroux, A., & Bretagnolle, V. (2000). Responses of an avian predator to

variations in prey density at a temperate latitude. *Ecology*, *81*(9), 2428-2441.

https://doi.org/10.1890/0012-9658(2000)081[2428:ROAAPT]2.0.CO;2

Sam, K., Koane, B., & Novotny, V. (2015). Herbivore damage increases avian and ant predation of

caterpillars on trees along a complete elevational forest gradient in Papua New Guinea.

Ecography, *38*(3), 293-300.

Sánchez-Giraldo, C., Correa Ayram, C., & Daza, J. M. (2021). Environmental sound as a mirror of

landscape ecological integrity in monitoring programs. *Perspectives in Ecology and*

Conservation, *19*(3), 319-328. https://doi.org/10.1016/j.pecon.2021.04.003

Schemske, D. W., Mittelbach, G. G., Cornell, H. V., Sobel, J. M., & Roy, K. (2009). Is There a Latitudinal

Gradient in the Importance of Biotic Interactions? *Annual Review of Ecology, Evolution, and*

Systematics, *40*(1), 245-269. https://doi.org/10.1146/annurev.ecolsys.39.110707.173430

Sekercioglu, C. (2006). Increasing awareness of avian ecological function. *Trends in Ecology &*

Evolution, *21*(8), 464-471. https://doi.org/10.1016/j.tree.2006.05.007

Sinclair, A. R. E., Mduma, S., & Brashares, J. S. (2003). Patterns of predation in a diverse predator–

prey system. *Nature*, *425*(6955), 288-290. https://doi.org/10.1038/nature01934

Singer, M. S., Farkas, T. E., Skorik, C. M., & Mooney, K. A. (2011). *Tritrophic Interactions at a*

Community Level : Effects of Host Plant Species Quality on Bird Predation of Caterpillars.

Smith, D. G., Truskinger, A., Roe, P., & Watson, D. M. (2020). Do acoustically detectable species

reflect overall diversity? A case study from Australia's arid zone. *Remote Sensing in Ecology*

and Conservation, *6*(3), 286-300. https://doi.org/10.1002/rse2.173

Southwood, T. R. E., Wint, G. W., Kennedy, C. E., & Greenwood, S. R. (2005). The composition of the

arthropod fauna of the canopies of some species of oak (Quercus). *European Journal of*

Entomology, *102*(1), 65-72.

Speakman, J. R., Rydell, J., Webb, P. I., Hayes, J. P., Hays, G. C., Hulbert, I. A. R., & McDevitt, R. M.

(2000). Activity patterns of insectivorous bats and birds in northern Scandinavia (69° N),

during continuous midsummer daylight. *Oikos*, *88*(1), 75-86. https://doi.org/10.1034/j.1600-

0706.2000.880109.x

Steen, J. (1958). Climatic Adaptation in Some Small Northern Birds. *Ecology*, *39*(4), 625-629.

https://doi.org/10.2307/1931602

Sueur, J., Aubin, T., & Simonis, C. (2008). Seewave, a free modular tool for sound analysis and

synthesis. *Bioacoustics*, *18*(2), 213-226. https://doi.org/10.1080/09524622.2008.9753600

Sueur, J., Farina, A., Gasc, A., Pieretti, N., & Pavoine, S. (2014). Acoustic Indices for Biodiversity

Assessment and Landscape Investigation. *Acta Acustica United with Acustica*, *100*(4),

772-781. https://doi.org/10.3813/AAA.918757

Sueur, J., Pavoine, S., Hamerlynck, O., & Duvail, S. (2008). Rapid Acoustic Survey for Biodiversity

Appraisal. *PLoS ONE*, *3*(12), e4065. https://doi.org/10.1371/journal.pone.0004065

Sugai, L. S. M., Silva, T. S. F., Ribeiro, J. W., & Llusia, D. (2019). Terrestrial Passive Acoustic

Monitoring : Review and Perspectives. *BioScience*, *69*(1), 15-25.

https://doi.org/10.1093/biosci/biy147

Tobias, J. A., Sheard, C., Pigot, A. L., Devenish, A. J. M., Yang, J., Sayol, F., Neate-Clegg, M. H. C.,

Alioravainen, N., Weeks, T. L., Barber, R. A., Walkden, P. A., MacGregor, H. E. A., Jones, S. E.

I., Vincent, C., Phillips, A. G., Marples, N. M., Montaño-Centellas, F. A., Leandro-Silva, V.,

Claramunt, S., … Schleuning, M. (2022). AVONET : Morphological, ecological and geographical

data for all birds. *Ecology Letters*, *25*(3), 581-597. https://doi.org/10.1111/ele.13898

Towsey, M., Wimmer, J., Williamson, I., & Roe, P. (2014). The use of acoustic indices to determine

avian species richness in audio-recordings of the environment. *Ecological Informatics*, *21*,

110-119. https://doi.org/10.1016/j.ecoinf.2013.11.007

Valdés-Correcher, E., Moreira, X., Augusto, L., Barbaro, L., Bouget, C., Bouriaud, O., Branco, M.,

Centenaro, G., Csóka, G., Damestoy, T., Dobrosavljević, J., Duduman, M., Dulaurent, A.,

Eötvös, C. B., Faticov, M., Ferrante, M., Fürjes-Mikó, Á., Galmán, A., Gossner, M. M., …

Castagneyrol, B. (2021). Search for top-down and bottom-up drivers of latitudinal trends in

insect herbivory in oak trees in Europe. *Global Ecology and Biogeography*, *30*(3), 651-665.

https://doi.org/10.1111/geb.13244

Vellinga, W.-P., & Planque, R. (2015). *The Xeno-canto collection and its relation to sound recognition*

and classification. 11.

Vickery, J., & Arlettaz, R. (2012). The importance of habitat heterogeneity at multiple scales for birds

in European agricultural landscapes. *Fuller, R.J. (éd) Birds and Habitat : Relationships in*

Changing Landscapes, 177-204.

Villanueva-Rivera, L. J., & Pijanowski, B. C. (2018). *Soundscape Ecology* (package « sounecology »

1.3.3) [Logiciel]. http://ljvillanueva.github.io/soundecology/

Villanueva-Rivera, L. J., Pijanowski, B. C., Doucette, J., & Pekin, B. (2011). A primer of acoustic analysis

for landscape ecologists. *Landscape Ecology*, *26*(9), 1233-1246.

https://doi.org/10.1007/s10980-011-9636-9

Villéger, S., Mason, N. W. H., & Mouillot, D. (2008). New multidimensional functional diversity indices

for a multifaceted framework in functional ecology. *Ecology*, *89*(8), 2290-2301.

https://doi.org/10.1890/07-1206.1

Wansink, D., & Tinbergen, J. M. (1994). The Influence of Ambient Temperature on Diet in the Great

Tit. *Journal of Avian Biology*, *25*(4), 261. https://doi.org/10.2307/3677272

Willig, M. R., Kaufman, D. M., & Stevens, R. D. (2003). Latitudinal Gradients of Biodiversity : Pattern,

Process, Scale, and Synthesis. *Annual Review of Ecology, Evolution, and Systematics*, *34*(1),

273-309. https://doi.org/10.1146/annurev.ecolsys.34.012103.144032

Zeuss, D., Brunzel, S., & Brandl, R. (2017). Environmental drivers of voltinism and body size in insect

assemblages across Europe : Voltinism and body size in insect assemblages. *Global Ecology*

and Biogeography, *26*(2), 154-165. https://doi.org/10.1111/geb.12525

Zvereva, E. L., Castagneyrol, B., Cornelissen, T., Forsman, A., Hernández-Agüero, J. A., Klemola, T.,

Paolucci, L., Polo, V., Salinas, N., Theron, K. J., Xu, G., Zverev, V., & Kozlov, M. V. (2019).

Opposite latitudinal patterns for bird and arthropod predation revealed in experiments with

differently colored artificial prey. *Ecology and Evolution*, *9*(24), 14273-14285.

https://doi.org/10.1002/ece3.5862

Zvereva, E. L., & Kozlov, M. V. (2021). Latitudinal gradient in the intensity of biotic interactions in

terrestrial ecosystems : Sources of variation and differences from the diversity gradient

revealed by meta-analysis. *Ecology Letters*, *24*(11), 2506-2520.

https://doi.org/10.1111/ele.13851

Biosketch

Laura Schillé is a PhD candidate interested in the functional ecology of bird communities, which she studies at different scales. She also has an interest in acoustic ecology.

Co-authors are ornithologists and/or have interests in community ecology and functional ecology.

Author contribution: B.C., L.B. and E.V.C. conceptualized the study and developed the methodology.

E.V.C., F.B., M.C.B, M.Bo., M.Br., T.D., M.dG., J.D., M.L.D., A.M.D., S.G., C.B.E., M.F., P.F.C., E.F., A.G., M.M.G., S.G., J.H., D.H., R.H., R.I., A.K., M.V.K, V.L., B.L.T., C.L.V, S.M., E.M., S.M., X.M., A.M., D.L.M., A.P., A.V.P., A.P., A.S., K.S., T.S., A.T., R.T., D.T., G.V., I.V.H., Z.V., L.B. & B.C. collected the data. A.H., M.dG., T.B., A.P., V.L., L.P., F.A., D.F., T.P., M.M., N.S., L.B., V.M., E.M., J.G., K.T., L.G., A.M., M.C.B. & R.B. processed audio recordings for bird species identification. L.S. processed and analyzed the data with guidance from B.C. and L.B. L.S., B.C., L.B. led the writing and all authors contributed critically to the revisions.