Latitudinal gradient in avian insectivory: complementary effects of climate, habitat and bird diversity

Laura Schillé[1,*], Elena Valdés-Correcher[1], Frédéric Archaux[2], Flavius Bălăcenoiu[3,4], Mona Chor Bjørn[5], Michal Bogdziewicz[6], Thomas Boivin[7], Manuela Branco[8], Thomas Damestoy[9], Maarten de Groot[10], Jovan Dobrosavljević[11], Mihai-Leonard Duduman[12], Anne-Maïmiti Dulaurent[9], Samantha Green[13], Jan Grünwald[14], Csaba Béla Eötvös[15], Maria Faticov[16], Pilar Fernandez-Conradi[7], Elisabeth Flury[17], David Funosas[18], Andrea Galmán[19,20], Martin M. Gossner[21,22], Sofia Gripenberg[23], Lucian Grosu[12], Jonas Hagge[24], Arndt Hampe[1], Deborah Harvey[25], Rick Houston[26], Rita Isenmann[27], Andreja Kavčič[10], Mikhail V. Kozlov[28], Vojtech Lanta[29], Bénédicte Le Tilly[30], Carlos Lopez Vaamonde[31], Soumen Mallick[32], Elina Mäntylä[28,33,34], Anders Mårell[2], Slobodan Milanović[11], Márton Molnár[35], Xoaquín Moreira[36], Valentin Moser[37,38], Anna Mrazova[33,39], Dmitrii L. Musolin[40], Thomas Perot[2], Andrea Piotti[41], Anna V. Popova[42], Andreas Prinzing[32], Ludmila Pukinskaya[43], Aurélien Sallé[44], Katerina Sam[33,39], Nickolay V. Sedikhin[43,45], Tanja Shabarova[46], Ayco Tack[47], Rebecca Thomas[25], Karthik Thrikkadeeri[48], Dragoș Toma[3,4], Grete Vaicaityte[49], Inge van Halder[1], Zulema Varela[50], Luc Barbaro[51,52,¤] & Bastien Castagneyrol[1,¤]

[¤] Luc Barbaro and Bastien Castagneyrol share last co-authorship

[1] BIOGECO, INRAE, University Bordeaux, Cestas, France

[2] EFNO, INRAE, Nogent-sur-Vernisson, France

[3] National Institute for Research and Development in Forestry « Marin Drăcea », Voluntari, Romania

[4] Romania Faculty of Silviculture and Forest Engineering, Transilvania University of Brașov, Romania

[5] Department of Geosciences and Natural Resource Management, Section Forest, Nature and Biomas, University of Copenhagen, Denmark

[6] Forest Biology Center, Faculty of Biology, Adam Mickiewicz University in Poznan, Poland

[7] UR629, INRAE, Centre PACA, Avignon, France

[8] Centro de Estudos Florestais, Instituto Superior de Agronomia, Universidade de Lisboa, Lisbon,

Portugal

[9] UniLaSalle, AGHYLE, UP.2018.C101, Beauvais, France

[10] Slovenian Forestry Institute, Ljubljana, Slovenia

[11] Department of Forestry and Nature Protection, University of Belgrade Faculty of Forestry,

Belgrade, Serbia

[12] Forestry Faculty, Applied Ecology Lab, "Ştefan cel Mare" University of Suceava, Romania

[13] Centre for Agroecology, Water and Resilience, Coventry University, UK

[14] Institute for Environmental Studies, Faculty of Science, Charles University in Prague, Czech Republic

[15] University of Sopron, Forest Research Institute, Department of Forest Protection, Mátrafüred,

Hungary

[16] Université de Sherbrooke, 2500, boul. de l'Université, Sherbrooke, Canada

[17] Kinderlab Landquart, Landquart, Switzerland

[18] Laboratoire Ecologie Fonctionnelle et Environnement, University of Toulouse, CNRS, France

[19] Martin-Luther-Universität Halle-Wittenberg, Halle (Saale), Germany

[20] German Centre for Integrative Biodiversity Research (iDiv), Leipzig, Germany

[21] Forest Entomology, Swiss Federal Research Institute WSL, Birmensdorf, Switzerland

[22] Institute of Terrestrial Ecosystems, Department of Environmental Systems Science, ETH Zürich,

Switzerland

[23] School of Biological Sciences, University of Reading, UK

[24] Forest Nature Conservation, Northwest German Forest Research Institute, Hann. Münden,

Germany

[25] Royal Holloway University of London, Egham, UK

[26] Beckley School Oxford, UK

[27] Grimmelshausenschule, Renchen, Germany

[28] Department of Biology, University of Turku, Finland

[29] Department of Functional Ecology, Institute of Botany CAS, Průhonice, Czech Republic

[30] Collège Saint Joseph, Paimpol, France

[31] INRAE Centre Val de Loire, Unité Zoologie Forestière, Orléans & IRBI, UMR 7261, CNRS-Université

de Tours, France

[32] Centre National de la Recherche Scientifique, Université de Rennes 1, Research Unit UMR 6553,

Ecosystèmes Biodiversité Evolution (ECOBIO), Campus de Beaulieu, 35042, Rennes, France

[33] Institute of Entomology, Biology Centre of the Czech Academy of Sciences, Ceske Budejovice, Czech

Republic

[34] Faculty of Science, Department of Zoology, University of South Bohemia, Ceske Budejovice, Czech

Republic

[35] Magyar Madártani és Természetvédelmi Egyesület (Birdlife Hungary), Szuha, Hungary

[36] Misión Biológica de Galicia (MBG-CSIC), Apdo. 28, 36080 Pontevedra, Galicia, Spain

[37] Community Ecology, Swiss Federal Research Institute WSL, Birmensdorf, Switzerland

[38] Aquatic Ecology, Swiss Federal Institute of Aquatic Science and Technology Eawag, Dübendorf,

Switzerland

[39] University of South Bohemia, Ceske Budejovice, Czech Republic

[40] European and Mediterranean Plant Protection Organization, Paris, France

[41] Institute of Biosciences and BioResources, National Research Council of Italy, Sesto Fiorentino, Italy

[42] A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Moscow, Russia

[43] Saint Petersburg State Forest Technical University, Lisinsky Forest College, Saint Petersburg, Russia

[44] LBLGC, Université d'Orléans, France

[45] Zoological institute of Russian Academy of Science, Saint Petersburg, Russia

[46] Biology Centre CAS, Institute of Hydrobiology, Ceske Budejovice, Czech Republic

[47] Department of Ecology, Environment and Plant Sciences, Stockholm University, Sweden

[48] Nature Conservation Foundation, Bengaluru, India

[49] Lithuanian Centre of Non-Formal Youth Education, Vilnius, Lithuania

[50] CRETUS,Ecology Unit, Department of Functional Biology, Universidade de Santiago de Compostela,

Spain

[51] DYNAFOR, University of Toulouse, INRAE, France

[52] LTSER ZA Pyrenees Garonne, Auzeville-Tolosane, France

* Correspondance: *laura.schille@hotmail.fr*

Acknowledgements

This study was permitted by the financial support of the BNP Paribas Foundation through its Climate & Biodiversity Initiative for the 'Tree bodyguards' citizen science project. FB and DT was supported by the PN 23090102 and 34PFE./30.12.2021 'Increasing the institutional capacity and performance of INCDS "Marin Drăcea" in the activity of RDI - CresPerfInst' funded by the Ministry of Research, Innovation and Digitalization of Romania.MB is supported by the Forest Research Centre (CEF) (UIDB/00239/2020) and the Laboratory for Sustainable Land Use and Ecosystem Services—TERRA (LA/P/0092/2020) funded by FCT, Portugal. MdG and AK were supported by the core research group "Forest biology, ecology and technology" (P4-0107) of the Slovenian Research Agency. MVK was supported by the Academy of Finland (project 316182). VL was funded by the Czech Science Foundation (project 23-07533S) and Academy of Sciences (RVO 67985939). EM and KS were supported by the Grant Agency of the Czech Republic (19-28126X). KS and AM were supported by ERC StG BABE 805189. XM was supported by a grant from the Spanish National Research Council (2021AEP082) and a grant from the Regional Government of Galicia (IN607A 2021/03). ZV was supported by a grant awarded by the Autonomous Government of Galicia (Spain; Modalidade B-2019), and Maria Zambrano program from the Spanish Ministry of Universities. LB was supported by fundings from LTSER ZA Pyrenees Garonne. We thank the personnel of Centro Nazionale Carabinieri Biodiversità "Bosco Fontana" for help in data collection. We thank Tarja Heinonen for help in field work in Berlin, Germany.

101 We thank Roman Hrdlička for bird voice identification. We thank E. L. Zvereva for help in data
102 collection.

Abstract

Aim

According to the Latitudinal Biotic Interaction Hypothesis (LBIH), the general increase in biodiversity towards lower latitudes can be partially explained by an increase in the intensity of biotic interactions. While LBIH received some support for plant-herbivores interactions, much less is known about how higher trophic levels may contribute to shape biotic interactions across latitudinal gradients. We hypothesized that the diversity of insectivorous birds increases towards lower latitude, leading to higher predation rates on insect herbivores.

Location

Europe.

Taxon

Insectivorous birds and pedunculate oaks.

Methods

We deployed plasticine caterpillars in 138 oak trees in 47 sites along a 19° latitudinal gradient in Europe to quantify bird insectivory through predation attempts. In addition, we used passive acoustic monitoring to (i) characterize the acoustic diversity of surrounding soundscapes; and (ii) infer both taxonomic and functional diversity of insectivorous birds from recordings.

Results

The functional diversity of insectivorous birds increased towards lower latitude. Bird predation

increased with latitude, forest cover and bird acoustic diversity but decreased with mean annual

temperature and functional richness of insectivorous birds. Contrary to our predictions, latitudinal

clines in bird predation attempts were not directly mediated by changes in insectivorous bird diversity

or acoustic diversity, but latitude and habitat still had independent effects on predation attempts.

Main conclusions

Our study does not fully support the predictions of the LBIH of more biotic interactions southwards

and advocates for better accounting for activity and abundance of insectivorous birds when studying

the large-scale variation in insect-tree interactions.

Keywords: Acoustic diversity, Functional diversity, Insectivorous birds, Latitudinal gradient, Plasticine

caterpillars, Predation function

133 Introduction

The general increase in terrestrial and marine biodiversity from high to low latitudes is one of the most

consistent patterns in biogeography (MacArthur, 1984; Schemske, Mittelbach, Cornell, Sobel & Roy,

2009). Of the numerous hypotheses that have been proposed to explain this ubiquitous pattern, the

Latitudinal Biotic Interactions Hypothesis (LBIH) states that biotic interactions are more intense and

are the main driver of biotic evolution under milder climates, at low latitudes (Dobzhansky, 1950).

Whereas LBIH received partial support from numerous studies on plant-herbivore interactions

(Anstett, Chen & Johnson, 2016; Kozlov, Lanta, Zverev & Zvereva, 2015), much less is known about

biotic interactions involving higher trophic levels. Given the importance of top-down factors in the

control of herbivory, this omission represents a critical gap in knowledge that needs to be addressed.

Latitudinal clines in predation rates have received much less attention than latitudinal clines in

herbivory, and remains inconclusive (see Zvereva & Kozlov, 2021 for a review). Some authors have

documented an increase in predation by arthropods toward the equator (Roslin et al., 2017; Zvereva

et al., 2019), but neutral (Lövei & Ferrante, 2017; Roslin et al., 2017; Valdés-Correcher et al., 2021) or

even opposite (Zvereva et al., 2019) patterns have been found for predation by birds. Local variation

in resources drives the diversity and foraging activity of predators, with cascading effects on predation

intensity. By controlling for these factors we can further test our understanding of the effect of

latitudinal clines on prey-predator interactions.

Current theory predicts a positive relationship between biodiversity and ecosystem functioning

regulation (Loreau et al., 2001), suggesting that greater predator diversity may, to some extent, lead

to greater predation (van Bael et al., 2008; Otto, Berlow, Rank, Smiley & Brose, 2008; Sinclair, Mduma

& Brashares, 2003). Beyond taxonomic diversity, functional diversity – that is the diversity, distribution

and complementarity of predator traits involved in predation – is a key driver of predation rates

(Barbaro et al., 2014; Greenop, Woodcock, Wilby, Cook & Pywell, 2018; Philpott et al., 2009). Birds are

among predators contributing the most to the control of insect herbivores in terrestrial ecosystems

(van Bael et al., 2008; Sekercioglu, 2006). Niche opportunities increase towards lower latitudes due to

climatic conditions that allow higher habitat heterogeneity and increased species coexistence, leading

to higher bird species richness and greater functional diversity (Blackburn & Gaston, 1996; Willig,

Kaufman & Stevens, 2003). Bird predation should in turn follow the same pattern as bird diversity

(Zvereva & Kozlov, 2021). However, whether the diversity-predation relationship ultimately

contributes to increased predation upon herbivores, and whether it cascades down to the primary

producer level, is still controversial (Mooney et al., 2010), partly due to bird predation on predatory

arthropods (i.e., intraguild predation).

Biological communities are shaped by a combination of biotic and abiotic factors acting at multiple,

nested scales (Charbonnier et al., 2016; Ikin et al., 2014; Kissling, Sekercioglu & Jetz, 2012). Thus, also

local factors can alter macroecological patterns by filtering the regional species pool (De la Mora,

García-Ballinas & Philpott, 2015; Kleijn, Rundlöf, Scheper, Smith & Tscharntke, 2011). Forest cover

proved to be a good predictor of biodiversity, including acoustic diversity, likely due to the complex

architecture of trees and diversity of resources and trophic interactions between trees, insect

herbivores and their natural enemies (Barbaro et al., 2022; Long & Frank, 2020). Birds are particularly

sensitive to forest cover at different scales, which influences both the amount of available prey and

nesting sites (Lee, Fahrig, Freemark & Currie, 2002; Mazerolle & Villard, 1999; Rega-Brodsky & Nilon,

2017). As a result, modeling the response of forest bird communities to large-scale bioclimatic drivers

benefits from using a combination of habitat variables and biotic predictors such as food amount and

availability (Barbaro et al., 2019; Speakman et al., 2000). At the local scale, bird foraging activity is

ultimately determined by the vertical and horizontal heterogeneity of the habitat, which influences

both where prey can be found, and where foraging birds can hide from predators (Vickery & Arlettaz,

2012). However, cross-continental studies exploring the relationship between latitude and the

strength of biotic interactions generally ignore local factors, which may partly explain inconsistencies

in their findings (but see Just, Dale, Long & Frank, 2019).

A major challenge to analyze latitudinal patterns in biotic interactions consists in simultaneously

characterizing changes in predator biodiversity and experimentally assessing the strength of predation,

while considering the effect of contrasting habitats. However, the recent development of passive

acoustic monitoring provides a standardized, low-cost and non-invasive approach for ecological

studies and biodiversity monitoring (Gibb, Browning, Glover-Kapfer, Jones & Börger, 2019). Acoustic

monitoring of a habitat firstly allows the identification of bird species. Secondly, through the

calculation of acoustic diversity indices, it allows to characterize the activity of the community as a

whole while also accounting for the importance of human disturbances (Gasc et al., 2013; Sueur,

Farina, Gasc, Pieretti & Pavoine, 2014). Should such indices consistently correlate with macro-scales

biotic interactions, ecoacoustics would be a promising complementary approach to existing methods

in macroecology and in functional ecology.

Here, we addressed the LBIH through the lens of predation by simultaneously characterizing changes

in insectivorous bird community diversity, bird predation, and the biodiversity-function relationship

while controlling for local factors throughout the European distribution range of the pedunculate oak

(*Quercus robur* L., 1753), a major forest tree species. Specifically, we predict the following (Fig. 1): (i)

bird diversity (including bird acoustic diversity, insectivorous bird species richness and functional

diversity) and predation rates increase toward lower latitudes; (ii) bird predation rates increase with

bird acoustic activity, species richness and greater functional diversity of insectivorous birds; (iii) bird

diversity, acoustic activity and bird predation rates increase with increasing forest cover at both local

(neighborhood) and larger spatial scales; (iv) large-scale variability in bird predation rates is primarily

driven by local changes in the diversity and activity of birds. To test these predictions, we quantified

bird predation attempts on plasticine caterpillars and estimated bird species richness, functional

diversity and activity through simultaneous passive acoustic monitoring. We eventually tested the

respective responses of these variables and their relationships at the pan-European scale.

Figure 1: Conceptual diagram of the predictions of this study and the relationships already established in the literature. Boxed

elements written in bold correspond to the main categories of variables tested, they are not variables as such. Variables used

in models are shown in regular font. Where several variables described the same category (e.g. ACI, NDSI, BI, ADI, H, NPIC,

MultiAcou, all describing acoustic indices), we used multi-model comparisons to identify the best variable. Items framed in

black on a white background represent untested variables. Black arrows indicate relationships well supported by the literature

(see Fig.2 Sánchez-Giraldo, Correa Ayram & Daza, 2021). Our specific predictions are represented with grey arrows, solid and

dashed lines representing positive and negative (predicted) relationships. Numbers refer to predictions as stated in the main

text.

Materials and methods

Study area

We focused on the pedunculate oak, *Quercus robur,* which is one of the keystone deciduous tree

species in temperate European forests, where it is of high ecological, economic and symbolic

importance (Eaton, Caudullo, Oliveira & de Rigo, 2016). The species occurs from central Spain (39°N)

to southern Fennoscandia (62°N) and thus experiences a huge gradient of climatic conditions (Petit et

al., 2002). A widely diverse community of specialist and generalist herbivorous insects is associated

with this species throughout its distributional range (Southwood, Wint, Kennedy & Greenwood, 2005).

Between May and July 2021, we studied 138 trees in 47 sites across 17 European countries covering

most of the pedunculate oak geographic range (Fig. 2). The sites were chosen with the minimal

constraint of being located in a wooded area of at least 1ha (Valdés-Correcher et al., 2021). We

haphazardly selected three mature oaks per site, with the exception of six sites (three sites with one

tree, one site with two trees and two sites with five trees, see Table S1.1 in Appendix S1 in Supporting

Information).

230

231 **Figure 2** *Locations of the 47 sites sampled in spring 2021. Average annual temperature (color scale) according to WorldClim*

232 *(Hijmans, Cameron, Parra, Jones & Jarvis, 2005) and Quercus robur distribution range are indicated.*

Bird predation attempts

We measured bird predation attempts in the field by exposing a total of 40 plasticine caterpillars (20

plasticine caterpillars twice) on each individual oak. We made plasticine caterpillars of green plasticine,

mimicking common lepidopteran larvae (3 cm long, 0.5 cm diameter, see Low, Sam, McArthur, Posa &

Hochuli, 2014). We secured them on twigs with a 0.3 mm metallic wire. We attached five plasticine

caterpillars to each of four branches facing opposite directions (i.e., 20 caterpillars per tree) at about

2 m from the ground.

We installed the plasticine caterpillars six weeks after budburst in each study area, thus synchronizing

the study with local oak phenology. We removed the plasticine caterpillars after 15 days and installed

another set of 20 artificial caterpillars per tree for another 15 days. At the end of each exposure period

(which varied from 10 to 20 (mean ± SD: 14.5 ± 1.23) days due to weather conditions, we carefully

removed the plasticine caterpillars from branches, placed them into plastic vials and shipped them to

the project coordinator. Plasticine caterpillars from six sites were either lost or altered during shipping,

preventing the extraction of relevant data.

A single trained observer (EVC) screened the surface of plasticine caterpillars with a magnifying lens to

search for the presence of bill marks on clay surface (Low et al., 2014). As we were ultimately interested

in linking bird diversity with bird predation rates, we did not consider marks left by arthropods and

mammals.

We defined *bird predation attempts index* as *p / d*, where *p* is the proportion of plasticine caterpillars

with at least one sign of attempted predation by birds and *d* is the number of days plasticine caterpillars

were exposed to predators in the field. We only considered as attacked those caterpillars that we

retrieved; missing caterpillars were not accounted for in the calculation of *p*. We calculated bird

predation attempts for each tree and survey period separately. Because other variables were defined

at site level (see below), we averaged bird predation attempts across trees and surveys in each site

(total: *n* = 41).

To assess the effect of temperature independently of other variables that could covary with latitude,

we also calculated a second bird predation attempts index by standardizing the predation attempts by

daylight duration in every site (see Fig. S2.1 in Appendix S2 for the calculation of this index). We run

the statistical models in exactly the same way as for the non-standardized bird predation attempts and

the results are presented in Table S3.2 in Appendix S3.

Acoustic monitoring and related variables

We used passive acoustic monitoring to characterize both the overall acoustic diversity (i.e., the

soundscape surrounding target oaks) and the bird species richness and functional diversity associated

with oaks, targeting the canopy insectivores that are expected to play a functional predation role on

oak herbivores (Fig. 2). In each site, we randomly chose one oak among those used to measure bird

predation rates in which we installed an AudioMoth device (Hill et al., 2018) to record audible sounds

for 30 min every hour. Automated recording started the day we installed the first set of 20 plasticine

caterpillars in trees and lasted until batteries stopped providing enough energy. The recording settings

are the following: *Recording period: 00.00-24.00 (UTC); Sample rate: 48 kHZ; Gain: Medium; Sleep*

duration: 1800 s, Recording duration: 1800 s.

In all 47 sites, Audiomoths were active on average (± S.D.) for 9 ± 3 days (range: 1-24), which

corresponded to 5920 h of recordings in total and from 70 to 335 (246 ± 65) 30 min continuous acoustic

samples per site. When Audiomoths ran out of battery, the recordings lasted less than 30 min

(between 1 and 56 recordings per site were affected).

We processed acoustic samples with functions in the *soundecology* (Villanueva-Rivera & Pijanowski,

2018) and *seewave* libraries (Sueur, Aubin & Simonis, 2008) in the R environment (R Core Team, 2020),

and a wrap-up function made available by A. Gasc in GitHub (https://github.com/agasc/Soundscape-

analysis-with-R). To summarize information on the acoustic diversity of each site, we first divided every

acoustic sample (regardless of its length) into non-overlapping 1 min samples. Then, we calculated six

common acoustic diversity indices for each 1 min sample: the Acoustic Complexity Index (ACI), the

Acoustic Diversity Index (ADI), the Bioacoustic Index (BI), the total acoustic entropy (H), the Normalized

Difference Sound Index (NDSI), and the Number of frequency Peaks NPIC (Bradfer-Lawrence et al.,

2019; Sueur et al., 2014). We further aggregated these indices into a multiacoustic index (MultiAcou)

as proposed by Barbaro et al. (2022). We calculated the median of each acoustic index per day and

then averaged median values across days for each site separately. We proceeded like this because 24

h cycles summarize the acoustic activity and account for all possible sounds of a given day.

Furthermore, other studies have previously shown that median values of acoustic indices for a given

day are more representative than mean values of the acoustic activity because they are less sensitive

to extreme values (Barbaro et al., 2022; Dröge et al., 2021). This procedure resulted in one single value

of each acoustic diversity index per site. It is important to note that being calculated over 24 h, these

indices do not represent bird-related variables only, but are rather seen as an integrative component

of the surrounding habitat around target oaks.

Bird species richness and functional diversity

We used acoustic samples to identify birds based on their vocalizations (songs and calls) at the species

level, from which we further computed functional diversity indices (Fig. 3).

Data processing – For each site, we subsampled the 30 min samples corresponding to the songbird

morning chorus (*i.e.,* the period of maximum singing activity), which incidentally also corresponds to

the time of the day when anthropic sounds were of the lowest intensity. Specifically, we selected

sounds recorded within a period running from 30 min before sunrise to 3 h 30 min after sunrise. We

then split each 30 min sample into up to three 10 min sequences, from which we only retained those

recorded on Tuesday, Thursday, Saturday, and Sunday. We chose these days on purpose to balance

the differences in anthropogenic noises between working days and weekends. For each sound sample,

we displayed the corresponding spectrogram with the '*seewave*' library in the *R* environment (Sueur,

Aubin & Simonis, 2008). We visually sorted sound samples thanks to spectrograms and discarded

samples with noise from anthropogenic sources, rain, or wind, which can be recognized as very low

frequency noise on the spectrogram. We also discarded samples with noise of very high frequency

corresponding to cicada chirps. We then randomly selected one sound sample per site and per day,

with the exception of four sites for which the four samples only covered two to three days. In total, we

selected 188 samples of 10 min (i.e., 4 samples per site).

Figure 3: Methodological pathway used to identify bird species (in light green) and calculate acoustic indices (in dark green) from automated recordings (see text for details)

Bird species identification – We distributed the samples among 21 expert ornithologists. Each expert performing aural bird species identifications from 4 (one site) to 52 samples (13 sites), primarily from her/his region of residence, for auditory acoustic detection of bird species. We established a presence/absence Site × Species matrix, from which we calculated species richness and functional diversity. Note that bird species abundance can not be reliably assessed with this approach.

Functional diversity – We defined 25 bird species as candidate insectivores for attacking plasticine caterpillars (Table S4.3 in Appendix S4) with those bird species meeting the following criteria: be insectivorous during the breeding season or likely to feed their offspring with insects, forage primarily in forested habitats, and are likely to use substrates such as lower branches or lower leaves of trees where caterpillars were attached to find their prey (Barbaro et al., 2021; Brambilla & Gatti, 2022). We calculated the functional diversity of these candidate insectivores by combining morphological, reproductive, behavioral and acoustic traits.

With the exception of acoustic traits, we extracted functional traits from different published sources,

listed in Table S4.4 in Appendix S4. Specifically, we used three continuous traits: *body mass*, *mean*

clutch size and *bill culmen length* (see Fig. 2 in Tobias et al., 2022) combined with four categorical traits:

foraging method (predominantly understory gleaner, ground gleaner, canopy gleaner), *diet*

(insectivores or mixed diet), *nest type* (open in shrub, open on ground, cavity or open in tree) and

migration (short migration, long migration or resident).

We derived acoustic traits calculations from the work of Krishnan & Tamma (2016). We first extracted

five pure recordings without sonic background for each of the 25 candidate insectivore species from

the online database Xeno-canto.org (Vellinga & Planque, 2015). We then calculated the *number of*

peaks (i.e., NPIC) in the audio signal (see § Acoustic diversity, above) as well as the *frequency of the*

maximum amplitude peaks for each vocal element using the *seewave* library (Sueur, Aubin & Simonis,

2008) and averaged these frequencies for each species. Being based on song and call frequency and

complexity, these indices inform on the adaptation of the vocal repertoire of these species to their

environment.

We summarized the information conveyed by the 9 traits categories into five indices representing

complementary dimensions of the functional diversity (FD) of a community (Mouillot, Graham,

Villéger, Mason & Bellwood, 2013): functional richness (FRic, i.e., convex hull volume of the functional

trait space summarized by a principal coordinates analysis), functional evenness (FEve, i.e., minimum

spanning tree measuring the regularity of trait abundance distribution within the functional space),

and functional divergence (FDiv, i.e., trait abundance distribution within the functional trait space

volume) (Villéger, Mason & Mouillot, 2008), as well as Rao's quadratic entropy (RaoQ, i.e., species

dispersion from the functional centroïd) (Botta-Dukát, 2005). These were calculated for each site with

the *dbFD* function of the *FD* library (Laliberté, Legendre & Shipley, 2014) in the R environment.

Environmental data

Environmental data refer to local temperature and forest cover. We used the high 10-m resolution GIS layers from the Copernicus open platform (Cover, 2018) to calculate forest cover for all European sites. We manually calculated the percentage of forest cover for the two sites located outside Europe using the "World imagery" layer of Arcgis ver. 10.2.3552. We calculated both the percentage of forest cover in a 20-m (henceforth called *local* forest cover) and 200-m (*landscape* forest cover) buffer around the sampled oaks. We chose two nested buffer sizes to better capture the complexity of habitat structure on the diversity and acoustic activity of birds. Local forest cover is particularly important for estimating bird occurrence probability (Melles, Glenn & Martin, 2003), whereas landscape forest cover is an important predictor of bird community composition in urban areas (Rega-Brodsky & Nilon, 2017). Moreover, both local and landscape habitat factors shape insect prey distribution (Barr, van Dijk, Hylander & Tack, 2021). Preliminary analyses revealed that results were qualitatively the same using 10-, 20- or 50-m buffers as predictors of local forest cover and 200- or 500-m buffers as predictors of landscape forest cover (see Table S5.5 in Appendix S5). Because other variables were defined at the site level, we averaged the percentage of forest cover for the sampled trees per site and per buffer size.

We extracted the mean annual temperature at each site from the WorldClim database (the spatial resolution is ~86 km^2, Hijmans et al., 2005).

Statistical analyses

We analyzed 14 response variables in separate linear models (LMs) (Table S3.2 in Appendix S3): bird predation attempts, species richness of the entire bird community and that of candidate insectivores, functional diversity (each of the 4 indices) and acoustic diversity (each of the 7 indices). For each response variable, we first built a full model including a minimum of two components: geography and habitat. The general model equation was (Eq. 1):

$Y_i = \beta_0 + \beta_1 \times Forest_{20,i} + \beta_2 \times Forest_{200,i} + \beta_3 \times Geography_i + \varepsilon_i$ (1)

where Y is the response variable, β_0 the model intercept, β_{is} model coefficient parameters, $Forest_{20}$ and

$Forest_{200}$ the effects of the local and landscape forest cover respectively, *Geography* the effect of mean

annual temperature or latitude (since the two variables are correlated, they were never tested

together in the same models) and ε the residuals.

When modeling the response of bird predation attempts (Eq. 2), we added two more variables to the

model, being any of the seven acoustic diversity indices (*Acoustic diversity*, Eq. 2) and the species

richness or any of the four indices describing the functional diversity of candidate insectivores (*Bird*

diversity, Eq. 2):

$Y_i = \beta_0 + \beta_1 \times Forest_{20,i} + \beta_2 \times Forest_{200,i} + \beta_3 \times Geography_i +$

$\beta_4 \times Bird\ diversity_i + \beta_5 \times Acoustic\ diversity_i + \varepsilon_i$ (2)

We used logarithmic transformations (for bird predation attempts and acoustic entropy (H) models)

or square rooted transformation (for species richness of the complete bird community) of some

response variables where appropriate to satisfy model assumptions. We scaled and centered every

continuous predictor prior to modeling to facilitate comparisons of their effect sizes, and made sure

that none of the explanatory variables were strongly correlated using the variance inflation factor (VIF)

(all VIFs < 5, the usual cutoff values used to check for multicollinearity issues (Miles, 2014)).

For each response variable, we ran the full model as well as every model nested within the full model

and then used Akaike's Information Criterion corrected for small sample size (AICc) to identify the most

parsimonious model(s) fitting the data the best. We simultaneously selected the best variable

describing the geography, habitat, diversity and acoustic component (variable selection) and the best

set of variables describing the variability of the response variable (model selection).

First, we ranked each model according to the difference in AICc between the given model and the

model with the lowest AICc (ΔAICc). Models within 2 ΔAICc units of the best model (i.e., the model

with the lowest AICc) are generally considered as likely (Burnham & Anderson, 2002). We computed

AICc weights for each model (w_i). w_i is interpreted as the probability of a given model being the best

model among the set of candidate models. Eventually, we calculated the relative variable importance

(RVI) as the sum of w_i of every model including this variable, which corresponds to the probability a

variable is included in the best model.

When several models competed with the best model (i.e., when multiple models were such that their

ΔAICc < 2), we applied a procedure of multimodel inference, building a consensus model including the

variables in the set of best models. We then averaged their effect sizes across all the models in the set

of best models, using the variable weight as a weighting parameter (i.e., model averaging). We

considered that a given predictor had a statistically significant effect on the response variable when its

confidence interval excluded zero.

Finally, we used confirmatory structural equation modeling (SEM) (Lefcheck, 2016) to infer direct and

indirect causality links between predictors and response variables. We first built a theoretical model

linking the temperature and forest cover with bird acoustic and functional diversity, and the latter with

bird predation attempts. We then ran a second model also including the direct paths between

temperature, landscape forest cover and bird predation attempts and compared models with *vs*

without direct paths between temperature or landscape forest cover and bird predation attempts. To

avoid model over-parameterization, we used the results of the model selection procedure described

above to identify which variable should represent the habitat, diversity and acoustic components. We

tested independence claims between temperature or forest cover and avian predation rates with *d-*

separation, and we evaluated the overall goodness of fit of the model with Fisher's C statistic.

425 We run all analyses in the R language environment (R Core Team, 2020) with libraries *MuMIn* (Bartoń,

426 2020), *lme4* (Bates, Mächler, Bolker & Walker, 2015), and *piecewiseSEM* (Lefcheck, 2020).

Results

Bird acoustic diversity

430 Of the seven acoustic diversity indices (see Fig. S6.2 in Appendix S6 for correlation between indices),

431 only Acoustic Diversity Index (ADI) and acoustic entropy (H) were significantly associated with any of

432 the predictors tested, i.e., temperature or latitude, local forest cover and landscape forest cover (Table

433 S3.2 in Appendix S3). ADI and H both increased with local forest cover (i.e., percentage of forest cover

434 in a 20-m buffer around recorders). Landscape-scale forest cover (i.e., percentage of forest cover in a

435 200-m buffer around recorders) was the only other predictor retained in the set of competing models

436 in a range of ΔAICc < 2 to explain acoustic entropy variation, but this predictor had little importance

437 (RVI < 0.5) and its effect was not statistically significant (Fig. 5b; Table S3.2 in Appendix S3).

Bird species richness and functional diversity

440 We identified a total of 87 bird species, among which 25 were classified as candidate functional

441 insectivores. Bird species richness varied from 8 to 23 species per recording site (mean ± SD: 15.2 ±

442 3.7, n = 47 sites) and richness of candidate insectivores from 2 to 9 species (5.7 ± 1.5). The null model

443 was among models competing in a range of ΔAICc < 2 for both total species richness and candidate

444 insectivores (Table S3.2 in Appendix S3).

445 Among the five bird functional diversity and species richness indices, only functional quadratic entropy

446 (Rao's Q) characterizing species dispersion from the functional centroid was significantly influenced by

447 the predictors tested (temperature or latitude, local and landscape forest cover, Table S3.2 in Appendix

448 S3). Specifically, Rao's Q decreased towards higher latitudes (Fig. 4a and Fig. 5c). Other predictors

449   retained in the set of competing models in a range of ΔAICc < 2 had little importance (RVI < 0.5) and

450   were not significant (Fig. 5c; Table S3.2 in Appendix S3).

453   *Figure 4*: *Scatter diagrams showing changes in (a) Rao's Q and (b) predation attempts with latitude. These relationships were*

454   *identified as significant in the linear models tested*

Figure 5: *Effects of geography (temperature or latitude) and habitat (percentage of forest cover at 20 or 200 m) on Acoustic*

Diversity Index (ADI) (a), Acoustic Entropy Index (H) (b), Rao's quadratic entropy (RaoQ) (c) and bird predation attempts (d)

and effects of acoustic (Bioacoustic Index), bird diversity (Functional Richness) on bird predation attempts (d). Circles and error

bars represent standardized parameter estimates and corresponding 95% confidence intervals (CI), respectively. The vertical

dashed line centered on zero represents the null hypothesis. Full and empty circles represent significant and non-significant

effect sizes, respectively. Circle size is proportional to RVI. The colors correspond to the different components tested (acoustic,

bird diversity, geography and habitat).

Bird predation attempts

Of the 4,860 exposed dummy caterpillars, 22.8% (*n* = 1,108) had bird bill marks. Model selection

retained two models in the set of competing models in a range of ΔAICc < 2 (Table S3.2 in Appendix

S3). Landscape forest cover (RVI = 1.00), bird functional richness (FRic) (RVI = 1.00) and bioacoustic

index (BI) (RVI = 1.00) were selected in all models. Latitude (RVI = 0.73) and temperature (RVI = 0.27)

were also each selected in one of the two best models.

Bird predation attempts increased towards higher latitudes and consistently decreased with increasing

mean annual temperature. Bird predation attempts further increased with bioacoustic index (BI), but

decreased with bird functional richness (FRic) (Fig. 4b and Fig. 5d).

The results were comparable when we incorporated latitudinal changes in diel phenology in the

calculation of predation attempts through the standardization with the daylight duration (see Table

S3.2 in Appendix S3). The only noticeable difference was that the importance of temperature as a

predictor was higher (RVI=1.00, compared with RVI = 0.27 for the first model with unstandardized rates

of predation attempts).

A first SEM was set up by specifying direct links between environmental variables (latitude and

landscape forest cover) and bird diversity variables (FRic and BI) on the one hand and between bird

diversity variables and predation attempts on the other. The model was however misspecified (C = 27,

P < 0.001, AIC = 51). We ran a second model also including the direct paths between latitude, landscape

forest cover and bird predation attempts. Model AIC dropped dramatically (ΔAIC = 19), indicating that

the full model with direct paths between latitude, forest cover and bird predation attempts had a

better fit (C = 4, P = 0.131, AIC = 32). Latitude, landscape forest cover and bioacoustic index had a

significant positive effect on bird predation attempts whereas FRic had a significant negative effect on

bird predation attempts. Altogether, these results indicate that latitude and forest cover had direct

effects on bird predation attempts that were not mediated by local changes in bird diversity (Fig. 6).

Figure 6. *Structural equation model. The thickness of the arrows is proportional to coefficient parameter estimate.*

*Standardized estimates for each response variable are indicated next to each arrow. *P < 0.05, **P < 0.01, ***P < 0.001.*

Continuous and dashed arrows represent significant (P < 0.05) and non-significant (P > 0.05) relationships respectively. Black

and gray arrows represent positive and negative relationships respectively. FRic refers to the functional richness of candidate

insectivorous birds.

Discussion

Bird predation and functional diversity of bird insectivores are both influenced by latitude

The Latitudinal Biotic Interactions Hypothesis (Dobzhansky, 1950) is a controversial idea, particularly

for biotic interactions between herbivores and higher trophic levels (Lövei & Ferrante, 2017; Zvereva

& Kozlov, 2021). Against this hypothesis and contrarily with our prediction ((i), Fig. 1), we found that

latitude was positively related to bird predation attempts. More bird predation attempts at higher

latitudes could be due to longer daylight duration in spring northwards, leading insectivorous birds to

have more time per day to find their prey and thus allowing high coexistence of predators during a

period of unlimited resource availability (Speakman et al., 2000). Alternatively, birds needing more

energy in colder temperatures to thermoregulate may need to feed more to maintain their metabolic

activity (Caraco et al., 1990; Kendeigh, 1969; Steen, 1958; Wansink & Tinbergen, 1994). Consistently

with this explanation, we found a negative relationship between temperature and bird predation

attempts. Moreover, temperature remained an important, significant predictor of bird predation

attempts when we controlled for the duration of daylight (Table S3.2 in Appendix S3), which further

supports this explanation. However, we cannot exclude the possibility that the lower predation rates

at lower latitudes was due to lower prey detectability.

In agreement with our prediction (i, Fig. 1), we provide evidence for a significant negative relationship

between latitude and the functional diversity of insectivorous birds. Despite substantial differences

among functional diversity indices, this suggests that more functionally diverse assemblages of

insectivorous birds are able to coexist locally in oak woods towards the south of Europe (Currie et al.,

2004; Hillebrand, 2004; Willig et al., 2003). Of the multiple functional diversity indices commonly used

to describe ecological communities, it is noticeable that only the quadratic entropy index responded

negatively to latitude, for it is a synthetic index which simultaneously takes into account the richness,

evenness, and divergence components of functional diversity (Mouillot et al., 2013).

Bird predation attempts are partly predicted by bird functional and acoustic diversity

We predicted that bird predation attempts would increase with bird functional diversity (ii, Fig. 1), but

found no or opposite effects, depending on the functional index considered. Only functional

insectivore richness was negatively correlated to predation rates. Negative relationships between

predation and predator functional diversity can arise from a combination of both intraguild predation

--- predators preying upon predators (Mooney et al., 2010) --- and intraguild competition (Houska

Tahadlova et al., 2022), although we can hardly tease them apart in the present study. An important

step forward would consist in testing whether predation patterns revealed with artificial prey are

representative of predation intensity as a whole (Zvereva & Kozlov, 2021). For example, functional

richness may be a proxy for dietary specialization such that functionally more diverse predator

communities would seek more prey of which they are specialists and thus predate less on artificial

caterpillars. It is possible that a higher diversity of insectivorous birds in warmer regions was linked to

higher diversity and abundance of arthropod prey (Kissling et al., 2012) and therefore to greater prey

availability (Charbonnier et al., 2016). If so, then the pattern we observed may merely be

representative of the 'dilution' of bird attacks on artificial prey among more abundant and diverse real

prey (Zeuss, Brunzel & Brandl, 2017; Zvereva et al., 2019). A follow-up of the present study should

therefore pay special attention on the real prey density pre-existing in each sampling site where

artificial prey are to be deployed as a standardized measure of predation rates across sites.

Bird predation attempts increased with the Bioacoustic Index (BI), which is regarded as a good proxy

for bird abundance (Sueur et al., 2014). This finding is thus consistent with previous studies having

reported positive relationships between bird abundance and predation attempts on artificial prey

(Roels, Porter & Lindell, 2018; Sam, Koane & Novotny, 2015). Acoustic diversity indices are

complementary to functional diversity indices and reflect both the composition of the avian

community and the nature of the habitat (Gasc, Francomano, Dunning & Pijanowski, 2017; Sugai, Silva,

Ribeiro & Llusia, 2019). We further show that the acoustic monitoring of the habitat could yield proxies

for the intensity of bird predation on invertebrates. In addition, our study is to our knowledge among

the first to detect significant relationships between acoustic indices and an ecosystem function

measured independently (Gasc et al., 2013), thus opening pathways for new research on the link

between functional and acoustic ecology.

Local forest cover predicts bird acoustic activity, whereas landscape forest cover increases bird

predation

Acoustic diversity increased with closeness of canopy cover in the immediate neighborhood (20m

radius) of sampled trees (iii, Fig. 1). The most responsive indices were the acoustic diversity (ADI) and

the acoustic entropy (H). Both are especially suitable at predicting acoustic diversity across different

habitats under various ambient sound conditions (Fuller, Axel, Tucker & Gage, 2015; Machado, Aguiar

& Jones, 2017). The former is related to a greater regularity of the soundscape and the latter is related

to the amplitude between frequency bands and time. They therefore correspond to soundscapes

containing multiple vocalizing species (Sueur et al., 2008; Villanueva-Rivera, Pijanowski, Doucette &

Pekin, 2011). Acoustic entropy is also known to respond significantly to local forest habitat (Barbaro et

al., 2022), which is generally a good predictor of bird occupancy probability (Morante-Filho, Benchimol

& Faria, 2021).

Bird predation attempts were best predicted by forest cover at the landscape level (Prediction (iii), Fig.

1). Indeed, it is likely that forest cover at the landscape level provides structural complexity with a

dense understorey and habitat heterogeneity that is both a source of food and niches for predatory

birds to exploit (Poch & Simonetti, 2013). As a result, forest cover at the landscape scale is often a key

predictor of avian insectivory in various study areas (Barbaro et al., 2014; González-Gómez et al., 2006;

Valdés-Correcher et al., 2021). This is also consistent with the results of Rega-Brodsky & Nilon, (2017)

who found greater abundance of insectivorous birds in mosaic urban or rural landscapes including a

significant part of semi-natural wooded habitats, such as those we studied here.

Latitudinal variation in avian predation is independent from large-scale changes in the diversity and

acoustic activity of birds

We found no evidence that the relationship between latitude and bird predation attempts was

mediated by changes in bird diversity or activity (iv, Fig. 1). On the contrary, geography and bird

diversity had independent and complementary effects on predation.

Latitude may directly drive both bird activity and abundance according to available resources

(Pennings & Silliman, 2005). Even changes in the abundance of a single, particularly active, predator

species along the latitudinal gradient to explain the observed pattern (Maas et al., 2015; Philpott et

al., 2009). For example, the blue tit *Cyanistes caeruleus* and the great tit *Parus major* are typical and

widespread canopy insectivores of European oak forests and are particularly prone to predate

herbivorous caterpillars while showing considerable adaptive behaviour to prey availability (Mols &

Visser, 2002; Naef-Daenzer & Keller, 1999). If the predation attempts on the plasticine caterpillars were

to be predominantly due to these species, then it would be their abundance and activity that would

play a role in predation attempts rather than the overall diversity of insectivores (Maas et al., 2015).

Here, we based our assessment of functional bird composition on candidate insectivore occurrences

obtained from standardized acoustic surveys, which on the one hand insures that we have no observer,

site, or phenological biases on species occurrences, but on the other hand also makes it difficult to

precisely account for each species' abundance. Other complementary methods to assess the relative

roles of each individual bird species on predation rates should be deployed further to better account

for actual predatory bird abundance and activity, including DNA sampling (Garfinkel, Minor & Whelan,

2022), camera traps (Martínez-Núñez et al., 2021) or species-specific bird surveys involving tape calls

or capture methods.

Conclusion

We found a negative association between latitude and bird functional diversity, but at the same time,

a positive relationship between latitude and avian predation. Our study therefore provides partial

support for the latitudinal clines in biodiversity hypothesis, but clearly conflicts with the Latitudinal

Biotic Interaction Hypothesis (Dobzhansky, 1950). As cross-continental studies exploring the large-

scale relationship between latitude and the strength of biotic interactions generally ignore local

factors, we argue that characterizing the contrasting habitats of the study sites is a good way to

circumvent some inconsistencies in the results. We identify pre-existing real prey density and single

key bird species abundances as two particularly important variables deserving further attention.

Furthermore, predicting ecosystem services — here, potential pest regulation service — on a large

scale by standardized proxies such as acoustic ecology for predator diversity and plasticine caterpillars

for predation function seem to be good ways to reduce methodological biases and strengthen our

understanding of the macro-ecology of biotic interactions.

Data availability statement

For the moment there is an embargo on data and codes which will be lifted after open acceptance.

Schille, Laura; Valdés-Correcher, Elena; Archaux, Frédéric; Bălăcenoiu, Flavius; Bjørn, Mona Chor;

Bogdziewicz, Michal; Boivin, Thomas; Branco, Manuela; Damestoy, Thomas; de Groot, Maarten;

Dobrosavljević, Jovan; Duduman, Mihai-Leonard; Dulaurent, Anne-Maïmiti; Green, Samantha;

Grünwald, Jan; Eötvös, Csaba Béla; Faticov, Maria; Fernandez-Conradi, Pilar; Flury, Elisabeth; Funosas,

David; Galmán, Andrea; Gossner, Martin; Gripenberg, Sofia; Grosu, Lucian; Hagge, Jonas; Hampe,

Arndt; Harvey, Deborah; Houston, Rick; Isenmann, Rita; Kavčič, Andreja; Kozlov, Mikhail V.; Lanta,

Vojtech; Le Tilly, Bénédicte; Lopez Vaamonde, Carlos; Mallick, Soumen; Mäntylä, Elina; Mårell, Anders;

Milanović, Slobodan; Molnár, Márton; Moreira, Xoaquín; Moser, Valentin; Mrazova, Anna; Musolin,

Dmitrii L.; Perot, Thomas; Piotti, Andrea; Popova, Anna V.; Prinzing, Andreas; Pukinskaya, Ludmila;

Sallé, Aurélien; Sam, Katerina; Sedikhin, Nickolay V.; Shabarova, Tanja; Tack, Ayco; Thomas, Rebecca;

Thrikkadeeri, Karthik; Toma, Dragoș; Vaicaityte, Grete; van Halder, Inge; Varela, Zulema; Barbaro, Luc;

Castagneyrol, Bastien, 2023, « Data and codes for the article "Latitudinal gradient in avian insectivory:

complementary effects of climate, habitat and bird diversity" », https://doi.org/10.57745/0E0JEA,

Recherche Data Gouv.

Copyright statement

document and will be applied to all subsequent versions up to the Author Accepted Manuscript arising

from this submission.

References

Anstett, D. N., Chen, W., & Johnson, M. T. J. (2016). Latitudinal Gradients in Induced and Constitutive

Resistance against Herbivores. *Journal of Chemical Ecology*, *42*(8), 772-781.

https://doi.org/10.1007/s10886-016-0735-6

Bael, S. A. V., Philpott, S. M., Greenberg, R., Bichier, P., Barber, N. A., Mooney, K. A., & Gruner, D. S.

(2008). Birds as predators in tropical agroforestry systems. *Ecology*, *89*(4), 928-934.

https://doi.org/10.1890/06-1976.1

Barbaro, L., Allan, E., Ampoorter, E., Castagneyrol, B., Charbonnier, Y., De Wandeler, H., Kerbiriou, C.,

Milligan, H. T., Vialatte, A., Carnol, M., Deconchat, M., De Smedt, P., Jactel, H., Koricheva, J.,

Le Viol, I., Muys, B., Scherer-Lorenzen, M., Verheyen, K., & van der Plas, F. (2019). Biotic

predictors complement models of bat and bird responses to climate and tree diversity in

European forests. *Proceedings of the Royal Society B: Biological Sciences*, *286*(1894),

20182193. https://doi.org/10.1098/rspb.2018.2193

Barbaro, L., Assandri, G., Brambilla, M., Castagneyrol, B., Froidevaux, J., Giffard, B., Pithon, J., Puig-

Montserrat, X., Torre, I., Calatayud, F., Gaüzère, P., Guenser, J., Macià-Valverde, F., Mary, S.,

Raison, L., Sirami, C., & Rusch, A. (2021). Organic management and landscape heterogeneity

combine to sustain multifunctional bird communities in European vineyards. *Journal of

Applied Ecology*, *58*(6), 1261-1271. https://doi.org/10.1111/1365-2664.13885

Barbaro, L., Giffard, B., Charbonnier, Y., van Halder, I., & Brockerhoff, E. G. (2014). Bird functional

diversity enhances insectivory at forest edges : A transcontinental experiment. *Diversity and

Distributions*, *20*(2), 149-159. https://doi.org/10.1111/ddi.12132

Barbaro, L., Sourdril, A., Froidevaux, J. S. P., Cauchoix, M., Calatayud, F., Deconchat, M., & Gasc, A.

(2022). Linking acoustic diversity to compositional and configurational heterogeneity in

mosaic landscapes. *Landscape Ecology*, *37*(4), 1125-1143. https://doi.org/10.1007/s10980-

021-01391-8

Barr, A. E., van Dijk, L. J. A., Hylander, K., & Tack, A. J. M. (2021). Local habitat factors and spatial

connectivity jointly shape an urban insect community. *Landscape and Urban Planning*, *214*,

104177. https://doi.org/10.1016/j.landurbplan.2021.104177

Bartoń, K. (2020). *MuMIn : Multi-model inference.* (R package version 1.43.17.) [Computer software].

https://CRAN.Rproj ect.org/packa ge=MuMIn

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using

lme4. *Journal of Statistical Software*, *67*(1). https://doi.org/10.18637/jss.v067.i01

Blackburn, T. M., & Gaston, K. J. (1996). Spatial patterns in the species richness of birds in the New

World. *Ecography*, *19*(4), 369-376.

Botta-Dukát, Z. (2005). Rao's quadratic entropy as a measure of functional diversity based on

multiple traits. *Journal of vegetation science*, *16*(5), 533-540.

Bradfer-Lawrence, T., Gardner, N., Bunnefeld, L., Bunnefeld, N., Willis, S. G., & Dent, D. H. (2019).

Guidelines for the use of acoustic indices in environmental research. *Methods in Ecology and*

Evolution, *10*(10), 1796-1807.

Brambilla, M., & Gatti, F. (2022). No more silent (and uncoloured) springs in vineyards? Experimental

evidence for positive impact of alternate inter-row management on birds and butterflies.

Journal of Applied Ecology, 1365-2664.14229. https://doi.org/10.1111/1365-2664.14229

Burnham, K. P., & Anderson, D. R. (2002). *Model Selection and Multimodel Inference : A Practical*

Information Theoretic Approach, 2nd edn. (Springer).

Caraco, T., Blanckenhorn, W. U., Gregory, G. M., Newman, J. A., Recer, G. M., & Zwicker, S. M. (1990).

Risk-sensitivity : Ambient temperature affects foraging choice. *Animal Behaviour*, *39*(2),

338-345. https://doi.org/10.1016/S0003-3472(05)80879-6

Charbonnier, Y. M., Barbaro, L., Barnagaud, J.-Y., Ampoorter, E., Nezan, J., Verheyen, K., & Jactel, H.

(2016). Bat and bird diversity along independent gradients of latitude and tree composition

in European forests. *Oecologia*, *182*(2), 529-537. https://doi.org/10.1007/s00442-016-3671-9

Cover, C. L. (2018). *European Union, Copernicus Land Monitoring Service.* European Environment

Agency.

Currie, D. J., Mittelbach, G. G., Cornell, H. V., Field, R., Guégan, J.-F., Hawkins, B. A., Kaufman, D. M.,

Kerr, J. T., Oberdorff, T., & O'Brien, E. (2004). Predictions and tests of climate-based

hypotheses of broad-scale variation in taxonomic richness. *Ecology letters*, *7*(12), 1121-1134.

De la Mora, A., García-Ballinas, J. A., & Philpott, S. M. (2015). Local, landscape, and diversity drivers

of predation services provided by ants in a coffee landscape in Chiapas, Mexico. *Agriculture,*

Ecosystems & Environment, *201*, 83-91. https://doi.org/10.1016/j.agee.2014.11.006

Dobzhansky, T. (1950). Evolution in the tropics. *American scientist*, *38*(2), 209-221.

Dröge, S., Martin, D. A., Andriafanomezantsoa, R., Burivalova, Z., Fulgence, T. R., Osen, K.,

Rakotomalala, E., Schwab, D., Wurz, A., Richter, T., & Kreft, H. (2021). Listening to a changing

landscape : Acoustic indices reflect bird species richness and plot-scale vegetation structure

across different land-use types in north-eastern Madagascar. *Ecological Indicators*, *120*,

106929. https://doi.org/10.1016/j.ecolind.2020.106929

Eaton, E., Caudullo, G., Oliveira, S., & de Rigo, D. (2016). *Quercus robur and Quercus petraea*. 4.

Fuller, S., Axel, A. C., Tucker, D., & Gage, S. H. (2015). Connecting soundscape to landscape : Which

acoustic index best describes landscape configuration? *Ecological Indicators*, *58*, 207-215.

https://doi.org/10.1016/j.ecolind.2015.05.057

Garfinkel, M., Minor, E., & Whelan, C. J. (2022). Using faecal metabarcoding to examine consumption

of crop pests and beneficial arthropods in communities of generalist avian insectivores. *Ibis*,

164(1), 27-43. https://doi.org/10.1111/ibi.12994

Gasc, A., Francomano, D., Dunning, J. B., & Pijanowski, B. C. (2017). Future directions for soundscape

ecology : The importance of ornithological contributions. *The Auk*, *134*(1), 215-228.

https://doi.org/10.1642/AUK-16-124.1

Gasc, A., Sueur, J., Jiguet, F., Devictor, V., Grandcolas, P., Burrow, C., Depraetere, M., & Pavoine, S.

(2013). Assessing biodiversity with sound : Do acoustic diversity indices reflect phylogenetic

and functional diversities of bird communities? *Ecological Indicators*, *25*, 279-287.

https://doi.org/10.1016/j.ecolind.2012.10.009

Gibb, R., Browning, E., Glover-Kapfer, P., & Jones, K. E. (2019). Emerging opportunities and challenges

for passive acoustics in ecological assessment and monitoring. *Methods in Ecology and*

Evolution, *10*(2), 169-185. https://doi.org/10.1111/2041-210X.13101

González-Gómez, P. L., Estades, C. F., & Simonetti, J. A. (2006). Strengthened insectivory in a

temperate fragmented forest. *Oecologia*, *148*(1), 137-143. https://doi.org/10.1007/s00442-

005-0338-3

Greenop, A., Woodcock, B. A., Wilby, A., Cook, S. M., & Pywell, R. F. (2018). Functional diversity

positively affects prey suppression by invertebrate predators : A meta-analysis. *Ecology*,

99(8), 1771-1782. https://doi.org/10.1002/ecy.2378

Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high resolution

interpolated climate surfaces for global land areas. *International Journal of Climatology*,

25(15), 1965-1978. https://doi.org/10.1002/joc.1276

Hill, A. P., Prince, P., Piña Covarrubias, E., Doncaster, C. P., Snaddon, J. L., & Rogers, A. (2018).

AudioMoth : Evaluation of a smart open acoustic device for monitoring biodiversity and the

environment. *Methods in Ecology and Evolution*, *9*(5), 1199-1211.

https://doi.org/10.1111/2041-210X.12955

Hillebrand, H. (2004). On the Generality of the Latitudinal Diversity Gradient. *The American*

Naturalist, *163*(2), 192-211. https://doi.org/10.1086/381004

Houska Tahadlova, M., Mottl, O., Jorge, L. R., Koane, B., Novotny, V., & Sam, K. (2022). Trophic

cascades in tropical rainforests : Effects of vertebrate predator exclusion on arthropods and

plants in Papua New Guinea. *Biotropica*.

Ikin, K., Barton, P. S., Stirnemann, I. A., Stein, J. R., Michael, D., Crane, M., Okada, S., & Lindenmayer,

D. B. (2014). Multi-Scale Associations between Vegetation Cover and Woodland Bird

Communities across a Large Agricultural Region. *PLoS ONE*, *9*(5), e97029.

https://doi.org/10.1371/journal.pone.0097029

Just, M. G., Dale, A. G., Long, L. C., & Frank, S. D. (2019). Urbanization drives unique latitudinal

patterns of insect herbivory and tree condition. *Oikos*, *128*(7), 984-993.

https://doi.org/10.1111/oik.05874

Kendeigh, S. C. (1969). *Energy Responses of Birds to Their Thermal Environments*. *81*(4), 10.

Kissling, W. D., Sekercioglu, C. H., & Jetz, W. (2012). Bird dietary guild richness across latitudes,

environments and biogeographic regions : Guild richness in birds. *Global Ecology and*

Biogeography, *21*(3), 328-340. https://doi.org/10.1111/j.1466-8238.2011.00679.x

Kleijn, D., Rundlöf, M., Scheper, J., Smith, H. G., & Tscharntke, T. (2011). Does conservation on

farmland contribute to halting the biodiversity decline? *Trends in Ecology & Evolution*, *26*(9),

474-481. https://doi.org/10.1016/j.tree.2011.05.009

Kozlov, M. V., Stekolshchikov, A. V., Söderman, G., Labina, E. S., Zverev, V., & Zvereva, E. L. (2015).

Sap-feeding insects on forest trees along latitudinal gradients in northern Europe : A climate-

driven patterns. *Global Change Biology*, *21*(1), 106-116. https://doi.org/10.1111/gcb.12682

Krishnan, A., & Tamma, K. (2016). Divergent morphological and acoustic traits in sympatric

communities of Asian barbets. *Royal Society Open Science*, *3*(8), 160117.

https://doi.org/10.1098/rsos.160117

Laliberté, E., Legendre, P., & Shipley, B. (2014). *Measuring Functional Diversity (FD) from Multiple*

Traits, and Other Tools for Functional Ecology : Package « FD » (1.0-12.1) [Computer

software].

Lee, M., Fahrig, L., Freemark, K., & Currie, D. J. (2002). Importance of patch scale vs landscape scale

on selected forest birds. *Oikos*, *96*(1), 110-118. https://doi.org/10.1034/j.1600-

0706.2002.960112.x

Lefcheck, J. (2020). *Piecewise Structural Equation Modeling* (package « piecewiseSEM » version 2.1.2)

[Computer software].

Lefcheck, J. S. (2016). PIECEWISESEM: Piecewise structural equationmodelling in R for ecology,

evolution, and systematics. *Methods in Ecology and Evolution*, *7*(5), 573-579.

https://doi.org/10.1111/2041-210X.12512

Long, L. C., & Frank, S. D. (2020). Risk of bird predation and defoliating insect abundance are greater

in urban forest fragments than street trees. *Urban Ecosystems*, *23*(3), 519-531.

https://doi.org/10.1007/s11252-020-00939-x

Loreau, M., Naeem, S., Inchausti, P., Bengtsson, J., Grime, J. P., Hector, A., Hooper, D. U., Huston, M.

A., Raffaelli, D., Schmid, B., Tilman, D., & Wardle, D. A. (2001). Biodiversity and Ecosystem

Functioning : Current Knowledge and Future Challenges. *Science*, *294*(5543), 804-808.

https://doi.org/10.1126/science.1064088

Lövei, G. L., & Ferrante, M. (2017). A review of the sentinel prey method as a way of quantifying

invertebrate predation under field conditions : Measuring predation pressure by sentinel

prey. *Insect Science*, *24*(4), 528-542. https://doi.org/10.1111/1744-7917.12405

Low, P. A., Sam, K., McArthur, C., Posa, M. R. C., & Hochuli, D. F. (2014). Determining predator

identity from attack marks left in model caterpillars : Guidelines for best practice.

Entomologia Experimentalis et Applicata, *152*(2), 120-126.

https://doi.org/10.1111/eea.12207

Maas, B., Tscharntke, T., Saleh, S., Dwi Putra, D., & Clough, Y. (2015). Avian species identity drives

predation success in tropical cacao agroforestry. *Journal of Applied Ecology*, *52*(3), 735-743.

https://doi.org/10.1111/1365-2664.12409

MacArthur, R. H. (1984). *Geographical ecology : Patterns in the distribution of species*. Princeton

University Press.

Machado, R. B., Aguiar, L., & Jones, G. (2017). Do acoustic indices reflect the characteristics of bird

communities in the savannas of Central Brazil? *Landscape and Urban Planning*, *162*, 36-43.

https://doi.org/10.1016/j.landurbplan.2017.01.014

Martínez-Núñez, C., Rey, P. J., Manzaneda, A. J., García, D., Tarifa, R., & Molina, J. L. (2021).

Insectivorous birds are not effective pest control agents in olive groves. *Basic and Applied*

Ecology, *56*, 270-280. https://doi.org/10.1016/j.baae.2021.08.006

Mazerolle, M. J., & Villard, M.-A. (1999). Patch characteristics and landscape context as predictors of

species presence and abundance : A review [1]. *Écoscience*, *6*(1), 117-124.

https://doi.org/10.1080/11956860.1999.11952204

Melles, S., Glenn, S. M., & Martin, K. (2003). Urban Bird Diversity and Landscape Complexity :

Species-environment Associations Along a Multiscale Habitat Gradient. *Conservation Ecology*,

7(1), art5. https://doi.org/10.5751/ES-00478-070105

Miles, J. (2014). Tolerance and variance inflation factor. *Wiley statsref: statistics reference online*.

Mols, C. M. M., & Visser, M. E. (2002). Great tits can reduce caterpillar damage in apple orchards :

Great tits reduce caterpillar damage. *Journal of Applied Ecology*, *39*(6), 888-899.

https://doi.org/10.1046/j.1365-2664.2002.00761.x

Mooney, K. A., Gruner, D. S., Barber, N. A., Van Bael, S. A., Philpott, S. M., & Greenberg, R. (2010).

Interactions among predators and the cascading effects of vertebrate insectivores on

arthropod communities and plants. *Proceedings of the National Academy of Sciences*,

107(16), 7335-7340. https://doi.org/10.1073/pnas.1001934107

Morante-Filho, J. C., Benchimol, M., & Faria, D. (2021). Landscape composition is the strongest

determinant of bird occupancy patterns in tropical forest patches. *Landscape Ecology*, *36*(1),

105-117. https://doi.org/10.1007/s10980-020-01121-6

Mouillot, D., Graham, N. A. J., Villéger, S., Mason, N. W. H., & Bellwood, D. R. (2013). A functional

approach reveals community responses to disturbances. *Trends in Ecology & Evolution*, *28*(3),

167-177. https://doi.org/10.1016/j.tree.2012.10.004

Naef-Daenzer, B., & Keller, L. F. (1999). The foraging performance of great and blue tits (Parus major

and P. caeruleus) in relation to caterpillar development, and its consequences for nestling

growth and fledging weight. *Journal of Animal Ecology*, *68*(4), 708-718.

Otto, S. B., Berlow, E. L., Rank, N. E., Smiley, J., & Brose, U. (2008). Predator diversity and identity

drive interaction strength and trophic cascades in a food web. *Ecology*, *89*(1), 134-144.

https://doi.org/10.1890/07-0066.1

Pennings, S. C., & Silliman, B. R. (2005). Linking biogeography and community ecology : Latitudinal

variation in plant–herbivore interaction strength. *Ecology*, *86*(9), 2310-2319.

https://doi.org/10.1890/04-1022

Petit, R. J., Brewer, S., Bordács, S., Burg, K., Cheddadi, R., Coart, E., Cottrell, J., Csaikl, U. M., van Dam,

B., & Deans, J. D. (2002). Identification of refugia and post-glacial colonisation routes of

European white oaks based on chloroplast DNA and fossil pollen evidence. *Forest ecology*

and management, *156*(1-3), 49-74.

Philpott, S. M., Soong, O., Lowenstein, J. H., Pulido, A. L., Lopez, D. T., Flynn, D. F. B., & DeClerck, F.

(2009). Functional richness and ecosystem services : Bird predation on arthropods in tropical

agroecosystems. *Ecological Applications*, *19*(7), 1858-1867. https://doi.org/10.1890/08-

1928.1

Poch, T. J., & Simonetti, J. A. (2013). Insectivory in Pinus radiata plantations with different degree of

structural complexity. *Forest Ecology and Management*, *304*, 132-136.

https://doi.org/10.1016/j.foreco.2013.04.044

R Core Team. (2020). *R: A Language and environment for statistical computing.* R Foundation for

Statistical Computing, Vienna, Austria. https://www.r-project.org/

Rega-Brodsky, C. C., & Nilon, C. H. (2017). Forest cover is important across multiple scales for bird

communities in vacant lots. *Urban Ecosystems*, *20*(3), 561-571.

https://doi.org/10.1007/s11252-016-0614-5

Roels, S. M., Porter, J. L., & Lindell, C. A. (2018). Predation pressure by birds and arthropods on

herbivorous insects affected by tropical forest restoration strategy. *Restoration Ecology*,

26(6), 1203-1211. https://doi.org/10.1111/rec.12693

Roslin, T., Hardwick, B., Novotny, V., Petry, W. K., Andrew, N. R., Asmus, A., Barrio, I. C., Basset, Y.,

Boesing, A. L., Bonebrake, T. C., Cameron, E. K., Dáttilo, W., Donoso, D. A., Drozd, P., Gray, C.

L., Hik, D. S., Hill, S. J., Hopkins, T., Huang, S., … Slade, E. M. (2017). Higher predation risk for

insect prey at low latitudes and elevations. *Science*, *356*(6339), 742-744.

https://doi.org/10.1126/science.aaj1631

Sam, K., Koane, B., & Novotny, V. (2015). Herbivore damage increases avian and ant predation of

caterpillars on trees along a complete elevational forest gradient in Papua New Guinea.

Ecography, *38*(3), 293-300.

Sánchez-Giraldo, C., Correa Ayram, C., & Daza, J. M. (2021). Environmental sound as a mirror of

landscape ecological integrity in monitoring programs. *Perspectives in Ecology and*

Conservation, *19*(3), 319-328. https://doi.org/10.1016/j.pecon.2021.04.003

Schemske, D. W., Mittelbach, G. G., Cornell, H. V., Sobel, J. M., & Roy, K. (2009). Is There a Latitudinal

Gradient in the Importance of Biotic Interactions? *Annual Review of Ecology, Evolution, and*

Systematics, *40*(1), 245-269. https://doi.org/10.1146/annurev.ecolsys.39.110707.173430

Sekercioglu, C. (2006). Increasing awareness of avian ecological function. *Trends in Ecology &*

Evolution, *21*(8), 464-471. https://doi.org/10.1016/j.tree.2006.05.007

Sinclair, A. R. E., Mduma, S., & Brashares, J. S. (2003). Patterns of predation in a diverse predator–

prey system. *Nature*, *425*(6955), 288-290. https://doi.org/10.1038/nature01934

Southwood, T. R. E., Wint, G. W., Kennedy, C. E., & Greenwood, S. R. (2005). The composition of the

arthropod fauna of the canopies of some species of oak (Quercus). *European Journal of*

Entomology, *102*(1), 65-72.

Speakman, J. R., Rydell, J., Webb, P. I., Hayes, J. P., Hays, G. C., Hulbert, I. A. R., & McDevitt, R. M.

(2000). Activity patterns of insectivorous bats and birds in northern Scandinavia (69° N),

during continuous midsummer daylight. *Oikos*, *88*(1), 75-86. https://doi.org/10.1034/j.1600-

0706.2000.880109.x

Steen, J. (1958). Climatic Adaptation in Some Small Northern Birds. *Ecology*, *39*(4), 625-629.

https://doi.org/10.2307/1931602

Sueur, J., Aubin, T., & Simonis, C. (2008). Seewave, a free modular tool for sound analysis and

synthesis. *Bioacoustics*, *18*(2), 213-226. https://doi.org/10.1080/09524622.2008.9753600

Sueur, J., Farina, A., Gasc, A., Pieretti, N., & Pavoine, S. (2014). Acoustic Indices for Biodiversity

Assessment and Landscape Investigation. *Acta Acustica United with Acustica*, *100*(4),

772-781. https://doi.org/10.3813/AAA.918757

Sueur, J., Pavoine, S., Hamerlynck, O., & Duvail, S. (2008). Rapid Acoustic Survey for Biodiversity

Appraisal. *PLoS ONE*, *3*(12), e4065. https://doi.org/10.1371/journal.pone.0004065

Sugai, L. S. M., Silva, T. S. F., Ribeiro, J. W., & Llusia, D. (2019). Terrestrial Passive Acoustic

Monitoring : Review and Perspectives. *BioScience*, *69*(1), 15-25.

https://doi.org/10.1093/biosci/biy147

Tobias, J. A., Sheard, C., Pigot, A. L., Devenish, A. J. M., Yang, J., Sayol, F., Neate-Clegg, M. H. C.,

Alioravainen, N., Weeks, T. L., Barber, R. A., Walkden, P. A., MacGregor, H. E. A., Jones, S. E.

I., Vincent, C., Phillips, A. G., Marples, N. M., Montaño-Centellas, F. A., Leandro-Silva, V.,

Claramunt, S., … Schleuning, M. (2022). AVONET : Morphological, ecological and geographical

data for all birds. *Ecology Letters*, *25*(3), 581-597. https://doi.org/10.1111/ele.13898

Valdés-Correcher, E., Moreira, X., Augusto, L., Barbaro, L., Bouget, C., Bouriaud, O., Branco, M.,

Centenaro, G., Csóka, G., Damestoy, T., Dobrosavljević, J., Duduman, M., Dulaurent, A.,

Eötvös, C. B., Faticov, M., Ferrante, M., Fürjes-Mikó, Á., Galmán, A., Gossner, M. M., …

Castagneyrol, B. (2021). Search for top-down and bottom-up drivers of latitudinal trends in

insect herbivory in oak trees in Europe. *Global Ecology and Biogeography*, *30*(3), 651-665.

https://doi.org/10.1111/geb.13244

Vellinga, W.-P., & Planque, R. (2015). *The Xeno-canto collection and its relation to sound recognition*

and classification. 11.

Vickery, J., & Arlettaz, R. (2012). The importance of habitat heterogeneity at multiple scales for birds

in European agricultural landscapes. *Fuller, R.J. (éd) Birds and Habitat : Relationships in*

Changing Landscapes, 177-204.

Villanueva-Rivera, L. J., & Pijanowski, B. C. (2018). *Soundscape Ecology* (package « sounecology »

1.3.3) [Computer software]. http://ljvillanueva.github.io/soundecology/

Villanueva-Rivera, L. J., Pijanowski, B. C., Doucette, J., & Pekin, B. (2011). A primer of acoustic analysis

for landscape ecologists. *Landscape Ecology*, *26*(9), 1233-1246.

https://doi.org/10.1007/s10980-011-9636-9

Villéger, S., Mason, N. W. H., & Mouillot, D. (2008). New multidimensional functional diversity indices

for a multifaceted framework in functional ecology. *Ecology*, *89*(8), 2290-2301.

https://doi.org/10.1890/07-1206.1

Wansink, D., & Tinbergen, J. M. (1994). The Influence of Ambient Temperature on Diet in the Great

Tit. *Journal of Avian Biology*, *25*(4), 261. https://doi.org/10.2307/3677272

Willig, M. R., Kaufman, D. M., & Stevens, R. D. (2003). Latitudinal Gradients of Biodiversity : Pattern,

Process, Scale, and Synthesis. *Annual Review of Ecology, Evolution, and Systematics*, *34*(1),

273-309. https://doi.org/10.1146/annurev.ecolsys.34.012103.144032

Zeuss, D., Brunzel, S., & Brandl, R. (2017). Environmental drivers of voltinism and body size in insect

assemblages across Europe : Voltinism and body size in insect assemblages. *Global Ecology*

and Biogeography, *26*(2), 154-165. https://doi.org/10.1111/geb.12525

Zvereva, E. L., Castagneyrol, B., Cornelissen, T., Forsman, A., Hernández-Agüero, J. A., Klemola, T.,

Paolucci, L., Polo, V., Salinas, N., Theron, K. J., Xu, G., Zverev, V., & Kozlov, M. V. (2019).

Opposite latitudinal patterns for bird and arthropod predation revealed in experiments with

differently colored artificial prey. *Ecology and Evolution*, *9*(24), 14273-14285.

https://doi.org/10.1002/ece3.5862

Zvereva, E. L., & Kozlov, M. V. (2021). Latitudinal gradient in the intensity of biotic interactions in

terrestrial ecosystems : Sources of variation and differences from the diversity gradient

revealed by meta-analysis. *Ecology Letters*, *24*(11), 2506-2520.

https://doi.org/10.1111/ele.13851

918 Biosketch

Laura Schillé is a PhD candidate interested in the functional ecology of bird communities, which she

studies at different scales. She also has an interest in acoustic ecology.

Co-authors are ornithologists and/or have interests in community ecology and functional ecology.

Author contribution: B.C., L.B. and E.V.C. conceptualized the study and developed the methodology.

E.V.C., F.B., M.C.B, M.Bo., M.Br., T.D., M.dG., J.D., ML.D., AM.D., S.G., CB.E., M.F., P.F.C., E.F., A.G.,

MM.G., S.G., J.H., D.H., R.H., R.I., A.K., MV.K, V.L., B.L.T., C.L.V, S.M., E.M., S.M., X.M., A.M., DL.M., A.P.,

AV.P., A.P., A.S., K.S., T.S., A.T., R.T., D.T., G.V., I.V.H., Z.V., L.B. & B.C. collected the data. A.H., M.dG.,

T.B., A.P., V.L., L.P., F.A., D.F., T.P., M.M., N.S., L.B., V.M., E.M., J.G., K.T., L.G., A.M., M.C.B. & R.B.

processed audio recordings for bird species identification. L.S. processed and analyzed the data with

guidance from B.C. and L.B. L.S., B.C., L.B. led the writing and all authors contributed critically to the

revisions.