Aging reduces motivation
through decreased *Bdnf* expression in the ventral tegmental area

Hanyue Cecilia Lei[1], Kyle E. Parker[2], Chao-Cheng Kuo[2], Carla M. Yuede[3],
Jordan G. McCall[2]*†, Shin-ichiro Imai[1]*†

[1]Department of Developmental Biology
[2]Department of Anesthesiology, Center for Clinical Pharmacology, Pain Center
[3]Department of Psychiatry
Washington University School of Medicine; Saint Louis, Missouri, USA.

†Equally contributing authors

Corresponding author:
Shin-ichiro Imai, M.D., Ph.D.
Theodore and Bertha Bryan Distinguished Professor
of Environmental Medicine
Department of Developmental Biology
Department of Medicine (Joint)
Washington University School of Medicine
Campus Box 8103
660 South Euclid Avenue
St. Louis, MO 63110
Tel (Office): 314-362-7228
Fax (Departmental): 314-362-7058
E-mail: imaishin@wustl.edu

Jordan G. McCall, Ph.D.
Associate Professor
Department of Anesthesiology
Center for Clinical Pharmacology
Pain Center
Washington University School of Medicine
Campus Box 8054
660 South Euclid Avenue
St. Louis, MO 63110
Tel (Office): 314-446-8157
E-mail: jordangmccall@wustl.edu

Abstract

Age-associated reduced motivation is a hallmark of neuropsychiatric disorders in the elderly. In

our rapidly aging societies, it is critical to keep motivation levels high enough to promote

healthspan and lifespan. However, how motivation is reduced during aging remains unknown.

Here, we used multiple mouse models to evaluate motivation and related affective states in

young and old mice. We also compared the effect of social isolation, a common stressor in aged

populations, to those of aging. We found that both social isolation and aging decreased

motivation in mice, but that *Bdnf* expression in the ventral tegmental area (VTA) was selectively

decreased during aging. Furthermore, VTA-specific *Bdnf* knockdown in young mice

recapitulated reduced motivation observed in old mice. These results demonstrate that

maintaining *Bdnf* expression in the VTA could promote motivation to engage in effortful

activities and potentially prevent age-associated neuropsychiatric disorders.

Introduction

In 2017, the World Health Organization reported that more than 20% of people aged 60 and over have a psychiatric or neurological disorder, including dementia, depression, and anxiety[1]. Older adults are also more susceptible to stressors such as social isolation, a common risk factor for many mental disorders and health issues particularly during the COVID-19 pandemic[2-4]. These age-associated psychiatric conditions are quite often associated with reduced motivation to engage in effortful activities[5]. Furthermore, maintaining high motivation levels is indicative of healthy mental conditions in old age[6]. How aging impacts mental functions, particularly motivation, and the underlying biological mechanisms, however, remains largely unknown.

To determine the neurobiological mechanisms of aging-induced decreased motivation, we adapted multiple mouse models thought to impact motivation. In particular, we used social isolation as a stressor to mimic a real-life condition in aging human population[7, 8]. We used sexually matured young adult (4 month-old) and old but still healthy (18 month-old) male mice. After one month of single or grouped housing, these mice were tested in a battery of motivational assays.

Results

Chronic social isolation and aging differentially impair motivational behaviors

Voluntary wheel-running activity (VWR) is innately rewarding to rodents even with no other extrinsic reward[9, 10]. Therefore, to test whether age impacts this endogenously motivated behavior, we compared VWR activities between young and old mice under grouped or socially isolated housing conditions. Old mice ran much less than young mice during the dark time, as reported previously[11, 12]. However, social isolation substantially reduced dark-time wheel-

running in both young and old mice (Fig. 1a and b). Most VWR reduction occurred in the early

dark time when mice reached their daily peaks of running (Extended Data Fig. 1a and b), and the

effect of social isolation was greater in old mice, compared to young mice (40.3% vs. 25.7%

reduction). Although VWR is a complex behavior that can be affected by both physiological

(e.g., reduced skeletal muscle strength and energy metabolism) and motivational changes, these

results suggest that both aging and social isolation undermine motivation.

To directly examine whether aging and social isolation reduce motivation levels in mice,

we next conducted an operant progressive ratio (PR) test to measure the amount of effort that

animals are willing to expend to gain a palatable food reward. We habituated mice to sugar

pellets in home cages two days before training and trained under fixed ratio-1 and fixed ratio-3

for five days each (Extended Data Fig. 1c). Following this task learning, we next tested the mice

under a progressive ratio (PR) for one day (Fig. 1c). In the PR test, socially isolated young mice

performed significantly fewer correct nose-pokes than group-housed controls (Fig. 1d).

Interestingly, group-housed old mice showed significantly lower correct nose-pokes than group-

housed young mice, but very similar to those of the socially isolated young mice (Fig. 1d).

Socially isolated old mice tend to show further reduction in correct nose-pokes (Fig. 1d).

Breakpoint analysis, which determines the maximum number of nose-pokes that an animal

achieves to gain a single reward[13], further demonstrates consistent reduction in effort to achieve

rewards in old and socially isolated mice (Fig. 1e). In both socially isolated groups, there were

tendencies of fewer entries into the food tray (Extended Data Fig. 1d). These results clearly

suggest that reduced motivation is a shared outcome of social isolation and aging, and old mice

seemingly responded less to the stress of social isolation perhaps due to the already reduced

effort induced by aging.

Both social isolation and aging induce depressive- and anxiety-like behaviors

Apathy and anhedonia are often comorbid with decreased motivation[14]. To determine whether

normal reward processing is impacted by social isolation and aging, we next conducted the

sucrose preference test (SPT). All groups of mice manifested robust preference to sucrose

solution over water in the sucrose preference test (Extended Data Fig. 1e), indicating no obvious

anhedonia. Nonetheless, the amount of sucrose solution consumed was selectively reduced by

social isolation and aging (Fig. 1f), although the extent of reduction by social isolation did not

reach statistical significance in either young or old mice. Reduction in energy demand was an

unlikely explanation because daily food intake was not different for any group on the test day

(Extended Data Fig. 1f). Notably, the results from the SPT showed very similar patterns to those

from the PR test, supporting the notion that social isolation and aging both reduce motivated

behavior in mice.

Finally, to determine whether approach-avoidance conflicts that underly some features of

an anxiety-like state might be altered by social isolation and aging in young and old mice, we

tested mice in the locomotor activity/open field test (LA/OFT) and elevated plus maze (EPM).

Socially isolated young mice spent less time exploring the center area than the group-housed

controls, whereas both group-housed and socially isolated old mice exhibited similarly lowered

center time and distance traveled (Fig. 1g and Extended Data Fig. 2a and b). Contrarily, socially

isolated young mice showed significant increases in time and distance traveled to explore the

peripheral area (Extended Data Fig. 2a and b). Additionally, in the EPM, social isolation

increased distance traveled and percentages of moving times in the close arm in both young and

old mice, as we observed previously[15], whereas there were no differences between young and old

mice (Extended Data Fig. 2c-e). We also conducted the hole-board olfactory preference test and

the social interaction test to assess other types of motivation-related social behavior between

young and old mice in both group-housed and socially isolated conditions (Extended Data Fig. 3).

Social isolation had no effect on either assay, except for a trend for isolated young mice to seek

more social cues. Therefore, when motivation was reduced by social isolation and aging, certain

anxiety- or depression-related behavioral states were also present. Together, these results suggest

that both aging and social isolation reduce motivation in mice, but perhaps not additively.

Aging impairs dark time-dependent enhancement of reward-seeking motivation

To determine the extent to which age alters motivation, we tested a new cohort of young and old

mice across different testing conditions. Remarkably, we found that when food-deprived and

tested during the dark time (Fig. 2a), differences in correct nose-pokes detected in the PR test

between young and old mice were greatly enhanced (Fig. 2b). Similar results were obtained for

break points (Fig. 2c). These results suggest that the reward-seeking motivation of young mice is

enhanced by fasting and heightened during the dark time, whereas old mice fail to show such

dark time-dependent enhancement of reward-seeking motivation, confirming that aging is a

critical contributor to reduced motivation.

BDNF mRNA and protein levels are decreased specifically in the ventral tegmental area

(VTA) during aging

Extensive literature has defined the role of the mesolimbic dopamine (DA) system in regulating

reward and motivation[16, 17]. Two major DA populations are identified in the ventral tegmental

area (VTA) and the substantia nigra (SN) that locate adjacently in the ventral midbrain[18]. To

determine the mechanism for this age-associated decline in motivation, we examined mRNA

expression levels of characteristic functional genes in these two regions by collecting tissue

samples from young and old brain sections with laser-capture microdissection and conducting

quantitative RT-PCR. We found that expression of brain-derived neurotrophic factor (*Bdnf*) was

significantly decreased in the aged VTA (Fig. 3a). Interestingly, genes related to DA turnover

did not show any significant changes. The expression of *Gria1*, which encodes the GluA1

AMPA glutamate receptor subunit, tended to decrease in aged VTA (Fig. 3a). On the other hand,

no genes examined in the SN were significantly altered during aging, except for a trend of

increase in the expression of *Maoa*, which can be indicative for augmented DA metabolism and

oxidative stress in this region (Fig. 3b)[19, 20]. We also confirmed this age-associated decrease in

Bdnf expression by fluorescent *in situ* hybridization. Lower *Bdnf* signals were detected across

eight consecutive bregma levels throughout the aged VTA, compared to the young VTA (Fig.

3c-d), whereas the expression of *Slc6a3*, a DA neuronal marker, did not show any significant

differences between young and old mice (Extended Data Fig. 4a-b). Age-associated decreased

Bdnf expression was present throughout the VTA in both *Slc6a3*-positive and *Slc6a3*-negative

cells (Extended Data Fig. 4c-f), suggesting widespread downregulation of *Bdnf* throughout the

VTA. Whereas aging selectively decreased *Bdnf* expression in the VTA, social isolation did not

decrease *Bdnf* expression in the VTA (Extended Data Fig. 4g, left panel). On the other hand,

social isolation tended to decrease *Bdnf* expression in the dorsal raphe nucleus (DRN) at least in

young mice (Extended Data Fig. 4g, right panel). Given that the activity of DRN DA neurons is

important to elicit a loneliness-like state and promote social preference[21,22], a commonality

between aging and social isolation appears to be decreased *Bdnf* expression, but in different areas

of the brain. Following from the decrease in *Bdnf* mRNA, we further observed that matured

BDNF (mBDNF) protein levels were moderately reduced in the aged VTA (Fig. 3 f-g),

suggesting a potential involvement of *Bdnf* in age-associated reduced motivation.

VTA-specific *Bdnf* knockdown in young mice recapitulates the reduced motivation

observed in aged mice

Following the observation that VTA *Bdnf* was decreased in aged mice, we attempted to restore

VTA *Bdnf* expression in old mice to determine if their motivation could be rescued. To do so, we

used a *Bdnf*-expressing lentivirus. This approach yielded significant BDNF protein

overexpression in primary mesencephalon neuronal culture, but it failed to increase BDNF

protein levels *in vivo* so that PR performances were not improved in old mice (Extended Data

Fig. 5). We suspect that this could be due to certain auto-regulatory mechanisms that suppress

Bdnf transcription and/or translation in the VTA neurons in response to overexpression[23-26]. It is

also possible that the lentiviral transgene was silenced in the host cells over time[27-29]. We next

sought to determine whether decreasing VTA *Bdnf* expression in young mice can recapitulate the

motivational dysfunction observed in aged mice. To do so, we knocked down *Bdnf* expression in

the VTA by stereotactically injecting a *Bdnf* shRNA-expressing lentivirus into the VTA of young

mice (Fig. 4a). *Bdnf* mRNA levels were reduced by ~30% (Fig. 4b-c), and the mBDNF protein

levels were also reduced to a similar extent to aged mice in the VTA, but not in the downstream

nucleus accumbens (NAc) (Fig. 4d-f). Two months after surgery, the VTA-specific *Bdnf*-

knockdown (VTA-*Bdnf*-KD) mice were tested for the extended PR paradigm with food

deprivation during light and dark times. Importantly, VTA-*Bdnf*-KD mice exhibited significant

reduction in correct nose-pokes and break points with food deprivation during both light and

dark times, successfully recapitulating motivational phenotypes observed in aged mice (Fig. 4g-h,

and Extended Data Fig. 6a-b). On the other hand, VTA-*Bdnf*-KD mice did not alter behaviors in

the VWR and OFT (Fig. 4i-l, and Extended Data Fig. 6c-d), supporting a selective role of *Bdnf*

in the VTA in reward-seeking motivational behavior. Prior work has shown that VTA BDNF can

play a role in regulating VTA dopamine neuron excitability[30, 31]. To determine whether the

downregulation of VTA *Bdnf* in aged mice altered excitability in these VTA neurons, we

examined the intrinsic neuronal electrical properties of both putative dopaminergic and non-

dopaminergic neurons in young and aged mice. For these purposes, we operationally defined

these neurons by the expression of hyperpolarization-activated cation current (I_h)[32] (Extended

Data Fig. 7a-b). No significant differences were found between young and aged mice, suggesting

that baseline neuronal excitability in the VTA does not drive the decreased motivation through

aging (Extended Data Fig. 7c-f). Taken together, these results demonstrate that specific

decreases in *Bdnf* expression in the aged VTA contribute, at least in part, to age-associated

reduced motivation in mice.

Discussion

Here we demonstrated that both social isolation and aging significantly decreased motivation in

mice along with the induction of certain anxiety- or depression-related behaviors. Aging is a

strong contributor to reduced motivation, and this may explain why aged mice respond

substantially less to social isolation compared to young mice. In this regard, we found that *Bdnf*

expression was significantly reduced throughout the aged VTA, in both DA and non-DA cells.

Remarkably, the VTA-specific knockdown of *Bdnf* in young mice recapitulated the reduced

motivation observed in old mice, suggesting that this VTA-specific reduction in *Bdnf* contributed

to the impaired reward-seeking motivation in aged mice. Whereas VTA BDNF levels have been

implicated in the development and progression of drug-induced reward processing or addiction[33,]

[34], our findings suggest that BDNF also plays an important role in maintaining motivated states

in a healthy, physiological conditions, which impair with age. However, given that VTA-*Bdnf*-

KD mice did not alter voluntary wheel-running and open field activities, the effect of age-

associated BDNF reduction is limited to palatable reward-seeking motivation. Furthermore,

because neuronal excitability in the VTA shows no significant changes during aging, it is

possible that endogenously evoked activity and synaptic transmission in this particular reward-

seeking circuit may be affected by age-associated BDNF reduction.

Importantly, while the high frequency of depression in Parkinson's disease appears

dependent on decreased DA innervation of the limbic system[35, 36], the age-associated decreases

we observed in VTA *Bdnf* expression occurred in both DA and non-DA cells, suggesting that

this is a distinct mechanism for age-associated decreased motivation. Interestingly, it has been

reported that leptin receptor-expressing neurons in the lateral hypothalamus (LH) mainly project

to non-DA VTA neurons and drive PR performance through the activation of these VTA neurons

[37]. We previously showed that specific neuronal populations in the dorsomedial hypothalamus

and LH play critical roles in counteracting age-associated functional decline and determining

longevity in mice [38-40]. Therefore, it will be of great interest to determine whether these neuronal

populations that fundamentally control aging and longevity also regulate motivation through

Bdnf-expressing VTA neurons. Because the LH shows significant reduction in its NAD^+ levels

during aging and may rely on extracellular nicotinamide phosphoribosyltransferase (eNAMPT),

an important NAD^+ biosynthetic enzyme counteracting aging, for its NAD^+ biosynthesis[41-43], it

will be important to examine whether eNAMPT supplementation could restore motivation in

aged mice.

In humans, it has been reported that BDNF in the brain decreases with age[44-47]. In

particular, Buchman *et al.* reported that in more than 500 older adults, high *BDNF* expression

levels in the brain were associated with slower rates of cognitive decline[45]. Additionally, Oh *et al.*

showed that age-associated *BDNF* decline was assocaited with the downregulation of excitatory

and inhibitory synaptic markers, implicating that *BDNF* reduction may affect synaptic

transmission in the aged brain[44]. Although previous studies did not assess the connection

between *BDNF* and motivation, it is likely that age-associated *BDNF* decline in the brain could

cause reduced motivation in humans. Interestingly, young adult human subjects (age 18-29)

perceive the highest levels of isolation during the pandemic of COVID-19, compared to old

subjects (age 50-69), and overall life satisfaction is significantly lower for people who perceived

greater social isolation[48]. This trend is qualitatively similar to our results in the PR test, showing

that aged mice are less responsive to social isolation, compared to young mice, most likely due to

already reduced levels of motivation. Therefore, our results in this study could provide an

important insight into age-associated alterations in mental states in humans.

In conclusion, the present study advances our understanding of the mechanism of age-

associated motivational changes and the physiological importance of Bdnf in the VTA. It also

stimulates development of interventions to keep our motivation levels high and healthy during

aging.

Methods

Animals

All animal procedures were approved by the Washington University Institutional Animal Care

and Use Committee and were in accordance with National Institutes of Health guidelines. Unless

otherwise noted, all mice were housed in groups of 2-5, fed *ad libitum* on a standard chow diet

(PicoLab Rodent Diet 20-5053; LabDiet, St. Louis, MO) and autoclaved water in a

temperature/humidity-controlled holding room within a barrier facility, maintained on a 12:12-

hour light/dark cycle. Food, beddings (Aspen Chip, Northeastern Products Corp.), and nesting

material (Nestlets, Ancare) were changed once per week; water bottles were changed as needed.

Mice were labeled on their tails with sharpie markers for easy identification and monitored

constantly for their health status. All handling, cage changing, measuring procedures, and

behavior tests were conducted by the same experimenters throughout the study. Only male mice

were used.

All aged mice and the corresponding young groups were obtained from the NIH aged rodent

colony (C57BL/6JN) or our in-house C57BL/6J colony. Due to the limited availability of aged

mice, this study only used male mice to limit the number of potential biological variables to

remain sufficiently powered. All group-housed cages were housed only with original cage mates

from weaning. Mice used for *Bdnf* knock-down analysis were purchased from Jackson

Laboratories (C57BL/6J) at 12-week-old and acclimated to the animal facility for ~2 months

before surgeries; upon arrival, mice were randomly assigned into cages of 4/5 and maintained

until sacrifice.

Social isolation procedure

All mice were originally group-housed in cages of 3-5 cage mates. After acclimation to the

holding rooms, cages containing around half of the animals were separated into individual cages

and maintained isolated until sacrifice; the rest of the control cages remained unchanged across

the study. All cages within a certain cohort were placed in proximity on the same rack to control

for traffic, noise levels, and light exposure.

Behavior tests

All behavioral tests were performed within a sound-attenuated room maintained at 20-26°C after

at least one week of acclimation to the local holding room. All tests were performed by female

experimenters. Unless otherwise noted, all mice in the same cohort underwent the same

test/procedure on the same day(s) from ZT3 to ZT9; if a test/procedure cannot be finished within

one day, the cohort will be divided into sub cohorts by counterbalancing grouping, home cages,

and other factors. For all experiments performed outside of the holding room, mice were brought

into the corresponding behavior space to habituate for at least 20 minutes before any

test/procedure started. All behavioral apparatuses were cleaned with 2% chlorhexidine in

between subjects.

Operant Progressive-Ratio Test

To expose the animals to the reward, mice fed *ad libitum* were given 3 g/mouse of the reward

sugar pellet (Dustless Precision Pellet, F0071, Bio-Serv) on the bedding of their home cages

when cages were changed at ZT12. Consumption of the pellets was confirmed the next day. One

day after the exposure, mice were gently placed in a mouse operant-conditioning box

(15.24×13.34×12.7 cm) within a sound attenuating cubicle (ENV-307A & ENV-020M, Med

306 Associates Inc., Fairfax, VT). The box was equipped with a stainless steel grid floor, two nose

307 holes (active and inactive) flanking a trough pellet receptacle (food tray) connected to a pellet

308 dispenser on the bottom of the right-hand side wall; a house light was installed at the top center

309 of the left-hand side wall. Pokes into the ports are monitored by infrared light beam sensors.

310 Upon starting of a fixed-ratio 1 (FR1) session, both nose holes present a cue light; one nosepoke

311 into the active hole results in a sugar pellet (20 mg) dispensed to the food tray, along with cue

312 lights turned off and house light turned on for 2 s, during which no action triggers event (time-

313 out period); poking the inactive hole has no consequence. Each session lasted 60 minutes. In

314 between trials, boxes were cleaned with 2% chlorhexidine, and leftover pellets in the dispenser

315 were replaced with fresh and intact ones. All trials took place from ZT3 to ZT8, unless otherwise

316 specified, and each animal was subjected to the sessions at the same time across days. In FR1,

317 mice were trained to discriminate the active hole from the inactive one and receive rewards from

318 the food tray. Following 5 consecutive days of FR1 sessions, mice were moved on to another 5

319 consecutive days of 60-minute fixed-ratio 3 (FR3) sessions, in which three active nosepokes

320 results in one reward delivery and 20 s time-out period. Mice were conditioned to repeatedly

321 perform the correct operant behavior until receipt of the reward during FR3. After completion of

322 the training, motivation levels of the mice were assessed using a progressive-ratio (PR) schedule

323 of reinforcement following the equation: response ratio $= 5e^{reward\ number/5} - 5$ (rounded to the

324 nearest integer), resulting in an exponential increase of the number of correct pokes required for

325 each subsequent reward: 1, 2, 4, 7, 9, 12, 15, 20, 25, 32, 40, 50, 62, 77, 95, 118, 145, 178, 219,

326 268, 328, 402…[49]. All events were programed, and actions were recorded using MED-PC

327 software (Med Associates) controlled by a PC. The number of active nosepokes counted toward

328 rewards during the 60-minute session was calculated as the readout of motivation; A break point

(BP) was calculated by taking the bigger number of the last completed ratio versus the number of

correct nosepokes after the last reward[13]. For the extended PR paradigm, following FR1 and FR3

training, two PR sessions were conducted on consecutive days, and the averaged readouts served

as the PR baseline. Mice then underwent a break day with no session, followed by a third PR test

(light fed) under the same conditions (*ad libitum* in the light time); after another break day and

overnight fasting from ZT12, a fourth PR test was conducted (light fasted); lastly, two days after,

the mice were fasted from ZT0, and a fifth PR test under fasting was conducted during ZT15-20

under dim red lighting (~10 Lux).

Sucrose Preference Test

Mice were habituated to the sipper tubes in their original home cages by replacing the drinking

bottle with a sipper tube for 4 days; daily water consumption was confirmed to be > 2

mL/mouse/day by the last day of the habituation. The sipper tube was home-constructed with a

cut 25-mL plastic serological pipettes and a stainless steel pin valve (Lixit) sealed together with

heat shrink tubing. Mice were then all separated into individual cages (if not already separated as

part of the social isolation procedure), given *ad libitum* regular chow and two identical sipper

tubes positioned side-by-side next to the chow, both containing drinking water. Baseline water

consumption was measured by recording volume left in each sipper tube daily for 4 days. After

that, water in one of the tubes was replaced with 2% sucrose solution. Sucrose solution and water

consumed per day were recorded for another 4 days, and the percentage of sucrose solution over

total liquid intake was calculated to represent sucrose preference. Tube positions (left vs right)

were switched daily after volume checking across the entire test. Mice were housed back with

their original cage mates after the procedure was finished.

Elevated-Plus Maze Test

The EPM was performed under dim warm lighting stabilized at ~5-6 lux. The apparatus was a

plus-shaped platform comprised of two open arms (25×5×0.5 cm) across from each other and

perpendicular to two closed arms (25×5×16 cm) with a center platform (5×5×0.5 cm), elevated

to be 50 cm above the floor (60140, Stoelting Co). After habituating to the room, each mouse

was individually placed onto the apparatus from one of the open arms and allowed to freely

explore. Activity was video recorded using a CCD camera for 20 minutes and videos analyzed

using Ethovision XT 13 (Noldus Information Technology, Wageningen, The Netherlands).

Behavior tracking after the mice made the first entry into the center area was included in the

analysis.

One-hour Locomotor Activity and Open Field Test

The 1-hour LA/OFT was conducted in the Animal Behavior Core at Washington University

School of Medicine in St. Louis. The test was performed under standard room lighting stabilized

at ~150 lux The apparatus consisted of a transparent (40.6×40.6×38.1 cm) polystyrene enclosure,

surrounded by a frame containing a 16×16 matrix of photocell pairs. Output was fed to an online

computer and analyzed by the system software (Hamilton-Kinder, LLC, Poway, CA). A center

area was defined as a 9x9 grid within the 15x15 grid in this chamber, with the peripheral area

extending to the sides of the chamber as the outermost boundary. After habituating to the room,

each mouse was individually placed in the center of the apparatus and allowed to freely explore

for one hour. The OFT for VTA-*Bdnf*-KD mice was performed under dim warm lighting

stabilized at ~5-6 lux. The apparatus consisted of an open-top opaque square enclosure

(50×50×31 cm). After habituating to the room, each mouse was individually placed at the center

of the apparatus and allowed to freely explore. Activity was video recorded using a Google Pixel

3 XL smartphone for 30 minutes and videos analyzed using Ethovision XT 13. The center was

defined as a square comprised of 36% the total area of the OFT (i.e., each side was 0.6 that of the

total arena).

Hole-board olfactory preference test

The HOP was conducted in the Animal Behavior Core at Washington University School of

Medicine in St. Louis. The test was performed under standard room lighting stabilized at ~150

lux. The apparatus consisted of a transparent (40.6×40.6×38.1 cm) polystyrene enclosure

equipped with a computerized hole-board apparatus of the same size, which contains 4 corner

and 4 side holes, with a side hole being equidistant between the corner holes (Learning

Holeboard; MotorMonitor, Kinder Scientific, LLC, Poway, CA). This task consisted of two

consecutive days. On day 1, mice were put into the apparatus for 20 min to habituate. On day 2,

mice were presented with one of 8 options for configuring of corner holes containing either

female urine, vanilla extract, distilled water, or empty. Pairs of photocells are equipped within

each hole (27 mm diameter) and were used to quantify the frequency and duration of pokes.

Specifically, a mouse was required to place its head into a hole at a depth of at least 35 mm and

break a photobeam for the response to be registered as a hole poke. All odorant-containing cups

were cleaned with mild soap and water, and the entire apparatus was cleaned with a 2%

chlorohexidine diacetate solution to remove the odor between trials.

Social Interaction Test

The test was performed in recording closets under lighting stabilized at ~ 10 lux. The apparatus

consisted of a homemade black opaque two-chamber box (52×26×26 cm) divided in half by

blockers on the sides, leaving the center open (W10×H26 cm). A metal, mesh pencil cup

(10H×8.5 radius cm) was invertedly placed in the center of each chamber, with extra weight and

a cone cup attached on top to prevent animals from moving or climbing on them. The mesh

allows for exchange of sensory information and prevents direct physical interactions between

animals inside and outside of the cup. The floor of the box was covered by a thin layer of

bedding material. After habituating to the room, to assess baseline activities, each subject mouse

was individually placed in the center of the box and allowed to freely explore for 10 min,

followed by 30 min back to their home cages. To test for sociability, a sex- and age-matched

stranger conspecific was placed under one of the cups in a counterbalanced and randomized

manner, and the subject mouse was then reintroduced to the same box and allowed to freely

explore for another 10 min. Bedding in the boxes was replaced with fresh one in-between

subjects. Activity was video-recorded using a CCD camera, and video recordings were analyzed

using Ethovision XT 13. Activities of the mice in either chamber during the baseline and the test

phases were analyzed.

Voluntary wheel-running activity

After concluding prior behavior test(s), mice were transferred to the VWR facilities and housed

individually in a Tecniplast polycarbonate cage (33.2×15×13 cm) (PT2-MCR, Phenome

Technologies) with a vertical running wheel (11 cm inside diameter, 5.4 cm wide), stainless steel

wire top, and a filter cage cover (Actimetrics). Wheel-running behavior was recorded and

analyzed using the ClockLab software (https://actimetrics.com/products/clocklab/). Mice were

allowed to habituate to the wheel cages for at least a week and acquire stable running activity

before data was collected.

Laser-captured microdissection

Mice were sacrificed by CO_2. Brains were removed, segmented into separate parts in a brain

matrix, and immediately embedded in OCT and snap frozen by powered dry ice. Brain samples

were stored at $-80°C$ until sliced into 30 μm coronal sections in a cryostat at $-14°C$. Sections

were thaw-mounted onto PEN-membrane slides (#11505158, Leica Microsystems, Buffalo

Grove, IL, USA) and kept frozen until the next step. After collecting all sections containing brain

regions of interest, slides were hydrated sequentially in 100%, 95%, 75%, and 50% ethanol for

30 sec each. The hydrated slides were stained with 0.5% Cresyl Violet (Sigma-Aldrich) for 2

minutes, and dehydrated with 50%, 75%, 95%, and 2 rounds of 100% ethanol for 30s each. The

dehydrated slides were then incubated in xylene twice for 1 minute each. After being air-dried,

VTA, SN, and DRN regions were microdissected using the Leica LMD 6000 laser-

microdissection system, following the Franklin and Paxinos mouse brain atlas. Laser-cut tissue

samples were collected in the extraction buffer of the Arcturus PicoPure RNA Isolation Kit

(12204-01, Applied Biosystems), and stored at $-80°C$ until RNA was isolated following the

manufacturer's protocol and DNase (79254, Qiagen) treatment.

Quantitative real-time RT-PCR

Concentrations of RNA samples were measured by NanoDrop and stored at $-80°C$ until being

reverse-transcribed into cDNA with the High Capacity cDNA Reverse Transcription kit

(#4368814, Thermo Fisher). qRT-PCR was conducted with the TaqMan[TM] Fast Universal PCR

Master mix (4352042, Applied Biosystems) and TaqMan primers for each gene (*Gapdh*:

Mm99999915_g1, *Th*: Mm00447557_m1, *Slc6a3*: Mm00438388_m1, *Slc18a2*:

Mm00553058_m1, *Ddc*: Mm00516688_m1, *Maoa*: Mm00558004_m1, *Bdnf*: Mm04230607_s1

and 4331348_APNKZZR, a customized primer for the overexpressed *Bdnf* mRNA:

ATAGCAAAAAGAGAATTGGCTGGCGATTCATAAGGATAGACACTTCCTGTGTATGT

ACACTGACCATTAAAAGGGGAAGA, *Gria1*: Mm00433753_m1) in the StepOnePlus^{TM}

Real-Time PCR System (Applied Biosystems). Relative expression levels were calculated for

each gene by normalizing to *Gapdh* CT values and then to the mean of the control/reference

group.

RNAscope Fluorescent *In Situ* Hybridization (FISH)

Mice were transcardinally perfused first with 0.01M phosphate-buffered saline (PBS) and then

the fixative: 2% paraformaldehyde (PFA) and 0.2% picric acid in 0.01M PBS, pH 7.4. Brains

were removed and post-fixed in the fixative at 4°C overnight, followed by cryoprotection in 20%

sucrose in 0.01M PBS at 4°C for at least 24□hours and snap freezing by powered dry ice. Brains

were stored at -80°C until sliced into 20 μm coronal sections containing the VTA region in a

cryostat at -20°C. Sections were thaw-mounted onto Superfrost^{TM} Plus Microscope Slides (12-

550-15, Fisher Scientific) and kept frozen until use. FISH was performed using RNAscope

Multiplex Fluorescent Reagent Kit v2 [Advanced Cell Diagnostics, Inc. (ACD)] as per the

manufacturer's protocol. Briefly, slides washed with 1x DPBS were fixed in pre-chilled 4% PFA

in 1x DPBS at 4°C for 15 minutes. Sections were then dehydrated through 50%, 70%, and two

rounds of 100% ethanol for 5 minutes each, followed by air-drying. Then, the sections were

treated with RNAscope® Hydrogen Peroxide for 10 minutes at room temperature. A

hydrophobic barrier was created around the sections using ImmEdge™ Hydrophobic Barrier

PAP Pen (H-4000, Vector Laboratories) and allowed to air-dry. The slides were then incubated

with Protease III for 30 minutes and hybridized with ACD probes for target genes (Mm-*Bdnf*-

CDS, #457761, probe region 662-1403; Mm-*Slc6a3*-C3, 315441-C3, probe region 1486-2525)

for 2 hours at 40°C using a HybEZ™ oven. The signals were amplified by subsequent

incubations with Amp-1, Amp-2 and Amp-3 reagents for 30, 30, and 15 minutes, and then with

the HRP reagents, depending on the channels used, in combination with the TSA Plus

(NEL760001KT, PerkinElmer) and Opal (FP1487001KT, Akoya) fluorophores at 1:1000 to

1:500 dilutions. Sections were washed with 1x RNAscope Washing Buffer in-between every

incubation. Slides were counterstained with DAPI, and coverslips were mounted with

FluorSave™ Reagent (#345789, Calbiochem). Images were obtained using fluorescent

microscopy (DMi8, Leica Microsystems) under 20x magnification. Fluorescent signals were

quantified using a customized ImageJ (NIH) pipeline. Briefly, after tiff images were exported, a

circular ROI (radius = 200 μm) was selected at the center of the VTA region, referring to the

shape covered by the *Slc6a3* signal; after subtracting background from all channels, the *Slc6a3*

channel underwent Gaussian Blur, binary conversion, and shape growth to approximate the area

covered by the dopaminergic neurons; the number of the *Bdnf* puncta was quantified using the

Find Maxima function, and the identified points were overlaid on the *Slc6a3* area to count the

Bdnf signals that are expressed in the dopaminergic neurons. For the *Bdnf* knock-down

experiment, signals were quantified where the virus caused infection in the VTA.

Brain punching

Mice were transcardinally perfused with 0.01M PBS, and brains were removed and snap frozen

by powered dry ice. Brain samples were stored at -80°C until sliced into 200um coronal sections

from anterior to posterior of the brain in a cryostat at -8°C. Section containing the nucleus

accumbens (NAc) and VTA regions were identified by bregma landmarks and punched to

acquire these regions using a Brain Punch Set (#57401, Stoelting). The NAc region were

collected by the 1.19 mm punch from one side of 4 consecutive brain sections (Bregma around

1.5 to 0.7); the VTA region were collected by the 0.74 mm punch from both sides of 5

consecutive brain sections (Bregma around -2.8 to -3.8). To proceed to western blotting, each

punched sample was homogenized in 60 μL 1x sample buffer containing 10% 2-

Mercaptoethanol by passing through a syringe with 28-gauge needle (102-SN1C2805P,

McKesson) for 5 times. The homogenate was then centrifuged at 16,000 xg for 2 minutes, and

the clear upper layer of 55 μL was carefully collected into a new tube.

Western blotting

Protein samples prepared in 1x sample buffer containing 10% 2-Mercaptoethanol were incubated

at 95°C for 10 minutes and allowed to cool down. 14 μL of each sample was electrophoresed on

a 4-15% Mini-PROTEAN® TGX™ Precast Protein Gels (#4561086, BioRAD) in a Tris-Glycine

running buffer, at 80-100V. The separated protein samples were then transferred to a

Immobilon®-P PVDF Membrane (IPVH00010, Millipore) in iced transfer buffer that contains

0.1% SDS and 10% methanol, at 0.2A for 50 minutes in a cold room. After being washed in 1x

Tris-buffered saline with 0.1% Tween-20 (TBS-T) and blocked in 5% w/v skim milk/TBS-T for

1 hour at room temperature (RT), the membrane was then incubated with rabbit anti-BDNF

(1:1000, ab108319, Abcam) or rabbit anti-β-actin (1:1000; PA1-183, Thermo Fisher) in 5% skim

milk overnight at 4°C. After the membranes were washed in TBS-T for 3 times, they were

incubated with HRP conjugated anti-rabbit IgG antibody (1:10,000, NA9340, Cytiva) in 5%

skim milk for 1 hour at RT. After washing the membrane in TBS-T for 3 times,

chemiluminescence signals were developed using Amersham ECL reagents (RPN2235 or

RPN2232, Cytiva) and visualized using the [647SP, no light] setting of the ChemiDoc MP

Imaging System (BioRAD) with exposure times of 1-20 minutes. When need to be reblotted,

membranes were stripped with a stripping buffer (#21059, Thermo Fisher) for 15-30 minutes at

RT. All incubations and washes were done with constant rocking. Signal intensity of the

appropriate bands was quantified using ImageJ (NIH).

Virus vector construction

To generate the shRNA expressing lentiviral constructs, 56-bp double-stranded oligonucleotides,

each of which contains a sense target sequence, a microRNA based loop sequence

(CTTCCTGTCA), an antisense sequence, and a 4-nucleotide sequence complimentary to the

vector cutting site of the restriction enzyme AarI at both ends, were generated for *Bdnf* and

firefly luciferase (*fLuc*) and cloned into the U6-PGK-eGFP vector, provided by the Hope Center

Viral Vectors Core at Washington University School of Medicine in St. Louis. Three shRNA

targets on the *Bdnf* coding sequence were cloned and tested in parallel, and the one that gave the

highest knock-down efficiency *in vitro* was selected for *in vivo* experiments. The sense sequence

used   for   *Bdnf*   was   5'-ACTATCCATTCTGGTTGATAA-3'   and   5'-

TACGCGGAATACTTCGAAATG-3' for *fLuc*.

Gibson Assembly designed using the NEBuilder™ website (https://nebuilder.neb.com)

was used to generate the *Bdnf* overexpressing lentiviral constructs. The *Bdnf* cDNA without stop

codon was cloned from total cDNA extracted from a wildtype mouse hippocampus sample using

the primer set: 5'-CCGTTACTAGTGGATCCACCGGTGCCACCATGACCATCCTTTTC-3'

and 5'-CGGAGCCTCTTCCCCTTTTAATGGTCAG-3'. A P2A-eGFP sequence was cloned

from the AAV-CamKIIa-hWFS1-P2A-eGFP vector, provided by the Hope Center Viral Vectors

Core, using the primer set: 5'-GGGGAAGAGGCTCCGGAGCCACGAAC-3' and 5'-

AATTCCGCGGGCCCGTCGACTTACTTGTACAGCTCGTCCATGC-3'. The two segments

were then assembled into the LV-SYN-eGFP vector, provided by the Hope Center Viral Vectors

Core, using the Gibson Assembly® Master Mix (E2611, NEB). The original LV-SYN-eGFP

vector was used as the control. After confirming the new constructs by sanger sequencing

provided by GENEWIZ, high concentrations of vector DNA samples were produced using

ZymoPURE II Plasmid Maxiprep Kit (D4203, ZYMO), following the manufacturer's protocol.

High-titer lentiviruses were produced by the Hope Center Viral Vectors Core. Briefly, HEK293T

cells were co-transfected with the shRNA-expressing vectors and three packaging vectors (pMD-

Lg/pRRE, pCMV-G, and RSV-REV) by using polyethylenimine (PEI) reagent. Six hours after

transfection, the medium was replaced with the complete medium containing 6 mM sodium

butyrate. Culture supernatant was collected 42 hours after transfection. The supernatant was

passed through a 0.2 μm filter, concentrated by ultracentrifugation through a 20% sucrose

cushion, and stored at -80°C until use. Virus titer was determined by transducing HT1080 cells

and assaying for vector sequence using qPCR. Virus with a titer of 1.1-1.7 x 10^9 IU/mL was used

for stereotaxic injection.

Primary mesencephalon culture

Primary mesencephalon culture was prepared as previously described with modifications[50, 51].

Briefly, embryonic-day 12.5 (E12.5) C57BL/6JN mouse embryos were removed from timed

pregnant females and placed in washing medium containing DMEM-High glucose (12800-017,

Gibco) with antibiotics (15240-062, Gibco) on ice. The embryos were dissected to isolate the

midbrain region, which was further trimmed to obtain the ventral mesencephalon tissue. The

collected tissue was stored in and rinsed with iced washing medium for 2-3 times in a 15 mL

conical tube before being digested in 1 mL (or 0.1 mL/embryo, which ever that is larger) 0.25%

trypsin (25200-056, Gibco) with 0.5 mg/mL DNase (DN25-100MG, Sigma-Aldrich) in a 37°C

water bath for 15 minutes. After gently swirling the mixture every 5 minutes, enzymatic activity

was neutralized by adding an equal volume of plating medium comprised of DMEM/F12 (1:1)

(11330-032, Gibco), 0.1 mg/mL glucose (25-037-CI, Corning), B27 supplement (17504-04,

Gibco), and antibiotics (15240-062, Gibco). After mixing the content 3-5 times and waiting until

tissue chunks settled down to the bottom of the tube, the supernatant was carefully removed with

a pipette and slowly added 0.7-1.0 mL plating medium. Tissue was dissociated by gently

pipetting the tissue ~15 times with a flamed glass pasture pipette. The tube was kept undisturbed

for 1-2 minutes to allow undissociated tissue debris to settle down, and the upper phase

containing dissociated cells was collected to a new tube. The cell density was calculated using a

hemocytometer (#1492, Hausser Scientific), and then the cell suspension was diluted with the

plating medium to 1×10^6 cells/mL. The cells were seeded at a density of 120,000 cells/cm^2 on a

poly-D-lysine (P6407, Sigma-Aldrich; 200 mg/mL)-coated culture plate. After 4-6 hours, the

medium was replaced to the complete culture medium comprised of Neurobasal medium (21103-

049, Gibco), GlutaMAX Supplement (35050-51, Gibco), B27 supplement (17504-04, Gibco),

and antibiotics (15240-062, Gibco). To maintain the culture, half of the medium was changed

every 3 days. Viruses were added to seeded wells at 1:200 ratio by volume after 7-8 days *in vitro*

(DIV). Cells were harvested 72 hours after infection. RNA samples were extracted using the

PureLink RNA Mini Kit (#12183025, Thermo Fisher) with DNase (#79254, Qiagen). Protein

extracts were prepared using 1x sample buffer containing 10% 2-mercaptoethanol (0.1 mL/cm^2)

and centrifuged at 16,000 xg for 2 minutes. Supernatants were collected into new tubes and

subjected to Western blotting.

Stereotaxic injection

Mice were anaesthetized in an induction chamber (2-4% isoflurane) and placed in a stereotaxic

frame (Model 942, Kopf Instruments) where they were maintained under 1-2% isoflurane on a

37°C heating pad throughout the procedure. Following craniotomy under aseptic techniques, all

injections were done using a Hamilton Neuros Syringe (65460-03, Hamilton) driven by a Micro4

microsyringe pump controller (World Precision Instruments, Sarasota, FL), with the beveled

opening facing posterior. The syringe was slowly inserted into the brain tissue and dwelled in

place for 3 minutes before the injection started; after injection, the syringe was dwelled for

another 1 min/100 nL virus injected and then slowly withdrawn. The skin incision was closed

with 6-0 nylon sutures. For the VTA *Bdnf* knock-down experiments, mice were injected

bilaterally with 1 μL of either lenti-shRNA-*Bdnf*-eGFP or lenti-shRNA-*fLuc*-eGFP at the rate of

200 nL/min into the VTA (coordinates relative to Bregma: AP −3.05, ML ±0.3, DV −4.6). For

the VTA *Bdnf* overexpression experiments, mice were injected bilaterally with 0.5 μL of either

lenti-syn-*Bdnf*-eGFP or lenti-syn-eGFP at the rate of 100 nL/min into the VTA (coordinates

relative to Bregma: AP -3, ML ±0.5, DV -4.8). Immediately after the surgery, mice were allowed

to recover on a heated table (37°C) until becoming fully awake and then returned to their home

cages. All mice in the same cage were injected on the same day. Mice were given Carprofen

(Rimadyl) chewable tablet (0.5 mg/mouse/day) as analgesics in home cages for 3 days after

surgeries.

Electrophysiology

Mice were anaesthetized via i.p. injection of ketamine/xylazine/acepromazine mixture (69.57

mg/ml; 4.35mg/ml; 0.87mg/ml), then perfused with ice-cold slicing-aCSF consisting of 92 mM

N-methyl-d-glucose (NMDG), 2.5 mM KCl, 1.25 mM NaH_2PO_4, 10 mM $MgSO_4$, 20 mM

HEPES, 30 mM $NaHCO_3$, 25 mM glucose, 0.5 mM $CaCl_2$, 5 mM sodium ascorbate and 3 mM

sodium pyruvate, oxygenated with 95% O_2 and 5% CO_2. The pH and osmolarity were adjusted

to 7.3-7.4 and 315–320 mOsm, respectively. The brain was dissected and embedded with 2%

agarose made in slice-aCSF, and 350μm transverse slices containing VTA were cut using

vibratome (VF310-0Z, Precisionary Instruments, MA). Slices were first incubated in warm

(32°C) slicing-aCSF for 30 mins then transferred to holding-aCSF consisting of 92 mM NaCl,

2.5 mM KCl, 1.25 mM NaH_2PO_4, 30 mM $NaHCO_3$, 20 mM HEPES, 25 mM glucose, 2 mM

$MgSO_4$, 2 mM $CaCl_2$, 5 mM sodium ascorbate and 3 mM sodium pyruvate, oxygenated with

95% O_2 and 5% CO_2. The pH and osmolarity were adjusted to 7.3-7.4 and 310–315 mOsm,

respectively. The procedure of acute brain slice preparation was modified from our previous

study[52]. Electrophysiological recordings were made in a recording chamber mounted on upright

microscope (BX51WI, Olympus Optical Co., Ltd, Tokyo, Japan) with continuous perfusion of

warm (29–31°C) recording-aCSF consisting of 124 mM NaCl, 2.5 mM KCl, 1.25 mM NaH_2PO_4,

24 mM $NaHCO_3$, 5 mM HEPES, 12.5 mM glucose, 2 mM $MgCl_2$, 2 mM $CaCl_2$, oxygenated with

95% O_2 and 5% CO_2. The pH and osmolarity were adjusted to 7.3-7.4 and 310–305 mOsm,

respectively. VTA neurons were identified through a highspeed camera (ORCA-Flash4.0LT,

Hamamatsu Photonics, Shizuoka, Japan) coupled with water immersion objective lens

(LUMPLFLN-40xW, Olympus, Tokyo, Japan). All recordings were made through glass pipettes

pulled from borosilicate glass capillary (GC150F-10, Warner Instruments, Hamden, CT) with a

resistance around 6-9 MΩ when filled with intra-pipette solution consisting of 120 mM

potassium gluconate, 5 mM NaCl, 10 mM HEPES, 1.1 mM EGTA, 15 mM Phosphocreatine, 2

mM ATP and 0.3 mM GTP, with pH 7.2–7.3 and osmolality adjusted to 300 mOsm. Recordings

were discarded if the membrane potential (Vm) of recorded cell was over −40 mV or action

potentials did not overshoot 0 mV. Recorded cells were continuously clamped at -70 to -75mV

under current-clamp mode and received one second current injections from -100 to 400 pA with

10pA steps. Input resistance was calculated using the linear relationship of responses upon

current injections between -30 to 0 pA and action potential threshold was measured using the

first AP firing upon rheobase current injection. All data were collected using a Multiclamp 700B

amplifier (Molecular Devices, San Jose, CA) with a lowpass filter at 2 kHz and digitized at 10k

Hz through Axon Digidata 1550B interface (Molecular Devices, CA) coupled with Clampex

software (Molecular Devices, CA). Data were further analyzed using MATLAB (MathWorks,

MA).

Statistical analysis and graphing

Data are presented as mean ± standard error of the mean (SEM). Statistical significance was

taken as *p≤0.05, **p≤0.01, ***p≤0.001, and ****p≤0.0001, as determined by Student's *t*-test,

Holm-Šídák multiple comparison test, or two-way repeated-measure ANOVA, followed by post-

hoc multiple comparisons as appropriate. Significant and close-to-significance main effects of

factors, interactions, and pair-wise comparisons are labeled on the graphs. Sample sizes (N) were

determined based on empirical estimations. Data sorting and statistical analysis were performed

in Microsoft Excel, R studio, and GraphPad Prism 9.0 (Graphpad, La Jolla, CA). Graphs are

made with GraphPad Prism 9.0. Cartoons are created with BioRender.com and Microsoft

PowerPoint.

Lei et al.

655 **Data availability**

656 All data generated or analyzed during this study are included in this manuscript. Any additional

657 materials are available from the corresponding authors on reasonable request.

Lei et al.

References

1.   WHO. Mental health of older adults. https://www.who.int/news-room/fact-sheets/detail/mental-health-of-older-adults, 2017.

2.   Luo Y, Hawkley LC, Waite LJ, Cacioppo JT. Loneliness, health, and mortality in old age: a national longitudinal study. *Soc Sci Med* 2012; **74**(6)**:** 907-914.

3.   Robb CE, de Jager CA, Ahmadi-Abhari S, Giannakopoulou P, Udeh-Momoh C, McKeand J *et al.* Associations of Social Isolation with Anxiety and Depression During the Early COVID-19 Pandemic: A Survey of Older Adults in London, UK. *Front Psychiatry* 2020; **11:** 591120.

4.   Sayin Kasar K, Karaman E. Life in lockdown: Social isolation, loneliness and quality of life in the elderly during the COVID-19 pandemic: A scoping review. *Geriatr Nurs* 2021; **42**(5)**:** 1222-1229.

5.   Soutschek A, Bagaini A, Hare TA, Tobler PN. Reconciling psychological and neuroscientific accounts of reduced motivation in aging. *Soc Cogn Affect Neurosci* 2022; **17**(4)**:** 398-407.

6.   Mortby ME, Maercker A, Forstmeier S. Midlife motivational abilities predict apathy and depression in Alzheimer disease: the aging, demographics, and memory study. *J Geriatr Psychiatry Neurol* 2011; **24**(3)**:** 151-160.

7.   Cacioppo S, Capitanio JP, Cacioppo JT. Toward a neurology of loneliness. *Psychol Bull* 2014; **140**(6)**:** 1464-1504.

8.   Ieraci A, Mallei A, Popoli M. Social Isolation Stress Induces Anxious-Depressive-Like Behavior and Alterations of Neuroplasticity-Related Genes in Adult Male Mice. *Neural Plast* 2016; **2016:** 6212983.

9.   Novak CM, Burghardt PR, Levine JA. The use of a running wheel to measure activity in rodents: relationship to energy balance, general activity, and reward. *Neurosci Biobehav Rev* 2012; **36**(3)**:** 1001-1014.

10.   Meijer JH, Robbers Y. Wheel running in the wild. *Proc Biol Sci* 2014; **281**(1786).

11.   Acosta-Rodriguez V, Rijo-Ferreira F, Izumo M, Xu P, Wight-Carter M, Green CB *et al.* Circadian alignment of early onset caloric restriction promotes longevity in male C57BL/6J mice. *Science* 2022; **376**(6598)**:** 1192-1202.

12.   Bartling B, Al-Robaiy S, Lehnich H, Binder L, Hiebl B, Simm A. Sex-related differences in the wheel-running activity of mice decline with increasing age. *Exp Gerontol* 2017; **87**(Pt B)**:** 139-147.

Lei et al.

704 13. Tsibulsky VL, Norman AB. Methodological and analytical issues of progressive ratio
705   schedules: Definition and scaling of breakpoint. *J Neurosci Methods* 2021; **356:** 109146.
706

707 14. Husain M, Roiser JP. Neuroscience of apathy and anhedonia: a transdiagnostic approach.
708   *Nat Rev Neurosci* 2018; **19**(8)**:** 470-484.
709

710 15. Parker KE, Arackal JS, Hunter SC, McCall JG. Punishment Risk Task: Monitoring
711   anxiogenic states during goal directed actions in mice. *bioRxiv* 2022**:**
712   2022.2010.2030.514458.
713

714 16. Lerner TN, Holloway AL, Seiler JL. Dopamine, Updated: Reward Prediction Error and
715   Beyond. *Curr Opin Neurobiol* 2021; **67:** 123-130.
716

717 17. Tryon VL, Baker PM, Long JM, Rapp PR, Mizumori SJY. Loss of Sensitivity to
718   Rewards by Dopamine Neurons May Underlie Age-Related Increased Probability
719   Discounting. *Front Aging Neurosci* 2020; **12:** 49.
720

721 18. Fu Y, Yuan Y, Halliday G, Rusznak Z, Watson C, Paxinos G. A cytoarchitectonic and
722   chemoarchitectonic analysis of the dopamine cell groups in the substantia nigra, ventral
723   tegmental area, and retrorubral field in the mouse. *Brain Struct Funct* 2012; **217**(2)**:** 591-
724   612.
725

726 19. Naoi M, Maruyama W, Shamoto-Nagai M. Type A monoamine oxidase and serotonin are
727   coordinately involved in depressive disorders: from neurotransmitter imbalance to
728   impaired neurogenesis. *J Neural Transm (Vienna)* 2018; **125**(1)**:** 53-66.
729

730 20. Raitsin S, Tong J, Kish S, Xu X, Magomedova L, Cummins C *et al.* Subchronic
731   glucocorticoids, glutathione depletion and a postpartum model elevate monoamine
732   oxidase a activity in the prefrontal cortex of rats. *Brain Res* 2017; **1666:** 1-10.
733

734 21. Matthews GA, Nieh EH, Vander Weele CM, Halbert SA, Pradhan RV, Yosafat AS *et al.*
735   Dorsal Raphe Dopamine Neurons Represent the Experience of Social Isolation. *Cell* 2016;
736   **164**(4)**:** 617-631.
737

738 22. Lee CR, Matthews GA, Lemieux ME, Wasserlein EM, Borio M, Miranda RL *et al.*
739   Separable Dorsal Raphe Dopamine Projections mediate the Facets of Loneliness-like
740   state. *bioRxiv* 2025**:** 2025.2002.2003.636224.
741

742 23. Keifer J, Zheng Z, Ambigapathy G. A MicroRNA-BDNF Negative Feedback Signaling
743   Loop in Brain: Implications for Alzheimer's Disease. *Microrna* 2015; **4**(2)**:** 101-108.
744

745 24. Shrestha S, Phay M, Kim HH, Pouladvand P, Lee SJ, Yoo S. Differential regulation of
746   brain-derived neurotrophic factor (BDNF) expression in sensory neuron axons by
747   miRNA-206. *FEBS Open Bio* 2019; **9**(2)**:** 374-383.
748

25. Wang M, Li D, Yun D, Zhuang Y, Repunte-Canonigo V, Sanna PP *et al.* Translation of BDNF-gene transcripts with short 3' UTR in hippocampal CA1 neurons improves memory formation and enhances synaptic plasticity-relevant signaling pathways. *Neurobiol Learn Mem* 2017; **138:** 121-134.

26. Lau AG, Irier HA, Gu J, Tian D, Ku L, Liu G *et al.* Distinct 3'UTRs differentially regulate activity-dependent translation of brain-derived neurotrophic factor (BDNF). *Proc Natl Acad Sci U S A* 2010; **107**(36)**:** 15945-15950.

27. Ellis J. Silencing and variegation of gammaretrovirus and lentivirus vectors. *Hum Gene Ther* 2005; **16**(11)**:** 1241-1246.

28. Cabrera A, Edelstein HI, Glykofrydis F, Love KS, Palacios S, Tycko J *et al.* The sound of silence: Transgene silencing in mammalian cell engineering. *Cell Syst* 2022; **13**(12)**:** 950-973.

29. Cooray S, Howe SJ, Thrasher AJ. Retrovirus and lentivirus vector design and methods of cell conditioning. *Methods Enzymol* 2012; **507:** 29-57.

30. Pu L, Liu QS, Poo MM. BDNF-dependent synaptic sensitization in midbrain dopamine neurons after cocaine withdrawal. *Nat Neurosci* 2006; **9**(5)**:** 605-607.

31. Cordeira JW, Frank L, Sena-Esteves M, Pothos EN, Rios M. Brain-derived neurotrophic factor regulates hedonic feeding by acting on the mesolimbic dopamine system. *J Neurosci* 2010; **30**(7)**:** 2533-2541.

32. Parker KE, Pedersen CE, Gomez AM, Spangler SM, Walicki MC, Feng SY *et al.* A Paranigral VTA Nociceptin Circuit that Constrains Motivation for Reward. *Cell* 2019; **178**(3)**:** 653-671 e619.

33. Vargas-Perez H, Ting AKR, Walton CH, Hansen DM, Razavi R, Clarke L *et al.* Ventral tegmental area BDNF induces an opiate-dependent-like reward state in naive rats. *Science* 2009; **324**(5935)**:** 1732-1734.

34. Vargas-Perez H, Bahi A, Bufalino MR, Ting AKR, Maal-Bared G, Lam J *et al.* BDNF signaling in the VTA links the drug-dependent state to drug withdrawal aversions. *J Neurosci* 2014; **34**(23)**:** 7899-7909.

35. Remy P, Doder M, Lees A, Turjanski N, Brooks D. Depression in Parkinson's disease: loss of dopamine and noradrenaline innervation in the limbic system. *Brain* 2005; **128**(Pt 6)**:** 1314-1322.

36. Chaudhuri KR, Schapira AH. Non-motor symptoms of Parkinson's disease: dopaminergic pathophysiology and treatment. *Lancet Neurol* 2009; **8**(5)**:** 464-474.

37. Schiffino FL, Siemian JN, Petrella M, Laing BT, Sarsfield S, Borja CB *et al.* Activation of a lateral hypothalamic-ventral tegmental circuit gates motivation. *PLoS One* 2019; **14**(7)**:** e0219522.

38. Satoh A, Brace CS, Rensing N, Cliften P, Wozniak DF, Herzog ED *et al.* Sirt1 extends life span and delays aging in mice through the regulation of Nk2 homeobox 1 in the DMH and LH. *Cell Metab* 2013; **18**(3)**:** 416-430.

39. Ito N, Takatsu A, Ito H, Koike Y, Yoshioka K, Kamei Y *et al.* Slc12a8 in the lateral hypothalamus maintains energy metabolism and skeletal muscle functions during aging. *Cell Rep* 2022; **40**(4)**:** 111131.

40. Tokizane K, Brace CS, Imai SI. DMH(Ppp1r17) neurons regulate aging and lifespan in mice through hypothalamic-adipose inter-tissue communication. *Cell Metab* 2024; **36**(2)**:** 377-392 e311.

41. Yoon MJ, Yoshida M, Johnson S, Takikawa A, Usui I, Tobe K *et al.* SIRT1-Mediated eNAMPT Secretion from Adipose Tissue Regulates Hypothalamic NAD+ and Function in Mice. *Cell Metab* 2015; **21**(5)**:** 706-717.

42. Yoshida M, Satoh A, Lin JB, Mills KF, Sasaki Y, Rensing N *et al.* Extracellular Vesicle-Contained eNAMPT Delays Aging and Extends Lifespan in Mice. *Cell Metab* 2019; **30**(2)**:** 329-342 e325.

43. Johnson S, Yoshioka K, Brace CS, Imai S. Quantification of localized NAD+ changes reveals unique specificity of NAD+ regulation in the hypothalamus. *npj Aging* 2023; **In press**.

44. Oh H, Lewis DA, Sibille E. The Role of BDNF in Age-Dependent Changes of Excitatory and Inhibitory Synaptic Markers in the Human Prefrontal Cortex. *Neuropsychopharmacology* 2016; **41**(13)**:** 3080-3091.

45. Buchman AS, Yu L, Boyle PA, Schneider JA, De Jager PL, Bennett DA. Higher brain BDNF gene expression is associated with slower cognitive decline in older adults. *Neurology* 2016; **86**(8)**:** 735-741.

46. Erickson KI, Miller DL, Roecklein KA. The aging hippocampus: interactions between exercise, depression, and BDNF. *Neuroscientist* 2012; **18**(1)**:** 82-97.

47. Erickson KI, Prakash RS, Voss MW, Chaddock L, Heo S, McLaren M *et al.* Brain-derived neurotrophic factor is associated with age-related decline in hippocampal volume. *J Neurosci* 2010; **30**(15)**:** 5368-5375.

48. Clair R, Gordon M, Kroon M, Reilly C. The effects of social isolation on well-being and life satisfaction during pandemic. *Hum Soc Sci Commun* 2021; **8**(1).

Lei et al.

840   49.   Richardson NR, Roberts DC. Progressive ratio schedules in drug self-administration
841   studies in rats: a method to evaluate reinforcing efficacy. *J Neurosci Methods* 1996; **66**(1)**:**
842   1-11.
843

844   50.   Thompson LH, Parish CL. Transplantation of fetal midbrain dopamine progenitors into a
845   rodent model of Parkinson's disease. *Methods in molecular biology (Clifton, NJ)* 2013;
846   **1059:** 169-180.
847

848   51.   Lotharius J, Dugan LL, O'Malley KL. Distinct mechanisms underlie neurotoxin-mediated
849   cell death in cultured dopaminergic neurons. *J Neurosci* 1999; **19**(4)**:** 1284-1293.
850

851   52.   Kuo CC, Hsieh JC, Tsai HC, Kuo YS, Yau HJ, Chen CC *et al.* Inhibitory interneurons
852   regulate phasic activity of noradrenergic neurons in the mouse locus coeruleus and
853   functional implications. *J Physiol* 2020; **598**(18)**:** 4003-4029.
854
855

Lei et al.

Acknowledgments

We thank Erik D. Herzog and Albert A. Davis for sharing their wheel-running facility and

providing critical suggestions on our study and for helping with the primary mesencephalon

culture, respectively. We also thank members of the Imai lab, particularly Kyohei Tokizane,

Cynthia S. Brace, and Brian V. Lananna, and of the Al-Hasani and McCall labs for technical

support and valuable discussions. Support was also provided by the Animal Behavior Core

(supported in part by the McDonnell Center for Systems Neuroscience and the Taylor Family

Institute) and the Hope Center Viral Vectors Core at Washington University School of Medicine.

This work was mainly supported by grants to S.I. from the National Institute on Aging

(AG037457, AG047902), and the Tanaka Fund at Washington University School of Medicine.

J.G.M. was supported by grants from the National Institute of Neurological Disorders and Stroke

(NS11789) and the Brain & Behavior Research Foundation (NARSAD YI-28565). H.C.L. was

supported by the LIFENAD Fellowship Fund.

Author information

Authors and Affiliations

Hanyue Cecilia Lei and Shin-ichiro Imai

Department of Developmental Biology, Washington University School of Medicine, St. Louis,

Missouri, USA

Kyle E. Parker, Chao-Cheng Kuo, and Jordan G. McCall

Department of Anesthesiology, Washington University School of Medicine, St. Louis, Missouri,

USA

Carla M. Yuede

Department of Psychiatry, Washington University School of Medicine; Saint Louis, Missouri,

USA.

Contributions

H.C.L. and S.I conceived the project. J.G.M. and S.I. acquired funding and supervised the project.

H.C.L., K.E.P., C.M.Y., C.C.K., J.G.M. and S.I. designed experiments, analyzed data, and wrote

the manuscript. H.C.L. conducted most experiments, H.C.L. and K.E.P. conducted the *Bdnf*

overexpression experiment *in vivo*, C.M.Y. conducted the 1-hour LA/OFT, and C.C.K.

conducted the VTA electrophysiology. All authors contributed to the final version of the

manuscript.

Corresponding author

Correspondence to Shin-ichiro Imai and Jordan G. McCall.

Ethics declarations

Competing interests

S.I. declares no Competing Non-Financial Interests but the following Competing Financial

Interests: S.I. receives a part of patent-licensing fees from MetroBiotech (USA) and the Institute

for Research on Productive Aging (Japan) through Washington University. S.I. also serves as the

President of the Institute for Research on Productive Aging (Japan) and a co-CEO of LongGen

Bioscience (Japan). S.I.'s External Professional Activities (EPAs) have already been reported to,

and a potential conflict of interest has been properly resolved through the Washington University

Lei et al.

901 Conflict of Interest Committee. The other authors have no Competing Financial or Non-

902 Financial Interests.

Fig. 1. Chronic social isolation and aging differentially impair motivational behaviors and induce depressive- and anxiety- like behaviors

Young (5-7 months) and old (19-21 months) mice were assessed by batteries of behavior tests after a month of social isolation or grouped housing. (**a**) Voluntary wheel-running activity recording in consecutive 48 hours, shaded area representing the dark phase. (**b**) Average VWR activity during the dark phase (ZT12-0). (**c**) Calendar illustrating the operant progressive ratio test protocol. Correct nosepokes performed (**d**) and break points reached (**e**) in the PR test. (**f**) Delta of sucrose solution intake to baseline water intake in the sucrose preference test. (**g**) Percentage of time spent in the center area in the 1-hour locomotor activity and open field test (LA/OFT). Data were analyzed by 3-way ANOVA (**a**) and 2-way ANOVA (**b**, **d**, **e**, **f**, **g**) with *post-hoc* multiple comparisons. n=7-20. Error bars indicate SEM. Significant pair-wise comparisons and main effects of factors and interactions are labeled on the graphs: *$p \leq 0.05$; **$p \leq 0.01$; ***$p \leq 0.001$; ****$p \leq 0.0001$.

Fig. 2. Aging leads to reduction of fasting-induced motivation for palatable food reward
Young (5 months) and old (19 months) mice were examined for motivation levels. (**a**) Calendar illustrating the extended operant progressive ratio test protocol. Correct nosepokes (**b**) and break points reached (**c**) normalized to individual baseline PR under different experimental conditions in the extended PR. Data were analyzed by 2-way ANOVA with *post-hoc* multiple comparisons. n=8-9. Error bars indicate SEM. Significant pair-wise comparisons and main effects of factors and interactions are labeled on the graphs: *p≤0.05; **p≤0.01; ***p≤0.001; ****p≤0.0001.

Fig. 3. *Bdnf* gene expression is decreased in the ventral tegmental area (VTA) of aged mice
RT-qPCR analysis of laser-capture-microdissected VTA (**a**) and SN (**b**) samples from young (7 months) and old (21 months) mice. Representative images of FISH (RNAscope) analysis of *Bdnf* in VTA, using dopamine active transporter (DAT/*Slc6a3*) mRNA as a marker for DA neurons in young (7 month) vs. old mice (22 months) are shown in (**c**) and quantified as the density of *Bdnf* mRNA signal puncta in the VTA region (**d**, **e**) across 8 bregma levels (80μm interval from -3.05). Western blots (**f**) and quantifications (**g**) of the precursor and matured forms of the BDNF protein (proBDNF and mBDNF, respectively) normalized to β-actin signals in punched VTA of young (4 months) vs. old (23 months) mice. *Gria1*: glutamate ionotropic receptor AMPA type subunit 1; *Slc6a3*: solute carrier family 6 member 3. Data were analyzed by 2-way ANOVA with *post-hoc* multiple comparisons (**a**, **b**, **d**, **g**) and student's t-test (**e**). n= 3-6. Error bars indicate SEM. Significant pair-wise comparisons and main effects of factors and interactions are labeled on the graphs: *p≤0.05; **p≤0.01; ***p≤0.001; ****p≤0.0001.

Fig. 4. VTA *Bdnf* KD in young mice partially recapitulates the age-associated reduction in palatable food motivation, not affecting other age-related phenotypes

(a) Calendar and schematics illustrating the experimental design. (**b**) Representative FISH images of *Bdnf* and Slc6a3 in VTA of *fLuc* vs *Bdnf*-shRNA virus injected mice after behavior tests, (**c**) quantified as the density of *Bdnf* mRNA signal puncta in infected regions (KD efficiency 27%).

Western blots (**d**) and quantifications (**e, f**) of proBDNF and mBDNF levels normalized to β-Actin signals in punched VTA and NAc of *fLuc* vs *Bdnf*-shRNA virus injected mice. Correct nosepokes

(**g**) and break points reached (**h**) normalized to individual baseline PR under different
experimental conditions in the extended PR. (**i**) Average VWR activity during the dark phase
(ZT12-0). Total distance traveled (**j**), percentage of time spent in the center area (**k**) and
percentage of time spent moving or resting in the center and peripheral areas (**l**) in the open field
test. Data were analyzed by 2-way ANOVA with *post-hoc* multiple comparisons (**e**, **f**, **g**, **h**,),
student's t-test (**c**, **i**, **j**, **k**), and Holm-Šídák multiple comparison test (**l**). n=4-18. Error bars
indicate SEM. Significant pair-wise comparisons and main effects of factors and interactions are
labeled on the graphs: *p≤0.05; **p≤0.01; ***p≤0.001; ****p≤0.0001.