TITLE:

Immunogenicity and safety in pigs of PHH-1V, a SARS-CoV-2 RBD fusion
heterodimer vaccine candidate

Authors: Alexandra Moros[1*], Antoni Prenafeta[1*], Antonio Barreiro[1], Eva Perozo[1], Alex
Fernández[1], Manuel Cañete[1], Luis González[1], Carme Garriga[1], Edwards Pradenas[2], Silvia
Marfil[2], Julià Blanco[2,3], Paula Cebollada Rica[4], Marta Sisteré-Oró[4], Andreas Meyerhans[4,5],
Teresa Prat Cabañas[1], Ricard March[1], Laura Ferrer[1]

*These authors contributed equally

Affiliations:

[1]HIPRA. Avda. La Selva, 135, 17170 Amer (Girona), Spain.

[2]IrsiCaixa. AIDS Research Institute, Germans Trias i Pujol Research Institute (IGTP), Can Ruti
Campus, UAB, 08916 Badalona, Spain.

[3]University of Vic–Central University of Catalonia (UVic-UCC), 08500, Vic, Catalonia, Spain.

[4]Infection Biology Laboratory, Department of Experimental and Health Sciences, Universitat
Pompeu Fabra (UPF), 08003 Barcelona, Spain.

[5]ICREA (Catalan Institution for Research and Advanced Studies), Pg. Lluís Companys 23,
08010 Barcelona, Spain.

Corresponding author: sandra.moros@hipra.com, antoni.prenafeta@hipra.com

WORD COUNT (Short communication):

ABSTRACT: 146 words

MANUSCRIPT (excluding abstract, captions, references): 1,762 words

REFERENCES: 18 references

FIGURES/TABLES: 3 figures

ABSTRACT

The continuing high global incidence of COVID-19 and the undervaccinated status of billions of persons strongly motivate the development of a new generation of efficacious vaccines. We have developed an adjuvanted vaccine candidate, PHH-1V, based on a protein comprising the receptor binding domain (RBD) of the Beta variant of SARS-CoV-2 fused in tandem with the equivalent domain of the Alpha variant, with its immunogenicity, safety and efficacy previously demonstrated in mouse models. In the present study, we immunized pigs with different doses of PHH-1V in a prime-and-boost scheme showing PHH-1V to exhibit an excellent safety profile in pigs and to produce a solid RBD-specific humoral response with neutralising antibodies to 7 distinct SARS-CoV-2 variants of concern, with the induction of a significant IFNγ^+ T-cell response. We conclude that PHH-1V is safe and elicits a robust immune response to SARS-CoV-2 in pigs, a large animal preclinical model.

KEYWORDS:

SARS-CoV-2; COVID-19; RBD; recombinant vaccine; humoral immune response; IFNγ.

INTRODUCTION

Coronavirus disease-19 (COVID-19) was declared a pandemic by the World Health Organization (WHO) in March 2020 due to its widespread transmission, prompting a global effort to develop safe and effective vaccines against SARS-CoV-2, with more than 240 candidate vaccines currently under clinical development [1]. These include messenger RNA, protein-based and viral vector-based vaccines targeting the viral Spike protein or its Receptor Binding Domain (RBD), and conventional inactivated vaccines targeting the entire virus [2]. Several vaccines have shown great efficacy in protecting from severe disease, and have been deployed and administered in an unprecedented global vaccination effort. However, the continued emergence of new viral variants with increasing immune escape and transmissibility characteristics compromises the intended control of the pandemic [3]. In addition, an unequal distribution of currently available vaccines is present, with over 80% of the population in low- and middle-income countries not having received any SARS-CoV-2 vaccine dose (as of July 2022) [1].

With these motivations, we have developed the adjuvanted PHH-1V vaccine candidate based on a fusion heterodimer protein containing, in tandem, the RBD of the B.1.351 (Beta) and B.1.1.7 (Alpha) SARS-CoV-2 variants at its N- and C-terminus, respectively. As a result, a key feature of PHH-1V distinguishing it from other adjuvanted protein-based subunit vaccines is the inclusion of sequences corresponding to two heterologous variants of concern (VOCs) as part of the same protein molecule. The safety, immunogenicity and efficacy of the PHH-1V vaccine candidate were previously tested in mouse models [4]. PHH-1V is currently under rolling review by the European Medicines Agency (EMA).

In the present study we have further investigated its safety and immunogenicity in pigs - a large animal model involving an immune system with physiologically relevant similarities to humans [5].

MATERIALS AND METHODS

PHH-1V vaccine

The PHH-1V vaccine is composed of a recombinant protein formed by the fusion of the RBD domain of the Beta variant of SARS-CoV-2 (B.1.351) at its N-terminus and the RBD domain of the Alpha variant (B.1.1.7) at its C-terminus, expressed in CHO cells with stable expression (clone 57), adjuvanted with SQBA, an oil-in-water emulsion based on squalene, as previously described elsewhere [4]. Phosphate-buffered saline (PBS) was inoculated as a negative control.

Animal study design

Large White x Landrace hybrid pigs were used as a model to assess immune responses to PHH-1V. Twenty-four pigs (12 females and 12 males) were distributed into three dosage groups (Group A: 10 µg; Group B: 20 µg; Group C: 40 µg), and one group of four pigs (2 females and 2 males) received PBS (Group D) **(Figure 1)**. In a second arm of the study for safety and cellular immune response analysis, 10 male pigs were distributed into one group of 6 pigs administered 40 µg PHH-1V per dose (Group 1) and a second group of 4 pigs administered PBS (Group 2). Pigs received an intramuscular injection in a total volume of 0.5 mL of test product on days 0 (D0) and 21 (D21) in both arms of the study **(Figure 1)**. Safety was assessed by monitoring general clinical signs (recorded daily) and rectal temperatures (recorded one day before each dosage, at vaccination, 4 and 6 hours after that, and daily for 5 days after the first injection and 2 days after the second injection) and local reactions (inflammation and nodules) at the injection site (observed daily) throughout the study.

***Ex vivo* assays**

Materials and methods for ELISA determinations of SARS-CoV-2 RBD-binding antibodies, pseudovirus-based neutralisation assays (PBNA), and RBD-induced IFNγ ELISpot assay are explained in detail in **Supplementary Data**.

Statistical analysis

Statistical analyses and plots were generated using R (version 4.0.5) or GraphPad Prism (version 9). Unless otherwise specified, all plots depict individual data points for each animal, along with the sample mean and standard deviation. When required, data was either $\log_{10}$- or arcsine-transformed (i.e. log-normal and percentage variables, respectively). For testing the effect of one or two factors, linear models were generally employed: generalized least squares or mixed effects models depending on the dependence of the observations. Parameter estimation was performed using restricted maximum likelihood and assumptions were tested graphically.

107   Statistically significant differences between groups are indicated with a line on top of each

108   group: ** $p<0.01$; * $p<0.05$; [+] $0.05<p<0.1$.

RESULTS

PHH-1V displays a good safety profile in pigs

Rectal temperatures showed no differences between groups. Local reactions at the injection site or general clinical signs were not observed in any animal throughout the study (data not shown).

PHH-1V elicits a robust humoral response

After acclimation, pigs were immunized (day 0) and boosted (day 21) with 3 different doses of PHH-1V. Seven days after the second dose (day 28), the sera from all vaccination groups had significant levels of RBD-specific IgG antibodies (**Figure 2A and Supplementary Table 1**). In all dosage groups, the antibody titres remained at stably elevated levels until the end of the study on day 42, with highly significant differences compared to control animals (**Figure 2A**).

PHH-1V elicits neutralising antibodies to all SARS-CoV-2 variants of concern

A prime-boost of PHH-1V (40 µg/dose) elicited significant titres of neutralising antibodies (nAbs) against all VOCs analysed compared to non-vaccinated animals in sera collected on day 42 post-prime dose (**Figure 2B**). When compared to the Alpha and Beta variants, the neutralising titre of the sera was slightly reduced against Gamma, Mu, Delta and Omicron BA.1 variants (**Figure 2B and Supplementary Table 2**).

PHH-1V elicits a robust cellular immune response

To determine the cellular immune response (second arm of the study), peripheral blood mononuclear cells (PBMCs) were isolated on day 40 and tested by IFNγ ELISpot *ex vivo* for their reactivity towards a pool of RBD peptides representing Alpha and Beta SARS-CoV-2 variant epitopes. The PBMCs from vaccinated pigs (40 µg RBD fusion heterodimer/dose) showed a significant higher percentage of IFNγ^+ T cells than in control animals after *in vitro* re-stimulation (**Figure 3A**).

In addition, in PHH-1V immunised pigs a tendency was observed in the *in vitro* stimulation of PBMCs by predominantly activating antigen-specific CD4$^+$ T helper cells and CD8$^+$ T cytotoxic cells expressing Th1 cytokines (IFNγ and/or IL-2 and/or TNF-α) at 19 days after the second dose (**Figures 3B and 3C**).

DISCUSSION

An unprecedented effort has enabled the development of a substantial number of vaccines and several therapies against SARS-CoV-2, a novel virus that has caused an enormous impact on lives [1]. Many vaccine candidates have been designed, using broadly diverse platforms and technologies, which has afforded the production of the large numbers of doses required for mass vaccination programs. Despite the phenomenal success and effectiveness of several of these vaccines, the continuing pandemic, the insufficient distribution of vaccines to many developing countries, and the emergence of new immune escape variants endowed with increasing transmissibility [6], must be urgently addressed with an end to the pandemic as a goal - including the development of a new generation of affordable vaccines featuring higher efficacy against emerging variants and improved safety [3].

In the present study we have examined the safety and immunogenic potential of the RBD subunit-based vaccine candidate, PHH-1V, in pigs - a large animal presenting an immune system with significant similarities to humans [5], and a useful pre-clinical model to for assessing the immunogenicity of upcoming SARS-CoV-2 vaccine candidates [7]. We have found that two vaccination doses are safe and induce high and specific humoral and cellular immune responses. As expected, PHH-1V elicited strong RBD-binding IgG antibodies after two doses. Notably, it also generated significant neutralisation activity, determined by pseudovirus-based assays, against all tested VOCs (Alpha, Beta, Gamma, Delta and Omicron BA.1). The results of ELISpot assaying showed that 40 µg of PHH-1V induces significantly higher percentage of T-cells expressing IFNγ than mock-vaccinated pigs - the 40-µg dose being a proper dose candidate to induce significant cellular responses.

Neutralising antibody levels correlate well with vaccine protective efficacy [8]. As the RBD bears immunodominant neutralisation epitopes [9], a vaccine consisting of RBD, rather than a full-length S protein, is expected to induce more effectively RBD-focused antibodies [10, 11]. A second critical feature of PHH-1V is the simultaneous presence in the same protein of RBDs for two heterologous VOCs, Beta and Alpha, in a two-domain tandem fusion configuration. [12]We have found that PHH-1V induces neutralising antibodies that are cross-reactive against all VOCs, including the highly immunoevasive Beta, Gamma and Omicron BA.1 variants, after two doses. The presence of known immune escape mutations on PHH-1V, such as K417N, E484K or N501Y [13, 14], may be partially responsible for the neutralisation cross-reactivity observed against variants sharing these mutations, such as Alpha (N501Y), Beta (K417N, E484K, N501Y), Gamma (E484K, N501Y), Mu (E484K) or Omicron (K417N, N501Y), although epitopes not containing such immune escape mutations must play a role in eliciting neutralising antibodies against the delta SARS-CoV-2 variant.

It has been shown that a third ("booster") vaccine dose, administered in homologous or heterologous immunization schedules, can at least partially restore neutralisation activity to Omicron [15]. The ability of a booster dose to restore the neutralising activity against Omicron

is not limited to mRNA-based or vector-based vaccines and has also been observed for
inactivated and subunit vaccines or a combination of infection with prior VOCs and vaccination
[16]. Such cross-reactive responses may be explained by the expansion of persisting clones of
memory B cells, including a response to more conserved regions of the RBD, after three
exposures to the SARS-CoV-2 Spike [17]. Given the relative maintenance of neutralisation
potency against Omicron BA.1 and Delta provided by PHH-1V we anticipate that a third or
booster dose of this vaccine, included as part of a heterologous vaccine scheme, is likely to elicit
potent neutralisation and protective immunity against these variants.

A limitation of the present study is that a challenging analysis after the PHH-1V vaccine is not
feasible, as the pig model is not susceptible to SARS-CoV-2 [18]. Nevertheless, our previous
challenge studies in mice models and non-human primates have demonstrated efficacy, safety
and protection from the viral challenge of PHH-1V [4]. Future studies should address additional
relevant aspects, such as the immune response to booster doses or the induction of B and T cell
memory.

192 CONCLUSIONS

In conclusion, PHH-1V exhibited an excellent safety profile and a strong and specific immune
response in pigs, showing a broadly cross-reactive neutralisation activity against heterologous
VOCs, and IFNγ^+ T-cells activation. These results and those observed in other pre-clinical models,
support the implementation of the PHH-1V vaccine as a primary or booster vaccination schedule
in humans to prevent the consequences of SARS-CoV-2 infection. Further Phase IIb and III clinical
trials are ongoing to support the potential of this candidate vaccine as a heterologous booster
(NCT05142553, NCT05246137).

FIGURE LEGENDS

Figure 1. Experimental procedure schematics. Two-dose prime-and-boost intra-muscular immunization on days 0 and 21 (0.5 mL/dose), with a 42-day follow-up. For safety and humoral response monitoring, 3 experimental groups (A-C) were inoculated (n = 6/group; 3 female and 3 male) with 10 μg, 20 μg or 40 μg of PHH-1V. A control group (D; n = 4; 2 female and 2 male) was inoculated with PBS. For safety and cellular response monitoring, pigs of group 1 (n = 6; 6 male) were inoculated with 40 μg of PHH 1V. Control group 2 (n = 4; 4 male) was inoculated with PBS.

Figure 2. Vaccination of pigs with PHH-1V elicits SARS-CoV-2 RBD-specific humoral and neutralising antibody responses against SARS-CoV-2 variants of concern (VOCs). A. Longitudinal analysis of RBD-specific IgG determined in samples collected on days 0, 21, 28, 35 and 42. Group A: 10 μg PHH-1V RBD fusion heterodimer; Group B: 20 μg PHH-1V RBD fusion heterodimer; Group C: 40 μg PHH-1V RBD fusion heterodimer; D: Control PBS. Each data point represents an individual pig with bars denoting the mean ± standard deviation per group. IgG longitudinal data have been analysed by means of a linear mixed effects model of repeated measures with the $\log_{10}$-transformed titres. * $p<0.05$; ** $p<0.01$.**B.** Neutralisation activity by sera collected on day 42 after the first vaccine dose from Group C (40 μg PHH-1V) vs. Group D (PBS) against 7 different VOCs: α (B.1.1.7), β (B.1.351), γ (P.1), μ (B.1.621), δ (B.1.617.2), δ+ (B.1.617.2.1) and o (BA.1). Pseudovirus neutralisation assays were conducted, yielding titres expressed as $\log_{10}$ IC_{50}. Each data point represents an individual pig with bars denoting the mean ± standard deviation per group. PBNA IC_{50} longitudinal data have been analysed by means of a linear mixed effects model of repeated measures with the $\log_{10}$-transformed titres. ** $p<0.01$. Titre values are provided in **Table S2**.

Figure 3. Vaccination of pigs with PHH-1V elicits cellular response. A. Induction of IFNγ$^+$ T-cells determined by ELISpot assays in PBMCs from vaccinated pigs, sampled on day 40 and stimulated *ex vivo* with an RBD peptide pool representing Alpha and Beta SARS-CoV-2 variant epitopes. Group 1: 40 μg PHH-1V RBD fusion heterodimer; Group 2: Control PBS. Each data point represents an individual pig with bars denoting the mean ± standard deviation per group. ELISpot data have been analysed by means of a generalised least squares model with the arcsine-square root-transformed proportion of IFN γ -positive cells * p<0.05. **B-C.** CD4$^+$ (**B**) and CD8$^+$ **(C)** T-cell response was measured by intracellular cytokine staining (ICS) after the stimulation of PMBCs with a pool of peptides from SARS-CoV-2 RBD of Alpha and Beta variants. Cytokine expression of PBMCs stimulated with medium was considered as background and subtracted from the responses measured from the RBD peptide pool for each individual pig. Group 1: PHH-1V vaccine 40 μg RBD fusion heterodimer/dose; Group 2: control group (PBS). Each data point represents an individual pig with bars denoting the mean ± standard deviation per group. ICS

238   data have been analysed by means of a generalised least squares model with the arcsine-square

239   root-transformed proportion of positive cells [+] $0.05 < p < 0.1$.

ACKNOWLEDGEMENTS

Anna Moya and Mireia Muntada for the ELISA analysis; Clara Panosa and Ester Puigvert for her assistance in the production of the vaccine antigen; Glòria Pujol and Eduard Fossas for their assistance in review of the manuscript; and Alicia Subtil-Rodríguez from Dynamic S.L.U. (Evidenze Clinical Research, Madrid, Spain) for providing medical writing support during the preparation of this paper funded by Hipra Scientific, S.L.U.

FUNDING

This project was partially funded by the Centre for the Development of Industrial Technology (CDTI, IDI20210115), a public organisation answering to the Spanish Ministry of Science and Innovation.

DECLARATION OF COMPETING INTEREST

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: A.Mo., A.P., A.B., E.P., A.F., M.C., L.G., C.G., T.P.C., R.M., and L.F. are employees of HIPRA. E.P. reports travel reimbursement from Gilead. J.B. reports consultancy fees from Albajuna Therapeutics; grants (outside the submitted work) from HIPRA, Grifols, Nesapor, and MSD; and speaking fees and travel reimbursement from Gilead. P.C.R., M.S.-O., and A.Me. reports grants (outside the submitted work) from HIPRA. S.M. declares no conflicts of interest.

REFERENCES

[1] WHO. WHO Coronavirus (COVID-19) Dashboard. 2022. https://covid19.who.int/. (Accesed July 4th, 2022)

[2] Mellet J, Pepper MS. A COVID-19 Vaccine: Big Strides Come with Big Challenges. Vaccines (Basel). 2021;9. https://doi.org/10.3390/vaccines9010039.

[3] Altmann DM, Boyton RJ. COVID-19 vaccination: The road ahead. Science. 2022;375:1127-32. https://doi.org/10.1126/science.abn1755.

[4] Barreiro A, Prenafeta A, Bech-Sabat G, Roca M, March R, González L, et al. Preclinical efficacy, safety, and immunogenicity of PHH-1V, a second-generation COVID-19 vaccine candidate based on a novel recombinant RBD fusion heterodimer of SARS-CoV-2. bioRxiv. 2021:2021.11.22.469117. https://doi.org/10.1101/2021.11.22.469117.

[5] Pabst R. The pig as a model for immunology research. Cell Tissue Res. 2020;380:287-304. https://doi.org/10.1007/s00441-020-03206-9.

[6] DeGrace MM, Ghedin E, Frieman MB, Krammer F, Grifoni A, Alisoltani A, et al. Defining the risk of SARS-CoV-2 variants on immune protection. Nature. 2022;605:640-52. https://doi.org/10.1038/s41586-022-04690-5.

[7] Lara-Puente JH, Carreño JM, Sun W, Suárez-Martínez A, Ramírez-Martínez L, Quezada-Monroy F, et al. Safety and Immunogenicity of a Newcastle Disease Virus Vector-Based SARS-CoV-2 Vaccine Candidate, AVX/COVID-12-HEXAPRO (Patria), in Pigs. mBio. 2021;12:e0190821. https://doi.org/10.1128/mBio.01908-21.

[8] Khoury DS, Cromer D, Reynaldi A, Schlub TE, Wheatley AK, Juno JA, et al. Neutralizing antibody levels are highly predictive of immune protection from symptomatic SARS-CoV-2 infection. Nat Med. 2021;27:1205-11. https://doi.org/10.1038/s41591-021-01377-8.

[9] Goel RR, Painter MM, Apostolidis SA, Mathew D, Meng W, Rosenfeld AM, et al. mRNA vaccines induce durable immune memory to SARS-CoV-2 and variants of concern. Science. 2021;374:abm0829. https://doi.org/10.1126/science.abm0829.

[10] Burton DR, Walker LM. Rational Vaccine Design in the Time of COVID-19. Cell Host Microbe. 2020;27:695-8. https://doi.org/10.1016/j.chom.2020.04.022.

[11] Liu Z, Xu W, Xia S, Gu C, Wang X, Wang Q, et al. RBD-Fc-based COVID-19 vaccine candidate induces highly potent SARS-CoV-2 neutralizing antibody response. Signal Transduct Target Ther. 2020;5:282. https://doi.org/10.1038/s41392-020-00402-5.

[12] Wang CY, Hwang KP, Kuo HK, Peng WJ, Shen YH, Kuo BS, et al. A multitope SARS-CoV-2 vaccine provides long-lasting B cell and T cell immunity against Delta and Omicron variants. J Clin Invest. 2022;132. https://doi.org/10.1172/jci157707.

[13] Jangra S, Ye C, Rathnasinghe R, Stadlbauer D, Krammer F, Simon V, et al. SARS-CoV-2 spike E484K mutation reduces antibody neutralisation. Lancet Microbe. 2021;2:e283-e4. https://doi.org/10.1016/s2666-5247(21)00068-9.

[14] Liu Y, Liu J, Plante KS, Plante JA, Xie X, Zhang X, et al. The N501Y spike substitution enhances SARS-CoV-2 infection and transmission. Nature. 2022;602:294-9. https://doi.org/10.1038/s41586-021-04245-0.

[15] Debes AK, Xiao S, Egbert ER, Caturegli P, Sitaras I, Pekosz A, et al. Comparison of total and neutralizing SARS-CoV-2 spike antibodies against omicron and other variants in paired samples after two or three doses of mRNA vaccine. medRxiv. 2022. https://doi.org/10.1101/2022.01.26.22269819.

[16] Wratil PR, Stern M, Priller A, Willmann A, Almanzar G, Vogel E, et al. Three exposures to the spike protein of SARS-CoV-2 by either infection or vaccination elicit superior neutralizing immunity to all variants of concern. Nat Med. 2022;28:496-503. https://doi.org/10.1038/s41591-022-01715-4.

[17] Goel RR, Painter MM, Lundgreen KA, Apostolidis SA, Baxter AE, Giles JR, et al. Efficient recall of Omicron-reactive B cell memory after a third dose of SARS-CoV-2 mRNA vaccine. Cell. 2022;185:1875-87.e8. https://doi.org/10.1016/j.cell.2022.04.009.

311   [18] Vergara-Alert J, Rodon J, Carrillo J, Te N, Izquierdo-Useros N, Rodríguez de la Concepción
312   ML, et al. Pigs are not susceptible to SARS-CoV-2 infection but are a model for viral
313   immunogenicity studies. Transbound Emerg Dis. 2021;68:1721-5.
314   https://doi.org/10.1111/tbed.13861.

Figure 1

Anti-SARS-CoV-2 RBD IgG titre

Anti-SARS-CoV-2 neutralising titre

- ● A: PHH-1V (10 μg RBD fusion heterodimer/dose)
- ● B: PHH-1V (20 μg RBD fusion heterodimer/dose)
- ● C: PHH-1V (40 μg RBD fusion heterodimer/dose)
- ● D: Control (PBS)

Figure 2

A ELISpot IFN-γ

B Th1 response of CD4+ T cells

C Th1 response of CD8+ T cells

- 1: PHH-1V (40 µg RBD fusion heterodimer/dose)
- 2: Control (PBS)

Figure 3