Long-term methylome changes after experimental seed demethylation and their interaction with recurrent water stress in *Erodium cicutarium* (Geraniaceae)

Francisco Balao[1*], Mónica Medrano[2], Pilar Bazaga[2], Ovidiu Paun[3], Conchita Alonso[2*]

[1] Departamento de Biología Vegetal y Ecología, Universidad de Sevilla, Apdo. 1095, 41080 Sevilla, Spain; [2] Estación Biológica de Doñana, CSIC, Avenida Américo Vespucio 26, 41092 Sevilla, Spain; [3] Department of Botany and Biodiversity Research, University of Vienna, 1030 Vienna, Austria

*Authors for correspondence: Conchita Alonso <conalo@ebd.csic.es >, phone: +34 954466700; Francisco Balao < fbalao@us.es >

Summary

- The frequency and length of drought periods are increasing in subtropical and temperate regions worldwide. Epigenetic responses to water stress could be key for plant resilience to this largely unpredictable challenge. Experimental DNA demethylation together with application of a stress factor stands as a suitable strategy to uncover the contribution of epigenetics to plant responses to stress.

- We analysed leaf cytosine methylation changes in adult plants of the Mediterranean weed, *Erodium cicutarium*, after seed demethylation with 5-Azacytidine and/or recurrent water stress in a greenhouse. We used bisulfite RADseq (BsRADseq) and a newly reported reference genome for *E. cicutarium* to characterize methylation changes in a 2x2 factorial design, controlling for plant relatedness.

- In the long-term, 5-Azacytidine treatment alone caused both hypo and hyper-methylation at individual cytosines, with increased hypomethylation in CG contexts. In control conditions, drought resulted in a decrease in methylation

29 level in all but CHH contexts. In contrast, the genome of plants that experienced

30 recurrent water stress and had been treated with 5-Azacytidine increased DNA

31 methylation level by ca. 5%.

32 • The two study factors exhibited a highly significant interaction between

33 demethylation and recurrent drought stress, with both agents stimulating a

34 general instability of DNA methylation, including around genic regions.

36 **Keywords:** abiotic stress, bisulfite sequencing, BsRADseq, differential methylation,

37 DNA methylation, drought, epigenetics, 5-Azacytidine

Introduction

Climate is changing. The period from 1983 to 2012 was likely the warmest 30-year period of the last 1400 years in the Northern Hemisphere (IPCC, 2014). Strong regional and temporal variability and an increased frequency of extreme events appear linked to global warming, together with a latitudinal redistribution of precipitation leading to increased flooding and drought events on different regions worldwide (Zhang *et al.*, 2007). The Mediterranean Basin is an especially vulnerable region to predicted global changes encompassing increase of summer temperature, reduction of winter precipitation and more frequent heat waves and drought events (Giorgi & Lionello, 2008). Plants in this region are generally used to inconsistent precipitation and to drought (Cowling *et al.*, 2005; Cook *et al.*, 2016) and could, thus, exhibit increased resilience to such stress factor (Matesanz & Valladares, 2014; Balao *et al.*, 2018; López□Rubio *et al.*, 2022). Analyzing the mechanisms behind adaptability to recurrent drought could thus be key to understand potential plant responses to the predicted effects of climate change.

Mounting experimental evidence substantiates that epigenetic mechanisms are involved in plant responses to stress (e.g., Gutzat & Mittelsten Scheid, 2012; Sun *et al.*, 2020). A comprehensive understanding of the interrelationships between chromatin configuration, DNA methylation, histone tail modification and the activity of transposable elements in response to stress is being acquired for model and crop species with well-annotated reference genomes (Mirouze & Paszkowski, 2011; Gutzat & Mittelsten Scheid, 2012; Bäurle, 2018). On the other hand, research on non-model plants has largely focused on changes in cytosine methylation, providing evidence this is a key element of plant epigenomes (Niederhuth & Schmitz, 2017). In plants, methylation can be found at CG, CHG and CHH contexts (where H = A, C or T) because of the action of different families of methyltransferases which can be independently regulated (Lyko, 2018). Experimental studies provide support for a relevant role of DNA methylation in response to several abiotic stressors and particularly to water deficit and salt excess (Peng & Zhang, 2009; Alonso *et al.*, 2016; Banerjee & Roychoudhury, 2017). In addition, experimental modification of DNA methylation profiles by applying inhibitors of DNA methyltransferases (DNMTs) such as Zebularine and 5-Azacytidine (Lopez *et al.*, 2016) has further substantiated the potential epigenetic regulation behind phenotypic responses to stress in non-model

plants (Verhoeven *et al.*, 2010; Herman *et al.*, 2016; Rendina González *et al.*, 2016).

Understanding the stability of epigenomic changes induced by artificial demethylation

at seed stage could be particularly relevant to better evaluate the ability to use epigenetic

information and seed priming for crop improvement (Gallusci *et al.*, 2017; Springer &

Schmitz, 2017).

Moving forward from anonymous markers and indirect evidence of epigenetic

responses to environmental stress in non-model species is timely to better understand

the ecological relevance of epigenetic variation in natural plant populations (Richards *et*

al., 2017). Intrinsic plant (epi)genomic features (Springer *et al.*, 2016), life-history traits

(Verhoeven & Preite, 2014) and the extrinsic factors associated to the environment in

which species have evolved (Balao *et al.*, 2018; Herrera *et al.*, 2019) might condition

the magnitude of epigenetic changes in response to specific stress factors and, thus,

generalizations from results obtained for model and crop species is not straightforward

(Richards *et al.*, 2017). Restriction site-associated DNA sequencing (RADseq hereafter)

has become a popular reduced representation method for non-model organisms.

RADseq uses restriction enzymes to guide complexity reduction to sequencing only a

representative fraction of the genome, and can be implemented with or without prior

genomic resources (Andrews *et al.*, 2016). Similarly, reduced representation bisulfite

sequencing (RRBS) methods provide a cost-effective alternative to Whole Genome

Bisulfite Sequencing (WGBS) for the analysis of DNA methylation differences between

groups of samples in species that lack a well-annotated genome (Paun *et al.*, 2018).

RRBS provide insights into the magnitude and the genomic location of substantial

methylation changes, a relevant step forward because the effects of DNA methylation

on gene expression, transposon activation and specific phenotypic traits depend on both

the sequence context and genomic location (Hirsch *et al.*, 2012; Niederhuth & Schmitz,

2017).

In this paper, we use BsRADseq (Trucchi *et al.*, 2016) to investigate changes in

DNA cytosine methylation in leaves of adult individuals of the annual herb *Erodium*

cicutarium, after experimental seed demethylation and recurrent drought along their

life-cycle. Addition of inhibitors of DNA methyltransferases (DNMTs) at seed

germination has been proofed to be effective in reducing global cytosine methylation in

seedlings of several species including this one (Alonso *et al.*, 2017). However, a

detailed analysis of the genome-wide effects of such demethylating agents is only

available for the model species *Arabidopsis thaliana*, which suggests a significant dose

dependent reduction in all cytosine contexts in 10 d seedlings (Griffin *et al.*, 2016). To

the best of our knowledge no study has applied genomic tools to assess if these early

effects are maintained later in the life cycle, although repeated foliar spraying and foliar

injection of DNMT inhibitors have been proposed as alternative methods to

continuously alter DNA methylation profiles along plant life-cycle (see Puy *et al.*, 2018;

Herrera *et al.*, 2019, respectively). Thus, evaluating the magnitude and specific patterns

of demethylation induced after seed treatment and their stability up to the reproductive

stage, is a critical point for assessing the utility of this method to reveal the associations

between DNA methylation and plant phenotypic variance, and its contribution to stress

response in a broad range of plant species (Richards *et al.*, 2017; Alonso *et al.*, 2019b)

as well as its potential relevance for epigenetic priming of seed crops (Gallusci *et al.*,

2017). In particular, only long-lasting demethylation effects that persist until the adult

reproductive stage, enduring pollen maturation and seed formation, might be relevant

for transgenerational transmission of induced epigenetic variants.

Stress exposure may prime plant individuals and/or their offspring for improved

response after similar or other stress events (Walter *et al.*, 2013; Pandey *et al.*, 2015;

Dangi *et al.*, 2018). However, enduring responses could also compromise plant fitness

under different conditions (Tricker, 2015; Crisp *et al.*, 2016; Latzel *et al.*, 2016; Douma

et al., 2017). In this study we focused on recurrent drought because it is characteristic of

the Mediterranean ecosystems in which *E. cicutarium* lives and also because we were

uncertain of the time lapse for eliciting strain and subsequent recovery after a single

extreme drought event (see e.g., Walter *et al.*, 2011; López-Jurado *et al.*, 2016). Our

specific questions were: do genomes of adult plant leaves exhibit any signature of

DNMT inhibition by 5-Azacytidine at seed stage? which are the genomic methylation

changes associated to recurrent water stress along individual *E. cicutarium* life cycle? is

there any interaction between these two factors?

Materials and Methods

Study species

Erodium cicutarium (L.) L'Hér. (Geraniaceae), is a diploid annual herb native to

Mediterranean Europe, North Africa and western Asia, that is currently distributed

globally in temperate areas with hot summers (Fiz-Palacios *et al.*, 2010; Francis *et al.*,

2012). It has a fast-growing cycle and autonomous self-pollination that make it a good
candidate for experimental studies in greenhouse.

Experimental design

Demethylation and recurrent water stress was applied to plants cultivated in the
greenhouse condition in the second generation. First generation was grown in order to
minimize the effects of heterogeneous growing conditions of maternal wild plants,
which could affect their offspring phenotype and DNA methylation profiles (Latzel,
2015). The parental generation (F_0) were adult plants at two *E. cicutarium* natural
populations located in Cazorla mountains (Jaén province, SE Spain): Puerto del Tejo
(PT, 1,590 m asl) and Nava de las Correhuelas (CH, 1,625 m asl), ca. 10 km distant in
straight-line. Puerto del Tejo site has a steeper slope and rockier substrate; individuals at
this site were larger in diameter but produced less leaves than those at CH site (Alonso
et al., 2017). In spring 2015, we collected mature fruits in these two locations and
stored them in darkness at room temperature. After 4-5 months, seeds were removed
from fruits, scarified, and germinated in universal substrate (COMPO SANA ®) mixed
in 3:1 with perlite (substrate hereafter). The first generation (F_1) seedlings were
subsequently transferred to 1L pots with the same substrate; pots were grouped in trays,
watered twice per week, and trays were periodically rotated within the greenhouse (16h
light; 25-20 °C) until the end of reproduction (ca. 6 months). Autonomously-pollinated
fruits were collected in paper bags, and stored at room temperature.

In October 2016, seeds were weighed and slightly scarified with sandpaper. A
demethylation treatment was applied to half of the seeds as described in Alonso *et al.*
(2017). In brief, scarified seeds were submerged in 150 µl of either Control (water with
DMSO 97:3, v:v) or a 0.5 mM solution of 5-Azacytidine (Sigma A2385-100mg) for 48
h at 4 °C. Immediately after, all seeds were individually transferred to seedling plug
trays filled up with substrate. Trays were saturated in water and placed under
greenhouse conditions (16h light; 25-20 °C). After 39 d of sowing, seedlings from 12
families (2 provenances * 6 F_1-mothers) were transplanted to 1L pots. Extra seedlings
were collected to confirm the effects of the demethylation treatment, which were
detected as a lower global cytosine methylation percentage in leaf DNA and production
of shorter and fewer leaves compared to controls at this developmental stage (Alonso *et*
al., 2017).

Three days later, when seedlings were 6 weeks old, the recurrent water-stress (WS
hereafter) treatment started. For each maternal line, half the offspring was watered at
field capacity twice per week and half the offspring was watered at field capacity once
every 10-11 d until the end of the experiment. Flowering started soon after
transplantation and reached the peak (75 % individuals with flowers) at week 10.
Demethylated plants tended to flower ca. five days later than their untreated relatives
(C. Alonso and M. Medrano unpub. data). The experiment finished when plants were 16
weeks old, and some plants started showing signs of senescence.

Sample processing

Methylation analyses were conducted on DNA extracted from leaves of a subset of
32 reproductive F_2 individuals grown in the greenhouse. At the end of the experiment a
sample of 2-3 full grown leaves without signs of damage or senescence of each F_2
individual was collected, placed in labeled paper bags and dried at ambient temperature
in sealed containers with abundant silica gel.

Dried samples were homogenized to a fine powder using a Retsch MM 200 mill.
Total genomic DNA was extracted using Bioline ISOLATE II Plant DNA Kit and
quantified using a Qubit fluorometer 2.0 (Thermo Fisher Scientific, Waltham, MA,
USA).

RADseq and BsRADseq library preparation

The RADseq and BsRADseq libraries were prepared following the laboratory
protocol detailed in Trucchi *et al*. (2016), with minor changes. In particular, we
modified the restriction enzyme used and employed the more frequent cutter *Pst*I (6 bp
restriction site), in order to maximize the density of markers for differential methylation
analysis. We applied the protocol to analyzing changes in leaf DNA methylation after
the two experimental treatments in 2 provenances * 4 F_1-mothers * 2 demethylation
levels * 2 watering levels = 32 F_2-individuals).

We used 20 units of *Pst*I HF restriction enzyme (New England Biolabs) to digest
800 ng DNA per individual and ligated 100 nM P1 barcoded methylated adapters over
night at 16°C. Groups of samples barcoded with different P1 barcodes were pooled,
sheared to a target peak of 400 bp (with a Covaris E220 focused-ultrasonicator) and

ligated to methylated P2 adapters. We employed a set of methylated P1 and P2

barcoded adapters to protect their sequence from modification during bisulfite

treatment. The fragments were constructed with double barcoding system: eight

different 5 bp barcodes were inserted with the P1 adaptor in combination with four

different P2 adaptor with 6 bp long barcodes. Our barcodes differed in at least four

bases from each other. At this step, one aliquot was separated and directly amplified by

PCR as a standard RADseq library. For the other aliquot, bisulfite conversion of non-

methylated Cs into Us was performed after P2 adapter ligation by treating the library

with MethylEdge Bisulfite Conversion System kit (Promega). PCR amplification was

executed using KAPA HiFi HotStart Uracil+ MasterMix (Kapa) for 23 cycles to convert

all Us into Ts. The final two libraries were sequenced at the VBCF Vienna

(https://www.viennabiocenter.org/vbcf/next-generation-sequencing/) as paired-end 125

bp reads in two lanes of an Illumina HiSeq 2500 machine.

Assembly of Erodium cicutarium draft genome

Erodium cicutarium has a genome size of 1C = 1.20 pg and CG content = 0.4

(Pustahija *et al.*, 2013). A draft genome for *E. cicutarium* was assembled from DNA

extracted from a single individual reared after two events of autonomous selfing under

greenhouse conditions with a paired-end strategy in Illumina HiSeq X (PE150) at

AllGenetics (www.allgenetics.eu). KmerGenie 1.7038 (Chikhi & Medvedev, 2014) was

used to estimate the best kmer in the pair reads and k = 71 was selected as the start point

for de novo assemble with ABySS 2.02 (Simpson *et al.*, 2009). Plastid and

mitochondrion genomes were handled separately. Reads were mapped back to the

assembled output to determine coverage depth using BWA 0.7.12 (Li & Durbin, 2010).

SAMtools 0.1.19 (Li *et al.*, 2009) was used to remove bad quality and secondary

alignments. To avoid possible inconsistencies or the presence of contaminant

sequences, contigs/scaffolds with a mean coverage < 10x were discarded, as well as

those shorter than 1,000 bp.

RepeatModeler 1.0.11 and RepeatMasker 4.0.7 were used to identify and mask the

predicted repeats in the filtered assembly. Geneid 1.4 (Alioto *et al.*, 2018) and SMA3s

(Muñoz-Mérida *et al.*, 2014) were used to perform the prediction of genes and their

subsequent functional annotation. The manually annotated protein database Swiss-Prot

(taxonomic division: plants) available in the UniProt database (The Uniprot

Consortium, 2007) was used as reference. In addition, the predicted genes were also

compared against GenBank's nr protein database including only the Embryophyta taxa

entries by using a BLAST+ 2.6.0 (Camacho *et al.*, 2009).

RADseq and BsRADseq data processing and alignment

Barcoded raw Illumina reads were processed with deML (Renaud *et al.*, 2015) and

STACKS v.2.0Beta8 (Rochette *et al.*, 2019). To demultiplex the individuals and remove

low quality data, the program *process_radtags* was run with the following settings: PstI

as restriction enzyme, restriction site check at the beginning of the reads disabled (due

to BS conversion affecting the sequence of the restriction site), discarding reads with

low quality scores according to default parameters, and rescuing barcodes and RADtags

with two sequencing error (following Trucchi *et al.*, 2016). The paired-end fastq files

obtained were filtered to retain only full-length reads (i.e., P1 120 bp and P2 125 bp

after barcode trimming), with no adaptor contamination, and unambiguous barcode

sites.

The standard RADseq sequences of each of the 32 samples were mapped to the

reference genome (see below) to identify variable sites (i.e., SNPs) involving a cytosine

in the reference sequence. C/T polymorphisms were subsequently filtered and masked

with vcftools for subsequent methylation analyses (see Trucchi *et al.*, 2016 for details).

BsRADseq reads of each of the 32 samples were mapped to the draft reference genome

using the mapping routine available in *Bismark* (Krueger & Andrews, 2011) specifically

designed for dealing with bisulfite converted reads with the core aligner Bowtie2

(Langmead & Salzberg, 2012) in a non-directional modus and allowing up to four

mismatches for a 120 bp read (options: --non_directional -L 32 -D 10 -R 1 --score_min

L,0,-0.2). The sodium bisulfite conversion efficiency rates were assessed by calculating

cytosine methylation levels in the chloroplast genome (Schmidt *et al.*, 2017) and were

found globally satisfactory with conversion ≥ 99.2% in all cases (Table S1). After

mapping, we checked the summary report for each individual, recording the mapping

efficiency, the number of cytosines screened, the distribution among the different

contexts (CG, CHG, and CHH) and the differential representation of original strands

versus complementary to original strands. As expected, the reads mainly mapped (>

99%) complementary to either the top or the bottom strand of the reference genome due

to peculiarities of the bsRADseq protocol (Trucchi *et al.*, 2016) and the stochastic

orientation of the contigs in the reference genome. At supplementary materials, Table

S1 presents a summary of sequencing and mapping results for each individual sample.

The next step was to extract the methylation information of each cytosine position using

the *Bismark_methylation_extractor* routine ignoring the first 4 bp in the reads, including

the "–no_overlap" flag to prevent counting the same cytosine if covered by both the

forward and reverse read and producing cytosine reports (CX_report files in the

Bismark output) for each individual sample with information on all cytosine contexts

and all strands. Output was then merged by sequence context (CG, CHG, CHH) for

downstream analysis.

Global methylation analyses

Differential methylation analyses were conducted in R software 4.2.1 (R Core

Team, 2022). Firstly, the global effects of WS and 5-Azacytidine agent (5-Aza

hereafter) on the genome wide cytosine methylation level were evaluated using a

generalized linear model (GLM) with binomial error and logit link. The dependent

variable was the proportion of methylated Cs to unmethylated Cs in each sample (i.e.,

independent of their genomic position) and the independent ones were WS and 5-Aza

and their interaction effect (5-Aza + WS), as well as the F_1-mother identity (as proxy of

genotype), all considered as fixed factors. The 3-way interaction was not assessed due

to absence of maternal replicates within each factor combination level, still the

experimental design controlled for any potential bias that maternal identity could

introduce by always including one plant from each maternal family in the four factorial

combinations. The analysis was run for the whole dataset and separately for each

sequence context (CG, CHG, CHH).

Differential methylation analyses

Furthermore, to gain insight into the specific effects of 5-Aza and WS on the

methylation patterns with an explicit genomic context, differentially methylated

cytosines (DMCs) at CG, CHG and CHH sites were estimated using the callDML

command in DSS (dispersion shrinkage for sequencing data) package v.2.34.0 (Feng *et*

al., 2014) with a 2 x 2 ANOVA design including the effects of WS, 5-Aza and their

interaction (DMLfit.multiFactor = WS + 5-Aza + WS:5-Aza). In brief, DSS allows

complex experimental designs (including interaction effects), based on a beta-binomial

regression model with arcsine link function, and it takes advantage of using a shrinkage

estimator of the dispersion parameter based on a Bayesian hierarchical model to reduce

the dependence of variance on mean. P-values were adjusted for multiple testing using

Benjamini and Hochberg false discovery rate (FDR) procedure. For each effect (WS, 5-

Aza and their interaction), DMCs with a corrected q-value < 0.05 and > 10% difference

in the methylation level were considered candidate DMCs.

To investigate the methylation changes of the DMCs with significant interaction

effects, we used the optimized K-mean clustering minimizing the sum-of-squares within

cluster (WCSS; Witten & Tibshirani, 2010) on the average methylation of each of the

four levels of treatment combination (Control, 5-Aza, WS, 5-Aza + WS). Methylation

of DMCs and clustered DMCs were visualized using a heatmap created with the R

package *pheatmap* v. 1.0.12 (Kolde, 2019).

Annotation of DMC-associated genes

DMC-associated genes were defined as genes with DMCs within the gene body

region (coding sequence region, CDS, and introns) and/or promoter regions (1 kb

upstream the CDS). In addition, DMCs located within regions annotated as transposable

elements were separately analyzed.

Results

Draft genome assembly and annotation

A total of 225 million pairs of reads have been used in the assembly giving a mean

coverage of 33.8x (SD = 33.5). The assembly of k = 97 was chosen as the best

reconstruction. A total of 65,050 contigs/scaffolds were kept in the final assembly (total

length 628,531,192 bp; N50 = 12,445; L50 = 15,770; GC% = 41.0). The BUSCO 3.0

(Waterhouse *et al.*, 2018) comparison to Embryophyta gene set yielded 75.5 %

complete, 8.5 % fragmented and 15.9 % missing BUSCO orthologs. SMA3s was able to

annotate a total of 43,537 genes with 99.9 % associated to a specific GO term.

Furthermore, a total of 2,135 repeat families were predicted by RepeatModeler and 38.5

% of the genome was masked as repetitive.

BsRADseq library output

We obtained between 2,654,652 and 31,840,142 pairs of reads for the individual

samples. After quality filtering, between 1,133,391 and 11,417,213 paired sequences

per individual were retained (Table S1), corresponding to 35.8% - 42.7% of the raw

reads. On average, the number of pair reads per sample was 4,486,477, the GC content

was 28.9 % (± 1.3), a low figure, as expected for bisulfite treated samples. The mapping

efficiency to our draft reference genome averaged 38.4 %, and ranged from 35 % to

42.5 % across study individuals, resulting in an average coverage of 98.6x +/- 2.1x

(Table S1).

Global DNA methylation changes

The global cytosine methylation level in the genomes of adult *E. cicutarium* leaves, as

estimated by *Bismark*, varied widely across study plants and conditions, ranging from

16.7% to 30.6% (Table 1), and averaged 22.8 % in the genomes of untreated plants (i.e.,

controls). Furthermore, the average cytosine methylation level was different at the three

sequence contexts analyzed, in CG context methylation was estimated as 66.0%,

whereas 30.8% of cytosines were methylated in CHG and only 9.3% in CHH contexts

(Table 1). CG sites were the rarest in the genome, but because of their high mean

methylation level, they represented a major proportion of methylated cytosines,

contributing the most to the overall cytosine methylation that globally averaged 23.2 %

(SD 3.7) in our full set of samples.

The wide range of variance in global methylation level across individuals was

congruent with the significant effect of the F_1-mother factor (as proxy of genotype) on

the proportion of methylated Cs (LR-χ^2= 5421658, P < 0.0001, Fig. S1) and the impact

of the two treatments, whose effects exhibited a significant interaction on this variable

(LR-χ2= 1036036, P < 0.0001. Whereas the 5-Aza seed exposure weakly affected the

Cs methylation in leaves of adult plants grown under optimal watering conditions, a

global and significant increment in methylation level (~5%) was observed in the

genome of leaves of plants that experienced recurrent WS and had been treated with 5-

Aza at seed stage (Fig. 1). Changes in methylation level of Cs at the three sequence

contexts followed similar patterns (Fig. 1). Interestingly, whereas drought stress

appeared to decrease the global methylation level and that at CG and CHG contexts, it

increased DNA methylation at CHH sites that tend to be associated with transposable
elements (e.g., Martin *et al.*, 2021).

Genomic methylation changes induced by 5-Aza and recurrent water stress in leaves of
E. cicutarium

A total of 3.1% of the 222,835 analyzed cytosine positions were identified as
differentially methylated cytosines (DMCs) between control plants and those
experiencing just 5-Aza, just WS or both treatments. An uneven distribution of DMCs
was observed along the genome, the number of DMCs per scaffold differed from a null
distribution (Wilcoxon' test $P < 0.001$, Fig. S2), mainly because the largest fraction of
scaffolds only got one or two DMCs regardless of their size and Cs abundance. Further,
the frequency of DMCs in the three sequence contexts did not resemble their relative
abundance, with the highest proportion being located in CHH (48.67%), followed by
CG (33.31%), and the lowest being in CHG (18.02%).

If we look at the changes induced by the two treatments separately, seed exposure
to 5-Aza had broader impact on the adult leaf methylome in terms of positions modified
(3,897 DMCs) than the WS treatment (1,417 DMCs). This trend was maintained across
the three sequence contexts (Fig. 2a) with 1,323 vs. 501 DMCs in CG context, 710 vs
235 in CHG and 1,864 vs 681 in CHH for 5-Aza and WS treatments, respectively.

As regards the direction of the observed changes, rather unexpectedly, seed
treatment with 5-Aza caused both hypo and hyperDMCs in adult leaves (Fig. 2). The
number of hypoDMCs (in comparison to control) was higher (773) to the number of
hyperDMCs (550) in the CG context, as expected by its predicted negative impact on
the activity of the methylase enzyme MET1, whereas the number of positions altered in
the two directions was similar at both CHG and CHH contexts (Fig. 2a). Furthermore,
WS caused similar number of hypo and hyperDMCs at CG, CHG and CHH contexts
(Fig. 2b-c).

Synergistic, antagonistic and transgressive effects of combined 5-Azacytidine and water
stress on leaf methylome

A total of 4,073 DMCs showed significant interactive effects between 5-Aza and
WS treatments. Most DMCs were located in CHH (48.3%), followed by CG (33.6%),

whereas the lowest proportion was identified in CHG context (18.1%). We clustered
these DMCs by the average methylation level across treatments highlighting groups
with distinct interaction patterns (Fig. 3a). The number of different patterns (i.e. k-
means groups) was eight (which minimized the within group sum of squares, Fig. S3).
The eight patterns could be resumed into two general trends (see dendrogram in Fig.
3a). The first one included DMCs with a very low methylation level in most of the
treatments (Clusters 1, 2, 3 and 4), usually including the control treatment. This "low-
methylation group" represented most of the DMCs analyzed with more than 600 DMC
per cluster. The second group (Clusters 5, 6, 7 and 8) showed the opposite trend
including DMCs with a high methylation level (~100%) in most of the treatments.

Furthermore, the k-mean clustering allowed us to unveil that many of the DMC
were fully methylated or non-methylated in the genomes of control plants identifying
hypo- and hypermethylation for the 5-Aza and WS treatments in comparison to the
control (Fig. 3b, clusters 4 and 8 respectively). In addition, distinctive hyper- and
hypomethylation was detected for 5-Aza + WS treatment (Fig. 3b, clusters 3 and 7,
respectively), singular hypermethylation and hypomethylation for the 5-Aza treatment
(clusters 5 and 1, respectively), increased hypermethylation (following methylation
increment with 5-Azacitidine) for the WS treatment (cluster 2), and the opposite trend
(i.e. increased demethylation) for the WS treatment (cluster 6). From these patterns, we
could discern the synergistic effects of 5-Aza and WS on the methylation (clusters 2, 4,
6 and 8). The antagonist effects of both treatments were also shown in the clusters 1 and
5. Furthermore, transgressive effects for treatments combination were detected (clusters
2, 3, 5, 6 and 7).

Annotation of genomic regions overlapping DMC

We further annotated the genomic regions that were found to be significantly
differentially methylated (Fig. 4). For the 5-Aza treatment, 43% of the CG DMCs, 36%
of the CHG DMCs and 38% of the CHH DMCs overlapped with gene regions (exons,
introns and promoters). These proportions were slightly higher for the WS treatment,
46%, 41% and 41% for CG, CHG and CHH DMCs, respectively. Remarkably, the
highest proportions of DMCs overlapping promoters and introns were found in the
CHH context for both treatments. In addition, on average, 21.3% of DMCs overlapped
transposable elements. Additionally, 36% of the DMCs overlapping TEs appeared

inserted in genic regions (introns and promoters, Fig. S4). The percentage of TEs with
DMCs (from total DMCs) inserted in genic regions increased from CG (4%) and CHG
(6.5%) to CHH context (11.5%). Furthermore, both treatments, 5-Aza and WS, showed
similar proportions of DMCs overlapping TE (without accounting those inserted in
genic regions) for each different context (Fig. 4). The proportions TE-overlapped DMCs
increased from GC context (8% in 5-Aza) to CHH context (19% in 5-Aza). In addition,
although a large amount of TE-overlapped DMC was uncategorized, the proportion of
TE classes significantly differed from the genome distribution ($\chi2 = 149.43$; df = 6; p <
0.001). *Copia* elements containing DMCs (234) showed a higher proportion than
expected (47.5% vs. 25.4%) whereas the LINE elements (61) showed a lower
proportion (12.3% vs. 27.9%; Fig. 5).

Finally, although the disperse nature of the BsRADseq DMCs (Fig. S2) limits
the possibility to identifying genomic regions with multiple DMCs (i.e., Differentially
Methylation Regions, DMR), we obtained 29 potential DMRs covering gene regions
(i.e., overlapping > 10 DMCs; Table 2). Eleven of these potential DMRs (37.9%)
overlapped just intronic regions, whereas 31.0% appeared just on promotor regions and
13.8% just exonic regions. The annotation of these DMR-associated candidate genes
included several with a potential role in stress response (Table 2).

Discussion

The broad interspecific variation in global DNA methylation level, its phylogenetic
signal, and its positive relationship with the haploid genome size (Alonso et al., 2016;
Niederhuth et al., 2016; Alonso et al., 2019), together with the evolution of plant
specific methyltransferases (Bewick et al., 2017; Lyko, 2018; Kenchanmane Raju et al.,
2019) support DNA methylation as a key component of plant epigenomes (Finnegan,
2010). DNA methylation status is regulated by *de novo* DNA methylation, maintenance
of methylation at DNA replication, and active demethylation; these three processes
together with the recruitment of histone modifiers and effectors contribute to genome
stability during development, and in response to genomic and environmental stress
(Law and Jacobsen, 2010; Boyko and Kovalchuk, 2011; Fitz-James and Cavalli, 2022).
The deep molecular understanding of stress response in model plant species with well-
annotated reference genomes contrasts the mostly indirect evidence gained for non-
model plant species based on experimental alteration of DNA methylation and

subsequent analysis of phenotypes, or from the analysis of methylation changes of

anonymous markers, such as MSAP (Richards *et al.*, 2017). Molecular insights gained

by RRBS methods provide a bridge between these two approaches (Paun *et al.*, 2018)

and could inform whether epigenomic regulation in response to stress might act

differently in species with contrasting epigenomic features relative to model organisms

(Springer *et al.*, 2016). In the following paragraphs we will discuss how our results can

help understand the genomic signature of experimental demethylation conducted by

application of 5-Aza at seed stage and its impact on plant methylome responses to

recurrent water stress in an annual plant.

Long-lasting methylation changes after seed demethylation detected by BsRADseq

5-Azacytidine is a structural analogue of 5-methyl-cytosine that can be incorporated

into DNA where it establishes a covalent bond with a methyltransferase enzyme that

can not be easily reversed. The enzyme is trapped reducing the number of active DNA

methyltransferase enzymes in the cells and therefore passively inducing DNA

demethylation (Griffin et al., 2016; Lopez et al., 2016; Lyko, 2018). Applied at seed

stage, 5-Aza reduced global DNA methylation of seedlings in a dose-dependent manner

(Griffin et al., 2016; Alonso et al., 2017). In *A. thaliana* seedlings the effect was more

significant in CG context, where it reduced the frequency of fully methylated loci,

particularly at highly methylated regions, whereas the loss of CHG and CHH

methylation was mainly evident in highly methylated areas of the pericentromeric

regions (Griffin et al., 2016).

Our analysis of changes in cytosine methylation in the DNA of *E. cicutarium* leaves

collected at the onset of flowering is to the best of our knowledge the first attempt to

characterize the long-lasting molecular signatures of seed demethylation treatment with

5-Aza. The abridged view of the methylome provided by BsRADseq analysis indicated

a mild overall reduction in global cytosine methylation that was most evident in CHG

contexts (Fig. 1). More interestingly, both hyper and hypoDMC were obtained in the

leaf genomes of flowering plants treated with 5-Aza at seed stage in comparison to

those grown from untreated seeds. In CG context the number of hypoDMCs was

slightly higher, whereas the number of positions altered in either direction was similar

at both CHG and CHH contexts. These findings altogether suggest that 5-Aza at seed

stage significantly altered the patterning of cytosine methylation in genomes of adult

plant, and might at least partially erase memories of past stress (see e.g., Akkerman *et*
al., 2016). Seed demethylation is, thus, a suitable tool for analyzing the impact of
deregulated epigenetic backgrounds on molecular and phenotypic response to stress
analyzed either in the treated individuals or their progenies (Burton & Metcalfe, 2014).
It is also important to highlight here the significant family effect detected by the
analyses of global methylation level which reveals that experimental demethylation
does not necessarily impact equally all treated individuals, an aspect that has been
already reported in replicated ecological studies analyzing phenotypic effects of seed
demethylation with different genetic lines or provenances (Herman *et al.*, 2016; Troyee
et al., 2022) in contrast to a single accession or genetic line (e.g., Akkerman *et al.*,
2016). Thus, a homogeneous impact of the treatment should not be taken as granted and
a screening of methylation changes across different genotypes, populations and species
would be highly desirable to assess the actual impact of the 5-Aza seed treatment.

Genomic methylation changes in response to recurrent water stress across the
individual life cycle

Epigenetic response to abiotic stress can involve complex regulation of histone
marks, small RNAs production and DNA methylation (Banerjee & Roychoudhury,
2017), that impinge transcription of metabolic pathways and, in some cases, can be
retained as memory marks of past events to improve individual (Fleta-Soriano &
Munné-Bosch, 2016) or transgenerational responses (Fitz-James & Cavalli, 2022). In
particular, significant DNA methylation changes in response to moderate to heavy water
stress treatments have been reported for several plant species (see e.g., Pardo *et al.*,
2020 for several grasses; Sow *et al.*, 2021 for poplar), including variable responses of
crop varieties expected to have contrasting drought tolerance in *Vicia faba* (Abid *et al.*,
2017) and *Oryza sativa* (Wang *et al.*, 2016), supporting its relevance as a mechanism
for both long-term wild plant adaptation and crop improvement. Mild water stress
applied at early seedling stage has been found to alter DNA methylation in the leaves of
A. thaliana Col-0 accession, mainly at CHH contexts and predominantly within TE
sequences, and significantly up- and down-regulate transcription of multiple genes (Van
Dooren *et al.*, 2020). Some stress-induced epigenetic variants are reversible and
recurrent stress is expected to promote molecular memory and increased tolerance in
subsequent events (Wang *et al.*, 2021). Our recurrent water stress, aimed to mimic
unpredictable rain in Mediterranean ecosystems, led to a significant overall increased

DNA cytosine methylation in all three contexts in genomes of *E. cicutarium* leaves at

flowering onset. Our analysis of DMCs indicated that the genomes of plants

experiencing WS were either hypo or hyper-methylated compared to those grown well-

watered. A large proportion of DMCs in all contexts overlapped in our study intergenic

regions (Fig. 4), an aspect in part potentially related to the reduced resolution of the

annotation of the newly reported reference genome. Apart from that, methylation

changes at CG contexts mainly overlapped exons, and those in CHH were found to be

comparatively more frequently associated to TEs and introns.

Combined effects of 5-Azacytidine and water stress on leaf methylome

Significant interactions between DNA demethylating agents and experimental

stress treatments have been proposed as a reliable approach to investigate the role of

DNA methylation in regulating plant responses to stress in non-model plant species

(Puy *et al.*, 2018; Alonso *et al.*, 2019a). Phenotypic evidence of significant interaction

effects in response to water stress has been gained in a few non-model plant species

(Herman *et al.*, 2016; Rendina González *et al.*, 2016). Our study confirms a significant

interaction at molecular level in the leaf genomes of *E. cicutarium* adult plants between

5-Aza-induced demethylation and water stress. This interaction appears to produce a

complex epigenomic instability making unpredictable the consequences of experimental

re-methylation and water stress on the genome-wide scale. On one hand, 5-Aza

application in seeds triggers widespread changes in DNA methylation in adult leaves,

but importantly it likely alters the expression of transcriptional and post-transcriptional

regulatory proteins as suggested by the DMRs overlapping the orthologous of

Methyltransferase 13, *RNA-dependent DNA polymerase* and *60S ribosomal protein*

(Litholdo & Bousquet-Antonelli, 2019; Sáez-Vásquez & Delseny, 2019). Interestingly,

5-Aza promoted the hypermethylation of the Ec-S10188_3 gene, which contained 28

differentially methylated cytosines in the promoter region (Table 2) and encodes a

putative disease resistance protein, with high identity to Resistance genes analogs 3

(RGA3) in *Theobroma cacao* (Sekhwal *et al.*, 2015). Likewise, the promoter of Ec-

S44030_4 gene, a *Cyclin-dependent protein kinase (CDKC)* orthologous, related to

stress tolerance mechanisms (Kitsios & Doonan, 2011), was also hypermethylated.

Although, an antagonist effect of water stress and 5-Aza exposure was found for these

genes (Cluster 1), the 5-Aza-induced hypermethylation may resemble a priming state

for a faster future responses to disease attacks and abiotic stress (Latzel *et al.*, 2016;

Alonso *et al.*, 2018).

On the other hand, environmental stresses, such as cold and drought, can also

affect the methylase and/or demethylase activity (Lucibelli *et al.*, 2022) explaining the

interaction with the demethylation agent 5-Azacytidine. For example, heat stress

induced downregulation of demethylases in *Triticum aestivum* (Gahlaut *et al.*, 2022) but

were significantly induced in *Camellia sinensis* under cold and drought (Zhu *et al.*,

2020). Demethylases were also significantly induced by cold in *Dendrobium officinale*

where methylases were also affected by drought (Yu *et al.*, 2021). In our study, water

stress, as well as the 5-Aza treatment, independently triggered the promoter

hypomethylation of the *Methyltransferase 13* which may activate or upregulate its

expression. However, the combination of treatments restored the control methylation

level of this cytosine methylase. In addition, the methylation stage of many abiotic

response-genes was directly altered by the water stress or its combination with the 5-

Aza treatment. For example, the promoter regions of the abiotic stress related genes

Phenylalanine ammonia-lyase (PAL, Jeong *et al.*, 2012) and *ABC transporter F family*

(ABCF, Dahuja *et al.*, 2021) were also affected. Most methylation changes were

observed at CHH context potentially linked to a genome-wide decrease in TE activity

repression, as previously reported for other stress studies (e.g., Van Dooren *et al.*,

2020).

Changes in methylation in Transposable Elements

In plants, various TEs are involved in environmental stress adaptation (Casacuberta &

González, 2013; Sahu *et al.*, 2013). We found a significant differential methylation of

Copia transposable elements in *Erodium cicutarium* under 5-Aza and WS treatments.

Copia superfamily of long terminal repeat (LTR) retrotransposons is known to be

activated by biotic and abiotic stresses (Ito, 2022). ONSEN elements are highly

expressed under heat stress in Brassicaceae (Pietzenuk *et al.*, 2016), whereas other

Copia elements are activated by several environmental stresses in different species (see

Table 1 in Ito, 2022). Transposons activation can affect gene regulation on a genome-

wide scale by acting as new cis-regulatory elements and conferring increased stress-

responsiveness to nearby protein-coding genes, which can facilitate acclimation or even

adaptation to stressful conditions (Horváth *et al.*, 2017; Dubin *et al.*, 2018).Transposon

activation under environmental stress can have negative effects on fitness (Horváth *et*
al., 2017), but increasing TE methylation, especially in genomic regions neighbouring
highly expressed stress-induced genes, could prevent TE transcription via RNA
polymerase II and hence avoid deleterious effects upon the plant (Secco *et al.*, 2015).

Conclusions

By using a 2x2 factorial design, combining recurrent drought stress and early life
demethylation with a reduced representation method of surveying genome-wide
methylation levels in adult plants, we show a significant interaction between
methylation and plant stress response in the Mediterranean *Erodium cicutarium*. We
uncover extensive effects of both seedling demethylation and drought stress on adult
DNA methylation, with a large portion of changes observed around genic regions and
within TEs. Altogether, our results provide useful information towards understanding
the role of DNA methylation in plant drought responses.

Acknowledgements

We thank Laura Cabral, Esmeralda López, Daisy Johnson, Pablo Martín, Alejandro
Mira, Noelia Zarza, and Elena Villa for assistance in greenhouse and laboratory; Maite
Lorenzo for her training with BsRADseq wet lab; Rocío Esteban and all the staff of
Allgenetics for their assistance with draft genome analyses; and Rubén Martín-Blázquez
for suggestions on a draft version of the manuscript. We are particularly indebted to
Carlos M. Herrera for his continuous encouragement and insightful discussions at all
phases of this work. Financial support was provided by grants EPIECOL-CGL2013-
43352-P, EPIENDEM-CGL2016-76605-P and EPINTER-PID2019-104365GB-I00
(Ministerio de Ciencia e Innovación, Spanish Government).

Author contributions

C.A., F.B. and M.M. designed the research; C.A. and M.M. conducted greenhouse
experiments; C.A., F.B. and O.P. designed the BsRADseq experiment and contributed
to data analyses; P.B. prepared the libraries; F.B. led data analyses; C.A. led the writing;
all authors contributed to refining the manuscript.

References

Abid G, Mingeot D, Muhovski Y, Mergeai G, Aouida M, Abdelkarim S, Aroua I, El Ayed M, M'hamdi M, Sassi K, *et al.* **2017**. Analysis of DNA methylation patterns associated with drought stress response in faba bean (*Vicia faba* L.) using methylation-sensitive amplification polymorphism (MSAP). *Environmental and Experimental Botany* **142**: 34–44.

Akkerman KC, Sattarin A, Kelly JK, Scoville AG. 2016. Transgenerational plasticity is sex-dependent and persistent in yellow monkeyflower (*Mimulus guttatus*). *Environmental Epigenetics* **2**: 1–8.

Alioto T, Blanco E, Parra G, Guigó R. 2018. Using geneid to identify genes. *Current Protocols in Bioinformatics* **64**: 1–32.

Alonso C, Medrano M, Pérez R, Bazaga P, Herrera CM. 2017. Tissue-specific response to experimental demethylation at seed germination in the non-model herb *Erodium cicutarium*. *Epigenomes* **1**: 1–11.

Alonso C, Medrano M, Pérez R, Canto A, Parra-Tabla V, Herrera CM. 2019a. Interspecific variation across angiosperms in global DNA methylation: phylogeny, ecology and plant features in tropical and Mediterranean communities. *New Phytologist* **224**: 949–960.

Alonso C, Pérez R, Bazaga P, Medrano M, Herrera CM. 2016. MSAP markers and global cytosine methylation in plants: A literature survey and comparative analysis for a wild-growing species. *Molecular Ecology Resources* **16**: 80–90.

Alonso C, Ramos-Cruz D, Becker C. 2018. The role of plant epigenetics in biotic interactions. *New Phytologist*.

Alonso C, Ramos-Cruz D, Becker C. 2019b. The role of plant epigenetics in biotic interactions. *New Phytologist* **221**: 731–737.

Andrews KR, Good JM, Miller MR, Luikart G, Hohenlohe PA. 2016. Harnessing the power of RADseq for ecological and evolutionary genomics. *Nature Reviews Genetics* **17**: 81–92.

Balao F, Paun O, Alonso C. 2018. Uncovering the contribution of epigenetics to plant phenotypic variation in Mediterranean ecosystems. *Plant Biology* **20**: 38–49.

Banerjee A, Roychoudhury A. 2017. Epigenetic regulation during salinity and drought stress in plants: Histone modifications and DNA methylation. *Plant Gene* **11**: 199–204.

Bäurle I. 2018. Can't remember to forget you: Chromatin-based priming of somatic

stress responses. *Seminars in Cell and Developmental Biology* **83**: 133–139.

Burton T, Metcalfe NB. **2014**. Can environmental conditions experienced in early life

influence future generations? *Proceedings of the Royal Society B: Biological Sciences*

281.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden

TL. **2009**. BLAST+: Architecture and applications. *BMC Bioinformatics* **10**: 1–9.

Casacuberta E, González J. **2013**. The impact of transposable elements in

environmental adaptation. *Molecular Ecology* **22**: 1503–1517.

Chikhi R, Medvedev P. **2014**. Informed and automated k-mer size selection for

genome assembly. *Bioinformatics* **30**: 31–37.

Cook BI, Anchukaitis KJ, Touchan R, Meko DM, Cook ER. **2016**. Spatiotemporal

drought variability in the mediterranean over the last 900 years. *Journal of Geophysical*

Research **121**: 2060–2074.

Cowling RM, Ojeda F, Lamont BB, Rundel PW, Lechmere-Oertel R. **2005**. Rainfall

reliability, a neglected factor in explaining convergence and divergence of plant traits in

fire-prone mediterranean-climate ecosystems. *Global Ecology and Biogeography* **14**:

509–519.

Crisp PA, Ganguly D, Eichten SR, Borevitz JO, Pogson BJ. **2016**. Reconsidering

plant memory: Intersections between stress recovery, RNA turnover, and epigenetics.

Science Advances **2**.

Dahuja A, Kumar RR, Sakhare A, Watts A, Singh B, Goswami S, Sachdev A,

Praveen S. **2021**. Role of ATP-binding cassette transporters in maintaining plant

homeostasis under abiotic and biotic stresses. *Physiologia Plantarum* **171**: 785–801.

Dangi AK, Sharma B, Khangwal I, Shukla P. **2018**. Combinatorial interactions of

biotic and abiotic stresses in plants and their molecular mechanisms: systems biology

approach. *Molecular Biotechnology* **60**: 636–650.

Van Dooren TJM, Silveira AB, Gilbault E, Jiménez-Gómez JM, Martin A, Bach L,

Tisné S, Quadrana L, Loudet O, Colot V. **2020**. Mild drought in the vegetative stage

induces phenotypic, gene expression, and DNA methylation plasticity in *Arabidopsis*

but no transgenerational effects. *Journal of Experimental Botany* **71**: 3588–3602.

Douma JC, Vermeulen PJ, Poelman EH, Dicke M, Anten NPR. **2017**. When does it

pay off to prime for defense? A modeling analysis. *New Phytologist* **216**: 782–797.

Dubin MJ, Mittelsten Scheid O, Becker C. **2018**. Transposons: a blessing curse.

Current Opinion in Plant Biology **42**: 23–29.

Feng H, Conneely KN, Wu H. **2014**. A Bayesian hierarchical model to detect

differentially methylated loci from single nucleotide resolution sequencing data. *Nucleic*

Acids Research **42**: 1–11.

Fitz-James MH, Cavalli G. **2022**. Molecular mechanisms of transgenerational

epigenetic inheritance. *Nature Reviews Genetics* **23**: 325–341.

Fiz-Palacios O, Vargas P, Vila R, Papadopulos AST, Aldasoro JJ. **2010**. The uneven

phylogeny and biogeography of *Erodium* (Geraniaceae): Radiations in the

Mediterranean and recent recurrent intercontinental colonization. *Annals of Botany* **106**:

871–884.

Fleta-Soriano E, Munné-Bosch S. **2016**. Stress memory and the inevitable effects of

drought: A physiological perspective. *Frontiers in Plant Science* **7**: 143.

Francis A, Darbyshire SJ, Légère A, Simard MJ. **2012**. The biology of Canadian

weeds. 151. *Erodium cicutarium* (L.) L'Hér. ex Aiton. *Canadian Journal of Plant*

Science **92**: 1359–1380.

Gahlaut V, Samtani H, Gautam T, Khurana P. **2022**. Identification and

characterization of DNA demethylase genes and their association with thermal stress in

Wheat (*Triticum aestivum* L .). *Frontiers in Genetics* **13**: 894020.

Gallusci P, Dai Z, Génard M, Gauffretau A, Leblanc-Fournier N, Richard-Molard

C, Vile D, Brunel-Muguet S. **2017**. Epigenetics for plant improvement: current

knowledge and modeling avenues. *Trends in Plant Science* **22**: 610–623.

Giorgi F, Lionello P. **2008**. Climate change projections for the Mediterranean region.

Global and Planetary Change **63**: 90–104.

Griffin PT, Niederhuth CE, Schmitz RJ. **2016**. A comparative analysis of 5-

azacytidine-and zebularine-induced DNA demethylation. *G3: Genes, Genomes,*

Genetics **6**: 2773–2780.

Gutzat R, Mittelsten Scheid O. **2012**. Epigenetic responses to stress: Triple defense?

Current Opinion in Plant Biology **15**: 568–573.

Herman JJ, Sultan SE, Herman JJ. **2016**. DNA methylation mediates genetic

variation for adaptive transgenerational plasticity. *Proceedings of the Royal Society B:*

Biological Sciences **283**: 20160988.

Herrera CM, Medrano M, Pérez R, Bazaga P, Alonso C. **2019**. Within-plant

heterogeneity in fecundity and herbivory induced by localized DNA hypomethylation in

the perennial herb *Helleborus foetidus*. *American Journal of Botany* **106**: 798–806.

Hirsch S, Baumberger R, Grossniklaus U. **2012**. Epigenetic variation, inheritance,

and selection in plant populations. *Cold Spring Harbor Symposia on Quantitative*

Biology **77**: 97–104.

Horváth V, Merenciano M, González J. **2017**. Revisiting the Relationship between

Transposable Elements and the Eukaryotic Stress Response. *Trends in Genetics* **33**:

832–841.

IPCC. **2014**. *Climate Change 2014: Synthesis Report. Contribution of Working Groups*

I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate

Change. Geneva, Switzerland.

Ito H. **2022**. Environmental stress and transposons in plants. *Genes & Genetic Systems*:

22–00045.

Jeong MJ, Choi BS, Bae DW, Shin SC, Park SU, Lim HS, Kim J, Kim JB, Cho

BK, Bae H. **2012**. Differential expression of kenaf phenylalanine ammonia-lyase (PAL)

ortholog during developmental stages and in response to abiotic stresses. *Plant OMICS*

5: 392–399.

Kitsios G, Doonan JH. **2011**. Cyclin dependent protein kinases and stress responses in

plants. *Plant Signaling and Behavior* **6**: 204–209.

Kolde R. **2019**. pheatmap: Pretty Heatmapsitle. : R package version 1.0.12.

Krueger F, Andrews SR. **2011**. Bismark: A flexible aligner and methylation caller for

Bisulfite-Seq applications. *Bioinformatics* **27**: 1571–1572.

Langmead B, Salzberg SL. **2012**. Fast gapped-read alignment with Bowtie 2. *Nature*

methods **9**: 357–9.

Latzel V. **2015**. Pitfalls in ecological research – transgenerational effects. *Folia*

Geobotanica **50**: 75–85.

Latzel V, Rendina González AP, Rosenthal J. **2016**. Epigenetic memory as a basis for

intelligent behavior in clonal plants. *Frontiers in Plant Science* **7**: 1–7.

Li H, Durbin R. **2010**. Fast and accurate long-read alignment with Burrows-Wheeler

transform. *Bioinformatics* **26**: 589–595.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis

G, Durbin R. **2009**. The Sequence Alignment/Map format and SAMtools.

Bioinformatics **25**: 2078–2079.

Litholdo CG, Bousquet-Antonelli C. **2019**. Chemical RNA modifications: The plant

epitranscriptome. In: Epigenetics in Plants of Agronomic Importance: Fundamentals

and Applications. Springer, 291–310.

López-Jurado J, Balao F, Mateos-Naranjo E. **2016**. Deciphering the ecophysiological

traits involved during water stress acclimation and recovery of the threatened wild

carnation, Dianthus inoxianus. *Plant Physiology and Biochemistry* **109**: 397–405.

López Rubio R, Pescador DS, Escudero A, Sánchez AM. **2022**. Rainy years

counteract negative effects of drought on taxonomic, functional, and phylogenetic

diversity: resilience in annual plant communities. *Journal of Ecology*: 1–13.

Lopez M, Halby L, Arimondo PB. **2016**. *DNA methyltransferase inhibitors:*

Development and applications.

Lucibelli F, Valoroso MC, Aceto S. **2022**. Plant DNA methylation : an epigenetic

mark in development , environmental interactions , and evolution. *International Journal*

of Molecular Sciences **23**: 8299.

Lyko F. **2018**. The DNA methyltransferase family: A versatile toolkit for epigenetic

regulation. *Nature Reviews Genetics* **19**: 81–92.

Martin GT, Seymour DK, Gaut BS. **2021**. CHH methylation islands: a nonconserved

feature of grass genomes that is positively associated with transposable elements but

negatively associated with gene-body methylation. *Genome Biology and Evolution* **13**:

1–16.

Matesanz S, Valladares F. **2014**. Ecological and evolutionary responses of

Mediterranean plants to global change. *Environmental and Experimental Botany* **103**:

53–67.

Mirouze M, Paszkowski J. **2011**. Epigenetic contribution to stress adaptation in plants.

Current Opinion in Plant Biology **14**: 267–274.

Muñoz-Mérida A, Viguera E, Claros MG, Trelles O, Pérez-Pulido AJ. **2014**.

Sma3s: A three-step modular annotator for large sequence datasets. *DNA Research* **21**:

341–353.

Niederhuth CE, Schmitz RJ. **2017**. Putting DNA methylation in context: from

genomes to gene expression in plants. *Biochimica et Biophysica Acta - Gene Regulatory*

Mechanisms **1860**: 149–156.

Pandey P, Ramegowda V, Senthil-Kumar M. **2015**. Shared and unique responses of

plants to multiple individual stresses and stress combinations: Physiological and

molecular mechanisms. *Frontiers in Plant Science* **6**: 1–14.

Pardo J, Wai CM, Chay H, Madden CF, Hilhorst HWM, Farrant JM, VanBuren

R. **2020**. Intertwined signatures of desiccation and drought tolerance in grasses.

Proceedings of the National Academy of Sciences of the United States of America **117**:

10079–10088.

Paun O, Verhoeven KJF, Richards CL. **2018**. Opportunities and limitations of
reduced representation bisulfite sequencing in plant ecological epigenomics. *New*
Phytologist.

Peng H, Zhang J. **2009**. Plant genomic DNA methylation in response to stresses:
Potential applications and challenges in plant breeding. *Progress in Natural Science* **19**:
1037–1045.

Pietzenuk B, Markus C, Gaubert H, Bagwan N, Merotto A, Bucher E, Pecinka A.
2016. Recurrent evolution of heat-responsiveness in Brassicaceae COPIA elements.
Genome Biology **17**: 1–15.

Pustahija F, Brown SC, Bogunić F, Bašić N, Muratović E, Ollier S, Hidalgo O,
Bourge M, Stevanović V, Siljak-Yakovlev S. **2013**. Small genomes dominate in plants
growing on serpentine soils in West Balkans, an exhaustive study of 8 habitats covering
308 taxa. *Plant and Soil* **373**: 427–453.

Puy J, Dvořáková H, Carmona CP, de Bello F, Hiiesalu I, Latzel V. **2018**. Improved
demethylation in ecological epigenetic experiments: Testing a simple and harmless
foliar demethylation application. *Methods in Ecology and Evolution* **9**: 744–753.

R Core Team. **2022**. A language and environment for statistical computing.

Renaud G, Stenzel U, Maricic T, Wiebe V, Kelso J. **2015**. DeML: Robust
demultiplexing of Illumina sequences using a likelihood-based approach.
Bioinformatics **31**: 770–772.

Rendina González AP, Chrtek J, Dobrev PI, Dumalasová V, Fehrer J, Mráz P,
Latzel V. **2016**. Stress-induced memory alters growth of clonal offspring of white
clover (*Trifolium repens*). *American Journal of Botany* **103**: 1567–1574.

Richards CL, Alonso C, Becker C, Bossdorf O, Bucher E, Colomé-Tatché M,
Durka W, Engelhardt J, Gaspar B, Gogol-Döring A, *et al.* **2017**. Ecological plant
epigenetics: Evidence from model and non-model species, and the way forward.
Ecology Letters **20**: 1576–1590.

Rochette NC, Rivera-Colón AG, Catchen JM. **2019**. Stacks 2: Analytical methods for
paired-end sequencing improve RADseq-based population genomics. *Molecular*
Ecology **28**: 4737–4754.

Sáez-Vásquez J, Delseny M. **2019**. Ribosome biogenesis in plants: From functional
45S ribosomal DNA organization to ribosome assembly factors. *Plant Cell* **31**: 1945–
1967.

Sahu PP, Pandey G, Sharma N, Puranik S, Muthamilarasan M, Prasad M. **2013**.

Epigenetic mechanisms of plant stress responses and adaptation. *Plant Cell Reports* **32**:

1151–1159.

Schmidt M, Van Bel M, Woloszynska M, Slabbinck B, Martens C, De Block M,

Coppens F, Van Lijsebettens M. **2017**. Plant-RRBS, a bisulfite and next-generation

sequencing-based methylome profiling method enriching for coverage of cytosine

positions. *BMC Plant Biology* **17**: 1–14.

Secco D, Wang C, Shou H, Schultz MD, Chiarenza S, Nussaume L, Ecker JR,

Whelan J, Lister R. **2015**. Stress induced gene expression drives transient DNA

methylation changes at adjacent repetitive elements. *eLife* **4**: 1–26.

Sekhwal MK, Li P, Lam I, Wang X, Cloutier S, You FM. **2015**. Disease resistance

gene analogs (RGAs) in plants. *International Journal of Molecular Sciences* **16**: 19248–

19290.

Simpson JT, Wong K, Jackman SD, Schein JE, Jones SJM, Birol I. **2009**. ABySS:

A parallel assembler for short read sequence data. *Genome Research* **19**: 1117–1123.

Sow MD, Le Gac AL, Fichot R, Lanciano S, Delaunay A, Le Jan I, Lesage-

Descauses MC, Citerne S, Caius J, Brunaud V, *et al.* 2021. RNAi suppression of

DNA methylation affects the drought stress response and genome integrity in transgenic

poplar. *New Phytologist* **232**: 80–97.

Springer NM, Lisch D, Li Q. **2016**. Creating order from chaos: epigenome dynamics

in plants with complex genomes. *The Plant Cell* **28**: 314–325.

Springer NM, Schmitz RJ. **2017**. Exploiting induced and natural epigenetic variation

for crop improvement. *Nature Publishing Group*.

Sun L, Jing Y, Liu X, Li Q, Xue Z, Cheng Z, Wang D, He H, Qian W. **2020**. Heat

stress-induced transposon activation correlates with 3D chromatin organization

rearrangement in Arabidopsis. *Nature Communications* **11**.

The Uniprot Consortium. **2007**. The Universal Protein Resource (UniProt). *Nucleic*

Acids Research **35**: 193–197.

Tricker PJ. **2015**. Transgenerational inheritance or resetting of stress-induced

epigenetic modifications: Two sides of the same coin. *Frontiers in Plant Science* **6**: 1–6.

Troyee AN, Medrano M, Müller C, Alonso C. **2022**. Variation in DNA Methylation

and Response to Short-Term Herbivory in *Thlaspi Arvense*. *Flora* **293**: 152106.

Trucchi E, Mazzarella AB, Gilfillan GD, Lorenzo MT, Schönswetter P, Paun O.

2016. BsRADseq: screening DNA methylation in natural populations of non-model

species. *Molecular Ecology* **25**: 1697–1713.

Verhoeven KJF, Jansen JJ, van Dijk PJ, Biere A. **2010**. Stress-induced DNA

methylation changes and their heritability in asexual dandelions. *New Phytologist* **185**:

1108–1118.

Verhoeven KJF, Preite V. **2014**. Epigenetic variation in asexually reproducing

organisms. *Evolution* **68**: 644–655.

Walter J, Jentsch A, Beierkuhnlein C, Kreyling J. **2013**. Ecological stress memory

and cross stress tolerance in plants in the face of climate extremes. *Environmental and*

Experimental Botany **94**: 3–8.

Walter J, Nagy L, Hein R, Rascher U, Beierkuhnlein C, Willner E, Jentsch A.

2011. Do plants remember drought? Hints towards a drought-memory in grasses.

Environmental and Experimental Botany **71**: 34–40.

Wang L, Cao S, Wang P, Lu K, Song Q, Zhao FJ, Chen ZJ. **2021**. DNA

hypomethylation in tetraploid rice potentiates stress-responsive gene expression for salt

tolerance. *Proceedings of the National Academy of Sciences of the United States of*

America **118**: 1–10.

Wang W, Qin Q, Sun F, Wang Y, Xu D, Li Z, Fu B. **2016**. Genome-wide differences

in DNA methylation changes in two contrasting rice genotypes in response to drought

conditions. *Frontiers in Plant Science* **7**: 1–13.

Waterhouse RM, Seppey M, Simao FA, Manni M, Ioannidis P, Klioutchnikov G,

Kriventseva E V., Zdobnov EM. **2018**. BUSCO applications from quality assessments

to gene prediction and phylogenomics. *Molecular Biology and Evolution* **35**: 543–548.

Witten DM, Tibshirani R. **2010**. A framework for feature selection in clustering.

Journal of the American Statistical Association **105**: 713–726.

Yu Z, Zhang G, Teixeira da Silva JA, Li M, Zhao C, He C, Si C, Zhang M, Duan J.

2021. Genome-wide identification and analysis of DNA methyltransferase and

demethylase gene families in *Dendrobium officinale* reveal their potential functions in

polysaccharide accumulation. *BMC Plant Biology* **21**: 1–17.

Zhang X, Zwiers FW, Hegerl GC, Lambert FH, Gillett NP, Solomon S, Stott PA,

Nozawa T. **2007**. Detection of human influence on twentieth-century precipitation

trends. *Nature* **448**: 461–465.

Zhu C, Zhang S, Zhou C, Chen L, Fu H, Li X, Lin Y, Lai Z, Guo Y. **2020**. Genome-

wide investigation and transcriptional analysis of cytosine-5 DNA methyltransferase

and DNA demethylase gene families in tea plant (*Camellia sinensis*) under abiotic stress

and withering processing. *PeerJ* **2020**.

Table 1. Genome-wide methylation levels as total and for each sequence context for the 32
samples. C, control; A, adult plants obtained after 5-Aza treatment of seeds; WW, optimal
watering; WS, drought stress.

Treatments		Family	Sample	No.	mC	mCG	mCHG	mCHH
5-Aza	Watering			cytosines	(%)	(%)	(%)	(%)
C	WW	PT_1031	CC_253	92353730	18.4	58.6	22.3	6.3
C	WW	PT_1001	CC_225	108547644	23.3	67.7	32.8	8.4
C	WW	PT_0969	CC_147	47011293	18.7	60.2	21.9	7.4
C	WW	PT_0925	CC_001	124533805	24.5	67.5	32.9	10.5
C	WW	CH_1307	CC_491	39099076	25.5	68.8	34.1	11.4
C	WW	CH_1245	CC_433	22052707	23.2	66.2	31.9	9.8
C	WW	CH_1233	CC_409	94508383	17.1	58.5	18.4	6.6
C	WW	CH_1187	CC_291	85841474	25.9	69.5	35.8	10.6
C	WS	PT_1031	CD_259	128071349	17.8	58.6	20.4	6.4
C	WS	PT_1001	CD_219	116121937	27.8	69.2	43.5	9.8
C	WS	PT_0969	CD_163	128356306	23.0	65.4	30.3	9.5
C	WS	PT_0925	CD_005	213718116	21.8	63.1	25.8	9.7
C	WS	CH_1307	CD_485	22299814	23.7	69.4	29.0	10.2
C	WS	CH_1245	CD_445	29859487	21.4	63.3	27.3	8.7
C	WS	CH_1233	CD_431	87893575	16.7	58.3	17.5	5.9
C	WS	CH_1187	CD_297	50225708	25.8	71.8	35.8	10.0
A	WW	PT_1031	AC_252	164753131	24.9	68.8	34.5	9.3
A	WW	PT_1001	AC_220	97316260	18.9	57.8	26	6.4
A	WW	PT_0969	AC_146	46070213	19.0	61.2	22.9	6.6
A	WW	PT_0925	AC_016	75973105	27.6	70.4	37.6	11.9
A	WW	CH_1307	AC_494	44354076	24.4	69.7	31.6	10.3
A	WW	CH_1245	AC_438	21496364	25.7	69.6	35.6	10.7
A	WW	CH_1233	AC_422	99564473	21.2	63.0	26.3	8.9
A	WW	CH_1187	AC_300	54846691	18.1	59.4	21.2	6.8
A	WS	PT_1031	AD_254	137114375	25.5	67.2	37.4	9.4
A	WS	PT_1001	AD_222	38996631	25.7	67.9	37.9	8.8
A	WS	PT_0969	AD_164	85184218	24.5	68.9	33.3	9.3
A	WS	PT_0925	AD_020	105676423	28.8	72.6	40	13.2
A	WS	CH_1307	AD_482	38691859	22.1	65.7	28.0	9.1
A	WS	CH_1245	AD_436	34188955	28.1	73.4	39.1	12.3

| A | WS | CH_1233 | AD_416 | 82021987 | 23.6 | 64.9 | 30.1 | 10.2 |
| A | WS | CH_1187 | AD_292 | 74545405 | 30.6 | 76.0 | 44.0 | 13.0 |

879   **Table 2**. Annotation of gene regions overlapping DMRs (i.e., >= 10 DMCs). We showed the

880   maximum number of DMCs, their genomic location, and the most frequent treatment (either 5-

881   Aza or WS) or interaction cluster (denoted by C1-C8 as in Fig. 3) associated to DMCs obtained.

882   See Table S2 for details.

Gene	Annotation	DMCs			Treatment/ Cluster
		Promoter	Exon	Intron	
Ec-S10188_3	CC-NBS-LRR resistance protein RGA3	28	0	0	C1
Ec-S25751_1	Methyltransferase-like protein 13	19	0	0	C5
Ec-S20359_1	ABC transporter F family member 4	17	0	0	C3
Ec-S27698_1	Phenylalanine ammonia-lyase PAM	17	0	0	C6
Ec-S71193_1	Uncharacterized protein	16	0	0	C1
Ec-S44030_4	Cyclin-dependent protein kinases 2 CDKC	13	0	0	C1
Ec-S06490_3	Uncharacterized protein	13	0	0	C2
Ec-S11263_1	Bpg2-like protein (Fragment)	11	0	0	5-Aza
Ec-S41184_1	1,4-alpha-D-glucan maltohydrolase	10	0	1	WS
Ec-S48520_2	Uncharacterized protein	6	6	0	C1
Ec-S09768_1	Uncharacterized protein	0	11	0	C8
Ec-S63962_1	Reverse transcriptase (RNA-dependent DNA polymerase)	0	10	0	5-Aza
Ec-S77906_2	Uncharacterized protein	0	10	0	C1
Ec-S53645_2	Apoptosis inhibitory protein 5 (API5)	0	10	0	WS
Ec-S29620_1	Amino acid permease 1	0	6	4	WS
Ec-S77915_1	ALE2 AtPERK1	0	1	18	WS
Ec-S76299_2	Agamous-like MADS-box protein AGL9 homolog	0	1	17	C6
Ec-S71479_1	Probable polygalacturonase	0	1	9	C3
Ec-S06731_4	DNA ligase 1	0	0	18	C2
Ec-S64428_1	(1->3)-beta-glucan endohydrolase 10	0	0	16	C1/C8
Ec-S64428_1	Glucan endo-1_3-beta-glucosidase 14	0	0	16	C8
Ec-S61275_5	Uncharacterized protein	0	0	14	WS
Ec-S08686_1	GDSL esterase/lipase At4g10955-like	0	0	13	WS
Ec-S63591_3	Uncharacterized protein	0	0	12	C4
Ec-S66538_1	Uncharacterized protein	0	0	12	C5
Ec-S76981_2	hypothetical protein CCACVL1_13302	0	0	11	C1
Ec-S72001_2	60S ribosomal protein L37-3	0	0	10	5-Aza
Ec-S62651_1	Uncharacterized protein	0	0	10	C4
Ec-S28865_5	Transcription factor WER-like	0	0	10	WS

Fig. 1. Boxplots of the percentage of methylated cytosines found on leaf genomes of adult plants obtained from seeds with no treatment (Control) and seeds treated with 5-Azacytidine (5-Azacytidine), and grown with optimal watering (blue, WW) or recurrent drought stress (yellow, WS). (a) Global methylation, (b) methylation percentage at CG, (c) methylation percentage at CHG, (d) methylation percentage at CHH. *** $P<0.001$. N=8 for each factor combination level. Note that in all cases the interaction between the two study factors was statistically significant (see Results for further details).

Fig. 2. a) Number of Differentially Methylated Cystosines (DMCs) by context induced by application of 5-Azacytidine at seed stage (5-Aza) and/or recurrent water stress (WS) that were hyper- or hypo-methylated compared to control plants. b) Heat map visualization of the 6848 DMCs obtained. Average methylation level for each experimental group (8 samples each) is shown. DMCs are clustered using the complete linkage clustering, and the scale is shown on the right, in which red and blue correspond to a higher and a lower averaged methylation status, respectively. c) The lower panels show the methylation difference between the two conditions of each study factor, blue for 5-Aza and orange for WS, at every DMC evaluated on each of the three sequence contexts analyzed. Each dot represents a certain DMC position for which a Y-axis value of 1 indicates complete lack of methylation in treated plants and full methylation in control condition.

Fig. 3. Visualization of 5-Azacytidine and water stress treatment interaction effects on the methylome. a) Hierarchical clustering of the 4073 DMCs whose methylation changes exhibited a significant interaction between the two study factors and heatmap of K-means clusters (K = 8) showing common DMC patterns. The number of cluster minimized the within cluster sum of squares. Size depicts the number of DMC within each cluster. b) Average methylation levels for each experimental group on each K-means cluster.

Fig. 4. Distribution of the DMCs identified in 5-Azacytidine treatment (top) and water stress treatment (bottom). Exons, introns, and promoters corresponds to DMCs overlapping with genes, and TE corresponds to DMCs overlapping with TEs. All other remaining DMCs are classified as intergenic

890 **Fig. 5**. Distribution of the TE classes along *Erodium cicutarium* genome and

891 distribution of TE classes for the overlapped-DMCs (N = 285,236 and 492,

892 respectively).