CONSTANS alters the circadian clock in *Arabidopsis thaliana*

Pedro de los Reyes[1], Francisco J Romero-Campero[1,2], He Gao[3], Gloria Serrano-
Bueno[1], Jose M Romero[1,4], and Federico Valverde[1]*

[1] Plant Development Group - Institute for Plant Biochemistry and Photosynthesis, Consejo Superior de
Investigaciones Científicas, Universidad de Sevilla, Seville, Spain.

[2] Department of Computer Science and Artificial Intelligence, Universidad de Sevilla, Seville, Spain.

[3] Department of Plant Developmental Biology, Max Planck Institute for Plant Breeding Research,
Cologne, Germany.

[4] Department of Biochemistry and Molecular Biology, Universidad de Sevilla, Seville, Spain.

25 *** Corresponding author**: federico.valverde@ibvf.csic.es

27 **Keywords**: CONSTANS, circadian clock, photoperiod.

29 **Running title:** Control of Arabidopsis clock by CONSTANS

Abstract

Plants are sessile organisms that have acquired highly plastic developmental strategies to adapt to the environment. Among these processes, the floral transition is essential to ensure reproductive success and is finely regulated by several internal and external genetic networks. The photoperiodic pathway, which controls the plant response to day length, is one of the most important pathways controlling flowering. In *Arabidopsis* photoperiodic flowering, *CONSTANS* (*CO*) is the central gene activating the expression of the florigen *FLOWERING LOCUS T* (*FT)* in the leaves at the end of a long day. *CO* expression is strongly regulated by the circadian clock. However, to date, no evidence has been reported regarding a feedback loop from the photoperiod pathway back to the circadian clock. Using transcriptional networks, we have identified relevant network motifs regulating the interplay between the circadian clock and the photoperiod pathway. Gene expression, chromatin immunoprecipitation experiments and phenotypic analysis allowed us to elucidate the role of CO over the circadian clock. Plants with altered *CO* expression showed a different internal clock period, measured by daily rhythmic movements in the leaves. We show that CO is able to activate key genes related to the circadian clock, such as *CCA1*, *LHY*, *PRR5* and *GI,* at the end of a long day by binding to specific sites on their promoters. Moreover, a significant number of PRR5 repressed target genes are upregulated by CO, and this could explain the phase transition promoted by CO. The CO-PRR5 complex interacts with the bZIP transcription factor HY5 and helps to localize the complex in the promoters of clock genes. Our results indicate that there may be a feedback loop in which CO communicates back to the circadian clock, feeding seasonal information to the circadian system.

Introduction

In order to interact with a regularly changing environment, plants have developed a great variety of morphological and physiological responses. The transcriptional programs that control these responses allow plants to respond to external stimuli with a high degree of plasticity throughout their life cycle (De Mendoza et al., 2013)☐. One of the most important decisions in the plant life cycle is flowering, which activates the molecular and metabolic changes that induce the transition from the vegetative to the reproductive growth programs (Piñeiro and Jarillo, 2013; Andrés and Coupland, 2010; Kinoshita and Richter, 2020)☐. This transition is finely regulated by multiple factors, since making this decision at the right moment ensures reproductive and evolutionary success. Thus, plants are able to recognize seasonal changes by measuring day length (photoperiod), allowing them to adapt to and grow at different latitudes. The photoperiodic pathway is one of the major components controlling flowering and is regulated by light sensors and an autonomous internal timekeeper called the circadian clock (Sanchez and Kay, 2016; Shim et al., 2017). The central gene of the photoperiod pathway is *CONSTANS* (*CO*), which acts as integrator of day length signals and is an output of the clock (Putterill et al., 1995; Suárez-López et al., 2001; Valverde, 2011).

   CO encodes a protein with two conserved protein domains: two B-boxes located at the amino terminus, involved in protein-protein interactions, and a carboxyl CCT (CONSTANS, CO LIKE, TIMING OF CAB1) domain, which includes a nuclear transport signal and mediates interactions with DNA and diverse proteins (Laubinger et al., 2006; Valverde, 2011)☐. Recently, a role for the central part of CO in regulating floral senescence has been proposed (Serrano-Bueno et al., 2022). *CO* is a hub in the photoperiodic regulatory network; it receives signals from a multitude of different pathways, such as day length and circadian clock (Serrano-Bueno et al., 2021), and triggers the expression of the florigen *FLOWERING LOCUS T* (*FT*), which induces flower differentiation (Ahn et al, 2008)☐. The circadian clock is an endogenous transcriptional clock that produces biological rhythms with periods of 24 h. This inner clock allows plants to synchronize and anticipate external changes and is one of the most important factors regulating flowering. In *Arabidopsis*, the circadian

clock consists of three transcriptional loops, the core loop (CCA1, LHY and TOC1),

the morning loop (PRR5, PRR7 and PRR9) and the evening loop (LUX, ELF3, ELF4

and GIGANTEA) (McClung, 2006; Sanchez and Kay, 2016)□. *CO* expression and

protein stability are highly regulated by the circadian clock and light inputs (Valverde

et al, 2014; Suárez-López et al., 2001; Mizoguchi et al., 2005)□. The GIGANTEA

(GI) and FLAVIN-BINDING, KELCH REPEAT, F-BOX 1 (FKF1) complex participates

in this control by regulating *CO* transcription by binding to its promoter in the late

afternoon and promoting the degradation of CYCLING DOF FACTORS (CDFs),

which act as repressors of *CO* expression (Imaizumi et al., 2005; Sawa et al.,

2007)□. Later, FLOWERING BHLH proteins (FBHs) can bind to these genomic sites

in the *CO* promoter to activate its expression (Ito et al., 2012)□. In addition, The

PSEUDO RESPONSE REGULATORs PRR9, PRR7 and PRR5 indirectly promote

photoperiodic flowering by repressing *CDF1* (Nakamichi et al., 2007)□.

Although the *CO* transcript is highly expressed from late afternoon to dawn, in

terms of protein stability, CO is unstable in the dark and stable during the daytime via

a process mediated by different photoreceptors and E3 ubiquitin ligases (Valverde et

al., 2004)□. During the daytime, the RING finger-containing E3 ubiquitin ligase

HOS1 (HIGH EXPRESSION OF OSMOTICALLY RESPONSIVE GENES1) interacts

with CO and targets it for degradation (Lazaro et al., 2012), thus□ allowing activation

of *FT* specifically at the right moment of the day. During the night time, COP1 and

SPA1 promote CO degradation (Laubinger et al., 2006; Jang et al., 2008)□. After

dawn, several photoreceptors begin to stabilize CO to promote its accumulation

during the first hours of the day. In response to blue light, CRY1 and CRY2 interact

with COP1 and SPA1 to avoid the degradation of CO by the COP1-SPA1 complex

(Yanovsky and Kay, 2002; Liu et al., 2008; Liu et al., 2011; Zuo et al., 2011)□.

Moreover, during the day, two phytochromes affect CO stability in different ways; in

response to red light, PHYB interacts with HOS1 to promote the degradation of CO,

but in the late afternoon, PHYA is able to disrupt the COP1-SPA1 complex to stabilize

CO in presence of far-red light (Valverde et al., 2004, Lazaro et al., 2012; Endo et al.,

2013; Hajdu et al., 2015; Lazaro et al., 2015; Sheerin et al., 2015)□.

In addition to the described photoreceptor-pathways, there are some

circadian-related proteins that regulate CO accumulation in response to blue light. In

the morning, ZEITLUPE (ZTL) forms a complex with GI to inhibit CO stability (Song et
al., 2014)☐, but in the afternoon, blue-light enhances CO accumulation by interacting
with FKF1 (Song et al., 2012). In fact, FKF1, GI and CO could form a complex that
stabilizes CO (Song et al., 2014)■. Recently, it has been described that PRRs
stabilize CO and promote flowering in response to day length (Hayama et al.,
2017)☐. CO stability and its role as transcriptional regulator are both affected.
DELLA proteins, the immunophilin FKBP12, the RING domain protein BOI, the B-
boxes BBX19 and miP1a, and the chromatin remodeling factor PICKLE can regulate
CO transcriptional function in different ways (Wang et al., 2014; Nguyen et al., 2015;
Graeff et al., 2016; Wang et al., 2016; Xu et al., 2016; Jing et al., 2019;
Serrano-Bueno et al., 2019)☐. Therefore, CO function is highly regulated by multiple
pathways, all of them ensuring the precise activation of *FT* and the initiation of
flowering. Regarding other biological processes regulated by CO, it has been
described that CO is able to modify starch metabolism through the activation of the
GRANULE BOUND STARCH SYNTHASE (*GBSS*) gene (Ortiz-Marchena et al.,
2014)☐. CO may also be involved in stomatal opening (Ando et al., 2013) and play a
role in floral senescence (Serrano-Bueno et al., 2022)☐. However, the involvement
of CO in other physiological processes is still largely unknown.

Although considered an output of the circadian machinery, it is not known if
CO is able to affect the clock. However, since CO presents a CCT domain in its
carboxyl terminus, similar to that present in PRRs, that may allow binding to circadian
genes promoters (Nakamichi et al., 2010), it was suggested that CO could participate
in circadian clock control (Strayer et al., 2000). Circadian network retrograde
signaling is often found, and in this study, we provide conclusive evidence that in
addition to being an output of the clock, CO can also regulate the expression of key
genes involved in the circadian time pacer. Therefore, the transcriptional programs
controlling the floral transition and the circadian clock are intertwined, constituting a
complex system exhibiting feedback loops. Recently, in order to unravel this system,
omics techniques based on high throughput sequencing have been applied.
However, these data are commonly fragmented, and molecular systems biology
approaches are necessary to integrate them and obtain a global comprehension of

the interactions between the transcriptional programs controlling the floral transition
and the circadian clock.

Network theory constitutes one of the central paradigms in data integration
and analysis in molecular systems biology (Barabási, 2015)□. The integration of
massive amounts of omics data has helped generate biological networks for
photosynthetic organisms that reveal emergent properties and improve the
understanding of processes such as the evolution of CO-like transcription factors
(TFs) in *Viridiplantae* (Romero-Campero et al., 2013)□, the conservation of circadian
patterns across the plant phylogenetic tree (de los Reyes et al., 2017)□, carbon
mobilization during the floral transition in *Arabidopsis* (Ortiz-Marchena et al., 2014)□
and the involvement of hormonal control (Serrano-Bueno et al., 2022). In this work,
we have constructed a transcriptional gene network by integrating deep sequencing
of chromatin immunoprecipitation (ChIP) data generated from key TFs involved in the
floral development/transition and circadian clock control. Feeding data on CO into the
analysis of this network allowed us to propose a new function for photoperiodic
signaling: the formation of a feed-back loop regulating the circadian clock in
Arabidopsis.
172
173
174

Results

Transcriptional networks reveal a connection between the circadian clock and photoperiodic signaling

To examine the interplay between the transcriptional programs controlling the circadian clock and photoperiod, we analyzed ChIP-seq data from 33 Arabidopsis TFs involved in light signaling, flowering and the circadian clock and built a transcriptional network called CircadianFloralNet. This network is composed of 20,601 nodes and 89,377 edges, comprising 75% of the *Arabidopsis* genome (Figure S1A). We also generated from CircadianFloralNet a transcriptional gene network containing only the analyzed TFs and their interactions (Figure S1B). This network constitutes the regulator core of CircadianFloralNet and was called CircadianFloralTFNet. Therefore, CircadianFloralTFNet is a transcriptional gene network that captures the transcriptional program governing the interaction between the circadian clock, light and the flowering process. CircadianFloralNet is a scale-free network showing a power-law node degree distribution (Figure S1C) according to the Kolmogorov-Smirnov test (Goldstein et al., 2004). Furthermore, it is a small-world network, implying that the distance between two random nodes is shorter than expected (Figure S1D). These topological properties indicate that transcriptional programs governing these processes are highly connected (Wagner and Fell., 2001; Tong et al., 2004; Yook et al., 2004).

To determine the possible input of photoperiod in this network, we carried out a genome-wide analysis of chromatin immunoprecipitation sequencing (ChIP-seq) experiments in 35S:*CO* and *co-10* 7-DAG seedlings around *Zeitgeber* Time (ZT) 14 under LD using specific CO antibodies (Serrano-Bueno et al., 2020). The analysis of CO ChIP-seq data identified 3,214 peaks and 2,418 putative target genes (Table S1), and we fed this information into CircadianFloralNet. Next, we generated CircadianFloralTFNet including CO and classified the nodes into light, flowering or clock genes based on their TAIR classifications (Fig 1A). CO was found to bind to the promoters of the *PIF1*, *PIF3* and *PIF4* genes related to light signals and to the promoters of *SOC1* and *SVP*, which are specifically associated with flowering time control. However, all genes clustering in the clock function in CircadianFloralTFNet,

including *CCA1*, *LHY*, *TOC1, PRR5, PRR7,* and *PRR9,* showed one or several peaks

of CO binding on their promoters (Fig 1A), indicating that CO significantly bound to

the promoters of genes involved in circadian rhythm regulation, including the core

components of the clock, CCA1/LHY and TOC1.

In order to identify relevant non-random components in the interactions

between the circadian clock and photoperiod, we performed a network motif analysis

on CircadianFloralTFNet. A network motif is a subgraph that appears in the network

of interest at a significantly higher number of times when compared to similar random

networks. Therefore, we tested the significance of all possible subgraphs consisting

of one, two or three nodes by generating 100,000 random graphs with the same

topological features as the original network. The most represented motifs identified

are shown in Figure 1B. One of the most significant and common motifs was the

'feedback loop with output' (p-value $<10^{-5}$). This meant that a high interconnection

existed between biological processes within the transcriptional programs that

integrate all processes analyzed in *Arabidopsis.*

The connectivity we identified in CircadianFloralNet confirmed the results of

other previous studies (Suárez-López et al., 2001; Imaizumi et al., 2005; Sawa et al.,

2007)☐ that established that *CO* expression was regulated by the circadian clock.

Specifically, PRR5 binds to cis elements in the *CO* promoter in order to regulate its

expression, and consequently, both genes are connected in CircadianFloralNet.

Additionally, a substantial number of genes of the *CO*-like (*COL*) family are regulated

by PRR5 and other central proteins of the circadian clock, such as *CCA1* and *PRR7*,

as shown in the subnetwork presented in Figure S1E. Since the feedback loop is a

network motif in CircadianFloralTFNet and PRR5 regulates *CO,* it was expected that

CO, in turn, would also regulate the expression of *PRR5*. We verified CO binding to

the *PRR5* promoter through ChIP-seq and ChIP-QPCR experiments (Figure S1F).

This result implies that one of the interactions between the flowering program and the

circadian clock is exerted by CO and PRR5.

In fact, based on ChIP-seq data, CO and PRR5 constitute a 'feedback loop

with multiple output' motif, represented in Figure 1C, that regulates a set of

common target genes related with the circadian clock (*CCA1, LHY, RVE1,8,*

TOC1, PRR7,9, LNK1,2) and photoperiod signaling (*GI, CDFs, BBXs, GBBS1*

and BAM1,9).

CONSTANS alters the expression of circadian clock genes

To investigate the proposed effect of CO on the clock, we carried out RNA-seq experiments from 35S:*CO* and Col-0 plants using RNA from LD (LD, 16 h light: 8 h night) grown seedlings collected in the evening at ZT14. Similarly, we included an available RNA-seq dataset of Col-0 and *co-10* mutant plants grown in continuous light (LL) (Gnesutta et al., 2017; GSE205675)□. After analyzing the data via standard pipelines (see methods), we found 637 genes with higher than two-fold expression in 35S:*CO* than in WT (Table S2) and 714 downregulated genes showing a logFC < -1.5 in *co-10* plants (Table S3). To obtain high-fidelity CO-regulated genes, we overlapped both sets of genes in a Venn diagram consisting of 148 intersecting genes (Figure 2A, Table S4). According to the GO-term enrichment analysis, these genes are involved in several biological processes, such as the response to light stimuli, developmental processes and, as we expected, rhythmic processes and circadian rhythm, with adjusted p-values of 7.84e-06 (Figure 2B, Figure S2A, Table S5). Specifically, several key circadian clock genes including *GI, CCA1* and *LHY* showed higher expression in *CO*-overexpressing plants and lower expression in *co-10* mutants (Figure S2B, Table 1).

To further asses if CO was able to modify the circadian pattern of expression of other genes, we analyzed RNA-seq data from a transcriptome of Col-0 plants under LD conditions in a 24 h period (GSE43865, Rugnone et al., 2013) and estimated gene expression levels at ZT2, ZT6, ZT10, ZT14, ZT18 and ZT22. Using this analysis, we identified genes exhibiting a circadian pattern of expression and classified them into clusters depending on the time point at which they reached their minimal and maximal expression (see methods). To identify the specific circadian patterns affected by *CO*, we studied the statistical significance of the intersections between the circadian gene clusters and the activated genes in 35S:*CO* at ZT14. Only three circadian clusters presented a significant intersection according to Fisher's exact test, as shown in Table S6. All the clusters showing a significant intersection presented a morning peak and an evening/afternoon trough. We concatenated these clusters into a single group called the "morning-phased" cluster and, again, the intersection of these genes with the activated genes in 35S:*CO* was significant

(Figure S3A), in agreement with previous studies (Gnesutta et al., 2017)□. This
result shows that CO activates the expression of genes in the evening that naturally
exhibit their minimum expression at that time of day, thus altering their circadian
patterns.

To confirm the effect of CO on the clock, using RT-qPCR, we measured the
expression of some of the genes regulated by CO every 4 h during a 24 h period in
WT, *co-10* mutant and 35S:*CO* plants (Figure 2C). We found that *PRR5*, *PRR7*, *GI*
and clock-output genes such as *BAM9* were activated in 35S:*CO*, especially at ZT14,
when CO protein is stabilized under LD (Suárez-López et al., 2001). This reflected
the fact that *CO* overexpression, and its absence, altered the daily expression of
clock genes.

To further asses the effect of photoperiod on the clock, we tested if CO was able
to alter some physiological aspects of the circadian behavior of *Arabidopsis*. In
plants, one of the most well-known and characterized circadian clock outputs is the
daily rhythmic movement of leaves. We tracked this movement in LD entrained 7-
DAG seedlings from WT, *co-10* and 35S:*CO* plants transferred to LL and estimated
the period and phase of circadian parameters. As shown in the graphic in Figure 2D
and its quantification in Figure 2E, 35S:*CO* plants exhibited a significant shorter
period than WT, while *co-10* mutant showed a longer period. However, this behavior
was not conserved in terms of phase (the time when the leaves are maximally
pointing upward), as both 35S:*CO* and *co-10* showed an early phase compared with
WT plants (Figure S3B). The change in nastic rhythms shown in *CO* mutants is a
strong evidence that CO is involved in circadian clock modification.

To determine if CO could be altering the circadian rhythms by binding to clock
gene promoters, we further analyzed the ChIP-seq data from 35S:*CO* and *co-10*
plants at DAG7. The CO ChIP peak distribution was found to be enriched mainly in
promoter regions near the transcription start sites (65%), and to a lesser degree,
regions up to 2 kb away (10%), downstream regions (6%) and distant intergenic
regions (7%), indicating that it was probably a *bona fide* transcription factor (Figure
3A). Next, we compared the ChIP-seq data with the gene expression profiles from
35S:*CO* (Figure S4A). We found that almost 60% (318 out of 540) of the upregulated
genes in 35S:*CO* were direct targets of CO, including its main target, the floral
integrator *FT*, as well as other photoperiodic growth/development related genes, such

as *CDF3* and *CDF6,* and key circadian clock genes, such as *PRR5*, *LHY, GI*, *RVE8*
and *RVE4* (Figure S4B). Binding of CO to clock gene promoters suggested that CO
was able to alter not only photoperiodic genes expression, as previously suggested
(Onouchi et al., 2000), but also the expression of circadian clock-related genes.

It was described that CO regulates *FT* expression by binding to its promoter in
association with NF-Y/HAP heterodimers to bind to CCAAT elements (Wenkel et al.,
2006) and to CORE sites (Tiwari et al., 2010)□. However, recently, it has been
shown that CO imparts DNA sequence specificity to the NF-Y heterodimer
resembling a histone pocket, thus allowing the complex to bind to the CCACA
element (Gnesutta et al., 2017, Shen et al., 2020). We performed unbiased DNA
motif discovery analysis in the CO CHiP-seq data and confirmed the presence of the
CCACA element ($p>10^{-19}$) and another element resembling a CORE-like element
($p>10^{-13}$) (Figure 3B); unexpectedly, we identified a G-box (CACGTG) that was
significantly enriched ($p>10^{-18}$) in CO binding peaks. Our analysis showed that CO
could bind to G-box sites in *CCA1, LHY* and other loci (Fig 3C) but not in the *FT*
promoter, where it was associated exclusively with CCACA sites, revealing a NF-
Y/HAP-independent mechanism.

Together, these results indicated that CO is not only an output of the circadian
clock pathway but also a factor exerting feedback regulation over key circadian clock
genes and circadian patterns, implying a novel function for CO in altering rhythmic
processes in *Arabidopsis*.

CO, PRRs and HY5 share genomic binding sites in circadian clock regulated
genes

G-box motifs are the binding sites of bZIP and bHLH TFs and have been described in
the transcriptional regulation exerted by PRRs, thus implying an association with
these TFs (Liu et al., 2016)□. Under short days (SD, 8 h light, 16 h dark), the bHLH
PIFs have been described to bridge the binding of PRRs to DNA to sequentially
repress shared target genes related to growth, such as *CDF5* (Martín et al., 2018)□.
Additionally, PIF4 is involved in the temperature-dependent activation of *FT* through
binding to CO under SD (Fernández et al., 2016), but no involvement on the clock

under LD was reported. On the other hand, the bZIP LONG HYPOCOTYL 5 (HY5) is

a light-inducible TF (Osterlund et al., 2000)☐ that binds to G-boxes (Lee et al., 2007;

Young et al., 2008; Zhang et al., 2011; Binkert et al., 2014), and HY5 mutations affect

flowering (Bhagat et al., 2021) and the clock (Hajdu et al., 2018). It has also been

described that HY5 can bind to BBX family proteins from group IV (BBX21-25)

through the b-box domain to activate or inhibit its role in photomorphogenesis (Datta

et al., 2007; Job et al. 2018; Bursch et al., 2020). Therefore, HY5 was a good

candidate to act as bridge mediating the binding of PRR5 and CO to clock-related

gene promoters under LDs. HY5 is also degraded by the COP1/SPA1 complex

during the night (Osterlund et al., 2000)☐, similar to CO (Jang et al., 2008), and it

has been shown that CO and HY5 interact with the same chromatin remodeling

factor, PKL (Jing et al., 2013; Jing et al., 2019)☐.

Indeed, when we compared the genome-wide analysis results for the ChIP-seq

data for PRR5 (Nakamichi et al., 2012)☐ and ChIP-on-chip data for HY5 (Lee et al.,

2007)☐ with the ChIP-seq data for CO, a significant overlap between the direct

targets of PRR5, HY5 and CO was found (Figure 4A). After carrying out a DNA motif

discovery analysis over the promoter sequences (1 kb) of the different gene sets

(Figure 4A, right; Table S7), we found a motif resembling the CCACA element

(Gnesutta et al., 2017) in the gene promoters regulated exclusively by CO (set 1),

indicating that it binds to these regions via an NF-CO trimer. A matrix containing a G-

box motif was enriched in the genes regulated by PRR5 (set 2), as described in

previous studies (Liu et al., 2016). Notice that the most significant enriched DNA

matrices come from bHLH TF ChIP-seq studies (Table S7), suggesting that PRR5

regulates the transcription of these genes in association with bHLHs (Martín et al.,

2018). However, a DNA matrix containing an E-box motif *ACGTG* was recovered

from the genes regulated by HY5 (set 3). This suggests a certain degree of specificity

imparted by the protein with which HY5 associates. We also found a G-box motif

enriched in the gene promoters bound by CO, HY5 and PRR5, indicating that this

DNA element plays a key role in the transcription of these clock-regulated genes.

However, the most significant DNA matrices were described in bZIP TF ChIP-seq

studies (Table S7).

To support our hypothesis, we followed a similar approach to analyze the
genome-wide binding sites of CO, HY5 and PRR5 (Huang et al., 2012; Nakamichi et
al., 2012; Liu et al., 2013; Liu et al., 2016) (Table S8). First, we found a significant
overlap between the three sets of binding regions according to permutation tests
(Table S9). Next, using the *intersectBed* function from *bedtools*, we retrieved the
overlapping peaks for the three TFs (a), the intersection between CO and PRR5 sites
(b), the intersection between HY5 and PRR5 sites (c) and the genome regions bound
by CO (d) (Figure 4B, left). In order to determine the DNA elements involved in the
regulation exerted by these TFs, we performed DNA motif discovery analyses over
the different binding site sets (Figure 4B, right). A CCACA element (Gnesutta et al.,
2017) was enriched exclusively in the CO binding sites (d), which is consistent with
previous studies based on transcriptomic data (Gnesutta et al., 2017). However, G-
boxes were enriched in the other intersections, a-c sites (Figure 4B, right). All these
results indicate that CO and PRR5 bind to a G-box to control the transcription of
clock-regulated genes, probably in association with a bZIP TF, such as HY5, as
shown in Figure 4C for the *CCA1* and *LHY* promoters.

To demonstrate HY5 binding to these sites, we performed ChIP experiments
using GFP-Trap antibodies in 35S:*GFP*:*HY5* (*hy5-215*) plants and observed a
significant enrichment of these amplicons compared with the control (WT plants) that
was absent in the *FT* promoter (Figure S5A, B). These results indicated that HY5
binds to the same sites as CO in these circadian clock promoters but not the CORE
site associated with CO in the promoter of *FT*, highlighting that CO has a different
function related to circadian rhythm that is independent of the photoperiodic flowering
pathway.

To further confirm the role of HY5 in CO-dependent clock gene transcriptional
modification, we crossed the *hy5-2* mutant with 35S:*CO* plants to obtain the double
35S:*CO hy5* mutant (Figure 5A). These plants showed a slight decrease in flowering
time compared to 35S:*CO* plants (Figure 5B). 35S:*CO hy5* plants showed high
expression of *CO* and no expression of *HY5* mRNA or protein, as confirmed by QRT-
PCR (Figure S6A) and Western blot experiments (Figure S6B), respectively.
Consequently, the expression of the clock-related genes *CCA1*, *PRR5, PRR7* and *GI*
activated by CO was reduced (Figure 5C), further confirming the role of HY5 in CO-
dependent clock gene expression.

CO, PRR5 and HY5 proteins interact *in planta*

The interaction between CO and some members of the PRR family was previously
shown; this binding stabilizes CO and enhances photoperiodic flowering under LD
conditions (Hayama et al., 2017)□. However, considering that CO and PRR5 bind to
the same sites in some promoters and that these two proteins seem to have the
opposite effect over transcription, it is possible that CO can gradually remove PRR5
from the DNA and promote transcription. Therefore, a transitory complex consisting
of HY5, PRR5 and CO during the evening could exist to alter gene expression. To
examine this possible ternary complex, we performed different protein-protein
interaction approaches.

First, we performed a bimolecular complementation (BiFC) assay in *Nicotiana*
between CO, HY5 and PRR5 in pairs, and we observed the interactions between
them in the nucleus and found different distribution patterns (Figure 6A). We also
performed yeast-two-hybrid experiments with the three main domains of CO
(BBOXes, middle domain and CCT domain) to confirm the interaction with both HY5
and PRR5. As can be seen in Figure 6B, and different from the previous results on
the binding of other BBXs to HY5 through the BBoxes (Bursch et al., 2020), CO
interacts with HY5 and PRR5 through the middle domain and, to a lesser extent, the
CCT domain. To further investigate if there could be a ternary complex, we tested the
colocalization of the three proteins in the nucleus of *Nicotiana* cells. To do this, we
reconstituted the YFP fused to HY5 and PRR5, and coinfiltrated a CO:CFP construct.
We detected a significantly higher colocalization in speckles, compared with the
control with CFP alone (Figure 6C). Following this approach, we measured
fluorescent resonance (FRET) between CFP and YFP through a sensitized emission
method, and we found higher values in the CO/HY5/PRR5 combination than in the
negative controls, confirming the close interaction among the three proteins (Figure
6D).

Since CO and HY5 physically interact and both proteins bind to the same
regions in clock gene promoters, we investigated if they could associate with DNA
when combined. To examine this possibility, the immunoprecipitated complex
resulting from the ChIP experiment was analyzed by immunoblot using anti-HY5

antibodies. We detected HY5 protein in the 35S:*CO* sample but not in the control
(Figure 6E), confirming the presence of a complex between CO-HY5 and DNA.

The clock-related leaf movement was also affected in *PRR5* and *HY5* mutants,
showing an acceleration of the period compared to *CO*ox plants (figure S7A). The
phase of the nastic movements of leaves was also affected in *prr5* and *hy5* mutant
plants, again showing a reduction in the phase (Figure S7B).

Together, these results indicated that these three proteins could form a
complex that binds to the DNA, acting HY5 as scaffold, since this type of TF, bZIP, is
largely described to be capable to directly binding to DNA (Chattopadhyay et al.,
1998; Abbas et al., 2014; Binkert et al., 2014)☐.

<u>Discussion</u>

A network science perspective is essential for understanding complex interconnected
systems (Barabási, 2015)☐. Transcriptional programs governing circadian clock and
photoperiodic pathways showed a high relatedness in CircadianFloralNet according
to its scale-free and small-world properties, which revealed many unexplored
connections. In the field of graph theory, network motifs are particularly interesting
because they confer particular functions. Network motifs are subgraphs that appear
in an observed network significantly more frequently than they do in other
comparable networks (Milo et al., 2002; Shen-Orr et al., 2002; Stone et al., 2019).
Despite the high complexity of gene networks, a certain level of simplicity can be
found in these sub-structures, which are present throughout biological networks and
across species (Alon, 2007)☐. Much debate has focused on the validity of analyzing
these network motifs because they are components of large and complex networks
(Mellis and Raj, 2015)☐. Indeed, CircadianFloralNet represents all potential
interactions under many conditions, tissues, growth stages and times of day;
furthermore, smaller and insulated networks have to be considered. Thus, the
network motif formed by *CO* and *PRR5* functions in the evening under LD conditions
during the floral transition.

In recent years, the circadian clock in *Arabidopsis* has emerged as a complex
regulatory system; it is not a simple molecular timer but a regulator of multiple

processes involving hormone, metabolism and stress pathways, some of which exert
feedback regulation on the clock (Sanchez and Kay, 2016)☐. Indeed, several TFs
that contribute to clock function have been identified (McClung, 2014)☐ and it has
been proposed that different tissues may show differences in the clock (Endo et al.,
2004).

CO function is strongly regulated at both the transcriptional and post-
translational levels in *Arabidopsis*, indicating that the photoperiodic pathway has
been subjected to high selective pressure (Romero and Valverde, 2009)☐. Other
than flowering transition and reproductive development, no other functions have been
described for CO, but recent studies suggested that it could regulate floral jasmonate
signaling (Serrano-Bueno et al., 2022) and circadian genes (Gnesutta et al., 2017)☐.
Here, we show that CO alters the circadian clock by binding to the promoters of key
clock-related genes, revealing an undescribed function. Therefore, there is an even
more complex regulation between the circadian clock and photoperiod pathways than
previously shown.

PRRs have been described as repressors of their direct target genes.
According to our network based on ChIP-seq data, PRR5 could bind to the *CO*
promoter. Nevertheless, PRRs cause a global positive effect over the photoperiod
pathway and CO-FT module regulation, since they stabilize CO (Hayama et al.,
2017)☐ and repress *CDF1* expression, triggering the net activation of *CO* and *FT*
expression (Nakamichi et al., 2007)☐. Then, the control of *PRR5* gene expression by
CO establishes a positive feedback loop between the circadian clock and the
photoperiodic pathway. In addition to PRRs, CO binds to the *GI* promoter. Since GI
exerts an activating role over CO (Imaizumi et al., 2005; Sawa et al., 2007; Song et
al., 2014)☐, this could also form a positive feedback loop to enhance and fix the
photoperiodic response in the afternoon under LDs. The crosstalk between flowering
and the circadian clock exerted by CO could help to explain the existence of tissue-
specific clocks (Endo et al., 2014)☐. These results support the already established
concept of complex interlocking loops instead of a unidirectional system composed of
input-oscillator-output.

Moreover, it seems that in addition to affecting clock genes such as *PRR5* or
GI, CO could regulate circadian clock target genes, including a significant number of

PRR5 targets. This set of genes would be repressed by PRR5 and activated by CO,

forming a transcriptional switch. In order to move from a vegetative phase to a

reproductive phase, massive changes in gene expression are needed in plants; thus,

CO could provide the activation of a set of repressed genes and promote the phase

transition (Figure 7).

There has been a long controversy about whether CO could act as a TF or as

a coactivator (Blackman and Michaels, 2010)□. Recently, it has been discovered

that CO can directly bind to CORE elements and associate with NF-Y/HAP

heterodimers to bind to CCACA elements, acting as an activator of gene expression

(Wenkel et al., 2006; Tiwari et al., 2010; Gnesutta et al., 2017)□. Indeed, CO activity

can be controlled by the formation of different types of protein complexes, even

repressor complexes formed by TOPLESS (Graeff et al., 2016)□. Here, we

demonstrate that CO is able to bind to G-box elements in DNA through the bZIP

transcription factor HY5, a TF involved in many processes and pathways (Gangappa

and Botto, 2016)□. Together, these findings show a great plasticity in CO function as

a transcriptional regulator and its ability to recruit circadian clock machinery at the

end of a long day. Depending on its partners, CO will exhibit different effects over

transcription and affinity for different genome binding sites; thus, conferring high

plasticity to plants in response to daylength changes.

Methods

Plant material and growth conditions

Plant lines in the Col-0 background used in this study included 35S:*CO* (Onouchi et

al., 2000), *co-10* (SAIL_24_H04), *hy5-2* (SALK_056405C), 35S:*GFP:HY5* (*hy5-211*)

(kindly provided by Dr Vicente Rubio), and *prr5-1* (SALK_006280; Michael et al.,

2003). 35S:*CO hy5* double mutant plants were generated by crossing 35S:*CO* with

hy5-2 to form homozygotes. *Arabidopsis* seedlings were grown in a SG-1400

phytotron (Radiber SA, Spain) under a LD light regime with temperatures ranging

from 22°C (day) to 18°C (night) and 100 µEm^{-2}s^{-1} light intensity. *Arabidopsis* seeds

were stratified for 3 days at 4°C in the dark and later grown in MS medium. For gene

expression analyses, *Arabidopsis* Col-0, 35S:*CO* and *co-10* seedlings were collected

every 4 h during a complete LD. For monitoring leaf movement, seedlings were
grown in soil for 10 days under LD conditions, transferred to LL conditions and
examined for 4 days.

Chromatin immunoprecipitation experiments

ChIP was carried out as described previously (Bowler et al., 2004)□ with minor
modifications to improve binding enrichment (Lau and Bergmann, 2015)□. Plants
were grown for 10 days under LD conditions and 9 g of seedlings were harvested at
ZT12-ZT16. Then, plant tissue was crosslinked in a vacuum concentrator in presence
of 1 mM DSG for 10 min, 1% (v/v) formaldehyde for 20 min and finally 0.125 M
glycine to stop crosslinking. Later, the tissue was ground in liquid nitrogen and
chromatin was isolated. Chromatin complex immunoprecipitation was performed
using αCO (Serrano-Bueno et al., 2020) and GFP-Trap agarose beads (ChromoTek)
to capture CO and HY5 proteins, respectively. After immunocomplex capture, DNA
was eluted from the beads, crosslinked with proteins, reverse transcribed, and
isolated. The subsequent qPCR was performed as previously described.
Immunoprecipitated samples were normalized to a 10% reverse crosslinked fraction
of each chromatin preparation. Primers used to measure DNA fragments enriched
during ChIP experiments are shown in Table S10. Two biological replicates were
processed for next-generation library preparation using the ThruPLEX DNA-seq kit
(Takara). Next, libraries were sequenced on NextSeq500 (Illumina) at the Cabimer
CSIC Genomics Unit. CO ChIP-seq analysis was implemented as described below.
The raw data has been deposited in Gene Expression Omnibus (GEO) under
accession GSE222657.

ChIP-seq data acquisition and analysis

We used ChIP-seq data generated for transcription factors involved in flowering and
the circadian clock that were available from the Sequence Read Archive (SRA,
https://www.ncbi.nlm.nih.gov/sra) (Wheeler et al., 2005)□. Specifically, data from 33
different transcription factors were downloaded in this study. These transcription
factors and their ChIP-seq data accession numbers are shown in Table S11. The
Arabidopsis reference genome TAIR10 and its corresponding annotation were

downloaded from the Ensembl Plants website (http://plants.ensembl.org/index.html).
The FASTQC software package was employed to examine the read quality of each
sample (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/). The *fastq* files
corresponding to each transcription factor were mapped to the reference genome
using the ultrafast and memory-efficient short read mapper *bowtie* (Langmead et al.,
2009). The read mappings for each sample were stored in *SAM* (Sequence
Alignment Map), *BAM* (Binary Alignment Map) and *bigWig* files. We performed the
"peak calling" step with the software tool *MACS2* (Zhang et al., 2008)□. Finally,
target gene selection was carried out using *PeakAnnotator*, part of the *PeakAnalyzer*
program (Salmon-Divon et al., 2010)□, and an *ad-hoc* R script developed in this
study according to the criterion of the nearest downstream gene. DNA motif
identification was carried out using *HOMER* software (Heinz et al., 2010)□. For
further peak annotation and visualization, *ChIPseeker* (Yu et al., 2015)□ and
ChIPpeakAnno (Zhu et al., 2010)□ R packages were employed. Finally, GO term
enrichment analysis was performed using the R package *clusterProfiler* (Wu et al.,
2021)□. For the CO ChIP-seq analysis, peaks detected in the 35S:*CO* experiment
were filtered based on the peaks found in *co-10* immunoprecipitation.

Transcriptional network construction and analysis
Our transcriptional network was generated and analyzed with the R package *igraph*
(Csardi, 2006)□. Graphical representations of the network were done using
Cytoscape (Smoot et al., 2011)□, a software package for network visualization and
data integration. Specifically, the prefuse force layout method was applied for network
visualization. Additionally, a network topology analysis was achieved. The network
node-degree distribution was calculated through degree function from the *igraph*
package. To analyze if the network adjusts to the scale-free property, the power.law.fit
function (*igraph* package) was used to test the node degree distribution and
determine whether it followed a power law. Furthermore, the network clustering
coefficient and the average path length was calculated through transitivity and
average.path.length functions, respectively. A network motif is a subgraph that
appears a significantly higher number of times in the network of interest when
compared to similar random networks. Therefore, we tested the significance of all

possible subgraphs consisting of one, two or three nodes by generating 100000
random graphs with the same topological features as CircadianFloralTFNet.
599

RNA isolation and gene expression analysis
Total RNA from *Arabidopsis* seedlings (0.1 g leaf tissue) was isolated by the Trizol
(Invitrogen) procedure following the recommendations of the manufacturer. The final
RNA sample was suspended in 21 µL of diethyl pyrocarbonate-treated water,
quantified in an ND-1000 spectrophotometer (Nanodrop) and stored at –80°C. Then,
500 ng of RNA was used to synthesize cDNA employing the QuantiTect Reverse Kit
(Qiagen) and stored at –20°C until qPCR was performed. qPCR was performed with
the iQTM5 Multicolor Real-Time PCR Detection System (Bio-Rad) in 10-µL reactions:
primer concentration of 0.2 µM, 10 ng of cDNA, and 5-µL of SensiFAST SYBR &
Fluorescein Kit (Bioline). Each sample was measured in triplicate. The qPCR
program consisted of (1) one cycle of 95°C for 2 min; (2) 40 cycles of 95°C for 5 s,
60°C for 10 s, and 72°C for 6 s; and (3) one cycle of 72°C for 6 s. Fluorescence was
measured at the end of each extension step, and melting curve analysis was
performed between 55°C and 95°C. The initial concentrations were calculated using
the ddCt Algorithm (Zhang et al., 2016)□. Primers used to measure gene expression
by qPCR are listed in Table S10.

RNA-seq data acquisition and processing
In this study, we used RNA-seq data from the *Arabidopsis* transcriptome publicly
available at the Sequence Read Archive (SRA). We analyzed different studies
(GSE43865, GSE204774 and GSE205675) and obtained 26 samples representing
three different genotypes, Col-0, *co-10* and 35S:*CO,* under LD and LL conditions
(see supplementary material: Table S12). The quality control of each sample was
carried out using the FASTQC software, and the read sequences were mapped to the
reference genome with the ultrafast universal aligner *STAR* (Dobin et al., 2013). The
number of reads that mapped to each gene was perform using the *featureCounts*
software of the *Subread* package (Liao et al., 2013). Genes with less than 10 reads
in all samples were removed, and TMM normalization and log2 transformation were
performed with *calcNormFactors* (*edgeR* package; Robinson et al. 2010) and *voom*

functions (*limma* package; Ritchie et al., 2015), respectively. Next, a linear model with
the variable 'genotype' was fitted for each gene using the *lmFit* function (*limma*
package) in order to carrying out the differential expression analyses.
632

Protein analysis
Proteins were isolated from seedlings grown on MS-agar using the TRIZOL
(Invitrogen) protocol, separated by SDS-PAGE by standard procedures, transferred
to nitrocellulose membranes, incubated with αCO (Serrano-Bueno et al., 2020) or
αHY5 (Agrisera) antibodies, and developed with a chemiluminescent substrate
(Clarity^{tm} Western ECL, Bio-Rad). Blots were visualized and quantified in an
Amersham Imager 680.
640

Determination of circadian genes and clustering techniques
The *bioconductor* R package *RAIN* (Rhythmicity Analysis Incorporating Non-
parametric methods) (Thaben and Westermark, 2014)☐ was used to determine
circadian genes, namely genes exhibiting rhythmic patterns of expression with a
period of 24 h. A time interval of 4 h and a p-value of 0.01 were used. Circadian
genes were classified into clusters based on the positions of their peak (maximal
level of expression) and trough (minimal level of expression) during a 24 h period.
648

Circadian leaf movement analysis
After transfer to LL, the first pair of leaves was monitored using a Raspberry Pi
camera. The leaf movement was tracked using *TRIP* software (Greenham et al.,
2015)☐ run in Octave, and the vertical motion was obtained as a function of time and
analyzed using *BioDare* software (which performs FFT-NLLS period analysis)
(Zielinski et al., 2014)☐ and the *CircaCompare* package (Parsons et al., 2020)☐
implemented in R.
656

657 **Transitory expression in *Nicotiana benthamiana***
658 To perform BiFC assays, *CO, HY5, PRR5* and *FLM* cDNAs were cloned into pYFN43
659 and pYFC43 Gateway vectors (Belda-Palazón et al., 2012)☐ producing a N-terminal
660 fusion with nYFP and cYFP protein, respectively. *A. tumefaciens* strain GVG3101

pmp90 were transformed with these constructs, and 4-week old *N. benthamiana* plants were infiltrated. Later, leaf disks were visualized with the Leica TCS SP2/DMRE confocal microscope. YFP was excited with the 514 nm line of an argon laser at 20% strength, and fluorescence was detected between 515-565 nm. For ternary complex colocalization and FRET experiments, *HY5* and *PRR5* cDNAs were fused to C- and N-terminal fragments of YFP as in the BiFC assay, and *CO* and *FLM* cDNAs were cloned into pGWB644 and pGWB642 vectors, which produces a C-terminal fusion of full-length CFP and YFP, respectively (Shyu et al., 2008)□. CFP was excited with the 458 line under a Leica TCS SP2 confocal microscope, and the emission was detected between 465-479 nm. YFP was excited with a 514 nm laser, and the band-pass filters were adjusted to 520-545 nm. Colocalization was measured using the Pearson coefficient of correlation implemented in the *coloc2* plugin of the Fiji distribution of *ImageJ* (Schindelin et al., 2012)□.

Y2H assays

The DupLEX system was used to detect protein-protein interactions between CO domains and PRR5 or HY5. CO domains (CCT, middle and Bbox domains) were cloned into the bait vector pJG4-5, while full-length CDSs of *PRR5* or *HY5* were cloned into the prey vector pEG202. Primers used to generate Y2H clones are listed in Table S10. EGY48 cells (MATα *trp1 ura3 his3 LEU2*::pLex *Aop6-LEU2*) were used as the host strain for Y2H experiments (Gyuris et al. 1993). Single colonies grown on selection plates were inoculated in 3 ml of SD-Ura-His-Trp and grown overnight at 28ºC. Saturated culture was then used to make serial dilutions with an optical density at 600 nm (OD600) of 4^{-1}. Then, 2.5 µl of the solution was then spotted on an SD-Ura-His-Trp plate, with an SGalRaf-Ura-His-Trp-Leu plate as a growth control. Plates were imaged after incubation for 60-72 h at 30ºC.

Statistical analysis

The mean comparison between two sets of data was performed using the Student's t-test. The difference between means was considered statistically significant when the p value was less than 0.05 (marked with a single asterisk), 0.01 (marked with two asterisks) or 0.001 (marked with three asterisks). Fisher's exact test was used to analyze the significance of Venn diagram overlaps. Precisely, t.test and fisher.test

functions from R package stats were employed. To determine the significance of
overlapping peaks, Monte Carlo permutation tests were carried out, in which one of
the peak sets was randomly shuffled and new intersections were calculated after
each shuffle; 10000 iterations were computed in order to estimate p values. These p
values were corrected for multiple testing using the Benjamini-Hochberg method.
Statistically significant differences between means were evaluated using ANOVA R
package.

Figure Legends

Figure 1. CONSTANS regulates key circadian clock-related genes in the
transcriptional network CircadianFloralNet. (A) Graphical representation of
CircadianFloralTFNet, a derived network including only the analyzed transcription
factors and their interactions. Nodes are clustered in CLOCK, LIGHT or
FLOWERING. CO target genes are colored dark orange. (B) Most representative
network motifs found in CircadianFloralNet. (C) Feedback loop with multiple outputs
containing *CO* and *PRR5* as regulators with a set of common target genes involved
in the circadian clock and photoperiodic signaling.

Figure 2. CONSTANS alters the circadian clock system in *Arabidopsis*. (A) Venn
diagram representing downregulated genes in the *co-10* mutant (green) and
upregulated genes in *CO*-overexpressing plants (blue). The intersecting 148 genes
are considered high-fidelity CO regulated genes. (B) GO-term enrichment analysis
over the intersection between downregulated genes in *co-10* mutants and
upregulated genes in 35S:*CO* plants, indicating that they are significantly involved in
the rhythmic process (q-value = 7.838x2e-06). (C) *PRR5, PRR7, GIGANTEA* and
BAM9 expression levels during a LD time course in Col-0, 35S:*CO* and *co-10* lines.
The grey rectangle indicates the coincidence window in which the CO protein is
stabilized. Error bars represent the standard deviation within two biological replicates.
Three measurements were carried out for each biological replicate. (D) Relative
vertical motion of leaves in 35S:*CO* (light blue), Col-0 (dark blue) and *co-10* (green)
plants. Shading indicates the standard deviation. Subjective nights are represented in

grey. (E) Period distribution of rhythmic leaf movement in 35S:*CO*, Col-0 and *co-10*
plants from the previous image.

Figure 3. CO binds to clock gene promoters. (A) Pie graph representing the
genome-wide distribution of CO peaks. Most of the binding sites (64.87%) are
located in promoters (<=1 kb). (B) *De novo* DNA motif identification over the CO
binding sites using *HOMER* software. (C) ChIP-seq visualizer views of CONSTANS
occupancy of circadian related genes. The binding profile of immunoprecipitation and
input experiments are colored red and blue, respectively. Peaks detected by *MACS*
are indicated by blue rectangles. CCACA and G-box elements located at peaks are
also showed.

Figure 4. HY5, PRR5 and CO genome binding analysis. (A) Left: Venn diagram
showing the significant intersection (p-value < -10^6) of CO (green), HY5 (grey) and
PRR5 (red) targets. Right: DNA motifs enriched in the promoters of the different gene
sets. (B) Left: Graphic scheme of the peak intersection analysis. Right: DNA motif
enrichment analysis performed in each retrieved peak set. (C) Peak visualizer views
of HY5 (grey), PRR5 (red) and CO (green) binding to *CCA1* and *LHY* promoters. G-
box and CCACA elements are indicated by blue and red rectangles.

Figure 5. Genetic interaction between *CO* and *HY5*. (A) Pictures of 35S:*CO*, *hy5-2*
and 35S:*CO hy5* plants at DAG15. (B) Flowering time of WT, 35S:*CO*, *hy5-2* and
35S:*CO hy5* plants measured as total leaf number. (C) Expression of *CO* target
genes *CCA1, PRR5, PRR7* and *GI* in WT, 35S:*CO*, *hy5-2* and 35S:*CO hy5* plants at
ZT14, DAG 15. Error bars indicate the standard deviation (s.d.) from three
independent experiments. P < 0.05, one-way ANOVA and Tukey's HSD.

Figure 6. Protein interaction between CO, HY5 and PRR5. (A) Confocal images of
CO, HY5 and PRR5 interactions using BiFC assays in *Nicotiana* leaves. (B) Y2H
assays showing the interaction between different parts of CO (BBOX, middle and
CCT domains) with HY5 and PRR5. Panels show 3-day-old colonies grown in control
and selective media. (C) Colocalization experiments of CO, HY5 and PRR5 proteins.
YFP was reconstituted by fusing N- and C- terminal parts to HY5 and PRR5; the full-

length sequence was fused to CO. The colocalization between both fluorophores was measured by Pearson correlation compared to a negative control. (D) FRET analysis between CO, HY5 and PRR5. CO was fused to CFP and YFP was reconstituted by YFN:HY5/YFC:PRR5. CFP:FLM and YFP:FLM were used as negative controls. The proteins were coexpressed in *N. benthamiana* leaves, and normalized FRET (nFRET) was measured in 10 cells. (E) Immunodetection of the HY5 protein in chromatin immunoprecipitation experiments of 35S:*CO* and *co-10* plants using anti-CO antibodies.

Figure 7. Proposed model of the regulation of the circadian clock by CO. Under noninductive conditions, PRR5 binds to G-boxes in the promoter of its target genes and represses their expression. When CO accumulates (i.e. long days), it binds to PRR5 to stabilize it and occupies the same genome binding sites to activate the expression of its target genes. HY5 or another bZIP transcription factor could mediate CO and PRR5 binding to the DNA.

Figure S1. Transcriptional network analysis. (A) *Cytoscape* representation of CircadianFloralNet, a transcriptional network integrating ChIP-seq data from transcription factors involved in the circadian clock, flowering transition and floral development. (B) CircadianFloralTFNet, the regulator core composed only of the analyzed transcription factors and their interactions. (C) In-Degree distribution of CircadianFloralNet, following a power law. (D) Path distribution of CircadianFloralNet showing that the average path distance between two random nodes is 2.089. (E) Subnetwork representing the transcriptional interactions between the *CO-like* gene family and central circadian regulators (*CCA1*, *PRR7* and *PRR5*). (F) ChIP-QPCR of amplicons located in the *FT* and *PRR5* promoter with the CO protein.

Figure S2. (A) Normalized expression level in 35S:*CO* (yellow) and Col-0 (green) based on the RNA-seq experiments for circadian clock- and photoperiod-related genes. (B) Expression fold changes in 35S:*CO* and *co-10* plants compared to Col-0 (red and blue, respectively) for key circadian clock genes.

Figure S3. (A) Significant overlap between morning-phased cluster genes and the overexpressed genes in 35S:*CO* plants. The significance of the intersection was tested using Fisher's Exact Test. (B) Phase distribution of leaf movement in 35S:*CO* (light blue), Col-0 (dark blue) and *co-10* (green) plants.

Figure S4. (A) Venn diagram of the CO targets from ChIP-seq data and upregulated genes in 35S:*CO* plants. (B) Genome browser views showing CO binding to the *FT*, *PRR5*, *LHY*, *GI*, *RVE8*, *RVE4*, *CDF3* and *CDF6* promoters.

Figure S5. (A) Peak Visualizer views of CO, HY5, and PRRs (TOC1, PRR5, PRR7 and PRR9) binding to G-box-containing regions within the *LHY* and *GI* promoters. (B) HY5 binding to *PRR5*, *GI* and *FT* promoters. Here, 12-DAG-old plants grown under LD conditions were employed to perform ChIP assays. The bars represent the standard deviation between two biological replicates. Three technical replicates were measured for each biological replicate.

Figure S6. (A) *CO* and *HY5* expression levels measured by qRT-PCR in Col-0 (WT), 35S:*CO*, *hy5-2* and 35S:*CO* hy5 lines. (B) Levels of CO and HY5 proteins in Col-0 (WT), 35S:*CO, co-10, hy5-2* and 35S:*CO* hy5 plant extracts by Western blot using αCo and αHY5. Ponceau staining is shown as the loading control. Molecular masses are indicated on the left. Error bars indicate the standard deviation (s.d.) from three independent experiments. P < 0.05, one-way ANOVA and Tukey's HSD.

Figure S7. (A) Period of leaf movement in 35S:*CO*, Col-0 (WT), *co-10*, *hy5-2* and *prr5-1* plants grown in LL. (B) Phase of leaf movement in the same experiment as above.

Acknowledgements

Work by G. S.-B. was supported by a European Union contract LONGFLOW, MSCAIF-2018- 838317 and CSIC LONGFLOW, CONV_EXT_014. The financial support of the Spanish Ministry for Science and Innovations (MICINN/FEDER) PID2020-117018RB-I00 to F.V. is also acknowledged. We thank prof George

Coupland (MPIZ, Cologne, Germany) for discussion and critical reading of the
manuscript.

The authors declare no competing interests.

References

Abbas N, Maurya JP, Senapati D, Gangappa SN, Chattopadhyaya S (2014)
Arabidopsis CAM7 and HY5 physically interact and directly bind to the HY5 promoter
to regulate its expression and thereby promote photomorphogenesis. Plant Cell **26**:
1036–1052

Alon U (2007) Simplicity in biology. Nature **446**: 497

**Ando E, Ohnishi M, Wang Y, Matsushita T, Watanabe A, Hayashi Y, Fujii M, Ma
JF, Inoue SI, Kinoshita T** (2013) TWIN SISTER OF FT, GIGANTEA, and
CONSTANS have a positive but indirect effect on blue light-induced Stomatal
opening in Arabidopsis. Plant Physiol **162**: 1529–1538

Barabási A-L (2015) Network Science.

**Belda-Palazón B, Ruiz L, Martí E, Tárraga S, Tiburcio AF, Culiáñez F, Farràs R,
Carrasco P, Ferrando A** (2012) Aminopropyltransferases Involved in Polyamine
Biosynthesis Localize Preferentially in the Nucleus of Plant Cells. PLoS One **7**:
e46907

Binkert M, Kozma-Bognár L, Terecskei K, De Veylder L, Nagy F, Ulm R (2014)
UV-B-Responsive association of the Arabidopsis bZIP transcription factor
ELONGATED HYPOCOTYL5 with target genes, including its own promoter. Plant
Cell **26**: 4200–4213

Blackman BK, Michaels SD (2010) Does CONSTANS act as a transcription factor
or as a co-activator? The answer may be - yes. New Phytol **187**: 1–3

**Bowler C, Benvenuto G, Laflamme P, Molino D, Probst A V., Tariq M,
Paszkowski J** (2004) Chromatin techniques for plant cells. Plant J **39**: 776–789

Bursch K, Toledo-Ortiz G, Pireyre M, Lohr M, Braatz C, Johansson H (2020)
Identification of BBX proteins as rate-limiting cofactors of HY5. Nat Plants **6**: 921-928

Chattopadhyay S, Ang LH, Puente P, Deng XW, Wei N (1998) Arabidopsis bZIP
protein HY5 directly interacts with light-responsive promoters in mediating light
control of gene expression. Plant Cell **10**: 673–683

Csardi G, Nepusz T (2006) The igraph software package for complex network
research. Inter J Complex Syst 1695: 1-9.

Datta S, Hettiarachchi C, Johansson H, Holm M (2007) SALT TOLERANCE
HOMOLOG2, a B-box protein in Arabidopsis that activates transcription and
positively regulates light-mediated development. Plant Cell **19**: 3242–3255

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut,
P., Chaisson, M. & Gingeras, T. R (2013). STAR: ultrafast universal RNA-seq
aligner. *Bioinformatics*, *29*(1), 15-21.

Endo M, Shimizu H, Nohales MA, Araki T, Kay SA (2014) Tissue-specific clocks in
Arabidopsis show asymmetric coupling. Nature **515**: 419–422

Endo M, Tanigawa Y, Murakami T, Araki T, Nagatani A (2013) Phytochrome-
dependent late-flowering accelerates flowering through physical interactions with
phytochrome B and CONSTANS. Proc Natl Acad Sci U S A **110**: 18017–18022

Fernández V, Takahashi Y, Le Gourrierec J, Coupland G (2016) Photoperiodic and
thermosensory pathways interact through CONSTANS to promote flowering at high
temperature under short days. Plant J **86**: 426-440.

Gangappa SN, Botto JF (2016) The Multifaceted Roles of HY5 in Plant Growth and
Development. Mol Plant **9**: 1353–1365

Gnesutta N, Kumimoto RW, Swain S, Chiara M, Siriwardana C, Horner DS, Holt
BF, Mantovani R (2017) CONSTANS imparts DNA sequence specificity to the
histone fold NF-YB/NF-YC Dimer. Plant Cell **29**: 1516–1532

Goldstein, M. L., Morris, S. A., & Yen, G. G. (2004). Problems with fitting to the
power-law distribution. *The European Physical Journal B-Condensed Matter and*
Complex Systems, *41*(2), 255-258.

Graeff M, Straub D, Eguen T, Dolde U, Rodrigues V, Brandt R, Wenkel S (2016)
MicroProtein-Mediated Recruitment of CONSTANS into a TOPLESS Trimeric
Complex Represses Flowering in Arabidopsis. PLoS Genet **12**: e1005959

Greenham K, Lou P, Remsen SE, Farid H, McClung CR (2015) TRiP: Tracking
Rhythms in Plants, an automated leaf movement analysis program for circadian
period estimation. Plant Methods. doi: 10.1186/S13007-015-0075-5

Hajdu A, Ádám É, Sheerin DJ, Dobos O, Bernula P, Hiltbrunner A, Kozma-
Bognár L, Nagy F (2015) High-level expression and phosphorylation of phytochrome
B modulates flowering time in Arabidopsis. Plant J **83**: 794–805

Hajdu A, Dobos O, Bernula P, Domijan M, Bálint B, Nagy I, Nagy F, Kozma-
Bognár L (2018) ELONGATED HYPOCOTYL 5 mediates blue light signalling to the
Arabidopsis circadian clock. Plant J **96**: 1242-1254

Hayama R, Sarid-Krebs L, Richter R, Fernández V, Jang S, Coupland G (2017)
PSEUDO RESPONSE REGULATORs stabilize CONSTANS protein to promote
flowering in response to day length. EMBO J e201693907

Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C,
Singh H, Glass CK (2010) Simple combinations of lineage-determining transcription
factors prime cis-regulatory elements required for macrophage and B cell identities.
Mol Cell **38**: 576–89

Huang W, Pérez-García P, Pokhilko A, Millar AJ, Antoshechkin I, Riechmann JL,
Mas P (2012) Mapping the core of the Arabidopsis circadian clock defines the
network structure of the oscillator. Science (80-) **335**: 75–79

Imaizumi T, Schultz TF, Harmon FG, Ho LA, Kay SA (2005) Plant science: FKF1 F-
box protein mediates cyclic degradation of a repressor of CONSTANS in Arabidopsis.
Science (80-) **309**: 293–297

Isabel Ortiz-Marchena M, Albi T, Lucas-Reina E, Said FE, Romero-Campero FJ,
Cano B, Teresa Ruiz M, Romero JM, Valverde F (2014) Photoperiodic control of
carbon distribution during the floral transition in Arabidopsis. Plant Cell **26**: 565–584

Ito S, Song YH, Josephson-Day AR, Miller RJ, Breton G, Olmstead RG,
Imaizumi T (2012) FLOWERING BHLH transcriptional activators control expression

of the photoperiodic flowering regulator CONSTANS in Arabidopsis. Proc Natl Acad
Sci U S A **109**: 3582–3587

Jang S, Marchal V, Panigrahi KCS, Wenkel S, Soppe W, Deng XW, Valverde F,
Coupland G (2008) Arabidopsis COP1 shapes the temporal pattern of CO
accumulation conferring a photoperiodic flowering response. EMBO J **27**: 1277–1288

Jing Y, Guo Q, Zha P, Lin R (2019) The chromatin-remodelling factor PICKLE
interacts with CONSTANS to promote flowering in Arabidopsis. Plant Cell Environ **42**:
2495–2507

Jing Y, Zhang D, Wang X, Tang W, Wang W, Huai J, Xu G, Chen D, Li Y, Lin R
(2013) Arabidopsis chromatin remodeling factor PICKLE interacts with transcription
factor HY5 to regulate hypocotyl cell elongation. Plant Cell **25**: 242–256

Job N, Yadukrishnan P, Bursch K, Datta S, Johansson H (2018) Two B-Box
proteins regulate photomorphogenesis by oppositely modulating HY5 through their
diverse C-terminal domains. Plant Physiol **176**: 2963–2976

Kinoshita A, Richter R (2020) Genetic and molecular basis of floral induction in
Arabidopsis thaliana. J Exp Bot **71**: 2490–2504

Langmead B, Trapnell C, Pop M, Salzberg SL (2009) Ultrafast and memory-
efficient alignment of short DNA sequences to the human genome. Genome Biol **10**:
R25

Lau OS, Bergmann DC (2015) MOBE-ChIP: a large-scale chromatin
immunoprecipitation assay for cell type-specific studies. Plant J **84**: 443–50

Laubinger S, Marchal V, Gentillhomme J, Wenkel S, Adrian J, Jang S, Kulajta C,
Braun H, Coupland G, Hoecker U (2006) Arabidopsis SPA proteins regulate
photoperiodic flowering and interact with the floral inducer CONSTANS to regulate its
stability. Development **133**: 3213–3222

Lazaro A, Mouriz A, Piñeiro M, Jarillo JA (2015) Red Light-Mediated Degradation
of CONSTANS by the E3 Ubiquitin Ligase HOS1 Regulates Photoperiodic Flowering
in Arabidopsis. Plant Cell **27**: 2437–54

Lazaro A, Valverde F, Piñeiro M, Jarillo JA (2012) The Arabidopsis E3 ubiquitin
ligase HOS1 negatively regulates CONSTANS abundance in the photoperiodic
control of flowering. Plant Cell **24**: 982–999

Lee J, He K, Stolc V, Lee H, Figueroa P, Gao Y, Tongprasit W, Zhao H, Lee I,
Deng XW (2007) Analysis of transcription factor HY5 genomic binding sites revealed
its hierarchical role in light regulation of development. Plant Cell **19**: 731–49

Liao, Y., Smyth, G. K., & Shi, W (2014). featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. *Bioinformatics*, *30*(7),
923-930.

Liu B, Zuo Z, Liu H, Liu X, Lin C (2011) Arabidopsis cryptochrome 1 interacts with
SPA1 to suppress COP1 activity in response to blue light. Genes Dev **25**: 1029–1034

Liu LJ, Zhang YC, Li QH, Sang Y, Mao J, Lian HL, Wang L, Yang HQ (2008)
COP1-mediated ubiquitination of CONSTANS is implicated in cryptochrome
regulation of flowering in Arabidopsis. Plant Cell **20**: 292–306

Liu T, Carlsson J, Takeuchi T, Newton L, Farré EM (2013) Direct regulation of
abiotic responses by the Arabidopsis circadian clock component PRR7. Plant J **76**:
101–14

Liu TL, Newton L, Liu M-J, Shiu S-H, Farré EM (2016) A G-Box-Like Motif Is
Necessary for Transcriptional Regulation by Circadian Pseudo-Response Regulators
in Arabidopsis. Plant Physiol **170**: 528–539

de los Reyes P, Romero-Campero FJ, Teresa Ruiz M, Romero JM, Valverde F
(2017) Evolution of daily gene co-expression patterns from algae to plants. Front
Plant Sci **8**: 1217

Salmon-Divon, M., Dvinge , H., Tammoja, K., Bertone, P (2010) PeakAnalyzer:
genome-wide annotation of chromatin binding and modification loci. BMC
Bioinformatics. doi: 10.1186/1471-2105-11-415

Martín G, Rovira A, Veciana N, Soy J, Toledo-Ortiz G, Gommers CMM, Boix M,
Henriques R, Minguet EG, Alabadí D, et al (2018) Circadian Waves of
Transcriptional Repression Shape PIF-Regulated Photoperiod-Responsive Growth in
Arabidopsis. Curr Biol **28**: 311-318.e5

McClung CR (2014) Wheels within wheels: New transcriptional feedback loops in the
Arabidopsis circadian clock. F1000Prime Rep. doi: 10.12703/P6-2

McClung CR (2006) Plant circadian rhythms. Plant Cell **18**: 792–803

Mellis IA, Raj A (2015) Half dozen of one, six billion of the other: What can small-
and large-scale molecular systems biology learn from one another? Genome Res **25**:
1466–1472

De Mendoza A, Sebé-Pedrós A, Šestak MS, Matejčić M, Torruella G, Domazet-
Lošo T, Ruiz-Trillo I (2013) Transcription factor evolution in eukaryotes and the
assembly of the regulatory toolkit in multicellular lineages. Proc Natl Acad Sci U S A
110: E4858–E4866

Michael, T. P., Salome, P. A., Yu, H. J., Spencer, T. R., Sharp, E. L., McPeek, M.
A., ... & McClung, C. R. (2003). Enhanced fitness conferred by naturally occurring
variation in the circadian clock. *Science*, *302*(5647), 1049-1053.

Milo R, Shen-Orr S, Itzkovitz S, Kashtan N, Chklovskii D, Alon U (2002) Network
motifs: Simple building blocks of complex networks. Science (80-) **298**: 824–827

Mizoguchi T, Wright L, Fujiwara S, Cremer F, Lee K, Onouchi H, Mouradov A,
Fowler S, Kamada H, Putterill J, et al (2005) Distinct roles of GIGANTEA in
promoting flowering and regulating circadian rhythms in Arabidopsis. Plant Cell **17**:
2255–2270

Mortazavi A, Williams BA, McCue K, Schaeffer L, Wold B (2008) Mapping and
quantifying mammalian transcriptomes by RNA-Seq. Nat Methods **5**: 621–628

Nakamichi N, Kiba T, Henriques R, Mizuno T, Chua N-H, Sakakibara H (2010)
PSEUDO-RESPONSE REGULATORS 9, 7, and 5 are transcriptional repressors in
the Arabidopsis circadian clock. Plant Cell **22**: 594–605

Nakamichi N, Kiba T, Kamioka M, Suzuki T, Yamashino T, Higashiyama T,
Sakakibara H, Mizuno T (2012) Transcriptional repressor PRR5 directly regulates
clock-output pathways. Proc Natl Acad Sci U S A **109**: 17123–8

Nakamichi N, Kita M, Niinuma K, Ito S, Yamashino T, Mizoguchi T, Mizuno T
(2007) Arabidopsis Clock-Associated Pseudo-Response Regulators PRR9, PRR7

and PRR5 Coordinately and Positively Regulate Flowering Time Through the
Canonical CONSTANS-Dependent Photoperiodic Pathway. Plant Cell Physiol **48**:
822–832

Nguyen KT, Park J, Park E, Lee I, Choi G (2015) The Arabidopsis RING Domain
Protein BOI Inhibits Flowering via CO-dependent and CO-independent Mechanisms.
Mol Plant **8**: 1725–1736

Onouchi, H., Igeño, M. I., Périlleux, C., Graves, K., & Coupland, G. (2000).
Mutagenesis of plants overexpressing CONSTANS demonstrates novel interactions
among Arabidopsis flowering-time genes. *The Plant Cell*, *12*(6), 885-900.

Osterlund MT, Hardtke CS, Ning W, Deng XW (2000) Targeted destabilization of
HY5 during light-regulated development of Arabidopsis. Nature **405**: 462–466

Parsons R, Parsons R, Garner N, Oster H, Rawashdeh O (2020) CircaCompare: A
method to estimate and statistically support differences in mesor, amplitude and
phase, between circadian rhythms. Bioinformatics **36**: 1208–1212

Pertea M, Kim D, Pertea GM, Leek JT, Salzberg SL (2016) Transcript-level
expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown.
Nat Protoc **11**: 1650–1667

Piñeiro M, Jarillo JA (2013) Ubiquitination in the control of photoperiodic flowering.
Plant Sci **198**: 98–109

Putterill J, Robson F, Lee K, Simon R, Coupland G (1995) The CONSTANS gene
of arabidopsis promotes flowering and encodes a protein showing similarities to zinc
finger transcription factors. Cell **80**: 847–857

Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK (2015) limma
powers differential expression analyses for RNA-sequencing and microarray studies.
Nucleic Acids Res **43**: e47–e47

Robinson, M. D., McCarthy, D. J., & Smyth, G. K (2010). edgeR: a Bioconductor
package for differential expression analysis of digital gene expression data.
bioinformatics, *26*(1), 139-140.

Romero-Campero FJ, Lucas-Reina E, Said FE, Romero JM, Valverde F (2013) A
contribution to the study of plant development evolution based on gene co-
expression networks. Front Plant Sci **4**: 291

Romero JM, Valverde F (2009) Evolutionarily conserved photoperiod mechanisms in
plants: When did plant photoperiodic signaling appear? Plant Signal Behav **4**: 642–
644

Rugnone ML, Faigón Soverna A, Sanchez SE, Schlaen RG, Hernando CE,
Seymour DK, Mancini E, Chernomoretz A, Weigel D, Más P, et al (2013) LNK
genes integrate light and clock signaling networks at the core of the Arabidopsis
oscillator. Proc Natl Acad Sci U S A **110**: 12120–5

Samach A, Onouchi H, Gold SE, Ditta GS, Schwarz-Sommer Z, Yanofsky MF,
Coupland G (2000) Distinct roles of constans target genes in reproductive
development of Arabidopsis. Science (80-) **288**: 1613–1616

Sanchez SE, Kay SA (2016) The Plant Circadian Clock: From a Simple Timekeeper
to a Complex Developmental Manager. Cold Spring Harb Perspect Biol **8**: a027748

Sawa M, Nusinow DA, Kay SA, Imaizumi T (2007) FKF1 and GIGANTEA complex
formation is required for day-length measurement in Arabidopsis. Science (80-) **318**:
261–265

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T,
Preibisch S, Rueden C, Saalfeld S, Schmid B, et al (2012) Fiji: An open-source
platform for biological-image analysis. Nat Methods **9**: 676–682

Serrano-Bueno G, Said FE, los Reyes P, Lucas-Reina EI, Ortiz-Marchena MI,
Romero JM, Valverde F (2020) CONSTANS–FKBP12 interaction contributes to
modulation of photoperiodic flowering in Arabidopsis. Plant J **101**: 1287-1302

Serrano-Bueno G, Sánchez de Medina Hernández V, Valverde F (2021)
Photoperiodic Signaling and Senescence, an Ancient Solution to a Modern Problem?
Front Plant Sci **12**: 634393

Serrano-Bueno G, de los Reyes P, Chini A, Ferreras-Garrucho G,Sánchez de
Medina Hernández V, Boter M, Solano R, Valverde F (2022) Regulation of floral

senescence in Arabidopsis by coordinated action of CONSTANS and jasmonate signaling. Mol Plant **15**: 1710-1724.

Sheerin DJ, Menon C, Oven-Krockhaus S Zur, Enderle B, Zhu L, Johnen P, Schleifenbaum F, Stierhof YD, Huq E, Hiltbrunner A (2015) Light-activated phytochrome A and B interact with members of the SPA family to promote photomorphogenesis in arabidopsis by reorganizing the COP1/SPA complex. Plant Cell **27**: 189–201

Shen C, Liu H, Guan Z, Yan J, Zheng T, Yan W, Wu C, Zhang Q, Yin P, Xing Y (2020) Structural Insight into DNA Recognition by CCT/NF-YB/YC Complexes in Plant Photoperiodic Flowering. Plant Cell **32**: 3469-3484

Shen-Orr SS, Milo R, Mangan S, Alon U (2002) Network motifs in the transcriptional regulation network of Escherichia coli. Nat Genet **31**: 64–68

Shim JS, Kubota A, Imaizumi T (2017) Circadian Clock and Photoperiodic Flowering in Arabidopsis: CONSTANS Is a Hub for Signal Integration. Plant Physiol **173**: 5-15.

Shyu YJ, Suarez CD, Hu CD (2008) Visualization of ternary complexes in living cells by using a BiFC-based FRET assay. Nat Protoc **3**: 1693–1702

Smoot ME, Ono K, Ruscheinski J, Wang P-L, Ideker T (2011) Cytoscape 2.8: new features for data integration and network visualization. Bioinformatics **27**: 431–2

Song YH, Estrada DA, Johnson RS, Kim SK, Lee SY, MacCoss MJ, Imaizumi T (2014) Distinct roles of FKF1, GIGANTEA, and ZEITLUPE proteins in the regulation of constans stability in Arabidopsis photoperiodic flowering. Proc Natl Acad Sci U S A **111**: 17672–17677

Song YH, Smith RW, To BJ, Millar AJ, Imaizumi T (2012) FKF1 conveys timing information for CONSTANS stabilization in photoperiodic flowering. Science (80-) **336**: 1045–1049

Stone L, Simberloff D, Artzy-Randrup Y (2019) Network motifs and their origins. PLOS Comput Biol **15**: e1006749

Suárez-López P, Wheatley K, Robson F, Onouchi H, Valverde F, Coupland G (2001) CONSTANS mediates between the circadian clock and the control of flowering in Arabidopsis. Nature **410**: 1116–1120

Thaben PF, Westermark PO (2014) Detecting rhythms in time series with RAIN. J Biol Rhythms **29**: 391–400

Tiwari SB, Shen Y, Chang H-C, Hou Y, Harris A, Ma SF, McPartland M, Hymus GJ, Adam L, Marion C, et al (2010) The flowering time regulator CONSTANS is recruited to the FLOWERING LOCUS T promoter via a unique cis-element. New Phytol **187**: 57–66

Tong, A. H. Y., Lesage, G., Bader, G. D., Ding, H., Xu, H., Xin, X., ... & Boone, C. (2004). Global mapping of the yeast genetic interaction network. *science*, *303*(5659), 808-813.

Valverde F (2011) CONSTANS and the evolutionary origin of photoperiodic timing of flowering. J Exp Bot **62**: 2453–63

Valverde F, Mouradov A, Soppe W, Ravenscroft D, Samach A, Coupland G (2004) Photoreceptor Regulation of CONSTANS Protein in Photoperiodic Flowering. Science (80-) **303**: 1003–1006

Wagner, A., & Fell, D. A. (2001). The small world inside large metabolic networks. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *268*(1478), 1803-1810.

Wang CQ, Guthrie C, Sarmast MK, Dehesh K (2014) BBX19 interacts with CONSTANS to repress FLOWERING LOCUS T transcription, defining a flowering time checkpoint in Arabidopsis. Plant Cell **26**: 3589–3602

Wang H, Pan J, Li Y, Lou D, Hu Y, Yu D (2016) The DELLA-CONSTANS transcription factor cascade integrates gibberellic acid and photoperiod signaling to regulate flowering. Plant Physiol **172**: 479–488

Wenkel S, Turck F, Singer K, Gissot L, Le Gourrierec J, Samach A, Coupland G (2006) CONSTANS and the CCAAT box binding complex share a functionally important domain and interact to regulate flowering of Arabidopsis. Plant Cell **18**: 2971–84

Wheeler DL, Barrett T, Benson DA, Bryant SH, Canese K, Church DM, DiCuccio M, Edgar R, Federhen S, Helmberg W, et al (2005) Database resources of the National Center for Biotechnology Information. Nucleic Acids Res **33**: D39-45

Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, Feng T, Zhou L, Tang W, Zhan L, et al (2021) clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. Innov **2**: 100141

Xu F, Li T, Xu P-B, Li L, Du S-S, Lian H-L, Yang H-Q (2016) DELLA proteins physically interact with CONSTANS to regulate flowering under long days in *Arabidopsis*. FEBS Lett **590**: 541–549

Yanovsky MJ, Kay SA (2002) Molecular basis of seasonal time measurement in Arabidopsis. Nature **419**: 308–312

Yook, S. H., Oltvai, Z. N., & Barabási, A. L. (2004). Functional and topological characterization of protein interaction networks. *Proteomics*, *4*(4), 928-942.

Young HS, Cheol MY, An PH, Seong HK, Hee JJ, Su YS, Hye JK, Yun DJ, Chae OL, Jeong DB, et al (2008) DNA-binding study identifies C-box and hybrid C/G-box or C/A-box motifs as high-affinity binding sites for STF1 and long hypocotyl5 proteins. Plant Physiol **146**: 1862–1877

Yu G, Wang L-G, He Q-Y (2015) ChIPseeker: an R/Bioconductor package for ChIP peak annotation, comparison and visualization. Bioinformatics **31**: 2382–2383

Zhang H, He H, Wang X, Wang X, Yang X, Li L, Deng XW (2011) Genome-wide mapping of the HY5-mediated genenetworks in Arabidopsis that involve both transcriptional and post-transcriptional regulation. Plant J **65**: 346–358

Zhang JD, Ruschhaupt M, Biczok R (2016) ddCt method for qRT–PCR data analysis.

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM, Brown M, Li W, et al (2008) Model-based analysis of ChIP-Seq (MACS). Genome Biol **9**: R137

Zhu LJ, Gazin C, Lawson ND, Pagès H, Lin SM, Lapointe DS, Green MR (2010)

ChIPpeakAnno: a Bioconductor package to annotate ChIP-seq and ChIP-chip data.

BMC Bioinforma 2010 111 **11**: 1–10

Zielinski T, Moore AM, Troup E, Halliday KJ, Millar AJ (2014) Strengths and

Limitations of Period Estimation Methods for Circadian Data. PLoS One **9**: e96462

Zuo Z, Liu H, Liu B, Liu X, Lin C (2011) Blue light-dependent interaction of CRY2

with SPA1 regulates COP1 activity and floral initiation in arabidopsis. Curr Biol **21**:

841–847

CONSTANS alters the circadian clock in *Arabidopsis thaliana*

Pedro de los Reyes[1], Francisco J Romero-Campero[1,2], He Gao[3], Gloria Serrano-Bueno[2], Jose M Romero[1,4] and Federico Valverde[1]*

[1] Plant Development Unit - Institute for Plant Biochemistry and Photosynthesis, Consejo Superior de Investigaciones Científicas, Universidad de Sevilla, Seville, Spain,
[2] Department of Computer Science and Artificial Intelligence, Universidad de Sevilla, Seville, Spain
[3] Department of Plant Developmental Biology, Max Planck Institute for Plant Breeding Research, Cologne, Germany
[4]Department of Biochemistry and Molecular Biology, Universidad de Sevilla, Seville, Spain.

A

B

Motif name	Auto-regulation	Feedback loop	Feedback loop with auto-regulation	Gene with multiple output	Feedback loop with output	Feedforward/ Feedback loop	Triple Feedback loop
Motif image							
Appearances on CFnet	24	62	56	494	235	187	50
Appearances on random graphs (mean ± sd)	10.32 ± 2.63	51.67 ± 5.03	5.025 ± 3.264	358 ± 29.96	73.93 ± 8.53	66.64 ± 12.58	4.98 ± 2.61
p-value	$<10^{-5}$	$1.65×10^{-2}$	$<10^{-5}$	$<10^{-5}$	$<10^{-5}$	$<10^{-5}$	$<10^{-5}$

C

A

Promoter (<=1kb) (64.87%)
Promoter (1-2kb) (9.4%)
5' UTR (0.12%)
3' UTR (3.48%)
1st Exon (4.2%)
Other Exon (3.14%)
1st Intron (0.65%)
Other Intron (0.9%)
Downstream (<=300) (6.25%)
Distal Intergenic (6.97%)

B

Motif	p-value	% of targets	% of background
CCACATCCATT	1e-19	38.24%	9.03%
ACACGTGGCT	1e-18	27.94%	4.64%
TTTGTGTGGATA	1e-13	13.97%	1.28%

C

CCA1

LHY

RVE8

PRR9

PRR5

FT

▲ G-box - CACGTG
▲ CCACA-box - CCACA

A

B

C

A

B

A

morning peak and
evening trough
cluster genes

p-value < 2.2e-16

1668   269   1064

35S:CO Activated
Genes at ZT16

B

A

B

A

B

A

B